Substrate ligand density modulates gap junction intercellular communication during mesenchymal cell condensation

Ignasi Casanellas[a,b,c], Anna Lagunas[c,a]*, Yolanda Vida[d,e], Ezequiel Pérez-Inestrosa[d,e], Cristina Rodríguez-Pereira[f], Joana Magalhaes[c,f], José A. Andrades[g,c,e], José Becerra[g,c,e], Josep Samitier[a,b,c]

[a]Institute for Bioengineering of Catalonia (IBEC), Barcelona Institute of Science and Technology (BIST). c/ Baldiri Reixac, 10-12, 08028 Barcelona, Spain.

[b]Department of Electronics and Biomedical Engineering, University of Barcelona (UB), Faculty of Physics. c/ Martí i Franquès, 1, 08028 Barcelona, Spain.

[c]Biomedical Research Networking Center in Bioengineering, Biomaterials, and Nanomedicine (CIBER-BBN). Av. Monforte de Lemos, 3-5. Pabellón 11. Planta 0, 28029 Madrid, Spain.

[d]Universidad de Málaga - IBIMA, Dpto. Química Orgánica. Campus de Teatinos s/n, 29071 Málaga, Spain.

[e]Centro Andaluz de Nanomedicina y Biotecnología-BIONAND. Parque Tecnológico de Andalucía, c/ Severo Ochoa 35, 29590 Campanillas, Málaga, Spain.

[f]Unidad de Medicina Regenerativa, Grupo de Investigación en Reumatología (GIR), Instituto de Investigación Biomédica de A Coruña (INIBIC), Complexo Hospitalario Universitario de A Coruña (CHUAC), Sergas, Universidade da Coruña (UDC). c/ Xubias de Arriba, 84, 15006 A Coruña, Spain.

[g]Department of Cell Biology, Genetics and Physiology, Universidad de Málaga (UMA), Instituto de Investigación Biomédica de Málaga (IBIMA). Av. Cervantes, 2, 29071 Málaga, Spain.

*Corresponding author: alagunas@ibecbarcelona.eu

Abstract

Gap junction intercellular communication (GJIC) provides a continuous and efficient flow of biological information during tissue formation and is essential to sustain homeostasis and function in living organisms. Cell-matrix interactions have been widely addressed and their influence on tissue organization is recognized. However, how extracellular matrix (ECM) adhesion affects intercellular communication during tissue formation remains elusive. Here we use substrates with uneven nanopatterns of adhesive ligand arginine-glycine-aspartic acid (RGD) to control cell adhesion during mesenchymal condensation, a prevalent morphogenetic transition. We show that the establishment of GJIC is an adhesion-gated mechanism, which is dynamically regulated. Substrate effects continuously propagate into the forming tissue through actomyosin contraction, affecting both the 3D architecture and functionality of the GJIC network developing in prechondrogenic condensates.

Keywords: Dendrimer-based nanopatterning, arginine-glycine-aspartic acid (RGD), cell adhesion, mesenchymal stem cells (MSCs), condensation, gap junction intercellular communication (GJIC).

1. Introduction

Mesenchymal cell condensation is a prevalent morphogenetic transition regulated by cell adhesion in which mesenchymal stem cells (MSCs) gather to form intimate cell-to-cell contacts [1]. The condensation phase was named "membranous skeleton" by Grüneberg to stress its distinctive function in skeletal development [2]. During skeletogenesis, cell density increases locally at the condensation sites by means of extracellular matrix (ECM)-driven active cell movement,[1] with an important role of the ECM ubiquitous protein fibronectin (FN), which is upregulated during mesenchymal condensation [3,4]. Cell condensates progressively evolve into cartilaginous nodules, setting the bases for cartilage formation.

In osteochondral development, mesenchymal cell condensation is concurrent with the formation of an extensive gap junction (GJ) communication network [5]. GJs are plasma membrane channels that provide intercellular and cell-matrix communication in almost all animal tissues, allowing cells to exchange ions and small molecules through a controlled gating mechanism [6]. During morphogenesis, an efficient network of GJs is an extremely versatile communication system that mediates the rapid synchronization between cells. GJs established during embryonic patterning allow multicellular groups to coordinate towards the formation of supracellular, tissue-level structures. Avascular tissues, such as cartilage, particularly rely on this form of intercellular communication for successful development [7–9].

Although previous studies have related gap junction intercellular communication (GJIC) with matrix-associated proteins [10–13], little is known about how extracellular inputs modulate the formation of intercellular GJ networks and the associated implications in tissue architecture and function. Integrin-mediated cell-matrix interactions have been shown to regulate many biological processes such as cell shape, proliferation, migration, differentiation and programmed cell death [14–16]. During morphogenesis, dynamic adhesion mechanisms, together with the associated regulatory signalling pathways, define tissue differentiation and architecture, and modulate collective cell behaviour [17–19].

Cell adhesion is governed at the nanoscale, as evidenced by the assembly of collagen fibres in the ECM or the folding/unfolding of fibronectin [20–22], and also by the compartmentalization of the cell membrane receptors into nanoclusters, which facilitates the allosteric regulation, increases the ligand rebinding probability and triggers the assembly of signalling complexes in the cytosol [23]. The nanopatterning of ECM motifs for the study of cell-surface interactions at the nanoscale highlighted the relevance of ECM ligand presentation to cells. Experiments using micellar lithography to create nanopatterns of the cell adhesive peptide arginine-glycine-aspartic acid (RGD) of fibronectin (FN), unveiled a nanospacing threshold of around 70 nm for an efficient cell adhesion on stiff substrates, and demonstrated that cell adhesion is more affected by the local rather than the global ligand concentrations

[24–27]. Since these first works in the field, others have emerged showing that not only cell adhesion, but

also many other cell responses are affected by the nanoscale ligand presentation [28].

In previous works, we have shown that a dendrimer-based RGD nanopatterning technique produces

uneven nanopatterns of tuneable local surface adhesiveness with liquid-like order and defined spacing on

large areas, thus being fully compatible with standard cell culture protocols [29]. Owing to steric

hindrance restrictions, each dendrimer of 4-5 nm in diameter, although bearing up to eight copies of the

RGD ligand, provides a single site for integrin receptor binding. Therefore, dendrimer nanopattern

configuration directly correlates with the RGD available for cell adhesion. The local surface adhesiveness

in the nanopatterns is obtained by quantifying the percentage of surface area with a minimum interparticle

distance below the 70 nm threshold for an efficient cell adhesion. These nanopatterns proved to sustain

cell adhesion more efficiently than the corresponding homogeneous surfaces [29] and improve cell

differentiation [30–32].

Taking chondrogenesis as a model [33], we used the RGD dendrimer-based nanopatterns to conduct a

systematic study of the influence of local surface adhesiveness on the establishment of GJIC network

during mesenchymal 3D condensation. Results show that local surface adhesiveness triggers cell

condensate responses from a threshold nanopattern configuration and we demonstrate that the adhesive

substrate information, which is transduced by integrins, propagates within cell condensates in a

continuous feedback mode through cytoskeletal contractility.

2. Methods

2.1. Production of nanopatterned substrates

All steps were performed in a sterile tissue culture hood, and only sterile materials, solutions and

techniques were used. All dendrimer solutions were sonicated and filtered through a Millex RB sterile

syringe filter (Merk Millipore) prior to use, and stock solutions were used within 6 months of preparation.

Nanopatterned substrates were prepared as previously described [30,31]. Briefly, a 95/5 L-lactide/DL-

lactide copolymer (Corbion) 2% m v^{-1} solution in dry 1,4-dioxane (Sigma-Aldrich, 296309) was spin-

coated at 3000 rpm for 30 s on 1.25×1.25 cm Corning® glass microslides (Sigma-Aldrich). Deionized

water (18 MΩ·cm Milli-Q, Millipore) was used for rinsing samples and to prepare RGD-functionalized

dendrimers' working solutions at 2.5 10^{-8}, 10^{-8} and 4 10^{-9}% w w^{-1} concentrations. Spin-coated on poly(L-

lactic acid) (PLLA) substrates were treated for 13 min under UV light and immersed in dendrimer

solutions for 16 h (pH = 5.6, T = 293 K). Then, the nanopatterned substrates were rinsed with copious

amounts of water and dried. Positive controls (S_{FN}) were obtained by incubating spin-coated PLLA
substrates with fibronectin (100 μg mL^{-1}) from bovine plasma-solution (Sigma-Aldrich, F1141).
Nanopatterned substrates were imaged by atomic force microscopy (AFM) in a Dimension 3100 AFM
instrument (Veeco Instruments) operated in tapping mode at room temperature in air. Silicon AFM probes
(Budget Sensors) with a spring constant k = 40 N m^{-1} and a resonant frequency ν = 300 kHz were used. At
least three images of 5x5 μm were taken per substrate of three independent substrates per condition
(initial dendrimer concentration in solution). The AFM height images were processed with WSxM 4.0
software [34]. Corresponding image thresholds were obtained manually and processed with ImageJ
software (NIH). The resulting particle positions were used to obtain the minimum interparticle distances
(d_{min}) using a custom-generated MATLAB code (The MATHWORKS, Inc.), and the corresponding
probability contour plots for d_{min} were constructed using an adapted MATLAB code [29]. At least three
images from independent samples were computed. Quantification of the percentage of the area with a d_{min}
threshold below 70 nm in the d_{min} probability contour plots was performed by manually selecting the
regions and processing them with ImageJ.
116
117 *2.2. Cell culture*
118
Human adipose-derived MSCs (ATCC, PCS-500-011) were cultured at 37°C and 5% CO_2 in MSC Basal
Medium (ATCC, PCS-500-030) supplemented with MSC Growth Kit Low Serum (ATCC, PCS-500-
040). Medium was replaced every 2 days. Passaging was carried out when cells reached 70-80%
confluence. For the experiments, cells were trypsinized at passages 3 to 4, counted, resuspended in
chondrogenesis-inducing medium Chondrocyte Differentiation Tool (ATCC, PCS-500-051) with 0.1% v
v^{-1} penicillin-streptomycin (Invitrogen, 15140), and seeded on nanopatterned and control substrates at a
density of 2,760 cells cm^{-2}. 3 replicates of each condition were seeded. Medium was replaced every 3
days.
127
128 *2.3. Immunostaining*
129
After 6 and 9 days of chondrogenic differentiation on the nanopatterns and controls, cells were carefully
rinsed with PBS (Gibco, 21600-10), fixed with Formalin Solution (Sigma, HT5011) for 20 min at room
temperature, and rinsed again twice with PBS. Aldehyde groups were blocked with 50 mM ammonium
chloride (Sigma, A9434) in PBS for 20 min. Samples were permeabilized with saponin (Sigma, 47036)
0.1% m v^{-1} in BSA (Sigma, A3059) 1% m/v in PBS for 10 min.

For connexin 43 (Cx43) and cell nuclei observation in cell condensates, samples were stained with rabbit

anti-Cx43 (Abcam, ab63851) antibody at 5 µg mL^{-1} in BSA 1% m v^{-1} in PBS for 1 h at room temperature,

then with anti-rabbit Alexa 568 (LifeTech, A11036) and Hoechst (Invitrogen, H3570) 1 µg mL^{-1} in BSA

1% m v-1 in PBS for 1 h. Samples were prepared with coverslips in Fluoromount mounting medium

(Sigma, HT5011).

2.4. Image acquisition and analysis

Samples were imaged with a Leica SPE Upright Confocal Microscope (Leica Microsystems) with a

40x/1.15 NA objective. The distance between imaged slices (z-size) was set at 1 µm. At least 3 cell

condensates were imaged for each sample.

Images were analyzed with ImageJ software. For condensate size measurements, a z-projection of each

condensate was created, and the whole condensate area was manually selected and measured. For the

measurement of Cx43 expression confocal z-projections were used (maximum stained area per sample).

Briefly, the background of z-projections was removed, and a threshold was applied to select areas of

Cx43 expression. The obtained total area was normalized against the area of the corresponding

condensate.

For the analysis of GJIC network, a threshold was applied to the Cx43 confocal stack and then it was

skeletonized with the ImageJ plugin (Fig. S1, Video S1). The resulting Cx43 network was analyzed to

retrieve the number of end-point voxels and the mean branch length in each condensate, which were

normalized to the Cx43 expression area and the number of slices taken for analysis. GJIC was calculated

as the inverse value of the end-point voxels.

2.5. RNA extraction and retrotranscription

Reverse transcription real-time PCR (RT-qPCR) was performed to measure Cx43 (*GJA1*) expression.

After 6 and 9 days of differentiation, mRNA was extracted from the samples and purified with a RNeasy

Micro Kit (Qiagen, 74004). Extracted mRNA was quantified in a Nanodrop ND-1000 Spectrophotometer

(Thermo Fisher Scientific). Reverse transcription for cDNA production was performed with an iScript

Advanced cDNA Synthesis Kit (Bio-Rad, 1725037) in a T100 Thermal Cycler (Bio-Rad). Three cell

culture replicates of each condition were obtained, with their RNA extracted and retrotranscribed. The

same procedure was performed on undifferentiated human mesenchymal stem cells (hMSCs) as a

reference.

2.6. qPCR and data analysis

qPCR was performed with the Sso Advanced Universal SYBR Green Supermix kit (Bio-Rad, 1725271)

in an Applied Biosystems StepOnePlus Real-Time PCR Machine (Thermo Fisher Scientific). Commercial

primer pairs were used for *GJA1* (Bio-Rad, qHsaCID0012977), as well as *B2M* (Bio-Rad,

qHsaCID0015347) and *RPL24* (Bio-Rad, qHsaCID0038677) as housekeeping genes. To prevent gDNA

amplification, a DNase digestion step was included during RNA extraction and intron-spanning primer

pairs were selected. The amplification program consisted of an initial activation step of 30 s at 95°C,

followed by 50 cycles of 10 s at 95°C for denaturation and 1 min at 60°C for annealing and extension, and

a final denaturation step of 15 s at 95°C. Melt curves were performed from 65°C to 95°C in steps of 0.5°C.

Technical duplicates of each sample were performed in the qPCR.

qPCR data were analyzed with qBase+ software version 3.1 (Biogazelle, Zwijnaarde, Belgium). Only

samples with a Ct below 40 were considered for analysis. The expression of each gene was calculated by

the $2^{-\Delta\Delta Ct}$ method, normalized to that of undifferentiated hMSCs (assigned value 1) and presented as

relative mRNA expression levels.

2.7. Neurobiotin (NB) assay

A tracer assay was performed to analyze the functionality of GJIC networks. Three replicates of each

condition were seeded in chondrogenesis-inducing medium, as described above. After 6 days of

differentiation, samples were washed with HBSS buffer without calcium or magnesium (Life

Technologies, 14175095) and treated with neurobiotin 2% m v^{-1} (Vector, SP-1120) in HBSS for 90 s at

37°C. Samples were then washed with HBSS, fixed with Formalin Solution, permeabilized with saponin

and stained with Streptavidin-Texas Red conjugate (Life Technologies, S872) and Hoechst 1 µg mL^{-1} in

BSA 1% m v^{-1} in PBS for 1 h at room temperature. Samples were imaged with a Leica SPE Upright

Confocal Microscope as described above.

Images were analyzed with ImageJ software. A z-projection of each condensate was created, and

background was removed. Distance of neurobiotin spread was measured in a straight line from the outer

rim of the condensates' inwards, in at least two separate locations for each condensate.

2.8. Condensate transplantation assay

Nanopatterns with 18% (S_{18}) and 90% (S_{90}) of local surface adhesiveness were used (Fig. S2 and Table

S1). A transplantation assay was performed to study the effects of RGD nanopatterned substrates on

formed condensates and the propagation of the adhesive information from the substrate into cell
condensates. Cells were cultured on the nanopatterns in chondrogenesis-inducing medium as described
above. After 3 days, cell condensates formed on the nanopatterns of S_{90} were removed by pipetting and
transferred to new S_{90} or S_{18} substrates. Transplanted condensates were cultured on the new substrates for
another 3 days, to a total of 6 days of differentiation. For each sample, around half of the condensates
were transplanted, whereas the other half were kept on the original substrate (not transplanted) as a
control of unaltered differentiation. Three replicates of each condition were seeded and transplanted.
Samples were fixed, immunostained, imaged and analyzed for Cx43 expression as described above.
Results were normalized to those of non-transplanted S_{90} condensates (assigned value 1) and presented as
relative values.
To visualize condensates from the side (transversal cuts), Z-stacks were resliced and one image from the
center of the condensate was selected. Cx43 production at the basal and top layers of condensates was
measured by the average staining intensity at each region on unprocessed images. For each condensate,
average basal intensity was divided by average top intensity to obtain the Basal/Top ratio, indicative of
Cx43 distribution within condensates.
218
219 *2.9. Integrin blocking and myosin inhibition assays*
220
Three replicates of each condition were seeded in chondrogenesis-inducing medium as described above.
For integrin blocking samples, medium was changed to fresh medium containing RGD dendrimers in
solution at $4\ 10^{-9}\%$ w w^{-1} after 5 days of differentiation. We selected this dendrimer concentration because
it yields S_{18} substrates. During substrate functionalization, equilibrium is reached between dendrimer
concentration in solution and adsorbed dendrimer density; hence, use of the concentration corresponding
to the substrates with the lowest density prevents further adsorption mid-assay**.**
For the myosin inhibition experiment, medium was changed to fresh medium with 50 µg mL^{-1}
blebbistatin (Sigma, B0560) 6 h before fixation.
All samples were fixed at day 6 of chondrogenesis, immunostained with anti-Cx43 antibody and Sir-
Actin (Tebu-bio, SC001), and imaged with a Zeiss LSM780 Confocal Microscope (Zeiss Microscopy)
with a 40x objective. Cx43 expression in condensates was quantified as described above and normalized
to corresponding non-treated samples.
233
234 *2.10.   Statistics*
235

Quantitative data are displayed, showing average and standard error of the mean (SEM). n is the sample

size. Significant differences were judged using the One-way ANOVA with Fisher LSD post-hoc test or T-

test when only two groups are compared, using OriginPro 8.5 or the Simple Interactive Statistical

Analysis (SISA) online tool [35]. Where data did not pass a normality test, a Kruskal-Wallis test with

Dunn means comparison was applied with GraphPad Prism 8.3. A p of less than 0.05 was considered

statistically significant.

3.   Results

3.1. *Gap junction intercellular communication: Network architecture and functionality*

We seeded hMSCs on the nanopatterns in chondrogenic medium. Due to the poor interfacing of PLLA

with living tissue [36], cells attached preferably through RGD-functionalized dendrimers. Pristine non-

patterned PLLA (S_0) and FN-coated (S_{FN}) substrates were the negative and positive controls for cell

adhesion, respectively. To control the local surface adhesiveness at the nanoscale during mesenchymal

condensation, we used nanopatterns with 18% (S_{18}), 45% (S_{45}) and 90% (S_{90}) of local surface

adhesiveness (Fig. S2 and Table S1). Under chondrogenic stimuli, mesenchymal cells aggregate into

three-dimensional condensates. While cells on pristine and nanopatterned substrates immediately start

aggregating, cells on fibronectin-coated substrates proliferate and adopt a monolayer configuration, from

which a few small condensates develop after 3 to 5 days (Fig. S3A) [30]. Condensates in S_{90} maintain

their structure in culture up to day 14 (Fig. S3B) [37], while condensates in the rest of the substrates

progressively collapse, with detached cells adopting a fibroblast-like morphology (Fig. S3C). We

measured the distance between adjacent cell nuclei and found that condensates on high-adherence

substrates (S_{90} and S_{FN}) were packed together more tightly than those on low- and mid- adherence ones

(Fig. S3D). In the case of large S_{90} condensates, this could be a factor explaining their resistance to

collapse through day 14.

Intercellular communication through gap junctions is crucial for correct cartilage development. We have

previously observed increased expression of chondrogenic markers on S_{90} substrates [30]; we thus

wondered if ligand density effects could improve cartilage formation beyond individual cell

differentiation. To examine cell-cell interconnectivity through GJ formation, we measured Cx43 in cell

condensates. Cx43 is the most widely expressed and studied GJ protein [38] and modulates cartilage

structure through its C-terminal cytosolic domain [39]. Connexin hemichannels, or connexons,

accumulate and dock with apposed connexons from adjacent cells to form dense GJ plaques. GJIC largely

depends on the size of GJ plaques [40] which are continuously regenerated by the addition of connexon

subunits at the edges and internalization from the center of the plaques [7,41,42]. Such a continuous

renewal may ensure the maintenance of the established GJIC network.

We analysed the expression of *GJA1* at day 6 and 9 of chondrogenesis (Fig. 1A). *GJA1* was

overexpressed on S_{90} to over twice the level of undifferentiated hMSCs at day 6 and reduced by half on

S_{FN}. However, expression on S_{90} levelled off at day 9 among the different nanopattern configurations and

S_0. *GJA1* expression in S_{18} and S_{45} nanopatterns is comparable to that of the negative control both at day 6

and 9 of chondrogenic induction.

Cx43 immunofluorescent images show Cx43 assembled in GJ plaques. Quantification of the percentage

of immunostained area showed higher values on S_{90} when compared to most other substrates at day 6 and

a slight decrease at day 9, mirroring mRNA expression (Fig. 1B). The fact that Cx43 in GJ plaques on S_{90}

is not as high as mRNA expression at day 6 could be explained by a rapid initial turnover [12,41].

Confocal z-projections (Fig. 1C, top) of immunostaining show that Cx43 GJ plaques were roughly

distributed within the cell condensates with a tendency to accumulate in the outer cell layers, particularly

in S_{45}.

Skeletonization of Cx43 immunostained images renders a 3D representation of the intercellular

communication network (Fig. 1C, bottom, Fig. S1 and Video S1), from which the average branch length

(Fig. S4) and number of branch terminations (end-point voxels) can be calculated. Shorter branches and

fewer end-point voxels indicate a more intricately shaped GJIC network, as shown in the zoomed-in

sections of the skeletonized images of condensates from S_{90} and S_0 nanopatterns. We thus took the

number of branch terminations as the inverse index for Cx43 architectural connectivity (Fig. 1D). At days

6 and 9 of chondrogenic induction, connectivity progressively increased with local surface adhesiveness

up to S_{90} and decreased for S_{FN}.

Observation of the spread of biotinylated or fluorescent tracers has become one of the most common

methods of demonstrating gap junctions network coupling [43]. Therefore, to further assess GJIC, we

conducted a neurobiotin (NB) tracer uptake assay in cell condensates (Fig. 1E). NB is a GJ/hemichannel

permeable dye that can penetrate from the exposed Cx43 connexons and diffuse inwards across the GJIC

network when Ca^{2+} is maintained below physiological levels (open channel conformation). Quantification

of NB diffusion into cell condensates showed NB uptake is significantly higher in S_{90} nanopatterns and

equal for all other substrates (Fig. 1F), thereby indicating that S_{90} condensates developed a more efficient

GJIC network.

Fig. 1: Ligand density modulates gap junction network architecture and function. (**A**) Expression of connexin 43 (*GJA1)* mRNA, relative to that of undifferentiated hMSCs (assigned value 1, not shown) (n=6). (**B**) Percentage of Cx43-stained areas in cell condensates (n≥3). (**C**) Representative confocal z-projections of Cx43 immunostaining (top) and the corresponding skeletonized images (bottom) revealing the branched architecture of the intercellular connectivity network. Scale bar = 40 µm. Zoomed-in sections for the skeletonized images of S_0 and S_{90} highlighting the differences in the GJIC network. Scale bar = 3 µm. (**D**) Quantification of Cx43 architectural network connectivity from skeletonized images normalized against Cx43 in GJs (n≥3). (**E**) Representative confocal z-projections showing neurobiotin (NB) tracer (red) in the mesenchymal condensates and cell nuclei stained with Hoechst (blue). White lines represent measured distances of neurobiotin uptake. Scale bar = 80 µm. (**F**) Quantification of neurobiotin tracer uptake in day 6 condensates after 90 s exposure (n≥24). Gray bars with green dots correspond to day 6 of chondrogenic induction; white bars with orange triangles correspond to day 9. Results are given as individual sample values with the mean ± SEM, *$p<0.05$, **$p<0.01$, ****$p<0.0001$

3.2. Matrix adhesion continually regulates GJIC

Results showed an effect of local surface adhesiveness on the architecture and functionality of GJIC

networks in pre-cartilaginous cell condensates. However, it was unclear whether substrate adhesion is

relevant only at the beginning of condensation as a memory effect or if it continuously influences the

tissue as it forms. To consider these two possibilities, we designed a transplantation assay in which

condensates formed on S_{90} and $S_{18,}$ (the nanopatterns of highest and lowest ligand density) were collected

at day 3, plated on new S_{90} and S_{18} substrates and maintained for 3 more days in culture (Fig. 2A). The

percentage of area immunostained for Cx43 (GJ plaques) was assessed at day 6 for the transplanted and

non-transplanted condensates. Transplantation produced a significant boosting effect for condensates

plated on new S_{90} substrates (Fig. 2B). In condensates transplanted from S_{90} to fresh S_{90}, Cx43 production

increased by 71±21%, while a 61±19% increase was observed for those transplanted from S_{18}.

Transplantation of either S_{90} or S_{18} condensates to fresh S_{18} substrates did not render significant changes

in Cx43 production.

Condensates formed on one substrate and then transplanted to another will sense the new input only from

the side in direct contact with the new substrate. We questioned if effects derived from changing substrate

conditions would be confined to cells in direct contact with it (those at the basal layer of condensates) or

instead propagate to the middle and top layers of condensates. Transversal views of transplanted

condensates presented increased Cx43 production at all heights (Fig. 2C), showing that fresh substrate

inputs spread through cells and modulate protein expression within the whole forming tissue. Moreover,

the proportion of Cx43 production between the basal versus top regions was equal in control and

transplanted condensates, indicating that transplantation did not alter the ratios of protein distribution

among layers (Fig. 2D).

Fig. 2: Cell condensates adapt to changing ligand density. (A) Schematics of the transplantation experiment. Condensates formed on S_{90} and S_{18} substrates were collected at day 3 and plated on fresh S_{90} or S_{18} substrates for 3 more days of chondrogenic induction. **(B)** Quantification of Cx43-stained percentage area for non-transplanted and transplanted condensates (n≥7). **(C)** Confocal side views of control and transplanted S_{90} condensates, stained for Cx43 (red) and Hoechst (blue). Substrate effect propagates to cells at all levels within condensates. Scale bar = 10 μm. **(D)** Ratio between Cx43 production at the basal and top regions of S_{90} condensates in control S_{90} substrates or transplanted to fresh S_{90} (n≥9). Results are given as mean ± SEM, **$p<0.01$

3.3. Mechanism of substrate sensing and input propagation

Integrin adhesions are the main mediators of cell-matrix adhesion and mechanical interactions [44,45]. To

evaluate their role in modulating GJIC on nanopatterns, we added RGD-functionalized dendrimers in

solution on formed condensates to block integrin receptors at the cell membrane. Disturbance of integrin

clustering at cell-substrate adhesion sites led to a Cx43 production decrease in S_{90} but not in S_{18} (Fig. 3A).

This indicates that integrin-mediated cell adhesion is responsible for the transduction and regulation of

substrate information that affects GJIC, and that cell response is triggered when integrins are engaged by

the RGD configuration provided by S_{90} nanopatterns.

Since perturbations at the cell membrane are transduced into chemical responses by propagation from

integrins through the cytoskeleton [16], we hypothesized that the substrate adhesion information

orchestrating GJIC during mesenchymal cell condensation propagates through the actin cortex. Cortical

actin assembly can be observed in early mesenchymal condensates [30]. Cortical tension gradients are

responsible for mesenchymal cell rearrangement during tissue formation and are essential in driving

tissue morphogenesis [46,47]. The cell cortex is formed by an actomyosin network, which is located

below the cell membrane. In the cortex, myosin-2 pulls on actin filaments and generates tension.

Inhibition of myosin-2 activity can cause a decrease in cortical tension of up to 80% [47]. Treatment of 6-

day condensates with the myosin-2 inhibitor blebbistatin decreased the percentage of area immunostained

for Cx43 of S_{90} condensates (Fig. 3A).

Adding dendrimers in solution affected cytoskeletal conformation in a similar manner as blebbistatin,

preventing polymerization and resulting in a diffuse distribution of actin in condensates, instead of

concentrating in clearly defined fibers as in control conditions (Fig. 3B). This shows that cytoskeletal

tension depends on integrin receptors engaging adherence sites at the substrates; we thus infer that cells at

the basal layer of condensates sense ligand inputs through integrins and propagate them by actin

contractility to adjacent cells (Fig. 3C).

Fig. 3: Signal transduction and propagation. **(A)** Relative Cx43 production in S_{90} (gray bars, red dots)
and S_{18} (white bars, blue triangles) (n≥12). Results are given as mean ± SEM, **p<0.01. **(B)** Actin
staining for S_{90} condensates at day 6 in control conditions, with RGD dendrimers in solution (integrin
block) or with blebbistatin (myosin inhibition). Scale bar = 25 µm. **(C)** Schematic representation of ligand
density sensing through integrin adhesions, followed by actomyosin contraction-guided propagation of
information in cell condensates.

4. Discussion

We studied the effects of ECM adhesion in the establishment of GJIC during mesenchymal cell
condensation, a prevalent morphogenetic transition. To this end, we used previously developed RGD
dendrimer-based nanopatterns of tunable local surface adhesiveness [29,30]. Larger and more stable cell
condensates are obtained on S_{90} nanopatterns, with cells packed closer together.

We analyzed the influence of substrate ligand density on the establishment of a gap junction intercellular
communication (GJIC) network in cell condensates. We measured gene expression and assembly at gap
junctions of Cx43, a connexin which is ubiquitously expressed in developing cartilage [39,48]. We found
Cx43 increased with local surface adhesiveness in the nanopatterns, a similar interdependence to what we
observed previously for the focal adhesion adaptor protein paxillin [30]. This is in agreement with Zhou *et
al*., who observed a direct interaction between FAK and paxillin with Cx43 in cells from the human apical
papilla, which depended on substrate compliance [13]. In our case, cell condensates on S_{90} nanopatterns

presented increased Cx43 expression with a higher protein turnover rate, revealing a direct effect of

substrate ligand density on gap junction protein regulation.

We analyzed the architectural connectivity of Cx43 in condensates to assess whether substrate ligand

density modulates the development of gap junction networks beyond protein expression. We found that

S_{90} condensates contain a Cx43 architecture that is more intricate, with less end-point voxels and shorter

branches, indicative of improved communication capacities. Moreover, a tracer uptake assay revealed that

GJIC is more efficient on S_{90} condensates. This shows that local surface adhesiveness given by the

nanopattern configuration determines not only protein expression but also its spatial disposition during

morphogenesis, affecting the functionality of the forming tissue. Given that S_{90} substrates also promote

cell differentiation towards chondrocytes [30], we conclude that nanopatterning at this particular ligand

density improves cartilage formation *in vitro*.

A condensate transplantation assay was designed to test whether cells in the condensate, previously in

contact with the substrate during cell recruitment, retain initial substrate information (memory effect) or

receive a continuous feedback of substrate input. Transplantation of cell condensates to a new S_{90}

substrate caused a significant increase in Cx43, suggesting that cell condensates still preserve a certain

level of mesenchymal plasticity at early stages of chondrogenic differentiation [49], thereby allowing

phenotype reconfiguration in response to the ECM input. No effects were observed for transplantation to

S_{18} substrates.

Effects of substrate transplantation on Cx43 expression were observed at all heights within condensates

and transplanted condensates maintained the proportion of total protein contained at the basal and top

regions, indicating that substrate inputs are propagated into the third dimension, across cells to the top.

These results show that changing environmental conditions during morphogenesis continually modulate

the characteristics of forming tissue, pointing to the importance of accounting for time variations in the

design of experimental setups for development studies, as well as biomaterials for tissue engineering.

Finally, we addressed the biological mechanism by which cells sense substrate ligand density and

propagate it into condensates. Blocking integrin-mediated interactions between the substrate and the basal

layer in cell condensates caused a Cx43 decrease in S_{90} but not in S_{18}. This observation, together with

results of condensate stability and GJIC, demonstrates that the establishment of GJIC during

mesenchymal condensation is an adhesion-gated mechanism, in which S_{90} nanopattern configuration

provides optimal ligand density and distribution to trigger cell response.

To investigate the propagation of substrate information within cell condensates, an inhibition experiment

for myosin-2 was conducted. Inhibition of myosin-2 activity by blebbistatin caused a decrease of Cx43 in

S_{90} cell condensates, in agreement with previous reports [50]. This indicates that once integrins are

engaged, the adhesion information from the substrate is transduced and propagates through the actin

filaments by myosin-2 mediated contraction, which in turn can regulate GJ accretion through ZO-1[12].
Results agree with those from Gowrishankar *et al*., who predicted that the dynamics of membrane
proteins are regulated by the cell cortex [51]. Therefore, we propose a contractility-based mechanism for
the propagation of ECM adhesion information within the condensates in which local substrate
adhesiveness induces a rearrangement and adaptation of the actin cortex and cell-cell junctions in the first
cell layers [52,53] and propagates through the tissue by myosin-2-mediated cortex contractions, regulating
Cx43 dynamics during the mesenchymal condensation process (Fig. 3C).

Our findings clearly reflect the impact that the fine tuning of materials interfaces has in cell response. By
tailoring the nanopattern configuration of the cell-adhesive motif RGD, we exerted control over the
architecture and function of a complex dynamic biological system. The results have an immediate
application to cartilage *in vitro* engineering [30,54], but they are also extensible to the study of other
biological processes in which active ECM remodelling and thus, changes in the adhesion requirements
and intercellular communication play an active role, such as in cancer progression [55].

CRediT author statement

I. Casanellas: Conceptualization, Methodology, Investigation, Formal analysis, Writing – Original Draft,
Visualization. **A. Lagunas:** Conceptualization, Methodology, Formal analysis, Writing – Original Draft,
Supervision, Project administration, Funding acquisition. **Y. Vida:** Methodology, Resources (design and
production of RGD-functionalized dendrimers), Writing – Review & Editing. **E. Pérez-Inestrosa**:
Methodology, Resources (design and production of RGD-functionalized dendrimers), Writing – Review
& Editing, Funding acquisition. **C. Rodríguez-Pereira:** Methodology, Investigation, Formal analysis
(RT-qPCR), Writing – Review & Editing. **J. Magalhaes:** Methodology (RT-qPCR), Supervision, Writing
– Review & Editing. **J. A. Andrades:** Writing – Review & Editing, Funding acquisition. **J. Becerra:**
Writing – Review & Editing, Funding acquisition. **J. Samitier:** Conceptualization, Methodology, Writing
– Review & Editing, Supervision, Project administration, Funding acquisition.

Funding

This work was supported by the Biomedical Research Networking Center (CIBER), Spain. CIBER is an
initiative funded by the VI National R&D&i Plan 2008–2011, Iniciativa Ingenio 2010, Consolider
Program, CIBER Actions, and the Instituto de Salud Carlos III (RD16/0006/0012; RD16/0011/0022),
with the support of the European Regional Development Fund (ERDF). This work was funded by the
CERCA Program and by the Commission for Universities and Research of the Department of Innovation,
Universities, and Enterprise of the Generalitat de Catalunya (2017 SGR 1079). This work has been
developed in the context of the AdvanceCat project (COMRDI15-1-0013) with the support of ACCIÓ

(Catalonia Trade and Investment; Generalitat de Catalunya) under the Catalonian ERDF operational

program 2014–2020. This work was funded by the Spanish Ministry of Economy and Competitiveness

(MINECO) through the projects MINDS (Proyectos I+D Excelencia + FEDER): TEC2015-70104-P-P

and BIOBOT (Programa Explora Ciencia / Tecnología): TEC2015- 72718-EXP, as well as the Spanish

Ministry of Science and Education (PID2019-104293GB-I00), and the Consejería de Salud, Junta de

Andalucía (UMA18-FEDERJA-007; UMA18□FEDERJA□133). The authors further acknowledge funds

from the INTERREG V cooperation program for Spain–Portugal (POCTEP) 2014–2020 project

(0245_IBEROS_1_E). I. C. acknowledges support from MINECO through the Subsidies for Predoctoral

Contracts for the Training of Doctors open call, co-funded by the European Social Fund (2016), grant

number: BES-2016-076682. C. R-P. acknowledges support from i-PFIS Doctoral Program (IFI15/00151)

funded by ISCIII.

469 Conflicts of interest

470 There are no conflicts to declare.

472 Acknowledgements

473 We thank N. Montserrat for help with the transplantation assay, R. Paoli for producing Video S1, and P.

474 Roca-Cusachs for the fruitful discussions.

476 References

[1] A.M. DeLise, L. Fischer, R.S. Tuan, Cellular interactions and signaling in cartilage development,

Osteoarthr. Cartil. 8 (2000) 309–334. https://doi.org/10.1053/joca.1999.0306.

[2] H. Grüneberg, The pathology of development: a study of inherited skeletal disorders in animals,

Blackwells Scientific Publications, Oxford, 1963.

[3] S. Christley, M.S. Alber, S.A. Newman, Patterns of mesenchymal condensation in a multiscale,

discrete stochastic model, PLoS Comput. Biol. 3 (2007) 743–753.

https://doi.org/10.1371/journal.pcbi.0030076.

[4] P. Singh, J.E. Schwarzbauer, Fibronectin and stem cell differentiation - lessons from

chondrogenesis, J. Cell Sci. 125 (2012) 3703–3712. https://doi.org/10.1242/jcs.095786.

[5] C.N.D. Coelho, R.A. Kosher, Gap junctional communication during limb cartilage differentiation,

Dev. Biol. 144 (1991) 47–53. https://doi.org/https://doi.org/10.1016/0012-1606(91)90477-K.

[6] J.C. Hervé, M. Derangeon, Gap-junction-mediated cell-to-cell communication, Cell Tissue Res.

352 (2013) 21–31. https://doi.org/10.1007/s00441-012-1485-6.

[7]   J.C. Hervé, M. Derangeon, D. Sarrouilhe, B.N.G. Giepmans, N. Bourmeyster, Gap junctional
492   channels are parts of multiprotein complexes, Biochim. Biophys. Acta - Biomembr. 1818 (2012)
493   1844–1865. https://doi.org/10.1016/j.bbamem.2011.12.009.

494   [8]   M.D. Mayan, R. Gago-Fuentes, P. Carpintero-Fernandez, P. Fernandez-Puente, P. Filgueira-
495   Fernandez, N. Goyanes, V. Valiunas, P.R. Brink, G.S. Goldberg, F.J. Blanco, Articular
496   chondrocyte network mediated by gap junctions: Role in metabolic cartilage homeostasis, Ann.
497   Rheum. Dis. 74 (2015) 275–284. https://doi.org/10.1136/annrheumdis-2013-204244.

498   [9]   H.J. Donahue, R.W. Qu, D.C. Genetos, Joint diseases: From connexins to gap junctions, Nat. Rev.
499   Rheumatol. 14 (2018) 42–51. https://doi.org/10.1038/nrrheum.2017.204.

500   [10]   P.D. Lampe, B.P. Nguyen, S. Gil, M. Usui, J. Olerud, W.G. Carter, Y. Takada, Cellular interaction
501   of integrin α3β1 with laminin 5 promotes gap junctional communication, J. Cell Biol. 143 (1998)
502   1735–1747. https://doi.org/10.1083/jcb.143.6.1735.

503   [11]   N. Batra, S. Burra, A.J. Siller-Jackson, S. Gu, X. Xia, G.F. Weber, D. DeSimone, L.F. Bonewald,
504   E.M. Lafer, E. Sprague, M.A. Schwartz, J.X. Jiang, Mechanical stress-activated integrin α5β1
505   induces opening of connexin 43 hemichannels, Proc. Natl. Acad. Sci. 109 (2012) 3359–3364.
506   https://doi.org/10.1073/pnas.1115967109.

507   [12]   A.E. Zemljic-Harpf, J.C. Godoy, O. Platoshyn, E.K. Asfaw, A.R. Busija, A.A. Domenighetti, R.S.
508   Ross, Vinculin directly binds zonula occludens-1 and is essential for stabilizing connexin-43-
509   containing gap junctions in cardiac myocytes, J. Cell Sci. 127 (2014) 1104–1116.
510   https://doi.org/10.1242/jcs.143743.

511   [13]   C. Zhou, D. Zhang, W. Du, J. Zou, X. Li, J. Xie, Substrate mechanics dictate cell-cell
512   communication by gap junctions in stem cells from human apical papilla, Acta Biomater. 107
513   (2020) 178–193. https://doi.org/10.1016/j.actbio.2020.02.032.

514   [14]   B.M. Gumbiner, Cell adhesion: The molecular basis of tissue architecture and morphogenesis,
515   Cell. 84 (1996) 345–357. https://doi.org/10.1016/S0092-8674(00)81279-9.

516   [15]   J.T. Parsons, A.R. Horwitz, M.A. Schwartz, Cell adhesion: Integrating cytoskeletal dynamics and
517   cellular tension, Nat. Rev. Mol. Cell Biol. 11 (2010) 633–643. https://doi.org/10.1038/nrm2957.

518   [16]   M. Bachmann, S. Kukkurainen, V.P. Hytönen, B. Wehrle-Haller, Cell adhesion by integrins,
519   Physiol. Rev. 99 (2019) 1655–1699. https://doi.org/10.1152/physrev.00036.2018.

520   [17]   D.C. Van Essen, A tension-based theory of morphogenesis and compact wiring in the central
521   nervous system, Nature. 385 (1997) 313–318. https://doi.org/10.1038/385313a0.

522   [18]   M. Tada, M.L. Concha, Vertebrate gastrulation: Calcium waves orchestrate cell movements, Curr.
523   Biol. 11 (2001) 470–472. https://doi.org/10.1016/S0960-9822(01)00284-6.

[19]   Hung Ping Shih, D. Panlasigui, V. Cirulli, M. Sander, ECM signaling regulates collective cellular

dynamics to control pancreas branching morphogenesis, Cell Rep. 14 (2016) 169–179.

https://doi.org/10.1016/j.celrep.2015.12.027.

[20]   F. Jiang, H. Hörber, J. Howard, D.J. Müller, Assembly of collagen into microribbons: Effects of

pH and electrolytes, J. Struct. Biol. 148 (2004) 268–278. https://doi.org/10.1016/j.jsb.2004.07.001.

[21]   M.L. Smith, D. Gourdon, W.C. Little, K.E. Kubow, R.A. Eguiluz, S. Luna-Morris, V. Vogel,

Force-induced unfolding of fibronectin in the extracellular matrix of living cells, PLoS Biol. 5

(2007) 2243–2254. https://doi.org/10.1371/journal.pbio.0050268.

[22]   W.C. Little, M.L. Smith, U. Ebneter, V. Vogel, Assay to mechanically tune and optically probe

fibrillar fibronectin conformations from fully relaxed to breakage, Matrix Biol. 27 (2008) 451–

461. https://doi.org/10.1016/j.matbio.2008.02.003.

[23]   M.F. Garcia-Parajo, A. Cambi, J.A. Torreno-Pina, N. Thompson, K. Jacobson, Nanoclustering as a

dominant feature of plasma membrane organization, J. Cell Sci. 127 (2014) 4995–5005.

https://doi.org/10.1242/jcs.146340.

[24]   M. Arnold, E.A. Cavalcanti-Adam, R. Glass, J. Blümmel, W. Eck, M. Kantlehner, H. Kessler, J.P.

Spatz, Activation of integrin function by nanopatterned adhesive interfaces, ChemPhysChem. 5

(2004) 383–388. https://doi.org/10.1002/cphc.200301014.

[25]   E.A. Cavalcanti-Adam, T. Volberg, A. Micoulet, H. Kessler, B. Geiger, J.P. Spatz, Cell spreading

and focal adhesion dynamics are regulated by spacing of integrin ligands, Biophys. J. 92 (2007)

2964–2974. https://doi.org/10.1529/biophysj.106.089730.

[26]   J.A. Deeg, I. Louban, D. Aydin, C. Selhuber-Unkel, H. Kessler, J.P. Spatz, Impact of local versus

global ligand density on cellular adhesion, Nano Lett. 11 (2011) 1469–1476.

https://doi.org/10.1021/nl104079r.

[27]   R. Oria, T. Wiegand, J. Escribano, A. Elosegui-Artola, J.J. Uriarte, C. Moreno-Pulido, I. Platzman,

P. Delcanale, L. Albertazzi, D. Navajas, X. Trepat, J.M. García-Aznar, E.A. Cavalcanti-Adam, P.

Roca-Cusachs, Force loading explains spatial sensing of ligands by cells, Nature. 552 (2017) 219–

224. https://doi.org/10.1038/nature24662.

[28]   A. Lagunas, D. Caballero, J. Samitier, Influence of Controlled Micro☐ and Nanoengineered

Environments on Stem Cell Fate, in: A. Tiwari, B. Garipcan, L. Uzun (Eds.), Adv. Surfaces Stem

Cell Res., 1st ed., Hoboken: Wiley, 2016: pp. 87–141.

[29]   A. Lagunas, A.G. Castaño, J.M. Artés, Y. Vida, D. Collado, E. Pérez-Inestrosa, P. Gorostiza, S.

Claros, J.A. Andrades, J. Samitier, Large-scale dendrimer-based uneven nanopatterns for the study

of local arginine-glycine-aspartic acid (RGD) density effects on cell adhesion, Nano Res. 7 (2014)

399–409. https://doi.org/10.1007/s12274-014-0406-2.

[30]   A. Lagunas, I. Tsintzou, Y. Vida, D. Collado, E. Pérez-Inestrosa, C. Rodríguez Pereira, J.

Magalhaes, J.A. Andrades, J. Samitier, Tailoring RGD local surface density at the nanoscale

toward adult stem cell chondrogenic commitment, Nano Res. 10 (2017) 1959–1971.

https://doi.org/10.1007/s12274-016-1382-5.

[31]   I. Casanellas, A. Lagunas, I. Tsintzou, Y. Vida, D. Collado, E. Pérez-Inestrosa, C. Rodríguez-

Pereira, J. Magalhaes, P. Gorostiza, J.A. Andrades, J. Becerra, J. Samitier, Dendrimer-based

uneven nanopatterns to locally control surface adhesiveness: A method to direct chondrogenic

differentiation, J. Vis. Exp. 131 (2018). https://doi.org/10.3791/56347.

[32]   I. Casanellas, A. Lagunas, Y. Vida, E. Pérez-Inestrosa, J.A. Andrades, J. Becerra, J. Samitier,

Matrix nanopatterning regulates mesenchymal differentiation through focal adhesion size and

distribution according to cell fate, Biomimetics. 4 (2019).

https://doi.org/10.3390/biomimetics4020043.

[33]   B.E. Bobick, F.H. Chen, A.M. Le, R.S. Tuan, Regulation of the chondrogenic phenotype in

culture, Birth Defects Res. Part C - Embryo Today Rev. 87 (2009) 351–371.

https://doi.org/10.1002/bdrc.20167.

[34]   I. Horcas, R. Fernández, J.M. Gómez-Rodríguez, J. Colchero, J. Gómez-Herrero, A.M. Baro,

WSXM: A software for scanning probe microscopy and a tool for nanotechnology, Rev. Sci.

Instrum. 78 (2007). https://doi.org/10.1063/1.2432410.

[35]   D.G. Uitenbroek, SISA Binomial, (1997). http://www.quantitativeskills.com/sisa/statistics/t-

test.htm (accessed January 30, 2020).

[36]   Y. Zhu, C. Gao, X. Liu, T. He, J. Shen, Immobilization of Biomacromolecules onto Aminolyzed

Poly(L-lactic acid) toward Acceleration of Endothelium Regeneration, Tissue Eng. 10 (2004) 53–

61. https://doi.org/10.1089/107632704322791691.

[37]   I. Casanellas, A. Lagunas, Y. Vida, P. Ezequiel, J.A. Andrades, J. Becerra, J. Samitier, The Janus

role of adhesion in chondrogenesis, Int. J. Mol. Sci. 21 (2020) 5269.

https://doi.org/10.3390/ijms21155269.

[38]   T.M. Ribeiro-Rodrigues, T. Martins-Marques, S. Morel, B.R. Kwak, H. Girão, Role of connexin

43 in different forms of intercellular communication - gap junctions, extracellular vesicles and

tunnelling nanotubes, J. Cell Sci. 130 (2017) 3619–3630. https://doi.org/10.1242/jcs.200667.

[39]   R. Gago-Fuentes, J.F. Bechberger, M. Varela-Eirin, A. Varela-Vazquez, B. Acea, E. Fonseca, C.C.

Naus, M.D. Mayan, The C-terminal domain of connexin43 modulates cartilage structure via

chondrocyte phenotypic changes, Oncotarget. 7 (2016) 73055–73067.

https://doi.org/10.18632/oncotarget.12197.

[40]   J.M. Rhett, J. Jourdan, R.G. Gourdie, Connexin 43 connexon to gap junction transition is regulated

by zonula occludens-1, Mol. Biol. Cell. 22 (2011) 1516–1528. https://doi.org/10.1091/mbc.E10-

06-0548.

[41] E. Leithe, S. Sirnes, T. Fykerud, A. Kjenseth, E. Rivedal, Endocytosis and post-endocytic sorting of connexins, Biochim. Biophys. Acta - Biomembr. 1818 (2012) 1870–1879. https://doi.org/10.1016/j.bbamem.2011.09.029.

[42] M. Piehl, C. Lehmann, A. Gumpert, J.-P. Denizot, D. Segretain, M.M. Falk, Internalization of Large Double-Membrane Intercellular Vesicles by a Clathrin-dependent Endocytic Process, Mol. Biol. Cell. 18 (2007) 337–347. https://doi.org/10.1091/mbc.E06.

[43] J. Jacoby, A. Nath, Z.F. Jessen, G.W. Schwartz, A Self-Regulating Gap Junction Network of Amacrine Cells Controls Nitric Oxide Release in the Retina, Neuron. 100 (2018) 1149-1162.e5. https://doi.org/10.1016/j.neuron.2018.09.047.

[44] Z. Sun, S.S. Guo, R. Fässler, Integrin-mediated mechanotransduction, J. Cell Biol. 215 (2016) 445–456.

[45] J.Z. Kechagia, J. Ivaska, P. Roca-Cusachs, Integrins as biomechanical sensors of the microenvironment, Nat. Rev. Mol. Cell Biol. (2019). https://doi.org/10.1038/s41580-019-0134-2.

[46] M. Nishikawa, S.R. Naganathan, F. Jülicher, S.W. Grill, Controlling contractile instabilities in the actomyosin cortex, Elife. 6 (2017) 1–21. https://doi.org/10.7554/eLife.19595.

[47] H. Tao, M. Zhu, K. Lau, O.K.W. Whitley, M. Samani, X. Xiao, X.X. Chen, N.A. Hahn, W. Liu, M. Valencia, M. Wu, X. Wang, K.D. Fenelon, C.C. Pasiliao, D. Hu, J. Wu, S. Spring, J. Ferguson, E.P. Karuna, R.M. Henkelman, A. Dunn, H. Huang, H.Y.H. Ho, R. Atit, S. Goyal, Y. Sun, S. Hopyan, Oscillatory cortical forces promote three dimensional cell intercalations that shape the murine mandibular arch, Nat. Commun. 10 (2019) 1–18. https://doi.org/10.1038/s41467-019-09540-z.

[48] R.A. Meyer, M.F. Cohen, S. Recalde, J. Zakany, S.M. Bell, W.J.J. Scott, C.W. Lo, Developmental regulation and asymmetric expression of the gene encoding Cx43 gap junctions in the mouse limb bud., Dev. Genet. 21 (1997) 290–300. https://doi.org/10.1002/(SICI)1520-6408(1997)21:4<290::AID-DVG6>3.0.CO;2-2.

[49] Y. Liu, N. Muñoz, A.C. Tsai, T.M. Logan, T. Ma, Metabolic Reconfiguration Supports Reacquisition of Primitive Phenotype in Human Mesenchymal Stem Cell Aggregates, Stem Cells. 35 (2017) 398–410. https://doi.org/10.1002/stem.2510.

[50] S. Olk, G. Zoidl, R. Dermietzel, Connexins, cell motility, and the cytoskeleton, Cell Motil. Cytoskeleton. 66 (2009) 1000–1016. https://doi.org/10.1002/cm.20404.

[51] K. Gowrishankar, S. Ghosh, S. Saha, C. Rumamol, S. Mayor, M. Rao, Active remodeling of cortical actin regulates spatiotemporal organization of cell surface molecules, Cell. 149 (2012) 1353–1367. https://doi.org/10.1016/j.cell.2012.05.008.

627   [52]   N. Wang, J.D. Tytell, D.E. Ingber, Mechanotransduction at a distance: Mechanically coupling the

628   extracellular matrix with the nucleus, Nat. Rev. Mol. Cell Biol. 10 (2009) 75–82.

629   https://doi.org/10.1038/nrm2594.

630   [53]   P. Chugh, E.K. Paluch, The actin cortex at a glance, J. Cell Sci. 131 (2018) 1–9.

631   https://doi.org/10.1242/jcs.186254.

632   [54]   C. Rodríguez-Pereira, A. Lagunas, I. Casanellas, Y. Vida, E. Pérez-Inestrosa, J.A. Andrades, J.

633   Becerra, J. Samitier, F.J. Blanco, J. Magalhães, RGD-dendrimer-poly(L-lactic) acid nanopatterned

634   substrates for the early chondrogenesis of human mesenchymal stromal cells derived from

635   osteoarthritic and healthy donors, Materials (Basel). 13 (2020).

636   https://doi.org/10.3390/ma13102247.

637   [55]   C. Bonnans, J. Chou, Z. Werb, Remodelling the extracellular matrix in development and disease,

638   Nat. Rev. Mol. Cell Biol. 15 (2014) 786–801. https://doi.org/10.1038/nrm3904.

A *GJA1*

B Cx43 in gap junctions

C

D Cx43 network connectivity

E

F GJIC

A

Cx43 relative production

B

Control  RGD in solution  Blebbistatin

C

Contractility-mediated information propagation → GJIC regulation

Integrin transduction

▮▮ Integrin
⬤ GJ plaque
— Contractile cortex
⬤ RGD dendrimer