Quantitative multidimensional phenotypes improve genetic analysis of laterality traits

Running title: Improving genetic analysis of laterality traits

Dr Judith Schmitz[1],

Dr Mo Zheng[2],

Dr Kelvin F. H. Lui[3],

Professor Catherine McBride[2],

Professor Connie S.-H. Ho[4],

Dr Silvia Paracchini[1]

[1]School of Medicine, University of St Andrews

[2]Department of Psychology, The Chinese University of Hong Kong

[3]Wofoo Joseph Lee Consulting and Counselling Psychology Research Centre, Lingnan University

[4]Psychology Department, The University of Hong Kong

Corresponding author:

Silvia Paracchini (sp58@st-andrews.ac.uk)

School of Medicine, Medical and Biological Sciences Building, North Haugh, St Andrews, KY16 9TF

Abstract

Handedness is the most commonly investigated lateralised phenotype and is usually

measured as a binary left/right category. Its links with psychiatric and neurodevelopmental

disorders prompted studies aimed at understanding the underlying genetics, while other

measures and side preferences have been less studied. We investigated the heritability of

hand, as well as foot, and eye preference by assessing parental effects ($n \leq 5\ 028$ family trios)

and SNP-based heritability (SNP-h^2, $n \leq 5\ 931$ children) in the Avon Longitudinal Study of

Parents and Children (ALSPAC). An independent twin cohort from Hong Kong ($n = 358$) was

used to replicate results from structural equation modelling (SEM). Parental left-side

preference increased the chance of an individual to be left-sided for the same trait, with

stronger maternal than paternal effects for footedness. By regressing out the effects of sex,

age, and ancestry, we transformed laterality categories into quantitative measures. The SNP-

h^2 for quantitative handedness and footedness was .21 and .23, respectively, which is higher

than the SNP-h^2 reported in larger genetic studies using binary handedness measures. The

heritability of the quantitative measure of handedness increased (.45) compared to a binary

measure for writing hand (.27) in the Hong Kong twins. Genomic and behavioural SEM

identified a shared genetic factor contributing to handedness, footedness, and eyedness, but

no independent effects on individual phenotypes. Our analysis demonstrates how

quantitative multidimensional laterality phenotypes are better suited to capture the

underlying genetics than binary traits.

1   Introduction

The cerebral hemispheres differ in function and structure underpinning specialisation for

cognition, perception, and motor control [1]. For instance, language is predominantly processed

in the left hemisphere in most individuals [2] and the *planum temporale* typically shows a

pronounced structural leftward asymmetry [3], although there is little evidence for a strong

association between the two forms of asymmetry [4]. Neurodevelopmental disorders such as

dyslexia [5,6], schizophrenia [7], or autism spectrum disorder (ASD) [8] have been associated with a

higher prevalence of atypical *planum temporale* asymmetry.

The most commonly studied lateralised trait is handedness. Worldwide, around 10% of the

general population is left-handed with slight geographical variation [9], likely influenced by

cultural factors [10,11]. Meta-analyses have confirmed higher rates of left- or non-right-

handedness in ASD [12] and schizophrenia [13]. A genetic influence on handedness has been

inferred from family and adoption studies [14]. For instance, the probability of left-handedness

increases with the number of left-handed parents [15]. Twin studies reported slightly higher

rates of concordance in monozygotic (MZ) compared to dizygotic (DZ) twins [16,17] and provided

heritability estimates of around .25 [18,19].

Family studies have suggested differential effects of fathers and mothers to their offspring's

handedness. A stronger maternal than paternal effect was repeatedly found in biologically

related parent-offspring trios [20,21] and a similar trend was observable in an adoption study [22].

A maternal effect on non-right-handedness was also found in 592 families, where a paternal

effect was only detectable in males [23].

A recent large-scale genome-wide association study (GWAS; $n \sim 2M$) estimated that up to 6%

of the variance in left-handedness and up to 15% of the variance in ambidexterity are

explained by common genetic markers [24]. As in most large-scale laterality studies, handedness

was assessed as hand preference for writing, leading to three categories: right, left or both.

The "both" category identifies individuals who say that they can write equally well with both

hands, referred to as ambidextrous. However, a single task cannot identify mixed-handed

individuals who prefer different hands for different activities. Instead, self-report

questionnaires such as the Edinburgh Handedness Inventory (EHI) [25] assess the preferred hand

for several manual activities and therefore capture both mixed-handed and ambidextrous

individuals. A GWAS on brain imaging parameters (n = 32 256) revealed that genetic markers

associated with structural brain asymmetries overlapped with markers previously associated

with writing hand preference. Moreover, genetic factors involved in brain asymmetry overlap

with neurodevelopmental and cognitive traits such as ASD, schizophrenia, educational

attainment (EA) [26], and intelligence (IQ) [27]. These data suggest a general mechanism for the

establishment of left/right asymmetry which is also important for neurodevelopmental

outcomes. Therefore, the analysis of other lateralised preferences will contribute to the

understanding of such general mechanisms.

Foot and eye preference have received considerably less attention, even though associations

with neurodevelopmental disorders have been reported as well. For example, we found an

increased prevalence of non-right-footedness in neurodevelopmental and psychiatric

disorders (n_{cases} = 2 431, $n_{controls}$ = 116 938) [28]. Smaller studies point to higher rates of left eye

preference in schizophrenia (n_{cases} = 88, $n_{controls}$ = 118 [29]; n_{cases} = 68, $n_{controls}$ = 944 [30]) and ASD

(n_{cases} = 37; $n_{controls}$ = 20) [31]. Warren et al. [32] reported heritability estimates for foot and eye

preference to be .12 and .13, respectively. In Japanese twins, Suzuki and Ando [33] provided

heritability estimates for foot preference ranging from .08 to .24 and having one left-footed

parent increased the probability of being left-footed [34]. These studies support a genetic

component for foot and eye preference although there is variability in heritability estimates,

probably resulting from small sample sizes.

We performed the largest heritability study to date for multiple side preferences in the Avon

Longitudinal Study of Parents and Children (ALSPAC) and a twin cohort from Hong Kong to

investigate the heritability of laterality phenotypes, their associations with one another, and

their links to neurodevelopmental and cognitive outcomes.

Materials and Methods

Cohorts

ALSPAC: ALSPAC is a population-based longitudinal cohort. Pregnant women living in Avon,

UK, with expected dates of delivery from 1st April 1991 to 31st December 1992 were invited

to take part, resulting in 14 062 live births and 13 988 children who were alive at 1 year of age

[35,36]. Informed consent for the use of data collected via questionnaires and clinics was

obtained from participants following the recommendations of the ALSPAC Ethics and Law

Committee at the time. Ethical approval for the study was obtained from the ALSPAC Ethics

and Law Committee and the Local Research Ethics Committees. Please note that the study

website contains details of all the data that is available through a fully searchable data

dictionary and variable search tool (http://www.bristol.ac.uk/alspac/researchers/our-data/).

Hong Kong: Study participants were recruited from the Chinese-English Twin Study of

Biliteracy, a longitudinal study of primary school twin children starting in 2014 [37]. Participating

children were recruited from Hong Kong primary schools and had Cantonese as their native

language. Language and cognitive ability tests have been conducted for over four waves with

a one-year interval between assessments. Laterality data were collected during the second

wave of assessment.

Participants and phenotypes

ALSPAC: Laterality phenotypes were assessed for children based on maternal reports and for

parents as self-report. Hand preference was assessed using eleven items for parents and six

items for children. Foot preference and eye preference were assessed using four and two

items, respectively, for parents and children. All items were rated on a 3-point scale (coded as

left = 1, either = 2, right = 3, see Table S1). Two summary items (one in a right-mixed-left [R-

M-L] classification and one in a right-left classification [R-L]) were derived from recoded mean

values across non-missing items for hand, foot, and eye preference (see supplementary

methods and Figures S1-S3 for details). Mean ages of mothers, fathers, and children were

32.54 (SD = 4.42), 34.42 (SD = 5.60) and 3.55 (SD = 0.07) at the time of assessment,

respectively.

Hong Kong: The overall sample comprised *n* = 366 twin children (183 twin pairs) with a mean

age of 8.67 years (SD = 1.23). This sample included 81 MZ pairs (37 male pairs and 44 female

pairs) and 102 DZ pairs (21 male pairs, 19 female pairs, and 62 opposite-sex pairs). Twin

zygosity of same-sex twins was determined by genotyping small tandem repeat (STR) markers

on chromosomes 13, 18, 21, X and Y by Quantitative Fluorescence-Polymerase Chain Reaction

(QF-PCR).

Hand, foot, and eye preference were assessed using a modification of the EHI [25]. The

questionnaire was translated into Chinese and included six hand preference items, one foot

preference item, and one eye preference item. All items were read to participants by a trained

research assistant as described in detail previously [38]. Items were coded to a 3-point scale and

a R-M-L summary item was created for hand preference (see supplementary methods and

Figure S4 for details).

Genotype quality control (QC)

ALSPAC: Children's genotypes were generated on the Illumina HumanHap550-quad array at

the Wellcome Trust Sanger Institute, Cambridge, UK and the Laboratory Corporation of

America, Burlington, NC, US. Standard QC was performed as described elsewhere [39]. In total,

9 115 children and 500 527 SNPs passed QC filtering.

1. *Hong Kong:* Genotyping was performed using Illumina Human Infinium OmniZhongHua-8 v1.3

2. Beadchip at the Prenatal Genetic Diagnosis Centre and the Pre-implantation Genetic Diagnosis

3. laboratory in the Prince of Wales Hospital and The Chinese University of Hong Kong, Hong

4. Kong SAR. Standard quality control measures were carried out. Genetic variants with missing

5. rate > 10%, minor allele frequency (MAF) < 0.01 and with significant deviation from Hardy-

6. Weinberg equilibrium ($p < 1 \times 10^{-6}$) were excluded. Individuals with genotyping rates < 90%

7. and outlying heterozygosity rates were excluded. In total, 911 178 SNPs passed QC filtering.

8. Among the $n = 366$ twin children, genotype data were available for $n = 358$ (81 MZ pairs and

9. 98 DZ pairs).

10. *Parental effects*

11. We included parent-child trios with complete phenotypic data on the summary items for

12. hand, foot or eye preference after excluding one of each twin pair ($n = 113$) and children with

13. physical disabilities ($n = 65$) or sensory impairments ($n = 50$), resulting in a sample size (number

14. of trios) of $n_{hand} = 5\ 028$, $n_{foot} = 4\ 960$ and $n_{eye} = 4\ 762$ (see Table S1).

15. For hand, foot, and eye preference, we first performed two logistic regression analyses using

16. both parents' sidedness as a predictor (coded as 0 = two right-sided parents, 1 = one mixed-

17. sided parent, 2 = one left-sided parent, 3 = two mixed-sided parents, 4 = one mixed- and one

18. left-sided parent, 5 = two left-sided parents). This analysis was performed for child sidedness

19. (coded as right = 0, left = 1) using both the A) R-M-L classification (excluding mixed-sided

20. children and their parents) and the B) R-L classification.

21. Next, we differentiated maternal and paternal effects by using maternal sidedness, paternal

22. sidedness (both coded as right = 0, mixed = 1, left = 2), and offspring sex, as well as interaction

23. terms between maternal and paternal sidedness with offspring sex as predictors. We used the

wald.test() function to test for a difference between maternal and paternal effects using the

R-M-L and the R-L classification.

As non-paternity could affect these analyses, we reran the logistic regression analyses

including only confirmed biological parent-offspring trios as confirmed by genotype data.

Genotypes were available for n = 1 719 fathers. We used the R package Sequoia [40], which

assigns parents to offspring based on Mendelian errors. Sequoia uses birth year and sex to

decrease the number of potential relationships between individuals and to correctly infer

parents and offspring. As the exact birth year of children and parents in ALSPAC was unknown

to us, children's birth year was set to 1992 and parents' birth year was roughly estimated from

the age of the assessment of laterality data. We selected 500 SNPs randomly from a subset

that had MAF > 0.45, high genotyping rate (missingness < 0.01) and low linkage disequilibrium

(LD; r^2 < 0.01 within a 50 kb window). The 500 SNPs were spread across chromosomes 1 to 22.

Sequoia confirmed paternity for n = 1 624 fathers. Among this subsample of 1 624 trios,

complete phenotypic data were available for 1 161 trios for handedness, 1 150 trios for

footedness, and 1 105 trios for eyedness (see Table S1).

To assess the reliability of maternal reports, we performed Spearman rank correlation analysis

between hand preference for drawing (left/right) assessed by maternal report at age 3.5 and

self-reported hand preference for writing at age 7.5 (M_{age} = 7.50 years; n = 3 129).

Phenotypic analysis

Unrelated children (genetic relationship < 0.05, n = 5 956) with genome-wide genetic and

phenotypic data were selected for Genome-wide Complex Trait Analysis (GCTA) [41]. The same

sample was used for phenotypic analysis. Sample sizes varied from n = 4 630 (foot used to pick

up a pebble) to n = 5 931 (summary item for hand preference).

Summary items in the R-M-L classification for hand, foot, and eye preference and 12 single

items were residualised for sex, age, and the two most significant principal components:

[1] $Y_i = \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \beta_3 X_{3i} + \beta_4 X_{4i} + \varepsilon_i$

Where Y_i is the laterality summary item (coded as right = 0, mixed = 1, left = 2), β_0 is the

intercept, β_1 is the regression weight for offspring sex, X_{1i} is offspring sex, β_2 is the regression

weight for offspring age in weeks, X_{2i} is offspring age in weeks, β_3 is the regression weight for

PC1, X_{3i} is PC1, β_4 is the regression weight for PC2, X_{4i} is PC2, and ε_i reflects random error.

Phenotypes were then inverse rank-transformed to achieve normally distributed phenotypes.

Principal components were calculated based on directly genotyped (MAF < 0.05) and LD

pruned ($r^2 < 0.01$ within a 50 kb window) SNPs (excluding high LD regions) using Plink v2. The

rationale for including PCs in the phenotype transformation was based on the Genetic-

relationship-matrix structural equation modelling (GRM-SEM) method which has been

developed using the ALSPAC cohort [42]. As there is little population stratification in ALSPAC, the

PC effect on the phenotypes is very small. Instead, higher scores indicated being left-sided,

being female [43,44], and younger age. Phenotypic correlations between rank-transformed items

were calculated with Pearson correlation, applying FDR correction for 105 comparisons using

the Benjamini Hochberg method [45].

Heritability estimates

SNP-h^2 was calculated for the transformed R-M-L summary items (3) and single items (12)

using restricted maximum-likelihood (REML) analysis in GCTA [46], which compares phenotypic

similarity and genotypic similarity based on a genetic-relationship matrix (GRM) in unrelated

individuals. A GRM was estimated based on directly genotyped SNPs for unrelated children

(genetic relationship < 0.05, n = 5 956) using GCTA.

As a comparison, SNP-h^2 was calculated for the untransformed categorical items using sex, age, and the first two principal components as covariates. We estimated SNP-h^2 separately for left-sidedness (left vs. right, excluding mixed-sided individuals) and mixed-sidedness (mixed vs. right, excluding left-sided individuals).

Next, we estimated heritability from parent-offspring data [47]. Among the subsample with genomic data and confirmed paternity, we selected those with information on age at the time of laterality assessment, resulting in a sample of 1 000 trios for handedness, 991 trios for footedness, and 957 trios for eyedness. Summary items in the R-M-L classification for hand, foot, and eye preference (coded as right = 0, mixed = 1, left = 2) were transformed following the same procedure described above for the ALSPAC children. We estimated heritability by performing linear regression analyses using mean parental laterality as predictor and child laterality as the outcome:

[2] $Y_i = \beta_0 + \beta_1 X_i + \varepsilon_i$

Where Y_i is the transformed offspring laterality item, β_0 is the intercept, β_1 is the regression weight (heritability index), X_i is the mean parental laterality, and ε_i reflects random error.

SEM

We applied GRM-SEM [42] to quantify shared and unique genetic factors among R-M-L summary items for hand, foot, and eye preference. This method has recently been used to study genetic associations among language and literacy skills in the ALSPAC cohort [48]. Equivalent to heritability analysis in twin research, GRM-SEM partitions phenotypic variance/covariance into genetic and residual components, but estimates genetic variance/covariance based on genome-wide genetic markers. We used the same GRM described above (based on directly genotyped SNPs for n = 5 956 unrelated children using GCTA). A GRM-SEM was fitted using

the grmsem library in R (version 1.1.0) using all children with phenotypic data for at least one

phenotype. Multivariate trait variances were modelled using a saturated model (Cholesky

decomposition). GRM-SEM was also used to estimate bivariate heritability, i.e. the

contribution of genetic factors to the phenotypic covariance.

The heritability of laterality phenotypes was additionally estimated using a classical twin

design that compares the similarity of MZ to that of DZ twins. Since MZ twins share nearly all

their genetic variants, whereas DZ twins share on average 50% of their genetic variants, any

excess similarity of MZ twins over DZ twins is the result of genetic influences. This method

partitions phenotypic variance into that due to additive genetic (A), shared environmental (C)

and non-shared environmental influence (E). The variance attributed to each component can

be estimated using the structural equation modelling (SEM) technique and the proportion of

variance explained by the genetic influence (A) is termed heritability. Phenotypes were

transformed following the same procedure described for ALSPAC above. We fit a multivariate

ACE model to the transformed phenotypes (handedness, footedness, and eyedness) and

compared ACE with its constrained models, such as the AE model. Analyses were performed

using the OpenMx software package 2.18.1 [49]. The script was adapted from the International

Workshop on Statistical Genetic Methods for Human Complex Traits [50].

Polygenic risk score (PRS) analysis

We conducted PRS analyses using summary statistics for handedness assessed as a binary

trait, psychiatric and neurodevelopmental conditions (ASD, ADHD, bipolar disorder (BIP),

schizophrenia (SCZ)), and cognitive measures (EA and IQ) using PRSice 2.3.3 [51]. PRS analyses

were performed for hand and foot preference (which showed significant SNP-h^2). The

summary statistics for hand preference (left vs. right) were calculated after excluding

individuals from 23andMe as well as ALSPAC from the original GWAS [24] sample. Summary

statistics for ADHD [52], ASD [53], BIP [54], and SCZ [55] were accessed from the Psychiatric Genomics

Consortium (PGC) website (https://www.med.unc.edu/pgc/data-index/). Summary statistics

for IQ [56] and EA [57] were accessed from the Complex Trait Genetics (CTG) lab website

(https://ctg.cncr.nl/software/summary_statistics), and the Social Science Genetic Association

Consortium (https://www.thessgac.org/data), respectively.

PRS were derived from LD-clumped SNPs ($r^2 < 0.1$ within a 250 kb window) as the weighted

sum of risk alleles according to the training GWAS summary statistics. No covariates were

included as phenotypes had been corrected for effects of age, sex, and ancestry. Results are

presented for the best training GWAS *p*-value threshold (explaining maximum phenotypic

variance) as well as GWAS *p*-value thresholds of .001, 0.05, .1, .2, .3, .4, .5, and 1. Results were

FDR-corrected for 126 comparisons (7 training GWAS; 2 target phenotypes; 9 *p*-value

thresholds) using the Benjamini-Hochberg method [45].

Code availability

Data preparation and visualization were performed using R v.4.0.0. Analysis scripts are

available through Github (https://github.com/Judith-Schmitz/heritability_hand_foot_eye).

Results

Parental effects

We tested parental effects by assessing the percentages of non-right-sided (R-M-L) and left-

sided (R-L) offspring as a function of parental sidedness in the whole sample and in trios with

confirmed biological paternity. As expected, the percentage of non-right-sidedness and left-

sidedness were highest in individuals with two non-right-sided or two left-sided parents,

respectively (Tables S2 [R-M-L] and S3 [R-L]). The percentage of non-right-sidedness and left-

sidedness were higher in individuals with a non-right-sided or left-sided mother and a right-

sided father than vice versa for all three traits. This effect was visible in both the whole sample

(e.g. 31.23% vs. 25.83% for non-right-handedness, see Table S2) and in the subset with

confirmed biological paternity (e.g. 33.33% vs. 25.37%, see Table S2).

Second, we ran logistic regression analyses in $n \leq 5\,028$ ALSPAC family trios. In the R-M-L

classification (n_{hand} = 4 248, n_{foot} = 3 242 and n_{eye} = 3 050), parental sidedness predicted hand,

$X^2(5)$ = 39.5, p = 1.9×10^{-7}, foot, $X^2(5)$ = 59.9, p = 1.3×10^{-11}, and eye preference, $X^2(5)$ = 27.4,

p = 4.8×10^{-5}. In the R-L classification (n_{hand} = 5 028, n_{foot} = 4 960 and n_{eye} = 4 762), parental

sidedness also predicted hand, $X^2(2)$ = 42.6, p = 5.5×10^{-10}, foot, $X^2(2)$ = 69.1, p = 1.0×10^{-15},

and eye preference, $X^2(2)$ = 14.6, p = 6.9×10^{-4}. ORs show that having one or two left-sided

parents increased one's chances to be left-sided for hand, foot, and eye preference in the R-

M-L classification (Figure 1A) and in the R-L classification (Figure 1B). Analysis in the subsample

with confirmed paternity ($n \leq 1\,161$ family trios) showed similar, although attenuated,

parental effects for hand (R-M-L: $X^2(4)$ = 14.9, p = .005; R-L: $X^2(2)$ = 12.1, p = .002) and foot (R-

M-L: $X^2(4)$ = 22.5, p = .0002; R-L: $X^2(2)$ = 19.1, p = 7.1×10^{-5}), but not for eye (R-M-L: $X^2(5)$ =

5.3, p = .380; R-L: $X^2(2)$ = 2.7, p = .250) preference (Figure S5). The full regression model

outputs for the whole sample and for trios with confirmed paternity can be found in Tables
S4-S7.

Figure 1: Parental effects on child sidedness. *ORs [95% CI], resulting from logistic regression analysis.*

Third, we investigated maternal and paternal effects and possible interactions with offspring

sex. In the whole sample, Wald tests revealed a significant maternal effect on hand (R-M-L:

$X^2(4) = 38.9$, $p = 7.4 \times 10^{-8}$; R-L: $X^2(2) = 31.7$, $p = 1.3 \times 10^{-7}$), foot (R-M-L: $X^2(4) = 52.7$, $p = 9.8 \times$

10^{-11}; R-L: $X^2(2) = 96.6$, $p < 2.2 \times 10^{-16}$), and eye preference (R-M-L: $X^2(4) = 38.3$, $p = 9.7 \times 10^{-8}$;

R-L: $X^2(2) = 34.1$, $p = 3.9 \times 10^{-8}$). Paternal sidedness predicted hand (R-M-L: $X^2(4) = 10.3$, $p =$

.036; R-L: $X^2(2) = 12.3$, $p = .002$) and foot (R-M-L: $X^2(4) = 15.1$, $p = .005$; R-L: $X^2(2) = 6.0$, $p =$

.049), but not eye preference (R-M-L: $X^2(4) = 4.6$, $p = .330$; R-L: $X^2(2) = 0.6$, $p = .760$). Wald tests

contrasting maternal and paternal effects revealed a stronger maternal than paternal effect

only for foot preference (R-M-L: $X^2(1) = 4.6$, $p = .033$; R-L: $X^2(1) = 23.9$, $p = 1.0 \times 10^{-6}$). This

effect was confirmed in the subsample with confirmed paternity (R-M-L: $X^2(1) = 8.4$, $p = .004$;

R-L: $X^2(1) = 10.0$, $p = .002$). Although attenuated in the smaller subsample with confirmed

paternity, this finding suggests a genuinely stronger maternal than paternal effect on

footedness. In the whole sample, interaction terms between maternal/paternal sidedness and

offspring sex revealed that in the R-L classification, maternal left-sidedness had a greater

effect on left-footedness in girls compared to boys (ß = 0.49, SE = 0.19, z = 2.55, p = .011),

which was confirmed in the smaller subsample (ß = 0.96, SE = 0.44, z = 2.17, p = .030). The full

regression model outputs for both the whole sample and the subsample with confirmed

paternity can be found in Tables S8-S11.

Besides non-paternity, the reliability of the maternal report on laterality phenotypes could

have affected our analysis. Correlation analysis showed a strong association between hand

preference for drawing collected at 3.5 years of age and the self-reported hand preferred for

writing at age 7.5 (r = .95, 95% CI = [.93, .97], p < 2.2×10^{-16}). Among the 2 838 children with a

right-hand preference at age 3.5, seven reported a left-hand preference for writing at age 7.5.

Of the 291 children with left-hand preference at age 3.5, 19 showed a right-hand preference

for writing at age 7.5. Overall, 99.2% of individuals showed stable hand preference (see Table

S12), demonstrating the reliability of the maternal report.

Transformed phenotypes

Phenotypic correlation and genomic analyses (SNP-h^2 estimates, GRM-SEM and PRS analysis)

were performed in unrelated children from the ALSPAC cohort (n ≤ 5 931). Multivariate

behavioural SEM analysis was performed In the Hong Kong twin sample (n ≤ 358). The absolute

numbers and percentages of children with left, mixed and right side preference for the three

summary items in both cohorts are shown in Table 1.

Table 1: Children with left, mixed and right side preference for each phenotype in ALSPAC (unrelated children) and the Hong Kong cohort (twin children).

	ALSPAC				Hong Kong			
	n	Left	Mixed	Right	*n*	Left	Mixed	Right
Hand preference	5 931	471 (7.9%)	893 (15.1%)	4567 (77.0%)	358	20 (5.6%)	37 (10.3%)	301 (84.1%)
Foot preference	5 860	344 (5.9%)	2070 (35.3%)	3446 (58.8%)	358	31 (8.7%)	106 (29.6%)	221 (61.7%)
Eye preference	5 650	730 (12.9%)	2012 (35.6%%)	2908 (51.5%)	357	95 (26.5%)	107 (29.9%)	155 (43.4%)

By regressing out the effects of sex, age, and ancestry, we transformed laterality categories into quantitative measures using formula [1]. We assessed phenotypic correlations for the transformed items in ALSPAC and the Hong Kong cohort. In ALSPAC, the single item that best captured the summary item was "hand used to draw" for hand preference ($r = .87$, $t_{(5920)} = 139.01$, $p < 2.2 \times 10^{-16}$), "foot used to stamp" for foot preference ($r = .78$, $t_{(5765)} = 95.78$, $p < 2.2 \times 10^{-16}$), and "eye used to look through a bottle" for eye preference ($r = .96$, $t_{(5469)} = 249.61$, $p < 2.2 \times 10^{-16}$). In both cohorts, summary items showed positive correlations with each other (Figure S6, Figure S7). These correlations support a general left/right directionality captured by the different items.

Heritability estimates

We then tested the heritability of the transformed phenotypes. SNP-h^2 of transformed laterality items ranged from .00 ($p = .500$) for "eye used to look through a bottle" to .42 ($p = 8 \times 10^{-13}$) for "hand used to cut" (Figure 2, Table S13). The highest heritability estimate for summary measures was observed for footedness (SNP-$h^2 = .23$; $p = 2 \times 10^{-5}$), followed by handedness (SNP-$h^2 = .21$; $p = 1 \times 10^{-4}$). There was no significant SNP-h^2 for eyedness (SNP-$h^2 = .00$; $p = .469$).

Figure 2: SNP-h² estimates for laterality measures after transformation into quantitative scores in

ALSPAC. Results are shown for individual items and summary measures (yellow). Bars represent

standard errors.

For comparison, we estimated the SNP-h² for the untransformed categorical items for left-

and mixed-side preference categories. SNP-h² for left-side preference ranged from .00 (p =

.500) for "foot used to climb a step" to .13 (p = .031) for "hand used to cut" (Figure S8A, Table

S14). SNP-h² for mixed-side preference ranged from .00 (p = .500) for the hand preference

summary item to .12 (p = .031) for "hand used to draw" (Figure S8B, Table S15).

Parent-offspring regression run on the transformed summary items suggested heritability

estimates of .27 for handedness (95% CI = [.11, .42], p = 5.6 × 10⁻⁴), .09 for footedness (95% CI

= [.01, .17], p = .030), and .08 for eyedness (95% CI = [-.04, .20], p = .198).

Univariate SEM analysis in the Hong Kong cohort resulted in heritability estimates of .45 for

handedness (95% CI = [.29, .63]), .08 for footedness (95% CI = [.00, .25]), and .08 for eyedness

(95% CI = [.00, .26]). Therefore, the heritability estimates for the quantitative phenotypes

were consistently higher than for categorical measures, both for SNP-h^2, parent-offspring, and

twin SEM estimates.

SEM

Multivariate GRM-SEM analysis was performed on the transformed R-M-L summary items for

handedness, footedness, and eyedness in ALSPAC (Figure 3A). The squared path coefficient of

genetic factor A1 explains genetic variance in hand preference (a_{11}) and genetic variance that

is shared with foot (a_{21}) and eye preference (a_{31}). A single genetic factor (A1) explained 20.36%

of the phenotypic variance in handedness (a_{11} = 0.45, p = 2.4 × 10^{-12}), 22.12% of the variance

in footedness (a_{21} = 0.47, p = 9.2 × 10^{-10}) and 3.84% of the variance in eyedness (a_{31} = 0.20, p =

9.2 × 10^{-3}). All other path coefficients were non-significant, suggesting that one shared genetic

factor (A1) contributes to a general left/right directionality across all three phenotypes.

Figure 3: Results of SEM analyses between laterality phenotypes. *A) Results of GRM-SEM in ALSPAC.*

B) Results of behavioural SEM in the Hong Kong cohort. Circles on top and bottom indicate genetic (A)

and environmental (E) factors, respectively. Coloured boxes indicate the phenotypes. Solid lines indicate

significant path coefficients, dotted lines indicate non-significant path coefficients. White boxes

indicate path coefficients and standard errors (SE) for significant genetic factors. The contour of the

white boxes indicates the genetic factor (A1 in all cases).

Bivariate heritability analysis confirmed that shared genetic influences accounted for 36.7%

of the phenotypic correlation between handedness and footedness ($p = 6.6 \times 10^{-6}$), 24.9% of

the correlation of between footedness and eyedness ($p = .020$), and 26.2% of the correlation

between handedness and eyedness ($p = .020$). We replicated these findings with multivariate

behavioural SEM in an independent cohort ($n = 358$). In the Hong Kong cohort, A1 explained

44.30% (95% CI = [28.50, 62.30]) of the phenotypic variance in handedness ($a_{11} = 0.67$, $p <$

.001), 5.00% (95% CI = [0.20, 15.30]) of the variance in footedness ($a_{21} = 0.22$, $p = .014$), and

7.00% (95% CI = [0.80, 18.20]) of the variance in eyedness ($a_{31} = 0.26$, $p = .003$) (Figure 3B). All

other path coefficients were non-significant, consistent with results for ALSPAC.

PRS analysis

None of the PRS associations survived correction for multiple comparisons. The strongest

association was found for PRS for IQ, suggesting that genetic predisposition towards higher IQ

is associated with a tendency towards right-handedness ($ß = -1159.21$, SE = 414.71, PRS $R^2 =$

0.13%, $p = .005$). PRS results for all *p*-value thresholds are reported in Table S16.

Discussion

We investigated the heritability of hand, foot, and eye preference using multiple approaches. To the best of our knowledge, this is the largest study conducted to date for multiple laterality measures in the same individuals. Our analysis of family trios showed that the probability of being left-sided increased for any left-sided parent on the same trait, with stronger effects for hand and foot, rather than eye preference, in line with previous reports [15,34]. Stronger maternal than paternal effects have been reported in studies focussing mainly on handedness [20,23]. In ALSPAC, we found a stronger maternal than paternal effect for foot, but not hand or eye preference. This stronger maternal effect was detected in the whole sample (n = 4 960 trios) and confirmed in the subset with genetically confirmed paternity (n = 1 150 trios). Maternal/paternal effects could be explained with sex-linked genetic or parent-of-origin effects. For example, the imprinted *LRRTM1* gene was found to be associated with handedness under a parent of origin effect [58]. Parent of origin effects might be more wide-spread than appreciated, but their detection requires family samples as opposed to the most commonly used singleton cohorts [59]. Few examples of parent-of-origin effects have been reported, for example for language-related measures [60–62]. Besides non-paternity, the reliability of the maternal report on laterality phenotypes could have affected our analysis. We confirmed strong correlation (r = .95) between the preferred hand for drawing assessed using maternal report at age 3.5 and self-reported preferred hand for writing in later childhood. The fact that more children switch hand preference from left to right [63] could indirectly suggest that switching attempts by parents or teachers have occurred at least until the mid 1990s. Overall, our analysis supports a genetic component underlying these laterality traits and highlights a specific maternal effect for footedness. The maternal effects could result from a higher genetic load required to manifest left-side preference in females. A similar

buffering effect has been proposed to explain the higher prevalence of neurodevelopmental

disorders in males [64].

Using transformed quantitative phenotypes [48], we estimated SNP-h^2 for handedness,

footedness, and eyedness to be .21, .23, and .00, respectively. The heritability estimate for

handedness is similar to what has been reported in behavioural twin studies (h^2 = .25) [18,19] but

higher than observed in GWAS (SNP-h^2 = .06) [24,65,66] for categorical handedness. Instead,

estimates for categorical phenotypes were non-significant, suggesting that the transformed

phenotypes are better suited to detect the genetic component underlying lateralised traits

than binary phenotypes. Accordingly, behavioural analysis in the Hong Kong twin cohort

revealed a heritability estimate of .45 for the quantitative handedness phenotype - much

higher than what has been observed for a categorical measure of writing hand (.27) in the

same cohort [38]. Parent-offspring regression in ALSPAC also showed significant heritability for

handedness and footedness when using the quantitative phenotypes. We conclude that the

quantitative phenotypes are better suited to capture the polygenic nature of handedness as

expected under a liability threshold model [67]. The lack of association between the PRS derived

from a recent large-scale GWAS for categorical handedness [24] suggests the influence of

separate genetic factors. Lack of heritability for eyedness could reflect the poor quality of

phenotype assessment, i.e. eyedness might be more difficult to assess and report accurately.

Another possibility is that human eye preference does not have particular functional

advantages and therefore the preferred side is less influenced by evolutionary forces and

genetic factors. This is in contrast to other vertebrates such as bird [68] or fish species [69], where

eye preference is involved in predator detection or social interaction.

Heritability estimates differed substantially between items used to assess handedness,

footedness, and eyedness. We found the highest SNP-h^2 for "hand used to cut" (with a knife).

Previously, this item showed the weakest phenotypic correlation with the other questionnaire items [70,71] and the highest heritability [33]. It has been proposed that summary items have reduced value to determine genetic factors involved in laterality [32]. This was true for the handedness measure, but conversely, we observed higher SNP-h^2 for the summary rather than single footedness items in ALSPAC, suggesting that in contrast to handedness, multiple items might better capture a genetic component for footedness. One possible interpretation is that multiple items will allow identifying mixed-footed rather than ambipedal individuals, who prefer both feet equally. Similar to Suzuki and Ando [33], our results suggest that the item "foot used to kick a ball", which is often used as the only assessment item, is not the optimal choice to investigate the heritability of footedness. We previously showed that assessing footedness in terms of kicking systematically under-estimates the prevalence of mixed-footedness when compared to assessment using footedness inventories [28]. Overall, there is no one correct measure for laterality items, however, our results demonstrate the importance of reporting data for single items [72] in addition to the aggregates and suggest the value of using multiple items.

All transformed items showed positive correlations on the phenotypic level. Previous research has shown a tendency towards a higher probability of left-sided lateral preferences in left-handers [28,73], suggesting that a common dimension of asymmetry underlies hand, foot, and eye preference [74]. Multivariate SEM analysis supported the presence of one shared genetic factor explaining variance in handedness, footedness, and eyedness, but no unique genetic factors explaining independent variance for individual phenotypes in ALSPAC and the Hong Kong cohort. In ALSPAC, bivariate heritability analysis suggested that up to 37% of the phenotypic correlation is due to shared genetic effects.

An association between laterality and psychiatric disorders, especially schizophrenia [75], has long been debated. Of the different traits tested, we found suggestive evidence that PRS for IQ were associated with a tendency towards right-handedness, but not with footedness. Similarly, a recent dyslexia GWAS found positive genetic correlation between dyslexia and ambidexterity [76]. A possible explanation for a specific link between cognitive measures and handedness is its association with language. It has been suggested that the higher prevalence of human right- than left-handedness has arisen from a left-hemispheric dominance for manual gestures that gradually incorporated vocalisation [77]. Indeed, right-handers produce more right- than left-handed gestures when speaking [78]. This would suggest that footedness and eyedness are phenotypically secondary to handedness, as has been suggested previously [79].

Conclusion

We assessed the heritability of multiple side preferences using family, genomic, and twin analyses. For footedness, stronger maternal than paternal effects highlight the necessity of examining parent-of-origin effects on the genetic level in future studies. SEM supports a shared genetic factor involved in all three phenotypes without independent genetic factors contributing to footedness and eyedness. The transformed quantitative phenotypes present a heritability that is higher than categorical measures in both molecular and behavioural analyses, suggesting that they might be better suited to identify the underlying genetic factors.

Acknowledgments

We are extremely grateful to all the families who took part in this study, the midwives for their

help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer

and laboratory technicians, clerical workers, research scientists, volunteers, managers,

receptionists and nurses. We also thank Andrew P. Morris and Beate St Pourcain for advising

on statistical analyses. The authors would also like to express their gratitude to Sarah

Medland, Gabriel Cuellar Partida, and David Evans for providing the handedness GWAS

summary statistics.

The UK Medical Research Council and Wellcome (Grant ref: 217065/Z/19/Z) and the University

of Bristol provide core support for ALSPAC. This publication is the work of the authors and SP

and JS will serve as guarantors for the analysis of the ALSPAC data presented in this paper.

GWAS data were generated by Sample Logistics and Genotyping Facilities at Wellcome Sanger

Institute and LabCorp (Laboratory Corporation of America) using support from 23andMe.

Support to the genetic analysis was provided by the St Andrews Bioinformatics Unit funded

by the Wellcome Trust [grant 105621/Z/14/Z]. The Hong Kong sample was funded through a

Collaborative Research Fund from the Hong Kong Special Administrative Region Research

Grants Council (CUHK8/CRF/13G, and C4054-17WF). JS is funded by the Deutsche

Forschungsgemeinschaft (DFG, German Research Foundation, 418445085) and supported by

the Wellcome Trust [Institutional Strategic Support fund, Grant number 204821/Z/16/Z]. For

the purpose of open access, the author has applied a CC BY public copyright licence to any

Author Accepted Manuscript version arising from this submission. SP is funded by the Royal

Society (UF150663).

Conflict of interest

The authors declare no competing financial interests in relation to the work described.

References

1. Ocklenburg S., Hirnstein M., Beste C. & Güntürkün O. Lateralization and cognitive systems. *Front Psychol* **5**, 1143 (2014).

2. Knecht S. *et al.* Handedness and hemispheric language dominance in healthy humans. *Brain* **123 Pt 12**, 2512–2518 (2000).

3. Tzourio-Mazoyer N. & Mazoyer B. Variations of planum temporale asymmetries with Heschl's Gyri duplications and association with cognitive abilities: MRI investigation of 428 healthy volunteers. *Brain Struct Funct* **222**, 2711–2726 (2017).

4. Tzourio-Mazoyer N., Crivello F. & Mazoyer B. Is the planum temporale surface area a marker of hemispheric or regional language lateralization? *Brain Struct Funct* **223**, 1217–1228 (2018).

5. Altarelli I. *et al.* Planum temporale asymmetry in developmental dyslexia: Revisiting an old question. *Hum Brain Mapp* **35**, 5717–5735 (2014).

6. Elnakib A., Soliman A., Nitzken M., Casanova M.F., Gimel'farb G. & El-Baz A. Magnetic resonance imaging findings for dyslexia: a review. *J Biomed Nanotechnol* **10**, 2778–2805 (2014).

7. Oertel-Knöchel V., Knöchel C., Matura S., Prvulovic D., Linden D.E.J. & van de Ven V. Reduced functional connectivity and asymmetry of the planum temporale in patients with schizophrenia and first-degree relatives. *Schizophr Res* **147**, 331–338 (2013).

8. Floris D.L. *et al.* Atypically rightward cerebral asymmetry in male adults with autism stratifies individuals with and without language delay. *Hum Brain Mapp* **37**, 230–253 (2016).

9. Papadatou-Pastou M. *et al.* Human handedness: A meta-analysis. *Psychol Bull* **146**, 481–524 (2020).

10. Schmitz J., Metz G.A.S., Güntürkün O. & Ocklenburg S. Beyond the genome-Towards an epigenetic understanding of handedness ontogenesis. *Prog Neurobiol* **159**, 69–89 (2017).

11. de Kovel C.G.F., Carrión-Castillo A. & Francks C. A large-scale population study of early life factors influencing left-handedness. *Sci Rep* **9**, 584 (2019).

12. Markou P., Ahtam B. & Papadatou-Pastou M. Elevated Levels of Atypical Handedness in Autism: Meta-Analyses. *Neuropsychol Rev* **27**, 258–283 (2017).

13. Hirnstein M. & Hugdahl K. Excess of non-right-handedness in schizophrenia: meta-analysis of gender effects and potential biases in handedness assessment. *Br J Psychiatry* **205**, 260–267 (2014).

14. Carter-Saltzman L. Biological and sociocultural effects on handedness: comparison between biological and adoptive families. *Science* **209**, 1263–1265 (1980).

15. McManus I.C. & Bryden M.P. in *Handbook of neuropsychology:* (eds Rapin, I. & Segalowitz, S. J.) pp 115–44 (Elsevier Science1992).

16. Pfeifer L.S., Schmitz J., Papadatou-Pastou M., Peterburs J., Paracchini S. & Ocklenburg S. Handedness in Twins: Meta-Analyses. *PsyArXiv*, https://10.31234/osf.io/gy2nx (2021).

17. Sicotte N.L., Woods R.P. & Mazziotta J.C. Handedness in twins: a meta-analysis. *Laterality* **4**, 265–286 (1999).

18. Medland S.E., Duffy D.L., Wright M.J., Geffen G.M. & Martin N.G. Handedness in twins: joint analysis of data from 35 samples. *Twin Res Hum Genet* **9**, 46–53 (2006).

19. Medland S.E. *et al.* Genetic influences on handedness: data from 25,732 Australian and Dutch twin families. *Neuropsychologia* **47**, 330–337 (2009).

20. Coren S. & Porac C. Family patterns in four dimensions of lateral preference. *Behav Genet* **10**, 333–348 (1980).

21. Reiss M. Genetic associations between lateral signs. *Anthropol Anz* **57**, 61–68 (1999).

22. Saudino K. & McManus I.C. Handedness, footedness, eyedness and earedness in the Colorado Adoption Project. *British Journal of Developmental Psychology* **16**, 167–174 (1998).

23. Tran U.S. & Voracek M. Evidence of Sex-Linked Familial Transmission of Lateral Preferences for Hand, Foot, Eye, Ear, and Overall Sidedness in a Latent Variable Analysis. *Behav Genet* **45**, 537–546 (2015).

24. Cuellar-Partida G. *et al.* Genome-wide association study identifies 48 common genetic variants associated with handedness. *Nat Hum Behav* **5**, 59–70 (2021).

25. Oldfield R.C. The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia* **9**, 97–113 (1971).

26. Sha Z. *et al.* The genetic architecture of structural left-right asymmetry of the human brain. *Nat Hum Behav*, https://10.1038/s41562-021-01069-w (2021).

27. Kong X.-Z. *et al.* Large-Scale Phenomic and Genomic Analysis of Brain Asymmetrical Skew. *Cereb Cortex*, https://10.1093/cercor/bhab075 (2021).

28. Packheiser J. *et al.* Four meta-analyses across 164 studies on atypical footedness prevalence and its relation to handedness. *Sci Rep* **10**, 14501 (2020).

29. Dane S. *et al.* Handedness, eyedness, and hand--eye crossed dominance in patients with schizophrenia: sex-related lateralisation abnormalities. *Laterality* **14**, 55–65 (2009).

30. Giotakos O. Crossed hand-eye dominance in male psychiatric patients. *Percept Mot Skills* **95**, 728–732 (2002).

31. Dane S. & Balci N. Handedness, eyedness and nasal cycle in children with autism. *Int J Dev Neurosci* **25**, 223–226 (2007).

32. Warren D.M., Stern M., Duggirala R., Dyer T.D. & Almasy L. Heritability and linkage analysis of hand, foot, and eye preference in Mexican Americans. *Laterality* **11**, 508–524 (2006).

33. Suzuki K. & Ando J. Genetic and environmental structure of individual differences in hand, foot, and ear preferences: a twin study. *Laterality* **19**, 113–128 (2014).

34. Ooki S. Nongenetic factors associated with human handedness and footedness in Japanese twin children. *Environ Health Prev Med* **11**, 304–312 (2006).

35. Boyd A. *et al.* Cohort Profile: the 'children of the 90s'--the index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol* **42**, 111–127 (2013).

36. Fraser A. *et al.* Cohort Profile: the Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. *Int J Epidemiol* **42**, 97–110 (2013).

37. Wong S.W.L., Ho C.S.-H., McBride C., Chow B.W.-Y. & Waye M.M.Y. Less is More in Hong Kong: Investigation of Biscriptal and Trilingual Development Among Chinese Twins in a (Relatively) Small City. *Twin Res Hum Genet* **20**, 66–71 (2017).

38. Zheng M., McBride C., Ho C.S.-H., Chan J.K.-C., Choy K.W. & Paracchini S. Prevalence and heritability of handedness in a Hong Kong Chinese twin and singleton sample. *BMC Psychol* **8**, 37 (2020).

39. Brandler W.M. *et al.* Common variants in left/right asymmetry genes and pathways are associated with relative hand skill. *PLoS Genet* **9**, e1003751 (2013).

40. Huisman J. Pedigree reconstruction from SNP data: parentage assignment, sibship clustering and beyond. *Mol Ecol Resour* **17**, 1009–1024 (2017).

41. Yang J., Lee S.H., Goddard M.E. & Visscher P.M. GCTA: a tool for genome-wide complex trait analysis. *Am J Hum Genet* **88**, 76–82 (2011).

42. St Pourcain B. *et al.* Developmental Changes Within the Genetic Architecture of Social Communication Behavior: A Multivariate Study of Genetic Variance in Unrelated Individuals. *Biol Psychiatry* **83**, 598–606 (2018).

43. Papadatou-Pastou M., Martin M., Munafò M.R. & Jones G.V. Sex differences in left-handedness: a meta-analysis of 144 studies. *Psychol Bull* **134**, 677–699 (2008).

44. Buenaventura Castillo C., Lynch A.G. & Paracchini S. Different laterality indexes are poorly correlated with one another but consistently show the tendency of males and females to be more left- and right-lateralized, respectively. *R Soc Open Sci* **7**, 191700 (2020).

45. Benjamini Y. & Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)* **57**, 289–300 (1995).

46. Yang J. *et al.* Common SNPs explain a large proportion of the heritability for human height. *Nat Genet* **42**, 565–569 (2010).

47. Fernandez G.C. & Miller J.C. Estimation of heritability by parent-offspring regression. *Theor Appl Genet* **70**, 650–654 (1985).

48. Verhoef E., Shapland C.Y., Fisher S.E., Dale P.S. & St Pourcain B. The developmental origins of genetic factors influencing language and literacy: Associations with early-childhood vocabulary. *J Child Psychol Psychiatry*, https://10.1111/jcpp.13327 (2020).

49. Boker S. *et al.* OpenMx: An Open Source Extended Structural Equation Modeling Framework. *Psychometrika* **76**, 306–317 (2011).

50. Neale M., Eaves L., Bartels M., Boomsma D.I., Posthuma D. & Bates T. The 2018 International Workshop on Statistical Genetic Methods for Human Complex Traits. *Behav Genet* **47**, 729–730 (2017).

51. Choi S.W. & O'Reilly P.F. PRSice-2: Polygenic Risk Score software for biobank-scale data. *Gigascience* **8** (2019).

52. Demontis D. *et al.* Discovery of the first genome-wide significant risk loci for attention deficit/hyperactivity disorder. *Nat Genet* **51**, 63–75 (2019).

53. Grove J. *et al.* Identification of common genetic risk variants for autism spectrum disorder. *Nat Genet* **51**, 431–444 (2019).

54. Mullins N. *et al.* Genome-wide association study of more than 40,000 bipolar disorder cases provides new insights into the underlying biology. *Nat Genet* **53**, 817–829 (2021).

55. Schizophrenia Working Group of the Psychiatric Genomics Consortium. Biological insights from 108 schizophrenia-associated genetic loci. *Nature* **511**, 421–427 (2014).

56. Savage J.E. *et al.* Genome-wide association meta-analysis in 269,867 individuals identifies new genetic and functional links to intelligence. *Nat Genet* **50**, 912–919 (2018).

57. Lee J.J. *et al.* Gene discovery and polygenic prediction from a genome-wide association study of educational attainment in 1.1 million individuals. *Nat Genet* **50**, 1112–1121 (2018).

58. Francks C. *et al.* LRRTM1 on chromosome 2p12 is a maternally suppressed gene that is associated paternally with handedness and schizophrenia. *Mol Psychiatry* **12**, 1129-39, 1057 (2007).

59. Hofmeister R.J., Rubinacci S., Ribeiro D.M., Kutalik Z., Buil A. & Delaneau O. *Parent-of-origin effects in the UK Biobank* (2021).

60. Nudel R. *et al.* Genome-wide association analyses of child genotype effects and parent-of-origin effects in specific language impairment. *Genes Brain Behav* **13**, 418–429 (2014).

61. Pettigrew K.A. *et al.* Further evidence for a parent-of-origin effect at the NOP9 locus on language-related phenotypes. *J Neurodev Disord* **8**, 24 (2016).

62. Hitchcock T.J., Paracchini S. & Gardner A. Genomic Imprinting As a Window into Human Language Evolution. *Bioessays* **41**, e1800212 (2019).

63. Fennell E.B., Satz P. & Morris R. The development of handedness and dichotic ear listening asymmetries in relation to school achievement: A longitudinal study. *Journal of Experimental Child Psychology* **35**, 248–262 (1983).

64. Polyak A., Rosenfeld J.A. & Girirajan S. An assessment of sex bias in neurodevelopmental disorders. *Genome Med* **7**, 94 (2015).

65. de Kovel C.G.F. & Francks C. The molecular genetics of hand preference revisited. *Sci Rep* **9**, 5986 (2019).

66. Wiberg A. *et al.* Handedness, language areas and neuropsychiatric diseases: insights from brain imaging and genetics. *Brain* **142**, 2938–2947 (2019).

67. Paracchini S. Recent Advances in Handedness Genetics. *Symmetry* **13**, 1792 (2021).

68. Koboroff A., Kaplan G. & Rogers L.J. Hemispheric specialization in Australian magpies (Gymnorhina tibicen) shown as eye preferences during response to a predator. *Brain Res Bull* **76**, 304–306 (2008).

69. De Santi A., Bisazza A. & Vallortigara G. Complementary left and right eye use during predator inspection and shoal-mate scrutiny in minnows. *Journal of Fish Biology* **60**, 1116–1125 (2002).

70. Veale J.F. Edinburgh Handedness Inventory - Short Form: a revised version based on confirmatory factor analysis. *Laterality* **19**, 164–177 (2014).

71. McFarland K. & Anderson J. Factor stability of the Edinburgh Handedness Inventory as a function of test-retest performance, age and sex. *British Journal of Psychology* **71**, 135–142 (1980).

72. Packheiser J. *et al.* Using Mobile EEG to Investigate Alpha and Beta Asymmetries During Hand and Foot Use. *Front Neurosci* **14**, 109 (2020).

73. Bourassa D.C., McManus I.C. & Bryden M.P. Handedness and eye-dominance: a meta-analysis of their relationship. *Laterality* **1**, 5–34 (1996).

74. Porac C. Eye preference patterns among left-handed adults. *Laterality* **2**, 305–316 (1997).

75. Crow T.J. The 'big bang' theory of the origin of psychosis and the faculty of language. *Schizophr Res* **102**, 31–52 (2008).

76. Doust C. *et al. Discovery of 42 Genome-Wide Significant Loci Associated with Dyslexia* (2021).

77. Corballis M.C. From mouth to hand: gesture, speech, and the evolution of right-handedness. *Behav Brain Sci* **26**, 199-208; discussion 208-60 (2003).

78. Kimura D. Manual activity during speaking— I. Right-handers. *Neuropsychologia* **11**, 45–50 (1973).

79. McManus I.C., Porac C., Bryden M.P. & Boucher R. Eye-dominance, writing hand, and throwing hand. *Laterality* **4**, 173–192 (1999).