Will Current Protected Areas Harbour Refugia for Threatened Arctic Vegetation Types until

2050?

Merin Reji Chacko[1,2,3*], Ariane K.A. Goerens[1], Jacqueline Oehri[1,4], Elena Plekhanova[1] and Gabriela

Schaepman-Strub[1*]

Abstract

Arctic vegetation is crucial for fauna and the livelihoods of Northern peoples, and tightly linked to

climate, permafrost soils, and water. Yet, a comprehensive understanding of climate change effects

on Arctic vegetation is lacking. Protected areas cannot halt climate change, but could reduce future

pressure from additional drivers, such as land use change and local industrial pollution. Therefore, it

is crucial to understand the contribution of protected areas in safeguarding threatened Arctic

vegetation types. We compare the 2003 baseline with existing 2050 predictions of circumpolar Arctic

vegetation type distributions and demonstrate an overrepresentation of dominant and

underrepresentation of declining vegetation types within protected areas. According to IUCN criteria,

five of eight assessed vegetation types were classified as threatened by 2050. Potential climate

change refugia, areas with the highest potential for safeguarding threatened vegetation types, were

also identified. This study provides an essential first step to assessing vegetation type vulnerability

based on predictions covering 46% of Arctic landscapes. The co-development of new protective

measures by policymakers and indigenous peoples at a pan-Arctic scale requires more robust and

spatially complete vegetation prediction, as increasing pressures from resource exploration and

infrastructure development threaten the sustainable development of the rapidly thawing and greening

Arctic.

Keywords: global change, arctic conservation, tundra vegetation, CAVM, vegetation shifts, climate

change refugia

[1] Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstrasse 190, 8057 Zürich,
Switzerland

[2] Department of Environmental Systems Science, Institute of Terrestrial Ecosystems, ETH Zürich, Universitätstrasse 16, 8092
Zürich, Switzerland

[3] Swiss Federal Research Institute WSL, Zürcherstrasse 111, 8903 Birmensdorf, Switzerland

[4] Department of Biology, McGill University, Montréal, QC, Canada

*Correspondence: merinreji.chacko@usys.ethz.ch, gabriela.schaepman@ieu.uzh.ch

1.   Introduction

The Arctic experiences climate warming at twice the global mean (Collins et al., 2013; Walsh et al.,

2011), leading to observed shifts in the distribution and composition of Arctic tundra vegetation

(Serreze et al., 2000) that are projected to intensify in the future (Pearson et al., 2013a). These

changes may threaten not only endemic plant species but entire vegetation types and the associated

ecosystem functions. Vegetation types provide habitats for sessile and migratory animal species

(Wheeler et al., 2018), support livelihoods of Northern peoples (CAFF, 2013) and take part in various

feedbacks involving climate (Juszak et al., 2014; Loranty et al., 2011; Myers-Smith et al., 2011;

Swann et al., 2010), permafrost soils (Bonfils et al., 2012; Juszak et al., 2014; Myers-Smith et al.,

2011), lakes, rivers, and the ocean through water, energy and carbon fluxes (Lunt et al., 2004; Woo &

Young, 2006).

Under the intermediate and high carbon emission scenarios (RCP 4.5 and 8.5), a recent long-term

modelling study predicted that the current tundra area in Siberia will be reduced to 5.7% by 2500

(Kruse & Herzschuh, 2022). The southern border to the Arctic tundra is the treeline, which has been

both observed and projected to advance northwards, decreasing the total extent of the tundra while

simultaneously displacing southern Arctic vegetation types (Bjorkman et al., 2020; Chapin & Starfield,

1997; Lloyd et al., 2002; Rundqvist et al., 2011). Due to decreasing snow cover and phenological

changes, primary productivity and vascular plant biomass have been increasing, specifically that of

tall shrubs (Myers-Smith et al., 2011). This shrubification of the tundra has been widespread, resulting

in a reduction in the abundance of less productive lichen- and moss-dominated vegetation types

(Bjorkman et al., 2020; Blok et al., 2010; Elmendorf et al., 2012; Loranty & Goetz, 2012; Myers-Smith

et al., 2011; Sturm et al., 2001). Widespread greening of the Arctic due to climate warming has long

been identified (Myers-Smith et al., 2020; Myneni et al., 1997). However, spectral browning and

increased heterogeneity of Arctic greening at the circumpolar scale have also been observed in

recent years (Bjorkman et al., 2020; Phoenix & Bjerke, 2016), further complicating our understanding

of changes in the highly heterogenous vegetation of the Arctic tundra.

Arctic vegetation is characterised by small vascular plants, bryophytes and lichens forming distinct

plant communities. The zonal vegetation can be classified into five physiognomic classes (barrens,

graminoid-dominated tundras, prostrate-shrub-dominated tundras, erect-shrub-dominated tundras,

and wetlands), further divided into fifteen vegetation types based on the dominant plant functional

types (Walker et al., 2005). The plant functional types are separated by plant traits such as growth

form (e.g. graminoids, shrubs), size (e.g. dwarf or low shrubs), taxonomical status (e.g. sedges,

rushes, grasses), and stature of woody shrubs. The resulting units include vegetation types such as

cryptogram-dominated barren, graminoid-dominated, prostrate dwarf-shrub, low-shrub tundra, and

sedge-dominated wetlands. They served as the fundamental units of the Circumpolar Arctic

Vegetation Map (2003 CAVM) (CAVM Team, 2003), a project within the Conservation of Arctic Flora

and Fauna (CAFF) group of the Arctic Council (CFG, 2022). Arctic vegetation types, as classified by

the CAVM approach, relate to their ecological function in the Arctic system, using a standardised

approach across the entire Arctic tundra biome. Hence, the CAVM approach has been widely applied

to study very diverse processes relying on plant functional types, including biogeochemical fluxes

(e.g. Virkkala et al., 2022; Zona et al., 2022), land surface energy flux (e.g. Oehri et al., 2022; Yu et

al., 2022), land surface modelling improvements (Sulman et al., 2021), Alaskan tundra wildfire activity

(Masrur et al., 2022) and studies investigating impacts of increasing shrub abundance on migratory

songbirds (e.g. Boelman et al., 2015). However, a modelling study forecasted that 48–84% of Arctic

vegetation types will have shifted by 2050 due to the effects of climate change (Pearson et al.,

2013a).

Novel pressures on Arctic vegetation beyond those posed by warming are arising in the Arctic due to

increasing anthropogenic presence. Historically, human land use and modification have been

relatively low or non-existent in the Arctic biome. The human footprint map classifies most of the

Arctic as under low pressure, and the human modification index demonstrates that the tundra remains

thus far one of the last true terrestrial wild places on Earth (Kennedy et al., 2019; Venter et al., 2016).

However, human interest in Arctic commodities such as oil and gas is increasing due to rising global

energy needs, and with it also the extent of disturbances that terrestrial ecosystems experience due

to exploration and infrastructure development (CAFF, 2013; Kumpula et al., 2011). The effects of

human disturbances and landscape processes such as climate and biota shifts on the distribution and

composition of vegetation over time must be explicitly included in effective conservation efforts. As

vegetation in the Arctic is more fragile and requires longer times to recover from perturbation in

comparison to southern vegetation (Kumpula et al., 2011), it is both an ecological, political and

economic imperative to have plans in place to mitigate and minimise these disturbances in order to

provide a path towards the sustainable development of the Arctic.

As of 2016, 20.2% of the terrestrial Arctic area is protected to some degree (CAFF & PAME, 2017).

Although conservation efforts in the Arctic are well-developed, their focus is generally at the species

level rather than the scale of ecosystems (Chapin et al., 2015). In other biomes, the impact of climate

change-induced biota shifts on conservation efforts has been recognised, and adaptive conservation

strategies have been developed, though not widely implemented (Heller & Zavaleta, 2009). A recent

systematic study identified areas with high potential for the persistence of multiple biodiversity

elements under climate change in North America, and demonstrated that at the biome-scale, ~80% of

areas within the top quintile of future conservation importance lacked formal protection, though this

study excluded the High Arctic due to lacking data (Stralberg et al., 2020). Additionally, conservation

needs in the North are influenced by rapid change at global scales and can no longer be addressed

solely by local actions (Chapin et al., 2015). Indeed, local studies of Arctic plant communities do not

always mirror observational trends over more extended temporal periods, which tend to be

heterogeneous and complex (Bjorkman et al., 2020; Myers-Smith et al., 2020). Conservation actions

which operate at various scales of space, time and biological organisation may be required to

effectively prevent the loss of potentially vulnerable ecosystems and their functions.

Refugia have become a focus of interest in conservation, as they have been demonstrated to enable

the persistence of biodiversity over longer temporal scales and changing climates while retaining

ecosystem and habitat functions (Morelli et al., 2016; Tzedakis et al., 2002). Vegetation refugia can

be defined as areas where existing vegetation will remain within current suitable climate conditions

(Thorne et al., 2020). In this study, we consider refugia as areas where threatened vegetation types

already occur, where the climatic envelope is predicted to remain suitable, and where competing

vegetation types might still be absent due to dispersal limitations. Refugia may serve as a means for

giving plant communities the time needed to allow for local adaptation to new environmental states by

providing habitats where the effects of climate change are least felt in the short term (Morelli et al.,

2020). The identification and protection of refugia may facilitate the persistence of retreating

vegetation types under projected anthropogenic climate change (Noss, 2001; Taberlet & Cheddadi,

2002).

Presently, protected areas in the Arctic have not been established with climate change-induced

vegetation shifts in mind (CAFF, 2013). Consequently, there is no consensus on the current state of

vegetation vulnerability in the Arctic. Additionally, as Arctic vegetation shifts, today's protected areas

may no longer protect the same vegetation to the same extent in the future. Vegetation changes lead

to trophic cascades, which alter the fundamental structures and functions of ecosystems (Wookey et

al., 2009). Vegetation distribution projections for the year 2050 (Pearson et al., 2013b) provide the

opportunity to locate vegetation refugia in the Arctic. Therefore, the conservation of potential

vegetation refugia could, in effect, protect species at higher trophic levels. However, a comprehensive

overview of vegetation refugia and their protected status in the Arctic has been unavailable to date.

This study aims to establish an overview of vegetation type abundance in Arctic protected areas and

assess their vulnerability. For this purpose, we utilised the 2003 CAVM as the baseline abundance of

Arctic vegetation types within and outside of current protected areas. We then determined the

distribution of Arctic vegetation types within protected areas using previously existing 2050 vegetation

scenarios (Pearson et al., 2013b). Furthermore, we assessed the risk of collapse of the vegetation

types following the IUCN Red List of Ecosystems criteria for the baseline and the future (Bland et al.,

2015). Lastly, to inform conservation efforts, we located potential refugia for the vegetation types that

had been identified as threatened.

2.   Materials and Methods

2.1.   Baseline (2003) status of vegetation type abundance in protected areas

We defined the network of protected areas in the Arctic according to their extent in the Map of Arctic

Protected Areas (MAPA) (CAFF & PAME, 2017). The protected areas were treated as one network

regardless of the six management categories within the map, as our focus was on the collective pan-

Arctic protection status. We utilised the 2003 CAVM to determine the baseline distribution of Arctic

vegetation types within protected areas, which included fifteen distinct vegetation types (Appendix A)

as well as the non-vegetative glacier class (CAVM Team, 2003; Walker et al., 2005). We intersected

the 2003 CAVM with MAPA and computed a zonal histogram. This resulted in the pixel count of the

protected area for each vegetation type in order to obtain the total and absolute areas of the

vegetation types present within protected areas.

2.2.   Future (2050) status of vegetation type abundance in protected areas

The bases for all analyses concerning future vegetation type distribution were the 36 Maps of Future

Arctic Vegetation Distribution (MFAVD), which predicted climate change-induced vegetation shifts for

2050 (Pearson et al., 2013a). Two machine learning methods for ecological niche modelling, three

climate models, two emissions scenarios, and three tree dispersal scenarios were applied to create

36 predictions (Pearson et al., 2013a). The MFAVDs are the result of all possible combinations of the

parameters named above. Areas covered by glaciers, barren lands and wetlands were excluded from

the predictions, but tree cover was incorporated to illustrate the northward treeline shift of the taiga.

We intersected all 36 MFAVDs with MAPA to obtain the complete range of possible outcomes. We

selected an MFAVD that we deemed to be most realistic (hereafter 'realistic model'): random forest

machine learning because it was the one with slightly higher accuracy (Pearson et al., 2013a); the

intermediate of the three global climate models (i.e. CSIRO); the higher of the two emissions

scenarios, A2a (IPCC, 2000); and the intermediate tree dispersal of 20 km, as it represents the

highest rate at which trees have been observed to disperse northward in Alaska and Canada

(Lescop-Sinclair & Payette, 1995; Lloyd et al., 2002).

We analysed the eight vegetation types available in the MFAVDs out of the fifteen originally present in

the 2003 CAVM, as barren (four classes) and wetlands (three classes) had been excluded from the

predictions (Pearson et al., 2013a). To assess the future distribution and protection status of Arctic

vegetation, we converted MAPA from a polygon to a 4.5-km resolution raster to match the spatial

resolution of the MFAVDs and attributed the protection status to every pixel. Finally, we summed the

pixels and obtained the number of pixels per total as well as per protected area for each vegetation

type.

2.3.   Vulnerability of Arctic vegetation types following IUCN criteria

We applied the "IUCN Red List of Ecosystems Categories and Criteria" to quantify the vulnerability of

Arctic vegetation types (Bland et al., 2015). The IUCN provides guidelines to assess the risk of

ecosystem collapse, designed to be applicable to various ecosystems. Here, we defined ecosystems

as the CAVM vegetation types, representing over 400 plant communities classified into broader

functional groups (Walker et al., 2005). The IUCN assigns one of three graded categories to

threatened ecosystems: "vulnerable", "endangered", and "critically endangered". We did not

differentiate here between the two graded categories for unthreatened ecosystems ("not threatened"

and "least concern") and instead simply summarised them as "not threatened".

Five criteria are used to determine the IUCN risk category of the ecosystem: "reduction in geographic

distribution", "restricted in geographic distribution", "environmental degradation", "disruption of biotic

processes or interactions", and "quantitative analysis that estimates the probability of ecosystem

collapse" (Bland et al., 2015). It is recommended that as many criteria as possible are assessed, as

the ultimate classification of risk is determined as the highest of the five criteria (Bland et al., 2015). At

the pan-Arctic scale, data were only available for "restricted distribution" and "decline of distribution";

therefore, this study could only assess these two criteria. Hence, our results represent the minimum

risk status that each vegetation type can be assigned. Analyses of the remaining three criteria could

potentially result in a higher risk category.

To determine the "restricted distribution" criterion classification, the number of 10×10 km grid cells

occupied by the ecosystem must be calculated. An ecosystem is classified as "vulnerable",

"endangered", or "critically endangered" if it occupies at most 50, 20 or 2 grid cells, respectively

(Dudley, 2008). After the total extent of each vegetation type within the 2003 CAVM was calculated in

square kilometres, the extents were divided by 100 to determine the number of 10×10 grid cells

occupied by the vegetation type across the pan-Arctic extent. The "decline of distribution" criterion is

determined by the predicted relative reduction in the distribution of an ecosystem over fifty years. An

ecosystem is classified as "vulnerable", "endangered", or "critically endangered" if the reduction in its

spatial extent is at least 30%, 50% or 80%, respectively (Dudley, 2008). In this analysis, the relative

changes in area were calculated according to the changes between the 2003 CAVM (CAVM Team,

2003) and the 2050 MFAVDs (Pearson et al., 2013a); therefore, the classification is based on the

predicted relative reduction over 47 rather than 50 years.

2.4.   Identification of potential refugia for threatened vegetation types

For the threatened vegetation types, we identified areas which may serve as refugia, i.e. where a

vegetation type is present in the baseline and is predicted to persist at least until 2050. We overlaid

the realistic model with the 2003 CAVM and selected the areas where the pixels remained unchanged

to create a map of areas with persistent vegetation types. We did not consider areas where these

vegetation types had been predicted to shift to and only considered the areas where the vegetation

remained the same, as a) the predictions of future vegetation distribution contain a measure of

uncertainty, b) there is evidence of lag effects in species-scale responses to climate change (Stewart

et al., 2016), and c) considerable variability in their ability to track new climatic niches exists (La Sorte

& Jetz, 2012).

Spatial data were analysed using ArcMap 10.5.1 (ESRI, 2017) and RStudio 1.2.5033 (R Development

Core Team, 2017; Team RStudio, 2020). Maps were converted into the Lambert Azimuthal Equal

Area Polar Projection to preserve area, a crucial feature for this analysis.

3.   Results

3.1.   Baseline (2003) status of vegetation type abundance in protected areas

The terrestrial pan-Arctic region covers over 7.1 million km^2, 20.2% of which is protected. Of the

vegetated areas (4.7 million km^2), 21% — or approximately 977,000 km^2 — fall within protected

areas. Additionally, a significant portion of the protected areas encompasses glaciated areas (777,000

km^2) but could not be considered within the scope of this study due to a lack of predictions.

Our analysis demonstrates that the fifteen vegetation types have substantially different absolute and

relative spatial abundances in the pan-Arctic tundra (Figure 1). The most abundant vegetation type

(S1, erect dwarf-shrub tundra) covers over six times more area than the least abundant vegetation

type (W1, sedge/grass, moss wetland), 626,000 km^2, and 102,000 km^2, respectively. The abundances

of the vegetation types within protected areas generally mirror their abundance at the pan-Arctic

scale. For example, 11.4% of Arctic vegetation is composed of the G3 vegetation type (nontussock

sedge, dwarf-shrub, moss tundra), which accordingly covers 11.3% of protected areas. Notable

exceptions do occur, such as the wetland types. In comparison to their total extents (W1:

sedge/grass, moss wetland, 2.0%; W2: sedge, moss, dwarf-shrub wetland, 2.7%; W3: sedge, moss,

low-shrub wetland, 3.7%), they are overrepresented within protected areas (W1, 3.0%; W2, 4.8%;

W3, 6.7%). Contrastingly, though 7.5% of Arctic vegetation is composed of the vegetation type B2

(cryptogam barren complex, bedrock), it spans only 2.5% of the protected area. In terms of absolute

extents, the most abundant types at pan-Arctic scales are overrepresented in protected areas, while

the least abundant types are underrepresented.

3.2.   Predicted vegetation type abundance in protected areas by 2050

Compared to the baseline abundances of vegetation types, the assessment of predicted abundances

across protected areas by 2050 includes only eight vegetation types (see section 2.2). Predictions for

the vegetation types within the barren (25.6% of non-glaciated terrestrial protected areas, 260,000

km^2) and wetland (14.5% of non-glaciated terrestrial protected areas,148,000 km^2) classes were

unavailable. Additionally, as the nearly 777,000 km^2 of glaciated protected areas were lacking in

predictions, they were excluded from the study. The results differed considerably between the 36

MFAVDs: between 48,000 (0.1%) and 1,826,000 km^2 (39.9%) of the total area covered by tundra

vegetation in the 2003 CAVM is predicted to be replaced by taiga (Pearson et al., 2013a). Hence,

considerable differences also resulted in the predictions of single vegetation types as a function of

MFAVDs. We restrict the presentation of results to our selected realistic model and present the results

of the other MFAVDs as an envelope around this model result. Of the eight vegetation types, five are

predicted to decline in their total area according to the realistic model, accompanied by a mirrored

decline in their abundance within the protected area network (Figure 2). In contrast, two southern

vegetation types (S2: low shrub tundra and G3: nontussock sedge, dwarf-shrub moss tundra) are

predicted to gain in area, both within and outside of protected areas. Vegetation type G4 (tussock

sedge, dwarf shrub, moss tundra) showed a decrease in total area, yet its abundance within the

protected area network increased.

3.3.   Vulnerability of Arctic vegetation types following IUCN criteria

To determine the risk of collapse for the eight vegetation types, we analysed two spatial criteria:

"restriction of distribution" and "decline of distribution"; the final assigned category of risk was

determined as the higher of these two. The "restriction of distribution" criterion diagnosed none of the

eight vegetation types as threatened because each vegetation type has an extent of at least 5,000

km^2. The "decline in distribution" criterion assigned different results (Figure 3). Depending on the

MFAVD, the specific risk status assigned to each type varied. Under the realistic model, one

vegetation type (P1: prostrate dwarf-shrub, herb tundra) was classified as critically endangered. Four

vegetation types were classified as endangered (G1: rush/grass forb, cryptogam tundra; G2:

graminoid, prostrate dwarf-shrub, forb tundra; P2: prostrate/hemiprostrate dwarf-shrub tundra; S1:

erect dwarf-shrub tundra), and three as not threatened (G3: nontussock sedge, dwarf-shrub, moss

tundra; G4: tussock sedge, dwarf-shrub, moss tundra; S2: low shrub tundra). The threatened

vegetation types (i.e., those classified at least as endangered) are generally the more northern types

with sparse, low-growing, non-vascular vegetation.

3.4.   Potential refugia for threatened vegetation types

For the five vegetation types (G1, G2, P1, P2, S1) classified as threatened under the realistic model,

we determined refugia, regions where these vegetation types are predicted to persist until 2050

(Figure 4, Appendix C). The total area of refugia ranges from < 9,000 km^2 (P2) to > 100,000 km^2 (S1),

accounting for 2.1% and 8.7% of currently protected areas. The refugia are scattered over Canada,

Greenland, Norway, Russia, and the USA (Figures 4a-d).

4. Discussion

We evaluated the capability of the current Arctic network of protected areas for the conservation of

Arctic vegetation types. Our results demonstrate that all Arctic vegetation types are found within the

protected areas, and their relative abundance significantly varied but generally followed the same

pattern as seen at the pan-Arctic scale. Noticeably, wetlands tend to be protected to an above-

average extent; this is likely due to conservation efforts focusing on Wetlands of International

Importance (Ramsar sites) related to their protection as breeding areas for migrating birds (CAFF &

PAME, 2017).

The impacts of climate change may demand the adaptation of protected areas (Hannah et al., 2002;

Heller & Zavaleta, 2009), as they may no longer harbour the same vegetation types. Thus, we

analysed the protection and threat status of the vegetation type abundances predicted for the year

2050. The two most productive vegetation types (G3 and S2) occupying the lower southern

bioclimatic zones were demonstrably increasing in extent. The five vegetation types which were

predicted to decline drastically currently occupy the three more northern bioclimatic zones (Pearson et

al., 2013b) and are characterised by sparse and low-growing vegetation (Raynolds et al., 2019). Our

results demonstrate that these northern vegetation types also decrease in their representation within

protected areas. We attribute this to two reasons. First, protected areas are generally situated in more

southern regions rather than the northernmost edges of the Arctic, where these northern vegetation

types can find sanctuary. This interpretation is validated by the fact that the refugia we located are

clustered around the northernmost edges of the terrestrial Arctic. In our realistic model, we observe

them being especially abundant in the Canadian Archipelago, which contains most of the terrestrial

High Arctic. The climate will stay unsuitable for southern vegetation types in these regions at least

until 2050. Secondly, Arctic protected areas were not established with the conservation of vegetation

types in mind (CAFF, 2013); therefore, it stands to reason that vegetation types of conservation

importance have not been recognised and hence prioritised.

4.1.   Implications for flora, fauna and indigenous people if vegetation types are left unprotected

Vegetation shifts have been observed to have outcomes for species and communities across trophic

levels (Myers-Smith et al., 2011; Wheeler et al., 2018). Threatened vegetation types may lead to the

increased vulnerability of endemic flora and fauna, especially those that occur only in the High Arctic,

which are most at risk. For example, there is evidence that vegetation exerts bottom-up control on

caribou populations, as increasing shrubification decreases pasture quality (Fauchald et al., 2017).

This decline is exacerbated by the loss of lichen-rich vegetation types, which negatively affect caribou

populations due to a loss of winter foraging options (Cornelissen et al., 2001; Joly et al., 2009) and

energy through increased methane emission if lichens are reduced in their diet (Hansen et al., 2018).

Indigenous knowledge also shows that caribou populations have already adopted more northern

migration patterns (Ksenofontov et al., 2019). This can, in turn, have negative consequences for

indigenous communities that depend on caribou for food and as economic resources (Joly et al.,

2009; Nuttall et al., 2005). Without focused conservation efforts on maintaining these vegetation

types, the species depending on them will also become threatened with extinction.

4.2.   Implications of land use change on threatened vegetation types if left unprotected

Climate warming and technological advances have opened up the Arctic as a new frontier in

economic development. Increasing interest in commodities such as oil, gas, and mineral resources

will naturally lead to increased infrastructure requirements in the Arctic, which may, in turn, intensify

disturbances in Arctic ecosystems. Surface disturbances such as road networks and settlements have

led to permafrost degradation through soil warming (Forbes et al., 2001). Off-road vehicles like tundra

tractors can destroy endemic vegetation and leave the tracks visible for decades as vegetation

struggles to recover (Forbes et al., 2001). A study of the Alaska North Slope demonstrated that 34%

of the area was affected by oil development by 2010 (Raynolds et al., 2014). Further exploration has

already been proposed, such as a 3-D seismic survey covering 63,000 km of trails within the Arctic

National Wildlife Refugia in Alaska (SAExploration, 2018). A recent study demonstrated that this could

result in mid-to-high level impacts on 122 km^2 of the area in question, leading to increased

thermokarst formation and erosion, and negatively impacting moist vegetation types (Raynolds et al.,

2020). Additionally, vehicle tires can significantly increase the dispersal distance of southern invaders,

such as *Salix lanata,* which has been found to occur in many roadside areas, replacing lichen and

moss cover (Raynolds et al., 2014). Local warming due to the settlements also generates pockets of

suitable habitats in latitudes where southern vegetation could not otherwise persist (Forbes et al.,

2001).

Permafrost thaw is critically threatening Arctic infrastructures (Hjort et al., 2018), and their collapse

can lead to dramatic consequences for terrestrial ecosystems and indigenous livelihoods. As the

MFAVDs did not account for human land use modification, this study severely understates their

effects on vegetation shifts. At the pan-Arctic scale, a combination of economic, geopolitical, climatic,

infrastructure and ecological factors lead to uncertainty in the future spatial distribution of

development pressures and their degree of impact (Scibilia et al., 2015). Therefore, it is essential to

predefine conservation areas for the currently unprotected and threatened vegetation types before

widespread development begins in the increasingly accessible Arctic. Complementing protected

areas, biodiversity conservation schemes developed for other regions of the world, such as around

mining sites, need to be adapted to Arctic conditions to prevent loss of vegetation where industrial

development outside of protected areas is allowed.

4.3.   Implications of using climate change refugia for land conservation and management

Globally protected areas are 10.6% richer in species diversity than non-protected areas (Gray et al.,

2016). Nevertheless, conserving biodiversity using protected areas poses to be difficult under climate

change. A study assessing the protected areas of Canada found that climate change will lead to over

40% of the protected areas experiencing a change in biome type, with the total extent of the Canadian

tundra standing to decline by 38–79% (Lemieux & Scott, 2005). In order to most efficiently use limited

resources and protect threatened vegetation, efforts in in situ management must be focused on areas

where the vegetation is likely to persist under global change, such as climate change refugia for

vegetation. There is scepticism over the use of refugia as areas where vegetation may retreat until

conditions in the surrounding environment become more favourable. Vegetation shifts due to climate

warming are projected to continue and intensify; therefore, the time periods required until favourable

conditions return may be longer than the existence of these refugia. However, they may serve a

powerful purpose by buying time for the climate adaptation of vulnerable species and ecological

communities (Morelli et al., 2020). Adjusting Arctic protection efforts in mitigating the consequences of

global change with a focus on relatively transient climate change refugia may aid in a long-term

transformation of these threatened vegetation types into novel community assemblages (i.e.

ecological replacements) that have adapted to these environmental changes while performing the

same ecological and habitat functions (Morelli et al., 2020). This process has been documented in the

late Quaternary (Jackson & Overpeck, 2000).

4.4.   Limitations and urgent need for comprehensive predictions of Arctic vegetation type

357   distribution

The aim of this study is to construct a first assessment of the protected status of Arctic vegetation

types in light of climate change based on existing vegetation predictions. These predictions (Pearson

et al., 2013b) were based on the 2003 CAVM, a vector map with polygons of 14 km minimum

diameter (Walker et al., 2005); thus, for the sake of consistency, it was utilised by this study instead of

the more recent raster version (2019 CAVM) that is resolved at 1 km (Raynolds et al., 2019).

Nevertheless, these disparities did not affect the results regarding the baseline risk status of the

vegetation types; the general pattern of abundant vegetation types being better represented within

protected areas holds, regardless of the map used (Appendix B).

Increasing spatial resolution in future predictions could reveal pockets of vegetation occurring outside

of the main macroclimatic niche of a vegetation type. Such pockets could accelerate the spread of

vegetation types into surrounding areas should the climate become more favourable. Potentially,

southern vegetation types could then disperse northward at faster rates than modelled (Pearson et

al., 2013a), increasing the displacement of northern vegetation types and potentially our risk

classifications. This displacement may be balanced by potential northern micropockets, which would

increase the total extent of refugia. Nevertheless, these disparities did not affect the results regarding

the present risk status of the vegetation types. Though the "decline in distribution" criterion could not

be tested as MFAVDs do not exist for the 2019 CAVM, none of the vegetation types of the 2019

CAVM showed a restricted distribution and were classified as unthreatened under the IUCN

"restriction of distribution" criterion.

Due to limited data availability in the Arctic, we could only analyse two of the five IUCN criteria.

Therefore, the assigned risk category only represents the minimal risk and could potentially be higher,

both in the present and in the future. Though it is recommended that as many criteria as possible are

assessed, the IUCN Red List of Ecosystems was created with the purpose of flexibility in the use of

data containing variations in quality and coverage (Bland et al., 2015). Therefore, as data availability

increases, this risk assessment still provides valuable information as a starting point for future

assessments.

The MFAVDs only rendered predictions for 46% of the pan-Arctic area. Specifically, they excluded

glaciers (including nunataks), barren lands, and wetlands (Pearson et al., 2013a). Subsequently, we

were unable to assess the future distribution of wetland and barren vegetation types, classify their risk

of collapse, and locate potential refugia. Therefore, our results do not consider the potential of the

assessed vegetation types to disperse into areas currently covered by these unassessed vegetation

types and vice versa.

Glaciers currently cover nearly one-third of the Arctic (Walker et al., 2005), but extensive climate

change-induced glacier melt is predicted (Alley et al., 2010; Hinzman et al., 2005). Vegetation

succession following retreating glaciers in the Arctic follows the same patterns as elsewhere;

however, these processes occur at larger temporal scales and may be less relevant for the next thirty

years (Hodkinson et al., 2003; Raynolds & Walker, 2009). Vegetation growth response to warming is

slower in the High Arctic than in the Low Arctic due to lower growing season temperatures, seasonal

length and nutrient availability (Walker et al., 2006). In Svalbard, for instance, vascular plant

establishment was highly limited for the first century after glacial retreat (Hodkinson et al., 2003).

Additionally, vegetation can take up to 500 years to reach equilibrium after glacial retraction in

Greenland (Lunt et al., 2004). Therefore, the inclusion of glaciers would have likely had limited

impacts on the results for threatened vegetation types shown here for 2050. Nevertheless, they would

be crucial for imminent studies which investigate the future of the terrestrial Arctic at longer

timescales.

One-quarter of the terrestrial Arctic consists of barren lands (Walker et al., 2005), which were

excluded in this analysis. Therefore, the northern vegetation types may be less threatened than

indicated by this study because they may expand into currently barren areas. Including barren lands

in future studies would likely lead to predictions of northern vegetation types being slightly more

abundant and increasing over centennial periods. However, due to the slow response in the

development of soils and vegetation, the time scales at which this northern expansion into barren

lands occurs may be longer than the 47-year span of our study, and this expansion may be further

limited as barren lands lack in soil sufficient enough to support vegetation (Pearson et al., 2013a).

Wetlands constitute 7% of the Arctic vegetated area (Walker et al., 2005). Predictions for future

wetland distribution are particularly difficult due to the complexity of the hydrological processes

regulating them (Walvoord & Kurylyk, 2016). Permafrost thaw, increasing precipitation, ice melt, and

evaporation due to warming influence the abundance and distribution of surface water, indirectly

affecting wetlands (White et al., 2007; Woo & Young, 2006). In the southern areas with discontinuous

permafrost, ponds are generally in decline as permafrost thaw increases drainage (Riordan et al.,

2006; Smith, 2005; Yoshikawa & Hinzman, 2003). Conversely, the thawing of continuous permafrost

in the High Arctic has increased abundance of thaw ponds (Smith, 2005). In the Canadian Arctic, this

trend of thermokarst pond formation has been shown to occur at rapid paces (Farquharson et al.,

2019). This pattern of wetland development means that generally, southern vegetation types would

have more potential areas of expansion, whereas northern areas may become further displaced by

expanding wetland. We hypothesise that the inclusion of wetlands in this study could consequently

have a reinforcing effect on the observed patterns of threatened vegetation types. While wetlands are

crucial because of the abundance of migratory birds dependent on them, they make up a relatively

small portion of the vegetated pan-Arctic and are generally better protected than other vegetation

types. However, this may change significantly in the future due to climate change. Our study

highlights the need for further focus on these ill-understood vegetation types in longer-term

predictions.

The future vegetation shifts (Pearson et al., 2013a) were predicted under various assumptions about

climate change and tree dispersal rate scenarios, which, though necessary, carry over to this study.

Uncertainties in future vegetation distributions are inherent to predictions based on climate models

with many uncertain assumptions, where one model is no more or less valid than the other (IPCC,

2000). Our selection of the realistic future model with regard to climate and emission scenario is,

therefore, subjective. There is a large spread of the vegetation type abundances across the models,

making the results, especially for vegetation types G1, G2, G4 and S1, more uncertain (Figure 2 and

Appendix D). However, the general trend of increase in southern vegetation types and decrease in

northern vegetation types still holds within and outside of protected areas, as demonstrated for most

results by the minimum and maximum scenario envelope.

While land surface schemes for earth system models and vegetation development models are being

adapted to Arctic tundra conditions, parameterisations at the Arctic tundra vegetation type level,

including all vegetation types, need to be implemented, and biotic and abiotic interactions integrated

with holistic modelling approaches, at pan-Arctic scale. Furthermore, the inclusion of human land use

scenarios (e.g. reindeer herding densities, potential mining sites, road development) will be necessary

for future modelling efforts to allow for informed policies and decisions.

5.   Conclusions

This study identifies baseline and future abundances of Arctic vegetation types within protected areas

as well as potential climate change refugia for threatened vegetation types using the 2003

Circumpolar Arctic Vegetation Map and existing predictions of future Arctic vegetation distribution.

Though uncertainties exist within the maps provided here, the general trends seen in this study are

valuable in guiding future conservation efforts. Further studies on the development of Arctic

vegetation types which include the whole Arctic — especially wetlands, barrens and ice-covered

areas — are urgently needed. They could provide complete and precise findings to improve and

adapt conservation efforts to climate change-induced biota shifts. Moreover, we recommend

independent validation of these hypothetical predictions to better incorporate climate change refugia

into designated protected areas. This could potentially be achieved through recent validation methods

which compare predictions of refugia to local measures of species richness, endemic species

persistence, genetic diversity, plant functional traits and/or demographic variables (Barrows et al.,

2020).

This study aims to shift the focus of pan-Arctic protection efforts towards higher scales of biological

organisation, from species to communities and ecosystems, or in this case, vegetation types.

Endemic tundra vegetation types face the threat of ecosystem collapse due to global change. The

establishment of refugia for the vegetation types identified here could protect them as refugia have

done over the history of life on earth (Alley et al., 2010; Jackson & Overpeck, 2000). The protection of

vegetation refugia would additionally have positive effects on Arctic fauna and climate regulation

through bottom-up biotic and abiotic interactions. However, the integration of vegetation types for

climate change-adapted conservation in the Arctic requires urgent collaboration between

policymakers and indigenous peoples as the area becomes increasingly under pressure from

exploration and rapid infrastructure development. As experienced in the recent decade, Arctic change

is rapid. Extreme events, including extreme winter precipitation and summer drought, are already

affecting Arctic ecosystems through major disturbance events such as extensive flooding and fires,

adding yet another dimension of abrupt change to mitigate in the future. Without a plan already in

place for the protection of these critically important Arctic landscapes, we cannot enable the

sustainability of economic and structural development in what is increasingly no longer one of the

world's last truly wild places.

6. Acknowledgements

This work was supported by the University Research Priority Programme Global Change and Biodi-

versity of the University of Zurich.

7. Conflict of Interest

The authors declare that there is no conflict of interest.

References

Alley, R. B., Andrews, J. T., Brigham-Grette, J., Clarke, G. K. C., Cuffey, K. M., Fitzpatrick, J. J.,

Funder, S., Marshall, S. J., Miller, G. H., Mitrovica, J. X., Muhs, D. R., Otto-Bliesner, B. L.,

Polyak, L., & White, J. W. C. (2010). History of the Greenland Ice Sheet: paleoclimatic insights.

Quaternary Science Reviews. https://doi.org/10.1016/j.quascirev.2010.02.007

Barrows, C. W., Ramirez, A. R., Sweet, L. C., Morelli, T. L., Millar, C. I., Frakes, N., Rodgers, J., &

Mahalovich, M. F. (2020). Validating climate-change refugia: empirical bottom-up approaches to

support management actions. *Frontiers in Ecology and the Environment*, *18*(5), 298–306.

https://doi.org/10.1002/fee.2205

Bjorkman, A. D., García Criado, M., Myers-Smith, I. H., Ravolainen, V., Jónsdóttir, I. S., Westergaard,

K. B., Lawler, J. P., Aronsson, M., Bennett, B., Gardfjell, H., Heiðmarsson, S., Stewart, L., &

Normand, S. (2020). Status and trends in Arctic vegetation: Evidence from experimental

warming and long-term monitoring. *Ambio*. https://doi.org/10.1007/s13280-019-01161-6

Bland, L. M., Keith, D. A., Miller, R. M., Murray, N. J., & Rodriguez, J. P. (2015). Guidelines for the

application of IUCN Red List of ecosystems categories and criteria. In *Guidelines for the*

application of IUCN Red List of ecosystems categories and criteria.

https://doi.org/10.2305/iucn.ch.2016.rle.1.en

Blok, D., Heijmans, M. M. P. D., Schaepman-Strub, G., Kononov, A. V., Maximov, T. C., & Berendse,

F. (2010). Shrub expansion may reduce summer permafrost thaw in Siberian tundra. *Global*

Change Biology. https://doi.org/10.1111/j.1365-2486.2009.02110.x

Boelman, N. T., Gough, L., Wingfield, J., Goetz, S., Asmus, A., Chmura, H. E., Krause, J. S., Perez, J.

H., Sweet, S. K., & Guay, K. C. (2015). Greater shrub dominance alters breeding habitat and

food resources for migratory songbirds in Alaskan arctic tundra. *Global Change Biology*.

https://doi.org/10.1111/gcb.12761

Bonfils, C. J. W., Phillips, T. J., Lawrence, D. M., Cameron-Smith, P., Riley, W. J., & Subin, Z. M.

(2012). On the influence of shrub height and expansion on northern high latitude climate.

Environmental Research Letters. https://doi.org/10.1088/1748-9326/7/1/015503

CAFF. (2013). Arctic Biodiversity Assessment. Status and trends in Arctic biodiversity. Conservation

of Arctic Flora and Fauna. In *Arctic Biodiversity Assessment*.

CAFF, & PAME. (2017). *Arctic Protected Areas: Indicator Report, 2017: Conservation of Arctic Flora*

and Fauna and Protection of the Arctic Marine Environment.

CAVM Team. (2003). Circumpolar Arctic Vegetation Map. (1:7,500,000 scale), Conservation of Arctic

Flora and Fauna (CAFF) Map No. 1. In *US Fish and Wildlife Service.(Anchorage, AK)*. ISBN: 0-

9767525-0-6, ISBN-13: 978-0-9767525-0-9.

CFG. (2022). *CAFF - About the Circumpolar Flora Group (CFG)*. https://www.caff.is/flora-cfg/about-

cfg

Chapin, F. S., Sommerkorn, M., Robards, M. D., & Hillmer-Pegram, K. (2015). Ecosystem

stewardship: A resilience framework for arctic conservation. *Global Environmental Change*.

https://doi.org/10.1016/j.gloenvcha.2015.07.003

Chapin, F. S., & Starfield, A. M. (1997). Time lags and novel ecosystems in response to transient

climatic change in arctic Alaska. *Climatic Change*. https://doi.org/10.1023/A:1005337705025

Collins, M., Knutti, R., Arblaster, J., Dufresne, J. L., Fichefet, T., Friedlingstein, P., Wehner, M., &

IPCC. (2013). Long-term climate change: Projections, commitments and irreversibility. In

Climate Change 2013: The Physical Science Basis. IPCC Working Group I Contribution to AR5.

https://doi.org/10.1017/CBO9781107415324.024

Cornelissen, J. H. C., Callaghan, T. V., Alatalo, J. M., Michelsen, A., Graglia, E., Hartley, A. E., Hik, D.

S., Hobbie, S. E., Press, M. C., Robinson, C. H., Henry, G. H. R., Shaver, G. R., Phoenix, G. K.,

Jones, D. G., Jonasson, S., Chapin, F. S., Molau, U., Neill, C., Lee, J. A., … Aerts, R. (2001).

Global change and arctic ecosystems: Is lichen decline a function of increases in vascular plant

biomass? *Journal of Ecology*. https://doi.org/10.1046/j.1365-2745.2001.00625.x

Dudley, N. (2008). Guidelines for applying protected area management categories. In *Guidelines for*

applying protected area management categories. https://doi.org/10.2305/iucn.ch.2008.paps.2.en

Elmendorf, S. C., Henry, G. H. R., Hollister, R. D., Björk, R. G., Boulanger-Lapointe, N., Cooper, E. J.,

Cornelissen, J. H. C., Day, T. A., Dorrepaal, E., Elumeeva, T. G., Gill, M., Gould, W. A., Harte,

J., Hik, D. S., Hofgaard, A., Johnson, D. R., Johnstone, J. F., Jónsdóttir, I. S., Jorgenson, J. C.,

… Wipf, S. (2012). Plot-scale evidence of tundra vegetation change and links to recent summer warming. *Nature Climate Change.* https://doi.org/10.1038/nclimate1465

ESRI. (2017). *ArcMap 10.5.1 [GIS Software]* (10.5.1). Environmental Systems Research Institute, Inc.

Farquharson, L. M., Romanovsky, V. E., Cable, W. L., Walker, D. A., Kokelj, S. V., & Nicolsky, D. (2019). Climate Change Drives Widespread and Rapid Thermokarst Development in Very Cold Permafrost in the Canadian High Arctic. *Geophysical Research Letters.* https://doi.org/10.1029/2019GL082187

Fauchald, P., Park, T., Tømmervik, H., Myneni, R., & Hausner, V. H. (2017). Arctic greening from warming promotes declines in caribou populations. *Science Advances.* https://doi.org/10.1126/sciadv.1601365

Forbes, B. C., Ebersole, J. J., & Strandberg, B. (2001). Anthropogenic disturbance and patch dynamics in circumpolar arctic ecosystems. In *Conservation Biology.* https://doi.org/10.1046/j.1523-1739.2001.015004954.x

Gray, C. L., Hill, S. L. L., Newbold, T., Hudson, L. N., Boïrger, L., Contu, S., Hoskins, A. J., Ferrier, S., Purvis, A., & Scharlemann, J. P. W. (2016). Local biodiversity is higher inside than outside terrestrial protected areas worldwide. *Nature Communications*, *7*(1), 1–7. https://doi.org/10.1038/ncomms12306

Hannah, L., Midgley, G. F., & Millar, D. (2002). Climate change-integrated conservation strategies. *Global Ecology and Biogeography.* https://doi.org/10.1046/j.1466-822X.2002.00306.x

Hansen, K. K., Sundset, M. A., Folkow, L. P., Nilsen, M., & Mathiesen, S. D. (2018). Methane emissions are lower from reindeer fed lichens compared to a concentrate feed. *Polar Research*, *37*(1). https://doi.org/10.1080/17518369.2018.1505396

Heller, N. E., & Zavaleta, E. S. (2009). Biodiversity management in the face of climate change: A review of 22 years of recommendations. In *Biological Conservation.* https://doi.org/10.1016/j.biocon.2008.10.006

Hinzman, L. D., Bettez, N. D., Bolton, W. R., Chapin, F. S., Dyurgerov, M. B., Fastie, C. L., Griffith, B., Hollister, R. D., Hope, A., Huntington, H. P., Jensen, A. M., Jia, G. J., Jorgenson, T., Kane, D.

L., Klein, D. R., Kofinas, G., Lynch, A. H., Lloyd, A. H., McGuire, A. D., … Yoshikawa, K. (2005).

Evidence and implications of recent climate change in Northern Alaska and other Arctic regions.

Climatic Change. https://doi.org/10.1007/s10584-005-5352-2

Hjort, J., Karjalainen, O., Aalto, J., Westermann, S., Romanovsky, V. E., Nelson, F. E., Etzelmüller,

B., & Luoto, M. (2018). Degrading permafrost puts Arctic infrastructure at risk by mid-century.

Nature Communications, *9*(1). https://doi.org/10.1038/s41467-018-07557-4

Hodkinson, I. D., Coulson, S. J., & Webb, N. R. (2003). Community assembly along proglacial

chronosequences in the high Arctic: Vegetation and soil development in north-west Svalbard.

Journal of Ecology. https://doi.org/10.1046/j.1365-2745.2003.00786.x

IPCC. (2000). *IPCC Special Report on Emissions Scenarios (SRES) of Working Group III*.

https://doi.org/citeulike-article-id:9904924

Jackson, S. T., & Overpeck, J. T. (2000). Responses of plant populations and communities to

environmental changes of the late Quaternary. *Paleobiology*.

https://doi.org/10.1017/s0094837300026932

Joly, K., Jandt, R. R., & Klein, D. R. (2009). Decrease of lichens in Arctic ecosystems: The role of

wildfire, caribou, reindeer, competition and climate in north-western Alaska. In *Polar Research*.

https://doi.org/10.1111/j.1751-8369.2009.00113.x

Juszak, I., Erb, A. M., Maximov, T. C., & Schaepman-Strub, G. (2014). Arctic shrub effects on NDVI,

summer albedo and soil shading. *Remote Sensing of Environment*.

https://doi.org/10.1016/j.rse.2014.07.021

Kennedy, C. M., Oakleaf, J. R., Theobald, D. M., Baruch-Mordo, S., & Kiesecker, J. (2019). Managing

the middle: A shift in conservation priorities based on the global human modification gradient.

Global Change Biology. https://doi.org/10.1111/gcb.14549

Kruse, S., & Herzschuh, U. (2022). Regional opportunities for tundra conservation in the next 1000

years. *ELife*, *11*. https://doi.org/10.7554/elife.75163

Ksenofontov, S., Backhaus, N., & Schaepman-Strub, G. (2019). 'There are new species': indigenous

knowledge of biodiversity change in Arctic Yakutia. *Polar Geography*, *42:1*, 34–57.

https://doi.org/https://doi.org/10.1080/1088937X.2018.1547326

Kumpula, T., Pajunen, A., Kaarlejärvi, E., Forbes, B. C., & Stammler, F. (2011). Land use and land

cover change in Arctic Russia: Ecological and social implications of industrial development.

Global Environmental Change. https://doi.org/10.1016/j.gloenvcha.2010.12.010

La Sorte, F. A., & Jetz, W. (2012). Tracking of climatic niche boundaries under recent climate change.

Journal of Animal Ecology. https://doi.org/10.1111/j.1365-2656.2012.01958.x

Lemieux, C. J., & Scott, D. J. (2005). Climate change, biodiversity conservation and protected area

planning in Canada. *Canadian Geographer*. https://doi.org/10.1111/j.0008-3658.2005.00103.x

Lescop-Sinclair, K., & Payette, S. (1995). Recent Advance of the Arctic Treeline Along the Eastern

Coast of Hudson Bay. *The Journal of Ecology*. https://doi.org/10.2307/2261175

Lloyd, A. H., Rupp, T. S., Fastie, C. L., & Starfield, A. M. (2002). Patterns and dynamics of treeline

advance on the Seward Peninsula, Alaska. *Journal of Geophysical Research D: Atmospheres*.

https://doi.org/10.1029/2001jd000852

Loranty, M. M., & Goetz, S. J. (2012). Shrub expansion and climate feedbacks in Arctic tundra. In

Environmental Research Letters. https://doi.org/10.1088/1748-9326/7/1/011005

Loranty, M. M., Goetz, S. J., & Beck, P. S. A. (2011). Tundra vegetation effects on pan-Arctic albedo.

Environmental Research Letters. https://doi.org/10.1088/1748-9326/6/2/029601

Lunt, D. J., de Noblet-Ducoudré, N., & Charbit, S. (2004). Effects of a melted greenland ice sheet on

climate, vegetation, and the cryosphere. *Climate Dynamics*. https://doi.org/10.1007/s00382-004-

0463-4

Masrur, A., Taylor, A., Harris, L., Barnes, J., & Petrov, A. (2022). Topography, Climate and Fire

History Regulate Wildfire Activity in the Alaskan Tundra. *Journal of Geophysical Research:*

Biogeosciences, *127*(3), e2021JG006608. https://doi.org/10.1029/2021JG006608

Morelli, T. L., Barrows, C. W., Ramirez, A. R., Cartwright, J. M., Ackerly, D. D., Eaves, T. D.,

Ebersole, J. L., Krawchuk, M. A., Letcher, B. H., Mahalovich, M. F., Meigs, G. W., Michalak, J.

L., Millar, C. I., Quiñones, R. M., Stralberg, D., & Thorne, J. H. (2020). Climate-change refugia:

biodiversity in the slow lane. *Frontiers in Ecology and the Environment*, *18*(5), 228–234.

https://doi.org/10.1002/fee.2189

Morelli, T. L., Daly, C., Dobrowski, S. Z., Dulen, D. M., Ebersole, J. L., Jackson, S. T., Lundquist, J.

D., Millar, C. I., Maher, S. P., Monahan, W. B., Nydick, K. R., Redmond, K. T., Sawyer, S. C.,

Stock, S., & Beissinger, S. R. (2016). Managing climate change refugia for climate adaptation.

PLoS ONE, *11*(8), 1–17. https://doi.org/10.1371/journal.pone.0159909

Myers-Smith, I. H., Forbes, B. C., Wilmking, M., Hallinger, M., Lantz, T., Blok, D., Tape, K. D., MacIas-

Fauria, M., Sass-Klaassen, U., Lévesque, E., Boudreau, S., Ropars, P., Hermanutz, L., Trant,

A., Collier, L. S., Weijers, S., Rozema, J., Rayback, S. A., Schmidt, N. M., … Hik, D. S. (2011).

Shrub expansion in tundra ecosystems: Dynamics, impacts and research priorities.

Environmental Research Letters. https://doi.org/10.1088/1748-9326/6/4/045509

Myers-Smith, I. H., Kerby, J. T., Phoenix, G. K., Bjerke, J. W., Epstein, H. E., Assmann, J. J., John,

C., Andreu-Hayles, L., Angers-Blondin, S., Beck, P. S. A., Berner, L. T., Bhatt, U. S., Bjorkman,

A. D., Blok, D., Bryn, A., Christiansen, C. T., Cornelissen, J. H. C., Cunliffe, A. M., Elmendorf, S.

C., … Wipf, S. (2020). Complexity revealed in the greening of the Arctic. In *Nature Climate*

Change. https://doi.org/10.1038/s41558-019-0688-1

Myneni, R. B., Keeling, C. D., Tucker, C. J., Asrar, G., & Nemani, R. R. (1997). Increased plant growth

in the northern high latitudes from 1981 to 1991. *Nature*. https://doi.org/10.1038/386698a0

Noss, R. F. (2001). Beyond Kyoto: Forest management in a time of rapid climate change. In

Conservation Biology. https://doi.org/10.1046/j.1523-1739.2001.015003578.x

Nuttall, M., Berkes, F., Forbes, B., Kofinas, G., Vlassova, T., & Wenzel, G. W. (2005). Ch.12 Hunting,

herding, fishing, and gathering: Indigenous peoples and renewable resource use in the Arctic.

Arctic Climate Impact Assessment.

Oehri, J., Schaepman-Strub, G., Kim, J.-S., Grysko, R., Kropp, H., Grünberg, I., Zemlianskii, V.,

Sonnentag, O., Euskirchen, E. S., Reji Chacko, M., & the ArcticSEB-Synthesis Team. (2022).

Vegetation Type is an Important Predictor of the Arctic Terrestrial Summer Surface Energy

Budgethttps. *EGU General Assembly 2022*. https://doi.org/10.5194/egusphere-egu22-7636

Pearson, R. G., Phillips, S. J., Loranty, M. M., Beck, P. S. A., Damoulas, T., Knight, S. J., & Goetz, S.

J. (2013b). Shifts in Arctic vegetation and associated feedbacks under climate change. *Nature*

Climate Change. https://doi.org/10.1038/nclimate1858

Pearson, R. G., Phillips, S. J., Loranty, M. M., Beck, P. S. A., Damoulas, T., Knight, S. J., & Goetz, S.

J. (2013a). *Predicted Arctic vegetation distribution shifts under future climate change*. Arctic

Data Center. https://doi.org/10.18739/A2V935.

Phoenix, G. K., & Bjerke, J. W. (2016). Arctic browning: extreme events and trends reversing arctic

greening. In *Global Change Biology*. https://doi.org/10.1111/gcb.13261

R Development Core Team. (2017). R: A language and environment for statistical computing. In

Vienna, Austria. https://doi.org/R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-

900051-07-0, URL http://www.R-project.org.

Raynolds, M. K., Jorgenson, J. C., Jorgenson, M. T., Kanevskiy, M., Liljedahl, A. K., Nolan, M., Sturm,

M., & Walker, D. A. (2020). Landscape impacts of 3D☐seismic surveys in the Arctic National

Wildlife Refuge, Alaska. *Ecological Applications*, *0*(0), 1–20. https://doi.org/10.1002/eap.2143

Raynolds, M. K., & Walker, D. A. (2009). Effects of deglaciation on circumpolar distribution of arctic

vegetation. *Canadian Journal of Remote Sensing*. https://doi.org/10.5589/m09-006

Raynolds, M. K., Walker, D. A., Ambrosius, K. J., Brown, J., Everett, K. R., Kanevskiy, M., Kofinas, G.

P., Romanovsky, V. E., Shur, Y., & Webber, P. J. (2014). Cumulative geoecological effects of 62

years of infrastructure and climate change in ice-rich permafrost landscapes, Prudhoe Bay

Oilfield, Alaska. *Global Change Biology*. https://doi.org/10.1111/gcb.12500

Raynolds, M. K., Walker, D. A., Balser, A., Bay, C., Campbell, M., Cherosov, M. M., Daniëls, F. J. A.,

Eidesen, P. B., Ermokhina, K. A., Frost, G. V., Jedrzejek, B., Jorgenson, M. T., Kennedy, B. E.,

Kholod, S. S., Lavrinenko, I. A., Lavrinenko, O. V., Magnússon, B., Matveyeva, N. V.,

Metúsalemsson, S., … Troeva, E. (2019). A raster version of the Circumpolar Arctic Vegetation

Map (CAVM). *Remote Sensing of Environment*. https://doi.org/10.1016/j.rse.2019.111297

Riordan, B., Verbyla, D., & McGuire, A. D. (2006). Shrinking ponds in subarctic Alaska based on

1950-2002 remotely sensed images. *Journal of Geophysical Research: Biogeosciences*.

https://doi.org/10.1029/2005JG000150

Rundqvist, S., Hedenås, H., Sandström, A., Emanuelsson, U., Eriksson, H., Jonasson, C., &

Callaghan, T. V. (2011). Tree and shrub expansion over the past 34 years at the tree-line near

Abisko, Sweden. In *Ambio*. https://doi.org/10.1007/s13280-011-0174-0

SAExploration. (2018). *Marsh Creek 3D plan of operationswinter seismic survey*.

https://eplanning.blm.gov/public_projects/nepa/111085/153349/187888/Marsh_Creek_Plan_of_

Operations_Submitted_May2018.pdf

Scibilia, E., Walker, D. A., Kofinas, G., Raynolds, M., Kanevskiy, M., Shur, Y., Ambrosius, K.,

Buchhorn, M., Matyshak, G., Romanovsky, V., Wirth, L., Kumpula, T., Forbes, B., Khumotov, A.,

Leibman, M., Khitun, O., Lemay, M., Allard, M., Lamoureux, S., & Shiklomanov, N. (2015). *Rapid*

Arctic Transitions due to Infrastructure and Climate (RATIC): A contribution to ICARP III.

https://doi.org/10.13140/RG.2.1.4089.0006

Serreze, M. C., Walsh, J. E., Chapin, F. S., Osterkamp, T., Dyurgerov, M., Romanovsky, V., Oechel,

W. C., Morison, J., Zhang, T., & Barry, R. G. (2000). Observational evidence of recent change in

the northern high-latitude environment. *Climatic Change*.

https://doi.org/10.1023/A:1005504031923

Smith, L. C. (2005). Disappearing Arctic Lakes. *Science*. https://doi.org/10.1126/science.1108142

Stewart, L., Alsos, I. G., Bay, C., Breen, A. L., Brochmann, C., Boulanger-Lapointe, N., Broennimann,

O., Bültmann, H., Bøcher, P. K., Damgaard, C., Daniëls, F. J. A., Ehrich, D., Eidesen, P. B.,

Guisan, A., Jónsdóttir, I. S., Lenoir, J., le Roux, P. C., Lévesque, E., Luoto, M., … Pellissier, L.

(2016). The regional species richness and genetic diversity of Arctic vegetation reflect both past

glaciations and current climate. *Global Ecology and Biogeography*.

https://doi.org/10.1111/geb.12424

Stralberg, D., Carroll, C., & Nielsen, S. E. (2020). Toward a climate-informed North American

protected areas network: Incorporating climate-change refugia and corridors in conservation

planning. *Conservation Letters*. https://doi.org/10.1111/conl.12712

Sturm, M., Racine, C., & Tape, K. (2001). Increasing shrub abundance in the Arctic. *Nature*.

https://doi.org/10.1038/35079180

Sulman, B. N., Salmon, V. G., Iversen, C. M., Breen, A. L., Yuan, F., & Thornton, P. E. (2021).

Integrating Arctic Plant Functional Types in a Land Surface Model Using Above- and

Belowground Field Observations. *Journal of Advances in Modeling Earth Systems*, *13*(4),

e2020MS002396. https://doi.org/10.1029/2020MS002396

Swann, A. L., Fung, I. Y., Levis, S., Bonan, G. B., & Doney, S. C. (2010). Changes in arctic vegetation

amplify high-latitude warming through the greenhouse effect. *Proceedings of the National*

Academy of Sciences of the United States of America. https://doi.org/10.1073/pnas.0913846107

Taberlet, P., & Cheddadi, R. (2002). Ecology: Quaternary refugia and persistence of biodiversity. In

Science. https://doi.org/10.1126/science.297.5589.2009

Team RStudio. (2020). RStudio: Integrated Development for R. In *Rstudio Team, PBC, Boston, MA*

URL http://www.rstudio.com/.

Thorne, J. H., Gogol-Prokurat, M., Hill, S., Walsh, D., Boynton, R. M., & Choe, H. (2020). Vegetation

refugia can inform climate-adaptive land management under global warming. *Frontiers in*

Ecology and the Environment, *18*(5), 281–287. https://doi.org/10.1002/fee.2208

Tzedakis, P. C., Lawson, I. T., Frogley, M. R., Hewitt, G. M., & Preece, R. C. (2002). Buffered tree

population changes in a Quaternary refugium: Evolutionary implications. *Science*.

https://doi.org/10.1126/science.1073083

Venter, O., Sanderson, E. W., Magrach, A., Allan, J. R., Beher, J., Jones, K. R., Possingham, H. P.,

Laurance, W. F., Wood, P., Fekete, B. M., Levy, M. A., & Watson, J. E. M. (2016). Global

terrestrial Human Footprint maps for 1993 and 2009. *Scientific Data*.

https://doi.org/10.1038/sdata.2016.67

Virkkala, A. M., Natali, S. M., Rogers, B. M., Watts, J. D., Savage, K., Connon, S. J., Mauritz, M.,

Schuur, E. A. G., Peter, D., Minions, C., Nojeim, J., Commane, R., Emmerton, C. A., Goeckede,

M., Helbig, M., Holl, D., Iwata, H., Kobayashi, H., Kolari, P., … Zyryanov, V. I. (2022). The

ABCflux database: Arctic-boreal CO2flux observations and ancillary information aggregated to

monthly time steps across terrestrial ecosystems. *Earth System Science Data*, *14*(1), 179–208.

https://doi.org/10.5194/essd-14-179-2022

Walker, D. A., Raynolds, M. K., Daniëls, F. J. A., Einarsson, E., Elvebakk, A., Gould, W. A., Katenin,

A. E., Kholod, S. S., Markon, C. J., Melnikov, E. S., Moskalenko, N. G., Talbot, S. S., Yurtsev, B.

A., Bliss, L. C., Edlund, S. A., Zoltai, S. C., Wilhelm, M., Bay, C., Gudjónsson, G., … Vairin, B. A.

(2005). The Circumpolar Arctic vegetation map. *Journal of Vegetation Science*.

https://doi.org/10.1111/j.1654-1103.2005.tb02365.x

Walker, M. D., Wahren, C. H., Hollister, R. D., Henry, G. H. R., Ahlquist, L. E., Alatalo, J. M., Bret-

Harte, M. S., Calef, M. P., Callaghan, T. V., Carroll, A. B., Epstein, H. E., Jónsdóttir, I. S., Klein,

J. A., Magnússon, B., Molau, U., Oberbauer, S. F., Rewa, S. P., Robinson, C. H., Shaver, G. R.,

… Wookey, P. A. (2006). Plant community responses to experimental warming across the

tundra biome. *Proceedings of the National Academy of Sciences of the United States of*

America. https://doi.org/10.1073/pnas.0503198103

Walsh, J. E., Overland, J. E., Groisman, P. Y., & Rudolf, B. (2011). Ongoing climate change in the

arctic. *Ambio*. https://doi.org/10.1007/s13280-011-0211-z

Walvoord, M. A., & Kurylyk, B. L. (2016). Hydrologic Impacts of Thawing Permafrost-A Review.

Vadose Zone Journal. https://doi.org/10.2136/vzj2016.01.0010

Wheeler, H. C., Høye, T. T., & Svenning, J. C. (2018). Wildlife species benefitting from a greener

Arctic are most sensitive to shrub cover at leading range edges. *Global Change Biology*.

https://doi.org/10.1111/gcb.13837

White, D. M., Craig Gerlach, S., Loring, P., Tidwell, A. C., & Chambers, M. C. (2007). Food and water

security in a changing arctic climate. *Environmental Research Letters*.

https://doi.org/10.1088/1748-9326/2/4/045018

Woo, M. K., & Young, K. L. (2006). High Arctic wetlands: Their occurrence, hydrological

characteristics and sustainability. *Journal of Hydrology*.

https://doi.org/10.1016/j.jhydrol.2005.07.025

Wookey, P. A., Aerts, R., Bardgett, R. D., Baptist, F., Bråthen, K., Cornelissen, J. H. C., Gough, L.,

Hartley, I. P., Hopkins, D. W., Lavorel, S., & Shaver, G. R. (2009). Ecosystem feedbacks and

cascade processes: Understanding their role in the responses of Arctic and alpine ecosystems

to environmental change. *Global Change Biology*. https://doi.org/10.1111/j.1365-

2486.2008.01801.x

Yoshikawa, K., & Hinzman, L. D. (2003). Shrinking thermokarst ponds and groundwater dynamics in

discontinuous permafrost near Council, Alaska. *Permafrost and Periglacial Processes*.

https://doi.org/10.1002/ppp.451

Yu, L., Leng, G., & Python, A. (2022). Attribution of the spatial heterogeneity of Arctic surface albedo

feedback to the dynamics of vegetation, snow and soil properties and their interactions.

Environmental Research Letters, *17*(1), 014036. https://doi.org/10.1088/1748-9326/ac4631

Zona, D., Lafleur, P. M., Hufkens, K., Bailey, B., Gioli, B., Burba, G., Goodrich, J. P., Liljedahl, A. K.,

Euskirchen, E. S., Watts, J. D., Farina, M., Kimball, J. S., Heimann, M., Göckede, M., Pallandt,

M., Christensen, T. R., Mastepanov, M., López-Blanco, E., Jackowicz-Korczynski, M., …

Oechel, W. C. (2022). Earlier snowmelt may lead to late season declines in plant productivity

and carbon sequestration in Arctic tundra ecosystems. *Scientific Reports*, *12*(1), 1–10.

https://doi.org/10.1038/s41598-022-07561-1

Figures
767

768
Figure 1. **The 2003 baseline abundance and protection status of Arctic vegetation types.** The Arctic tundra encompasses
approximately 4.7 million km^2. The height of the bars represents the absolute extent of each vegetation type within and outside
of protected areas. The percentages above the bars represent the relative abundance of the vegetation type within the pan-
Arctic tundra. The percentages within the darker coloured bars (protected area) represent the relative abundance of each
vegetation type in the protected area in comparison to the total protected area. The vegetation types can be summarised into
barren tundra (B1: cryptogam herb barren; B2: cryptogam barren complex (bedrock); B3: noncarbonate mountain complex and
B4: carbonate mountain complex), graminoid tundra (G1: rush/grass forb, cryptogam tundra; G2: graminoid, prostrate dwarf-
shrub, forb tundra; G3: nontussock sedge, dwarf-shrub, moss tundra; G4: tussock sedge, dwarf-shrub, moss tundra), prostrate-
shrub tundra (P1: prostrate dwarf shrub, herb tundra; P2: prostrate/hemiprostrate dwarf-shrub tundra), erect-shrub tundra (S1:
erect dwarf-shrub tundra; S2: low shrub tundra) and wetlands (W1: sedge/grass, moss wetland; W2: sedge, moss, dwarf-shrub
wetland; W3: sedge, moss, low-shrub wetland).

Figure 2. **A comparison of the abundance and protection status of Arctic vegetation types at the baseline and in the future.** The data presented refers to the realistic model scenario; error bars represent the envelope of the minimal and maximal abundance predicted by the different MFAVDs. The minimum and maximum values of the envelope are not always the result of the same MFAVDs. Depending on the vegetation type, different models predict the highest or lowest abundance. The vegetation types can be summarised into graminoid tundra (G1: rush/grass forb, cryptogam tundra; G2: graminoid, prostrate dwarf-shrub, forb tundra; G3: nontussock sedge, dwarf-shrub, moss tundra; G4: tussock sedge, dwarf-shrub, moss tundra), prostrate-shrub tundra (P1: prostrate dwarf shrub, herb tundra; P2: prostrate/hemiprostrate dwarf-shrub tundra), and erect-shrub tundra (S1: erect dwarf-shrub tundra; S2: low shrub tundra). Barren and wetland vegetation types were not assessed as predictions for 2050 were unavailable.

Figure 3. **Risk status according to the 'decline in distribution' criterion of the IUCN Red List of Ecosystems Categories**
and Criteria under the realistic model. The vegetation types are ordered along the bioclimatic subzones they occur in, from
low to high summer temperatures (left to right). The data presented refers to the realistic model scenario; error bars represent
the envelope of the minimal and maximal abundance predicted by the different MFAVDs. The minimum and maximum values
of the envelope are not always the result of the same MFAVDs. Depending on the vegetation type, different models predict the
highest or lowest abundance. The vegetation types can be summarised into graminoid tundra (G1: rush/grass forb, cryptogam
tundra; G2: graminoid, prostrate dwarf-shrub, forb tundra; G3: nontussock sedge, dwarf-shrub, moss tundra; G4: tussock
sedge, dwarf-shrub, moss tundra), prostrate-shrub tundra (P1: prostrate dwarf shrub, herb tundra; P2: prostrate/hemiprostrate
dwarf-shrub tundra), and erect-shrub tundra (S1: erect dwarf-shrub tundra; S2: low shrub tundra). Barren and wetland
vegetation types were not assessed as no predictions for 2050 were available.

Figure 4. **Refugia maps of threatened vegetation types in the Arctic (a), and magnified subsets centred on the Yamal peninsula, Russia (b); the Canadian Archipelago (c); and Alaska, USA, and North-eastern Siberia, Russia (d).** The vulnerable vegetation types presented here are G1: rush/grass forb, cryptogam tundra; G2: graminoid, prostrate dwarf-shrub, forb tundra; P1: prostrate dwarf shrub, herb tundra; P2: prostrate/hemiprostrate dwarf-shrub tundra and S1: erect dwarf-shrub tundra.