Title page

Sleep-dependent upscaled excitability and saturated neuroplasticity in the human brain: From brain physiology to cognition

Mohammad Ali Salehinejad [1,2], Elham Ghanavati [1,3], Joerg Reinders [4], Jan G. Hengstler [4], Min-Fang Kuo [1], and Michael A. Nitsche [1,5*]

1 Department of Psychology and Neurosciences, Leibniz Research Centre for Working Environment and Human Factors, Dortmund, Germany
2 International Graduate School of Neuroscience, Ruhr-University Bochum, Bochum, Germany
3 Department of Neuropsychology, Institute of Cognitive Neuroscience, Faculty of Psychology, Ruhr University Bochum, Bochum, Germany
4 Department of Toxicology, Leibniz Research Centre for Working Environment and Human Factors, Dortmund, Germany
5 University Medical Hospital Bergmannsheil, Department of Neurology, Bochum, Germany

*Corresponding author: Prof. Dr. Michael A. Nitsche

Department of Psychology and Neurosciences, Leibniz Research Centre for Working Environment and Human Factors, Ardeystr. 67, 44139 Dortmund, Germany
Phone: +49 231 1084 301
Email: nitsche@ifado.de

Abstract word count: 156
Main text word count (excluding the Materials and methods, References, and Figure legends): 5130
Display items (Figures + Tables): 7
Supplementary material display items: 3 figure, 5 table

Abstract

Sleep and cognition are strongly linked via their impact on synaptic strength. Whether and *how* sleep deprivation modulates human brain physiology and cognition is, however, not well understood. Here we examined how cortical excitability, inducibility of LTP- and-LTD-like plasticity, learning and memory formation and higher-order cognition are affected by overnight sleep deprivation. We show that sleep deprivation upscales cortical excitability due to enhanced glutamate-related cortical facilitation and decreased and/or reversed GABAergic cortical inhibition. We furthermore demonstrate that non-invasive brain stimulation-induced LTP-like plasticity is abolished while LTD-like plasticity converts to excitatory LTP-like under sleep deprivation, likely caused by synaptic saturation and upscaled brain excitability. This was associated with increased synaptic strength measured by EEG theta activity. Finally, we show that learning and memory formation, behavioral counterparts of plasticity, and working memory and attention, which rely on cortical excitability, are impaired during sleep deprivation. Our data show converging effects of sleep deprivation on human brain physiology and cognition.

Keywords: sleep, cortical excitability, neuroplasticity, synaptic homeostasis, learning and memory, non-invasive brain stimulation.

1. Introduction

Over the past decade, a strong link has been established between sleep and cognition (Deak and Stickgold, 2010; Lim and Dinges, 2010). Adequate sleep is critical for optimal cognitive functions across the lifespan (Carskadon, 2011; Lo et al., 2016) and findings from experimental settings support this critical role of sleep for cognition in animals (Walker and Stickgold, 2006), and humans (Krause et al., 2017) especially for memory consolidation and sequence learning (Stickgold, 2005; Chouhan et al., 2021). As a ubiquitous *physiological* phenomenon, sleep has extensive impacts on brain physiology and especially on parameters relevant for cognition such as brain excitability and plasticity.

Previous experimental studies, mostly in nonhuman animals, have linked sleep and synaptic homeostasis. Specifically, extended wakefulness (or sleep deprivation) is associated with the expression of long-term potentiation (LTP)-related molecular changes and plasticity-related genes (e.g. BDNF, CREB) in the brain, leading to saturation of synaptic potentiation (Tononi and Cirelli, 2003). Sleep, on the other hand, desaturates synapses that have been potentiated during wakefulness, resulting in an improved signal-to-noise ratio and a renewed capacity for encoding new information and cognitive processing (Kuhn et al., 2016). A recent study confirmed this sleep-dependent synaptic downscaling by showing reduced or weakened synaptic connections in the primary motor and somatosensory cortex during sleep (de Vivo et al., 2017). This demonstrates that sleep is required for preparing the brain for proper cognitive, motor, and physiological functioning, however, the effect of sleep on specific parameters of *human* brain physiology and their association with cognition and behaviour remains to be further determined.

In humans, molecular mechanisms of synaptic homeostasis cannot be directly studied, however, non-invasive (indirect) markers of brain physiology can be used for studying the impact of sleep and extended wakefulness on synaptic potentiation and cortical excitability (Kuhn et al., 2016). Non-invasive brain stimulation (NIBS) techniques are safe methods for directly monitoring, and modifying brain functions in humans providing a means for studying the causality of brain-behaviour relationships (Polanía et al., 2018). Several NIBS techniques, including transcranial magnetic stimulation (TMS) and transcranial electrical stimulation (tES), are widely used to non-invasively monitor and induce changes in cortical excitability, and neuroplasticity (Nitsche and Paulus, 2000; Huang et al., 2017; Polanía et al., 2018). Recently, these techniques have been used for studying the contribution of sleep to brain physiological and cognitive functions in humans. It has been shown that cortical excitability

increases after sleep deprivation (Kuhn et al., 2016; Ly et al., 2016). This increase of cortical

excitability after sleep deprivation is linked to increased net synaptic strength in humans (i.e.,

TMS intensity required to elicit a predefined motor-evoked potential, EEG activity) (Kuhn et

al., 2016), animals (i.e, synaptic potentiation) (Vyazovskiy et al., 2008), and reduced

inhibitory intracortical mechanisms (Kreuzer et al., 2011; Placidi et al., 2013). This can

reduce the ability of the brain to induce neuroplasticity (as a result of synaptic saturation). In

this line, decreased LTP has been shown in rats after sleep deprivation shown in both, *in vivo*

and *in vitro* (McDermott et al., 2003; Kopp et al., 2006; Vyazovskiy et al., 2008), and a recent

human study also showed decreased LTP-like plasticity, induced by paired associative

stimulation (Kuhn et al., 2016).

While these studies added novel insight into sleep-dependent effects on human brain

physiology, a *comprehensive* investigation of how brain physiology and cognition interact

under sleep deprivation is missing. Different excitatory and inhibitory neurotransmitter

systems (e.g. glutamatergic, dopaminergic, GABAergic, cholinergic) are involved in cortico-

cortical and corticospinal *excitability* which are also closely related to neuroplasticity and

cognition. These specific parameters of brain excitability can be monitored with different

TMS protocols. Furthermore, the suggested role of the sleep-wake cycle for the inducibility of

synaptic plasticity, which appears to be dependent on an optimal sleep-dependent temporal

window (Kuhn et al., 2016), can be investigated via induction of both LTP- and/or LTD-like

plasticity. Very few studies in humans have investigated induced neuroplasticity under sleep

deprivation conditions so far, and these studies are mostly limited to LTP-like plasticity

(Kuhn et al., 2016), leaving LTD-like plasticity untouched. Gaining knowledge about LTP-

and LTD-like plasticity mechanisms of action can extend the sleep synaptic hypothesis

(Tononi and Cirelli, 2014) for the human brain. Moreover, *whether* and *how* these

physiological parameters are associated with cognitive functions remains to be investigated.

A common research paradigm for studying the effect of sleep on these parameters of brain

physiology and cognition is sleep deprivation. In this context, a strong and strict control is

exerted over all possible exogenous (body posture, light, temperature) and endogenous (stress

level, digestion, motivation) factors that might affect the sleep-wake cycle, avoiding a

confounding influence of these parameters on the factors of interest (Schmidt et al., 2007). In

this paradigm, participants are kept in an extended wakefulness condition and are deprived of

sleep for a certain amount of time. Using this paradigm, we systematically investigated the

impact of one-night sleep deprivation, compared to one-night sufficient sleep on converging

measures of human brain physiology and cognition. Specifically, we monitored cortical

excitability of the brain via TMS protocols that measure cortical inhibition and facilitation (i),

induced both LTP-like and LTD-like plasticity (ii), and measured synaptic strength via the

resting-EEG theta/alpha pattern (iii). We also examined learning and memory formation,

behavioral indices of brain plasticity (iv), and higher-order cognitive functions (attention,

working memory) which depend on cortical excitability (v) and their electrophysiological

correlates. Cortisol and melatonin levels were also assessed (vi). All measurements were

conducted under sufficient sleep vs one-night sleep deprivation at fixed times (Figure 1).

Briefly, we demonstrate that after sleep deprivation, the human brain displays a hyperexcited

state marked by exaggerated intracortical facilitation and diminished or converted

intracortical inhibition. Inducing neuroplasticity at this state, diminishes anodal stimulation-

induced changes in the motor-evoked potentials (LTP-like plasticity) while converts cathodal

stimulation-induced LTD-like plasticity into LTP-like excitatory state. EEG theta and beta

activity in the same line show increased synaptic net strength after sleep deprivation. These

physiological findings are associated with comprised learning, memory and attentional

functioning at behavioral and electrophysiological levels.

Fig.1: **The course of the experiment. a,** Using single-pulse and double-pulse TMS protocols, corticospinal and corticocortical excitability was measured after sleep deprivation or sufficient sleep. RMT = Resting motor threshold, AMT = Active motor threshold, SAI = Short-latency afferent inhibition, SICI-ICF = short-latency intracortical inhibition and facilitation, I-O curve = Input-output curve. **b,**

Neuroplasticity was induced with anodal and cathodal stimulation after sufficient sleep (SS) vs sleep deprivation (SD). SI1 mv=Stimulation intensity to elicit an MEP amplitude of 1 mV, M1=primary motor cortex, Fp2=right supraorbital area. **c,** Saliva samples were taken at 8:45 in each session. Following the resting-EEG acquisition, participants performed motor learning, working memory, and attention tasks at the beginning of each experimental session (sufficient sleep vs sleep deprivation) while their EEG was recorded. SRTT=serial reaction time task, AX-CPT=AX continuous performance task.

2. Results

2.1. Sleep deprivation upscales cortical excitability

We monitored corticospinal and intracortical excitability of the motor cortex after "sufficient

sleep" and "sleep deprivation" sessions with different TMS protocols. Input-Output curve (I-

O curve) and intracortical facilitation (ICF) were used as measures of global corticospinal

excitability and cortical facilitation, respectively. Short-interval cortical inhibition (SICI), I-

wave facilitation, and short-latency afferent inhibition (SAI) were applied as cortical

inhibition protocols. These TMS protocols are based on different predominant

neurotransmitter systems related to cortical facilitation (glutamatergic) and inhibition

(GABAergic, cholinergic) (Chen, 2000; Di Lazzaro et al., 2000; Di Lazzaro et al., 2005a) (see

Methods). Baseline MEP values of control conditions in TMS protocols did not significantly

differ across sleep conditions (Tables S1, S2) and the changes in protocol-specific MEPs

cannot be due to baseline MEP differences across sleep conditions.

Input-output curve (I/O curve). I-O curve is a global measure of *corticospinal*

excitability (Boroojerdi et al., 2001) and the slope of the I-O curve reflects excitability of

corticospinal neurons modulated by glutamatergic activity at higher TMS intensities (see

Methods). The results of the 2×4 ANOVA showed a marginally significant interaction of

sleep condition×TMS intensity ($F_{1.71}=3.41$, $p=0.048$; $\eta p^2=0.10$), and significant main effects

of sleep condition ($F_1=4.95$, $p=0.034$; $\eta p^2=0.14$) and TMS intensity ($F_{1.22}=100.13$, $p<0.001$;

$\eta p^2=0.77$) on the slope of the I-O curve. MEP amplitudes were numerically larger at all TMS

intensities after sleep deprivation vs sufficient sleep, however, these differences were not

significant based on the Bonferroni-corrected post hoc comparisons (Fig. 2a).

SICI-ICF. In this double-pulse TMS protocol, the interstimulus interval (ISI) between a

subthreshold conditioning stimulus and a suprathreshold test stimulus determines inhibitory

(ISIs 2 and 3 ms) or facilitatory (ISIs 10 and 15 ms) effects on cortical excitability (Kujirai et

al., 1993). The results of the 2×5 ANOVA showed a significant interaction of sleep

condition×ISI ($F_{3.69}=14.85$, $p<0.001$, $\eta p^2=0.34$), and significant main effects of sleep

condition ($F_1=13.81$, $p<0.001$, $\eta p^2=0.72$), and ISI ($F_{3.10}=93.77$, $p<0.001$, $\eta p^2=0.76$) on MEP

values. Bonferroni-corrected post hoc comparisons revealed a significant intracortical inhibition shown by decreased MEPs in the ISI 2 and 3 ms conditions only after sufficient sleep, and significant differences of MEPs obtained with these ISIs across sleep conditions (Fig. 2b). This indicates that intracortical inhibition was significantly lower after sleep deprivation vs sufficient sleep. For intracortical facilitation, MEP amplitudes were significantly enhanced only at an ISI of 15 ms when compared with single pulse-elicited MEP amplitudes (baseline) after sufficient sleep, while they were significantly increased at ISIs of 10 and 15 ms after sleep deprivation. These MEPs were also significantly larger after sleep deprivation vs respective MEPs after sufficient sleep (Fig. 2b). Together, these results demonstrate a significantly *lower* cortical inhibition and *higher* cortical facilitation after sleep deprivation.

I-wave facilitation. In this double-pulse TMS protocol, cortical inhibition is reflected by I-wave peaks which are mainly observed at three ISIs occurring at 1.1-1.5 ms (early), 2.3-2.9 ms (middle), and 4.1-4.4 (late) ms after test pulse application. The results of the 2×3 ANOVA showed a significant interaction of sleep condition×ISI ($F_{1.74}=14.59$, $p<0.001$, $\eta p^2=0.33$), and main effects of sleep condition ($F_1=20.36$, $p<0.001$, $\eta p^2=0.41$) and ISI ($F_{1.67}=47.39$, $p<0.001$, $\eta p^2=0.62$) on I-wave peak MEP amplitudes. Bonferroni-corrected post hoc comparisons showed a significant increase of I-wave peaks for early and middle ISIs, as compared to single-pulse MEPs after both sleep conditions. The I-wave peaks for late ISI were significant only after sleep deprivation. Importantly, the peaks (at all ISIs) were significantly larger after sleep deprivation vs sufficient sleep (Fig. 2c). These results indicate reduced GABAergic inhibition, resulting in I-wave facilitation, after sleep deprivation.

SAI. In this protocol, the TMS stimulus is coupled with peripheral nerve stimulation that has an inhibitory effect on motor cortex excitability at ISIs of 20 and 40 ms. Smaller MEPs indicate cortical inhibition. A significant interaction of sleep condition×ISI ($F_{1.81}=27.51$, $p<0.001$, $\eta p^2=0.48$) and a significant main effect of sleep condition ($F_1=70.18$, $p<0.001$, $\eta p^2=0.71$), but not ISI ($F_1=1.58$, $p<0.217$, $\eta p^2=0.05$) were observed on MEP amplitudes. Bonferroni-corrected post hoc comparisons revealed a significantly pronounced inhibitory effect of peripheral stimulation on cortical excitability after *sufficient sleep,* compared to the single TMS pulse at both ISIs. However, respective MEPs were significantly *converted* to excitatory effects after sleep deprivation. Moreover, cortical inhibition was significantly reduced after sleep deprivation vs sufficient sleep at the respective ISIs (Fig. 2d). In line with the SICI and I-wave protocols, this suggests a reduction of cortical inhibition and its *conversion* to excitatory effects after sleep deprivation.

Fig 2. **Corticospinal and corticocortical excitability after sufficient sleep vs sleep deprivation**. **a,** There is a trend of higher corticospinal excitability after the sleep deprivation session compared to sufficient sleep especially at 150% of RMT intensity. **b,** Cortical inhibition

significantly decreased after sleep deprivation as compared with sufficient sleep (t_{ISI2}=4.24, p<0.001; t_{ISI3}=4.50, p<0.001). In contrast, cortical facilitation is significantly upscaled after sleep deprivation compared with sufficient sleep (t_{ISI10}=7.69, p<0.001; t_{ISI15}= 6.66, p<0.001). **c,** I-wave peaks were significantly facilitated for early and middle ISIs after both, sufficient sleep and sleep deprivation, and for late ISIs only after sleep deprivation. For all ISIs, I-wave peaks were significantly more upscaled after sleep deprivation vs sufficient sleep (t_{early}=3.90, p<0.001; t_{middle}=3.91, p<0.001; t_{late}=4.40, p<0.001), indicative of less cortical inhibition. **d,** Cortical inhibitory effect of peripheral nerve stimulation on motor cortical excitability was observed only after sufficient sleep (t_{ISI20}=4.53, p<0.001; t_{ISI40}=4.25, p<0.001), whereas the inhibitory effect of peripheral stimulation turned to excitatory effects after sleep deprivation (t_{ISI20}=2.54, p=0.035; t_{ISI40}=4.55, p<0.001). MEP amplitude were significantly upscaled after sleep deprivation vs sufficient sleep (t_{ISI20}=7.08, p<0.001; t_{ISI40}=8.83, p<0.001). All pairwise comparisons were calculated using the Bonferroni correction for multiple comparisons (n=30). All error bars represent the standard error of means (s.e.m.). Filled symbols represent a significant difference in MEP amplitudes compared to the respective test pulses (for SICI-ICF, I-Wave, SAI) or MEP at RMT intensity (for I-O curve). Asterisks represent statistically significant comparisons between sleep conditions. *Note*: MEP = motor evoked potential; RMT = resting motor threshold; ms = milliseconds.

Taken together, our results demonstrate that corticocortical and corticospinal excitability are

upscaled after sleep deprivation. Interestingly, cortical inhibition was decreased or turned into

facilitation which is again indicative of higher cortical excitability. Cortical excitability is

closely related to LTP/LTD plasticity in the brain and is expected to be related to changes of

synaptic potentiation after lack of sleep (Tononi and Cirelli, 2003; Kuhn et al., 2016).

Accordingly, in the next step, we investigated the impact of non-invasively inducing

LTP/LTD-like neuroplasticity under sleep deprivation vs sufficient sleep conditions.

171
2.2. Sleep deprivation saturates induction of LTP-like plasticity and converts the direction
of LTD-like plasticity

Here, we were interested in determining how sleep deprivation, and the resultant upscaled

cortical excitability, affect LTP- and-LTD-like plasticity in the brain. The sleep synaptic

hypothesis proposes that synaptic strength is saturated during long awake times and restored

after sleep (Tononi and Cirelli, 2014). Saturation can lead to decreased LTP-like plasticity in

humans (Kuhn et al., 2016). Accordingly, we expected that induction of LTP-like plasticity is

decreased due to saturated synaptic strength and hyperexcited brain state identified in the

previous section. We were also interested in determining how induction of LTD-like plasticity

is affected under this brain state, which has not been investigated so far. To this end,

participants received "anodal vs sham" and "cathodal vs sham" transcranial direct current

stimulation (tDCS) over the motor cortex after sufficient sleep and sleep deprivation (see

Methods). The MEPs (7 timepoints) obtained from each tDCS state (active, sham) in each group (anodal, cathodal) across sleep conditions (sufficient sleep, sleep deprivation) were analyzed with a mixed model 2×2×2×7 ANOVA. A significant four-way interaction of sleep condition×group×tDCS state×timepoint was found ($F_{5.35}$=12.14, p<0.001, ηp^2=0.30), indicating that tDCS-induced LTP/LTD-like neuroplasticity was differentially affected in the sleep conditions. Other interactions and main effects are summarized in Table 1. Baseline MEPs did not significantly differ across sleep and stimulation conditions (Table S3). Reported side effects and analyses of blinding efficacy can be found in the supplementary material (Tables S4-5).

Anodal LTP-like-induced plasticity. Bonferroni-corrected post hoc *t*-tests reveal that after sufficient sleep, anodal tDCS significantly increased MEP amplitudes immediately after, 5, 10, 15, 20, and 25 min after the intervention. The increase of MEP amplitudes significantly larger at all timepoints when compared to sleep deprivation and against the sham intervention (Fig. 3a,b). In contrast, sleep deprivation prevented induction of LTP-like plasticity via anodal tDCS at all timepoints. No significant effect of anodal tDCS was observed when compared to the baseline and against the sham intervention after sleep deprivation for all timepoints.

Cathodal LTD-like-induced plasticity. Here, post hoc analyses show that after sufficient sleep, LTD-like plasticity was induced (decreased MEP amplitudes) via cathodal tDCS at 10, 15, and 20 min timepoints compared to baseline MEP. The MEP amplitudes at 10 and 15 min timepoints were significantly different from respective timepoints in the sham condition. Importantly, the MEP decrease was significantly larger at all timepoints when compared to MEP size after sleep deprivation. Sleep deprivation, interestingly, *reversed* the inhibitory LTD-like aftereffects of cathodal stimulation into excitatory LTP-like aftereffects. Specifically, sleep deprivation led to an increase of MEP amplitudes (LTP-like) at all timepoints when compared to the baseline, against the sham intervention, and compared to the respective timepoints after sufficient sleep (Fig. 3c,d). This excitability-enhancing effect of cathodal stimulation was longer-lasting too, as shown by a sustained MEP amplitude enhancement at the 30 the min timepoint when MEP amplitudes are expected to be back at baseline levels.

Fig. 3. LTP/LTD-like plasticity induction after sufficient sleep vs sleep deprivation. **a,** Cortical excitability alterations after inducing LTP-like plasticity with anodal tDCS sleep conditions. Post hoc comparisons (Bonferroni-corrected) of MEP amplitudes to respective baseline values, the sham condition, and sleep conditions are marked by symbols in the figures. **b,** Cortical excitability alterations after inducing LTD-like plasticity with cathodal tDCS under sleep conditions. Sham stimulation after both, sufficient sleep and sleep deprivation did not induce any significant change in cortical excitability. Filled symbols indicate a significant difference of cortical excitability against the respective baseline values. The black asterisks [*] indicate a significant difference between the real vs sham tDCS conditions, and the red asterisks [*] indicate a significant difference between respective timepoints of tDCS condition after sufficient sleep vs sleep deprivation. All error bars represent the standard error of means (s.e.m.). **c,d,** Individual mean MEPs variability obtained from tDCS conditions after sufficient sleep and sleep deprivation. The X-axis represents timepoint (Bl, 0, 5, 10, 15, 20, 25, 30 min) and the Y-axis represents mean MEP amplitudes. *n*=30 (15 per group). *Note*: MEP = motor-evoked potential; tDCS = transcranial direct current stimulation.

216   **Table 1**. Mixed-model ANOVA results for the effect of tDCS on MEP amplitudes after sufficient sleep

217   and sleep deprivation

Factor	*df*	*F*	*p*	ηp^2
group	1	11.74	**0.002**	0.296
sleep condition	1	0.735	0.399	0.026
stimulation state	1	37.09	**<0.001**	0.570

timepoint	4.90	5.76	**<0.001**	0.171
sleep condition×group	1	61.97	**<0.001**	0.689
stimulation state×group	1	3.600	0.068	0.114
timepoint×group	4.90	6.48	**<0.001**	0.188
sleep condition×timepoint	5.18	0.445	0.822	0.016
stimulation state×timepoint	5.08	2.91	**0.015**	0.094
sleep condition×stimulation state	1	5.28	**0.029**	0.159
sleep condition×stimulation state×group	1	57.99	**<0.001**	0.674
sleep condition×timepoint×group	5.18	9.40	**<0.001**	0.251
stimulation state×timepoint×group	5.08	3.68	**0.003**	0.116
sleep condition×stimulation state×timepoint	5.35	1.09	0.369	0.037
group×sleep condition×stimulation state×timepoint	5.35	12.14	**<0.001**	0.301

218 *Note*: tDCS = transcranial direct current stimulation; MEP = motor-evoked potentials.
219 Significant effects are marked in **bold** (where $P<0.05$), $n = 30$ (15 per group).
220

2.3. Electrophysiological evidence of upscaled cortical excitability and saturated synaptic potentiation

So far, we found that sleep deprivation upscales cortical excitability, prevents induction of

LTP-like plasticity, presumably due to saturated synaptic potentiation, and converts LTD-like

plasticity to the LTP-like state. Previous studies in animals and humans have shown that EEG

theta activity is a marker for homeostatic sleep pressure, net synaptic potentiation and

increased cortical excitability due to sleep pressure (Vyazovskiy et al., 2008; Kuhn et al.,

2016). In line with this, we investigated how sleep deprivation affects resting-state brain

oscillations at the theta band (4-7 Hz), beta band (15-30 Hz) as another marker for cortical

excitability, vigilance and arousal (Eoh et al., 2005; Fischer et al., 2008) and alpha band (8-14

Hz) which is important for cognition (e.g. memory, attention) (Klimesch, 2012). To this end,

we analyzed EEG spectral power at mid-central electrodes (Fz, Cz, Pz) using a 3×2 mixed

ANOVA. For theta activity, a significant main effect of location ($F_{1.86}=14.52$, $p<0.001$;

$\eta p^2=0.34$) and sleep condition ($F_1=13.07$, $p<0.001$; $\eta p^2=0.31$) but not their interaction was

observed indicating that theta oscillations at all target electrodes were similarly affected.

Bonferroni-corrected post hoc tests revealed theta oscillations, grand averaged at mid-central

electrodes, were significantly *increased* after sleep deprivation ($p<0.001$) (Fig. 4a). Analyses

of beta spectral power showed a significant interaction of location×sleep condition

($F_{1.97}=5.66$, $p=0.006$; $\eta p^2=0.16$) and the main effect of sleep condition ($F_1=7.25$, $p=0.012$;

$\eta p^2=0.20$) as well. Beta oscillations, grand averaged at mid-central electrodes, were

significantly *increased* after sleep deprivation ($p=0.012$) (Fig. 4b). Finally, for the alpha band,

the main effects of location ($F_{3.15}=12.94$, $p<0.001$; $\eta p^2=0.31$) and sleep condition ($F_1=8.22$,

$p=0.008$; $\eta p^2=0.22$) but not their interaction were significant. Alpha oscillations, grand

averaged at mid-central electrodes, were significantly *decreased* after sleep deprivation

(*p*=0.008) (Fig. 4c). These electrophysiological data support findings from the previous

sections that sleep deprivation upscales cortical excitability and leads to synaptic saturation.

Fig. 4. Resting-state theta, beta and alpha oscillations at electrodes Fz, Cz, and Pz. **a,** Theta band activity was significantly *higher* after sleep deprivation vs sufficient sleep condition (t_{Fz}=4.61, *p*<0.001; t_{Cz}=2.24, *p*=0.034; t_{Pz}=2.59, *p*=0.015). **b,** Beta band activity was significantly *higher* at electrodes Fz and Cz after sleep deprivation compared with the sufficient sleep condition (t_{Fz}=3.25, *p*=0.003; t_{Cz}=2.66, *p*= 0.012). **c**, Alpha band activity was significantly *lower* at all electrodes after sleep deprivation vs the sufficient sleep condition (t_{Fz}=2.39, *p*=0.023; t_{Cz}=2.65, *p*=0.013; t_{Pz}=2.92, *p*=0.007). Data of one participant was excluded due to excessive noise. All pairwise comparisons were calculated via *post hoc* Student's t-tests (paired, *p*<0.05). *n*=29. Error bars represent s.e.m. ns = nonsignificant; Asterisks [*] indicate significant differences. Boxes indicate the interquartile range that contains 50% of values (range from the 25th to the 75th percentile) and whiskers show the 1 to 99 percentiles.

2.4. Sleep deprivation compromises learning, memory formation, and cognitive

performance

LTP and LTD are the primary mechanisms mediating learning and memory. Concentration of

GABA (Kolasinski et al., 2019) and glutamate (Stagg, 2014) is important for motor learning

and synaptic strengthening as well. Results of the resting-EEG data also showed decreased

alpha activity which is critically involved in cognition (e.g. memory, attention) (Klimesch,

2012). Showing these converging effects of sleep deprivation on brain physiology, we were

interested in determining how sleep deprivation affects sequence learning and cognitive

functions as behavioral indices of cortical excitability and neuroplasticity. To this end, we

measured motor sequence learning using the serial reaction time task (SRTT), working

memory with a 3-back letter task, and attentional functioning with the Stroop and AX

continuous performance test (AX-CPT). Electrophysiological correlates of task performance

(e.g. ERP) were measured as well (see Methods).

Motor sequence learning.   The differences in the standardized reaction time (RT) of block

5 vs 6, indicative of learning *acquisition*, and block 6 vs 7, indicative of learning *retention*,

were analyzed with a 3 (block)×2 (sleep condition) repeated-measures ANOVA. The results

showed a significant interaction of block×sleep condition ($F_{1.95}=7.03$, $p=0.002$, $\eta p^2=0.19$) and

the main effects of sleep condition ($F_1=21.47$, $p<0.001$, $\eta p^2=0.42$) and block ($F_{1.93}=41.63$,

$p<0.001$, $\eta p^2=0.58$) as well. Post hoc comparisons revealed a significantly larger RT

difference at blocks 6-5 and blocks 6-7 only after sufficient sleep and lower committed errors

(Fig. 5a,b). Absolute RT, error rate, and RT variability were similarly affected by sleep

deprivation (Fig. S1). Next, we explored electrophysiological correlates of motor learning.

The P300 component is evoked in response to stimuli of low probability and stimulus

sequence (Squires et al., 1976). We expected a higher-amplitude P300 component, when the

learned sequence of stimuli is violated (at block 6), after having sufficient sleep. We analyzed

the P-300 amplitudes (250-500 ms) in blocks (5-7) and the results revealed only a significant

main effect of block on the amplitude at electrodes Pz ($F_{1.78}=15.88$, $p<0.001$, $\eta p^2=0.35$) and

P3 ($F_{1.90}=6.63$, $p=0.003$, $\eta p^2=0.18$), which are among regions of interest in this task. The P-

300 amplitude in block 6 vs blocks 5 and 7 was significantly larger at electrode Pz after both

sleep conditions, but respective comparisons between blocks at electrode P3 were significant

only after sufficient sleep (Fig. 5c,d). A similar trend was observed for the other electrodes of

interest (Fig. S2).

Working memory.   For working memory performance, the ANOVA results revealed a

significant main effect of sleep condition on the N-back hits ($F_1=12.36$, $p<0.001$; $\eta p^2=0.30$),

and *d* prime ($F_1=11.77$, $p=0.002$; $\eta p^2=0.278$) as the primary outcomes of interest, but not on

RT of hits ($F_1=0.01$, $p=0.894$). Post hoc analyses showed significantly enhanced WM

performance with significantly more RT variability after sufficient sleep, which could be due

to an accuracy-RT trade-off (Fig. 5e,f). Furthermore, the P300 ERP component was

investigated across sleep conditions. No significant main effect of sleep condition was

observed on the P300 component at electrodes Fz ($F_1=1.66$, $p=0.208$), Pz ($F_1=0.88$, $p=0.364$),

and Cz ($F_1=1.01$, $p=0.310$). Yet, a trend-wise increase of the P300 amplitude was identified at

electrodes Fz and Cz after sufficient sleep compared to sleep deprivation (Fig. 5g,h).

Selective attention.   The RT difference of congruent and incongruent trials in the Stroop

task was analyzed with a 2 (sleep condition) ×2 (congruency) factorial ANOVA. Sleep

condition ($F_1=23.77$, $p<0.001$; $\eta p^2=0.45$) and congruency ($F_1=106.15$, $p<0.001$; $\eta p^2=0.78$)

had significant effects on Stroop interference but they did not interact ($F_1=0.69$, $p=0.413$).

Post hoc comparisons revealed a significant Stroop effect (slower RT of incongruent trials vs

congruent trials) after both, sufficient sleep ($t=3.01$, $p=0.009$) and sleep deprivation ($t=3.47$,

$p<0.001$). However, the interference effect was significantly stronger after sleep deprivation

vs sufficient sleep in overall trials ($t=2.82$, $p=0.015$), congruent trials ($t=2.71$, $p=0.021$) and

incongruent trials ($t=3.19$, $p=0.005$) (Fig. 5i,j). Reduced Stroop effects are associated with

higher N200 and N450 amplitudes, which are indicative of higher selective attention and

better detection of conflicting stimuli. We analyzed these ERP components too. The results of

the 2 (congruency) ×2 (sleep condition) ANOVA showed a significant interaction of sleep

condition×congruency on the N200 ($F_1=3.90$, $p=0.05$; $\eta p^2=0.12$) and N450 amplitudes

($F_1=6.43$, $p=0.017$; $\eta p^2=0.19$) for the electrode Fz. The main effect of sleep condition was not

significant, and the main effect of congruency was significant only for N450 component

($F_1=4.29$, $p=0.045$; $\eta p^2=0.14$). Sleep deprivation was related to a significantly *smaller* N200

amplitude, but not N450, for the incongruent trials only, at electrode Fz ($t=2.75$, $p=0.010$).

Both N200 and N450 amplitudes of incongruent trials were significantly *larger* compared to

congruent trials after sufficient sleep ($t_{N200}=2.51$, $p=0.018$; $t_{N450}=3.63$, $p=0.001$), but not sleep

deprivation ($t_{N200}=0.24$, $p=0.810$; $t_{N450}=0.48$, $p=0.634$), indicating that conflict detection was

more clearly processed after having sufficient sleep (Fig. 5k,l). Results of the electrode Cz can

be found in supplementary materials (Fig. S3b).

Sustained attention.   The AX-CPT was used for measuring sustained attention. We found a

significant main effect of sleep condition on performance accuracy ($F_1=28.12$, $p<0.001$;

$\eta p^2=0.49$) as the primary outcome of interest, RT of hit trials ($F_1=10.85$, $p=0.003$; $\eta p^2=0.27$),

and variability of RT ($F_1=9.85$, $p=0.004$; $\eta p^2=0.25$). Participants responded significantly less

accurately, with slower RT, and more variable RT after sleep deprivation compared to

sufficient sleep (Fig 5m,n). Here again, the P300 serves as an attentional index of the target

stimulus and memory storage. Analysis of this ERP component showed a significant main

effect of the sleep condition on the P300 component at electrodes Fz ($F_1=20.25$, $p<0.001$;

$\eta p^2=0.43$), Cz ($F_1=20.57$, $p<0.001$; $\eta p^2=0.44$), but not Pz ($F_1=0.72$, $p=0.402$). Post hoc

analyses indicated that sleep deprivation was related to a significantly smaller P300 amplitude

in these (Fig 5o,p), and other electrodes of interest (Fig S3c).

Relevant correlations.   We found several relevant correlations between behavioral

learning *vs* plasticity and cognition *vs* excitability indices. LTP-like plasticity effects after

sufficient sleep were correlated with better sequence learning acquisition ($r=-0.558$, $p=0.031$)

and retention ($r=-0.734$, $p=0.002$). Enhanced working memory and sustained attention after

    having sufficient sleep were also correlated with higher cortical facilitation and lower cortical

    inhibition (supplementary materials).

Fig. 5. Impact of sleep deprivation on sequence learning, working memory and attention. **a,** The RT difference of BL6-5 (learning acquisition) and BL6-7 (learning retention) were significant only after sufficient sleep (BL6-5: t=3.73, p<0.001; BL6-7: t=2.95, p=0.003) but not sleep deprivation (BL6-5: t=1.67, p=0.094; BL6-7: t=0.95, p=0.337). **b,** Performance was more erroneous after sleep deprivation. Asterisks [*] represent significant differences between learning block RTs [BL 6-5, BL 6-7]. n=30. **c,** For both P3 and C2 electrodes, the P300 amplitude (300-600 ms) was significantly larger in block 6 vs BL5 and 7 only after sufficient sleep (P3: t_{6-5}=3.50, p<0.001, t_{6-7}=3.30, p=0.003; C2: t_{6-5}=2.74, p=0.010, t_{6-7}=2.64, p=0.013) marked by the filled symbol. n=30. **d,** P300 schematic at electrode P3. **e,** Participants had more correct responses (t=3.56, p<0.001) and a higher d index (t=3.43, p=0.002) after having sufficient sleep vs sleep deprivation. **f,** Performance speed was not significantly different but was more variable after sufficient sleep. n=30. **g,** The P300 amplitude (300-600 ms) did not significantly differ across sleep conditions at electrodes Fz, Pz, and Cz. n=29. **h,** P300 schematic at electrode Cz. **i,** RT of the congruent, incongruent, and overall trials in the Stroop task was significantly slower after sleep deprivation. **j,** Participants displayed a significantly stronger Stroop interference effect (RT$_{incongruent}$-RT$_{congruent}$) after sleep deprivation vs sufficient sleep (t=2.63, p=0.009). n=29. **k,** The N200 at electrode Fz was significantly

larger for incongruent trials, but not congruent trials, after sufficient sleep vs sleep deprivation. Both N200 and N450 were significantly larger for incongruent vs congruent trials only after sufficient sleep. **l**, N200-N450 schematic at electrode Fz. **m,n**, Participants were less accurate in identifying AX trials (t=5.30, p<0.001), had slower RT (t=3.29, p=0.003), and showed a larger variability of RT (t=3.13, p=0.004) after sleep deprivation vs sufficient sleep. **o**, The P300 amplitude was significantly larger after sufficient sleep at electrodes Fz and Cz. n=27. **p**, P300 schematic at electrode Cz. All Pairwise comparisons were calculated via post hoc Student's t-tests (paired, p<0.05). Error bars represent s.e.m. ns=nonsignificant; Asterisks [*] indicate significant differences. Boxes indicate interquartile range that contains 50% of values (25th-75th) and whiskers show 1 to 99 percentiles. *Note*: In working memory ERP analysis (n=1), Stroop behavioral (n=1) and ERP analyses (n=2) and AX-CPT ERP analyses (n=3), the data of some participants were excluded from the analysis for excessive noise. BL = block, RT = reaction time. See also Figure S1-S3.

2.5. Demographics, subjective sleepiness and cortisol level

The mean age of participants was 24.62±4.16 years. 50 percent of the participants were male.

Age and gender did not correlate with the dependent variables discussed in previous sections.

Ratings of sleepiness and alertness at 9:00 AM showed a significantly higher sleep pressure,

as measured by the Karolinska and Stanford Sleepiness Scales (KSS, SSS) after sleep

deprivation (Mean$_{KSS}$=7.10±1.76, Mean$_{SSS}$=4.90±1.32), compared to the sufficient sleep

condition (Mean$_{KSS}$=2.96±0.764, Mean$_{SSS}$=2.23±0.62). Sleep condition had a significant

effect on KSS (F_1=159.02, p<0.001; ηp^2=0.84) and SSS (F_1=122.10, p<0.001; ηp^2=0.81)

ratings, and a significantly higher sleep pressure was observed after the sleep deprivation vs

the sufficient sleep. The average levels of cortisol and melatonin were lower after sleep

deprivation vs sufficient sleep (cortisol: 3.51±2.20 vs 4.85±3.23, p=0.05; melatonin

10.50±10.66 vs 16.07±14.94, p=0.16).

3. Discussion

In the last two decades, a strong link has been established between the function of sleep and

synaptic strength (Tononi and Cirelli, 2014; de Vivo et al., 2017). These studies, mostly in

nonhuman animals, show the critical role of sleep in neuronal plasticity (mostly LTP) and

thereby learning and memory. Whether and how the need for sleep affects relevant parameters

of human brain physiology including neuroplasticity (both LTP and LTD), cortical

excitability, and their behavioral counterparts (learning, memory, cognition) is not fully

investigated. We found that sleep deprivation upscales cortical excitability. It furthermore has

a nonlinear effect on neuroplasticity inducibility and saturates induction of LTP-like plasticity

while converts induced LTD-like to LTP-like plasticity. These physiological changes are

associated with compromised sequence learning, working memory, and attentional

functioning and their electrophysiological correlates.

Changes of cortical excitability following sleep deprivation were observed in neurotransmitter

systems involved in intracortical facilitation and inhibition (Chen, 2000; Di Lazzaro et al.,

2000; Di Lazzaro et al., 2005a), and brain oscillations. Specifically, glutamate-related cortical

facilitation (measured by ICF) was upscaled while GABA- and acetylcholine-related cortical

inhibition (measured by SICI, I-wave, SAI) were disinhibited or reversed after sleep

deprivation. These sleep-dependent changes of cortical excitability are in line with the

synaptic homeostasis hypothesis and resultant changes in the brain. According to this

hypothesis, synaptic strength is increasingly potentiated during wakefulness and saturated, if

wakefulness is extended (Tononi and Cirelli, 2003; Vyazovskiy et al., 2008). Animal studies

have shown that at this state, molecular and electrophysiological markers of synaptic strength

increase including AMPA receptors, cortical spine density, slope and amplitude of cortical

evoked responses, and even the size and number of synapses (Tononi and Cirelli, 2003;

Vyazovskiy et al., 2008; Bushey et al., 2011; Maret et al., 2011). Our findings also

complement those of human studies that show an increase and decrease of intracortical

facilitation and inhibition respectively (Huber et al., 2012; Kuhn et al., 2016), and increase of

theta and beta band activity, markers of homeostatic sleep pressure, after sleep deprivation

(Finelli et al., 2000; Vyazovskiy et al., 2008; Kuhn et al., 2016). Together, these findings

argue for an upscaled state of cortical excitability mediated by the need for sleep. Cortical

excitability is a basic physiological response of cortical neurons to an input and is, therefore, a

fundamental aspect of human brain functioning, neuroplasticity, and cognition (Kuhn et al.,

2016; Ly et al., 2016; Gaggioni et al., 2019). Changes in brain excitability will therefore

affect neuroplasticity and cognition.

Previous studies in animals (Campbell et al., 2002; de Vivo et al., 2017) and humans (Kuhn et

al., 2016) have shown that synaptic LTP is inevitably affected by sleep deprivation. LTD

changes are not, however, fully understood in the human brain. We non-invasively induced

LTP-, and LTD-like plasticity by tDCS to investigate how a synaptic saturation state due to

sleep deprivation affects the inducibility of LTP- and LTD-like plasticity. For LTP-like

plasticity, the results demonstrate diminished plasticity in the motor cortex after sleep

deprivation which can be explained via the synaptic strength perspective. Sleep deprivation

saturates (or upscales) synaptic strength which eventually leads to deficient LTP inducibility.

In contrast, sufficient sleep downscales synaptic strength and makes it responsive to LTP-like

plasticity induction. The presence of anodal tDCS-induced LTP-like plasticity after sufficient

sleep and its diminution after sleep deprivation fits with this model. This is moreover in

agreement with *in vivo* and *in vitro* studies in rats (McDermott et al., 2003; Kopp et al., 2006;

Vyazovskiy et al., 2008), and in line with the results of a recent human study that showed

decreased LTP-like plasticity after sleep deprivation (Kuhn et al., 2016). Regarding LTD, our

results showed that sleep deprivation *reversed* the LTD-like inhibitory effect of cathodal

stimulation into LTP-like excitatory effects which is the first evidence for the effect of sleep

deprivation on LTD-like plasticity induction in humans. This conversion of LTD-like into

LTP-like plasticity is indeed in line with the saturating effect of sleep deprivation on synaptic

strength rather than cortical underactivation.

One mechanism that can explain the diminished LTP-like and conversion of LTD-like to

LTP-like plasticity during sleep deprivation is intracellular calcium concentration (Fig. 6). It

is known that the directionality of plasticity (LTP or LTD) depends on the level of calcium

concentration (Lisman, 2001) with LTP and LTD being linked to higher and lower

intracellular Ca^{2+} concentration respectively. There are, however, certain limit zones for

induction of plasticity. In this line, anodal LTP-like plasticity is linked to largely enhanced

intracellular calcium concentration in animals and humans (Islam et al., 1995; Nitsche et al.,

2005; Biabani et al., 2017), while cathodal tDCS-induced LTD-like plasticity is assumed

linked to lower intracellular Ca^{2+} concentration (Nitsche et al., 2003b). Our excitability results

earlier showed a glutamate-controlled NMDA receptor facilitation during sleep deprivation,

which enhances calcium influx. In this state, applying anodal tDCS, which enhances calcium

concentration, can then result in abolishment of LTP-induction due to calcium overflow

(Lisman, 2001; Misonou et al., 2004; Grundey et al., 2018). On the other hand, inducing

LTD-like plasticity might downregulate upscaled cortical excitability at the network level,

and open some synaptic space for LTP-related plasticity induction. This is in line with

previous studies that show calcium enhancement can convert the direction of cathodal-

induced LTD-like to LTP-like plasticity (Batsikadze et al., 2013).

Fig 6: **Proposed mechanism for plasticity induction**. The intracellular calcium concentration (x-axis) determines directionality of plasticity (Lisman, 2001). It can be assumed that intracellular calcium concentration under no sleep pressure (sufficient sleep) is at an optimal level leading to stronger tDCS-induced LTP/D-like plasticity. Under sleep deprivation, LTP-like plasticity induction via anodal tDCS is prevented due to calcium overflow, and LTD-like plasticity via cathodal tDCS is converted to LTP-like plasticity possibly via (a) enhanced baseline calcium (due to upscaled excitability) which makes minor calcium increase obtained from cathodal stimulation to be sufficient to induce LTP-like plasticity and (b) gradual downregulating upscaled cortical excitability and opening some synaptic space for LTP-related plasticity induction.

Changes in synaptic strength are the primary mechanisms mediating learning and memory

(Feldman, 2009). The important question here is whether upscaled excitability and saturated

plasticity after sleep derivation, influence learning, memory formation and higher-order

cognition that depend on brain plasticity and cortical excitability (Nitsche et al., 2003a; Huang

et al., 2017). The sleep-dependent global synaptic downscaling spares neuronal assemblies

crucially involved in the encoding of information (Niethard and Born, 2019) which results in

an improved signal-to-noise ratio and a renewed capacity for encoding new information

(Kuhn et al., 2016). Accordingly, a better cognitive/behavioural performance is expected after

sleep, compared to sleep deprivation. In this line, our results show enhanced motor sequence

learning after sufficient sleep and impaired sequence learning and retention after sleep

deprivation. Similarly, our results show compromised working memory and attentional

functioning after sleep deprivation. In support of this, we noticed specific alterations of task-

dependent ERP components, such as lower amplitudes of the P300 after sleep deprivation in

the sustained attention, and motor learning tasks, and suppressed N200 and N450 in the

Stroop conflict condition, which measures selective attention and interference control.

To converge our findings, it is important to consider the links between the physiological

parameters of the brain with motor learning, memory formation and cognitive functions. In

sequence learning, a specific link can be made between behavioural learning performance and

plasticity results. The behavioural and electrophysiological markers of motor sequence

learning and retention were associated with facilitated LTP/LTD-like plasticity in the motor

cortex after sufficient sleep, which supports the suggested link between inducibility of

neuroplasticity and learning and memory formation. This observation makes sense, as tDCS-

induced neuroplasticity in the motor cortex and behavioural motor learning share intracortical

mechanisms (Stagg, 2014). For the cognitive functions measured in our study, which do not

require plasticity, cortical excitability alterations are more relevant. The increase of cortical

excitability parameters and the resultant synaptic saturation following sleep deprivation can

explain the respective cognitive performance decline.

The timing of the experimental measurements should be taken into account for interpreting

the results. Although we strictly controlled for external and environmental factors that could

affect cortical excitability, especially in the sleep-wake cycle, the measurement time of

cognitive functions, cortical excitability monitoring, and neuroplasticity induction differed by

a few hours (24, 26, 27 h sleep deprivation respectively). By fixing the time of the specific

protocols, we however controlled for potential confounding effects of different measurement

times *within* a specific protocol. Nevertheless, for the between protocol comparisons, this

procedure could have gradually affected subjective sleepiness, and probably brain physiology

(Kuhn et al., 2016). Furthermore, the physiological measures were based on the motor system

and indirect measures of the involved neurotransmitters, and the cognitive tasks under study

are more closely related to prefrontal regions, with the exception of the motor learning task.

In conclusion, the results of this experiment show a negative impact of sleep deprivation on

learning and cognitive processes, including learning, memory formation and attentional

functions, as well as specific alterations of brain physiology underlying these cognitive

processes and their electrophysiological correlates. These findings complement the current

knowledge about the critical role of sleep in neuroplasticity and cognition, especially in

humans.

4. Materials and Methods

4.1. Participants

Thirty healthy adult volunteers (15 females, mean age = 24.44±3.93) who met the inclusion criteria were recruited from the TU Dortmund University, Ruhr-University Bochum, and the surrounding community. Power analysis showed that for a medium effect size (Minarik et al., 2016) (f = 0.35 equivalent to partial eta squared = 0.10), a minimum of 30 subjects are required to achieve 95% power at an alpha of 0.05 for the primary applied statistical test. All participants had to be right-handed, non-smokers, with a regular sleep-wake pattern (determined by sleep diary) and underwent a medical screening to verify no history of neurological diseases, epilepsy or seizures, central nervous system-acting medication, metal implants, and current pregnancy. As gender and age may affect the sleep-wake cycle and brain excitability, we balanced participants' gender and kept the age range to early adulthood. Each participant took part in a test TMS session to become acquainted with experiencing stimulation and the study protocol. Female subjects were not examined during the menstrual period to ensure hormonal changes would not interfere with the measurements. This study conformed to the Declaration of Helsinki guidelines and was approved by the Institutional Review Board of the xx. Participants gave informed consent and received monetary compensation.

4.2. Study design and course of study

This was a randomized, sham-controlled, cross-over study. Participants completed a sleep diary two weeks before the beginning of the experiment and attended two experimental sessions after having "sufficient sleep [23:00-8:00]" and "sleep deprivation [23:00-8:00]" overnight in counterbalanced order. The experimental sessions took place at a fixed time in the morning (9:00 AM). The interval between the sessions was at least 2 weeks. In each session, participants underwent the same experimental protocol. Each experimental session began with saliva sampling at 8:45 AM, followed by behavioural and cognitive task performance during EEG recording (1.5 h), followed by cortical excitability monitoring sessions (1 h), and finally followed by neuroplasticity induction with active and sham tDCS (1.5 h). The reason for starting the session with behavioural/EEG measures was to take advantage of 1 hour between 8-9 AM for preparing the EEG cap, which took about 45 min on average. The order of measurements (behavioural/EEG, cortical excitability, tDCS) was

identical across participants, to keep the start time of each measurement and the number of

hours deprived from sleep fixed for each measure.

4.3. Sleep conditions

In the "sufficient sleep" condition, participants had to go to bed at around 23:00 and have at

least 8 hours of uninterrupted sleep. The experiment was scheduled to start at 9:00.

Participants were refrained from drinking alcohol and coffee 12 h before sleep time and

afterwards until the experiment. In the case of poor sleep quality for any reason (measured by

sleepiness rating scales) or unregular sleep pattern (sleep onset, wake-up time) informed from

the sleep diary (±2.5 h), the scheduled session was canceled and postponed until when

sufficient sleepcondition requirements were met. In the "sleep deprivation" condition,

participants spent the night in a specific lounge at the local institute where the data was

collected. The lounge was equipped with an unrecording live camera and prepared for

participants' stay overnight. Participants spent all night awake in the lounge (23:00-8:00) and

were supervised by a scientific staff member. Additionally, their sleep-waking status was

recorded via a wrist-worn Actigraphy (MotionWatch 8.0, CamTech, Cambridge, UK). Food

and drinks were provided (the consumption of coffee, caffeine-containing soft drinks, black

tea, and alcohol was not allowed), watching TV programs, reading and working on the

computer were also allowed. Participants were prevented from any sleep-related activities

during the night, such as lying down, or closing their eyes for a prolonged time. They were

refrained from drinking caffeine-containing drinks and sleeping or taking a nap from the

afternoon before they joined the sleep deprivation session at 23:00. Before starting each

experimental session, subjective sleepiness of participants and their alertness were evaluated

with the Karolinska sleepiness scale (KSS) (Akerstedt and Gillberg, 1990) and the Stanford

Sleepiness Scale (SSS) (Hoddes et al., 1972). All external factors that could affect circadian

rhythmicity such as light and food intake were controlled during the experiment.

4.4. Determination of cortisol and melatonin from saliva

Saliva samples were collected using salivettes (Sarstedt AG & Co. KG, Germany). After

centrifugation at 3,000 xg for 2 min at 4 °C, samples were aliquoted at 500 µL each and

stored at -20 °C until measurement. Cortisol and melatonin were measured using Cortisol

Saliva ELISA and Melatonin direct Saliva ELISA (both IBL International GmbH, Germany)

according to manufacturer´s instructions. Duplicate measurements were conducted for all

samples in a range from 0.05 µg/mL - 30 µg/mL for cortisol using 50 µL of sample and from

1.0 pg/mL – 50 pg/mL for melatonin using 100 µL of sample. In case of values exceeding the

measurement range for melatonin samples were diluted 1:10 and remeasured.

4.5. Cortical excitability

Different protocols of single-pulse and paired-pulse TMS were used to monitor cortico-spinal

and intracortical excitability in the motor cortex. These protocols included: resting motor

threshold (RMT), active motor threshold (AMT), I-O curve, short intracortical inhibition

and facilitation (SICI-ICF), intracortical I-wave facilitation, and Short-latency afferent

inhibition (SAI). RMT, AMT and I-O curve examine corticospinal excitability, SICI-ICF

measures both, intracortical facilitation and inhibition, and intracortical I-wave facilitation

and SAI are measures of intracortical inhibition of the human motor cortex (Kujirai et al.,

1993; Chen, 2000; Di Lazzaro et al., 2000).

4.5.1. Single-pulse MEP, resting and active motor threshold

Single-pulse biphasic TMS at 0.25 Hz ± 10% (random) was through a figure-of-eight

magnetic coil (diameter of one winding, 70 mm; peak magnetic field, 2T) held 45° to the

midline and applied over the left primary motor cortex. Surface motor-evoked potentials

(MEPs) were recorded from the right abductor digiti minimi muscle (ADM) with gold cup

electrodes in a belly-tendon montage. RMT was examined using the TMS Motor Threshold

Assessment Tool (MTAT 2.0, http://www.clinicalresearcher.org/software.htm) (Awiszus,

2003) and was determined as the lowest stimulator intensity required to evoke a peak-to-peak

MEP of 50 µV in the relaxed ADM muscle in at least five out of ten consecutive trials. The

AMT was determined as the lowest stimulator intensity required to elicit MEP response of

∼200–300 µV during moderate tonic contraction of the right ADM muscle (∼20% of the

maximum muscle strength) (Rothwell et al., 1999) in at least three of six consecutive trials.

4.5.2. Input-output curve (I-O curve)

The I-O curve is a TMS single-pulse protocol that reflects excitability of cortico-spinal

neurons. It is modulated by glutamatergic activity and refers to the increase of MEP

amplitudes with increasing TMS intensity (Chen, 2000). The slope of the recruitment curve

increases at higher TMS intensities with higher glutamatergic and adrenergic transmission and

decreases by drugs that enhance effects of GABA (Chen, 2000; Paulus et al., 2008). In the I-O

curve protocol, MEP-amplitudes in the relaxed right ADM muscle were measured in four

blocks with different stimulus intensities (100%, 110%, 130%, and 150% RMT)(Batsikadze

et al., 2013), each block with 15 pulses, and a mean (MEP amplitudes) was calculated for
each intensity.

4.5.3. Short latency intracortical inhibition and intracortical facilitation (SICI-ICF)

The SICI-ICF is a TMS paired-pulse protocol for monitoring of GABAergic-mediated cortical
inhibition and the glutamate-mediated cortical facilitation (Chen, 2000). In this protocol, a
subthreshold conditioning stimulus (determined as 70% of AMT) is followed by a
suprathreshold test stimulus which was adjusted to evoke a baseline MEP of ∼1 mV. The
paired stimuli are presented in inter-stimulus intervals (ISI) of 2, 3, 5, 10 and 15 ms(Kujirai et
al., 1993). ISIs of 2 and 3 ms represent short-latency intracortical inhibition (SICI) and have
inhibitory effects on test pulse MEP amplitudes, and ISIs of 10 and 15 ms represent
intracortical facilitation (ICF) and have enhancing effects on single pulse TMS-elicited MEP
amplitudes (Kujirai et al., 1993; Lazzaro et al., 1998; Lazzaro et al., 2003). The stimuli
(subthreshold and suprathreshold stimuli) were organized in blocks in which each ISI and one
single test stimulus were applied once in pseudorandomized order. Each block was repeated
15 times, which resulted in a total of 90 single-pulse or paired-pulse MEP per session. The
exact interval between the paired pulses was randomized (4 ± 0.4 s).

4.5.4. Short-interval intracortical I-Wave facilitation

This TMS protocol is based on I (indirect) waves which refer to high-frequency repetitive
discharges of corticospinal neurons produced by single-pulse stimulation of the motor cortex
(Di Lazzaro et al., 2012) (for a detailed review see (Ziemann et al., 1998; Di Lazzaro et al.,
2012)). In this protocol, two successive stimuli (supra- and subthreshold) are separated by
short ISIs, but this protocol involves a suprathreshold first stimulus and a subthreshold second
stimulus (Ziemann et al., 1998). The ISIs range from 1.1 ms to 4.5 ms latency and are
presented in pseudorandomized order. We grouped ISIs to early (mean MEP at ISIs 1.1, 1.3,
1.5 ms), middle (mean MEP at ISIs 2.3, 2.5, 2.7, 2.9 ms), and late (mean MEP at ISIs 4.1, 4.3,
4.5 ms) epochs. The intensity of the first conditioning suprathreshold stimulus (S1) is adjusted
to produce a baseline MEP of ∼1 mV when given alone and is followed by a second
subthreshold stimulus (S2) that was set to 70 % of RMT) (Batsikadze et al., 2013). For each
ISI, 15 pulses were recorded. Another 15 pulses were recorded for the control MEPs, in which
the suprathreshold stimulus (S1) was given alone and adjusted to achieve a baseline MEP of
∼1 mV. The pairs of stimuli were organized in blocks in which each ISI and one test pulse
was represented once and were pseudorandomized. This TMS paired-pulse protocol (a first

suprathreshold stimulus and a second subthreshold stimulus) has facilitatory effects on MEP

peaks (Ziemann et al., 1998) that occur at ISIs of about 1.3, 2.6, and 4.2 ms. This effect is

suggested to be produced as a result of elicited I-waves (indirect waves: descending volleys

produced by indirect activation of pyramidal tract neurons via presynaptic neurons) by the

subthreshold S2 and is controlled by GABA-related neural circuites (Ziemann et al., 1998;

Hanajima et al., 2002; Paulus et al., 2008)

4.5.5. Short-latency afferent inhibition (SAI)

SAI is a TMS protocol coupled with peripheral nerve stimulation and is based on the concept

that peripheral somatosensory inputs have an inhibitory effect on motor cortex excitability at

short intervals (e.g. 20-40 ms) (Di Lazzaro et al., 2005b). SAI has been linked with

cholinergic(Di Lazzaro et al., 2000) and GABAergic systems (Di Lazzaro et al., 2005a) at the

cortical level. In this protocol, single-pulse TMS serves as test stimulus and is adjusted to

evoke a MEP response with a peak-to-peak amplitude of approximately 1 mV. The

conditioning afferent stimuli were single pulses (200 µs) of electrical stimulation applied to

the right ulnar nerve at the wrist level (cathode proximal) through bipolar electrodes

connected to a Digitimer D185 stimulator (Digitimer Ltd., Welwyn Garden City, UK). The

conditioning afferent stimuli were applied with an intensity of ~ 2.5–3 times of perceptual

threshold adjusted to evoke a minimal visible twitch of the thenar muscles (Di Lazzaro et al.,

2000), followed by a single TMS pulse (test stimulus) applied over the motor cortical

representation of the right ADM. The stimuli were applied in blocks containing the test

stimulus alone (control condition) and two paired-stimuli blocks with ISIs of 20 and 40 ms in

pseudorandomized order. Each block was repeated 20 times, resulting in a total of 60 trials.

4.5.6. Experimental procedure

Cortical excitability was monitored right after the behavioural/EEG measurements at a same

fixed time in both, *sufficient sleep*and *sleep deprivation* sessions. In each session, participants

were seated comfortably in a reclining chair, with a pillow resting under the right arm and a

vacuum-pillow around the neck to prevent head movement. First, the hotspot (the coil

position over the primary motor area that produces the largest MEP in the right ADM with a

given medium TMS intensity) was identified with TMS and marked with a water-proof pen.

The stimulation intensity was then adjusted to evoke MEPs with a peak-to-peak amplitude of

an average of 1 mV. Following this step, RMT and AMT were obtained. A 10 min break was

allowed after recording AMT in order to avoid an effect of muscle contraction on the next

measurements. After the break, the following TMS protocols were measured to monitor

cortical excitability: SAI, SICI-ICF, I-wave facilitation, and I/O curve. The order of measures

was randomized except for the I/O curve, which was always the last measure as it required

applying high intensities of TMS, which might induce after-effects on excitability. In the case

of single test pulse-generated MEP alterations of >20% during the session in the double pulse

conditions, stimulation intensities were adjusted (Kuo et al., 2017). Participants were visually

monitored to prevent them from closing the eyes due to sleep pressure. Each cortical

excitability session took 60-70 min. All TMS protocols were conducted with a PowerMag

magnetic stimulator (Mag & More, Munich, Germany) through a figure-of-eight magnetic

coil (diameter of one winding, 70 mm; peak magnetic field, 2T), held 45° to the midline and

applied over the left primary motor cortex.

4.6. Neuroplasticity

Electrical direct current was applied through a pair of saline-soaked surface sponge electrodes

(35 cm^2) and delivered through a battery-driven constant current stimulator (neuroConn

GmbH, Ilmenau, Germany). The target electrode was fixed over the motor-cortical

representation area of the right ADM as identified by TMS, and the reference electrode was

placed over the contralateral supraorbital area. The distance on the scalp between the edges of

the electrodes was kept at a minimum of 6 cm to reduce shunting of current through the scalp

(Nitsche et al., 2007). Based on the randomized condition, anodal, cathodal, or sham tDCS

with 1 mA intensity were applied for 7 min with 15 s ramp up/down at the beginning and end

of stimulation. For the sham condition, stimulation was delivered for 30 s, with a 30 s ramp

up and down. Using this procedure, participants are not able to distinguish between real and

sham tDCS (Ambrus et al., 2012). TMS intensity was set to evoke MEPs of approximately 1-

mV peak-to-peak amplitude and single-pulse MEPs were then obtained.

4.6.1. Experimental procedure

Participants were randomly assigned to the anodal tDCS (*N*=15) or cathodal tDCS (*N*=15)

groups. Each participant attended four sessions of tDCS (active and sham tDCS after

"sufficient sleep" and "sleep deprivation") in a counterbalanced order. Order of stimulation

was similar across sessions for each participant. TDCS sessions were conducted after the

cortical excitability measurement at a fixed starting time and took roughly 90 min in total. In

each session, participants were seated comfortably in a reclining chair, with a pillow

positioned under the right arm and a vacuum-pillow around the neck to prevent head

movement. First, baseline cortical excitability was measured by inducing MEPs over the left

M1 representation of the target muscle (right ADM) with a given TMS intensity. The hotspot

region was already identified and marked in the cortical excitability part. Stimulation intensity

was adjusted to reach a peak-to-peak MEP amplitude of 1 mV (SI1mV), which was then used

for the remaining measurements. Following a baseline measurement of 25 MEPs, 7 min of

active (anodal or cathodal, depending on group assignment) or sham stimulation was

delivered. Right after, MEP measurements were conducted immediately in epochs of every 5

min for up to 30 min after tDCS (7 total epochs). Five mine after the last MEP measurement

(30 min following tDCS), the second tDCS intervention (active or sham) started with the

same experimental procedure. Based on the previous works, this tDCS protocol induces

polarity-specific short aftereffects up to 30 min after stimulation (Nitsche and Paulus, 2001),

and thus applying the second intervention after 30 min is feasible. At the end of each session,

participants completed a side effect survey to rate the presence and severity of potential

adverse effects during stimulation and also asked to guess the stimulation intensity they

received (i.e., 0 mA intensity or 1 mA intensity) to evaluate blinding efficacy. The tDCS

intervention including both, active and sham stimulation took around 90 min in each session.

4.7. Behavioural measures

4.7.1. Motor learning. The Serial Reaction Time Task (SRTT) was used to measure implicit

motor learning in participants. Performance on this task is associated with increased activity

and cortical excitability of the motor, premotor and Supplementary motor areas and early

learning affects primarily the primary motor cortex (Honda et al., 1998; Nitsche et al., 2003a;

Schendan et al., 2003). In brief, the SRTT consisted of 8 blocks in which participants should

respond to a visually-cued stimuli sequence on a computer screen with the respective finger

positioned on a keyboard as fast and accurately as possible. Participants are instructed to push

the respective button with the respective finger of the right hand (index finger for Button 1,

middle finger for Button 2, ring finger for Button 3, and little finger for Button 4). In blocks 1

and 6, the sequence of dots followed a pseudorandom order and in the other blocks, the order

of stimuli follows an implicit sequence (e.g. A–B–A–D–B–C–D–A–C–B–D–C). The averaged

RT difference in block 5 (sequence order) vs block 6 (random order) is the primary measure of

motor *learning acquisition* as it indicates response to sequence learning vs sequence learning-

independent performance. The RT difference between block 6 (random order) and block 7

(sequenced order) is suggested to indicate additionally *learning retention*. In addition to RT,

which is the major indicator of *implicit* motor learning, RT variability and accuracy were also

calculated as outcome variables. Participants were not told about the repeating sequence and at

the end of the session, they were asked whether they noticed a sequence and if so, to write the

sequence in order to assess explicit learning of the task. In such a case, the data were excluded

from the final analysis. Two different sequences of the task, with no overlapping parts, and

comparable difficulties, were presented in the two sessions in a counterbalanced order.

4.7.2. Working memory.   A 3-back version letter of the task (Mull and Seyal, 2001) was

used to measure working memory. In this task, participants should indicate whether a letter

presented on the screen (the "target letter") matched the letter previously presented (the "cue"

letter). "Hits" (correct responses) were defined as any letter identical to the one presented three

trials back. Stimuli were pseudorandom sequences of 10 letters (A–J) presented at a fixed

central location on a computer screen. Each letter was visible for 30 ms with a 2000 ms inter-

stimulus interval, making the difficulty level of the task high. The letters were presented in

black on a white background and subtended 2.4 cm (when viewed at 50 cm eye to screen

distance). Participants completed 2 blocks consisting of 44 (practice block) and 143 trials (main

block) respectively, resulting in a total number of 187 trials. A short break (5–20 s) between

blocks was provided to allow participants to rest. Two different versions of the task were

employed in two sessions (sufficient sleepvs sleep deprivation sessions) and condition order

was randomized across participants. Accuracy, *d* prime (the proportion of hits rate minus the

proportion of false alarm rate) and "Hits" reaction times measures were the putcome measures.

The Stroop color-word task. The Stroop interference task is a neuropsychological test

extensively used for measuring selective attention, cognitive inhibition, and information

processing speed (Treisman and Fearnley, 1969; Grundey et al., 2015). We used a computerized

Stroop color/word test similar to the Victoria version, based on the previous studies (Grundey

et al., 2015). This task includes three blocks, the Stroop word, the Stroop color, and the Stroop

color-word task. In the Stroop word, the color names were written in black, and in the Stroop

color, capital XXXs were presented in red, green, yellow, and blue ink, and participants had to

respond with the corresponding keys. In the Stroop color-word task, participants were presented

with either "congruent" or "incongruent" color words. In the incongruent trials, the color of the

ink in which the word was displayed was different from the meaning of the word (for example,

the word "red" was written in blue) while in the congruent trials both, word and color of the

ink, were identical. Stimuli were presented on a screen with black background for 2000 ms with

a 500 ms inter-stimulus interval. The size of the stimuli was 1.4 cm at approximately 50 cm

eye-to-screen distance. A response box with only four keys, colored in red, blue, yellow, and

green, was placed in front of the subjects and they had to press the corresponding key of the

color in which the word was written. The Stroop interference block included 40 congruent and

120 incongruent trials, resulting in a total of 160 trials. The reason for the higher number of

trials in the Stroop block was to increase the power of the EEG analyses.

4.7.3. AX-Continuous Performance Test (AX-CPT). The AX-CPT is used for assessing

attentional functioning (sustained or transient attention), or executive control, depending on

the applied versions, which include baseline, proactive control, and reactive control (Smid et

al., 2006; Gonthier et al., 2016). A baseline version of the task was used, which is shorter

(around 15 min), less demanding, and measures transient attention (Smid et al., 2006). In this

task, visual stimuli were white letters on a dark background appearing one at a time on a

computer screen for 150 ms each with a 2000 ms inter-stimulus interval. Subjects were

instructed to press a button with the right index finger whenever the letter A (correct cue) was

followed by the letter X (correct target) as quickly and accurately as possible. All other

sequences were to be ignored, including sequences in which an incorrect cue (designated 'B',

but comprising all letters other than A or X) was followed by the target letter (X), or

sequences in which a correct cue (A) was followed by an incorrect target (designated 'Y', but

comprising all letters other than A or X). The AX sequences are presented with a high

probability, to guarantee a strong response bias. The tasks consisted of 240 pairs of letters

(480 trials) with 40% "AX", 40% "BY", 10% "BX" and 10% of "AY". Accuracy and RT

were recorded for the target trials.

4.7.4. Procedure

Participants performed the tasks in two versions randomized across the sufficient sleepand

sleep-deprivation sessions with at least two weeks intervals. The order of tasks was

counterbalanced across participants. All tasks (SRTT, N-back, Stroop, and AX-CPT) were

presented on a computer screen (15.6″in. Samsung) via E-prime software (Schneider et al.,

2002), the viewing distance from the monitor was approximately 50 cm. The tasks were

conducted in a soundproof electro-magnetic shielded room during EEG recording. The

session took about 1 h and 45 min including cap-preparation time and cleaning the head.

Following this part, participants were instructed to remove the gel from their hair and head

skin and were guided to the TMS lab for monitoring cortical excitability and neuroplasticity

induction.

4.8. EEG

4.8.1. EEG recording

EEG recording included resting-state measurements, which consisted of eyes open and closed states alternating every 2 min for 4 minutes, and task-based measurements. EEG was recorded from 64 scalp electrodes with two additional horizontal and vertical electro-oculogram electrodes (HEOG, VEOG) to measure horizontal and vertical eye movements. The electrodes were positioned according to the international 10–20 system using the NeurOne Tesla EEG amplifier (Bittium, NeurOne, Bittium Corporation, Finland) with a sampling rate of 1000 Hz. The scalp electrodes sites included: Fp1, Fp2, F7, F3, Fz, F4, F8, FC5, FC1, FC2, FC6, T7, C3, Cz, C4, T8, FPz, CP5, CP1, CP2, CP6, PO9, PO5, P7, P3, Pz, P4, P8, FCz, O1, Oz, O2, AF7, AF3, AF4, AF8, F5, F1, F2, F6, TP9, FT7, FC3, FC4, FT8, TP10, C5, IZ, PO10, C6, TP7, CP3, CPz, CP4, TP8, P5, P1, P2, P6, PO7, PO3, POz, PO4, and PO8, and were mounted on the head with a cap (EASYCAP GmbH, Herrsching, Germany). The reference electrode was positioned on FCz, and the ground electrode was placed at the AFz position. The electrodes were connected to the head using high-viscosity electrolyte gel (SuperVisc, Easycap, Herrsching, Germany). All impedances were kept below 10 kΩ throughout the experimental sessions. EEG data were collected in a shielded room, and no spectral peaks at 50 Hz were observed. Raw EEG data were recorded and stored for offline analysis using BrainVision Analyzer 2.1 (Brain Products GmbH, München, Germany).

4.8.2. EEG data analysis

EEG recordings were band-pass filtered offline between 1 and 30 Hz (48dB/Octave) and re-referenced to an average reference. The VEOG and HEOG signals were used to correct for eye movement artifacts in ERP recordings using the Gratton and Coles method (Gratton et al., 1983) embedded in the BrainVision Analyzer 2.1. EEG data were then time-locked to the stimulus of interest onset in each task. Epochs started 100 ms before stimulus onset and ended 700 ms after stimulus onset in the SRTT, 100 ms before the target onset and ended 1000 ms after target onset in the 3-back and AX-CPT tasks, and 100 ms before stimulus onset and ended 1000 ms after stimulus onset in the Stroop task (both congruent and incongruent trials). Epochs were baseline-corrected using a -100-0 ms time window. Artifacts were identified using a combination of automated (artifacts greater than 100 µV peak-to-peak) and manual selection processes. Segments were removed based on this automatic selection, and visual inspection to identify artifacts due to sources of non-neurogenic activity. The remaining epochs were averaged for calculating the average ERP. Average ERP of blocks 5, 6, 7 in the SRTT task were based on 120 trials per block. In the N-back and AX-CPT tasks, average ERP of hits (correct response) was based on 40 and 96 trials respectively. In the Stroop task, the average ERP of congruent and incongruent trials was based on 40 and 120 trials respectively. For the analyses,

the following averaged components were investigated: (1) the P300 at electrodes Pz, Cz and P3 within a time window of 300-600 ms after stimulus onset in the SRTT learning blocks (block 5,6,7), (2) the P300 at electrodes Fz and Cz within a time window of 300-600 ms (Picton, 1992; Kok, 2001) after target stimulus onset in the 3-back task, (3) the N200 and N450 at electrodes Fz and Cz within time windows of 200-300 ms and 400-550 ms respectively after congruent and incongruent trials onset (Feroz et al., 2017) in the Stroop task, and (4) the P300 at electrodes Fz and Cz within a time window of 300-600 ms (Picton, 1992; Tekok-Kilic et al., 2001) after target onset (when target letter X was preceded by cue A) in the AX-CPT task. The time windows were selected based on previous studies and designated as the maximum positive or negative deflection occurring at the post-stimulus latency window. A fast Fourier transform analysis (Hanning window length: 10%) was performed on the epochs to obtain spectral power levels in the beta (13-30 Hz), alpha (7-13 Hz), theta (4-7 Hz) and delta (1-4 Hz) range.

4.9. Statistical Analysis

4.9.1. Cortical excitability

For the TMS protocols with a double-pulse condition (i.e., SICI-ICF, I-wave facilitation, SAI), the resulting mean values were normalized to the respective single-pulse condition. First, mean values were calculated individually and then inter-individual means were calculated for each condition. For the I-O curves, absolute MEP values were used. To test for statistical significance, repeated-measures ANOVAs were performed with ISIs, TMS intensity (in I-O curve only), and condition (sufficient sleep vs sleep deprivation) as within-subject factors and MEP amplitude as the dependent variable. In case of significant results of the ANOVA, *post hoc* comparisons were performed using Bonferroni-corrected *post hoc* t-tests to compare mean MEP amplitudes of each condition against the baseline MEP and to contrast sufficient sleepvs sleep deprivation conditions. To determine if single-pulse conditions differed across sleep conditions, they were entered as dependent variables in an ANOVA with sleep condition (sufficient sleep vs sleep deprivation) as a within-subject factor.

4.9.2. Neuroplasticity

The mean peak-to-peak amplitude of the 25 MEPs obtained for each time-point (BL, 0, 5, 10, 15, 20, 25, 30 min after tDCS) was calculated and averaged for active and sham tDCS in the sufficient sleepand sleep deprivation conditions. To determine if individual baseline measures differed within and between sessions, SI1mV and Baseline MEP were entered as dependent variables in a mixed-model ANOVA with session (4 levels) and condition (sufficient sleep vs

sleep deprivation) as within-subject factors and group (anodal vs cathodal) as between-subject factor. The mean MEP amplitude for each measurement time-point was normalized to the session's baseline (individual quotient of the mean from the baseline mean) resulting in values representing either increased (> 1.0) or decreased (< 1.0) excitability. Individual averages of the normalized MEP from each time-point were then calculated and entered as dependent variables in a mixed-model ANOVA with stimulation condition (active, sham), time-point (8 levels), and sleep condition (normal vs deprivation) as within-subject factors and group (anodal vs cathodal) as between-subject factor. In case of significant ANOVA results, *post hoc* comparisons of MEP amplitudes at each time point were performed using Bonferroni-corrected *post hoc* t-tests to examine if active stimulation resulted in a significant difference relative to sham (comparison 1), baseline (comparison 2), the respective stimulation condition at sufficient sleepvs sleep deprivation (comparison 3), and the between-group comparisons at respective timepoints (comparison 4).

4.9.3. Behavioural task performance

Means of RT, RT variability, and accuracy for SRTT blocks 5, 6, and 7 were calculated. Trials with wrong responses, as well as those with RTs of less than 150 ms (Collins and Long, 1996; Mella et al., 2015) or more than 3000 ms, and trials which deviated by 3 standard deviations or more from the average individual response time, were discarded (Tzvi et al., 2016). The mean RT, RT variability and accuracy of blocks were entered as dependent variables in repeated-measures ANOVAs with block (5, vs 6, 6 vs 7) and condition (sufficient sleep vs sleep deprivation) as within-subject factors. Because the RT differences between blocks 5 vs 6 and 6 vs 7 were those of major interest, *post hoc* comparisons were performed on RT differences between these blocks using paired-sample t-tests (two-tailed, $p<0.05$). For 3-back, Stroop and AX-CPT tasks, mean and standard deviation of RT and accuracy were calculated and entered as dependent variables in repeated-measures ANOVAs with sleep condition (sufficient sleep vs sleep deprivation) as the within-subject factor. For significant ANOVA results, *post hoc* comparisons of dependent variables were performed using paired-sample t-tests (two-tailed, $p<0.05$).

4.9.4. Correlational analyses

To assess the relationship between induced neuroplasticity and motor sequence learning, and the relationship between cortical excitability and cognitive task performance we used bivariate linear regression analysis (Pearson's correlation, two-tailed). For the first correlation, we used

individual grand-averaged MEP amplitudes obtained from anodal and cathodal tDCS pooled for the time-points between 0, and 20 min after interventions, and individual motor learning performance (i.e. BL6-5 and BL6-7 RT difference) across sleep conditions. For the second correlation, we used individual grand-averaged MEP amplitudes obtained from each TMS protocol and individual accuracy/RT obtained from each task across sleep conditions.

4.9.5. EEG

EEG data preprocessing and analysis were described in the previous section. For the resting state data, brain oscillations at mid-central electrodes (Fz, Cz, Pz) were analyzed with a $3{\times}2$ ANOVA with location (Fz, Cz, Pz) and sleep condition (sufficient sleep vs sleep deprivation) as the within-subject factors. For all tasks, individual ERP means were grand-averaged and entered as dependent variables in repeated-measures ANOVAs with sleep condition (sufficient sleep vs sleep deprivation) as the within-subject factor. No correction was used for investigating multiple electrode locations.

5. Acknowledgments

We appreciate Nicole Rück, Ensiyeh Ghasemian-Shirvan, Elham Ghanavati, and Mohsen Mosaybi-Samani for the stay overnight during the course of the experiment. We also thank Tobias Klimek and Tobias Blanke for their technical support. MAN is supported by a grant from the German Ministry of Research and Education (GCBS grant 01EE1403C).

6. Competing interests

MAN is a member of the Scientific Advisory Boards of Neuroelectrics and NeuroDevice. All other authors declare no competing interests

7. Author contributions

MAS: Investigation, Data Curation, Formal analysis, Data Curation, Methodology, Software, Visualization, Writing - Original Draft, Writing - Review & Editing. **E.GH:** Data Curation, Visualization, Writing - Review & Editing. **JR & JH**: Data Curation, Formal analysis (Saliva sampling), Writing - Review & Editing. **M-FK & MAN:** Conceptualization, Methodology, Project administration, Resources, Supervision, Writing - Review & Editing.

Supplementary material

1. Cortical excitability supplementary results

1.1.TMS protocols threshold values

Baseline MEP values of single-pulse conditions and other control conditions, as well as subject-
specific baseline sensitivity to TMS (SI1mV), are summarized in Table S.1. For each TMS
protocol, single-pulse condition MEPs (control condition) and the SI1mV obtained at each sleep
condition were compared by a repeated-measures ANOVA with sleep condition (sufficient
sleep, sleep deprivation) as the within-subject factor. The results of the respective ANOVAs
show that the mean values of the single pulse-elicited MEP, %MSO for RMT and AMT, and
SI_{1mv} did not significantly differ between the sleep conditions (Table S.2). A trendwise
difference was found between SI_{1mv} values of baseline MEP, with lower values after sleep
deprivation compared to sufficient sleep ($p = 0.054$). Furthermore, for the I-O curve protocol,
a significant difference between MEP amplitudes at RMT intensity (t = 2.42, $p = 0.022$) was
revealed.

Table S1. Baseline measurements of the cortical excitability protocols

Protocol	measurement	Experimental session	
		Sufficient sleep	**Sleep deprivation**
Single-pulse MEP	SI 1_{mV} (%)	50.40 ± 9.419	49.633 ± 8.973
RMT	%MSO	40.733 ± 7.319	40.466 ± 7.560
AMT	%MSO	34.60 ± 7.308	34.566 ± 6.941
I-O curve	RMT intensity MEP	0.176 ± 0.097	0.260 ± 0.171
SICI-ICF	Single-pulse MEP	1.033 ± 0.160	1.073 ± 0.190
I-wave facilitation	Single-pulse MEP	1.052 ± 0.216	1.096 ± 0.154
SAI	Single-pulse MEP	1.081 ± 0.143	1.089 ± 0.137

904 *Note*: MEP = motor-evoked potentials; SI_{1mv} (%) = maximum stimulator output (%MSO) required for
905 the SI1mV MEP amplitude; I-O curve = input-output curve; SICI-ICF = short latency intracortical
906 inhibition and facilitation; SAI = short-latency afferent inhibition; RMT = resting motor threshold; AMT
907 = Active motor threshold.

Table S2. Results of repeated-measures ANOVAs for control MEPs (RMT intensity MEP for I-O curve, single-pulse MEP for other protocols), SI1mV, and %MSO for RMT/AMT.

Protocol	Measurement	Factor	*df*	*F*	*p*
Single-pulse MEP	SI$_{1mV}$	Sleep condition	1	2.72	0.109
RMT	%MSO	Sleep condition	1	0.231	0.634
AMT	%MSO	Sleep condition	1	0.012	0.915
I-O curve	RMT intensity MEP	Sleep condition	1	5.89	**0.022**
SICI-ICF	Single-pulse MEP	Sleep condition	1	1.01	0.322
I-wave facilitation	Single-pulse MEP	Sleep condition	1	1.01	0.323
SAI	Single-pulse MEP	Sleep condition	1	0.09	0.766

Note: MEP = motor-evoked potentials; sleep condition = sufficient sleep and sleep deprivation; SI$_{1mv}$ (%) = maximum stimulator output (%MSO) required for the SI1mV MEP amplitude; I-O curve = input-output curve; SICI-ICF = short latency intracortical inhibition and facilitation; SAI = short-latency afferent inhibition; RMT = resting motor threshold; AMT = active motor threshold.

2. Neuroplasticity supplementary results

2.1. Baseline MEP difference

The average baseline MEPs (absolute value) as well as the TMS stimulus intensity required for 1 mV amplitude (SI1mV) obtained for each tDCS condition (anodal, cathodal, sham) for each group are summarized in Table S3. The baseline MEP values obtained from each sleep condition (sufficient sleep, sleep deprivation) for both groups (anodal, cathodal) from each tDCS state (active, sham) were analyzed with a 2×2×2 factorial ANOVA. The results show no interaction of sleep condition×group×tDCS state ($F_1 = 1.48$, $p = 0.234$), or sleep condition×group ($F_1 = 0.49$, $p = 0.486$), or sleep condition×tDCS state ($F_1 = 0.65$, $p = 0.426$), or group×tDCS state ($F_1 = 0.36$, $p = 0.550$). The main effects of group ($F_1 = 0.05$, $p = 0.819$), sleep condition ($F_1 = 0.06$, $p = 0.811$) and tDCS state ($F_1 = 0.17$, $p = 0.680$) were not significant neither. The same 2×2×2 factorial ANOVA was conducted for SI1mV values, and the results show no interaction of sleep condition×group×tDCS state ($F_1 = 0.25$, $p = 0.616$), sleep condition×group ($F_1 = 1.06$, $p = 0.311$), sleep condition×tDCS state ($F_1 = 0.45$, $p = 0.505$), and main effect of group ($F_1 = 0.88$, $p = 0.355$). The interaction of group×tDCS state ($F_1 = 4.73$, $p = 0.038$), and the main effects of sleep condition ($F_1 = 6.65$, $p = 0.015$), and tDCS state ($F_1 = 5.48$, $p = 0.026$) were, however, significant. Pairwise comparisons of stimulation state (active vs sham) in the sleep conditions with *post hoc* t-tests revealed significant differences of SI1mV

amplitudes between sufficient sleep vs sleep deprivation conditions for both, active tDCS ($t = 2.58$, $p = 0.015$) and sham tDCS ($t = 2.46$, $p = 0.020$). When stimulation states were compared at the same sleep condition (active vs sham, sufficient sleep; active vs sham, sleep deprivation), a marginal significant difference in SI1mV amplitudes was observed between active and sham tDCS in the sufficient sleep ($t = 2.06$, $p = 0.048$), but not sleep deprivation ($t = 1.17$, $p = 0.252$) conditions.

Table S3: Baseline measurements of the neuroplasticity sessions

tDCS	Sleep condition	baseline MEP		SI_{1mV} (%)	
		Active	**sham**	**Active**	**sham**
Anodal stimulation	sufficient sleep	1.002±0.071	1.019±0.071	48.666±8.050	48.533±7.827
	sleep deprivation	1.020±0.081	1.026±0.068	46.80 ±6.667	46.90±6.644
Cathodal stimulation	sufficient sleep	1.027±0.037	1.005±0.065	51.333±10.244	50.866±10.183
	sleep deprivation	1.001±0.088	1.018±0.056	50.566±10.051	50.133±10.043

Note: tDCS = transcranial direct current stimulation; MEP = motor-evoked potentials; SI_{1mv} (%) = maximum stimulator output (%MSO) required for the SI1mV MEP amplitude.

2.2.Reported tDCS side effects

The reported side effects during each tDCS session (average ± SD) after sufficient sleep and sleep deprivation are summarized in Table S4. The results of the 2 (group: anodal, cathodal) × 2 (sleep condition) 2 × (tDCS state: active, sham) factorial ANOVA conducted for each side effect showed no interaction or main effects, except for a significant main effect of tDCS state for itching, and tingling (Table S5). Pairwise comparisons of itching and tingling ratings with *post hoc* t-tests revealed a significantly higher rating for itching sensation during anodal tDCS compared to the sham condition only after sleep deprivation ($t = 2.41$, $p = 0.030$). When the ratings were compared regardless of stimulation polarity (i.e., active tDCS vs sham tDCS), a significantly higher rating of the itching sensation was observed between active and sham tDCS after both sufficient sleep ($t = 2.85$, $p = 0.008$) and sleep deprivation ($t = 3.06$, $p = 0.005$). The intensity of the reported side effects was in general low.

Table S4. Reported side effects of tDCS during stimulation after sufficient sleep and sleep deprivation

Side effects	Sleep condition	Anodal-tDCS	Sham-tDCS	Cathodal-tDCS	Sham-tDCS
Visual	sufficient sleep	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
	sleep deprivation	0.20±0.77	0.20±0.56	0.06±0.25	0.06±0.25
Itching	sufficient sleep	1.66±1.67	0.86±1.18	1.40±1.18	0.93±0.96

	sleep deprivation	2.26±1.57	1.26±1.43	1.66±1.58	0.93±1.16
Tingling	sufficient sleep	1.20±1.37	1.06±1.09	1.40±0.98	0.93±1.09
	sleep deprivation	1.46±1.12	1.00±0.92	1.20±1.26	1.00±1.25
Burning	sufficient sleep	1.26±1.43	1.26±1.16	0.73±0.96	0.73±1.27
	sleep deprivation	1.53±1.64	1.40±1.50	1.46±1.69	1.00±1.25
Pain	sufficient sleep	0.20±0.77	0.20±0.56	0.13±0.35	0.66±1.39
	sleep deprivation	0.46±1.24	0.26±0.70	0.60±1.05	0.33±0.61

The presence and intensity of the side-effects were rated on a numerical scale ranging from zero to five, zero representing no and five extremely strong sensations. Data are presented as mean ± SD.

Table S5. Repeated-measures ANOVA results for the presence and intensity of reported tDCS side-effects

Side effects	Source	df	F	p	ηp^2
Visual phenomenon	sleep condition	1	3.89	0.058	0.122
	group	1	0.97	0.332	0.034
	tDCS state	1	0.00	1.000	0.000
	sleep condition × group	1	0.97	0.33	0.034
	group × tDCS state	1	0.00	1.000	0.000
	sleep condition × tDCS state	1	0.00	1.000	0.000
Itching	sleep condition	1	2.23	0.146	0.074
	group	1	0.57	0.453	0.020
	tDCS state	1	15.14	**0.001**	0.351
	sleep condition × group	1	0.75	0.394	0.026
	group × tDCS state	1	0.60	0.443	0.021
	sleep condition × tDCS state	1	0.47	0.499	0.017
Tingling	sleep condition	1	0.01	0.924	0. 001
	group	1	0.02	0.870	0.001
	tDCS state	1	4.38	**0.046**	0.135
	sleep condition × group	1	0.23	0.633	0.008
	group × tDCS state	1	0.01	0.913	0.001
	sleep condition × tDCS state	1	0.01	0.926	0.001
Burning	sleep condition	1	2.95	0.141	0.076
	group	1	1.06	0.311	0.037
	tDCS state	1	1.60	0.190	0.050
	sleep condition × group	1	0.36	0.549	0.013
	group × tDCS state	1	0.18	0.668	0.007
	sleep condition × tDCS state	1	0.44	0.512	0.16
Pain	sleep condition	1	0.59	0.448	0.021
	group	1	0.41	0.525	0.015
	tDCS state	1	0.02	0.892	0.001
	sleep condition × group	1	0.10	0.744	0.004
	group × tDCS state	1	0.92	0.344	0.032
	sleep condition × tDCS state	1	3.69	0.065	0.117

The presence and intensity of reported side-effects during tDCS were analyzed by repeated-measures mixed-model ANOVAs with sleep condition (sufficient sleep vs sleep deprivation) and tDCS state (active vs sham) as the within-subject factors and group (anodal, cathodal) as the between-subject factor. Pairwise comparisons are calculated using *Student's* t-test. *n* = 30 (15 per group).

2.3.tDCS Blinding efficacy

To explore blinding efficacy we asked participants to guess whether they received real tDCS (1 mA) or sham tDCS (0 mA) after each stimulation condition across sleep conditions. Using the Chi-square Test for Associations, we explored whether participants in each group (anodal, cathodal) could correctly discern each real stimulation condition from its respective sham condition in the sufficient sleep and sleep deprivation sessions. The results of the respective Chi-square tests show no significant differences of participants' guesses between each real stimulation vs sham stimulation in both, anodal ($\chi^2 = 0.682$, $p=0.409$; $\chi^2 = 0.085$, $p=0.770$) and cathodal groups ($\chi^2 = 0.00$, $p=1.000$; $\chi^2 = 0.268$, $p=0.605$), and the whole group ($\chi^2 = 0.557$, $p=0.448$; $\chi^2 = 0.007$, $p=0.993$) (All) (Table 3.6).

3. *Implicit motor learning*

3.1. Absolute Reaction Time

We also analyzed SRTT task performance based on the absolute RT values. The results of the 2×3 ANOVA showed a significant interaction of sleep condition × block ($F_{1.89} = 3.43$, $p=0.042$, $\eta p^2 = 0.11$), and significant main effects of sleep condition ($F_1 = 51.95$, $p < 0.001$, $\eta p^2=0.64$), and block ($F_{1.94} = 30.17$, $p < 0.001$, $\eta p^2=0.51$). *Post hoc* comparisons of blocks revealed a significantly faster RT at blocks 5 and 7 and longer RT at block 6 after sufficient sleep, but not sleep deprivation (Fig. 3.9b). Baseline block and block 6 RT (2 values), which contain stimuli in random order were also compared, and the results of the 2×2 ANOVA showed a significant main effect of sleep condition ($F_1 = 2.27$, $p<0.001$), baseline block ($F_1 = 10.16$, $p=0.003$) and their interaction ($F_1 = 12.72$, $p=0.002$). These results show a generally slower RT after sleep deprivation, compared to sufficient sleep (Fig. S1a).

3.2.Error rate

The number of errors in the learning blocks in the respective sleep conditions was analyzed as well. The results of the 2×3 ANOVA showed a significant interaction of sleep condition×block ($F_{1.96} = 8.49$, $p = 0.001$, $\eta p^2 = 0.23$) and a significant main effect of sleep condition ($F_1 = 6.49$, $p =0.016$, $\eta p^2 = 0.18$), but not block ($F_{1.73} = 1.02$, $p = 0.365$). Post hoc t-tests showed a significantly higher number of committed errors at block 6 compared to block 5 only after sleep deprivation (Fig. S1b). Furthermore, when every single block was compared across sleep conditions, the number of committed errors was significantly higher at BL 4, 6, 7, and 8 after sleep deprivation (Fig. S1c).

1002

Fig. S1. The impact of sleep deprivation on motor learning performance. **a**, BL 5-6 absolute RT difference represents sequence learning and was significant only after sufficient sleep (t=2.78, *p*=0.005) but not sleep deprivation (t=1.47, *p*=0.141). The BL 6-7 RT difference represents learning retention and was again significant only after sufficient sleep (t=2.16, *p*=0.031) but not sleep deprivation (t=0.86, *p*=0.392). Asterisks [*] represent statistically significant differences between learning blocks RT (BL 6-5, BL 6-7]. The brackets refer to RT difference between blocks 6 vs 5 and 6 vs 7. **b,** After sleep deprivation, participants committed more errors at block 6 compared to block 5 (*t*=2.38, *p*=0.024) but not 7 (*t*=0.70, *p*=0.489). **c**, Block-specific error rate was however, significantly higher after sleep deprivation in BL 6 (*t*=3.80, *p*<0.001), 7 (*t*=3.12, *p*=0.004), and also BL 4 (*t*=2.41, *p*=0.022) and 8 (*t*=3.09, *p*=0.004), as compared to the sufficient sleep condition. **d**, Participants showed higher RT variability after sleep deprivation, in all learning block including block 5 (*t*=3.17, *p*=0.004), block 6 (*t* = 4.43, *p*<0.001), and block 7 (*t*=3.89, *p*<0.001). All pairwise comparisons are calculated using *post hoc* Student's t-tests (paired, *p*<0.05). *n*=30. Error bars represent s.e.m. BL = block; Asterisks [*] indicate significant differences.

1003
1004
1005

 3.3.RT variability

 The results of the 2×3 ANOVA showed a significant interaction of sleep condition × block

 ($F_{1.55} = 4.57$, $p = 0.023$, $\eta p^2 = 0.13$) and main effects of sleep condition ($F_1 = 16.72$, $p < 0.001$,

 $\eta p^2 = 0.36$) and block ($F_{1.81} = 4.64$, $p = 0.016$) on RT variability. Post hoc t-tests showed no

 significant difference between RT variation of at block 6 compared to block 5 across sleep

 conditions. The RT variability from BL 6 to 7 was, however, significantly higher after sleep

 deprivation. Furthermore, when every single block was compared across sleep conditions, RT

 variability was significantly higher at each block (BL 5, 6, 7) after sleep deprivation compared

 to sufficient sleep (Fig. S1d).

 3.4. EEG supplementary results

 Further analyses were conducted for electrodes of cento-parietal regions close to the electrodes

 of interest (C1, C2, P1, P2). These revealed larger P300 amplitudes at block 6 vs block 5 and 7

 after sufficient sleep. For the C1 electrode, the results of the respective ANOVA showed a

 significant main effect of sleep condition (F_1=9.54, p=0.001, ηp^2=0.25) and block ($F_{1.48}$=3.49,

 p=0.050, ηp^2=0.11), but no interaction of these factors on P300 amplitudes. Post hoc

 comparisons showed a significantly larger P-300 amplitude in all learning blocks, including

 block 6, after sufficient sleep compared to sleep deprivation (t_{BL5}=2.10, p=0.044, t_{BL6}=3.23,

 p=0.003, t_{BL7}=2.48, p=0.019) (Fig. S2). No significant difference between BL 6-5 and BL 6-7

 was observed in either condition. For the electrode C2, the results of the ANOVA showed a

 significant interaction of sleep condition × block (F_2=3.32, p=0.043; ηp^2=0.10), and main

 effects of sleep condition (F_1=5.02, p=0.033, ηp^2=0.15) and block ($F_{1.42}$=4.76, p=0.023,

 ηp^2=0.14). Post hoc comparisons of learning block showed a significantly higher P300

 amplitude at block 6 compared to block 5 and 7 only after sufficient sleep (t_{6-5}=2.74, p=0.010,

 t_{6-7}=2.64, p=0.013) but not sleep deprivation (t_{6-5}=0.70, p=0.485, t_{6-7}=1.92, p=0.064) . The P300

 amplitudes were also significantly larger at blocks 6 and 7 after sufficient sleep compared to

 the sleep deprivation (t_{BL6}=2.23, p=0.034, t_{BL7}=2.21, p=0.035) (Fig. S2). Finally, the results of

 the ANOVA conducted for the electrode P1 showed a significant main effect of learning blocks

 ($F_{1.40}$=5.68, p=0.013, ηp^2=0.16) but not sleep condition (F_1=1.54, p=0.22) or interaction of

 sleep condition×block ($F_{1.15}$=2.12, p=0.152) on the P300 amplitude. Post hoc comparisons of

 P300 amplitudes within and between conditions showed a significantly larger component at

 block 6 vs 5 (t=4.21, p<0.001) and 6 vs 7 (t=4.71, p<0.001) only after sufficient sleep. Similarly,

 for electrode P2, a significant main effect of learning blocks ($F_{1.39}$=7.79, p=0.004, ηp^2=0.21),

 but not sleep condition (F_1 =1.78=1, p=0.21), or interaction of sleep condition×block

1039 ($F_{1.20}=2.73$, $p=0.073$) were found for the P300 amplitude. Post hoc comparisons of the P300

1040 amplitude within and between conditions showed that the P300 amplitude was significantly

1041 larger at block 6 vs 5 ($t=4.27$, $p<0.001$) and 6 vs 7 ($t=5.17$, $p<0.001$) only after sufficient sleep

1042 (Fig S2).

Fig. S2. **P300 amplitudes of electrodes C1, C2, P1, and P2 during motor sequence learning across sleep conditions.** For electrode C1, there was no significant difference between learning blocks across sleep conditions. However, between-condition comparisons show significantly larger P300 amplitudes in all learning blocks after sufficient sleep compared to sleep deprivation. For electrode C2, Pairwise comparisons show a significantly larger P300 amplitude in block 6 compared to blocks 5 and 7 only after sufficient sleep as compared to sleep deprivation. Between-condition comparisons of respective blocks show a significantly higher P300 amplitude at blocks 6 and 7 in the sufficient sleep condition. For electrode P1, there was no significant difference between individual learning blocks across sleep conditions (i.e., sufficient sleep vs sleep deprivation). However, within-condition comparisons show a significantly higher P300 amplitude in block 6 vs 5 and block 6 vs 7 only after sufficient sleep. The same pattern of response was found for electrode P2. All pairwise comparisons are calculated using post hoc t-tests (paired, $p <0.05$). $n=30$. [*/ns] indicates significant/non-significant differences between each block across sleep conditions. Filled symbols represent significant differences between BL 6-5 and BL 6-7. ns = nonsignificant.

1043

4. Working memory and attention tasks

For working memory performance, we also calculated variability of RT at a secondary outcome

measure. The result of the within-subject design ANOVA revealed a significant main effect of

sleep conditions RT variability of hits ($F_1=4.78$, $p=0.037$). Post hoc Student's t-tests showed a

significantly enhanced WM performance with significantly more RT variability after sufficient

sleep, which could be due to an accuracy-RT trade-off. In the Stroop task, we investigated

performance accuracy and ERP components at electrode Cz as well. The results of respective

ANOVAs showed a significant main effect of sleep condition on the overall accuracy of the

Stroop stage (F_1=6.32, p=0.018; ηp^2=0.18), accuracy of congruent trials (F_1=4.77, p=0.037;

ηp^2=0.14), and accuracy of incongruent trials (F_1=5.03, p=0.029; ηp^2=0.16). Post hoc

comparisons of accuracy rate revealed that participants had a significantly higher number of

accurate responses to trials in the Stroop stage as well as incongruent and congruent trials (Fig.

S3a). For the electrode Cz, the results of the 2 (congruency)×2 (sleep condition) ANOVA

showed only a significant main effect of sleep condition on the N200 (F_1=9.03, p=0.006;

ηp^2=0.25) but not N450 component (Fig. S3b). Similarly, *post hoc Student´s* t-tests indicated a

significantly smaller N200 amplitude, for the incongruent trials only, for the Cz electrode after

sleep deprivation as compared to sufficient sleep (Fig. S3b). Finally for the AX-CPT task, we

also analyzed ERP components at other potentially relevant electrodes (F3, F4, C3, C4), and a

comparable main effect of sleep condition was found on P300 amplitude for electrodes F3 (F_1

$= 4.77$, $p = 0.038$; $\eta\text{p}^2 = 0.15$), F4 ($F_1 = 7.82$, $p = 0.011$; $\eta\text{p}^2 = 0.21$), C3 ($F_1 = 24.31$, $p < 0.001$;

$\eta\text{p}^2 = 0.48$), and C4 ($F_1 = 7.60$, $p = 0.011$; $\eta\text{p}^2 = 0.22$). Post hoc Student´s t-tests indicated that

sleep deprivation was related to a significantly smaller P300 amplitude in the F3, F4, C3 and

C4 electrodes (Fig. S3c).

Fig. S3. The impact of sleep deprivation on Stroop accuracy. a, Participants had less correct responses in overall trials of the Stroop stage (t=2.51, p=0.018), incongruent trials (t=2.30, p=0.029) and congruent trials (t=2.18, p=0.037) after sleep deprivation compared to sufficient sleep. n=29. **b,** N200 and N450 ERP components of Stroop task performance across sleep conditions for electrode Cz. The N200 component was significantly larger for the incongruent trials, but not congruent trials, after sufficient sleep vs sleep deprivation for the electrodes Cz (t=3.51, p=0.002). The N450 did not significantly differ during incongruent vs congruent trials across sleep sessions. n=28. **c,** The P300 amplitude was significantly larger after sufficient sleep at electrodes F3 (t=2.18, p=0.038), F4 (t=2.66, p=0.013), C3 (t=4.93, p<.001), and C4 (t=2.75, p=0.011). The temporal window of 250-650 ms including the P300 amplitude (300-600 ms). n=27. Error bars represent s.e.m. ns = nonsignificant; Asterisks [*] indicate significant differences. All pairwise comparisons were calculated via *post hoc* Student's t-tests (paired, p<0.05).

1067

5. Correlational analyses

5.1. Correlation between sequence learning and plasticity induction

To explore the association between motor learning and plasticity, we calculated the correlation between the respective parameters (Pearson's correlation, two-tailed). We found a significant negative correlation between enhanced anodal LTP-like plasticity after sufficient sleep and enhanced motor learning (indicated by reduced RT at learning blocks). Specifically, MEP amplitude enhancement after anodal tDCS was negatively correlated with both sequence learning acquisition (block 6 - 5 RT difference) (r=-0.558, p=0.031) and sequence learning retention (block 6 - 7 RT difference) (r=-0.734, p=0.002). This indicated that LTP-like plasticity effects after sufficient sleep were associated with better sequence learning. No correlation was found between cathodal LTD-like plasticity and sequence learning.

5.2. Correlation between cortical excitability, working memory, and attention

To explore the association between physiological parameters of cortical excitability, and cognitive performance, we correlated performance in the 3-back letter task, Stroop test and AX-CPT with the respective cortical excitability results. In the 3-back letter task, enhanced d prime index (a measure of performance accuracy) was positively correlated with cortical facilitation measured by ICF at ISI of 15 ms (i.e., larger MEP at ICF) after having sufficient sleep (r= 0.425, p=0.019). Conversely, lower accurate response during sleep deprivation was negatively correlated with converted intracortical inhibition to facilitation (i.e., larger MEP amplitude) measured by SAI at ISI of 40 ms (r= -0.386, p=0.035). This indicates that upscaled cortical facilitation was associated with poor working memory performance.

No correlation was observed between Stropp task outcome measures and cortical excitability measures. For AX-CPT task performance, there was only a significant *negative* correlation between enhanced performance accuracy after sufficient sleep and reduced intracortical inhibition (measured by averaged MEPs of SICI) at the same time (r= -0.372, p=0.043). This indicates that improved task performance (i.e., higher accuracy) were associated with decreased intracortical inhibition after having sufficient sleep. In the sleep deprivation condition, lower performance accuracy and negatively correlated with higher corticospinal excitability (i.e., enhanced MEP at 150% of RMT intensity) (r= -0.429, p=0.018).

References

Akerstedt T, Gillberg M (1990) Subjective and objective sleepiness in the active individual. Int J Neurosci 52:29-37.

Ambrus GG, Al-Moyed H, Chaieb L, Sarp L, Antal A, Paulus W (2012) The fade-in – Short stimulation – Fade out approach to sham tDCS – Reliable at 1 mA for naïve and experienced subjects, but not investigators. Brain Stimulation 5:499-504.

Awiszus F (2003) TMS and threshold hunting. In: Supplements to Clinical Neurophysiology (Paulus W, Tergau F, Nitsche MA, Rothwell JG, Ziemann U, Hallett M, eds), pp 13-23: Elsevier.

Batsikadze G, Moliadze V, Paulus W, Kuo M-F, Nitsche MA (2013) Partially non-linear stimulation intensity-dependent effects of direct current stimulation on motor cortex excitability in humans. The Journal of Physiology 591:1987-2000.

Biabani M, Aminitehrani M, Zoghi M, Farrell M, Egan G, Jaberzadeh S (2017) The effects of transcranial direct current stimulation on short-interval intracortical inhibition and intracortical facilitation: a systematic review and meta-analysis. Rev Neurosci 29:99-114.

Boroojerdi B, Battaglia F, Muellbacher W, Cohen LG (2001) Mechanisms influencing stimulus-response properties of the human corticospinal system. Clinical Neurophysiology 112:931-937.

Bushey D, Tononi G, Cirelli C (2011) Sleep and Synaptic Homeostasis: Structural Evidence in Drosophila. Science 332:1576-1581.

Campbell IG, Guinan MJ, Horowitz JM (2002) Sleep Deprivation Impairs Long-Term Potentiation in Rat Hippocampal Slices. Journal of Neurophysiology 88:1073-1076.

Carskadon MA (2011) Sleep's effects on cognition and learning in adolescence. In: Progress in Brain Research (Van Dongen HPA, Kerkhof GA, eds), pp 137-143: Elsevier.

Chen R (2000) Studies of human motor physiology with transcranial magnetic stimulation. Muscle Nerve 23:S26-S32.

Chouhan NS, Griffith LC, Haynes P, Sehgal A (2021) Availability of food determines the need for sleep in memory consolidation. Nature 589:582-585.

Collins LF, Long CJ (1996) Visual reaction time and its relationship to neuropsychological test performance. Archives of Clinical Neuropsychology 11:613-623.

de Vivo L, Bellesi M, Marshall W, Bushong EA, Ellisman MH, Tononi G, Cirelli C (2017) Ultrastructural evidence for synaptic scaling across the wake/sleep cycle. Science 355:507-510.

Deak MC, Stickgold R (2010) Sleep and cognition. Wiley Interdisciplinary Reviews: Cognitive Science 1:491-500.

Di Lazzaro V, Pilato F, Dileone M, Tonali PA, Ziemann U (2005a) Dissociated effects of diazepam and lorazepam on short-latency afferent inhibition. The Journal of Physiology 569:315-323.

Di Lazzaro V, Profice P, Ranieri F, Capone F, Dileone M, Oliviero A, Pilato F (2012) I-wave origin and modulation. Brain Stimulation 5:512-525.

Di Lazzaro V, Oliviero A, Profice P, Pennisi MA, Di Giovanni S, Zito G, Tonali P, Rothwell JC (2000) Muscarinic receptor blockade has differential effects on the excitability of intracortical circuits in the human motor cortex. Experimental Brain Research 135:455-461.

Di Lazzaro V, Oliviero A, Saturno E, Dileone M, Pilato F, Nardone R, Ranieri F, Musumeci G, Fiorilla T, Tonali P (2005b) Effects of lorazepam on short latency afferent inhibition and short latency intracortical inhibition in humans. The Journal of Physiology 564:661-668.

Eoh HJ, Chung MK, Kim S-H (2005) Electroencephalographic study of drowsiness in simulated driving with sleep deprivation. Int J Ind Ergon 35:307-320.

Feldman DE (2009) Synaptic Mechanisms for Plasticity in Neocortex. Annual Review of Neuroscience 32:33-55.

Feroz FS, Leicht G, Steinmann S, Andreou C, Mulert C (2017) The Time Course of Activity within the Dorsal and Rostral-Ventral Anterior Cingulate Cortex in the Emotional Stroop Task. Brain Topogr 30:30-45.

Finelli L, Baumann H, Borbély A, Achermann P (2000) Dual electroencephalogram markers of human sleep homeostasis: correlation between theta activity in waking and slow-wave activity in sleep. Neuroscience 101:523-529.

Fischer T, Langner R, Birbaumer N, Brocke B (2008) Arousal and Attention: Self-chosen Stimulation Optimizes Cortical Excitability and Minimizes Compensatory Effort. Journal of Cognitive Neuroscience 20:1443-1453.

Gaggioni G, Ly JQM, Muto V, Chellappa SL, Jaspar M, Meyer C, Delfosse T, Vanvinckenroye A, Dumont R, Coppieters 't Wallant D, Berthomier C, Narbutas J, Van Egroo M, Luxen A, Salmon E, Collette F, Phillips C, Schmidt C, Vandewalle G (2019) Age-related decrease in cortical excitability circadian variations during sleep loss and its links with cognition. Neurobiology of Aging 78:52-63.

Gonthier C, Macnamara BN, Chow M, Conway ARA, Braver TS (2016) Inducing Proactive Control Shifts in the AX-CPT. Front Psychol 7.

Gratton G, Coles MGH, Donchin E (1983) A new method for off-line removal of ocular artifact. Electroencephalogr Clin Neurophysiol 55:468-484.

Grundey J, Amu R, Ambrus GG, Batsikadze G, Paulus W, Nitsche MA (2015) Double dissociation of working memory and attentional processes in smokers and non-smokers with and without nicotine. Psychopharmacology 232:2491-2501.

Grundey J, Barlay J, Batsikadze G, Kuo M-F, Paulus W, Nitsche M (2018) Nicotine modulates human brain plasticity via calcium-dependent mechanisms. The Journal of Physiology 596:5429-5441.

Hanajima R, Ugawa Y, Terao Y, Enomoto H, Shiio Y, Mochizuki H, Furubayashi T, Uesugi H, Iwata NK, Kanazawa I (2002) Mechanisms of intracortical I-wave facilitation elicited with paired-pulse magnetic stimulation in humans. The Journal of Physiology 538:253-261.

Hoddes E, Dement W, Zarcone V (1972) Stanford-sleepiness-scale. Enzyklopädie der Schlafmedizin:1184.

Honda M, Deiber MP, Ibáñez V, Pascual-Leone A, Zhuang P, Hallett M (1998) Dynamic cortical involvement in implicit and explicit motor sequence learning. A PET study. Brain 121:2159-2173.

Huang Y-Z, Lu M-K, Antal A, Classen J, Nitsche M, Ziemann U, Ridding M, Hamada M, Ugawa Y, Jaberzadeh S, Suppa A, Paulus W, Rothwell J (2017) Plasticity induced by non-invasive transcranial brain stimulation: A position paper. Clinical Neurophysiology 128:2318-2329.

Huber R, Mäki H, Rosanova M, Casarotto S, Canali P, Casali AG, Tononi G, Massimini M (2012) Human Cortical Excitability Increases with Time Awake. Cerebral Cortex 23:1-7.

Islam N, Aftabuddin M, Moriwaki A, Hattori Y, Hori Y (1995) Increase in the calcium level following anodal polarization in the rat brain. Brain research 684:206-208.

Klimesch W (2012) Alpha-band oscillations, attention, and controlled access to stored information. Trends in Cognitive Sciences 16:606-617.

Kok A (2001) On the utility of P3 amplitude as a measure of processing capacity. Psychophysiology 38:557-577.

Kolasinski J, Hinson EL, Divanbeighi Zand AP, Rizov A, Emir UE, Stagg CJ (2019) The dynamics of cortical GABA in human motor learning. The Journal of Physiology 597:271-282.

Kopp C, Longordo F, Nicholson JR, Lüthi A (2006) Insufficient Sleep Reversibly Alters Bidirectional Synaptic Plasticity and NMDA Receptor Function. The Journal of Neuroscience 26:12456-12465.

Krause AJ, Simon EB, Mander BA, Greer SM, Saletin JM, Goldstein-Piekarski AN, Walker MP (2017) The sleep-deprived human brain. Nature Reviews Neuroscience 18:404-418.

Kreuzer P, Langguth B, Popp R, Raster R, Busch V, Frank E, Hajak G, Landgrebe M (2011) Reduced intra-cortical inhibition after sleep deprivation: A transcranial magnetic stimulation study. Neuroscience Letters 493:63-66.

Kuhn M, Wolf E, Maier JG, Mainberger F, Feige B, Schmid H, Bürklin J, Maywald S, Mall V, Jung NH, Reis J, Spiegelhalder K, Klöppel S, Sterr A, Eckert A, Riemann D, Normann C, Nissen C (2016) Sleep recalibrates homeostatic and associative synaptic plasticity in the human cortex. Nature Communications 7:12455.

Kujirai T, Caramia MD, Rothwell JC, Day BL, Thompson PD, Ferbert A, Wroe S, Asselman P, Marsden CD (1993) Corticocortical inhibition in human motor cortex. The Journal of Physiology 471:501-519.

Kuo H-I, Paulus W, Batsikadze G, Jamil A, Kuo M-F, Nitsche MA (2017) Acute and Chronic Noradrenergic Effects on Cortical Excitability in Healthy Humans. International Journal of Neuropsychopharmacology 20:634-643.

Lazzaro VD, Restuccia D, Oliviero A, Profice P, Ferrara L, Insola A, Mazzone P, Tonali P, Rothwell JC (1998) Magnetic transcranial stimulation at intensities below active motor threshold activates intracortical inhibitory circuits. Experimental Brain Research 119:265-268.

Lazzaro VD, Oliviero A, Profice P, Pennisi MA, Pilato F, Zito G, Dileone M, Nicoletti R, Pasqualetti P, Tonali PA (2003) Ketamine Increases Human Motor Cortex Excitability to Transcranial Magnetic Stimulation. The Journal of Physiology 547:485-496.

Lim J, Dinges DF (2010) A meta-analysis of the impact of short-term sleep deprivation on cognitive variables. Psychological bulletin 136:375.

Lisman JE (2001) Three Ca2+ levels affect plasticity differently: the LTP zone, the LTD zone and no man's land. The Journal of physiology 532:285-285.

Lo JC, Groeger JA, Cheng GH, Dijk D-J, Chee MWL (2016) Self-reported sleep duration and cognitive performance in older adults: a systematic review and meta-analysis. Sleep Med 17:87-98.

Ly JQM, Gaggioni G, Chellappa SL, Papachilleos S, Brzozowski A, Borsu C, Rosanova M, Sarasso S, Middleton B, Luxen A, Archer SN, Phillips C, Dijk D-J, Maquet P, Massimini M, Vandewalle G (2016) Circadian regulation of human cortical excitability. Nature Communications 7:11828.

Maret S, Faraguna U, Nelson AB, Cirelli C, Tononi G (2011) Sleep and waking modulate spine turnover in the adolescent mouse cortex. Nat Neurosci 14:1418-1420.

McDermott CM, LaHoste GJ, Chen C, Musto A, Bazan NG, Magee JC (2003) Sleep Deprivation Causes Behavioral, Synaptic, and Membrane Excitability Alterations in Hippocampal Neurons. The Journal of Neuroscience 23:9687-9695.

Mella N, Fagot D, Lecerf T, de Ribaupierre A (2015) Working memory and intraindividual variability in processing speed: A lifespan developmental and individual-differences study. Mem Cognit 43:340-356.

Minarik T, Berger B, Althaus L, Bader V, Biebl B, Brotzeller F, Fusban T, Hegemann J, Jesteadt L, Kalweit L, Leitner M, Linke F, Nabielska N, Reiter T, Schmitt D, Spraetz A, Sauseng P (2016) The Importance of Sample Size for Reproducibility of tDCS Effects. Frontiers in Human Neuroscience 10.

Misonou H, Mohapatra DP, Park EW, Leung V, Zhen D, Misonou K, Anderson AE, Trimmer JS (2004) Regulation of ion channel localization and phosphorylation by neuronal activity. Nat Neurosci 7:711-718.

Mull BR, Seyal M (2001) Transcranial magnetic stimulation of left prefrontal cortex impairs working memory. Clinical Neurophysiology 112:1672-1675.

Niethard N, Born J (2019) Back to baseline: sleep recalibrates synapses. Nat Neurosci 22:149-151.

Nitsche M, Paulus W (2000) Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. The Journal of Physiology 527:633-639.

Nitsche M, Paulus W (2001) Sustained excitability elevations induced by transcranial DC motor cortex stimulation in humans. Neurology 57:1899-1901.

Nitsche MA, Schauenburg A, Lang N, Liebetanz D, Exner C, Paulus W, Tergau F (2003a) Facilitation of Implicit Motor Learning by Weak Transcranial Direct Current Stimulation of the Primary Motor Cortex in the Human. Journal of Cognitive Neuroscience 15:619-626.

Nitsche MA, Doemkes S, Karaköse T, Antal A, Liebetanz D, Lang N, Tergau F, Paulus W (2007) Shaping the Effects of Transcranial Direct Current Stimulation of the Human Motor Cortex. Journal of Neurophysiology 97:3109-3117.

Nitsche MA, Fricke K, Henschke U, Schlitterlau A, Liebetanz D, Lang N, Henning S, Tergau F, Paulus W (2003b) Pharmacological Modulation of Cortical Excitability Shifts Induced by Transcranial Direct Current Stimulation in Humans. The Journal of Physiology 553:293-301.

Nitsche MA, Seeber A, Frommann K, Klein CC, Rochford C, Nitsche MS, Fricke K, Liebetanz D, Lang N, Antal A (2005) Modulating parameters of excitability during and after transcranial direct current stimulation of the human motor cortex. The Journal of physiology 568:291-303.

Paulus W, Classen J, Cohen LG, Large CH, Di Lazzaro V, Nitsche M, Pascual-Leone A, Rosenow F, Rothwell JC, Ziemann U (2008) State of the art: Pharmacologic effects on cortical excitability measures tested by transcranial magnetic stimulation. Brain Stimulation 1:151-163.

Picton TW (1992) The P300 wave of the human event-related potential. J Clin Neurophysiol 9:456-479.

Placidi F, Zannino S, Albanese M, Romigi A, Izzi F, Marciani MG, Palmieri MG (2013) Increased cortical excitability after selective REM sleep deprivation in healthy humans: A transcranial magnetic stimulation study. Sleep Med 14:288-292.

Polanía R, Nitsche MA, Ruff CC (2018) Studying and modifying brain function with non-invasive brain stimulation. Nat Neurosci 21:174-187.

Rothwell J, Hallett M, Berardelli A, Eisen A, Rossini P, Paulus W (1999) Magnetic stimulation: motor evoked potentials. The International Federation of Clinical Neurophysiology. Electroencephalogr Clin Neurophysiol Suppl 52:97.

Schendan HE, Searl MM, Melrose RJ, Stern CE (2003) An FMRI study of the role of the medial temporal lobe in implicit and explicit sequence learning. Neuron 37:1013-1025.

Schmidt C, Collette F, Cajochen C, Peigneux P (2007) A time to think: circadian rhythms in human cognition. Cognitive neuropsychology 24:755-789.

Schneider W, Eschman A, Zuccolotto A (2002) E-Prime reference guide: Psychology Software Tools, Incorporated.

Smid HGOM, de Witte MR, Homminga I, van den Bosch RJ (2006) Sustained and Transient Attention in the Continuous Performance Task. Journal of Clinical and Experimental Neuropsychology 28:859-883.

Squires K, Wickens C, Squires N, Donchin E (1976) The effect of stimulus sequence on the waveform of the cortical event-related potential. Science 193:1142-1146.

Stagg CJ (2014) The Physiological Basis of Brain Stimulation. In: The Stimulated Brain (Cohen Kadosh R, ed), pp 145-177. San Diego: Academic Press.

Stickgold R (2005) Sleep-dependent memory consolidation. Nature 437:1272-1278.

Tekok-Kilic A, Shucard JL, Shucard DW (2001) Stimulus modality and Go/NoGo effects on P3 during parallel visual and auditory continuous performance tasks. Psychophysiology 38:578-589.

Tononi G, Cirelli C (2003) Sleep and synaptic homeostasis: a hypothesis. Brain Research Bulletin 62:143-150.

Tononi G, Cirelli C (2014) Sleep and the Price of Plasticity: From Synaptic and Cellular Homeostasis to Memory Consolidation and Integration. Neuron 81:12-34.

Treisman A, Fearnley S (1969) The Stroop Test: Selective Attention to Colours and Words. Nature 222:437-439.

Tzvi E, Verleger R, Münte TF, Krämer UM (2016) Reduced alpha-gamma phase amplitude coupling over right parietal cortex is associated with implicit visuomotor sequence learning. NeuroImage 141:60-70.

Vyazovskiy VV, Cirelli C, Pfister-Genskow M, Faraguna U, Tononi G (2008) Molecular and electrophysiological evidence for net synaptic potentiation in wake and depression in sleep. Nat Neurosci 11:200-208.

Walker MP, Stickgold R (2006) Sleep, Memory, and Plasticity. Annual Review of Psychology 57:139-166.

Ziemann U, Tergau F, Wassermann EM, Wischer S, Hildebrandt J, Paulus W (1998) Demonstration of facilitatory I wave interaction in the human motor cortex by paired transcranial magnetic stimulation. The Journal of Physiology 511:181-190.

a Theta (4-7 Hz) band

b Beta (15-30 Hz) band

c Alpha (8-14 Hz) band

Supplementary material

1. *Cortical excitability supplementary results*

1.1. TMS protocols threshold values

Baseline MEP values of single-pulse conditions and other control conditions, as well as subject-specific baseline sensitivity to TMS (SI1mV), are summarized in Table S.1. For each TMS protocol, single-pulse condition MEPs (control condition) and the SI1mV obtained at each sleep condition were compared by a repeated-measures ANOVA with sleep condition (sufficient sleep, sleep deprivation) as the within-subject factor. The results of the respective ANOVAs show that the mean values of the single pulse-elicited MEP, %MSO for RMT and AMT, and SI_{1mv} did not significantly differ between the sleep conditions (Table S.2). A trendwise difference was found between SI_{1mv} values of baseline MEP, with lower values after sleep deprivation compared to sufficient sleep ($p = 0.054$). Furthermore, for the I-O curve protocol, a significant difference between MEP amplitudes at RMT intensity ($t = 2.42$, $p = 0.022$) was revealed.

Table S1. Baseline measurements of the cortical excitability protocols

Protocol	measurement	Experimental session	
		Sufficient sleep	**Sleep deprivation**
Single-pulse MEP	SI 1_{mV} (%)	50.40 ± 9.419	49.633 ± 8.973
RMT	%MSO	40.733 ± 7.319	40.466 ± 7.560
AMT	%MSO	34.60 ± 7.308	34.566 ± 6.941
I-O curve	RMT intensity MEP	0.176 ± 0.097	0.260 ± 0.171
SICI-ICF	Single-pulse MEP	1.033 ± 0.160	1.073 ± 0.190
I-wave facilitation	Single-pulse MEP	1.052 ± 0.216	1.096 ± 0.154
SAI	Single-pulse MEP	1.081 ± 0.143	1.089 ± 0.137

Note: MEP = motor-evoked potentials; SI_{1mv} (%) = maximum stimulator output (%MSO) required for the SI1mV MEP amplitude; I-O curve = input-output curve; SICI-ICF = short latency intracortical inhibition and facilitation; SAI = short-latency afferent inhibition; RMT = resting motor threshold; AMT = Active motor threshold.

Table S2. Results of repeated-measures ANOVAs for control MEPs (RMT intensity MEP for I-O curve, single-pulse MEP for other protocols), SI1mV, and %MSO for RMT/AMT.

Protocol	Measurement	Factor	df	F	p
Single-pulse MEP	SI_{1mV}	Sleep condition	1	2.72	0.109
RMT	%MSO	Sleep condition	1	0.231	0.634
AMT	%MSO	Sleep condition	1	0.012	0.915
I-O curve	RMT intensity MEP	Sleep condition	1	5.89	**0.022**
SICI-ICF	Single-pulse MEP	Sleep condition	1	1.01	0.322
I-wave facilitation	Single-pulse MEP	Sleep condition	1	1.01	0.323
SAI	Single-pulse MEP	Sleep condition	1	0.09	0.766

Note: MEP = motor-evoked potentials; sleep condition = sufficient sleep and sleep deprivation; SI_{1mv} (%) = maximum stimulator output (%MSO) required for the SI1mV MEP amplitude; I-O curve = input-output curve; SICI-ICF = short latency intracortical inhibition and facilitation; SAI = short-latency afferent inhibition; RMT = resting motor threshold; AMT = active motor threshold.

2. *Neuroplasticity supplementary results*

2.1. Baseline MEP difference

The average baseline MEPs (absolute value) as well as the TMS stimulus intensity required for 1 mV amplitude (SI1mV) obtained for each tDCS condition (anodal, cathodal, sham) for each group are summarized in Table S3. The baseline MEP values obtained from each sleep condition (sufficient sleep, sleep deprivation) for both groups (anodal, cathodal) from each tDCS state (active, sham) were analyzed with a 2×2×2 factorial ANOVA. The results show no interaction of sleep condition×group×tDCS state ($F_1 = 1.48$, $p = 0.234$), or sleep condition×group ($F_1 = 0.49$, $p = 0.486$), or sleep condition×tDCS state ($F_1 = 0.65$, $p = 0.426$), or group×tDCS state ($F_1 = 0.36$, $p = 0.550$). The main effects of group ($F_1 = 0.05$, $p = 0.819$), sleep condition ($F_1 = 0.06$, $p = 0.811$) and tDCS state ($F_1 = 0.17$, $p = 0.680$) were not significant neither. The same 2×2×2 factorial ANOVA was conducted for SI1mV values, and the results show no interaction of sleep condition×group×tDCS state ($F_1 = 0.25$, $p = 0.616$), sleep condition×group ($F_1 = 1.06$, $p = 0.311$), sleep condition×tDCS state ($F_1 = 0.45$, $p = 0.505$), and main effect of group ($F_1 = 0.88$, $p = 0.355$). The interaction of group×tDCS state ($F_1 = 4.73$, $p = 0.038$), and the main effects of sleep condition ($F_1 = 6.65$, $p = 0.015$), and tDCS state ($F_1 = 5.48$, $p = 0.026$) were, however, significant. Pairwise

comparisons of stimulation state (active vs sham) in the sleep conditions with *post hoc* t-tests revealed significant differences of SI1mV amplitudes between sufficient sleep vs sleep deprivation conditions for both, active tDCS ($t = 2.58$, $p = 0.015$) and sham tDCS ($t = 2.46$, $p = 0.020$). When stimulation states were compared at the same sleep condition (active vs sham, sufficient sleep; active vs sham, sleep deprivation), a marginal significant difference in SI1mV amplitudes was observed between active and sham tDCS in the sufficient sleep ($t = 2.06$, $p = 0.048$), but not sleep deprivation ($t = 1.17$, $p = 0.252$) conditions.

Table S3: Baseline measurements of the neuroplasticity sessions

tDCS	Sleep condition	baseline MEP		SI_{1mV} (%)	
		Active	sham	Active	sham
Anodal stimulation	sufficient sleep	1.002±0.071	1.019±0.071	48.666±8.050	48.533±7.827
	sleep deprivation	1.020±0.081	1.026±0.068	46.80 ±6.667	46.90±6.644
Cathodal stimulation	sufficient sleep	1.027±0.037	1.005±0.065	51.333±10.244	50.866±10.183
	sleep deprivation	1.001±0.088	1.018±0.056	50.566±10.051	50.133±10.043

Note: tDCS = transcranial direct current stimulation; MEP = motor-evoked potentials; SI_{1mv} (%) = maximum stimulator output (%MSO) required for the SI1mV MEP amplitude.

2.2. Reported tDCS side effects

The reported side effects during each tDCS session (average ± SD) after sufficient sleep and sleep deprivation are summarized in Table S4. The results of the 2 (group: anodal, cathodal) × 2 (sleep condition) 2 × (tDCS state: active, sham) factorial ANOVA conducted for each side effect showed no interaction or main effects, except for a significant main effect of tDCS state for itching, and tingling (Table S5). Pairwise comparisons of itching and tingling ratings with *post hoc* t-tests revealed a significantly higher rating for itching sensation during anodal tDCS compared to the sham condition only after sleep deprivation ($t = 2.41$, $p = 0.030$). When the ratings were compared regardless of stimulation polarity (i.e., active tDCS vs sham tDCS), a significantly higher rating of the itching sensation was observed between active and sham tDCS after both sufficient sleep ($t = 2.85$, $p = 0.008$) and sleep deprivation ($t = 3.06$, $p = 0.005$). The intensity of the reported side effects was in general low.

Table S4. Reported side effects of tDCS during stimulation after sufficient sleep and sleep deprivation

Side effects	Sleep condition	Anodal-tDCS	Sham-tDCS	Cathodal-tDCS	Sham-tDCS
Visual	sufficient sleep	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
	sleep deprivation	0.20±0.77	0.20±0.56	0.06±0.25	0.06±0.25
Itching	sufficient sleep	1.66±1.67	0.86±1.18	1.40±1.18	0.93±0.96
	sleep deprivation	2.26±1.57	1.26±1.43	1.66±1.58	0.93±1.16
Tingling	sufficient sleep	1.20±1.37	1.06±1.09	1.40±0.98	0.93±1.09
	sleep deprivation	1.46±1.12	1.00±0.92	1.20±1.26	1.00±1.25
Burning	sufficient sleep	1.26±1.43	1.26±1.16	0.73±0.96	0.73±1.27
	sleep deprivation	1.53±1.64	1.40±1.50	1.46±1.69	1.00±1.25
Pain	sufficient sleep	0.20±0.77	0.20±0.56	0.13±0.35	0.66±1.39
	sleep deprivation	0.46±1.24	0.26±0.70	0.60±1.05	0.33±0.61

The presence and intensity of the side-effects were rated on a numerical scale ranging from zero to five, zero representing no and five extremely strong sensations. Data are presented as mean ± SD.

Table S5. Repeated-measures ANOVA results for the presence and intensity of reported tDCS side-effects

Side effects	Source	df	F	p	ηp^2
Visual phenomenon	sleep condition	1	3.89	0.058	0.122
	group	1	0.97	0.332	0.034
	tDCS state	1	0.00	1.000	0.000
	sleep condition × group	1	0.97	0.33	0.034
	group × tDCS state	1	0.00	1.000	0.000
	sleep condition × tDCS state	1	0.00	1.000	0.000
Itching	sleep condition	1	2.23	0.146	0.074
	group	1	0.57	0.453	0.020
	tDCS state	1	15.14	**0.001**	0.351
	sleep condition × group	1	0.75	0.394	0.026
	group × tDCS state	1	0.60	0.443	0.021
	sleep condition × tDCS state	1	0.47	0.499	0.017
Tingling	sleep condition	1	0.01	0.924	0. 001
	group	1	0.02	0.870	0.001
	tDCS state	1	4.38	**0.046**	0.135
	sleep condition × group	1	0.23	0.633	0.008
	group × tDCS state	1	0.01	0.913	0.001
	sleep condition × tDCS state	1	0.01	0.926	0.001
Burning	sleep condition	1	2.95	0.141	0.076
	group	1	1.06	0.311	0.037
	tDCS state	1	1.60	0.190	0.050
	sleep condition × group	1	0.36	0.549	0.013
	group × tDCS state	1	0.18	0.668	0.007
	sleep condition × tDCS state	1	0.44	0.512	0.16

Pain	sleep condition	1	0.59	0.448	0.021
	group	1	0.41	0.525	0.015
	tDCS state	1	0.02	0.892	0.001
	sleep condition × group	1	0.10	0.744	0.004
	group × tDCS state	1	0.92	0.344	0.032
	sleep condition × tDCS state	1	3.69	0.065	0.117

The presence and intensity of reported side-effects during tDCS were analyzed by repeated-measures mixed-model ANOVAs with sleep condition (sufficient sleep vs sleep deprivation) and tDCS state (active vs sham) as the within-subject factors and group (anodal, cathodal) as the between-subject factor. Pairwise comparisons are calculated using *Student's* t-test. $n = 30$ (15 per group).

2.3. tDCS Blinding efficacy

To explore blinding efficacy we asked participants to guess whether they received real tDCS (1 mA) or sham tDCS (0 mA) after each stimulation condition across sleep conditions. Using the Chi-square Test for Associations, we explored whether participants in each group (anodal, cathodal) could correctly discern each real stimulation condition from its respective sham condition in the sufficient sleep and sleep deprivation sessions. The results of the respective Chi-square tests show no significant differences of participants' guesses between each real stimulation vs sham stimulation in both, anodal ($\chi^2 = 0.682$, $p=0.409$; $\chi^2 = 0.085$, $p=0.770$) and cathodal groups ($\chi^2 = 0.00$, $p=1.000$; $\chi^2 = 0.268$, $p=0.605$), and the whole group ($\chi^2 = 0.557$, $p=0.448$; $\chi^2 = 0.007$, $p=0.993$) (All) (Table 3.6).

3. Implicit motor learning

3.1. Absolute Reaction Time

We also analyzed SRTT task performance based on the absolute RT values. The results of the 2×3 ANOVA showed a significant interaction of sleep condition × block ($F_{1.89} = 3.43$, $p=0.042$, $\eta p^2 = 0.11$), and significant main effects of sleep condition ($F_1 = 51.95$, $p < 0.001$, $\eta p^2 = 0.64$), and block ($F_{1.94} = 30.17$, $p < 0.001$, $\eta p^2 = 0.51$). *Post hoc* comparisons of blocks revealed a significantly faster RT at blocks 5 and 7 and longer RT at block 6 after sufficient sleep, but not sleep deprivation (Fig. 3.9b). Baseline block and block 6 RT (2 values), which contain stimuli in random order were also compared, and the results of the 2×2 ANOVA showed a significant main effect of sleep condition ($F_1 = 2.27$, $p<0.001$), baseline block ($F_1 = 10.16$, $p=0.003$) and their interaction ($F_1 = 12.72$, $p=0.002$). These results show a generally slower RT after sleep deprivation, compared to sufficient sleep (Fig. S1a).

Fig. S1. The impact of sleep deprivation on motor learning performance. **a**, BL 5-6 RT difference between represents sequence learning and was significant only after sufficient sleep (t=2.78, *p*=0.005) but not sleep deprivation (t=1.47, *p*=0.141). The BL 6-7 RT difference represents learning retention and was again significant only after sufficient sleep (t=2.16, *p*=0.031) but not sleep deprivation (t=0.86, *p*=0.392). Asterisks [*] represent statistically significant differences between learning blocks RT (BL 6-5, BL 6-7]. The brackets refer to RT difference between blocks 6 vs 5 and 6 vs 7. **b,** After sleep deprivation, participants committed more errors at block 6 compared to block 5 (*t*=2.38, *p*=0.024) but not 7 (*t*=0.70, *p*=0.489). **c**, Block-specific error rate was however, significantly higher after sleep deprivation in BL 6 (*t*=3.80, *p*<0.001), 7 (*t*=3.12, *p*=0.004), and also BL 4 (*t*=2.41, *p*=0.022) and 8 (*t*=3.09, *p*=0.004), as compared to the sufficient sleep condition. **d**, Participants showed higher RT variability after sleep deprivation, in all learning block including block 5 (*t*=3.17, *p*=0.004), block 6 (*t* = 4.43, *p*<0.001), and block 7 (*t*=3.89, *p*<0.001). All pairwise comparisons are calculated using *post hoc* Student's t-tests (paired, *p*<0.05). *n*=30. Error bars represent s.e.m. BL = block; Asterisks [*] indicate significant differences.

1.1. Error rate

The number of errors in the learning blocks in the respective sleep conditions was analyzed as well. The results of the 2×3 ANOVA showed a significant interaction of sleep condition×block ($F_{1.96} = 8.49$, $p = 0.001$, $\eta p^2 = 0.23$) and a significant main effect of sleep condition ($F_1 = 6.49$, $p =0.016$, $\eta p^2 = 0.18$), but not block ($F_{1.73} = 1.02$, $p = 0.365$). Post hoc t-tests showed a significantly higher number of committed errors at block 6 compared to block 5 only after sleep deprivation (Fig. S1b). Furthermore, when every single block was compared across sleep conditions, the number of committed errors was significantly higher at BL 4, 6, 7, and 8 after sleep deprivation (Fig. S1c).

1.2. RT variability

The results of the 2×3 ANOVA showed a significant interaction of sleep condition × block ($F_{1.55} = 4.57$, $p = 0.023$, $\eta p^2 = 0.13$) and main effects of sleep condition ($F_1 = 16.72$, $p < 0.001$, $\eta p^2 = 0.36$) and block ($F_{1.81} = 4.64$, $p = 0.016$) on RT variability. Post hoc t-tests showed no significant difference between RT variation of at block 6 compared to block 5 across sleep conditions. The RT variability from BL 6 to 7 was, however, significantly higher after sleep deprivation. Furthermore, when every single block was compared across sleep conditions, RT variability was significantly higher at each block (BL 5, 6, 7) after sleep deprivation compared to sufficient sleep (Fig. S1d).

1.3. EEG supplementary results

Further analyses were conducted for electrodes of cento-parietal regions close to the electrodes of interest (C1, C2, P1, P2). These revealed larger P300 amplitudes at block 6 vs block 5 and 7 after sufficient sleep. For the C1 electrode, the results of the respective ANOVA showed a significant main effect of sleep condition ($F_1=9.54$, $p=0.001$, $\eta p^2=0.25$) and block ($F_{1.48}=3.49$, $p=0.050$, $\eta p^2=0.11$), but no interaction of these factors on P300 amplitudes. Post hoc comparisons showed a significantly larger P-300 amplitude in all learning blocks, including block 6, after sufficient sleep compared to sleep deprivation ($t_{BL5}=2.10$, $p=0.044$, $t_{BL6}=3.23$, $p=0.003$, $t_{BL7}=2.48$, $p=0.019$) (Fig. S2). No significant difference between BL 6-5 and BL 6-7 was observed in either condition. For the electrode C2, the results of the ANOVA showed a significant interaction of sleep condition × block ($F_2=3.32$, $p=0.043$; $\eta p^2=0.10$), and main effects of sleep condition ($F_1=5.02$, $p=0.033$, $\eta p^2=0.15$) and block ($F_{1.42}=4.76$, $p=0.023$, $\eta p^2=0.14$). Post hoc comparisons of learning block

showed a significantly higher P300 amplitude at block 6 compared to block 5 and 7 only after sufficient sleep (t_{6-5}=2.74, p=0.010, t_{6-7}=2.64, p=0.013) but not sleep deprivation (t_{6-5}=0.70, p=0.485, t_{6-7}=1.92, p=0.064) . The P300 amplitudes were also significantly larger at blocks 6 and 7 after sufficient sleep compared to the sleep deprivation (t_{BL6}=2.23, p=0.034, t_{BL7}=2.21, p=0.035) (Fig. S2). Finally, the results of the ANOVA conducted for the electrode P1 showed a significant main effect of learning blocks ($F_{1.40}$=5.68, p=0.013, ηp^2=0.16) but not sleep condition (F_1=1.54, p=0.22) or interaction of sleep condition×block ($F_{1.15}$=2.12, p=0.152) on the P300 amplitude. Post hoc comparisons of P300 amplitudes within and between conditions showed a significantly larger component at block 6 vs 5 (t=4.21, p<0.001) and 6 vs 7 (t=4.71, p<0.001) only after sufficient sleep. Similarly, for electrode P2, a significant main effect of learning blocks ($F_{1.39}$=7.79, p=0.004, ηp^2=0.21), but not sleep condition (F_1=1.78=1, p=0.21), or interaction of sleep condition×block ($F_{1.20}$=2.73, p=0.073) were found for the P300 amplitude. Post hoc comparisons of the P300 amplitude within and between conditions showed that the P300 amplitude was significantly larger at block 6 vs 5 (t=4.27, p<0.001) and 6 vs 7 (t=5.17, p<0.001) only after sufficient sleep (Fig S2).

Fig. S2. **P300 amplitudes of electrodes C1, C2, P1, and P2 during motor sequence learning across sleep conditions.** For electrode C1, there was no significant difference between learning blocks across sleep conditions. However, between-condition comparisons show significantly larger P300 amplitudes in all learning blocks after sufficient sleep compared to the sleep deprivation. For electrode C2, Pairwise comparisons show a significantly larger P300 amplitude in block 6 compared to blocks 5 and 7 only after sufficient sleep as compared to sleep deprivation. Between-condition comparisons of respective blocks show a significantly higher P300 amplitude at blocks 6 and 7 in the sufficient sleep condition. For electrode P1, there was no significant difference between individual learning blocks across sleep conditions (i.e., sufficient sleep vs sleep deprivation). However, within-condition comparisons show a significantly higher P300 amplitude in block 6 vs 5 and block 6 vs 7 only after sufficient sleep. The same pattern of response

was found for electrode P2. All pairwise comparisons are calculated using post hoc t-tests (paired, $p < 0.05$). $n=30$. [*/ns] indicates significant/non-significant differences between each block across sleep conditions. Filled symbols represent significant differences between BL 6-5 and BL 6-7. ns = nonsignificant.

2. Working memory and attention tasks

For working memory performance, we also calculated variability of RT at a secondary outcome measure. The result of the within-subject design ANOVA revealed a significant main effect of sleep conditions RT variability of hits ($F_1=4.78$, $p=0.037$). Post hoc Student's t-tests showed a significantly enhanced WM performance with significantly more RT variability after sufficient sleep, which could be due to an accuracy-RT trade-off. In the Stroop task, we investigated performance accuracy and ERP components at electrode Cz as well. The results of respective ANOVAs showed a significant main effect of sleep condition on the overall accuracy of the Stroop stage ($F_1=6.32$, $p=0.018$; $\eta p^2=0.18$), accuracy of congruent trials ($F_1=4.77$, $p=0.037$; $\eta p^2=0.14$), and accuracy of incongruent trials ($F_1=5.03$, $p=0.029$; $\eta p^2=0.16$). Post hoc comparisons of accuracy rate revealed that participants had a significantly higher number of accurate responses to trials in the Stroop stage as well as incongruent and congruent trials (Fig. S3a). For the electrode Cz, the results of the 2 (congruency)×2 (sleep condition) ANOVA showed only a significant main effect of sleep condition on the N200 ($F_1=9.03$, $p=0.006$; $\eta p^2=0.25$) but not N450 component (Fig. S3b). Similarly, *post hoc Student´s* t-tests indicated a significantly smaller N200 amplitude, for the incongruent trials only, for the Cz electrode after sleep deprivation as compared to sufficient sleep (Fig. S3b). Finally for the AX-CPT task, we also analyzed ERP components at other potentially relevant electrodes (F3, F4, C3, C4), and a comparable main effect of sleep condition was found on P300 amplitude for electrodes F3 ($F_1 = 4.77$, $p = 0.038$; $\eta p^2 = 0.15$), F4 ($F_1 = 7.82$, $p = 0.011$; $\eta p^2 = 0.21$), C3 ($F_1 = 24.31$, $p < 0.001$; $\eta p^2 = 0.48$), and C4 ($F_1 = 7.60$, $p = 0.011$; $\eta p^2 = 0.22$). Post hoc Student´s t-tests indicated that sleep deprivation was related to a significantly smaller P300 amplitude in the F3, F4, C3 and C4 electrodes (Fig. S3c).

Fig. S3. The impact of sleep deprivation on Stroop accuracy. a, Participants had less correct responses in overall trials of the Stroop stage (t=2.51, p=0.018), incongruent trials (t=2.30, p=0.029) and congruent trials (t=2.18, p=0.037) after sleep deprivation compared to sufficient sleep. n=29. **b**, N200 and N450 ERP components of Stroop task performance across sleep conditions for electrode Cz. The N200 component was significantly larger for the incongruent trials, but not congruent trials, after sufficient sleep vs sleep deprivation for the electrodes Cz (t=3.51, p=0.002). The N450 did not significantly differ during incongruent vs congruent trials across sleep session. n=28. **c**, The P300 amplitude was significantly larger after sufficient sleep at electrodes F3 (t=2.18, p=0.038), F4 (t=2.66, p=0.013), C3 (t=4.93, p<.001), and C4 (t=2.75, p=0.011). The temporal window of 250-650 ms including the P300 amplitude (300-600 ms). n=27. Error bars represent s.e.m. ns = nonsignificant; Asterisks [*] indicate significant differences. All pairwise comparisons were calculated via *post hoc* Student's t-tests (paired, p<0.05).

3. Correlational analyses

3.1. Correlation between sequence learning and plasticity induction

To explore the association between motor learning and plasticity, we calculated the correlation between the respective parameters (Pearson's correlation, two-tailed). We found a significant negative correlation between enhanced anodal LTP-like plasticity after sufficient sleep and enhanced motor learning (indicated by reduced RT at learning blocks). Specifically, MEP amplitude enhancement after anodal tDCS was negatively correlated with both sequence learning acquisition (block 6 - 5 RT difference) (r=-0.558, p=0.031) and sequence learning retention (block 6 - 7 RT difference) (r=-0.734, p=0.002). This indicated that LTP-like plasticity effects after sufficient sleep were associated with better sequence learning. No correlation was found between cathodal LTD-like plasticity and sequence learning.

3.2. Correlation between cortical excitability, working memory, and attention

To explore the association between physiological parameters of cortical excitability, and cognitive performance, we correlated performance in the 3-back letter task, Stroop test and AX-CPT with the respective cortical excitability results. In the 3-back letter task, enhanced *d* prime index (a measure of performance accuracy) was positively correlated with cortical facilitation measured by ICF at ISI of 15 ms (i.e., larger MEP at ICF) after having sufficient sleep ($r= 0.425$, $p=0.019$). Conversely, lower accurate response during sleep deprivation was negatively correlated with converted intracortical inhibition to facilitation (i.e., larger MEP amplitude) measured by SAI at ISI of 40 ms ($r= -0.386$, $p=0.035$). This indicates that upscaled cortical facilitation was associated with poor working memory performance.

No correlation was observed between Stropp task outcome measures and cortical excitability measures. For AX-CPT task performance, there was only a significant *negative* correlation between enhanced performance accuracy after sufficient sleep and reduced intracortical inhibition (measured by averaged MEPs of SICI) at the same time ($r= -0.372$, $p=0.043$). This indicates that improved task performance (i.e., higher accuracy) were associated with decreased intracortical inhibition after having sufficient sleep. In the sleep deprivation condition, lower performance accuracy and negatively correlated with higher corticospinal excitability (i.e., enhanced MEP at 150% of RMT intensity) ($r= -0.429$, $p=0.018$).