DAXX safeguards heterochromatin formation in embryonic stem cells

Authors: Antoine Canat (1,2), Adeline Veillet (1), Renaud Batrin (1), Clara Dubourg (1,4), Robert Illingworth (3), Emmanuelle Fabre (1), Pierre Therizols (1,4)[*]

Affiliation:

(1) Université de Paris, Laboratoire Génomes, Biologie Cellulaire et Thérapeutiques, CNRS UMR7212, INSERM U944, Institut de Recherche St Louis, F- 75010 Paris, France

(2) Present address: Institute of Epigenetics and Stem Cells, Helmholtz Zentrum München, München, Germany

(3) Centre for Regenerative Medicine, Institute for Regeneration and Repair, The University of Edinburgh, Edinburgh BioQuarter, 5 Little France Drive, Edinburgh EH16 4UU, United Kingdom.

(4) Present address: Université de Paris, Epigenetics and Cell Fate, CNRS, 75013 Paris, France

[*] Correspondence: pierre.therizols@cnrs.fr. Tel: +33-1-5727-8926

Abstract

Genomes comprise a large fraction of repetitive sequences folded into constitutive heterochromatin to protect genome integrity and cell identity. *De novo* formation of heterochromatin during preimplantation development is an essential step for preserving the ground-state of pluripotency and the self-renewal capacity of embryonic stem cells (ESCs). Yet, the molecular mechanisms responsible for the remodeling of constitutive heterochromatin are largely unknown. Here, we find that DAXX, an H3.3 chaperone, is essential for ESCs maintenance in the ground-state of pluripotency. DAXX accumulates at pericentromeric regions, and recruits PML and SETDB1, thereby promoting heterochromatin formation. In absence of DAXX or PML, the 3D-architecture and physical properties of pericentric and peripheral heterochromatin are disrupted, resulting in derepression of major satellite DNA, transposable elements and genes associated with the nuclear lamina. Using epigenome editing tools, we observe that H3.3, and specifically H3.3K9 modification, directly contribute to maintaining pericentromeric chromatin conformation. Altogether, our data reveal that DAXX is crucial for the maintenance and 3D-organization of the heterochromatin compartment and protects ESCs viability.

Keywords: Heterochromatin; Pericentromere; Chromocenters; Nuclear periphery; histone variant H3.3; DAXX; SETDB1; PML; Embryonic Stem Cell

Introduction

Eukaryotic nuclei generally contain two major genomic compartments, referred to as A and B, which respectively contain the active and inactive fractions of the genome. At the chromosome level, these compartments are characterized by alternating gene-rich and gene-poor domains. The B compartment is composed primarily of repetitive sequences, with satellite tandem repeats comprising one of the most abundant classes. Pericentromeric regions in mouse are composed of tandem repeats of a 234 bp sequence, named Major Satellite, that constitute approximately 8 Mb on each chromosome (Consortium et al., 2002; Guenatri et al., 2004). Most genes and repeated sequences of the B compartment are transcriptionally repressed by cooperative epigenetic mechanisms leading to the formation of constitutive heterochromatin. The three-dimensional organization of constitutive heterochromatin is an important layer regulating transcription (Falk et al., 2019; Lu et al., 2021). In some species, including the mouse, the pericentromeric heterochromatin (PCH) of different chromosomes aggregates to form large DAPI-dense heterochromatin clusters, called chromocenters. Disruption of satellite clustering is often associated with increased DNA damage and defects in chromosome segregation that are seen in pathologies such as Alzheimer's disease and breast cancer (Hahn et al., 2013; Jagannathan et al., 2018; Mansuroglu et al., 2016; Zhu et al., 2011). Gene-poor domains are preferentially found beneath the nuclear edge and often referred as Lamina-Associated Domains (LADs) (Guelen et al., 2008). The tethering to the nuclear envelope is stochastic, and LADs are also found associated with other nuclear landmarks such as chromocenters and nucleolus (Koningsbruggen et al., 2010; Wijchers et al., 2015).

At the molecular level, constitutive heterochromatin exhibits covalent modifications including DNA methylation, H3K9 and H4K20 trimethylation (H3K9me3 and H4K20me3). Maintenance of H3K9me3 at pericentromeric satellites relies on the SUV39H1/2 methyltransferases, whereas a distinct H3K9me3 methyltransferase, SETDB1, operates at dispersed DNA repeats and telomeres (Fukuda et al., 2018; Gauchier et al., 2019; Martens et al., 2005; Matsui et al., 2010).

Heterochromatin maintenance is intrinsically linked to chromatin replication during S phase,

but also requires the incorporation of histone variants, such as macroH2A and H3.3, into

chromatin independently of DNA synthesis (Buschbeck and Hake, 2017; Mendiratta et al.,

2019). The histone variant H3.3 differs from the canonical histone H3 by 4 or 5 residues and

is deposited throughout the cell cycle in active and inactive chromatin by different histone

chaperones. These histone-binding proteins prevent promiscuous incorporation into chromatin

(Szenker et al., 2011). Together with ATRX, DAXX forms a histone chaperone complex

responsible for H3.3 incorporation into heterochromatin regions, including PCH,

retrotransposons and telomeres (Drané et al., 2010; Elsässer et al., 2015; He et al., 2015;

Lewis et al., 2010; Sadic et al., 2015). At telomeres and retrotransposons, DAXX promotes

heterochromatin formation by recruiting SUV39H1/2 or SETDB1, hereby facilitating the

deposition of H3K9me3 (Elsässer et al., 2015; Gauchier et al., 2019; He et al., 2015; Hoelper

et al., 2017). In addition to its role in H3.3 deposition, DAXX also prevents promiscuous

incorporation of H3.3 by recruiting soluble H3.3-H4 dimers to PML nuclear bodies, membrane-

less hubs for post-translational modification of proteins (Delbarre et al., 2013; Delbarre et al.,

2017; Lallemand-Breitenbach and Thé, 2018).

Heterochromatin is *de novo* established during early embryogenesis, involving an important

remodeling of its molecular composition and 3D-organization (Burton and Torres-Padilla,

2014). However, the underlying mechanisms and their functional relevance for early

development remain elusive. During preimplantation development, the DNA methylation

inherited from parental gametes is erased and reaches its minimal level in the blastocyst,

defining the ground-state of pluripotency (Leitch et al., 2013; Ying et al., 2008). This wave of

DNA demethylation directly leads to the transcriptional upregulation of several repetitive

sequences, a process that is essential for the remodeling of constitutive heterochromatin

(Jachowicz et al., 2017; Lu et al., 2021; Probst et al., 2010). For instance, the upregulation of

major satellite transcripts is required for PCH reorganization into chromocenters and normal

developmental progression (Casanova et al., 2013; Probst et al., 2010). Yet, derepression of

DNA repeats correlates with high DNA damage signaling and their prolonged expression

results in developmental arrest, suggesting that silencing through a DNA methylation-

independent mechanism is required for normal embryogenesis (Jachowicz et al., 2017;

Ziegler-Birling et al., 2009). Polycomb group proteins were previously shown to bind and

facilitate silencing of repetitive sequences upon DNA hypomethylation (Saksouk et al., 2014;

Tosolini et al., 2018; Walter et al., 2016). However, knockout of most polycomb-related genes

does not impact pre-implantation embryogenesis (Aloia et al., 2013), implying that other factors

maintain heterochromatin in the absence of DNA methylation.

In mouse, both DAXX and H3.3 deletions cause genomic instability resulting in early embryonic

lethality with most knockout embryos failing to reach the blastocyst stage (Jang et al., 2015;

Liu et al., 2020; Michaelson et al., 1999). DAXX binds major satellite regions during the earliest

stages of development, just before PCH reorganizes to form chromocenters (Arakawa et al.,

2015; Liu et al., 2020). While the importance of DAXX for chromocenter formation remains

unknown, the viability of $Daxx^{-/-}$ embryos drops after chromocenter formation (Liu et al., 2020;

Probst et al., 2010), suggesting a direct link between DAXX function at chromocenters and

developmental progression. Surprisingly, *Daxx* is not required for *in vitro* culture of blastocyst-

derived Embryonic Stem Cells (ESCs) (Elsässer et al., 2015; He et al., 2015; Hoelper et al.,

2017). In ESCs, DAXX is not recruited to PCH, but rather to retrotransposons and telomeres

(Elsässer et al., 2015; Gauchier et al., 2019; He et al., 2015; Saksouk et al., 2014). However,

the recruitment of DAXX to chromatin is tightly associated with the low level of DNA methylation

that characterize preimplantation embryogenesis (Arakawa et al., 2015; He et al., 2015; Liu et

al., 2020). ESCs are typically cultured in serum-based media and accumulate high levels of

DNA methylation (Leitch et al., 2013), which could explain the discrepancies with *in vivo*

observations. ESCs can maintain low levels of DNA methylation and reach the ground-state

of pluripotency when grown in a serum-free medium, named 2i, but the role of DAXX for their

survival and the formation of PCH has yet to be addressed.

Here, we decipher the role played by DAXX at PCH in pluripotent ESCs. We find that DAXX is

essential for ground-state ESC survival. We provide evidence that DAXX localizes to

pericentric heterochromatin and recruits PML and SETDB1, facilitating heterochromatin

formation and organization through H3.3K9 modification. The deletion of *Daxx* or *Pml* impacts

pericentromeric heterochromatin formation altering its biophysical and clustering properties.

Thus, our results identify DAXX as an essential regulator preserving heterochromatin integrity

and maintaining the viability of ground-state ESCs.

Results

DAXX is essential for ESC survival upon ground-state conversion.

To assess the role of DAXX in pluripotent cells, we generated a *Daxx* knock-out (KO) ESC

line. We targeted exon 3 of *Daxx* using CRISPR/cas9 technology. While we confirmed the

absence of DAXX mRNA and protein (Fig. 1A; Fig. S1A), there were no obvious changes in

H3.3 protein levels in the resulting *Daxx* KO ESC line (Fig. 1A). Consistent with previous

reports, loss of DAXX did not impact the growth of ESCs cultured with a serum-based medium

(Elsässer et al., 2015). Likewise, neural differentiation induced by leukemia inhibitory factor

withdrawal and retinoic acid addition did not impact on cell viability in *Daxx* KO ESC (Diff., Fig.

1B). When the *Daxx* KO and WT ESCs were converted to the ground state, using a medium

with two small kinase inhibitors and vitamin C (hereafter denoted 2iV), no growth defect was

detected for *Daxx* KO cells after 4 days of conversion, but prolonged culture in 2iV medium

induced a drastic decrease in cell viability (Fig. 1B). Compared to the parental WT ESCs, only

12.6% of *Daxx* KO ESCs remained after 8 days of culture in 2iV (Fig. 1B). No surviving cells

could be detected after 9 to 10 days of conversion. We confirmed that under serum conditions,

ESCs have higher levels of DNA methylation than the blastocyst cells from which they are

derived (Fig. S1B) (Borgel et al., 2010). In contrast, ESCs cultured in 2iV, exhibit very low

levels of DNA methylation (Blaschke et al., 2013). Altogether, these results indicate an

essential role for DAXX in the maintenance of pluripotent cell survival upon ground-state

conversion.

To further investigate the impact of *Daxx* knockout, we monitored the transcriptional

landscapes under different conditions using RNA sequencing (RNA-seq). *Daxx* KO and WT

cells show only negligible differential expression of protein coding genes upon differentiation

with retinoic acid (Fig. S1C; Fig. S1D). We concluded that the absence of DAXX only has a

minor impact on transcriptome changes upon neural differentiation. This result is consistent

with the absence of morphological or cell growth defects observed upon differentiation and

confirms that DAXX is dispensable for neuronal differentiation. After 4 days of 2iV conversion

a substantial differential expression of genes in *Daxx* KO and WT ESCs was observed (Fig.

S1C). However, the global transcriptional response of *Daxx* KO ESCs to 2iV medium was

similar to WT ESCs (Fig. S1D). We compared the expression of 166 markers previously

identified as differentially expressed upon 2iV conversion (Blaschke et al., 2013). For both cell

lines most of these markers changed expression upon conversion according to the expected

pattern (Fig. S1E) supporting the conclusion that *Daxx* KO ESCs can reach the ground-state

of pluripotency after 4 days of 2iV conversion.

Overall, our data show that *Daxx* is essential for the survival of ground-state ESCs but their

loss of viability is unlikely caused by a conversion defect. Despite the mis-regulation of over

2500 genes after 4 days of 2iV conversion, no significant gene network that could explain the

subsequent loss of viability in *Daxx* KO ESCs could be identified.

DAXX contributes to the transcriptional repression of DNA repeats and peripheral

heterochromatin.

As DAXX is known to be responsible for H3.3 deposition in heterochromatic regions, we

investigated the impact of *Daxx* deletion on the transcriptional repression of the silenced

portion of the genome (Elsässer et al., 2015).

LADs make up a significant portion of the silenced compartment. To investigate the effect of

Daxx deletion on the expression of genes located in the 1180 LADs described in ESCs (Peric-

Hupkes et al., 2010), we compared their differential expression to that of genes not interacting

with the lamina (CEN). In serum and differentiated conditions, we found that LAD genes had

very similar differential expression to genes not located in LADs (Fig. 1C; Fig. 1D). However,

we observed an increase in the mRNA levels of several LAD genes normally silenced such as

Casp1, *Casp4*, and *Casp12* on chromosome 9 and *Naip6* and *Naip5* on chromosome 13, in

Daxx KO ESCs under 2iV condition (Fig. 1C; Fig. 1D; Fig. S2A). Genome-wide, the differential

expression of genes located in LADs was significantly increased in the 2iV condition,

suggesting that the presence of DAXX is necessary for proper silencing of these genes in

ground-state ESCs.

Histone modification H3K9me2 is a hallmark of LADs and generally accumulates beneath the

nuclear envelope (Kind et al., 2013; Peric-Hupkes et al., 2010). To assess whether the global

upregulation of LAD genes observed in *Daxx* KO ESCs upon 2iV conversion was associated

with a defect in peripheral heterochromatin assembly, we examined the accumulation of

H3K9me2 by immunofluorescence. In WT and *Daxx* KO ESCs grown in serum condition, the

H3K9me2 signal was irregularly diffuse in the nucleoplasm with a significant accumulation at

the nuclear edge (Fig. 1E). Quantification of H3K9me2 distribution by linescan density did not

show any significant difference in the enrichment of signal at the nuclear periphery between

WT and *Daxx* KO ESCs in serum (Fig. 1F; Fig. S2B). However, after 4 days of 2iV conversion,

the H3K9me2 signal remained enriched at the nuclear rim in WT converted cells, but only small

patches along the nuclear periphery were observed in the absence of DAXX (Fig. 1E).

Linescan quantification confirmed a significant decrease in H3K9me2 enrichment at the

nuclear periphery in *Daxx* KO ESCs (Fig. 1F). In contrast, LaminB1 accumulated normally at

the nuclear edge in both cell lines regardless of growing conditions, indicating no global nuclear

lamina assembly defects in *Daxx* KO ESCs (Fig. S2C; Fig. S2D). We therefore conclude that

DAXX is necessary for the maintenance of peripheral heterochromatin.

Overall, our observations show that *Daxx* deletion impacts the regulation of the nuclear

periphery by altering the tethering of peripheral heterochromatin and the transcriptional

silencing of genes located in LADs.

DAXX relocates to chromocenters upon ground-state conversion.

We additionally found that many endogenous retrovirus families were upregulated in *Daxx* KO

ESCs in serum and 2iV (Fig. 2A; Fig. S3), consistent with previous observations (Elsässer et

al., 2015; He et al., 2015). Several LINE1 elements, notably the L1MdT and L1MdA families,

not previously reported as DAXX targets, were also upregulated in pluripotent *Daxx* KO ESCs

(Fig. 2A; Fig. S3). The role of DAXX on transcriptional silencing was not limited to interspersed

repeats as we detected a strong upregulation of major satellites RNA in both serum and 2iV

condition in the *Daxx* KO cells (Fig. 2A; Fig.S3).

To investigate whether DAXX binds to major satellites in ESCs, we used immunofluorescence

with H3K9me3 staining to track PCH after confirming the strong correlation between H3K9me3

and major satellites using immuno-FISH (Fig. S4A; Fig. S4B). In both serum and 2iV

conditions, DAXX signal concentrated in the nucleoplasm as round foci (Fig. 2B; Fig. S4C).

The distribution of foci sizes was similar in both growing conditions, but a small fraction of

larger foci could be observed under 2iV condition. We therefore set a threshold defining small

and larger DAXX foci (Fig. S4D). The large DAXX foci exhibited higher levels of DAPI and

H3K9me3 specifically under 2iV condition, supporting that DAXX can relocate to PCH in

ground-state ESCs (Fig. S4E; Fig. S4F). To confirm this observation, we segmented nuclei

and PCH foci using DAPI and H3K9me3 signals, respectively, allowing us to measure the

enrichment of DAXX signal at PCH per nuclei (Fig. S4G). The level of DAXX observed at PCH

upon 2iV conversion was significantly higher than in the serum condition (Fig. 2C). Notably,

the increase in DAXX enrichment in 2iV was not homogenous among the H3K9me3 foci

population, but rather the result of a subset of clusters showing strong DAXX enrichment,

allowing us to determine a threshold to define DAXX-positive PCH (Fig. 2D). While only 14%

of cells showed one DAXX-positive PCH cluster in the serum condition, 83% of 2iV-converted

cells had at least one DAXX-positive H3K9me3 focus (Fig. 2E). A small proportion of cells

showed 2 or 3 or more DAXX-enriched PCH foci (16% and 5%, respectively). Interestingly, our

analysis revealed that DAXX-enriched PCH foci were bigger in size and had higher H3K9me3

staining intensities, suggesting a potential link between DAXX and H3K9me3 regulation (Fig.

2F; Fig. 2G).

Altogether, our data show that DAXX accumulates in a subset of PCH clusters in ground-state

ESCs, suggesting that it may play a role in the maintenance of PCH and the transcriptional

silencing of major satellite sequences.

DAXX maintains pericentric heterochromatin organization in pluripotent cells.

To investigate the role that DAXX might play at PCH in ground-state ESCs, we next

investigated whether DAXX facilitates PCH formation in ESCs. Since the loss of

pericentromeric silencing is generally caused by defective heterochromatin assembly and

often correlates with impaired clustering of chromocenters (Hahn et al., 2013; Healton et al.,

2020; Pinheiro et al., 2012; Zhu et al., 2011), we performed DNA FISH experiments against

the major satellites and segmented the 3D signal to determine the number of clusters (Fig.

3A). In serum or differentiated cells, the deletion of *Daxx* had no significant impact on the

number of clusters (Fig. 3B). However, after 4 days of 2iV conversion, the average number of

major satellite foci was increased by 37% in the *Daxx* KO ESCs. The presence of an increased

number of PCH foci indicates that DAXX is required for their clustering in ESCs, specifically

under 2iV condition and correlates with the loss of viability observed in these cells at later

stages of conversion (Fig. 1B). To confirm the role of DAXX in PCH clustering, we used the

TALE (transcription activator–like effector) epigenome editing tool to ask whether direct DAXX

recruitment to major satellites was sufficient to reduce the number of foci. We generated a

FLAG-tagged TALE engineered to specifically target major satellite sequences (hereafter called TALE$_{MajSat}$), following the design of a previous study (Miyanari et al., 2013). Immunofluorescence against the FLAG tag of the TALE$_{MajSat}$ confirmed its colocalization with the major satellite FISH signal (Fig. 3C). When bound to pericentromeres in pluripotent *Daxx* KO cells, TALE$_{MajSat}$-DAXX restored chromocenter clustering with the formation of fewer PCH foci (Fig. 3D). In WT ESCs, artificial tethering of DAXX to chromocenters had a similar impact, increasing endogenous PCH clustering (Fig. S5A). These results indicate that DAXX enhances physical interactions between major satellite sequences and contributes to the 3D organization of PCH.

We hypothesized that the decreased chromocenter clustering seen in *Daxx* KO ESCs might result from defective heterochromatin formation. Functional heterochromatin forms a self-segregating subcompartment with limited protein exchange with the rest of the nucleoplasm (Hinde et al., 2015; Strom et al., 2017). To test whether DAXX is important for heterochromatin segregation, we followed GFP-HP1α using live-imaging in WT and *Daxx* KO ESCs (Fig. 3E). We monitored the mobility of GFP-HP1α by measuring its fluorescence recovery rate at chromocenters after photobleaching (Fig. 3E; Fig. S5B). Compared to WT, the half-recovery time was significantly shorter in *Daxx* KO cells both in serum and 2iV conditions, suggesting higher protein exchange between chromocenter and the nucleoplasm in the absence of DAXX (Fig. 3F). However, the mobile fraction of GFP-HP1α remained constant in both cell lines, suggesting that the same amount of protein was bound to chromatin (Fig. S5B).

The boundary property of PCH is known to cause high variance in HP1α signal over time at the edge of a chromocenter (Strom et al., 2017). We reasoned that this higher HP1α recovery rate might arise from altered heterochromatin acting as a barrier to protein diffusion. We thus quantified temporal changes in GFP-HP1α signal intensity variation at unbleached chromocenters (Fig. 3G). We found that variance levels were low in the nucleoplasm of WT ESCs and only increased at chromocenters, peaking at their borders (Fig. 3G; Fig. 3H), confirming previous observations (Strom et al., 2017). However, in *Daxx* KO cells, the pattern

of the variance from HP1α signal was altered. In both serum and 2iV conditions, the peak of variance at the edge of chromocenters was significantly lower in the absence of DAXX, suggesting a compromised heterochromatin barrier (Fig. 3H). Moreover, molecular analysis using MNase digestion revealed that *Daxx* deletion in pluripotent ESCs increased the accessibility of pericentric chromatin, indicating that PCH exists in an abnormal state in these cells (Fig. S5C).

Altogether, our data suggests that DAXX plays a crucial role in regulating PCH compaction state, chromocenter boundary properties, and major satellite clustering. In conclusion, DAXX is essential for proper assembly and spatial organization of pericentromeric heterochromatin in pluripotent ESCs.

PML and DAXX share similar roles in ESCs.

PML, identified as a partner of DAXX, has been recently shown to contribute to the transcriptional repression of transposable elements in ESCs (Tessier et al., 2022). To investigate the role of PML in the organization of the heterochromatic compartment in ground-state ESCs, we first analyzed its localization in relation to DAXX and H3K9me3 by immunofluorescence (Fig. 4A; Fig. S6A). Independently of culture conditions, PML formed small round foci within the nucleoplasm that colocalized with DAXX, known as nuclear bodies (Fig. 4A; Fig. S6A). Upon 2iV conversion, a fraction of larger PML foci were observed with higher levels of H3K9me3 and DAPI (Fig. S6B; Fig. S6C; Fig. S6D). These larger PML foci accumulated around PCH in the form of either complete or partial structures, referred to as PML-rim or PML-arc, respectively (Fig. 4A; Fig. S6E). Both the PML-rim and PML-arc structures were consistently associated with the presence of DAXX at PCH and were observed in similar proportions as DAXX-positive PCH clusters (Fig. 2E; Fig. 4B). This indicates that PML is recruited to PCH alongside DAXX in ground-state ESCs.

To further examine the role played by PML at PCH clusters, we analyzed the distribution of

major satellites in *Pml* KO ESC using FISH (Tessier et al., 2022). In *Pml* KO ESCs, no protein

was detectable (Fig. S7A). Similar to the *Daxx* KO line, *Pml* KO ESCs showed a significant

increase in the number of major satellite foci only after 4 days of 2iV conversion (Fig. 4C; Fig.

4D), suggesting that PML, like DAXX, is required for proper PCH clustering in ground-state

ESCs.

The integrity of peripheral heterochromatin in *Pml* KO ESCs was analyzed.

Immunofluorescence revealed that H3K9me2 accumulated at the nuclear periphery to a

greater extent in serum conditions in *Pml* KO cells compared to WT cells (Fig. 4E; Fig. 4F).

However, after 4 days of 2iV conversion, the H3K9me2 signal in *Pml* KO cells was similar to

that observed in *Daxx* KO cells, with only small patches of H3K9me2 present beneath the

nuclear envelope and a significant reduction in peripheral enrichment of the signal (Fig. 4F).

In contrast, LaminB1 signal was not altered by Pml deletion supporting the conclusion that it is

specific to peripheral heterochromatin (Fig. S7B; Fig. S7C). To determine the effect of this

alteration of peripheral heterochromatin on gene expression, we analyzed the transcriptional

repression of LAD genes that were upregulated after *Daxx* deletion. Out of the six selected

genes, five showed higher mRNA levels in 2iV-converted *Pml* KO ESCs compared to WT cells

(Fig. 4G). Although the transcriptional upregulation was less severe in *Pml* KO cells than in

Daxx KO cells, our results suggest that PML, like DAXX, is involved in the transcriptional

silencing of LAD genes.

The deletion of *Pml* and *Daxx* have very comparable consequences on the heterochromatin

of ESCs after 4 days of 2iV conversion. We thus wondered how *Pml* deletion would impact the

viability of ESCs under prolonged 2iV conversion (Fig. 4H). After 4 days of conversion, the

viability of *Pml* KO was not impacted, but was decreased by nearly 75% after 8 days. Like the

deletion of *Daxx*, this effect was specific to prolonged 2iV conversion as the absence of PML

had no consequence on the viability under serum condition or after retinoic acid differentiation.

Overall, our data show that PML and DAXX play similar roles in ground-state ESCs. Like

DAXX, PML is recruited to PCH and its absence impairs the proper organization of the

heterochromatic compartment, significantly impacting cell viability.

DAXX recruits SETDB1 to chromocenters in ground-state ESCs.

We have shown that in ground-state ESCs, DAXX and PML are recruited to major satellites

thereby facilitating PCH formation. To further investigate the mechanism underlying the

function of DAXX and PML, we searched for DAXX- and PML-interacting factors that could be

recruited specifically to chromocenters. Amongst the DAXX interacting partners, SETDB1, an

H3K9me3 methyltransferase, colocalizes with DAXX at PML nuclear bodies and is important

for repetitive DNA transcriptional silencing (Cho et al., 2011; Karimi et al., 2011). SETDB1

partially colocalized with PML in a DAXX-independent manner in serum conditions (Fig. S8A).

Under 2iV treatment, SETDB1 relocalized to PCH with DAXX and PML in 70% of WT ESCs

(Fig. 5A; Fig. 5B; Fig. S8B). By contrast, PML and SETDB1 were only associated with PCH in

5 and 17% of *Daxx* KO cells, respectively (Fig. 5B). We conclude that DAXX is critical for PML

and SETDB1 recruitment to chromocenters. To confirm that DAXX was sufficient for SETDB1

recruitment to PCH, we expressed TALE$_{MajSat}$, fused to DAXX in *Daxx* KO ESCs. TALE-

mediated DAXX binding to major satellite repeats drastically increased the amount of SETDB1

signal observed at chromocenters (Fig. S8C; Fig. S8D).

The presence of DAXX at PCH correlated with higher amounts of H3K9me3 in 2iV condition

(Fig. 2G). However, DAXX deletion had no effect on H3K9me3 levels at chromocenters

independently of culture conditions (Fig. S9A; Fig. S9B). Since DAXX deletion prevented

SETDB1 recruitment to PCH, it suggests that the majority of H3K9me3 deposition at this locus

relies on, or can be rescued by other lysine methyl transferases, such as SUV39H1/2 (Lehnertz

et al., 2003; Peters et al., 2001).

To further investigate the conflicting results of DAXX-mediated SETDB1 recruitment and the

unchanged amount of H3K9me3 in *Daxx* KO ESCs, we examined the individual roles of DAXX,

SETDB1, and H3K9me3 in PCH spatial organization. We used Suv39H1/2 double knock-out

(Suv39dKO) ESCs, in which H3K9me3 can only result from SETDB1 activity. Suv39dKO cells

did not show any growth defects in 2iV medium, indicating the dispensability of both enzymes

for pluripotency. Unlike WT ESCs, Suv39dKO did not accumulate H3K9me3 at major satellite

foci, (Fig. S9A; Fig. 5C). However, we observed a strong accumulation of H3K9me3 in some

major satellite foci surrounded by PML upon 2iV conversion, suggesting that the fraction of

SETDB1 recruited to PCH in ground state is indeed functionally active (Fig. 5C; Fig. 5D). We

next used the TALE$_{MajSat}$ strategy to specifically target DAXX, SETDB1 or SUV39H1 to

chromocenters in Suv39dKO ESCs (Fig. 5E). Both histone methyltransferases were able to

individually restore H3K9me3 at chromocenters, when targeted by TALE$_{MajSat}$ (Fig. 5E; Fig

S9C). The specific recruitment of DAXX to major satellite foci also promoted H3K9me3

deposition via SETDB1, at 70% of chromocenters. The impact of the different TALE$_{MajSat}$

fusions on PCH clustering was further assessed by quantifying the number of TALE-bound foci

(Fig. 5F). Consistent with our observations in *Daxx* KO and WT ESCs (Fig. 3D, S5A) targeting

DAXX to major satellite also enhanced PCH clustering in Suv39dKO ESCs as evidenced by

the reduced number of PCH foci (Fig. 5F). However, the recruitment of either SETDB1 or

SUV39H1 did not affect the number of chromocenters, supporting the conclusion that

H3K9me3 modification does not influence PCH clustering (Fig. 5F).

Overall, our results demonstrate that DAXX recruits a catalytically active SETDB1 to PCH upon

2iV conversion. However, SETDB1 and H3K9me3 modification alone are not sufficient to

mediate PCH clustering, confirming that DAXX plays a central role in PCH organization in

ground-state ESCs.

H3.3K9 modification is crucial for PCH organization

Since DAXX, unlike SETDB1, had a direct impact on the spatial organization of chromocenters,

we examined the role of H3.3 chaperone activity of DAXX by expressing a mutated form,

DAXX^{Y222A}, that lowers its interaction with H3.3 (Elsässer et al., 2012). We confirmed that co-

immunoprecipitation of TALE$_{MajSat}$ DAXX^{Y222A} fusion with H3.3 was drastically reduced

compared to its wild-type counterpart (Fig. S9D). While targeting DAXX to chromocenters

could recruit a tagged version of H3.3, neither the control TALE$_{MajSat}$ nor TALE$_{MajSat}$ DAXX^{Y222A}

fusion were able to mobilize H3.3 to PCH (Fig. S9E; Fig. S9F). When expressed in Suv39dKO

ESCs, the TALE$_{MajSat}$ DAXX^{Y222A} fusion did not increase H3K9me3 levels at chromocenters, as

it failed to recruit SETDB1 (Fig. 5E; Fig. S8C; Fig. S8D). Importantly, expressing the DAXX^{Y222A}

mutant in *Daxx* KO ESCs did not change the number of chromocenters arguing that H3.3 is

crucial for DAXX-mediated PCH clustering (Fig. 5F). To further confirm a function for H3.3 in

chromocenter organization, we depleted endogenous H3.3 using a pool of siRNAs targeting

the 3'UTR of H3f3a/b mRNAs and expressed TALE$_{MajSat}$-DAXX in *Daxx* KO ESCs (Fig. 5G;

Fig. S9E). The H3.3 knock-down was rescued by the expression of different HA-tagged

versions of H3.3, which were assessed for their impact on PCH clustering by immuno-FISH

(Fig. 5G, S9G). The binding of DAXX at major satellite did not enhance PCH clustering when

H3.3 was knocked down. The expression of a wild-type H3.3-HA significantly reduced the

number of major satellite foci and rescued the PCH hyperclustering phenotype mediated by

TALE$_{MajSat}$-DAXX. In contrast, expression of H3.3^{G90M}, a mutant unable to bind DAXX (Elsässer

et al., 2012; Hoelper et al., 2017), did not decrease the number of major satellite foci (Fig. 5H).

This result is consistent with the absence of a chromocenter clustering phenotype in cells

transfected by TALE$_{MajSat}$- DAXX^{Y222A} (Fig. 5F) and confirms the important role of H3.3 in PCH

clustering. Since DAXX can recruit SETDB1 to PCH, we asked whether a H3.3K9 mutant that

cannot be methylated impacts PCH clustering and used H3.3^{K9A}. Like H3.3^{G90M}, H3.3^{K9A} failed

to rescue the hyper-clustering phenotype in TALE$_{MajSat}$-DAXX expressing cells, suggesting that

the modification of this residue is essential for the spatial organization of chromocenters (Fig.

5H).

Altogether, we conclude that the role of DAXX in PCH organization is intrinsically linked to its

chaperone activity. Interaction with H3.3 and H3.3K9 modifications are important in facilitating

the spatial organization of pericentromeres in pluripotent ESCs.

Discussion

During early embryogenesis, adapting heterochromatin to compensate for the wave of DNA

demethylation is essential to maintain transcriptional repression of repetitive DNA and protect

genome integrity. The mechanisms and molecular factors responsible for heterochromatin

reorganization were, however, largely unknown. Here, we describe a novel and essential role

for the H3.3-chaperone DAXX and its partner PML in the survival of ground-state pluripotent

stem cells. Taken together, our results support a model in which DAXX and PML facilitate both

pericentric and peripheral heterochromatin formation in ground-state ESCs (Fig. 6). At PCH,

DAXX deposits H3.3 and recruits PML and SETDB1, which can methylate H3.3K9. In the

absence of either DAXX or PML, PCH is compromised (Fig. 6). In *Daxx* KO ESCs,

heterochromatin at major satellite is less compact and partially loses its boundary properties

leading to defective chromocenter clustering. At the nuclear periphery, PML and DAXX are

necessary for the maintenance of the heterochromatin positioned beneath the nuclear

envelope, contributing to the transcriptional silencing of genes tethered to the nuclear lamina.

The global failure in heterochromatin compartment formation could directly contribute to the

impossibility of *Daxx* KO and *Pml* KO ESCs to maintain the ground-state of pluripotency.

DAXX and H3.3 impact chromocenter organization

Both DAXX and H3.3 are essential for early embryogenesis, but their specific functions have

remained elusive (Jang et al., 2015; Lin et al., 2013; Liu et al., 2020; Michaelson et al., 1999).

we found that DAXX is necessary for maintaining the spatial organization and chromatin state

of pericentric heterochromatin (PCH), thereby protecting the viability of ground-state ESCs

(Fig. 3B; Fig. 3F; Fig. 3H). This finding is consistent with *in vivo* observations, which show that

DAXX and SETDB1 are recruited to PCH just before chromocenter formation, and the viability

of *Daxx* KO embryos drops quickly after the transition to the pluripotent stage (Arakawa et al.,

2015; Cho et al., 2012; Liu et al., 2020). These observations support our conclusion that the

role of DAXX and SETDB1 at PCH might be crucial for the ground-state of pluripotency. Since

defective PCH assembly is often associated with genomic instability, our model may also

explain why *H3f3a/b* double knockout ESCs exhibit severe chromosome segregation defects

(Jang et al., 2015).

While *Daxx* deletion had no impact on major satellite transcription and chromocenter formation

upon neuronal differentiation (Fig. 2A; Fig. 3B), DAXX and H3.3 are important for chromocenter

clustering during myoblast differentiation (Park et al., 2018; Salsman et al., 2017). In

myoblasts, DAXX is recruited to chromocenters, where H3.3 deposition stimulates

transcription of major satellites, suggesting a mechanism different to the one we observed in

ESCs. Different DAXX and partner recruitment pathways could explain the opposite impact of

DAXX and H3.3 on major satellite transcription. In ESCs, DAXX recruits SETDB1, thus favoring

a repressive chromatin environment. Upon myoblast differentiation, *Setdb1* is downregulated

and might not be available for interaction with DAXX, which is recruited to major satellite

repeats by the muscle-specific lncRNA ChRO1 (Park et al., 2018; Song et al., 2015).

Recent insights into chromocenter organization have come from the observation that the

intrinsically disordered regions of HP1α facilitate the compartmentalization of chromocenters

through liquid-liquid phase separation (Larson et al., 2017; Strom et al., 2017). Since the first

fifty amino-acids and the C-terminal half of DAXX are intrinsically disordered (Escobar-Cabrera

et al., 2010), it is possible that DAXX may also undergo liquid-liquid phase separation.

Multivalent interactions between chromodomains recognizing H3K9me3 have also been

proposed to contribute to phase separation of heterochromatin (Wang et al., 2019). Yet, *Daxx*

deletion impacted the boundary properties of pericentromeric heterochromatin without

affecting H3K9me3 (Fig. 3H; Fig. S9A). Since DAXX-mediated chromocenter clustering relies

on H3.3K9 modification (Fig. 5H), future work will compare the interaction of HP1α and

additional heterochromatin-related proteins to H3.3K9me3 vs. H3K9me3.

Mechanism of DAXX targeting to pericentric heterochromatin

In ground-state ESCs, DAXX binds only a subset of chromocenters in approximatively 80% of

the population (Fig. 2E). Given that chromocenters replicate synchronously during mid-S

phase (Guenatri et al., 2004), the recruitment of DAXX is unlikely to rely on replicative

chromatin remodeling. DNA hypomethylation could be one of the factors responsible for DAXX

mobilization to PCH. Polycomb group proteins have been shown to be recruited to PCH

specifically under hypomethylated conditions (Pailles et al., 2022; Saksouk et al., 2014;

Tosolini et al., 2018). The fact that DAXX was shown to rely on PRC1 to bind major satellites

in zygotes would support a polycomb-mediated recruitment of DAXX in ground-state ESCs

(Liu et al., 2020). However, the presence of H3K27me3, deposited by PRC2, is relatively

homogenous amongst the PCH foci within a nucleus, suggesting a different mechanism

(Tosolini et al., 2018).

DNA damage may also be a factor in the recruitment of DAXX to PCH. Several studies have

described the incorporation of H3.3 after DNA damage (Adam et al., 2013; Fortuny et al., 2021;

Juhász et al., 2018; Li and Tyler, 2016; Luijsterburg et al., 2016). Both HIRA and DAXX-ATRX

complexes have been proposed to deposit H3.3 after DNA damage, but DAXX has been more

specifically observed at chromocenters damaged by UVC (Fortuny et al., 2021). Active DNA

demethylation may be a source of DNA damage responsible for DAXX recruitment. DAXX

relocation to PCH was specific to the 2iV culture condition (Fig. 2B), a medium that was shown

to stimulate TET enzymes activity (Blaschke et al., 2013). *In vivo*, DAXX binds preferentially

paternal chromosomes which are actively demethylated by the TET enzymes, a pathway that

can result ultimately in a DSB (Nakatani et al., 2015; Wossidlo et al., 2011).

Role of SETDB1 at chromocenters in ground-state ESCs

SETDB1 has been shown to be responsible for H3K9me3 deposition at transposable elements and telomeres (Elsässer et al., 2015; Gauchier et al., 2019; Karimi et al., 2011). Yet, knocking down SETB1 in Suv39h1/h2 double knock-out cells destabilizes chromocenters suggesting that SETDB1 could be involved in PCH formation (Pinheiro et al., 2012). Our data show that the chaperone activity of DAXX is necessary to recruit SETDB1 to pericentromeric heterochromatin (Fig. S8C). In contrast to SUV39H1/H2, SETDB1 contains a triple Tudor domain recognizing the double modification K14 acetylation and K9 methylation that may facilitate its binding to hyperacetylated, newly incorporated histone H3.3 (Jurkowska et al., 2017). Using ESCs lacking Suv39h1/h2, we found that SETDB1 can deposit H3K9me3 at chromocenters (Fig. 5C; Fig. 5D). This function of SETDB1 is consistent with its role at transposable element in ESCs (Elsässer et al., 2015; Hoelper et al., 2017). Yet, it contrasts with the observation that SETDB1 promotes H3K9me1 at major satellites in mouse embryonic fibroblasts (Loyola et al., 2009). However, SETDB1 was specifically recruited during S-phase by the H3.1/H3.2 chaperone, CAF1, suggesting that SETDB1 substrate specificity might change depending on its recruitment pathway. Similar to our data in ESCs, CAF1 recruited SETDB1 to chromocenters only in a subset of S-phase cells. Since pericentromeric satellites were shown to be particularly sensitive to replicative stress, SETDB1 recruitment could also result from DNA damage (Crosetto et al., 2013).

The role of DAXX and H3.3 upon DNA hypomethylation

This study reveals that DAXX is recruited to pericentromeres in the context of DNA hypomethylation. While PML nuclear bodies are generally devoid of chromatin, PML formed a shell around DAXX-positive chromocenters in hypomethylated ground-state ESCs (Fig. 4A). Similar structures have been observed in patients with immunodeficiency, centromeric instability and facial dysmorphia (ICF) syndrome associated with mutations in DNA methyltransferases (Luciani et al., 2006). In patient lymphocytes, DAXX and DNA repair proteins accumulate at hypomethylated pericentromeric satellites, suggesting that our proposed model could apply to other pathologies (Fig. 6).

In conclusion, our study reveals DAXX as an important factor for different heterochromatic compartments in ground-state ESCs. At the nuclear periphery, DAXX and its partner PML are necessary for the clustering of peripheral heterochromatin at the nuclear edge. At PCH, DAXX deposits H3.3 and recruits PML and SETDB1 to facilitate chromocenter formation by enhancing their clustering properties. It would be interesting to characterize whether DAXX also contributes to the 3D-organization of other heterochromatin domains, such as the clustering of LINE1 elements (Lu et al., 2021). Beyond early development, our work could also provide critical insights into the molecular pathways that overcome DNA hypomethylation transitions in pathological contexts, such as cancers.

Material and Methods

Cell lines

Feeder-free E14 mES cells were used for most experiment excepts otherwise notified. E14 mES cells were kindly provided by Pablo Navaro (Pasteur Institute). *Daxx* KO cell line was constructed using CRISPR/cas9 editing in E14 mES cells. Guide-RNA was designed using the online CRISPOR tool. Oligos were designed with a BbsI site on 5' to clone them into the pSpCas9(BB)-2A-Puro(pX459) v2.0 vector (Ran et al., 2013). Guide was designed carrying two guanines on 5' of the sequence to avoid off-target effects, as previously described (S. W. Cho et al., 2014). Sg-mDaxx: gGACCTCATCCAGCCGGTTCA. *Pml* KO ESCs were generated by CRISPR from E14 cells and described in a previous study (Tessier et al., 2022). Feeder-free Suv39HdKO and corresponding WT R1 ES cells were provided by Alice Jouneau and generated in Antoine Peters's laboratory (Lehnertz et al., 2003).

Culture conditions

Pluripotent cells were cultured either in a serum condition, defined as follow: DMEM (Gibco), supplemented with 10% of ESC-certified FBS (Gibco), 2-mercaptoethanol (0.05mM, Gibco), Glutamax (Gibco), MEM non-essential amino acids (0.1mM, Gibco), Penn-Strep (100units/mL,

Gibco) and LIF (1000units/mL,) for serum condition. Serum-cultivated cells were grown on

0.1% gelatin-coated plates or stem cell plates (Stem Cell technology) at 37°C with 5% CO2.

Medium was changed every day and cells were passaged every 2 to 3 days. The other culture

condition is the chemically defined serum-free 2i condition defined as follow:

Neurobasal:DMEM/F-12 (50:50, Gibco) medium, supplemented with N2 and B-27

supplements (Gibco), BSA fraction V (0.05%Gibco), 1-thioglycerol ($1.5x10^{-4}$M, Sigma) and

ESGRO 2 inhibitors (GSK3i and MEKi) and LIF (Merck). Vitamin C (L-ascorbic acid 2-

phosphate, Sigma) was added at a concentration of 100µg/mL (Blaschke et al., 2013). Cells

were grown on 0.1% gelatin-coated plates. Medium was changed daily. Cells were passed

every 2 days at 1:4 ratio for the first passage then at a 1:6 ratio. Differentiation of mES cells

was done by LIF removal for the first 24h. Then, non-LIF medium was supplemented with

retinoic acid (10^{-6}M,) for 4 days. For cell growth quantification, cells were counted at each

passage for at least 2 different experiments.

Vectors and transfections

Cells were harvested using trypsin and one million was plated in a 0.1% gelatin-coated plate

and transected with 0.2 to 2.5µg of DNA using the Lipofectamine 2000 reagent (Thermo),

following manufacturer's protocol. TALE vectors were constructed using previously described

methods (Ding et al., 2013). A TALE specific DNA binding domain targeting *Major Satellite*

repeats was created by the modular assembly of individual TALE repeats inserted into a

backbone vector containing TALE-Nrp1-VP64 previously described (Therizols et al., 2014).

The BamHI-NheI fragment containing VP64 was replaced by PCR products encoding the

DAXX protein, corresponding DAXX mutants or additional proteins such as SUV39H1 and

TET1CD. CDS of the different proteins were amplified by PCR from cDNAs obtained after RNA

extraction of serum E14 mES cells. DAXX mutants were generated by PCR from WT DAXX.

Oligos for major satellites targeting were designed with a BbsI site on 5' to clone them into the

pSpCas9(BB)-2A-Puro(pX459) v2.0 vector (Ran et al., 2013). Catalytically dead Cas9 (dCas9)

was inserted into the pSpCas9-2A-Puro(pX459) v2.0 to generate a SpdCas9-2A-Puro.

Target sequences for the TALE repeat domains and dCas9 associated guides RNA are listed

in supplementary table.

RNA extraction for RNAseq or RT-qPCR

Total RNA was extracted using RNeasy extraction kit (Qiagen) according with manufacturer's

protocol including DNAseI treatment for 15min at room temperature (Qiagen). Complementary

DNA were generated from 1µg of RNA using the Maxima first strand cDNA synthesis kit

(Thermo Fisher), with a second round of DNAseI from the Maxima kit for 15min. Real-time

qPCR was carried out using a LightCycler 480 instrument (Roche) and the LightCycler 480

SYBR green master mix (Roche). The qRT-PCR primers used in this study are listed in

supplementary table. Three independent biological repeats were obtained for each sample.

For RNAseq experiment, RNA quality was assessed using the Agilent 2100 bioanalyzer.

Libraries were prepared using oligo(dT) beads for mRNA enrichment, then fragmented and

reverse transcribed using random hexamers primer. After adaptor ligation, the double-stranded

cDNA is completed through size selection of 250-300bp and PCR amplification, then quality of

the library is assessed by the Agilent 2100 bioanalyzer. Sequencing was performed in 150bp

paired-end reads using an Illumina sequencer platform.

RNA-seq Mapping and Processing

FASTQ files generated by paired end sequencing were aligned to the mouse genome using

bowtie2 v2.2.6 (parameters: --local --threads 3; mm9 genome build). Mapped RNA-seq data

was processed using tools from the HOMER suite (v4.8). SAM files were converted into tag

directories using 'makeTagDirectory' (parameters: -format sam -sspe). Genomic intervals

which extended beyond the end of the chromosomes was removed using

'removeOutOfBoundsReads.pl'. bigWig browser track files were generated using

'makeUCSCfile' (parameters: -fsize 1e20 -strand + -norm 1e8). For gene expression analysis,

read depths were quantified for all annotated refseq genes using analyzeRepeats.pl

(parameters: rna mm9 -strand both -count exons -rpkm -normMatrix 1e7 -condenseGenes).

For repeat analysis, read coverage was quantified for each repeat and then condensed to a

single value for each named entry (parameters: repeats mm9 -strand both -rpkm -normMatrix

1e7 -condenseL1). Read depths were then corrected for the number of instances of each

repeat prior to expression analysis.

Expression Analysis

Quantified RNA-seq data was processed using the limma package (R/Bioconductor)(Team,

2017). Following the addition of an offset value (1 RPKM) to each gene or repeat, data was

normalised across all samples using 'normalizeBetweenArrays' with method='quantile'. Fold-

changes and p-values for differential expression of genes and repeats were determined using

empirical Bayes statistics. Briefly, data was fit to a linear model using 'lmFit' and specified

contrasts were applied using 'makeContrasts' and 'contrasts.fit'. Data was processed using the

'topTable' function with adjust.method="BH" (Benjamini-Hochberg multiple-testing correction).

Differential expression was defined as log2 fold change ≤-1 or ≥ 1 and an adjusted p-value of

≤ 0.01. Three biological replicates for each condition represent independently cultured pools

of cells.

Data visualization

Heatmaps and boxplots were generated using Prism GraphPad (v9). Histograms were drawn

using either Prism GraphPad or Excel. Volcano plots were generated using the plot function in

R.

Immunofluorescence

Murine ES cells were harvested with trypsin (Gibco) and plated for 4-12h onto either 0.1%

gelatin-coated or ECMatrix-coated (Sigma) glass cover slips. Cells were fixed with 4%

paraformaldehyde for 10min at room temperature, then rinsed three times with PBS. Cells

were permeabilized with 0.1X triton for 12min at room temperature, then rinsed three times

with PBS. Blocking was done in 3% BSA solution for 30min at room temperature. All

incubations with primary antibodies were performed for either 1h at room temperature or

overnight at 4°C with the following antibodies for H3K9me3 (Active Motif, 1:1000), DAXX (Santa Cruz, 1:500), H3K9me2 (Active Motif, 1:1000), LaminB1 (Abcam, 1:1000), PML () and SETDB1 (Proteintech, 1:100). Incubation with secondary antibodies (fluorescently labeled anti-mouse or anti-rabbit, 1:1000) were performed for 1h at room temperature. Mounting was performed using ProLong Diamond with DAPI mounting media (Thermo). Antibodies are listed in supplementary table.

Fluorescent in situ Hybridization

Murine ES cells were harvested with trypsin (Gibco) and plated for 4-6h onto 0.1% gelatin-coated glass cover slips. Cells were fixed with 4% paraformaldehyde (PFA) for 10min at room temperature, then rinsed three times with PBS. Cells were permeabilized with 0.5X triton for 12min at room temperature, then rinsed three times with PBS. Cells were briefly washed in 2X SSC, then treated with RNAseA (100µg/mL, Sigma) for 1h at 37°C. Cells were briefly washed in 2X SSC, then denatured by serial 2min incubation into 70,90 and 100% ethanol. Cover slips were air dried for 15min. Cover slips are incubated with 200nM of PNA probe, placed for 10min at 95°C for denaturation, then placed for 1h at room temperature in the dark for hybridization. Cover slips were washed twice in 2X SSC 0.1%Tween-20 for 10min at 60°C. Cover slips were immerged at room temperature in 2X SSC 0.1%Tween-20 for 2 min, then in 2X SSC for 2min and 1X SSC for 2min. Mounting is performed using ProLong Diamond with DAPI mounting media (Thermo).

Dot blot experiments

DNA extraction was performed using the Wizard genomic DNA extraction kit (Promega). Genomic DNA (1µg) was then denatured in 0.1M NaOH for 10min at 95°C before neutralization with 1M NH4OAc on ice for 5min. DNA samples were spotted on a nitrocellulose membrane. Blotted membrane was washed in 2X SSC and dried at 80°C for 5min before UV cross-linking at 120,000µJ/cm2. Membrane was then blocked using PBS, 5% milk 0.1% tween for 30min at room temperature. Membrane was incubated with 5mC antibody (Diagenode, 1:1000) for 3h

at room temperature. After 3 washes of 10min each in PBS, membrane was blocked again for

30min and then incubated with secondary anti-HRP antibody for 1h at room temperature.

Membrane was washed 3 times for 10min in PBS and visualized by chemiluminescence with

ECL Plus.

Fluorescent Recovery After Photobleaching and variance analysis

Murine ES cells transfected with HP1α-GFP were harvested with trypsin (Gibco) and plated

for 4-6h onto 0.1% gelatin-coated glass live-cell Nunc slides (Thermo). FRAP experiment was

carried with an LSM 800 confocal microscope (Zeiss). The 488nm laser was used to bleach

and acquire GFP signal. 1 image was taken before a bleach pulse of 5ms. Bleaching area was

set to target a single pericentric domain. Images were acquired every second during 35s post-

bleach. FRAP analysis was performed using a FRAP analysis ImageJ Jython script, that

generated FRAP curves and the associated half-recovery time and mobile fraction parameters.

GFP-HP1α variance along time was obtained from ImageJ analysis using standard deviation

z-projection along time for the whole duration of the movie. Quantification of heterochromatin

barriers were performed using a 1µm line across individual non-bleached chromocenter

borders, for which the variance intensity along time was measured with the ImageJ software.

Image acquisition and analysis

Images for immunofluorescence and FISH experiments were obtained with an inverted Nikon

Ti Eclipse widefield microscope using a 60X immersion objective and LED sources. Z-stacks

images were taken and then deconvoluted using a custom ImageJ deconvolution script.

Quantifications of images were performed using custom Icy scripts in Icy and ImageJ. To

quantify DAXX enrichment at PCH and the number of PCH foci, chromocenters were

segmented using ICY the spot detector function on either H3K9me3 or major satellite FISH

signal. Nuclei were isolated using the ICY k-means segmentation script. The nuclear periphery

enrichment was measured in ImageJ, briefly linescans of LaminB1 and H3K9me2 at the

periphery were isolated using the straighten command and subsequently used to generated intensity profiles.

H3K9me3 intensities at chromocenters were measured using à 2μm line across individual DAPI-dense chromocenters. Two to three chromocenters per nucleus were analyzed.

Statistical analysis

Number of objects counted, and statistical tests performed are indicated in the text, figure or figure legends. All statistical analysis results are listed in supplementary file. Pvalues are represented as follow: * <0.05; **<0.01; ***<0.001; ****<0.0001.

Western Blotting

Total protein extracts were prepared in RIPA buffer with protease inhibitor cocktail (Roche). Samples were sonicated for 3 minutes alterning 30seconds ON/ 30seconds OFF. Proteins were separated by electrophoresis in 8-15% poly-acrylamide gels then transferred onto nitrocellulose membranes. Membranes were incubated in ponceau then washed in PBS, 0.1% tween. Membranes were blocked in PBS, 0.1% tween, 5% milk for 30min at room temperature before incubation with primary antibody overnight at 4°C. After 3 washed in PBS, 0.1% tween, membranes were incubated with secondary HRP-conjugated antibody for 1h at room temperature. Membranes were washed 3 times in PBS, 0.1% tween and visualized by chemoluminescence using ECL Plus.

Acknowledgements

We would like to thank A. Bonnet-Garnier and A. Jouneau for the gift of Suv39dKO KO ESCs and advice. We are also grateful to image core microscopy facilities of IRSL, Paris, in particular N. Setterblad for his assistance with live imaging. We warmly thank V. Lallemand-Breitenbach and H. de Thé (Collège de France, Paris, France) for the sharing of PML antibody and helpful advice. We are thankful to Y. Khalil for her help in microscopy analysis. We kindly

688 thank P. Lesage and Annia Carré Simon for their critical readings of the manuscript, and other

689 members of the team for helpful advice. We finally thank support services of IRSL. This work

690 was funded by Agence Nationale pour la Recherche (ANR-16-CE12-0003), IDEX SLI, La Ligue

691 Nationale Contre le Cancer (LNCC)-Comité Ile de France and the labex Who am I. AC was

692 supported by MESR and LNCC. The authors declare no conflict of interest.

Author Contributions

695   AC, AV, CD, PT, performed all experiments, RI analyzed RNAseq data. AC, AV, PT

696 designed experiments, interpreted data. RB wrote the scripts for image analysis. AC, EF, PT

697 contributed to the writing of the manuscript. All authors reviewed the manuscript.

Declaration of interest

700 The authors declare no competing interests.

References

702
703

Adam, S., Polo, S. E. and Almouzni, G. (2013). Transcription recovery after DNA damage requires
chromatin priming by the H3.3 histone chaperone HIRA. *Cell* 155, 94–106.

Aloia, L., Stefano, B. D. and Croce, L. D. (2013). Polycomb complexes in stem cells and embryonic
development. *Development* 140, 2525–2534.

Arakawa, T., Nakatani, T., Oda, M., Kimura, Y., Sekita, Y., Kimura, T., Nakamura, T. and Nakano, T.
(2015). Stella controls chromocenter formation through regulation of Daxx expression in 2-cell
embryos. *Biochemical and biophysical research communications* 466, 60–65.

Blaschke, K., Ebata, K. T., Karimi, M. M., Zepeda-Martínez, J. A., Goyal, P., Mahapatra, S., Tam, A., Laird,
D. J., Hirst, M., Rao, A., et al. (2013). Vitamin C induces Tet-dependent DNA demethylation and a
blastocyst-like state in ES cells. *Nature* 500, 222–226.

Borgel, J., Guibert, S., Li, Y., Chiba, H., Schübeler, D., Sasaki, H., Forné, T. and Weber, M. (2010). Targets
and dynamics of promoter DNA methylation during early mouse development. *Nature Genetics* 42,
1093–1100.

Burton, A. and Torres-Padilla, M.-E. (2014). Chromatin dynamics in the regulation of cell fate allocation
during early embryogenesis. *Nat Rev Mol Cell Bio* 15, 723–735.

Buschbeck, M. and Hake, S. B. (2017). Variants of core histones and their roles in cell fate decisions,
development and cancer. *Nature Reviews Molecular Cell Biology* 18, 299–314.

Casanova, M., Pasternak, M., Marjou, F. E., Baccon, P. L., Probst, A. V. and Almouzni, G. (2013).
Heterochromatin reorganization during early mouse development requires a single-stranded
noncoding transcript. *Cell reports* 4, 1156–1167.

Cho, S., Park, J. S. and Kang, Y.-K. (2011). Dual functions of histone-lysine N-methyltransferase Setdb1
protein at promyelocytic leukemia-nuclear body (PML-NB): maintaining PML-NB structure and
regulating the expression of its associated genes. *Journal of Biological Chemistry* 286, 41115–
41124.

Cho, S., Park, J. S., Kwon, S. and Kang, Y.-K. (2012). Dynamics of Setdb1 expression in early mouse
development. *Gene expression patterns : GEP* 12, 213–218.

Consortium, M. G. S., Waterston, R. H., Lindblad-Toh, K., Birney, E., Rogers, J., Abril, J. F., Agarwal, P.,
Agarwala, R., Ainscough, R., Alexandersson, M., et al. (2002). Initial sequencing and comparative
analysis of the mouse genome. *Nature* 420, 520–562.

Crosetto, N., Mitra, A., Silva, M. J., Bienko, M., Dojer, N., Wang, Q., Karaca, E., Chiarle, R., Skrzypczak,
M., Ginalski, K., et al. (2013). Nucleotide-resolution DNA double-strand break mapping by next-
generation sequencing. *Nature methods* 10, 361–365.

Delbarre, E., Ivanauskiene, K., Küntziger, T. and Collas, P. (2013). DAXX-dependent supply of soluble
(H3.3-H4) dimers to PML bodies pending deposition into chromatin. *Genome Research* 23, 440–
451.

Delbarre, E., Ivanauskiene, K., Spirkoski, J., Shah, A., Vekterud, K., Moskaug, J. Ø., Bøe, S. O., Wong, L.
H., Küntziger, T. and Collas, P. (2017). PML protein organizes heterochromatin domains where it
regulates histone H3.3 deposition by ATRX/DAXX. *Genome Research* 27, 913–921.

Ding, Q., Lee, Y.-K., Schaefer, E. A. K., Peters, D. T., Veres, A., Kim, K., Kuperwasser, N., Motola, D. L.,
Meissner, T. B., Hendriks, W. T., et al. (2013). A TALEN genome-editing system for generating
human stem cell-based disease models. *Cell stem cell* 12, 238–251.

Drané, P., Ouararhni, K., Depaux, A., Shuaib, M. and Hamiche, A. (2010). The death-associated protein
DAXX is a novel histone chaperone involved in the replication-independent deposition of H3.3.
Genes & Development 24, 1253–1265.

Elsässer, S. J., Huang, H., Lewis, P. W., Chin, J. W., Allis, C. D. and Patel, D. J. (2012). DAXX envelops a
histone H3.3-H4 dimer for H3.3-specific recognition. *Nature* 491, 560–565.

Elsässer, S. J., Noh, K.-M., Diaz, N., Allis, C. D. and Banaszynski, L. A. (2015). Histone H3.3 is required for
endogenous retroviral element silencing in embryonic stem cells. *Nature* 522, 240–244.

Escobar-Cabrera, E., Lau, D. K. W., Giovinazzi, S., Ishov, A. M. and McIntosh, L. P. (2010). Structural
characterization of the DAXX N-terminal helical bundle domain and its complex with Rassf1C.
Structure (London, England : 1993) 18, 1642–1653.

Falk, M., Feodorova, Y., Naumova, N., Imakaev, M., Lajoie, B. R., Leonhardt, H., Joffe, B., Dekker, J.,
Fudenberg, G., Solovei, I., et al. (2019). Heterochromatin drives compartmentalization of inverted
and conventional nuclei. *Nature* 40, 47.

Fortuny, A., Chansard, A., Caron, P., Chevallier, O., Leroy, O., Renaud, O. and Polo, S. E. (2021). Imaging
the response to DNA damage in heterochromatin domains reveals core principles of
heterochromatin maintenance. *Nature communications* 12, 2428–16.

Fukuda, K., Okuda, A., Yusa, K. and Shinkai, Y. (2018). A CRISPR knockout screen identifies SETDB1-
target retroelement silencing factors in embryonic stem cells. *Genome Research* 28, 846–858.

Gauchier, M., Kan, S., Barral, A., Sauzet, S., Agirre, E., Bonnell, E., Saksouk, N., Barth, T. K., Ide, S.,
Urbach, S., et al. (2019). SETDB1-dependent heterochromatin stimulates alternative lengthening of
telomeres. *Science advances* 5, eaav3673.

Guelen, L., Pagie, L., Brasset, E., Meuleman, W., Faza, M. B., Talhout, W., Eussen, B. H., Klein, A. de,
Wessels, L., Laat, W. de, et al. (2008). Domain organization of human chromosomes revealed by
mapping of nuclear lamina interactions. *Nature* 453, 948–951.

Guenatri, M., Bailly, D., Maison, C. and Almouzni, G. (2004). Mouse centric and pericentric satellite
repeats form distinct functional heterochromatin. *The Journal of cell biology* 166, 493–505.

Hahn, M., Dambacher, S., Dulev, S., Kuznetsova, A. Y., Eck, S., Wörz, S., Sadic, D., Schulte, M., Mallm,
J.-P., Maiser, A., et al. (2013). Suv4-20h2 mediates chromatin compaction and is important for
cohesin recruitment to heterochromatin. *Genes & Development* 27, 859–872.

He, Q., Kim, H., Huang, R., Lu, W., Tang, M., Shi, F., Yang, D., Zhang, X., Huang, J., Liu, D., et al. (2015).
The Daxx/Atrx Complex Protects Tandem Repetitive Elements during DNA Hypomethylation by
Promoting H3K9 Trimethylation. *Cell stem cell* 17, 273–286.

Healton, S. E., Pinto, H. D., Mishra, L. N., Hamilton, G. A., Wheat, J. C., Swist-Rosowska, K., Shukeir, N.,
Dou, Y., Steidl, U., Jenuwein, T., et al. (2020). H1 linker histones silence repetitive elements by
promoting both histone H3K9 methylation and chromatin compaction. *Proceedings of the National*
Academy of Sciences of the United States of America 117, 14251–14258.

Hinde, E., Cardarelli, F. and Gratton, E. (2015). Spatiotemporal regulation of Heterochromatin Protein
1-alpha oligomerization and dynamics in live cells. *Scientific reports* 5, 12001–11.

Hoelper, D., Huang, H., Jain, A. Y., Patel, D. J. and Lewis, P. W. (2017). Structural and mechanistic
insights into ATRX-dependent and -independent functions of the histone chaperone DAXX. *Nature*
communications 8, 1193.

Jachowicz, J. W., Bing, X., Pontabry, J., Bošković, A., Rando, O. J. and Torres-Padilla, M.-E. (2017). LINE-
1 activation after fertilization regulates global chromatin accessibility in the early mouse embryo.
Nature Genetics 11, 778.

Jagannathan, M., Cummings, R. and Yamashita, Y. M. (2018). A conserved function for pericentromeric
satellite DNA. *eLife* 7, 1218.

Jang, C.-W., Shibata, Y., Starmer, J., Yee, D. and Magnuson, T. (2015). Histone H3.3 maintains genome
integrity during mammalian development. *Genes & Development* 29, 1377–1392.

Juhász, S., Elbakry, A., Mathes, A. and Löbrich, M. (2018). ATRX Promotes DNA Repair Synthesis and
Sister Chromatid Exchange during Homologous Recombination. *Molecular Cell* 71, 11-24.e7.

Jurkowska, R. Z., Qin, S., Kungulovski, G., Tempel, W., Liu, Y., Bashtrykov, P., Stiefelmaier, J., Jurkowski,
T. P., Kudithipudi, S., Weirich, S., et al. (2017). H3K14ac is linked to methylation of H3K9 by the
triple Tudor domain of SETDB1. *Nature communications* 8, 2057.

Karimi, M. M., Goyal, P., Maksakova, I. A., Bilenky, M., Leung, D., Tang, J. X., Shinkai, Y., Mager, D. L.,
Jones, S., Hirst, M., et al. (2011). DNA methylation and SETDB1/H3K9me3 regulate predominantly
distinct sets of genes, retroelements, and chimeric transcripts in mESCs. 8, 676–687.

Kind, J., Pagie, L., Ortabozkoyun, H., Boyle, S., Vries, S. S. de, Janssen, H., Amendola, M., Nolen, L. D.,
Bickmore, W. A. and Steensel, B. van (2013). Single-cell dynamics of genome-nuclear lamina
interactions. *Cell* 153, 178–192.

Koningsbruggen, S. van, Gierlinski, M., Schofield, P., Martin, D., Barton, G. J., Ariyurek, Y., Dunnen, J. T.
den and Lamond, A. I. (2010). High-resolution whole-genome sequencing reveals that specific
chromatin domains from most human chromosomes associate with nucleoli. *Molecular biology of*
the cell 21, 3735–3748.

Lallemand-Breitenbach, V. and Thé, H. de (2018). PML nuclear bodies: from architecture to function.
Current Opinion in Cell Biology 52, 154–161.

Larson, A. G., Elnatan, D., Keenen, M. M., Trnka, M. J., Johnston, J. B., Burlingame, A. L., Agard, D. A.,
Redding, S. and Narlikar, G. J. (2017). Liquid droplet formation by HP1α suggests a role for phase
separation in heterochromatin. *Nature* 547, 236–240.

Lehnertz, B., Ueda, Y., Derijck, A. A. H. A., Braunschweig, U., Perez-Burgos, L., Kubicek, S., Chen, T., Li,
E., Jenuwein, T. and Peters, A. H. F. M. (2003). Suv39h-mediated histone H3 lysine 9 methylation
directs DNA methylation to major satellite repeats at pericentric heterochromatin. *Current Biology*
13, 1192–1200.

Leitch, H. G., McEwen, K. R., Turp, A., Encheva, V., Carroll, T., Grabole, N., Mansfield, W., Nashun, B.,
Knezovich, J. G., Smith, A., et al. (2013). Naive pluripotency is associated with global DNA
hypomethylation. *Nature structural & molecular biology* 20, 1–9.

Lewis, P. W., Elsaesser, S. J., Noh, K.-M., Stadler, S. C. and Allis, C. D. (2010). Daxx is an H3.3-specific
histone chaperone and cooperates with ATRX in replication-independent chromatin assembly at
telomeres. *Proceedings of the National Academy of Sciences of the United States of America* 107,
14075–14080.

Li, X. and Tyler, J. K. (2016). Nucleosome disassembly during human non-homologous end joining
followed by concerted HIRA- and CAF-1-dependent reassembly. *eLife* 5, 94.

Lin, C.-J., Conti, M. and Ramalho-Santos, M. (2013). Histone variant H3.3 maintains a decondensed
chromatin state essential for mouse preimplantation development. *Development* 140, 3624–3634.

Liu, Z., Tardat, M., Gill, M. E., Royo, H., Thierry, R., Ozonov, E. A. and Peters, A. H. (2020). SUMOylated
PRC1 controls histone H3.3 deposition and genome integrity of embryonic heterochromatin. *The*
EMBO Journal e103697.

Loyola, A., Tagami, H., Bonaldi, T., Roche, D., Quivy, J.-P., Imhof, A., Nakatani, Y., Dent, S. Y. R. and
Almouzni, G. (2009). The HP1alpha-CAF1-SetDB1-containing complex provides H3K9me1 for Suv39-
mediated K9me3 in pericentric heterochromatin. *EMBO reports* 10, 769–775.

Lu, J. Y., Chang, L., Li, T., Wang, T., Yin, Y., Zhan, G., Han, X., Zhang, K., Tao, Y., Percharde, M., et al.
(2021). Homotypic clustering of L1 and B1/Alu repeats compartmentalizes the 3D genome. *Cell*
research 1, 26.

Luciani, J. J., Depetris, D., Usson, Y., Metzler-Guillemain, C., Mignon-Ravix, C., Mitchell, M. J.,
Megarbane, A., Sarda, P., Sirma, H., Moncla, A., et al. (2006). PML nuclear bodies are highly
organised DNA-protein structures with a function in heterochromatin remodelling at the G2 phase.
Journal of Cell Science 119, 2518–2531.

Luijsterburg, M. S., Krijger, I. de, Wiegant, W. W., Shah, R. G., Smeenk, G., Groot, A. J. L. de, Pines, A.,
Vertegaal, A. C. O., Jacobs, J. J. L., Shah, G. M., et al. (2016). PARP1 Links CHD2-Mediated Chromatin
Expansion and H3.3 Deposition to DNA Repair by Non-homologous End-Joining. *Molecular Cell* 61,
547–562.

Mansuroglu, Z., Benhelli-Mokrani, H., Marcato, V., Sultan, A., Violet, M., Chauderlier, A., Delattre, L.,
Loyens, A., Talahari, S., Bégard, S., et al. (2016). Loss of Tau protein affects the structure,
transcription and repair of neuronal pericentromeric heterochromatin. *Scientific reports* 6, 33047.

Martens, J. H. A., O'Sullivan, R. J., Braunschweig, U., Opravil, S., Radolf, M., Steinlein, P. and Jenuwein,
T. (2005). The profile of repeat-associated histone lysine methylation states in the mouse
epigenome. *The EMBO Journal* 24, 800–812.

Matsui, T., Leung, D., Miyashita, H., Maksakova, I. A., Miyachi, H., Kimura, H., Tachibana, M., Lorincz,
M. C. and Shinkai, Y. (2010). Proviral silencing in embryonic stem cells requires the histone
methyltransferase ESET. *Nature* 464, 927–931.

Mendiratta, S., Gatto, A. and Almouzni, G. (2019). Histone supply: Multitiered regulation ensures
chromatin dynamics throughout the cell cycle. *J Cell Biol* 218, 39–54.

Michaelson, J. S., Bader, D., Kuo, F., Kozak, C. and Leder, P. (1999). Loss of Daxx, a promiscuously
interacting protein, results in extensive apoptosis in early mouse development. *Genes &*
Development 13, 1918–1923.

Miyanari, Y., Ziegler-Birling, C. and Torres-Padilla, M.-E. (2013). Live visualization of chromatin
dynamics with fluorescent TALEs. *Nat Biotechnology* 20, 1321–1324.

Nakatani, T., Yamagata, K., Kimura, T., Oda, M., Nakashima, H., Hori, M., Sekita, Y., Arakawa, T.,
Nakamura, T. and Nakano, T. (2015). Stella preserves maternal chromosome integrity by inhibiting
5hmC-induced γH2AX accumulation. *EMBO reports* 16, 582–589.

Pailles, M., Hirlemann, M., Brochard, V., Chebrout, M., Oudin, J.-F., Marks, H., Jouneau, A. and Bonnet-
Garnier, A. (2022). H3K27me3 at pericentromeric heterochromatin is a defining feature of the early
mouse blastocyst. *Sci Rep-uk* 12, 13908.

Park, J., Lee, H., Han, N., Kwak, S., Lee, H.-T., Kim, J.-H., Kang, K., Youn, B. H., Yang, J.-H., Jeong, H.-J., et
al. (2018). Long non-coding RNA ChRO1 facilitates ATRX/DAXX-dependent H3.3 deposition for
transcription-associated heterochromatin reorganization. *Nucleic acids research* 46, 11759–11775.

Peric-Hupkes, D., Meuleman, W., Pagie, L., Bruggeman, S. W. M., Solovei, I., Brugman, W., Gräf, S.,
Flicek, P., Kerkhoven, R. M., Lohuizen, M. van, et al. (2010). Molecular maps of the reorganization
of genome-nuclear lamina interactions during differentiation. *Molecular Cell* 38, 603–613.

Peters, A. H., O'Carroll, D., Scherthan, H., Mechtler, K., Sauer, S., Schöfer, C., Weipoltshammer, K.,
Pagani, M., Lachner, M., Kohlmaier, A., et al. (2001). Loss of the Suv39h histone methyltransferases
impairs mammalian heterochromatin and genome stability. *Cell* 107, 323–337.

Pinheiro, I., Margueron, R., Shukeir, N., Eisold, M., Fritzsch, C., Richter, F. M., Mittler, G., Genoud, C.,
Goyama, S., Kurokawa, M., et al. (2012). Prdm3 and Prdm16 are H3K9me1 methyltransferases
required for mammalian heterochromatin integrity. *Cell* 150, 948–960.

Probst, A. V., Okamoto, I., Casanova, M., Marjou, F. E., Baccon, P. L. and Almouzni, G. (2010). A strand-
specific burst in transcription of pericentric satellites is required for chromocenter formation and
early mouse development. *Developmental Cell* 19, 625–638.

Ran, F. A., Hsu, P. D., Wright, J., Agarwala, V., Scott, D. A. and Zhang, F. (2013). Genome engineering
using the CRISPR-Cas9 system. *Nature protocols* 8, 2281–2308.

Sadic, D., Schmidt, K., Groh, S., Kondofersky, I., Ellwart, J., Fuchs, C., Theis, F. J. and Schotta, G. (2015).
Atrx promotes heterochromatin formation at retrotransposons. *EMBO reports* 16, 836–850.

Saksouk, N., Barth, T. K., Ziegler-Birling, C., Olova, N., Nowak, A., Rey, E., Mateos-Langerak, J., Urbach,
S., Reik, W., Torres-Padilla, M.-E., et al. (2014). Redundant mechanisms to form silent chromatin at
pericentromeric regions rely on BEND3 and DNA methylation. *Molecular Cell* 56, 580–594.

Salsman, J., Rapkin, L. M., Margam, N. N., Duncan, R., Bazett-Jones, D. P. and Dellaire, G. (2017).
Myogenic differentiation triggers PML nuclear body loss and DAXX relocalization to chromocentres.
Cell death & disease 8, e2724.

Song, Y. J., Choi, J. H. and Lee, H. (2015). Setdb1 is required for myogenic differentiation of C2C12
myoblast cells via maintenance of MyoD expression. *Molecules and cells* 38, 362–372.

Strom, A. R., Emelyanov, A. V., Mir, M., Fyodorov, D. V., Darzacq, X. and Karpen, G. H. (2017). Phase
separation drives heterochromatin domain formation. *Nature* 547, 241–245.

Szenker, E., Ray-Gallet, D. and Almouzni, G. (2011). The double face of the histone variant H3.3. *Cell*
research 21, 421–434.

Tessier, S., Ferhi, O., Geoffroy, M.-C., González-Prieto, R., Canat, A., Quentin, S., Pla, M., Niwa-
Kawakita, M., Bercier, P., Rérolle, D., et al. (2022). Exploration of nuclear body-enhanced
sumoylation reveals that PML represses 2-cell features of embryonic stem cells. *Nat Commun* 13,
5726.

Therizols, P., Illingworth, R. S., Courilleau, C., Boyle, S., Wood, A. J. and Bickmore, W. A. (2014).
Chromatin decondensation is sufficient to alter nuclear organization in embryonic stem cells.
Science 346, 1238–1242.

Tosolini, M., Brochard, V., Adenot, P., Chebrout, M., Grillo, G., Navia, V., Beaujean, N., Francastel, C.,
Bonnet-Garnier, A. and Jouneau, A. (2018). Contrasting epigenetic states of heterochromatin in the
different types of mouse pluripotent stem cells. *Scientific reports* 8, 5776.

Walter, M., Teissandier, A., Pérez-Palacios, R. and Bourc'his, D. (2016). An epigenetic switch ensures
transposon repression upon dynamic loss of DNA methylation in embryonic stem cells. *eLife* 5, R87.

Wang, L., Gao, Y., Zheng, X., Liu, C., Dong, S., Li, R., Zhang, G., Wei, Y., Qu, H., Li, Y., et al. (2019). Histone
Modifications Regulate Chromatin Compartmentalization by Contributing to a Phase Separation
Mechanism. *Molecular Cell* 76, 646-659.e6.

Wijchers, P. J., Geeven, G., Eyres, M., Bergsma, A. J., Janssen, M., Verstegen, M., Zhu, Y., Schell, Y.,
Vermeulen, C., Wit, E. de, et al. (2015). Characterization and dynamics of pericentromere-
associated domains in mice. *Genome Research* 25, 958–969.

Wossidlo, M., Nakamura, T., Lepikhov, K., Marques, C. J., Zakhartchenko, V., Boiani, M., Arand, J.,
Nakano, T., Reik, W. and Walter, J. (2011). 5-Hydroxymethylcytosine in the mammalian zygote is
linked with epigenetic reprogramming. *Nature communications* 2, 241.

Ying, Q.-L., Wray, J., Nichols, J., Batlle-Morera, L., Doble, B., Woodgett, J., Cohen, P. and Smith, A.
(2008). The ground state of embryonic stem cell self-renewal. *Nature* 453, 519–523.

Zhu, Q., Pao, G. M., Huynh, A. M., Suh, H., Tonnu, N., Nederlof, P. M., Gage, F. H. and Verma, I. M.
(2011). BRCA1 tumour suppression occurs via heterochromatin-mediated silencing. *Nature* 477,
179–184.

Ziegler-Birling, C., Helmrich, A., Tora, L. and Torres-Padilla, M.-E. (2009). Distribution of p53 binding
protein 1 (53BP1) and phosphorylated H2A.X during mouse preimplantation development in the
absence of DNA damage. *The International journal of developmental biology* 53, 1003–1011.

Legends

Fig. 1 | DAXX is essential for ESC survival upon ground-state conversion and is required for transcriptional repression of peripheral heterochromatin. (A) Schematic of the *Daxx* locus and the of the guide RNA targeting exon 3. Left, RT-qPCR experiment on DAXX RNA in WT cells or cells after CrispR/cas9 editing on *Daxx*. The mean of two biological replicates with SEM are shown. Right, western blot with anti-DAXX antibody, anti-LAMIN-B1 and anti-H3.3 served as loading control. One experiment from at least 2 biological replicates is shown. **(B)** Quantification of the number of cells (in %) compared to WT in each culture condition, serum, after 4 days (4d) or 8 days (8d) of 2iV conversion and after 5 days of Retinoic Acid differentiation (Diff.) Histogram represents mean with SEM of 3 biological replicates for each condition except Diff. (mean of two biological replicates). T-tests were used for statistical analysis. **(C)** RNA-seq tracks of 2 Lamin Associated Domains (LAD) of WT ESCs (blue) and *Daxx* KO (orange) under serum and 2iV conditions. **(D)** Boxplots of the log2 fold change of mRNA levels of genes located in LAD (LAD, orange) and non-LAD (CEN, blue) regions in <u>*Daxx*</u> KO over WT ESCs, grown in 2iV, serum and diff. media. Two-sided Mann-Whitney tests was used for statistical analysis **(E)** Representative nuclei after immuno-detection of H3K9me2 (red) and counterstaining with DAPI (blue) in WT and *Daxx* KO ESCs in serum and 2iV (scale bar = 2µm). Linescans below represent straighten H3K9me2 signal at the nuclear periphery (black). **(E)** Boxplots of H3K9me2 signal quantification enrichment at the nuclear periphery from 2 independent replicates. Two-sided Mann-Whitney tests was used for statistical analysis.

Fig. 2 | DAXX is recruited to PCH in 2iV converted ESCs. (A) Heatmap for different classes of transposable elements. Data are mean from RNA-seq experiments on 3 biological replicates. **(B)** Representative nuclei of ESCs grown in serum or 2iV immuno-detected with H3K9me3 (red) and DAXX (green). Nuclei were counterstained with DAPI (cyan). Scale bar = 2µm. **(C)** Boxplot of mean DAXX levels at pericentromeric heterochromatin (PCH) in WT ESCs grown in serum (white) or in 2iV (red) from 3 biological replicates. **(D)** Distribution curve of DAXX intensity observed at individual PCH foci in WT ESCs grown in serum (white) or in 2iV

(red) from 3 independent replicates. Dashed line indicates the threshold used to defined DAXX-

positive PCH. **(E)** Histogram of the average fraction of the cell population with 0, 1, 2 and 3 or

more, DAXX-positive PCH foci in WT ESCs in serum and 2iV. SEM are shown. **(F)** Boxplots

of DAXX-negative (neg) and DAXX-positive (pos) PCH foci surface in WT ESCs grown in

serum (white) or in 2iV (red). **(G)** Boxplots of H3K9me3 intensity in DAXX-negative (neg) and

DAXX-positive (pos) PCH foci in WT ESCs grown in serum (white) or in 2iV (red). Mann-

Whitney tests were used for statistical analysis in C, F and G.

Fig. 3 | DAXX maintains heterochromatin organization in pluripotent cells. (A)

Representative major satellite (MajSat) DNA FISH pictures of WT and Daxx KO ESCs in

serum, 2iV and differentiation (Diff.) conditions. **(B)** Quantification of the number of major

satellite foci per nucleus in WT (blue) or *Daxx* KO (yellow) ESC. n=total number of nuclei

analyzed from at least 3 biological replicates. **(C)** Representative immunoFISH pictures of

major satellite and Flag for *Daxx* KO cells transfected with TALE$_{MajSat}$-Δ or TALE$_{MajSat}$-DAXX.

(D) Quantification of the number of major satellite foci detected in the focal plane. n= total

number of nuclei analyzed from 4 biological replicates. **(E)** Top, scheme depicting the two

different analyses used in live-imaging experiments. Bottom, representative pictures from

FRAP experiments performed on WT and *Daxx* KO cells transfected with GFP-HP1α. t=-5''

corresponds to pre-bleach fluorescence. t=0 corresponds to the laser bleach pulse. t=4'' and

t=23'' correspond to post-bleach recovery images after 4 and 23 seconds respectively. **(F)**

Quantification of half-recovery times in seconds for individual nuclei. n=total number of nuclei

analyzed from 2 biological replicates. **(G)** Representative pictures of GFP-HP1α and its

corresponding variance in fluorescence intensities along time in serum or 2iV, WT or *Daxx* KO

ESCs. White dashed squares display magnification of individual chromocenters. Color

correspondence of signal variance strength (from high to low) is shown. Scale bar, 5µm. **(H)**

Graph displaying the variance intensities along measured on 1µm lines traced above non-

bleached foci. Two-sided Mann-Whitney tests were used for statistical analysis in B, D and F.

Fig. 4 | PML protects heterochromatin in ground-state ESCs. (A) Representative field of WT ESCs grown in serum or 2iV immuno-detection of H3K9me3 (red), PML (green), DAXX (purple) and counterstaining with DAPI (cyan). White squares highlight the absence (serum) or the presence (2iV) of PML-rim around H3K9me3 foci. **(B)** Histogram of the average fraction of cell population showing either 0, 1, 2 and 3 or more, PML-rims, with SEM, in 2i converted WT ESCs (n= total number of nuclei). **(C)** Representative major satellite DNA FISH pictures of *Pml* KO ESCs in serum, 2iV and differentiation (dif.) conditions. **(D)** Quantification of the number of major satellite foci per nucleus. WT values (blue) are taken from Fig. 3B. n=total number of nuclei analyzed from at least 3 biological replicates. **(E)** Representative nuclei immuno-detection of H3K9me2 (red) and counterstaining with DAPI (blue) in *Pml* KO EScs grown in serum and 2iV (scale bar = 2µm). Linescan represent straighten H3K9me3 signal at the nuclear periphery (black). **(F)** Boxplots of H3K9me2 signal quantification enrichment at the nuclear periphery. WT values (blue) are taken from Fig. 1F. n=total number of nuclei analyzed from at least 3 biological replicates. **(G)** RT-qPCR RNA quantification of LAD genes in WT and *Pml* KO ESCs in serum and upon 2iV conversion. Histogram represents means normalized to WT-serum with SEM of 3 biological replicates. T-tests were used for statistical analysis **(H)** Quantification of the number of cells compared to WT in each culture condition. Histogram represents mean with SEM of 3 biological replicates. Two-sided Mann-Whitney tests were used for statistical analysis in D and F.

Fig. 5 | DAXX recruits Setdb1 and mediates chromocenter clustering via H3.3K9me3 modification. (A) Representative immunofluorescence pictures for SETDB1, DAXX and PML in WT or *Daxx* KO ESCs upon 2iV conversion. **(B)** Quantification of the mean number of PML or SETDB1 foci at chromocenters in WT and *Daxx* KO ESCs. n= total number of nuclei analyzed from two independent replicates. Chi-square tests were used for statistical analysis. **(C)** Representative immunoFISH pictures for H3K9me3, PML and major satellites in 2iV and serum of Suv39H1/2 dKO (double knock out) ESCs. White dashed squares highlight a major satellite focus. **(D)** Cumulated fluorescence intensities profile of H3K9me3 positive PCH foci in

cells grown in serum or 2iV for either H3K9me3, PCH or PML signal. n=total number of nuclei

analyzed from at least 2 biological replicates. **(E)** Representative immunofluorescence pictures

for H3K9me3 and Flag in Suv39dKO serum ESCs transfected with TALE$_{MajSat}$-Δ, TALE$_{MajSat}$-

SUV39H1, TALE$_{MajSat}$-SETDB1, TALE$_{MajSat}$-DAXX or TALE$_{MajSat}$-DAXX^{Y222A}. **(F)** quantification

of the number of PCH foci observed in the different transfection conditions described in E. **(G)**

Representative immunoFISH pictures for major satellite and Flag in *Daxx* KO serum ESCs co-

transfected with siRNA against H3.3, TALE$_{MajSat}$-DAXX and either no additional construct or

H3.3WT, H3.3G90M or H3.3K9A. **(H)** Quantification of the number of major satellite foci

detected of medium focal plane observed in the different transfection conditions described in

G. For comparison, distribution T$_{MS}$-DAXX without H3.3 knockdown data comes from Fig. 3D.

n=total number of nuclei analyzed from 2 biological replicates. Two-sided Mann-Whitney tests

were used for statistical analysis in F and H.

Fig. 6 | Model for heterochromatin maintenance by DAXX in ESCs.

During ground-state conversion of ESCs, DAXX accumulates at PCH and is surrounded by a

PML shell. DAXX recruits the histone methyltransferase SETDB1 to promote H3.3K9me3 to

maintain the pericentromeric heterochromatin. In *Daxx* and *Pml* KO cells, the 3D-organization

of heterochromatin is impaired. The peripheral heterochromatin H3K9me2 fails to accumulate

properly at the nuclear edge leading to promiscuous expression of genes located in LADs. The

clustering and the physical properties of pericentromeric heterochromatin is impacted,

impairing transcriptional repression of major satellites. The global defect of heterochromatin

observed in these cells prevents prolonged culture of these cells in ground-state condition.

1031

1032 **Supplemental material legends**

1033 **Fig. S1 | Impact of *Daxx* deletion on ground-state conversion. (A)** Representative field of

1034 WT and *Daxx* KO ESCs after IF against DAXX (green) and counterstaining with DAPI (cyan).

1035 Scale bar = 5µm. **(B)** Dot blot using 5mC antibody to assess global DNA methylation levels

1036 upon 2iV conversion. One experiment is shown. Histogram displays mean with SEM from 3

independent replicates. **(C)** Volcano plot representing the number of differentially expressed genes in ground-state, serum or differentiated cells from RNA-seq experiments as in C. Differentially expressed genes in red for down-regulated or blue for up-regulated correspond to Log2(FoldChange(KO/WT)) <-1 or >1 with p-value<0.05. **(D)** Heatmaps representing genes displaying strongest transcriptional changes upon 2iV conversion (left) or upon differentiation (right) normalized by transcription in serum conditions. Same set of genes for the *Daxx* KO ESCs. Data are from 3 biological replicates of RNA-seq experiments. **(E)** Volcano plots representing the number of differentially expressed genes upon 2iV conversion of WT and *Daxx* KO ESCs. Previously identified 2iV conversion markers are in red for up-regulated or blue for down-regulated (Blaschke et al., 2013).

Fig. S2 | Impact of *Daxx* deletion on nuclear periphery. (A) RT-qPCR RNA quantification LAD genes in WT and *Daxx* KO ESCs in serum and upon 2iV conversion. Histogram represents mean normalized to WT-serum with SEM of 3 biological replicates. **(B)** Left, representative image of ESCs after IF against H3K9me2. Top right, linescan represents straighten H3K9me2 signal at the nuclear periphery (black). Bottom right, Intensity profiles of IF signals measured on linescans at the periphery (blue) or in the center (orange) of the nucleus. Ratio between mean intensity at the periphery and in the center was used to determined peripheral enrichment. **(C)** Representative nuclei after immuno-detection of LaminB1 (red) and counterstaining with DAPI (blue) in WT and *Daxx* KO ESCs in serum and 2iV (scale bar = 2μm). Linescan represent straighten LaminB1 signal at the nuclear periphery (black). **(D)** Boxplots of LaminB1 signal quantification enrichment at the nuclear periphery from 2 independent replicates. Mann-Whitney-tests were used for statistical analysis.

Fig. S3 | Impact of *Daxx* deletion on DNA repeats silencing. RT-qPCR RNA quantification of DNA repeats. L1Tf , IAP elements and major satellites in WT and *Daxx* KO ESCs in serum, and upon differentiation or 2i+VitC (2iV) conversion. Histogram represents mean with SEM of 3 biological replicates.

Fig. S4 | Quantification of DAXX at chromocenters. (A) Representative immunoFISH picture with H3K9me3 and major satellites DNA. **(B)** Dot plots of Pearson correlation between H3K9me3 and major satellite signals. **(C)** Representative field of ESCs grown in serum or 2iV immuno-detection of H3K9me3 (red), DAXX (green) and counterstaining with DAPI (cyan). White squares and arrows highlight the presence (2iV) of larger DAXX foci with higher levels of H3K9me3 and DAPI. Scale bar = 10µm. **(D)** Distribution of DAXX foci volume observed in WT ESCs grown in serum (black) or in 2iV (red) from 3 independent replicates. Dashed line indicates the threshold used to defined large or small DAXX foci. **(E)** Boxplots of DAPI signal intensity observed under small and large DAXX foci in serum (white) and 2iV (red) conditions. **(F)** Boxplots of H3K9me3 signal intensity observed under small and large DAXX foci in serum (white) and 2iV (red) conditions. **(G)** Top, ESCs nucleus after immuno-detection of H3K9me3, DAXX and counterstaining with DAPI. Bottom, for each nucleus, H3K9me3 and DAPI signals were segmented to PCH and nuclear ROIs. DAXX enrichment at PCH was defined by yhe ratio of DAXX intensity under PCH ROIs. Mann-Whitney-tests were used for statistical analysis in B, E and F.

Fig. S5 | Role of DAXX on pericentromeric heterochromatin formation. (A) Left, representative immunoFISH pictures for major satellites and Flag for WT ESCs, grown in serum, transfected with TALE$_{MajSat}$-Δ or TALE$_{MajSat}$-DAXX. Right, quantification of the number of major satellite foci detected in the focal plane and corresponding surface area. n=total number of nuclei for number and foci for sizes analyzed. Two-sided Mann-Whitney tests were used for statistical analysis. **(B)** Left, mobile fractions of GFP-HP1α calculated from data presented in Fig. 4D. n=total number of cells analyzed from 2 biological replicates. Right, fluorescence recovery curves generated for FRAP experiments in WT or *Daxx* KO cells in serum or ground-state condition. **(C)** MNAse digestion for indicated times in minutes (min.) in WT or *Daxx* KO cell lines. Total DNA is revealed on a BET agarose gel and Major satellites via a Southern blot experiment. Right, intensity values of total DNA or major satellite signals,

according to fragment size, from 4min of MNase digestion. One experiment is shown from 2 biological replicates.

Fig. S6 | Quantification of PML at chromocenters. (A) Representative field of ESCs grown in serum or 2iV after immuno-detection of H3K9me3 (red), PML (green) and counterstaining with DAPI (cyan). Scale bar = 10µm. **(B)** Distribution of PML foci volume observed in WT ESCs grown in serum (black) or in 2iV (red) from 3 independent replicates. Dashed line indicates the threshold used to defined large PML foci. **(C)** Boxplots of DAPI signals intensities observed under small and large PML foci in serum (white) and 2iV (red) conditions. **(D)** Boxplot of H3K9me3 signals intensities observed under small and large PML foci in serum (white) and 2iV (red) conditions. **(E)** Representative nucleus of ESCs grown in 2iV after immuno-detection of H3K9me3 (red), PML (green) and counterstaining with DAPI (cyan) presenting a PML-arc around a PCH cluster (white arrow). Scale bar = 2µm. Two-sided Mann-Whitney tests were used for statistical analysis in B and D.

Fig. S7 | Impact of *Pml* deletion on nuclear periphery. (A) Representative field of WT, *Daxx* KO and *Pml* KO ESCs, in serum or 2iV after IF against PML (green) and counterstaining with DAPI (cyan). **(B)** Representative nuclei after immuno-detection of LaminB1 (red) and counterstaining with DAPI (blue) in WT and *Pml* KO ESCs in serum and 2iV (scale bar = 2µm). Linescans represent straighten LaminB1 signal at the nuclear periphery (black). **(C)** Boxplots of LaminB1 signal quantification enrichment at the nuclear periphery from 2 independent replicates. WT values are taken from Fig. S2D. Two-sided Mann-Whitney tests were used for statistical analysis.

Fig. S8 | DAXX recruits SETDB1 to pericentromeres. (A) Representative immunofluorescence pictures for SETDB1, DAXX and PML in WT or *Daxx* KO ESCs in serum condition. **(B)** Top, representative immunofluorescence pictures for SETDB1, DAXX and PML in WT ESCs in serum. (**C**) Representative immunofluorescence pictures for SETDB1 and Flag-TALE in *Daxx* KO serum ESCs transfected with TALE$_{MajSat}$ (t$_{MS}$-Δ), TALE$_{MajSat}$-DAXX (t$_{MS}$-DAXX) or TALE$_{MajSat}$-DAXX^{Y222A} (t$_{MS}$-DAXXY222A). White dashed squares highlight DAPI-

dense Flag-positive chromocenters. **(D)** Quantification of SETDB1 recruitment to Flag-positive

chromocenters. Histogram represents mean with SEM from at least 2 biological replicates.

n=total number of nuclei analyzed. Chi-square tests were used for statistical analysis.

Fig. S9 | Role of H3.3 and H3K9me3 modifications at PCH. (A) Representative

immunoFISH pictures for H3K9me3 and DNA FISH of major satellites (MajSat) in both WT and

Daxx KO ESCs in serum or ground-state (2iV) conditions. **(B)** Mean and SEM of fluorescence

intensity of H3K9me3 measured over major satellite signal. n=total number of nuclei analyzed

from two independent replicates. **(C)** Quantification of the percentage of cells displaying

H3K9me3 recruitment at Flag-DAPI-dense PCH in Suv39dKO serum ESCs transfected with

$TALE_{MajSat}$-Δ, $TALE_{MajSat}$-SUV39H1, $TALE_{MajSat}$-SETDB1, $TALE_{MajSat}$-DAXX or $TALE_{MajSat}$-

$DAXX^{Y222A}$ grown in serum transfected with $TALE_{MajSat}$-Δ, $TALE_{MajSat}$-DAXX or $TALE_{MajSat}$-

DAXXY222A. T-tests were used for statistical analysis. **(D)** Represents signal after Flag-TALE

immunoprecipitation. Bottom line represents input line from total extract. **(E)** Representative

immunofluorescence pictures of HA-H3.3 in WT cells grown in serum transfected with

$TALE_{MajSat}$-Δ, $TALE_{MajSat}$-DAXX or $TALE_{MajSat}$-$DAXX^{Y222A}$. **(F)** quantification of H3.3 recruitment

at DAPI-dense chromocenters in the indicated Tale-transfected cells. Histogram represent

mean with SEM from at least 2 biological replicates. n=total number of nuclei analyzed. T-tests

were used for statistical analysis. **(G)** Western blot showing the amount of TALE and H3.3

variants in *Daxx* KO ESCs grown in serum transfected with the indicated constructions.

Figure 1. Canat et al. 2023

Figure 2. Canat et al. 2023

Figure 3. Canat et al. 2023

Figure 4. Canat et al. 2023

Figure 5. Canat et al. 2023

Figure 6. Canat et al. 2023