Title: Using total RNA quality metrics for time since deposition estimates in degrading
bloodstains.

Authors: Colin I. Elliott[a,b*], Theresa E. Stotesbury[c], Aaron B.A. Shafer[a,d]

[a] Department of Forensic Science, Trent University, 1600 West Bank Drive, Peterborough, ON,
K9L 0G2, Canada

[b] Applied Bioscience Graduate Program, Faculty of Science, Ontario Tech University, Oshawa,
ON L1G 0C5, Canada

[c] Faculty of Science, Forensic Science, Ontario Tech University, 2000 Simcoe St N, Oshawa, ON,
L1G 0C5, Canada

[d] Environmental and Life Sciences Graduate Program, Trent University, 1600 West Bank Drive,
Peterborough, ON, K9L 0G2, Canada

* **Corresponding author:** colinelliott@trentu.ca

Declaration of Competing Interest

The authors report no declarations of interest.

Acknowledgements

The authors would like to thank Reni Verasztó for helping with laboratory work and data

collection. The authors would also like to acknowledge Alon Gabriel, Jesse Wolf, and Marie-

Laurence Cossette of the Wildlife and Applied Genomics laboratory group at Trent University

for their insights and feedback on writing, visualizations, and statistical analyses. The authors

also thank Otonabee Meat Packers for providing the blood samples.

HIGHLIGHTS

- Amplification-free analysis of total RNA in degrading bloodstains.

- Short-term RNA degradation assessment using high-resolution size measurements.

- Total RNA quality and quantity metrics were assessed across a one-week.

- Total RNA quality metrics demonstrated the strongest timewise trends.

- Biological replicates produced similar results for RNA quality metrics.

ABSTRACT

Determining the time since deposition (TSD) of bloodstains would provide forensic scientists with critical information regarding the timeline of the events involving bloodshed. The physicochemical changes occurring to biomolecules as a bloodstain degrades can be used to approximate the TSD of bloodstains. Our study aims to quantify the timewise degradation trends and temperature dependence found in total RNA from bloodstains without the use of amplification, expanding the scope of the RNA TSD research which has previously targeted mRNA molecules. Whole bovine blood was stored in plastic microcentrifuge tubes at 21˚C or 4˚C and tested over different timepoints spanning one week. Nine RNA metrics were visually assessed and quantified using linear and mixed models; the RNA Integrity Number equivalent (RINe) and the DV200 demonstrated strong negative trends over time and statistical independence. The RINe model fit was high ($R^2 = 0.60$), and while including the biological replicate as a random effect increased the fit for all RNA metrics, no significant differences were found between biological replicates stored at the same temperature for the RINe and DV200 metrics. Importantly, this suggests that these standardized metrics can likely be directly

compared between scenarios and individuals, with DV200 having an inflection point at ~28 hrs.

This study provides a novel approach for blood TSD estimates, producing metrics that are not

affected by inter-individual variation and improving our understanding of the rapid degradation

occurring in bloodstains.

KEYWORDS: RNA Integrity Number equivalent and RINe, Time since deposition and TSD,

DV200, Bloodstain, Amplification-free, Aging

Introduction

Accurately determining the age of biological evidence, including soft tissue and bodily

fluids, remains a complex task in forensic investigations (1,2). Bloodstained evidence is

commonly observed at the scenes of crimes involving bloodshed, and determining the age, or

time since deposition (TSD) of a bloodstain, would provide context in criminal investigations,

including the development of investigative leads and the assessment of alibis (1–4). Currently,

bloodstains observed at crime scenes can be subjected to bloodstain pattern analysis, which

provides potential interpretations of the physical events that gave rise to the bloodshed (5).

DNA profiling, used for source attribution purposes, can be carried out from collected

bloodstains (6–9); however, these tools do not provide TSD estimates of blood, requiring

additional analyses. Bloodstains degrade in a series of complex processes, physical and

chemical changes are influenced by many factors, starting from the point of their formation on

a surface onwards (10–16). Previous studies have developed chemometric (2,13,14,17) and

genetic methods (3,18–20) that estimate the TSD of bloodstains in controlled laboratory

conditions. While these methods produce time-dependent responses, studies generally report

high variability in responses between samples, biological replicates and environmental

conditions, suggesting that the accuracy of TSD estimates might be increased by combining

metrics obtained from different biomolecules and/or using methods that are insensitive to

biological replicate variation (1,3,13,14,17–19,21).

Ex-vivo, blood undergoes a series of time-dependent physicochemical and

morphological changes, such as hemoglobin oxidation and conformational changes (13,22),

drying (12,23), clotting (24,25), and DNA and RNA degradation (3,15,26–28). Total RNA

encompasses ribosomal RNAs (rRNA), messenger RNAs (mRNA), transfer RNAs (tRNA) and

microRNAs, among others (29,30). Eukaryotic rRNA makes up for 80% of the total RNA in a cell

and consists of 5S, 5.8S, 18S and 28S rRNA (31). Messenger RNA expression patterns have been

used to identify body fluids found at crime scenes (32,33) and might also correlate with time

(34). Fu *et al.* (20) found that, when analyzing blood-specific mRNA transcripts, TSD for

bloodstains could be estimated to within a period of 2 to 6 weeks, depending on the TSD of the

sample. Anderson *et al.* (3) distinguished 6-day old blood from fresh and aged (30 days) blood,

while Lech *et al.* (18) classified deposition by three time-of-day periods with high prediction

accuracy. High throughput sequencing (HTS) by Weinbrecht *et al.* (19) demonstrated that

mRNA transcripts decreased over time, with blood-specific transcripts detected up to 12

months after deposition. Salzmann *et al.* (35) also used HTS to investigate the timewise changes

in the quality of the amplified RNA; the authors observed a marked decrease in mRNA

transcript integrity six months after deposition, highlighting the potential applicability of total

RNA quality metrics for TSD purposes (35). Collectively, these studies demonstrated the

timewise degradation of small RNA species in bloodstains and highlighted the surprising long-

term survival of RNA, supporting the applicability and implementation of techniques targeting

nucleic acid degradation in time series. We ask if features of quick RNA degradation can be

captured in the same way.

Most RNA studies to date have used PCR amplification to target specific mRNAs and

how they change in a degrading bloodstain over a long period of time (3,19,20,35). The vast

majority of mRNA molecules have a half-life of just minutes (36); therefore, genomic DNA

contamination is a concern in experiments where amplification is used to target specific mRNA

(37). Amplification-free analysis of RNA might provide insight into the naturally occurring RNA

degradation in blood without the risk of introducing any amplification biases or stochastic

effects seen while using RT-PCR and RT-qPCR (36,38,39). Accordingly, our study takes a holistic,

non-targeted and amplification-free approach by analyzing the quality (ribosomal ratios, RNA

integrity number equivalent; RINe, DV200) and quantity metrics (RNA size ranges, total

concentration, rRNA peak concentration) of total RNA in bloodstains, while simultaneously

quantifying the effect of temperature on RNA degradation and TSD models. Not only does

focusing on total RNA have the advantage of eliminating the stochastic effects and

amplification biases caused by reverse transcriptase and PCR, it can also circumvent potential

PCR inhibition from the heme in blood (38–40).

Materials and Methods

Blood sample collection

Bovine blood was collected from Otonabee Meat Packers, an abattoir in Peterborough,

Ontario, Canada. Acid dextrose anticoagulant solution A (ACD-A) was added to bovine blood at

a concentration of 12.5% v/v (41). Blood treated with ACD anticoagulants has demonstrated

better nucleic acid extraction efficiencies than heparin-based anticoagulants and has not been

shown to affect the cell counts of the different blood components (42). The ACD-A used in this

study was not shown to impact RNA metrics such as concentration and quality over time (Table

S1) and allowed for more accurate pipetting of the blood. ACD-A was made by dissolving 6.6

grams of sodium citric dextrose, 2.4 grams of citric acid, and 6.68 grams of dextrose anhydrous,

all purchased from ACP Chemicals Inc., in 500 mL of distilled water. Bovine blood was collected

in an amber Nalgene bottle containing the anticoagulant, which was subsequently placed on ice

for transportation back to the laboratory (approximately 15 minutes).

Blood deposition

50 µL of blood was deposited into 1.5 mL plastic microcentrifuge tubes. Samples were

left to air dry by keeping the microcentrifuge tubes lids open until RNA extraction. Samples

were stored at one of two temperatures: room temperature (~ 21˚C) or refrigerated at 4˚C for a

designated amount of time. The room temperature storage condition had an approximate 40%

relative humidity (RH), while the RH was likely slightly greater for samples stored at 4˚C. Five

time-series experiments were performed, three at 21˚C and two at 4˚C; blood from a different

cow (biological replicate) was used for each experiment. RNA extractions were completed at 0

hours (time of initial pipetting; ~ 30 minutes post-blood collection) and across various

timepoints (Table 1); the goal here was to compare the timewise degradation trends for blood

deposited at different temperatures. Five technical replicates were used for each of the

timepoints, resulting in a total of 200 assayed samples.

Table 1. Overview of the time series conducted and each experiment's storage temperature for

the bloodstains. Each biological replicate represents a different cow.

Biological Replicate	Temperature	Sample Size	Timepoints (hours after deposition)
A	21˚C	50	0,1.5,3,6,9,24,48,72,96,168
B	21˚C	50	0,1.5,3,6,9,24,36,48,72,96
C	21˚C	30	0,6,20,24,28,36
D	4˚C	35	0,3,6,9,24,48,72
E	4˚C	35	0,3,9,24,48,72,96

RNA extraction

Total RNA was extracted from the deposited blood using the PureLink Total RNA Blood

Purification kit by Invitrogen (43). The volume of lysis buffer (L5) recommended in the

manufacturer's protocol was doubled to minimize blood clotting. An optional on-column DNase

incubation was completed following the manufacturer's protocol. Total RNA was eluted with 30

μL of RNase-free water and prepared for analysis on the Agilent Technologies 4200 TapeStation

(44). 1 μL of the high sensitivity RNA screen tape sample buffer was mixed with 2 μL of the total

RNA extracts following the manufacturer's protocol for the *High Sensitivity RNA ScreenTape*

Assay (44).

Automated gel electrophoresis

RNA was analyzed using high-sensitivity automated gel electrophoresis with the Agilent

Technologies 4200 TapeStation (44). The TapeStation is a highly sensitive instrument that

detects RNA down to 100 pg/μl while offering a quantitative range beginning at 500 pg/μl (45).

It separates different-sized fragments that are tagged with an intercalating fluorescent dye by

automated electrophoresis, allowing for the quantification of specifically-sized RNA (45,46). A

ladder containing fragments of known size was also loaded next to the samples; the software

compared band locations and assigned 18S and 28S rRNA peaks to the RNA samples (47).

RNA metrics

A series of RNA metrics were collected and examined using the TapeStation Analysis Software

A.02.02 (Table S2). Of note, the RINe metric was generated by an algorithm that calculates the

ratio of the height of the 18S rRNA peak (in fluorescent units; FU) to the background RNA signal

(FU), which represents the signal for fragments found in the fast zone ranging from 200 bp to

1.8 kb (45,47). The RINe metric is a dimensionless number ranging from 1 to 10, with 10

representing intact and high-quality RNA, while a value of 1 represents low-quality and highly

fragmented RNA (45,48). Another RNA quality metric, the DV200, represents the percentage of

RNA fragments greater than 200 nucleotides (49). A custom RNA bin was made to estimate

concentration from 200 bp to 500 bp fragments, presumably representing the shorter mRNAs

extracted from our samples (50).

Statistical analyses

Correlation matrices were created prior to building models to ensure that the response

variables (RNA metrics) included in the analyses were not highly correlated to each other

(Pearson's r < |0.70|). Variables showing independence and a correlation to time were used to

build a series of linear mixed-effects models, with time (natural log-transformed) and

temperature as fixed effects and biological replicate (individual cow) as a random effect. An

interaction term between log-transformed time and the temperature was also included in our

models. We assigned timepoint zero a value of 0.1 (6 minutes) prior to the logarithm

transformation. Effect size, R^2, and p-values were recorded for each model.

We ran an unpaired two-sample Wilcoxon test, a non-parametric test, to determine

whether there was a difference in the biological replicates undergoing the same experimental

conditions across all timepoints (i.e. test for between-individual variance in RNA metrics). The

non-linearity of the RNA metrics over time was also explored by fitting non-linear curves and

assessing the impact on the residual standard error. We note the regressions do not require

shared time points between biological replicates to estimate slope, intercept, and inflection

points. All statistical analyses and data visualizations were created using R Version 4.0.3. Raw

data and scripts are publicly available and deposited at https://gitlab.com/WiDGeT_TrentU.

Results

We collected nine RNA metrics on 200 deposited bloodstains; four RNA metrics were

retained for statistical analyses as they were not correlated to each other (Pearson's r < |0.70|;

Fig. 1). The retained metrics included the RINe, the DV200, 28S/18S rRNA peak area ratio, and

total RNA concentration (Table 2).

Fig. 1 - Correlations between the nine RNA metrics obtained from the 4200 TapeStation RNA

analysis. The size of the circle increases with a greater correlation between variables. Blue-

coloured circles indicate a positive correlation between RNA metrics, while red-coloured circles

indicate a negative correlation between them. The shading of the circle increases with a greater

correlation, with light colours indicating a weak correlation and darker colours indicating a

strong correlation.

Table 2. Statistics for the mixed models of the four quantified RNA metrics. Note that there was

an interaction between log-transformed time and temperature. Biological replicate was used as

a random effect. Each regression was computed individually for each RNA metric and each row

represents a separate model. Because of the interaction term, log(Time) in the table represents

the relationship between the response variable and time at 21˚C; Temperature (4˚C) represents

the difference between the intercept at both temperatures; and log(Time)*Temperature

reflects the difference in correlation between retained RNA metrics and time between 21˚C and

4˚C. P-values below 0.05 are bolded.

RNA metrics	Marginal R^2	Conditional R^2	log(Time)			Temperature (4˚C)			log(Time)*Temperature		
			Estimate	Confidence Interval	p-value	Estimate	Confidence Interval	p-value	Estimate	Confidence Interval	p-value
RINe	0.51	0.60	-0.51	-0.58 – -0.43	**<0.001**	0.12	-0.74 – 0.98	0.78	0.17	0.04-0.29	**0.01**
DV200	0.31	NA	-2.4	-2.99 – -1.80	**<0.001**	-0.47	-3.59 – 2.64	0.77	1.99	1.03 – 2.96	**<0.001**
28S/18S Peak Area Ratio	0.061	0.24	0.34	-0.21 – 0.89	0.22	0.35	-3.50 – 4.20	0.86	0.19	-0.54 – 0.91	0.62
Concentration	0.069	0.30	-885.42	-1305.24 – -465.60	**<0.001**	-2142.06	-7554.02 – 3269.90	0.44	789.89	116.22 – 1481.56	**0.022**

RNA Integrity Number equivalent

The RINe metric demonstrated the best fit model (R^2 = 0.60; Table 2) and a relationship

with log-transformed time (Table 2). RINe values were influenced by storage temperature,

however, the effect of temperature on RINe varied according to time, as seen by the significant

interaction effect in Table 2. The addition of biological replicate (donor) as a random effect

slightly increased the model's goodness-of-fit (marginal vs conditional R^2 in Table 2); this was

driven by a temperature-donor effect, as RINe values obtained from the different biological

replicates at the same temperature were not different from each other (Table 3),

demonstrating consistency across individuals in the same treatment. The timewise changes in

RINe at different temperatures (Fig. 2) showed 21˚C caused more degradation (steeper slope)

than the samples stored at 4˚C (Fig. 2).

Table 3. Unpaired two-sample Wilcoxon tests between the different biological replicates for the

RINe, DV200 and total RNA concentration. Significant p-values are bolded.

RNA metric	Biological replicate pairs	Temperature (˚C)	Test statistic (W)	p-value
RINe				
	A-B	21	1030	0.17
	A-C	21	574	0.67
	B-C	21	625	0.37
	D-E	4	335	0.36
DV200				
	A-B	21	1352	0.38
	A-C	21	658	0.14
	B-C	21	619	0.40
	D-E	4	367	0.69
Concentration				
	A-B	21	1592	**0.010**
	A-C	21	742	**0.012**
	B-C	21	626	0.36
	D-E	4	94	**<0.001**

Fig. 2 - Relationship between RINe and log-transformed time (hours). RINe values are grouped

by storage temperature for comparison purposes; points and regression lines are coloured

according to storage temperature. Shaded areas around the regression lines represent the

standard error. The interaction effect between log-transformed time and temperature is not

shown as data is already separated by storage temperature.

DV200

The DV200 demonstrated the second-best model fit ($R^2 = 0.31$); the random effect of

biological replicate did not improve the model (Table 2). DV200 showed a statistically significant

relationship with time and was influenced by storage temperature, the effect of which varied

according to time (Table 2). The positive estimate for the interaction effect indicated that over

time, bloodstains kept at 4˚C provided greater DV200 values relative to those stored at 21˚C

(Table 2, Fig. 3). DV200 values obtained from the different biological replicates at the same

temperature were not different from each other (Table 3), but there was a difference between

DV200 values recovered from bloodstains deposited at 21˚C and 4˚C (W = 2572, p-value =

0.015; Fig. 4A-B). Bloodstains stored at room temperature exhibited greater levels of

fragmentation over time (Fig. S1, Fig. 4A-B). DV200 values from bloodstains deposited at room

temperature also exhibited a non-linear trend over time; we determined that the log-logistic

curve – akin to a dose-response curve – provided the best fit at 21˚C. The residual standard

error (RSE) was smaller (5.9) in the log-logistic curve than the linear model (7.7). The inflection

point of the curve was at 28.1 hours (95% CI: 24.6 – 31.5 hours), suggesting a key change at this

time (Fig 3).

Fig. 3 - Relationship between the DV200 and time and temperature. **A)** Linear regression curve

for the DV200 against log-transformed time at 4˚C. A negative slope is observed. **B)** Log-logistic

curve for the DV200 against time at 21˚C. The curve displayed an inflection point at 28.1 hours,

indicated by the vertical dashed line. Shaded areas in **A)** and **B)** represent the standard error.

Note that the y-axis is scaled differently for **A)** and **B)**. The interaction effect between log-

transformed time and temperature is not shown.

Total RNA concentration

The concentration of total RNA was shown to have a lower model fit than the RINe or

DV200 (Table 2); the addition of biological replicate as a random effect greatly improved the

model's fit, suggesting high inter-individual variation (Table 2). We detected a significant

relationship between total RNA concentration and time and temperature (Table 2, Fig 4).

Importantly, we uncovered differences between concentrations from biological replicates

originating from the same temperature (Table 3, Fig. S2). Density plots demonstrated bimodal

distributions for temperature data, concordant with the result that RNA concentrations from

different biological replicates at the same temperature were different (Table 3, Fig. S3). Most

concentrations recorded for bloodstains deposited at early timepoints were above the limit of

quantification (500 pg/µL), while many concentrations at room temperature fell below this

threshold as time progressed (Fig. 4).

Fig. 4 – Relationships between total RNA concentration and log-transformed time (hours).

Concentrations are grouped by storage temperature for comparison purposes; points and

regression lines are coloured according to storage temperature. Shaded areas around the

regression lines represent the standard error. Concentration was log-transformed for

visualization purposes. The red dashed line represents the 4200 TapeStation's limit of

quantification (500 pg/μL) and the navy dashed line represents the instrument's limit of

detection (100 pg/μL).

28S/18S rRNA peak area ratios

The 28S/18S rRNA peak area ratio was the retained RNA metric that displayed the

weakest model fit and little relationship to time or temperature (Table 2). Peak area ratios were

only obtained for timepoints up to 24 hours for bloodstains deposited at room temperature

and 72 hours for those stored at 4˚C (Fig. S4); the 28S/18S peak area ratios could not be

obtained for the later timepoints, as the 18S rRNA peak dropped out of the electropherograms.

Discussion

Estimating the TSD of biological fluids found at crime scenes, such as bloodstains,

remains a challenge within the forensic community (1,2). Intra and inter-individual variation in

blood composition, substrate type and storage conditions all influence bloodstain degradation

and must be accounted for to accurately estimate TSD (2). This study provided a novel

approach to the TSD of blood, by using a non-targeted, amplification-free technique within a

short time since deposition period (168 hours). This is contrary to most nucleic acid studies that

used targeted approaches (mRNA transcripts), relied on amplification, and explored longer time

periods (3,19,20). Similarly to what Salzmann *et al.*(35) demonstrated using the integrity of

mRNA transcripts (TINe – transcript integrity number equivalent), we found that total RNA

quality metrics, such as the RINe and DV200, both provided clear relationships with time, with a

minimal effect of biological replicate on the model. A key observation here is that the limited

effect of biological replicate and inflection point at 28 hours of DV200 suggests these could be

universal diagnostic metrics.

309   The RINe is an RNA quality metric that is equal to the ratio of the 18S rRNA peak height

310  to the signal present in the fast zone (approximately 200 bp - 1.8 kb), which represents

311  fragmented RNA (45). This ratio normalized the discrepancies observed in total RNA

312  concentration and had a clear relationship to time (Fig. 2, Table 2). The addition of biological

313  replicate as a random effect in the model only marginally increased the model fit (Table 2),

314  which was driven by a temperature-donor effect (Table2, Table 3). This is important because

315  previous degradation models found a large effect of biological replicate (28), but the

316  standardized nature of RINe makes it a universally comparable metric. Blood composition varies

317  by individual, for example, research on human blood showed that the amount of RNA in

318  healthy subjects varied from 6.7 to 22.7 µg/ml (51). This variability could lead to significant

319  differences in concentrations of total RNA obtained from biological replicates, as we observed

320  during our study (Fig S2). The RINe metric was not affected by this variation (Table 2, Table 3);

321  additionally, the RINe was not highly correlated to total RNA concentration and was

322  independent of sample volume, highlighting the potential for wide-scale applicability. The

323  independent and negative correlation with time suggests the utility of RINe as a useful TSD

324  metric. We also have no reason to suspect that the patterns observed herein would not be

325  present in human blood; bovine blood has previously been optimized for forensic research

326  when acid dextrose anticoagulant solution A (ACD-A) is added at a concentration of 12.5% v/v,

327  as it exhibits similar fluid properties to that of human blood (41). Bovine blood also has a similar

328  genome size and composition to humans (52) and has similar RNA concentration (51,53), white

329  blood cell (WBC) count (human: 3.4-11.6 x 10^3/µL, bovine: 4.9-13.3 x 10^3/µL), and subtypes of

330  WBCs as human blood (neutrophils and lymphocytes are the most abundant in each organism)

(51,54). Confirmatory analyses validating our results using human blood could make this

technique applicable to crime scene samples in the future.

A common theme observed is the preservation effect of cold temperatures. The

bloodstains at 4˚C consistently demonstrated improved quality metrics when compared to the

bloodstains stored at 21˚C. Colder temperatures have been shown to preserve nucleic acids and

slow their degradation (15,16,55), a trend that was reproduced herein by comparing the slopes

observed in the linear models for the RINe and the DV200 (Fig. 2, Fig. 4A, Fig. S1). In both

instances, room temperature produced steeper slopes, indicative of faster degradation.

Previous studies have shown that the DV200 provides an alternative to the RINe when

analyzing highly degraded samples, where electropherograms lack the 18S and 28S rRNA peaks

due to fragmentation (49,56). The DV200 is often used as a quality indicator before library

preparation for high-throughput sequencing, as it is not contingent on rRNA being present to

provide measurements (49,57). Our results indicated that the DV200 was influenced by the

dropout of 18S rRNA with time (Fig. 2); specifically, the values obtained at room temperature

exhibited non-linear behaviour over time that appeared to be directly driven by the

degradation of the 18S rRNA peak occurring after one full day of exposure (Fig. 2B, Fig. S5).

Non-linear relationships between response variables and time have been observed in previous

blood TSD studies (13–15,20), indicating that non-linearity, potentially resulting from

preferential degradation of certain RNA fragments (or parts therein) (20), oxidation and

dehydration (13), is not uncommon. The non-linearity and inflection point of the DV200 values

at room temperature could be exploited as a diagnostic tool to differentiate bloodstains aged

for less than 28 hours from older ones as there is a precipitous loss of RNA that occurs at this

time (Fig. 2B), with most measurements approaching the instrument's limit of quantification

(Fig. 4), implying that little RNA is present in the sample. It is conceivable that the same curve

and diagnostic signature would be present at 4°C over longer periods. Future work should

extend the 4°C sampling period as this could prove useful for crime scene investigation in cold

climates (58).

As this approach analyzed total RNA rather than mRNA, direct comparisons between

these results and those obtained in previous studies can be difficult. Low RNA concentrations

were observed throughout the study, often below the limit of quantification, as time

progressed. The size bin ranging from 200 bp – 500 bp, which should represent the smaller

mRNAs in the samples, followed the same trend as total RNA and produced concentrations

near the limit of detection (Fig. S6). This is consistent with known half-lives of mRNA (59).

Although trends in total RNA degradation tend to follow those found using mRNA transcripts,

the latter is more sensitive as it targets a specific gene and can determine actual copy numbers

via amplification (3,16,20). It is also susceptible to genomic DNA contamination (37). Fu *et* al.

(20), and Weinbrecht *et* al. (19), both detected blood-specific transcripts from bloodstains

deposited for up to one year, while the total RNA concentration in this study fell below the

quantitative range after ~24 hours of deposition. Here, we suggest the faster degradation of

total RNA observed in our study allows for better discrimination at early timepoints than mRNA

studies.

Conclusion

This study demonstrated significant changes in the RINe values and a potentially

diagnostic inflection point in DV200 of RNA extracted from bloodstains deposited for up to 168

hours. The biological replicates included in this study produced similar results for the quality

Moving forward, in addition to an increase in sample size, longer time series at colder

temperatures and experiments using human blood, the effects of humidity and substrate

should be evaluated. These factors have been shown to affect the RNA degradation and drying

times of bloodstains, which could cause the clear drop in RNA metrics to occur at earlier or later

timepoints (16,25,60). Timewise changes in the rheological properties of blood, DNA and RNA

could also be explored and combined with previously developed TSD models (61,62). This type

of combinatorial research could bring the forensic community one step closer to developing a

sensitive and robust TSD model for bloodstains that could one day be applied to crime scene

samples.

Funding

This work was supported by two NSERC Undergraduate Student Research Awards (Canada) to

Colin I. Elliott, Natural Sciences and Engineering Research Council of Canada, Grant/Award

Number: RGPIN-2017-03934 to Aaron B.A. Shafer and RGPIN-2020-05816 to Theresa E.

Stotesbury and Canadian Foundation for Innovation-JELF, Grant/Award Number: #36905 to

Aaron B.A. Shafer.

References

1. Bremmer RH, Bruin KG De, Gemert MJC Van, Leeuwen TG Van, Aalders MCG. Forensic quest for age determination of bloodstains. Forensic Sci Int 2012;216(1–3):1–11. doi:10.1016/j.forsciint.2011.07.027.

2. Zadora G, Menzyk A. In the pursuit of the holy grail of forensic science – Spectroscopic studies on the estimation of time since deposition of bloodstains. Trends Anal Chem 2018;105(5):137-65. doi:10.1016/j.trac.2018.04.009.

3. Anderson SE, Hobbs GR, Bishop CP. Multivariate analysis for estimating the age of a bloodstain. J Forensic Sci 2011;56(1):186–93. doi:10.1111/j.1556-4029.2010.01551.x.

4. Steelberg R, Pietro DS. A preliminary assessment of the correlation of drying time and the peripheral rim thickness of perimeter bloodstains. J Forensic Res 2019;10(2):1-5.

5. Peschel O, Kunz SN, Rothschild MA, Mutzel E. Blood stain pattern analysis. Forensic Sci Med Pathol 2011;7:257–70. doi:10.1007/s12024-010-9198-1.

6. Lynch M. God's signature: DNA profiling, the new gold standard in forensic science. Endeavour 2003;27(2):93–7. doi:10.1016/S0160-9327(03)00068-1.

7. Butler JM. Genetics and genomics of core short tandem repeat loci used in human identity testing. J Forensic Sci 2006;51(2):253–65.doi:10.1111/j.1556-4029.2006.00046.x.

8. Hares DR. Selection and implementation of expanded CODIS core loci in the United States. Forensic Sci Int Genet 2015;17:33–4. doi:10.1016/j.fsigen.2015.03.006.

9. Blackman S, Dawnay N, Ball G, Stafford-Allen B, Tribble N, Rendell P, et al. Developmental validation of the ParaDNA® Intelligence System - A novel approach to DNA profiling. Forensic Sci Int Genet 2015;17:137–48. doi:10.1016/j.fsigen.2015.04.009.

10. de Castro TC, Taylor MC, Carr DJ, Athens J, Kieser JA. Storage life of whole porcine blood used for bloodstain pattern analysis. J Can Soc Forensic Sci 2016;49(1):26–37. doi:10.1080/00085030.2015.1108770.

11. Pal A, Gope A, Iannacchione G. Temperature and concentration dependence of human whole blood and protein drying droplets. Biomolecules 2021;11(2):1–18. doi:10.3390/biom11020231.

12. Pal A, Gope A, Obayemi JD, Iannacchione GS. Concentration-driven phase transition and self-assembly in drying droplets of diluting whole blood. Sci Rep 2020;10(1):1–12. doi:10.1038/s41598-020-76082-6.

13. Laan N, Smith F, Nicloux C, Brutin D. Morphology of drying blood pools. Forensic Sci Int 2018;267:104–9.

14. Hanson EK, Ballantyne J. A blue spectral shift of the hemoglobin soret band correlates with the age (time since deposition) of dried bloodstains. PLoS One 2010;5(9):1–11. doi:10.1371/journal.pone.0012830.

15.   Doty KC, McLaughlin G, Lednev IK. A Raman "spectroscopic clock" for bloodstain age
determination: the first week after deposition. Anal Bioanal Chem 2016;408(15):3993–4001.
doi:10.1007/s00216-016-9486-z.

16.   Cossette ML, Stotesbury T, Shafer ABA. Quantifying visible absorbance changes and DNA
degradation in aging bloodstains under extreme temperatures. Forensic Sci Int
2021;318:110627. doi:10.1016/j.forsciint.2020.110627.

17.   Heneghan N, Fu J, Pritchard J, Payton M, Allen RW. The effect of environmental conditions on
the rate of RNA degradation in dried blood stains. Forensic Sci Int Genet 2021;51:102456.
doi:10.1016/j.fsigen.2020.102456.

18.   Agudelo J, Huynh C, Halámek J. Forensic determination of blood sample age using a bioaffinity-
based assay. Analyst 2015;140(5):1411–5. doi:10.1039/c4an02269f.

19.   Lech K, Liu F, Ackermann K, Revell VL, Lao O, Skene DJ, et al. Evaluation of mRNA markers for
estimating blood deposition time: Towards alibi testing from human forensic stains with
rhythmic biomarkers. Forensic Sci Int Genet 2016;21:119–25. doi:10.1016/j.fsigen.2015.12.008.

20.   Weinbrecht KD, Fu J, Payton M, Allen R. Time-dependent loss of mRNA transcripts from forensic
stains. Res Reports Forensic Med Sci 2017;7:1–12. doi:10.2147/rrfms.s125782.

21.   Fu J, Allen RW. A method to estimate the age of bloodstains using quantitative PCR. Forensic Sci
Int Genet 2019;39:103–8. doi:10.1016/j.fsigen.2018.12.004.

22.   Strasser S, Zink A, Kada G, Hinterdorfer P, Peschel O, Heckl WM, et al. Age determination of
blood spots in forensic medicine by force spectroscopy. Forensic Sci Int 2007;170(1):8–14.
doi:10.1016/j.forsciint.2006.08.023.

23.   Thom CS, Dickson CF, Gell DA, Weiss MJ. Hemoglobin variants: biochemical properties and
clinical correlates. Cold Spring Harb Perspect Med 2013; 3:1-22.
doi:10.1101/cshperspect.a011858.

24.   Brutin D, Sobac B, Loquet B, Sampol J. Pattern formation in drying drops of blood. J Fluid Mech
2011;667:85–95. doi:10.1017/S0022112010005070.

25.   Riddel Jr JP, Aouizerat BE, Miaskowski C, Lillicrap DP. Theories of blood coagulation. J Pediatr
Oncol 2007;24(3):123–31. doi:10.1177/1043454206298693.

26.   Laber TL, Epstein BP. Substrate effects on the clotting time of human blood. J Can Soc Forensic
Sci 2001;34(4):209–14. doi:10.1080/00085030.2001.10757531.

27.   Hansen J, Lesnikova I. DNA and RNA analysis of blood and muscle from bodies with variable
postmortem intervals. Forensic Sci Med Pathol 2014;10(3):322–8. doi:10.1007/s12024-014-
9567-2.

28.   Dissing J, Søndervang A, Lund S. Exploring the limits for the survival of DNA in blood stains. J
Forensic Leg Med 2010;17(7):392–6. doi:10.1016/j.jflm.2010.08.001.

29.   Stotesbury T, Cossette ML, Newell-Bell T, Shafer ABA. An exploratory time since deposition
analysis of whole blood using metrics of DNA degradation and visible absorbance spectroscopy.
Pure Appl Geophys 2020;178:735-43. doi: 10.1007/s00024-020-02494-0.

30.   Wu J, Xiao J, Zhang Z, Wang X, Hu S, Yu J. Ribogenomics: The science and knowledge of RNA.
Genom Proteom Bioinf 2014;12(2):57–63. doi: 10.1016/j.gpb.2014.04.002.

31.   Nowakowski J, Tinoco I. RNA structure and stability. Semin Virol 1997;165:153–65.

32.   Natsidis P, Schiffer PH, Salvador-Martínez I, Telford MJ. Computational discovery of hidden
breaks in 28S ribosomal RNAs across eukaryotes and consequences for RNA Integrity Numbers.
Sci Rep 2019;9(1):1–10. doi:10.1038/s41598-019-55573-1.

33.   Setzer M, Juusola J, Ballantyne J. Recovery and stability of RNA in vaginal swabs and blood,
semen, and saliva stains. J Forensic Sci 2008;53(2):296–305. https://doi.org/10.1111/j.1556-
4029.2007.00652.x.

34.   Hanson E, Ingold S, Haas C, Ballantyne J. Messenger RNA biomarker signatures for forensic body
fluid identification revealed by targeted RNA sequencing. Forensic Sci Int Genet 2018;34:206–
21. doi:10.1016/j.fsigen.2018.02.020.

35.   Bauer M. RNA in forensic science. Forensic Sci Int Genet 2007;1(1):69–74.
doi:10.1016/j.fsigen.2006.11.002.

36.   Salzmann AP, Russo G, Kreutzer S, Haas C. Degradation of human mRNA transcripts over time as
an indicator of the time since deposition (TsD) in biological crime scene traces. Forensic Sci Int
Genet 2021;53:1-10. doi:10.1016/j.fsigen.2021.102524.

37.   Chan LY, Mugler CF, Heinrich S, Vallotton P. Non-invasive measurement of mRNA decay reveals
translation initiation as the major determinant of mRNA stability. eLife 2018;7:1–32. doi:
10.7554/eLife.32536.

38.   Hashemipetroudi SH, Nematzadeh G, Ahmadian G, Yamchi A, Kuhlmann M. Assessment of DNA
contamination in RNA samples based on ribosomal DNA. J Vis Exp 2018;131:1–11.
doi:10.3791/55451.

39.   Minshall N, Git A. Enzyme- and gene-specific biases in reverse transcription of RNA raise
concerns for evaluating gene expression. Sci Rep 2020;10(1):1–7. doi:10.1038/s41598-020-
65005-0.

40.   Acinas SG, Sarma-Rupavtarm R, Klepac-Ceraj V, Polz MF. PCR-induced sequence artifacts and
bias: Insights from comparison of two 16s rRNA clone libraries constructed from the same
sample. Appl Environ Microbiol 2005;71(12):8966–9. doi:10.1128/AEM.71.12.8966-8969.2005.

41.   Sidstedt M, Hedman J, Romsos EL, Waitara L, Wadsö L, Steffen CR, et al. Inhibition mechanisms
of hemoglobin, immunoglobulin G, and whole blood in digital and real-time PCR. Anal Bioanal
Chem 2 018;410(10):2569–83. doi:10.1007/s00216-018-0931-z.

42.   Orr A, Gualdieri R, Cossette ML, Shafer A, Stotesbury T. Whole bovine blood use in forensic
research: Sample preparation and storage considerations. Sci Justice 2021;61(3):214-20.
doi:10.1016/j.scijus.2021.02.004.

43.   Giraldo CE, Álvarez ME, Carmona JU. Effects of sodium citrate and acid citrate dextrose solutions
on cell counts and growth factor release from equine pure-platelet rich plasma and pure-
platelet rich gel. BMC Vet Res 2015;11(60):1–7. doi:10.1186/s12917-015-0370-4.

44.   Invitrogen Life Technologies. PureLink ™ total RNA blood purification kit: for isolating total RNA
from whole blood. United States, 2005.

45.   Agilent Technologies. Agilent high sensitivity RNA screentape system quickguide. Germany
2015:1-4.

46.   Agilent Technologies. Agilent 4200 TapeStation system: system manual, Germany 2019:1-212.

47.   Agilent Technologies. Comparison of RIN and RINe algorithms for the Agilent 2100 Bioanalyzer
and the Agilent 2200 TapeStation systems. United States 2012:1–4.

48.   Schroeder A, Mueller O, Stocker S, Salowsky R, Leiber M, Gassmann M, et al. The RIN: an RNA
integrity number for assigning integrity values to RNA measurements. BMC Mol Biol
2006;7(3):1–14. doi:10.1186/1471-2199-7-3.

49.   Wimmer I, Tröscher AR, Brunner F, Rubino SJ, Bien CG, Weiner HL, et al. Systematic evaluation
of RNA quality, microarray data reliability and pathway analysis in fresh, fresh frozen and
formalin-fixed paraffin-embedded tissue samples. Sci Rep 2018;8(1):1–17. doi:10.1038/s41598-
018-24781-6.

50.   Piovesan A, Caracausi M, Antonaros F, Pelleri MC, Vitale L. GeneBase 1.1: A tool to summarize
data from NCBI gene datasets and its application to an update of human gene statistics.
Database 2016;2016:1–13.doi:10.1093/database/baw153.

51.   Chomczynski P, Wilfinger WW, Eghbalnia HR, Kennedy A, Rymaszewski M, Mackey K. Inter-
individual differences in RNA levels in human peripheral blood. PLoS One 2016;11(2):1–15.
doi:10.1371/journal.pone.0148260.

52.   Zimin A V., Delcher AL, Florea L, Kelley DR, Schatz MC, Puiu D, et al. A whole-genome assembly
of the domestic cow, Bos taurus. Genome Biol 2009;10(4):1-10. doi:10.1186/gb-2009-10-4-r42.

53.   Hammerle-Fickinger A, Riedmaier I, Becker C, Meyer HHD, Pfaffl MW, Ulbrich SE. Validation of
extraction methods for total RNA and miRNA from bovine blood prior to quantitative gene
expression analyses. Biotechnol Lett 2010;32(1):35–44. doi:10.1007/s10529-009-0130-2.

54.   Roland L, Drillich M, Iwersen M. Hematology as a diagnostic tool in bovine medicine. J Vet
Diagnostic Investig 2014;26(5):592–8. doi:10.1177/1040638714546490.

55.   Huang L, Lin P, Tsai K, Wang L. The effects of storage temperature and duration of blood
samples on DNA and RNA qualities. PLoS One 2017;12(9):1–13. doi:10.1371/journal.
pone.0184692.

56. Matsubara T, Soh J, Morita M, Uwabo T, Tomida S, Fujiwara T, et al. DV200 index for assessing
RNA integrity in next-generation sequencing. BioMed Res Int 2020;2020:1-6.doi:
10.1155/2020/9349132.

57. Illumina. Evaluating RNA quality from FFPE Samples. Guidelines for obtaining high-quality RNA
sequencing results from degraded RNA with Illumina RNA enrichment assays, 2016:1–4.

58. Historical Data - Climate - Environment and Climate Change Canada.
https://climate.weather.gc.ca/historical_data/search_historic_data_stations_e.html?searchTyp
e=stnProv&timeframe=1&lstProvince=&optLimit=yearRange&StartYear=1990&EndYear=2021&
Year=2021&Month=3&Day=3&selRowPerPage=25 (accessed June 30, 2021).

59. Baudrimont A, Voegeli S, Viloria EC, Stritt F, Lenon M, Wada T, et al. Multiplexed gene control
reveals rapid mRNA turnover. Sci Adv 2017;3(7):1-12. doi:10.1126/sciadv.1700006.

60. Sirker M, Schneider PM, Gomes I. A 17-month time course study of human RNA and DNA
degradation in body fluids under dry and humid environmental conditions. Int J Legal Med
2016;130(6):1431–8. https://doi.org/10.1007/s00414-016-1373-9

61. Baieth HEA. Physical parameters of blood as a non-Newtonian fluid. Int J Biomed Sci
2008;4(4):323–9.

62. Boukany PE, Wang S-Q. Shear banding or not in entangled DNA solutions depending on the level
of entanglement. J Rheol 2009;53(1):73–83. doi:10.1122/1.3009299.