Extracellular adenosine induces hypersecretion of IL-17A by T-helper 17 cells through the

adenosine A2a receptor to promote neutrophilic inflammation

Mieko Tokano[1,2], Sho Matsushita[1,3], Rie Takagi[1], Toshimasa Yamamoto[2], and Masaaki

Kawano[1,*]

[1]Department of Allergy and Immunology, Faculty of Medicine, Saitama Medical University, 38

Morohongo, Moroyama, Saitama 350-0495, Japan

[2]Department of Neurology, Saitama Medical University, 38 Morohongo, Moroyama, Saitama,

350-0495, Japan

[3]Allergy Center, Saitama Medical University, 38 Morohongo, Moroyama, Saitama 350-0495,

Japan

[*]Correspondence: Department of Allergy and Immunology, Faculty of Medicine, Saitama

Medical University, 38 Morohongo, Moroyama, Saitama 350-0495, Japan

Tel.: +81 49 276 1173; Fax: +81 49 294 2274

E-mail address: mkawano@saitama-med.ac.jp

Running title: Adenosine modulates neutrophilic inflammation

Abstract

Extracellular adenosine, produced from ATP secreted by neuronal or immune cells, may play a role in endogenous regulation of inflammatory responses. However, the underlying molecular mechanisms are largely unknown. Here, we show that adenosine primes hypersecretion of interleukin (IL)-17A by CD4$^+$ T cells via T cell receptor activation. This hypersecretion was also induced by an adenosine A2a receptor (A2aR) agonist, PSB0777. In addition, an A2aR antagonist (Istradefylline), and inhibitors of adenyl cyclase, protein kinase A, and cAMP response element binding protein (which are signaling molecules downstream of the Gs protein coupled with the A2aR), suppressed IL-17A production, suggesting that activation of A2aR induces IL-17A production by CD4$^+$ T cells. Furthermore, immune subset studies revealed that adenosine induced hypersecretion of IL-17A by T-helper (Th)17 cells. These results indicate that adenosine is an endogenous modulator of neutrophilic inflammation. Administration of an A2aR antagonist to mice with experimental autoimmune encephalomyelitis led to marked amelioration of symptoms, suggesting that suppression of adenosine-mediated IL-17A production is an effective treatment for Th17-related autoimmune diseases.

Introduction

T-helper (Th)17 cells are a subset of T-helper cells induced by stimulation of naïve CD4$^+$ T cells with both tumor growth factor (TGF)-β and interleukin (IL)-6 in the presence of T cell receptor signaling. IL-17A production by Th17 cells increases neutrophilic inflammation (1-3); however, not all neutrophilic inflammatory diseases are explained by known Th17 responses (4). Indeed, it is likely that as-yet-unknown Th17 or neutrophilic inflammatory responses occur. Here, we show that adenosine induces IL-17A production by CD4$^+$ T cells directly. Extracellular adenosine is one of the first "signals" identified during regulation of a large number of physiological and pathological processes, including bulging of an artery (5), sleep promotion (6), and regulation of nerve action (7,8). Extracellular adenosine is produced from secreted ATP that undergoes rapid stepwise dephosphorylation by ectonucleotidases such as the E-NTPDase CD39, which converts ATP or ADP to ADP or AMP, respectively, and the 5'-nucleotidase CD73, which dephosphorylates AMP to adenosine (9); both CD39 and CD73 are expressed by activated CD4$^+$ T cells and antigen presenting cells (APCs) (10,11). Extracellular adenosine stimulates adenosine receptors (A1R, A2aR, A2bR, and A3R) belonging to a superfamily of membrane proteins called the G protein-coupled receptor family of class A seven-transmembrane domain receptors. A2aR and A2bR signal the Gs protein to trigger cAMP synthesis, which in turn activates adenyl cyclase (AC), protein kinase A (PKA), and cAMP response element binding protein (CREB). By contrast, A1R and A3R signal the Gi protein to trigger cAMP degradation. In addition, A2bR also signals the Gq protein, which in turn activates phospholipase C. In an immunological context, adenosine receptors are expressed by various immune cells, including T cells and APCs (12). It is also suggested that adenosine stimulates neutrophil chemotaxis and phagocytosis via

A1R and A3R (13). In addition, adenosine induces Th17 differentiation by activating A2bR on

CD4$^+$ T cells (14). By contrast, several reports suggest that adenosine suppresses Th17

differentiation via activation of A2aR on CD4$^+$ T cells (15-17). Considering that the G protein

downstream of A2aR is Gs, and those of A2bR are Gs and Gq (5), it is assumed that induction of

A2bR-mediated Th17 differentiation is induced through simultaneous activation of Gq and

suppression of Gs, or via other unknown mechanisms. Therefore, the precise molecular

mechanism(s) underlying the effect of adenosine on Th cells is unclear. Here, we show that

adenosine promotes IL-17A production in a two-way mixed lymphocyte reaction (MLR). In

addition, an A2aR agonist (PSB0777) induced IL-17A production, and an A2aR antagonist

(Istradefylline) inhibited production, induced by adenosine, suggesting that activation of A2aR

plays a role in adenosine-mediated IL-17A production. This notion was further supported by the

observation that inhibitors of AC, PKA, and CREB, which are signaling molecules downstream

of the Gs protein (18), also suppressed adenosine-mediated IL-17A production. Immune subset

studies suggested that Th17 cells play a role in adenosine-mediated hypersecretion of IL-17A.

Administration of an A2R antagonist to mice with experimental autoimmune encephalomyelitis

(EAE) (19,20) markedly ameliorated symptoms. Taken together, the data indicate that

adenosine-dependent hypersecretion of IL-17A by Th17 cells contributes not only to

antibacterial defense but also to neutrophilic autoimmune diseases, and that suppressing this

process may be an effective therapy for the latter.

Methods

Mice

BALB/c mice were obtained from Japan SLC, Inc. SJL/J mice were obtained from Charles River

Laboratories Japan, Inc. Mice were housed in appropriate animal care facilities at Saitama

Medical University and handled according to international guidelines for experiments with

animals. All experiments were approved by the Animal Research Committee of Saitama Medical

University.

Two-way MLR

Splenic lymphocytes were collected by lyzing tissue in a Dounce homogenizer, followed by

layering over Ficoll Paque (GE health care, Chicago, IL, USA), as described previously (21).

BALB/c splenic lymphocytes (3×10^6) were mixed with SJL/J splenic lymphocytes (3×10^6) in

2 mL of DMEM medium containing 10% FCS, 100 U/ml penicillin, 100 μg/mL streptomycin, 2

mM L-glutamine, 1 mM sodium pyruvate, and 50 μM 2-mercaptoethanol (D10 medium) in 12

well plates in the presence of adenosine (0–1 mM) (Sigma, St. Louis, MO, USA); in the presence

of each adenosine receptor agonist (0–10 μM) (A1R: 2-Chloro-N6-cyclopentyladenosine

(CCPA), Tocris, Bristol, UK; A2aR: PSB0777, Tocris; A2bR: BAY 60-6583, Tocris; A3R:

HEMADO, Tocris); in the presence of A2aR antagonist (0–1 nM) (Istradefylline, Sigma) plus

adenosine (100 μM); in the presence of an AC inhibitor (0–1 μM) (MDL-12330A, Enzo Life

Sciences, Farmingdale, NY, USA), a PKA inhibitor (0–1 μM) (H-89, Tocris) plus adenosine

(100 μM), or a CREB inhibitor (0.1–0.6 μM) (666-15, Cayman Chemical Company, Ann Arbor,

MI, USA) plus adenosine (100 μM) (to inhibit A2aR signaling); in the presence of an activator

of AC (0.06–0.6 μM) (Forskolin, Sigma); or in the presence of a CD39 inhibitor (0–1 μM)

(ARL67156, Tocris) or a CD73 inhibitor (0–1 μM) (adenosine 5'-(α, β-methylene) diphosphate

(AMP-CP; Tocris) plus ATP (100 μM) (GE healthcare). After mixing, the plates were incubated

for 7 days at 37°C. The supernatants were collected for use in cytokine ELISAs.

Isolation of CD4$^+$, CD4$^+$CD62L$^+$ T cells, and B cells

CD4$^+$ T cells within the BALB/c and SJL/J splenocyte populations were isolated from the

mixture prepared previously after rupturing red blood cells (22). Cells were isolated by positive

selection of CD4$^+$ T cells using magnetic-activated cell sorting (MACS) (Miltenyi Biotec,

Bergisch Gladbach, Germany), according to the manufacturer's instructions. CD4$^+$CD62L$^+$ T

cells were isolated from BALB/c splenocytes by a combination of negative and positive selection

by MACS. During positive selection of CD62L$^+$ T cells, negatively isolated CD4$^+$ T cells were

collected as the flow through fraction. B cells were isolated from BALB/c splenocytes by

positive selection using MACS. Cells were resuspended in 1 mL of D10 medium and counted.

The purity and viability of CD4$^+$ T cells, CD4$^+$CD62L$^+$ T cells, and B cells were >90% (Sup. Fig.

1). The purity and viability of cells in the flow through fraction collected during isolation of

CD4$^+$CD62L$^+$ T cells are shown in Supplementary Figure 1.

Cell sorting

BALB/c CD4$^+$ T cells (prepared as described above) were labeled for 30 min at 4°C with

PE-conjugated anti-mouse CCR3, CCR5, CCD6, CD25, or CD62L antibodies (BioLegend). The

cells were then washed and sorted using a FACS Aria II flow cytometer (BD Biosciences). The

purity and viability of the sorted cells are Supplementary Figure 1.

CD3/CD28 stimulation

CD4$^+$ T cells (1×10^6) were stimulated for 7 days at 37°C with anti-mouse CD3 (BioLegend; 1

μg/mL) and CD28 (BioLegend; 0.5 μg/mL) antibodies (CD3/CD28) in the presence of adenosine

(0–1 mM), each adenosine receptor agonist (0–10 μM; A1R: 2-Chloro-N6-cyclopentyladenosine

(CCPA), Tocris, Bristol, UK; A2aR: PSB0777, Tocris; A2bR: BAY 60-6583, Tocris; and A3R:

HEMADO, Tocris), an A2aR antagonist (Istradefylline; 0–1 nM) plus adenosine (600 μM), an

AC inhibitor (0–1 μM; MDL-12330A), a PKA inhibitor (0–1 μM) (H-89), a CREB inhibitor

(0.1–0.6 μM; 666-15) plus adenosine (600 μM), or an activator of AC (0.06–0.6 μM; Forskolin)

in 500 μL of D10 medium. After cell sorting, cells (3×10^5) were plated in 24 well plates and

stimulated for 7 days at 37°C with anti-mouse CD3 (1 μg/mL) and CD28 (0.5 μg/mL) antibodies

in the presence of adenosine (600 μM) in 100 μL of D10 medium. After stimulation, the

supernatants were collected for use in cytokine ELISAs.

Adenosine or ATP ELISAs

MLR was performed by mixing BALB/c lymphocytes (6×10^6) with SJL/J lymphocytes ($6 \times$

10^6) in a 15 mL tube for 0–24 h at 37°C in the presence of a CD39 inhibitor (ARL67156) (0–1

μM) and a CD73 inhibitor (AMP-CP) (0–1 μM) in 200 μL of D10 medium. CD3/CD28

stimulation was performed for 24 h at 37°C in a 15 mL tube by incubating BALB/c CD4$^+$ T cells

(1×10^7) with anti-mouse CD3 (1 μg/mL) and CD28 (0.5 μg/mL) antibodies plus ARL67156 (1

μM) or AMP-CP (1 μM) in 200 μL of D10 medium. Simulation with LPS or an anti-mouse

CD40 antibody (Ab) was performed for 24 h at 37°C in a 15 mL tube; briefly, BALB/c B cells (1

$\times 10^7$) or BALB/c bone marrow (BM)-derived dendritic cells (BM-DCs) (1×10^7), generated

from mouse bone marrow cells as described previously (23), were incubated with LPS (Sigma;

0.5 μg/mL) or anti-mouse CD40 Ab (5 μg/mL; BioLegend) plus ARL67156 (1 μM) or AMP-CP

(1 μM) in 200 μL of D10 medium. The purity and viability of BM-DCs were >90% (Sup. Fig. 1).

After incubation, the supernatants were collected and tested in adenosine or ATP ELISAs

(Biovision, Milpitas, CA, USA).

Differentiation of naïve CD4$^+$ T cells

Naïve CD4$^+$ T cells (3×10^5) in 500 μL of D10 medium in 24 well plates were stimulated for 7

days at 37°C with anti-mouse CD3 (1 μg/mL) and CD28 (0.5 mg/mL) antibodies plus mouse

IL-6 (20 ng/mL) (Peprotech, Rocky Hill, NJ), and human TGF-β1 (2 ng/mL) (Peprotech) in the

presence of an A2aR antagonist (Istradefylline) (0–1 nM). After incubation, the supernatants

were collected for use in cytokine ELISAs, and total RNA was extracted from the cells using a

Sepasol-RNA I Super G (Nacalai Tesque, Kyoto, Japan) for quantitative PCR. In another

experiment, cells in 500 μL of D10 medium were stimulated for another 7 days at 37°C with

anti-mouse CD3 (1 μg/mL) and CD28 (0.5 μg/mL) antibodies in the presence of Istradefylline

(0.1–1 nM) plus adenosine (600 μM). Supernatants were collected for use in cytokine ELISAs.

Quantitative PCR

cDNA was synthesized from total RNA prepared using Superscript IV VILO master mix

(Invitrogen, Waltham, MA, USA), according to the manufacturer's instructions. Quantitative

PCR (qPCR) was performed as described previously (24) using the primer pairs described below,

Power SYBR green PCR Master mix (Applied Biosystems, Waltham, MA, USA), and a

QuantStudio 12K Flex (Applied Biosystems) equipped with QuantStudio 12K Flex software

(Applied Biosystems). Then, the ΔCt was calculated. The following forward and reverse primers,

respectively, were used (all from Eurofins Scientific, Luxembourg): GAPDH,

5'-AGCTTGTCATCAACGGGAAG-3' and 5'-TTTGATGTTAGTGGGGTCTCG-3'; IL-17A,

5'-TGTGAAGGTCAACCTCAAAGTCT-3'   and

5'-GAGGGATATCTATCAGGGTCTTCAT-3'.

EAE model

EAE was induced as described previously (19). Briefly, SJL/J mice received a subcutaneous

inguinal injection (100 μg/mouse) of the proteolipid protein (PLP) peptide (PLP139-151, Tocris)

emulsified in complete Freund's adjuvant (CFA) containing mycobacterium tuberculosis H37Ra

(100 μg/mouse; Difco, Detroit, MI, USA). Mice also received oral water or an A2aR antagonist

(Istradefylline; 6 μg/mouse) once every 2 days from Day −7 to Day +18 after immunization with

PLP peptide (Day 0). Mice were examined daily for signs of EAE, which were graded as

described (25).

Histological analysis of spinal cord samples

On Day 18 post-induction, histological analysis was performed by excising the lumbar, thoracal,

and cervical parts of the spinal cord. Paraffin-embedded spinal cord sections were stained with

hematoxylin and eosin. T cells were labeled with an anti-mouse CD3ε Ab (Cell Signaling

technology, Danvers, MA, USA). Tissue sections were observed using the light microscopy

mode of an inverted fluorescence phase-contrast microscope (BZ-9000, Keyence, Osaka, Japan).

Peptide pulse assay

At 7 days post-subcutaneous immunization with PLP peptide emulsified in CFA, splenocytes (2

$\times 10^6$ in 200 μL of D10 medium) were seeded in 96-well round plates and pulsed for 3 days at

37°C with PLP peptide (10 μM) in the presence of adenosine (600 μM) and an A2aR antagonist

(Istradefylline; 0–1 nM). For the oral administration experiments using the A2a antagonist,

splenocytes (2×10^6 in 200 μL of D10 medium) or inguinal lymph node lymphocytes, prepared

as described previously (26) (1×10^6 in 200 μL of D10 medium) were seeded in 96-well round

plates and pulsed for 3 days at 37°C; this was done at 7 days post-subcutaneous immunization

with PLP peptide emulsified in CFA. Mice also received oral water, an A2aR agonist (PSB0777)

(6 μg/mouse), or an A2aR antagonist (Istradefylline; 6 μg/mouse) once every 2 days from Day –

7 to Day +7 after immunization with PLP peptide (Day 0). At 18 days post-subcutaneous

immunization with PLP peptide emulsified in CFA, spinal cord cells prepared as described

previously (26) (100 μL of resuspended in 500 μL of D10 medium/mouse) were seeded in

96-well round plates and pulsed for 3 days at 37°C with PLP peptide (10 μM). The supernatants

were collected for use in cytokine ELISAs.

Flow cytometry analysis

The MLR was performed for 7 days in the presence or absence of adenosine (100 μM) (as

described above). For the $CD4^+IL\text{-}17A^+$ study, cells were blocked with anti-mouse CD16/CD32

antibodies (BioLegend, San Diego, CA, USA) and stained for 30 min at 4°C with a fluorescein

isothiocyanate (FITC)-conjugated anti-mouse CD4 Ab (BioLegend). The cells were then fixed,

permeabilized, and stained with a phycoerythrin (PE)-conjugated anti-mouse IL-17A Ab

(BioLegend). For the Th subset study, after 0, 3, and 7 days of the MLR assay, cells were

blocked with anti-mouse CD16/CD32 antibodies (BioLegend) and then stained for 30 min at 4°C

with a FITC-conjugated anti-mouse CD4 Ab (BioLegend) and PE-conjugated anti-mouse CCR5

(a Th1 marker (27)), CCR3 (a Th2 marker (28)), or CCR6 (a Th17 marker (29,30)) antibodies

(BioLegend). For the MHC, costimulatory molecule, and A2aR expression studies in APCs,

BALB/c B cells (3×10^5) or BALB/c BM-DCs (3×10^5) seeded in the wells of a 96-well in 200

μL of D10 medium were incubated with LPS (0.5 μg/mL; Sigma) or an anti-mouse CD40 Ab (5

μg/mL) (BioLegend) for 1 day. Next, supernatants were collected for cytokine ELISAs, or

blocked with anti-mouse CD16/CD32 antibodies (BioLegend) and then stained for 30 min at 4°C

with FITC-conjugated anti-mouse CD80, CD86, or CD40 Abs and corresponding isotype

controls (BioLegend) (for the costimulatory study), with PE-conjugated anti-mouse H-2 or

I-A/I-E Ab and corresponding isotype controls (BioLegend) (for the MHC study), or fixed,

permeabilized, and stained with a PE-conjugated anti-A2aR Ab (Novus Biologicals, Minneapolis,

MN, USA) and corresponding isotype control (for the A2aR expression study). For the A2aR

expression study in CD4$^+$ T cells, CD4$^+$ T cells (1×10^6) were stimulated for 7 days at 37°C with

anti-mouse CD3 (1 μg/mL) and CD28 (0.5 μg/mL) antibodies in the presence of adenosine (0–1

mM). At 0–7 days post-stimulation, cells were blocked with anti-mouse CD16/CD32 antibodies

(BioLegend) and then stained for 30 min at 4°C with a FITC-conjugated anti-mouse CD4 Ab

(BioLegend) before fixation, permeabilization, and staining with a PE-conjugated anti-A2aR Ab

(Novus Biologicals). For EAE analysis at 18 days post-subcutaneous immunization with PLP

peptide emulsified in CFA, splenocytes (1×10^6), inguinal lymph node lymphocytes (1×10^6),

and spinal cord cells (100 μL of resuspended in 500 μL of D10 medium/mouse) were blocked

with anti-mouse CD16/CD32 antibodies (BioLegend) and then stained for 30 min at 4°C with a

FITC-conjugated anti-mouse CD4 Ab (BioLegend) and PE-conjugated anti-mouse CCR5 (a Th1

marker (27)), CCR3 (a Th2 marker (28)), or CCR6 (a Th17 marker (29,30)) antibodies

(BioLegend). Finally, the cells were washed and analyzed on a FACSCanto II flow cytometer

(BD Biosciences, Franklin Lakes, NJ) using FACSDiva acquisition software (BD Biosciences).

Cytokine ELISAs

The concentrations of IFN-γ, IL-5, IL-6, IL-17A, IL-17F, IL-22, IL-23, and TGF-β1, in cell

supernatants were measured using specific ELISA kits (DuoSet Kit, R&D, Minneapolis, MN,

USA). Any value below the lower limit of detection (15.6 pg/mL) was set to 0. No cytokine

cross-reactivity was observed within the detection ranges of the kits. If necessary, samples were

diluted appropriately so that the measurements fell within the appropriate detection range for

each cytokine.

Statistical analysis

Differences between two groups were analyzed using an unpaired Student's t-tests. Differences

between three or more groups were analyzed using one-way ANOVA with Tukey's post-hoc test.

Clinical scores were analyzed using a non-parametric Mann-Whitney *U*-test. All calculations

were performed using KaleidaGraph software (Synergy software, Reading, PA, USA). A P value

< 0.05 was considered statistically significant.

Results

Adenosine promotes IL-17A production by CD4$^+$ T cells in an MLR

First, we analyzed the effect of adenosine on CD4$^+$ T cells during T cell-APC interactions in an

MLR (31). We found that CD4$^+$ T cells exposed to adenosine secreted IL-17A in a

dose-dependent manner (Fig. 1A–C). Since both agonist-mediated IL-17A production and

antagonist-mediated suppression of adenosine-mediated IL-17A production were observed in the

presence of an adenosine A2aR agonist (PSB0777) and an antagonist (Istradefylline),

respectively (Fig. 1D and Sup. Fig. 2A), we hypothesized that IL-17A in the MLR was produced

by CD4$^+$ T cells stimulated via the A2aR. This notion was supported by the finding that AC,

PKA, and CREB inhibitors of signaling molecules downstream of the A2aR also suppressed

IL-17A production (Fig. 1E). Consistent with this notion, activation of A2aR downstream of AC

upregulated IL-17A production (Fig. 1E). In the MLR, the Th1 and Th17 cell populations

increased slightly but significantly (by approximately 1.7-fold) in the presence of 100 μM

adenosine at Day 7 (Sup. Fig. 2C). Considering that 100 μM adenosine increased IL-17A

secretion by approximately 10-fold compared with medium (Fig. 1B), it appears that

adenosine-mediated upregulation of IL-17A is the main contributor to upregulation of the

IL-17A production by CD4$^+$ T cells. Furthermore, ATP induced IL-17A production in the MLR,

which was suppressed by the A2aR antagonist and by inhibitors of CD39 and CD73 (Fig. 1F),

suggesting that adenosine plays a role in IL-17A production. Since the A2aR antagonist and

inhibitors of CD39 and CD73 also inhibited production of IL-17A in the MLR (Fig. 1G), we

postulate that *de novo* adenosine production is induced in the MLR.

Adenosine production in the MLR

CD39 and CD73 expressed on the surface of endothelial cells (32,33) and immune cells (10,11),

including T cells and DCs, are critical for production of adenosine from ATP. Since we found

that inhibitors of CD39 and CD73 inhibited basal IL-17A production in the MLR (Fig. 1G), we

next addressed the source of adenosine production in the MLR. As shown in Figure 2A,

production of adenosine and ATP was time-dependent. In accordance with suppression of

IL-17A production, inhibitors of CD39 and CD73 suppressed adenosine production at 24 h after

the start of the MLR (Fig. 2B and C). Since the MLR induces activation of CD4$^+$ T cells by

APCs (31), we also addressed adenosine production by activated CD4$^+$ T cells, B cells, and

BM-DCs (Fig. 2D–F). Production of both adenosine and ATP by activated CD4$^+$ T cells, B cells,

and BM-DCs was observed; inhibitors of CD39 and CD73 suppressed production by activated

CD4$^+$ T cells although production of both adenosine and ATP by CD4$^+$ T cells, B cells, and

BM-DCs was observed without LPS or anti CD40 Ab stimulation (Fig. 2D). This indicates the

possibility that adenosine stimulation is mediated by the CD4$^+$ T cell to other immune cell

contact and adenosine produced by CD4$^+$ T cells or other immune cells such as B cells may

induce adenosine-mediated IL-17A hypersecretion by CD4$^+$ T cells in the MLR.

Th17 cells hypersecrete IL-17A in the presence of anti-CD3/CD28 antibodies and adenosine

Next, we tried to identify the Th subset that generated IL-17A in the presence of adenosine. First,

we confirmed that CD4$^+$ T cells expressed the A2aR and secreted IL-17A by stimulating them

with agonistic anti-CD3/CD28 antibodies in the presence of adenosine (Fig. 3A). Time course

studies showed that adenosine-mediated IL-17A production was detected from 3 days

post-CD3/CD28 stimulation (Fig. 3B and Sup. Fig. 3). Administration of adenosine within 6 h of

Ab stimulation triggered IL-17A production; however, administration at 24 h post-stimulation

did not (Fig. 3C). We noticed that A2aR expression by CD4$^+$ T cells was upregulated

significantly (2–4.7 fold) by anti-mouse CD3 and CD28 Abs at 1–7 days after stimulation, and

that this upregulation was further (slightly but significantly) enhanced (by approximately

1.5-fold) by increasing amounts of adenosine (Sup. Fig. 4A). As in the MLR, CD4$^+$ T cells also

produced IL-17A after stimulation with anti-CD3/CD28 antibodies in the presence of an A2aR

agonist (Fig. 3D and Sup. Fig. 2B). Adenosine-mediated IL-17A production by

CD3/CD28-stimulated CD4$^+$ T cells was suppressed by an A2aR antagonist (Fig. 3D and Sup.

Fig. 2B). This was supported by data showing that inhibitors of signaling molecules downstream

of the A2R also suppressed IL-17A production, whereas activation of A2aR downstream of AC

upregulated the production (Fig. 3E). Production of other Th17-related cytokines was also

induced by adenosine through the A2aR (Fig. 4). This again suggests that activated CD4$^+$ T cells

produce IL-17A upon A2aR activation. Furthermore, we noticed that CD4$^+$CD62L$^-$, but not

CD4$^+$CD62L$^+$, cells produced IL-17A after CD3/CD28 stimulation in the presence of adenosine,

suggesting that adenosine induces IL-17A production by effector Th cells (Fig. 3F). Therefore,

we performed immune subset studies after separating CD4$^+$ T cells using anti-chemokine

receptor (CCR) antibodies. As shown in Figure 3G, CD4$^+$CCR6^{hi} T cells produced IL-17A upon

stimulation of CD3/CD28, and production was strongly upregulated in the presence of adenosine.

Since CCR6 is a typical marker of Th17 cells (29,30), this suggests that activated Th17 cells

hypersecrete IL-17A in the presence of adenosine.

To address the effect of adenosine on APCs, we analyzed production of IL-6, IL-23,

and, TGF-β1 by B cells and BM-DCs (Sup. Fig. 4B). Also, we analyzed the effect of adenosine

on expression of CD80/CD86/CD40, MHC class I/class II, and A2aR by B cells and BM-DCs

(Sup. Fig. 4C). Increasing amounts of adenosine suppressed IL-6 production by LPS-stimulated

B cells. and IL-6 and IL-23 production by LPS-stimulated BM-DCs. Expression of

CD80/CD86/CD40, MHC class I/class II, and A2aR by LPS or anti-CD40 Ab-stimulated-B cells

and -BM-DCs in the presence of 100 μM and 600 μM adenosine was 0.70- and 1.31-fold higher

than that by cells stimulated by LPS or anti-CD40 Ab alone. These data indicate that adenosine

mainly affects CD4$^+$ T cells to activate Th17 function.

An adenosine A2aR antagonist ameliorates IL-17A-related autoimmune EAE responses

The above results raise the possibility that adenosine-mediated hypersecretion of IL-17A by

Th17 cells contributes to Th17-related autoimmune diseases. This hypothesis is supported by a

report showing that CD73 knockout mice are resistant to EAE (34), a Th17-mediated

autoimmune disease (20). We expected, therefore, that A2aR antagonist-mediated suppression of

Th17 responses should improve EAE. To address this, we examined the efficacy of an A2aR

antagonist in EAE model SJL/J mice (19). EAE was induced by immunization of mice with an

I-A^s restricted helper peptide derived from a myelin PLP peptide comprising amino acids 139–

151 (HSLGKWLGHPDKF). The peptide was emulsified in CFA. First, we confirmed that the

A2aR antagonist suppressed adenosine-mediated IL-17A production by CD3/CD28-stimulated

CD4$^+$ T cells from SJL/J strain mice (Fig. 5A). The A2aR antagonist also significantly

suppressed adenosine-mediated IL-17A production after differentiation of Th17 cells from naïve

CD4$^+$ T cells (Fig. 5B, *right*). The A2aR antagonist did not suppress IL-17A expression or

production during differentiation of Th17 cells from naïve CD4$^+$ T cells (Fig. 5B, *left and center*).

By contrast, and in agreement with Figure 3G, adenosine administration did not induce IL-17A

production during and after differentiation of Th1, Th2, and Treg cells from naïve CD4$^+$ T cells

(data not shown). Next, we pulsed splenocytes with the PLP peptide after immunization to

confirm that IL-17A production was induced in a peptide-dependent manner, and that production

was upregulated by adenosine. As expected, IL-17A production occurred in a peptide-dependent

manner and was upregulated by adenosine (Fig. 5C). Furthermore, the A2aR antagonist

suppressed production of IL-17A, suggesting that the A2aR antagonist inhibits IL-17A

production by CD4$^+$ T cells induced by immunization with the PLP peptide. We next addressed

the effect of an oral A2aR antagonist on production of IL-17A by CD4$^+$ T cells induced by

immunization with the PLP peptide (Fig. 5D). Oral administration of PSB0777 or Istradefylline

did not significantly induce or suppress IL-17A production by PLP peptide-pulsed splenocytes

and inguinal lymph node lymphocytes. These results indicate that the A2aR antagonist may not

inhibit PLP peptide-specific Th17-generation but rather suppress adenosine-mediated IL-17A

hypersecretion by PLP peptide-specific Th17 cells. Since oral administration of PSB0777 did not

up-regulate IL-17A production, adenosine-mediated IL-17A production may be saturated by *de*

novo extracellular adenosine in the body. Hence, we expect that the A2aR antagonist may

suppress adenosine-mediated IL-17A hypersecretion by PLP peptide-specific Th17 cells during

migration to the inflammatory site, which may suppress EAE symptoms. Finally, the A2aR

antagonist was administered orally to mice before and during EAE induction (Fig. 5E and F). As

shown, the clinical scores of mice receiving the A2aR antagonist were markedly lower than

those of control mice (receiving water) at 18 days post-immunization with the PLP peptide (Fig.

5E). Accordingly, histological studies showed that the numbers of central nervous

system-infiltrating CD3$^+$ cells in mice receiving the A2aR antagonist were much lower than

those in mice receiving water (Fig. 5F). Consistent with these data, flow cytometry analysis

showed that the proportion of CD4$^+$ cells in total counts in the spinal cord were reduced

significantly by the oral A2aR antagonist; in particular, the percentage of Th17 cells in the spinal

cord was reduced significantly (Fig. 5G), suggesting the A2aR antagonist suppressed migration

of $CD4^+$ T cells to, and retained Th17 cells within, the spinal cord during EAE induction.

Accordingly, PLP peptide-dependent production of IL-17A was observed in spinal cord cells

after oral administration of water; production was suppressed significantly by the oral A2aR

antagonist (Fig. 5H). These results suggest that the A2aR antagonist suppresses hypersecretion of

IL-17A by PLP peptide-specific Th17-cells, resulting in impaired migration from the draining

lymph nodes to the inflammatory site, and subsequent retention at the inflammatory site. These

phenomena may suppress IL-17A-mediated EAE symptoms.

Discussion

Here, we showed that adenosine induces hypersecretion of IL-17A by Th17 cells. Addition of adenosine (1 mM) to a two-way MLR increased IL-17A production to > 25 times the basal level; however, both basal production and increased production of IL-17A were suppressed by an A2aR antagonist, and by CD39/CD73 inhibitors. This indicates that hypersecretion of IL-17A in the presence of adenosine occurs by other mechanisms in addition to T cell-APC interactions. Since endothelial cells and nervous system cells also express CD39/CD73 (32,33,35,36) and produce adenosine (8,37), and activated CD4$^+$ T cells in the present study hypersecreted IL-17A at 6 h post-CD3/CD28 stimulation, it is possible that activated Th17 cells also receive adenosine from endothelial and neuronal cells to induce hypersecretion of IL-17A.

We hypothesize that adenosine affects multiple immunological events as follows: After APCs take up antigen they move to the secondary lymphoid organs. Followed by an innate immune response in the secondary lymphoid organs, APCs induce Th differentiation from naïve CD4$^+$ T cells. In particular, APCs induce Th17 differentiation by secreting IL-6 and TGF-β1. Then, the transcription factor RORγt switches on the IL-17A gene by binding to the IL-17A promoter (38). During this response, adenosine rarely stimulates naïve CD4$^+$ T cells due to low levels of A2aR expression (Sup. Fig. 4A). Following an adaptive immune response, APCs in the secondary lymphoid organs stimulates effector Th17 cells and drives expression of the IL-17A gene by recruiting transcription factors such as NF-κB and NFAT downstream of the TCR signaling pathway (39). In this response, A2aR expression is also upregulated (Sup. Fig. 4A), thereby increasing and sensitivity of A2aR signaling in response to adenosine. After stimulation, Th17 cells migrate toward the inflammatory site. During migration, activated Th17 cells receive adenosine from immune cells, neural cells, endothelial cells, or in an autocrine manner. During

this process, Th17 cells further up-regulate expression of the IL-17A gene by recruiting

transcription factors such as CREB downstream of the A2aR-cAMP-PKA pathway (40). Then,

Th17 cells induce IL-17A hypersecretion at the inflammatory site. In humans, the transcription

factor cAMP-responsive element modulator α increases transcription of the human *Il-17* gene by

binding to the promoter region (41). The A2aR antagonist may suppress adenosine-mediated

IL-17A hypersecretion by adenosine-stimulated Th17 cells at the inflammatory site and

subsequently suppress neutrophil-mediated inflammatory responses and Th cell migration. This

hypothesis is consistent with a previous EAE study showing that CD73 expression and adenosine

receptor signaling are required for efficient entry of lymphocytes into the central nervous system

during EAE development (34); our own data suggest that CD73 expression and adenosine

receptor signaling may be required to induce IL-17A hypersecretion by Th17 cells around the

inflammatory site. γδ T cells, which are categorized mainly into two subpopulations (CD8αα^{+}

and CD8^{-} cells), express RORγt and are also a source of IL-17A (42,43). We hypothesize that

the A2aR antagonist will also suppress adenosine-mediated IL-17A production by

adenosine-stimulated γδ T cells through A2aR signaling (44).

During the sequence of immunological events described above, TCR signaling

up-regulates phosphodiesterase activity and suppresses cAMP signaling during the adaptive

responses in the secondary lymphoid organs (45). The cAMP–PKA signaling pathway plays a

major role in regulating immune responses, and cAMP is the most potent and acute inhibitor of T

cell activation (45). Our data indicate that up-regulation of A2aR expression is induced after

TCR stimulation (Sup. Fig. 4A); therefore, adenosine-mediated IL-17A hypersecretion is

induced after stimulation of TCR signaling as seen in Fig. 3C. Therefore, extracellular adenosine

may not block TCR signaling in the body. Also, a specific amount of Forskolin may not block

TCR signaling, leading to subsequent upregulation of IL-17A secretion by TCR-activated $CD4^+$

T cells (Fig. 1E and 3E). We hypothesize that extracellular adenosine-mediated IL-17A

hypersecretion is induced after activation of TCR signaling during migration from the secondary

lymphoid organs to the inflammatory site. In the MLR and the PLP peptide pulse experiment,

both activated Th17 cells and adenosine-stimulated Th17 cells might produce IL-17A during the

T-APC interaction. Thus, inhibitors of CD39/CD73 and an A2aR antagonist may suppress

adenosine-mediated hypersecretion by adenosine-stimulated activated Th17 cells.

It is suggested that physiological concentrations of adenosine are lower than 1 μM;

however, they can be increased by stimuli such as high K^+ levels, electrical stimulation,

glutamate receptor agonists, hypoxia, hypoglycemia, and ischemia (46). To obtain sufficient

adenosine (> 100 μM) to trigger hypersecretion of IL-17A, activated Th17 cells may need to

make contact with non-immune cells such as adenosine-producing endothelial cells (47) and

neuronal cells (48) to form a microenvironment with a high adenosine concentration (as

observed during T cell-APC interactions at immunological synapses) (49). Thus, A2aR

antagonists, rather than CD39/CD73 inhibitors, might be more effective at inhibiting *de novo*

adenosine-mediated hypersecretion of IL-17 by Th17 cells. A previous study suggests that

intracellular adenosine is transported out of cells by efficient equilibrative transporters (50);

CD39/CD73 inhibitors would not suppress this type of *de novo* adenosine production.

With regard to the effect of adenosine on other Th subsets, our observations were

different from those of previous reports (51,52); here, we observed that adenosine upregulated

IFN-γ (a Th1-related cytokine) secretion at 5 and 7 days and had no significant effect on IL-5 (a

Th2-related cytokine) secretion by $CD4^+$ T cells after CD3/CD28 stimulation with 600 μM of

adenosine, although IL-17A production was significant (Sup. Fig. 3). This suggests that

adenosine induces hypersecretion of IL-17A by Th17 cell but does not suppress Th1 and Th2

activity. However, previous studies report that adenosine-mediated suppression of IFN-γ and

IL-5 was observed 1 day after T cell receptor-mediated stimulation of CD4$^+$ T cells (51,52). This

may indicate that in the short term adenosine prioritizes stimulation of Th17 cell activity rather

than that of Th1 and Th2 cells, and that it does not suppress effector Th activity in the long term.

It is also suggested that the A2aR agonist, CGS21680, suppresses Th17 differentiation

(15-17). This result is opposite to ours; one reason for this may be differences in the source of

the A2aR agonist. The A2aR agonist CGS 21680 is much less selective than the A2aR agonist

that we used in this study (PSB0777); this is because CGS21680 binds not only to the A2aR but

also to A1R and A3R, which are associated with the Gi protein (which has opposite effects to the

Gs protein) (53,54). Therefore, it is probable that CGS21680 may cancel out any agonist effects

by activating A1R and A3R. Also, it is suggested that the A2aR antagonist, SCH58261,

up-regulates Th17 differentiation in mice (17). We hypothesized that up-regulation of Th17

differentiation by SCH58261 may be induced through relative downregulation of A2aR activity

compared with that of A2bR; this relative increase in A2bR activity induces Th17 differentiation

in mice (55,56). Also, it is probable that SCH58261 induces relative increase in the activity of G

protein-coupled receptors other than adenosine receptors (e.g., dopamine receptors) to induce

Th17 differentiation in mice (57). We also hypothesize that although SCH58261 may induce

Th17 differentiation *in vivo*, it may not stimulate Th17 activity; this is because our data show

that an A2a antagonist (Istradefylline) suppressed IL-17A secretion by differentiated Th17 cells

but did not suppress Th17 differentiation (Fig. 5B). This hypothesis is supported by previous

data showing that SCH58261 markedly suppresses symptoms of EAE, a typical Th17-mediated

disease (34).

Our data suggest that production of IL-17A is relatively higher after exposure to an

A2aR agonist, PSB0777, than after exposure to an A2bR agonist, BAY 60-6583. PSB0777 is a

potent adenosine A2aR agonist (Ki = 44.4 nM for rat brain striatal A2aR) (54), and BAY

60-6583 is a potent adenosine A2bR agonist (Ki = 100 nM for rat A2bR) (58). By assuming that

the Ki values of PSB0777 and BAY 60-6583 are comparable, we thought that production of

IL-17A mediated by activation of the A2aR might be higher than that mediated by activation of

the A2bR. This hypothesis is supported by the notion that the A2aR is a high affinity receptor

with activity in the low to mid-nanomolar range, whereas the A2bR has a much lower affinity for

adenosine (micromolar) (5); this suggests that adenosine activates the A2aR rather than the

A2bR.

A2aR antagonists have been developed for treatment of Parkinsonism (59) and

malignancies (60). In addition, inhibitors of CD39/CD73 have been developed as anti-tumor

drugs (61,62). Regarding the effects of adenosine on tumor immunity, a previous study suggests

that adenosine suppresses effector T cell function since tumor cells express both CD39 and

CD73 and secrete adenosine (63). By contrast, several reports suggest that IL-17A promotes

emergence of pro-tumorigenic neutrophil phenotypes (64,65). Neutrophils in mouse tumor

models promote tumor metastasis (66-68), and observations in cancer patients have linked

elevated neutrophil counts in blood with increased risk of metastasis (69). Therefore, it is

probable that tumor-produced adenosine induces IL-17A secretion by $CD4^+CCR6^{hi}$ T cells

followed by neutrophilic inflammation, which promotes tumor metastasis. Cancer vaccines may

need to be administered along with an A2aR antagonist to suppress hypersecretion of IL-17A by

tumor-specific Th17 cells induced by tumor-produced adenosine, and to inhibit neutrophilic

inflammation at the tumor site.

The results presented herein indicate that these drugs may also be effective treatments

for Th17-mediated diseases (4) such as psoriasis, neutrophilic bronchial asthma, severe atopic

dermatitis, and autoimmune diseases by suppressing hypersecretion of IL-17A by Th17 cells.

Moreover, these drugs might be effective treatments for diseases caused by neutrophilic

inflammation of unknown cause in the dermis; such diseases include Behcet uveitis (70) and

vasculitis of adenosine deaminase 2 deficiency (71). This is because endothelial cells express

CD39/CD73 and produce adenosine, which could induce hypersecretion of IL-17A by Th17 cells,

thereby contributing to inflammation.

References

1 Matsuzaki, G. and Umemura, M. 2007. Interleukin-17 as an effector molecule of innate
and acquired immunity against infections. *Microbiol Immunol* 51:1139.

2 Manel, N., Unutmaz, D., and Littman, D. R. 2008. The differentiation of human T(H)-17
cells requires transforming growth factor-beta and induction of the nuclear receptor
RORgammat. *Nat Immunol* 9:641.

3 Ivanov, II, Zhou, L., and Littman, D. R. 2007. Transcriptional regulation of Th17 cell
differentiation. *Semin Immunol* 19:409.

4 Tesmer, L. A., Lundy, S. K., Sarkar, S., and Fox, D. A. 2008. Th17 cells in human
disease. *Immunol Rev* 223:87.

5 Cronstein, B. N. and Sitkovsky, M. 2017. Adenosine and adenosine receptors in the
pathogenesis and treatment of rheumatic diseases. *Nat Rev Rheumatol* 13:41.

6 Jones, B. E. 2009. Glia, adenosine, and sleep. *Neuron* 61:156.

7 Boison, D., Chen, J. F., and Fredholm, B. B. 2010. Adenosine signaling and function in
glial cells. *Cell Death Differ* 17:1071.

8 Sebastiao, A. M. and Ribeiro, J. A. 2009. Tuning and fine-tuning of synapses with
adenosine. *Curr Neuropharmacol* 7:180.

9 Yegutkin, G. G. 2008. Nucleotide- and nucleoside-converting ectoenzymes: Important
modulators of purinergic signalling cascade. *Biochim Biophys Acta* 1783:673.

10 Whiteside, T. L., Mandapathil, M., and Schuler, P. 2011. The role of the adenosinergic
pathway in immunosuppression mediated by human regulatory T cells (Treg). *Curr Med*
Chem 18:5217.

11   Allard, B., Longhi, M. S., Robson, S. C., and Stagg, J. 2017. The ectonucleotidases CD39

and CD73: Novel checkpoint inhibitor targets. *Immunol Rev* 276:121.

12   Junger, W. G. 2011. Immune cell regulation by autocrine purinergic signalling. *Nat Rev*

Immunol 11:201.

13   Barletta, K. E., Ley, K., and Mehrad, B. 2012. Regulation of neutrophil function by

adenosine. *Arteriosclerosis, thrombosis, and vascular biology* 32:856.

14   Wilson, J. M., Kurtz, C. C., Black, S. G., Ross, W. G., Alam, M. S., Linden, J., and Ernst,

P. B. 2011. The A2B adenosine receptor promotes Th17 differentiation via stimulation of

dendritic cell IL-6. *Journal of immunology* 186:6746.

15   Alchera, E., Rolla, S., Imarisio, C., Bardina, V., Valente, G., Novelli, F., and Carini, R.

2017. Adenosine A2a receptor stimulation blocks development of nonalcoholic

steatohepatitis in mice by multilevel inhibition of signals that cause immunolipotoxicity.

Transl Res 182:75.

16   Wang, L., Wan, H., Tang, W., Ni, Y., Hou, X., Pan, L., Song, Y., and Shi, G. 2018.

Critical roles of adenosine A2A receptor in regulating the balance of Treg/Th17 cells in

allergic asthma. *Clin Respir J* 12:149.

17   Ansari, M. A., Nadeem, A., Attia, S. M., Bakheet, S. A., Raish, M., and Ahmad, S. F.

2017. Adenosine A2A receptor modulates neuroimmune function through

Th17/retinoid-related orphan receptor gamma t (RORgammat) signaling in a BTBR T(+)

Itpr3(tf)/J mouse model of autism. *Cell Signal* 36:14.

18   Hasko, G., Linden, J., Cronstein, B., and Pacher, P. 2008. Adenosine receptors:

therapeutic aspects for inflammatory and immune diseases. *Nat Rev Drug Discov* 7:759.

19   Nakano, K., Higashi, T., Hashimoto, K., Takagi, R., Tanaka, Y., and Matsushita, S. 2008.

Antagonizing dopamine D1-like receptor inhibits Th17 cell differentiation: preventive

and therapeutic effects on experimental autoimmune encephalomyelitis. *Biochem*

Biophys Res Commun 373:286.

20 Kleinewietfeld, M., Manzel, A., Titze, J., Kvakan, H., Yosef, N., Linker, R. A., Muller, D.

N., and Hafler, D. A. 2013. Sodium chloride drives autoimmune disease by the induction

of pathogenic TH17 cells. *Nature* 496:518.

21 Higashi, T., Shimojo, N., Suzuki, S., Nakaya, M., Takagi, R., Hashimoto, K., Nakagome,

K., Nakamura, K., Kohno, Y., and Matsushita, S. 2011. Coenzyme A contained in

mothers' milk is associated with the potential to induce atopic dermatitis. *Int Immunol*

23:741.

22 Kawano, M., Takagi, R., Saika, K., Matsui, M., and Matsushita, S. 2018. Dopamine

regulates cytokine secretion during innate and adaptive immune responses. *Int Immunol*

30:591.

23 Lutz, M. B., Kukutsch, N., Ogilvie, A. L., Rossner, S., Koch, F., Romani, N., and Schuler,

G. 1999. An advanced culture method for generating large quantities of highly pure

dendritic cells from mouse bone marrow. *Journal of immunological methods* 223:77.

24 Joseph, J., Bittner, S., Kaiser, F. M., Wiendl, H., and Kissler, S. 2012. IL-17 silencing

does not protect nonobese diabetic mice from autoimmune diabetes. *Journal of*

immunology 188:216.

25 Podojil, J. R., Padval, M. V., and Miller, S. D. 2011. Combination treatment of mice with

CRx-153 (nortriptyline and desloratadine) decreases the severity of experimental

autoimmune encephalomyelitis. *Cell Immunol* 270:237.

26 Barthelmes, J., Tafferner, N., Kurz, J., de Bruin, N., Parnham, M. J., Geisslinger, G., and

Schiffmann, S. 2016. Induction of Experimental Autoimmune Encephalomyelitis in Mice and Evaluation of the Disease-dependent Distribution of Immune Cells in Various Tissues. *J Vis Exp*.

Takeuchi, A., Usui, Y., Takeuchi, M., Hattori, T., Kezuka, T., Suzuki, J., Okunuki, Y., Iwasaki, T., Haino, M., Matsushima, K., and Usui, M. 2005. CCR5-deficient mice develop experimental autoimmune uveoretinitis in the context of a deviant effector response. *Invest Ophthalmol Vis Sci* 46:3753.

Ma, W., Bryce, P. J., Humbles, A. A., Laouini, D., Yalcindag, A., Alenius, H., Friend, D. S., Oettgen, H. C., Gerard, C., and Geha, R. S. 2002. CCR3 is essential for skin eosinophilia and airway hyperresponsiveness in a murine model of allergic skin inflammation. *J Clin Invest* 109:621.

Acosta-Rodriguez, E. V., Rivino, L., Geginat, J., Jarrossay, D., Gattorno, M., Lanzavecchia, A., Sallusto, F., and Napolitani, G. 2007. Surface phenotype and antigenic specificity of human interleukin 17-producing T helper memory cells. *Nat Immunol* 8:639.

Annunziato, F., Cosmi, L., Santarlasci, V., Maggi, L., Liotta, F., Mazzinghi, B., Parente, E., Fili, L., Ferri, S., Frosali, F., Giudici, F., Romagnani, P., Parronchi, P., Tonelli, F., Maggi, E., and Romagnani, S. 2007. Phenotypic and functional features of human Th17 cells. *J Exp Med* 204:1849.

McDevitt, H. O. 2000. Discovering the role of the major histocompatibility complex in the immune response. *Annu Rev Immunol* 18:1.

Kaczmarek, E., Koziak, K., Sevigny, J., Siegel, J. B., Anrather, J., Beaudoin, A. R., Bach, F. H., and Robson, S. C. 1996. Identification and characterization of CD39/vascular ATP

597   diphosphohydrolase. *J Biol Chem* 271:33116.

598   33   Jalkanen, S. and Salmi, M. 2008. VAP-1 and CD73, endothelial cell surface enzymes in

599   leukocyte extravasation. *Arteriosclerosis, thrombosis, and vascular biology* 28:18.

600   34   Mills, J. H., Thompson, L. F., Mueller, C., Waickman, A. T., Jalkanen, S., Niemela, J.,

601   Airas, L., and Bynoe, M. S. 2008. CD73 is required for efficient entry of lymphocytes

602   into the central nervous system during experimental autoimmune encephalomyelitis.

603   *Proceedings of the National Academy of Sciences of the United States of America*

604   105:9325.

605   35   Braun, N., Sevigny, J., Robson, S. C., Enjyoji, K., Guckelberger, O., Hammer, K., Di

606   Virgilio, F., and Zimmermann, H. 2000. Assignment of ecto-nucleoside triphosphate

607   diphosphohydrolase-1/cd39 expression to microglia and vasculature of the brain. *Eur J*

608   *Neurosci* 12:4357.

609   36   Kulesskaya, N., Voikar, V., Peltola, M., Yegutkin, G. G., Salmi, M., Jalkanen, S., and

610   Rauvala, H. 2013. CD73 is a major regulator of adenosinergic signalling in mouse brain.

611   *PLoS One* 8:e66896.

612   37   Gunther, G. R. and Herring, M. B. 1991. Inhibition of neutrophil superoxide production

613   by adenosine released from vascular endothelial cells. *Ann Vasc Surg* 5:325.

614   38   Ivanov, II, McKenzie, B. S., Zhou, L., Tadokoro, C. E., Lepelley, A., Lafaille, J. J., Cua,

615   D. J., and Littman, D. R. 2006. The orphan nuclear receptor RORgammat directs the

616   differentiation program of proinflammatory IL-17+ T helper cells. *Cell* 126:1121.

617   39   Gaud, G., Lesourne, R., and Love, P. E. 2018. Regulatory mechanisms in T cell receptor

618   signalling. *Nat Rev Immunol* 18:485.

40   Hernandez, J. B., Chang, C., LeBlanc, M., Grimm, D., Le Lay, J., Kaestner, K. H., Zheng,

Y., and Montminy, M. 2015. The CREB/CRTC2 pathway modulates autoimmune disease

by promoting Th17 differentiation. *Nat Commun* 6:7216.

41 Rauen, T., Hedrich, C. M., Juang, Y. T., Tenbrock, K., and Tsokos, G. C. 2011.

cAMP-responsive element modulator (CREM)α protein induces interleukin 17A

expression and mediates epigenetic alterations at the interleukin-17A gene locus in

patients with systemic lupus erythematosus. *J Biol Chem* 286:43437.

42 Khairallah, C., Chu, T. H., and Sheridan, B. S. 2018. Tissue Adaptations of Memory and

Tissue-Resident Gamma Delta T Cells. *Frontiers in immunology* 9:2636.

43 Kadivar, M., Petersson, J., Svensson, L., and Marsal, J. 2016. CD8αβ+ γδ T Cells: A

Novel T Cell Subset with a Potential Role in Inflammatory Bowel Disease. *Journal of*

immunology 197:4584.

44 Liang, D., Shao, H., Born, W. K., O'Brien, R. L., Kaplan, H. J., and Sun, D. 2018. High

level expression of A2ARs is required for the enhancing function, but not for the

inhibiting function, of γδ T cells in the autoimmune responses of EAU. *PLoS One*

13:e0199601.

45 Bjørgo, E., Moltu, K., and Taskén, K. 2011. Phosphodiesterases as targets for modulating

T-cell responses. *Handb Exp Pharmacol*:345.

46 Latini, S. and Pedata, F. 2001. Adenosine in the central nervous system: release

mechanisms and extracellular concentrations. *J Neurochem* 79:463.

47 Carman, C. V. and Martinelli, R. 2015. T Lymphocyte-Endothelial Interactions:

Emerging Understanding of Trafficking and Antigen-Specific Immunity. *Frontiers in*

immunology 6:603.

48 Siffrin, V., Radbruch, H., Glumm, R., Niesner, R., Paterka, M., Herz, J., Leuenberger, T.,

Lehmann, S. M., Luenstedt, S., Rinnenthal, J. L., Laube, G., Luche, H., Lehnardt, S.,

Fehling, H. J., Griesbeck, O., and Zipp, F. 2010. In vivo imaging of partially reversible

th17 cell-induced neuronal dysfunction in the course of encephalomyelitis. *Immunity*

33:424.

49   Monks, C. R., Kupfer, H., Tamir, I., Barlow, A., and Kupfer, A. 1997. Selective

modulation of protein kinase C-theta during T-cell activation. *Nature* 385:83.

50   Fredholm, B. B. 2007. Adenosine, an endogenous distress signal, modulates tissue

damage and repair. *Cell Death Differ* 14:1315.

51   Csoka, B., Himer, L., Selmeczy, Z., Vizi, E. S., Pacher, P., Ledent, C., Deitch, E. A.,

Spolarics, Z., Nemeth, Z. H., and Hasko, G. 2008. Adenosine A2A receptor activation

inhibits T helper 1 and T helper 2 cell development and effector function. *FASEB J*

22:3491.

52   Lappas, C. M., Rieger, J. M., and Linden, J. 2005. A2A adenosine receptor induction

inhibits IFN-gamma production in murine CD4+ T cells. *Journal of immunology*

174:1073.

53   Klotz, K. N. 2000. Adenosine receptors and their ligands. *Naunyn Schmiedebergs Arch*

Pharmacol 362:382.

54   El-Tayeb, A., Michael, S., Abdelrahman, A., Behrenswerth, A., Gollos, S., Nieber, K.,

and Muller, C. E. 2011. Development of Polar Adenosine A2A Receptor Agonists for

Inflammatory Bowel Disease: Synergism with A2B Antagonists. *ACS Med Chem Lett*

2:890.

55   Kenakin, T. 2001. Inverse, protean, and ligand-selective agonism: matters of receptor

conformation. *FASEB J* 15:598.

56 Prather, P. L. 2004. Inverse agonists: tools to reveal ligand-specific conformations of G

protein-coupled receptors. *Sci STKE* 2004:pe1.

57 Nakano, K., Yamaoka, K., Hanami, K., Saito, K., Sasaguri, Y., Yanagihara, N., Tanaka,

S., Katsuki, I., Matsushita, S., and Tanaka, Y. 2011. Dopamine induces IL-6-dependent

IL-17 production via D1-like receptor on CD4 naive T cells and D1-like receptor

antagonist SCH-23390 inhibits cartilage destruction in a human rheumatoid

arthritis/SCID mouse chimera model. *Journal of immunology* 186:3745.

58 Alnouri, M. W., Jepards, S., Casari, A., Schiedel, A. C., Hinz, S., and Muller, C. E. 2015.

Selectivity is species-dependent: Characterization of standard agonists and antagonists at

human, rat, and mouse adenosine receptors. *Purinergic Signal* 11:389.

59 Kanda, T., Jackson, M. J., Smith, L. A., Pearce, R. K., Nakamura, J., Kase, H., Kuwana,

Y., and Jenner, P. 1998. Adenosine A2A antagonist: a novel antiparkinsonian agent that

does not provoke dyskinesia in parkinsonian monkeys. *Ann Neurol* 43:507.

60 Ohta, A., Gorelik, E., Prasad, S. J., Ronchese, F., Lukashev, D., Wong, M. K., Huang, X.,

Caldwell, S., Liu, K., Smith, P., Chen, J. F., Jackson, E. K., Apasov, S., Abrams, S., and

Sitkovsky, M. 2006. A2A adenosine receptor protects tumors from antitumor T cells.

Proceedings of the National Academy of Sciences of the United States of America

103:13132.

61 Kobie, J. J., Shah, P. R., Yang, L., Rebhahn, J. A., Fowell, D. J., and Mosmann, T. R.

2006. T regulatory and primed uncommitted CD4 T cells express CD73, which

suppresses effector CD4 T cells by converting 5'-adenosine monophosphate to adenosine.

Journal of immunology 177:6780.

62 Deaglio, S., Dwyer, K. M., Gao, W., Friedman, D., Usheva, A., Erat, A., Chen, J. F.,

Enjyoji, K., Linden, J., Oukka, M., Kuchroo, V. K., Strom, T. B., and Robson, S. C. 2007.

Adenosine generation catalyzed by CD39 and CD73 expressed on regulatory T cells

mediates immune suppression. *J Exp Med* 204:1257.

63   Moesta, A. K., Li, X. Y., and Smyth, M. J. 2020. Targeting CD39 in cancer. *Nat Rev*

Immunol.

64   Tuting, T. and de Visser, K. E. 2016. CANCER. How neutrophils promote metastasis.

Science 352:145.

65   Coffelt, S. B., Kersten, K., Doornebal, C. W., Weiden, J., Vrijland, K., Hau, C. S.,

Verstegen, N. J. M., Ciampricotti, M., Hawinkels, L., Jonkers, J., and de Visser, K. E.

2015. IL-17-producing gammadelta T cells and neutrophils conspire to promote breast

cancer metastasis. *Nature* 522:345.

66   Bald, T., Quast, T., Landsberg, J., Rogava, M., Glodde, N., Lopez-Ramos, D., Kohlmeyer,

J., Riesenberg, S., van den Boorn-Konijnenberg, D., Homig-Holzel, C., Reuten, R.,

Schadow, B., Weighardt, H., Wenzel, D., Helfrich, I., Schadendorf, D., Bloch, W.,

Bianchi, M. E., Lugassy, C., Barnhill, R. L., Koch, M., Fleischmann, B. K., Forster, I.,

Kastenmuller, W., Kolanus, W., Holzel, M., Gaffal, E., and Tuting, T. 2014.

Ultraviolet-radiation-induced inflammation promotes angiotropism and metastasis in

melanoma. *Nature* 507:109.

67   El Rayes, T., Catena, R., Lee, S., Stawowczyk, M., Joshi, N., Fischbach, C., Powell, C.

A., Dannenberg, A. J., Altorki, N. K., Gao, D., and Mittal, V. 2015. Lung inflammation

promotes metastasis through neutrophil protease-mediated degradation of Tsp-1.

Proceedings of the National Academy of Sciences of the United States of America

112:16000.

68   Wculek, S. K. and Malanchi, I. 2015. Neutrophils support lung colonization of

metastasis-initiating breast cancer cells. *Nature* 528:413.

69   Templeton, A. J., McNamara, M. G., Seruga, B., Vera-Badillo, F. E., Aneja, P., Ocana,

A., Leibowitz-Amit, R., Sonpavde, G., Knox, J. J., Tran, B., Tannock, I. F., and Amir, E.

2014. Prognostic role of neutrophil-to-lymphocyte ratio in solid tumors: a systematic

review and meta-analysis. *J Natl Cancer Inst* 106:dju124.

70   Yazici, H., Seyahi, E., Hatemi, G., and Yazici, Y. 2018. Behcet syndrome: a

contemporary view. *Nat Rev Rheumatol* 14:107.

71   Meyts, I. and Aksentijevich, I. 2018. Deficiency of Adenosine Deaminase 2 (DADA2):

Updates on the Phenotype, Genetics, Pathogenesis, and Treatment. *J Clin Immunol*

38:569.

Abbreviations: Th, T-helper; TGF, tumor growth factor; IL, interleukin; APCs, antigen

presenting cells; AC, adenyl cyclase; PKA, protein kinase A; CREB, cAMP response element

binding protein; MLR, two-way mixed lymphocyte reaction; EAE, experimental autoimmune

encephalomyelitis; CCPA, 2-Chloro-N6-cyclopentyladenosine; AMP-CP, adenosine 5'-(α,

β-methylene) diphosphate; BM, bone marrow; BM-DC, bone-marrow-derived dendritic cell;

CD3/CD28, agonistic anti-CD3/CD28 antibodies; PLP, myelin proteolipid protein; PLP peptide,

I-A^s restricted helper peptide derived from the PLP; CFA, complete Freund's adjuvant; CCR,

chemokine receptor; FITC, fluorescein isothiocyanate; PE, phycoerythrin; MACS,

magnetic-activated cell sorting; Ab, antibody; n, number of repeat experiments; SD, standard

deviation.

Acknowledgments

This work was supported by a Grant-in-Aid for Scientific Research (C) (no. 19K07201),

awarded to M.K., a Grant-in-Aid for Young Scientists (B) (no. 18K15327) to R.T., and a

Grant-in-Aid for Scientific Research (C) (no. 19K08887) awarded to S.M. by the Japanese

Society for the Promotion of Science. This work was also supported by the 44th and 45th

Science Research Promotion Fund, awarded to M.K. by the Promotion and Mutual Aid

Corporation for Private Schools of Japan.

Author contributions

M.T., R.T., S.M., and M.K., performed the experiments. M.T., S.M., and M.K., conceived and

designed the experiments. M.T., S.M., T.Y., and M.K., wrote the manuscript. All authors

discussed the results and commented on the manuscript.

Conflicts of interest

750   Sho Matsushita is an employee of iMmno, Inc.

751   The other authors have no conflicts of interest to declare.

 Figure legends

 Fig. 1. Adenosine induces hypersecretion of IL-17A by CD4$^+$ T cells in an MLR. A–C, An

 MLR was performed for 7 days in the presence of adenosine (0–1 mM). After 7 days, cells were

 stained with anti-CD4 (x-axis) and IL-17A (y-axis) antibodies (Abs), followed by flow

 cytometry analysis (**A**, n (number of repeat experiments) = 4). The percentage of

 IL-17A-producing CD4$^+$ T cells within the total CD4$^+$ T cell population is shown (**B**, n = 6–9).

 Cells supernatants were analyzed in an IL-17A ELISA (**C**, n = 6–9). **D**, The effects of the A2aR

 on IL-17A production in the presence of PSB0777 (an A2aR agonist) (*left*, n = 6–9),

 Istradefylline (an A2aR antagonist) plus adenosine (100 μM) (*right*, n = 6–9). **E**, The effects of

 A2aR signaling on IL-17A production in the presence of MDL-12330A (an adenyl cyclase

 inhibitor) plus adenosine (100 μM) (*first panel*, n = 4–6), H-89 (a protein kinase A inhibitor)

 plus adenosine (100 μM) (*second panel*, n = 4–6), 666-15 (a cAMP response element binding

 protein inhibitor) plus adenosine (100 μM) (*third panel*, n = 4–6), and Forskolin (an activator of

 adenyl cyclase) (*fourth panel*, n = 4–6) were analyzed in an IL-17A ELISA. **F**, The effects of

 CD39/CD73 inhibitors on IL-17A production in the presence of ARL67156 (a CD39 inhibitor)

 plus ATP (100 μM) (*left*, n = 4 – 6) or AMP-CP (a CD73 inhibitor) plus ATP (100 μM) (*right*, n

 = 4–6) were analyzed in an IL-17A ELISA. **G**, The effects of an A2aR antagonist and

 CD39/CD73 inhibitors on basal IL-17A production in the presence of Istradefylline (*left*, n = 4–

 7), ARL67156 (*center*, n = 4–7), or AMP-CP (*right*, n = 4–7) were analyzed in an IL-17A

 ELISA. Data are expressed as the mean ± standard deviation (SD) and were compared using an

 unpaired Student's t-test (**B**) or one-way ANOVA with Tukey's post-hoc test (**C–G**). *P < 0.05

 and **P < 0.01, compared with medium (**C**, **D**, *left*; **E**, *fourth panel*; and **G**), adenosine (100 μM)

(**D**, *right*, **E**, *first–third panels*), or ATP (100 μM) (**F**).

Fig. 2. Adenosine production in the MLR. A, Concentrations of adenosine or ATP in MLR

supernatants were measured in an adenosine (*left*, n = 4–6) or ATP (*right*, n = 4–6) ELISA (0–24

h). **B** and **C**, The effects of CD39 (**B**, ARL67156, n = 4–6) and CD73 (**C**, AMP-CP, n = 4–6)

inhibitors on production of adenosine or ATP in the MLR were analyzed in an adenosine (*left*) or

ATP (*right*) ELISA. **D, E** and **F**, Levels of adenosine and ATP in the supernatants of

CD3/CD28-stimulated CD4$^+$ T cells (**D**, n = 4), LPS- or anti-mouse CD40 Ab-stimulated B cells

(**E**, n = 4), and LPS- or anti-mouse CD40 Ab-stimulated BM-DCs (**F**, n = 4) were analyzed in

adenosine and ATP ELISAs at 24 h post-stimulation in the presence of CD39 or CD73 inhibitors.

Data are expressed as the mean ± SD and were compared using one-way ANOVA with Tukey's

post-hoc test. *$P < 0.05$ and **$P < 0.01$, compared with medium (**B** and **C**), CD3/CD28

stimulation (**D**), or LPS or anti-mouse CD40 Ab stimulation (**E** and **F**).

Fig. 3. Adenosine induces hypersecretion of IL-17A by Th17 cells. A and **B**, CD4$^+$ T cells

were stimulated for 1–7 days with anti-CD3/CD28 antibodies in the presence of adenosine (0–1

mM). After stimulation, the supernatants were analyzed in an IL-17A ELISA (n = 4–6). **C**,

Adenosine (600 μM) was added at 0–3 days after CD3/CD28 stimulation. At 7 days

post-CD3/CD28 stimulation, supernatants were analyzed in an IL-17A ELISA (n = 4). **D**, Effects

of the A2aR on IL-17A production in the presence of PSB0777 (an A2aR agonist) (*left*, n = 4–6),

Istradefylline (an A2aR antagonist) plus adenosine (600 μM) (*right*, n = 4–6). **E**, Effects of

A2aR signaling on IL-17A production in the presence of MDL-12330A (an adenyl cyclase

inhibitor) plus adenosine (600 μM) (*first panel*, n = 4–6), or H-89 (a protein kinase A inhibitor)

plus adenosine (600 μM) (*second panel*, n = 4–6), 666-15 (a cAMP response element binding

protein inhibitor) plus adenosine (600 μM) (*third panel*, n = 4–6), or Forskolin (an activator of

adenyl cyclase) (*fourth panel*, n = 4–6) were analyzed in an IL-17A ELISA. **F**, CD4⁺CD62L⁺ and

CD4⁺CD62L⁺FT cells were stimulated for 7 days with anti-CD3/CD28 antibodies in the

presence of adenosine (0–1 mM). After 7 days, supernatants were analyzed in an IL-17A ELISA

(n = 4). **G**, Subsets of CD4⁺ T cells were stimulated for 7 days by anti-CD3/CD28 antibodies in

the presence of adenosine (600 μM) after isolation of each CCR cell type (high (hi) and low (lo)

expression) (n = 4). Data are expressed as the mean ± SD and were compared using an unpaired

Student's t-test (**G**) or one-way ANOVA with Tukey's post-hoc test (**A–F**). *P < 0.05 and **P <

0.01, compared with CD3/CD28 stimulation (**A–C**, **D**, *left;* **E**, *fourth panel;* and **F**) or CD3/28

stimulation plus adenosine (600 μM) (**D**, *right* and **E**, *first–third panels*).

Fig. 4. Effect of adenosine and an A2aR antagonist on production of Th17-related cytokines.

A, An MLR was performed for 7 days in the presence of adenosine (0–1 mM) (*top row*),

PSB0777 (an A2aR agonist) (*second row*), or Istradefylline (an A2aR antagonist) plus adenosine

(100 μM) (*bottom row*). At 7 days post-incubation, the supernatants were analyzed in an

IL-17A (*left*), IL-17F (*center*), or IL-22 (*right*) ELISA (n = 6–9). **B,** CD4⁺ T cells were

stimulated with an anti-CD3/CD28 antibody for 7 days in the presence of adenosine (600 μM)

(*top row*), PSB0777 (an A2aR agonist) (*second row*), or Istradefylline (an A2aR antagonist) plus

adenosine (600 μM) *(bottom row)*. After stimulation, supernatants were analyzed in IL-17A (*left*),

IL-17F (*center*), and IL-22 (*right*) ELISAs (n = 4–6). Data are expressed as the mean ± SD and

were compared using one-way ANOVA with Tukey's post-hoc test. *P < 0.05 and **P < 0.01,

compared with medium or CD3/CD28 stimulation (*top and second row of panels*) or adenosine

(100 μM) or CD3/CD28 stimulation plus adenosine (600 μM) (*third row of panels*).

Fig. 5. Suppression of adenosine-mediated hypersecretion of IL-17A ameliorates EAE. A

and **B**, SJL/J CD4$^+$ T cells were stimulated for 7 days by anti-CD3/CD28 antibodies and an

A2aR antagonist (**A**, n = 4). Naïve CD4$^+$ T cells were stimulated for 7 days by anti-CD3/CD28

antibodies in the presence of IL-6, TGF-β1, and Istradefylline (an A2aR antagonist) (0–1 nM) (**B**,

left, n = 4). Alternatively, naïve CD4$^+$ T cells were stimulated for 7 days by anti-CD3/CD28

antibodies in the presence of IL-6 and TGF-β1, followed by another 7 day incubation with

anti-CD3/CD28 antibodies and Istradefylline (0–1 nM) (**B**, *right*, n = 4). After the stimulation,

supernatants were analyzed in an IL-17A ELISA. **C and D,** Mice were immunized

subcutaneously with PLP peptide emulsified in CFA (PLP peptide/CFA). Mice received oral

water, an A2aR agonist (PSB0777; 6 μg/mouse), or an A2aR antagonist (Istradefylline; 6

μg/mouse) once every 2 days from Day −7 to Day +7 after immunization with PLP peptide/CFA

(Day 0) (**D**). At 7 days post-immunization, splenocytes were incubated for 3 days with PLP

peptide in the presence of Istradefylline and adenosine (600 μM) (**C**, n = 4) or in the presence or

absence of PLP peptide (**D**, n = 4). After the incubation, supernatants were analyzed in an

IL-17A ELISA. **E,** To induce EAE, SJL/J mice were immunized with PLP peptide/CFA. Before

and post-immunization with the PLP peptide (Days 0 to 18), mice received oral Istradefylline (6

μg/mouse) or water once every 2 days. Clinical scores were recorded every day during EAE

induction (**E**, n = 15). Data were obtained from five independent experiments (n = 3 mice/group).

F, At 18 days post-immunization, spinal cord sections from mice administered oral Istradefylline

(*right*) or water (*left*) were stained with hematoxylin and eosin (*upper panels*) or with an

anti-mouse CD3 antibody (*lower panels*) (Scale bar, 100 μm) (**F**, n = 5). **G and H,** At 18 days

post-immunization, splenocytes (spleen, *left*), inguinal lymph node lymphocytes (inguinal lymph

node, *center*), and spinal cord cells (spinal cord, *right*) from mice administered oral water (*left* of

each panel) or Istradefylline (*right* of each panel) were stained with anti-CD4 (x-axis) and CCR5

(a Th1 marker; top panels, y-axis), CCR3 (a Th2 marker; second panels, y-axis), or CCR6 (a

Th17 marker; bottom panels, y-axis) antibodies (Abs), followed by flow cytometry analysis (**G**,

n = 5). Values shown in the upper and lower right of each panel denote the percentage of

CCR-positive counts within the total CD4-positive counts, and the percentage of CD4-positive

counts within the total count, respectively. *$P < 0.05$. Also, spinal cord cells were incubated for 3

days in the presence or absence of PLP peptide (**H**, n = 4). After the incubation, supernatants

were analyzed in an IL-17A ELISA. Data are expressed as the mean ± SD. Data were compared

using one-way ANOVA with Tukey's post-hoc test (**A**–**D, G, and H**), or using a non-parametric

Mann-Whitney *U*-test (**E**). *$P < 0.05$ and **$P < 0.01$, compared with CD3/CD28 stimulation

plus adenosine (600 μM) (**A**), CD3/CD28 stimulation plus IL-6 and TGF-β1 (**B**), PLP peptide

pulse (**C**), oral water (**G**), or PLP peptide pulse (oral water) (**H**).

Legends for the supplementary figures

Sup. Fig. 1. Purity and viability of each immune cell subset. A, *First row (left)*: Splenocytes

or isolated CD4$^+$ T cells were stained with a FITC-conjugated anti-mouse CD4 antibody (Ab)

(BioLegend) (x-axis) and a PE-conjugated anti-mouse CD3 Ab (BioLegend) (y-axis). *Second*

row (left): Splenocytes or CD4$^+$CD62L$^+$ T cells isolated by MACS were stained with a

FITC-conjugated anti-mouse CD4 Ab (BioLegend) (x-axis) and a PE-conjugated anti-mouse

CD62L Ab (BioLegend) (y-axis). *First row (right)*: Splenocytes and B cells were stained with a

FITC-conjugated anti-mouse CD19 Ab (BioLegend) (x-axis) and cell numbers were counted

(y-axis). *Second row (right)*: Splenocytes, BM leukocytes, and BM-DCs were stained with a

FITC-conjugated anti-mouse CD11c Ab (BioLegend) (x-axis) and cell numbers were counted

(y-axis). The number in each panel represents the percentage ± SD of each immune subset within

the total cell population (n = 3). **B**, *First row*: Isolated CD4$^+$ (by MACS)-, CD4$^+$CD62L$^-$ (by cell

sorting)-, or CD4$^+$CD62L$^+$ (by cell sorting) T cells were analyzed with a PE-conjugated

anti-mouse CD62L Ab (BioLegend) (x-axis) and cell numbers were counted (y-axis). The

number in each panel represents the percentage of each immune subset (- or +) within the total

cell population (- plus +). Data are representative of at least three repeat experiments. *Second*

row: Isolated CD4$^+$ (by MACS), CD4$^+$CCR3$^{low\ (lo)}$ (by cell sorting), or CD4$^+$CCR3$^{high\ (hi)}$ (by cell

sorting) T cells were analyzed with a PE-conjugated anti-mouse CCR3 Ab (BioLegend) (x-axis)

and cell numbers were counted (y-axis). *Third row*: Isolated CD4$^+$ (by MACS), CD4$^+$CCR5^{lo} (by

cell sorting), or CD4$^+$CCR5^{hi} (by cell sorting) T cells were analyzed with a PE-conjugated

anti-mouse CCR5 Ab (BioLegend) (x-axis) and cell numbers were counted (y-axis). *Fourth row*:

Isolated CD4$^+$ (by MACS), CD4$^+$CCR6^{lo} (by cell sorting), or CD4$^+$CCR6^{hi} (by cell sorting) T

cells were analyzed with a PE-conjugated anti-mouse CCR6 Ab (BioLegend) (x-axis) and cell

numbers were counted (y-axis). *Fifth row*: Isolated CD4$^+$ (by MACS), CD4$^+$CD25^{lo} (by cell

sorting), or CD4$^+$CD25^{hi} (by cell sorting) T cells were analyzed with a PE-conjugated anti-mouse

CD25 Ab (BioLegend) (x-axis) and cell numbers were counted (y-axis). The number in each

panel represents the mean percentage ± SD of each immune subset (hi or lo) within the total cell

population (hi plus lo) (n = 3). **C**, After cell isolation, each immune cell type was mixed with

Trypan blue. Viability was calculated as the number of unstained cells/(stained cells + unstained

cells) × 100 (n = 3). The percentage represents the mean percentage ± SD.

Sup. Fig. 2. Effect of an adenosine receptor agonist and antagonist on production of IL-17A

in an MLR and by CD3/CD28-stimulated CD4$^+$ T cells, and the effect of adenosine on

proliferation of Th cells. A, An MLR was performed for 7 days in the presence of each

adenosine receptor agonist, or each adenosine receptor antagonist plus adenosine (100 μM) (n =

6–9). **B**, CD4$^+$ T cells were stimulated with an anti-CD3/CD28 antibody for 7 days in the

presence of each adenosine receptor agonist, or in the presence of each adenosine receptor

antagonist plus adenosine (600 μM) (n = 6–9). After 7 days, the supernatants were analyzed in

an IL-17A ELISA. *First row*: CCPA (an A1R agonist). *Second row*: PSB0777 (an A2aR agonist,

left) and Istradefylline (an A2aR antagonist, *right*). *Third row*: BAY 60-653 (an A2bR agonist).

Forth row: HEMADO (an A3R agonist). **C**, An MLR was performed in the presence or absence

of adenosine (100 μM). After 0, 3, and 7 days, splenocytes were stained with anti-CD4 (x-axis)

and CCR5 (a Th1 marker; top panels, y-axis), CCR3 (a Th2 marker; second panels, y-axis), or

CCR6 (a Th17 marker; bottom panels, y-axis) antibodies, followed by flow cytometry analysis

(n = 3). Values in the upper right of each panel denote the percentage of CCR-positive counts

within the CD4 positive counts. *P < 0.05 and **P < 0.01. Data are expressed as the mean ± SD

and are compared using one-way ANOVA with Tukey's post-hoc test. *P < 0.05 and **P < 0.01,

compared with medium (**A**, *left*; **B**, *left*; and **C**), adenosine (100 μM) (**A**, *right*), CD3/28

stimulation plus adenosine (600 μM) (**B**, *right*).

Sup. Fig. 3. Effect of adenosine on production of IFN-γ, IL-5, and IL-17A by

CD3/CD28-stimulated CD4$^+$ T cells. CD4$^+$ T cells were stimulated for 1–7 days with

anti-CD3/CD28 antibodies in the presence or absence of adenosine (600 μM). After stimulation,

supernatants were analyzed in IFN-γ (*top row*), IL-5 (*second row*), and IL-17A (*bottom row*)

ELISAs (n = 4–6). Data are expressed as the mean ± SD and were compared using one-way

ANOVA with Tukey's post-hoc test. *P < 0.05 and **P < 0.01, compared with CD3/CD28

stimulation.

Sup. Fig. 4. Effect of adenosine on A2aR expression by CD4$^+$ T cells, and on expression of

MHC, costimulatory molecules, and A2aR and production of IL-6, IL-23, and TGF-β by

APCs. A, CD4$^+$ T cells were stimulated for 0–7 days with anti-CD3/CD28 antibodies in the

presence of adenosine (0–1 mM). After stimulation, cells were stained with a FITC-conjugated

anti-mouse CD4 antibody (Ab) (BioLegend) (x-axis) and a PE-conjugated anti-A2aR Ab (Novus

Biologicals) (y-axis). *First row*: Splenocytes (*left*) or isolated CD4$^+$ T cells (*right*) were stained

with anti-CD4 (x-axis) and A2aR (y-axis) Abs. *Second row*: CD3/CD28-stimulated CD4$^+$ T cells

were stained with anti-CD4 (x-axis) and A2aR (y-axis) Abs**.** *Third row*: CD4$^+$ T cells were

stimulated with CD3/CD28 Abs in the presence of adenosine (0–1 mM) and then stained with

anti-CD4 (x-axis) and A2aR (y-axis) Abs. The number in each panel represents the percentage ±

SD of A2aR^{high} cells within the CD4^{+} population (n = 3). Data were compared using one-way

ANOVA with Tukey's post-hoc test. *P < 0.05 and **P < 0.01, compared with CD3/CD28

stimulation (second and third rows). **B**, B cells and BM-DCs were stimulated for 1 day with LPS

(upper panels) or anti-mouse CD40 Ab (CD40) (lower panels) in the presence of adenosine (0.1–

1 mM). After stimulation, the IL-6, IL-23, and TGF-β1 concentrations in the supernatants were

analyzed by ELISA (n = 3). Data are expressed as the mean ± SD and were compared using

one-way ANOVA with Tukey's post-hoc test. *P < 0.05 and **P < 0.01, compared with LPS or

anti-mouse CD40 Ab stimulation. **C**, B cells (upper panels) and BM-DCs (lower panels) were

stimulated for 1 day with LPS or anti-mouse CD40 Ab (CD40) in the presence of adenosine (100

or 600 μM). After stimulation, the cells were stained with a FITC-conjugated anti-mouse CD80

(BioLegend), CD86 (BioLegend), or CD40 (BioLegend) Ab, PE-conjugated anti-mouse H-2

(BioLegend) or I-A/I-E (BioLegend) Ab, or a PE-conjugated anti-A2aR (Novus Biologicals) Ab

(x-axis), and cell numbers were counted (y-axis) (n = 3). The number in each panel represents

the mean fluorescence intensity ± SD (n = 3). Data were compared using one-way ANOVA with

Tukey's post-hoc test. *P < 0.05 and **P < 0.01, compared with LPS or anti-mouse CD40 Ab

stimulation.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

MLR

CD3/CD28-stimulated CD4⁺ T cells

Fig. 5

A

C

Cell type	Viability
Splenocytes	100 ± 0
CD4+ T cells	100 ± 0
B cells	100 ± 0
BM-DCs	94.5 ± 1.67
Naive CD4+ T cells	100 ± 0
Flow through of CD4+CD62L+ T cells	100 ± 0

Cell type	Viability
CD4+CD62Llo CD4+CD62Lhi	95.6 ± 2.90 98.3 ± 2.50
CD4+CCR3lo CD4+CCR3hi	97.1 ± 1.91 91.1 ± 6.37
CD4+CCR5lo CD4+CCR5hi	98.8 ± 0.89 91.1 ± 6.40
CD4+CCR6lo CD4+CCR6hi	98.8 ± 0.80 94.0 ± 4.24
CD4+CD25lo CD4+CD25hi	97.1 ± 1.77 98.4 ± 1.62

Sup. Fig. 2

Sup. Fig. 3

Sup. Fig. 4