Enhancing the dark side: Asymmetric gain of cone photoreceptors underpins discrimination of visual scenes based on their skewness.

Matthew Yedutenko[1], Marcus H.C. Howlett[1], and Maarten Kamermans[1, 2]

[1]Retinal Signal Processing Lab, Netherlands Institute for Neuroscience, Amsterdam, Netherlands
[2]Department of Biomedical Physics and Biomedical Optics, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, Netherlands

Abbreviated Tittle: Cone responses to skewness

Corresponding Author: Maarten Kamermans, m.kamermans@nin.knaw.nl

10 Figures

1 Table

Funding

This work was supported by a ZonMW grant 91215062 (M.K.), a grant from Horizon 2020: "Switchboard" (M.K.), a grant of ODAS (M.K.) and a grant of the Foundation of Friends of the NIN (M.K.).

Abstract

Psychophysical data indicates humans can discriminate visual scenes based on their skewness – the ratio of dark and bright patches within a visual scene. It was also shown that on a phenomenological level this skew discrimination is described by the so-called Blackshot mechanism, which accentuates strong negative contrasts within a scene. Here we demonstrate that the underlying computation starts as early as the cone phototransduction cascade whose gain is higher for strong negative contrasts than for strong positive contrasts. We recorded from goldfish cone photoreceptors and found that the asymmetry in the phototransduction gain leads to higher amplitude of the responses to negatively than to positively skewed light stimuli. This asymmetry in the amplitude was present in the photocurrent, voltage response and cone synaptic output. These results highlight the importance of the early photoreceptor non-linearity for perception. Additionally, we found that stimulus skewness leads to a subtle change in photoreceptor kinetics. For negatively skewed stimuli, the cone's impulse response functions peak later than for positively skewed stimulus. However, stimulus skewness does not affect the cone's overall integration time.

Introduction

Psychophysical studies show that humans are sensitive to the ratio of negative (intensity lower than the mean) and positive (intensity higher than the mean) patches of contrast in visual scenes (Chubb *et al.*, 1994, 2004; Graham *et al.*, 2016). This ratio is described by the parameter known as skewness. Visual stimuli are called positively skewed if there is a predominance of negative contrasts with some infrequent patches of high positive contrast and are called negatively skewed when the situation is reversed. Figure 1 illustrates one's ability to discriminate visual scenes based on skewness by mimicking an experiment performed by Chubb et al. (1994, 2004). The textures were randomly drawn from two distributions equal in every aspect but

skewness (Bonin *et al.*, 2006). Yet, one can appreciate the clear difference between negatively

skewed images (right upper) and positively skewed images (remaining panels). Chubb et al.

(1994, 2004) showed that on a phenomenological level the sensitivity to skewness can be

described by the so-called Blackshot mechanism. This Blackshot mechanism does not react to

skewness per se, rather its sensitivity to strong negative contrasts is simply much higher than

to strong positive contrasts and so it effectively reports the fraction of strong negative contrasts

within the scene.

What are the neuronal correlates of the Blackshot mechanism? Studies using salamander retinal

ganglion cells (RGC) (Tkačik *et al.*, 2014) and cat lateral geniculate nucleus neurons (LGN)

(Bonin *et al.*, 2006) did not report any response differences associated with changes in stimulus

skewness. Therefore, both studies concluded that the discrimination between skewed stimuli

occurs in the visual cortex. On the other hand, it is well-established that the retinal

photoreceptor's gain is asymmetric: given an equal input magnitude, the response amplitude to

a strong (>0.4 Weber unit) negative contrast step is greater than it is to a strong positive contrast

step (Laughlin, 1981; van Hateren, 2005; Endeman & Kamermans, 2010; Baden *et al.*, 2013;

Angueyra *et al.*, 2021). Furthermore, this responses asymmetry is observed throughout the

Figure 1

The discrimination of skewed stimuli. An example of textures used by Chubb et al.(1994, 2004) to psychophysically probe the ability of humans to discriminate visual scenes based on their skewness. The textures were randomly drawn from the probability distributions adjacent to each panel following the approach described by Bonin et al. (2006) and differed only in terms of skewness. The upper right texture is negatively skewed (-0.4), the remainder are positively skewed (+0.4).

post-receptor retinal stages (Lee *et al.*, 2003; Zaghloul *et al.*, 2003), the LGN (Kremkow *et al.*,

2014), and the primary visual cortex (Zemon *et al.*, 1988; Jin *et al.*, 2008; Yeh *et al.*, 2009;

Kremkow *et al.*, 2014). The asymmetrical processing of positive and negative contrasts should

lead to different response amplitudes to negatively and positively skewed stimuli and thus

might underpin the discrimination of skewed stimuli.

A possible reason why the differences in responses to skewed stimuli were not found in RGC

(Tkačik *et al.*, 2014) and LGN (Bonin *et al.*, 2006) studies was the power spectra of the stimuli

used. In both cases, the researchers employed band-limited white noise, where large

proportions of the signal power are outside the photoreceptor frequency bandwidth. Thus, in

both studies temporal filtering discarded a significant portion of the signal, reducing the

skewness and amplitude of the "effective" light stimuli actually available to photoreceptors.

Consequently, Bonin et al. (2006) and Tkacik et al. (2014) may not have found significant

differences in the processing of skewed stimuli because a) their stimuli hardly differed in terms

of "effective" skewness and b) the "effective" amplitudes of the employed stimuli were too

low to drive cone photoreceptors outside their linear response range.

Here we address whether photoreceptors process positive and negative skewed stimuli

differently. To account for the kinetics of cone photoreceptors, we stimulated goldfish cones

with sets of skewed stimuli with bandwidth similar to those of the goldfish cones. We found

the asymmetry in goldfish photoreceptor gain does indeed entail skew-dependent changes in

the cone response, leading to greater response amplitudes to negatively than to positively

skewed stimuli. This asymmetry originates in the cone phototransduction cascade and is the

first step in implementing the Blackshot mechanism. Additionally, we found that the cone's

impulse response function peaks ≈ 3.6 ms later for negatively skewed stimuli whereas the

cone's integration time is unaffected by stimulus skewness.

Thus, we show that in contrast to what models of various retinal, thalamic and cortical neurons

(Carandini *et al.*, 2005; Schreyer & Gollisch, 2021) often imply, cone photoreceptors do not

simply linearly relay visual stimuli to downstream circuitry. Rather, they also emphasize

specific features of their stimuli. These results highlight the important of accounting for the

full non-linear character of photoreceptor signal processing when studying the visual system.

Materials and Methods

Recording procedures

All animal experiments were conducted under the responsibility of the ethical committee of

the Royal Netherlands Academy of Arts and Sciences (KNAW), acting in accordance with the

European Communities Council Directive of 22 July 2003 (2003/65/CE) under license number

AVD-801002016517, issued by the Central Comity Animal Experiments of the Netherlands.

In all experiments retinas of adult goldfish (*Carassius auratus*) were used.

Goldfish were first dark-adapted for 5-10 minutes, sacrificed and the eyes enucleated. Retinas

were isolated under dim red illumination then placed photoreceptor side up in a recording

chamber (300 μl, model RC-26G, Warner Instruments) mounted on a Nikon Eclipse 600FN

microscope. The preparation was viewed on an LCD monitor by means of a 60× water-

immersion objective (N.A. 1.0, Nikon), a CCD camera, and infrared ($\lambda > 800$ nm; Kodak

wratten filter 87c, United States) differential interference contrast optics. Tissue was

continuously superfused with oxygenated Ringer's solution at room temperature (20˚C). The

composition of Ringer's solution was (in mM): 102.0 NaCl, 2.6 KCl, 1.0 $MgCl_2$, 1.0 $CaCl_2$,

28.0 $NaHCO_3$, 5.0 glucose continuously gassed with 2.5% CO_2 and 97.5% O_2 to yield a pH of

7.8 (osmolarity 245–255 mOsm). For calcium current (I_{Ca}) measurements, 5 mM of NaCl was

substituted with 5mM of CsCl and 100 μM of niflumic acid added.

Measurements from goldfish cones were performed in current- (voltage response), and voltage-

(photocurrent), clamp configurations. Patch pipettes (resistance 7-8 MOhm, PG-150T-10;

Harvard Apparatus, Holliston, Massachusetts) were pulled with a Brown Flaming Puller

(Model P-87; Sutter Instruments Company). Patch pipette solution contained (in mM): 96 K-

gluconate, 10 KCl, 1 MgCl2, 0.1 CaCl2, 5 EGTA, 5 HEPES, 5 ATP-K2, 1 GTP-Na3, 0.1

cGMP-Na, 20 phosphocreatine-Na2, and 50 units ml^{-1} creatine phosphokinase, adjusted with

KOH to pH 7.27–7.3 (osmolarity 265–275 mOsm). The chloride equilibrium potential (E_{Cl})

was -55mV except when the calcium current (I_{Ca}) was studied. Here E_{Cl} was set at -41 mV by

changing the concentrations of K-gluconate and KCl to 87 and 19 mM, respectively. All

chemicals were supplied by Sigma-Aldrich (Zwijndrecht, the Netherlands), except for NaCl

(Merck Millipore, Amsterdam, the Netherlands).

Filled patch pipettes were mounted on a MP-85 Huxley/Wall-type manual micromanipulator

(Sutter Instrument Company) and connected to a HEKA EPC-10 Dual Patch Clamp amplifier

(HEKA Elektronik GmbH, Lambrecht, Germany). After obtaining a whole cell configuration,

cones were first spectrally classified then stimulated with a set of skewed stimuli of

corresponding chromaticity. Data were recorded at a sample rate of 1 kHz using Patchmaster

software package (HEKA Elektronik GmbH). In total, we recorded 14 cones in voltage-clamp

mode (8 light responses, 6 measurements of I_{Ca}) and 16 cones in current-clamp mode (all light

responses). The liquid junction potential (approximately -15 to -16 mV) was not compensated.

Light stimuli

The light stimulator was a custom-built LED stimulator with a three-wavelength high-intensity

LED (Atlas, Lamina Ceramics, Westhampton, New Jersey, US). The peak wavelengths were

465, 525 and 624 nm. Bandwidth was smaller than 25 nm. Linearity was ensured by an optical

feedback loop. The output of the LED stimulator was coupled to the microscope via an optic

fiber and focused on cone outer segments though a 60× water-immersion objective. The mean

light intensity of all stimuli was $1.2*10^4$ photons/µm^2/s, which is in the photopic level for

goldfish (Malchow & Yazulla, 1986). Stimuli were presented at 1 kHz.

Skew Stimulus Set#1. Skewed stimuli were based on the natural time series of chromatic intensities (NTSCI) from the Van Hateren library (Van Hateren *et al.*, 2002). Given that psychophysical studies have shown visual scene discrimination occurs when intensities within the spatial domain are skewed (Figure 1), one may wonder how appropriate it is to study the phenomena with time series of intensities. However, we argue such an approach is correct for the following two reasons. Firstly, saccadic eye movements convert spatial stimuli into time series. Hence, the photoreceptors of subjects in psychophysical studies 'perceive' the presented skewed textures as time series of intensities. Secondly, a photoreceptor's direct light response only depends on the light intensities falling upon its outer segment and not on the stimulus's spatial structure. Hence, one can reproduce the response of an array of cones to the presentation of a single spatial texture by recording the response of a single cone to a collection of time series of intensities that correspond to the differing retinal locations.

The NTSCI power spectra is typical of that of 'natural stimuli' in that power declines as a function of frequency(Van Hateren, 1997; Van Hateren *et al.*, 2002; Frazor & Geisler, 2006). As a result of the predominance of lower frequencies, most of the light intensity changes throughout the NTSCI occur over timescales accessible to goldfish cones and previously the NTSCI has been used to unlock several non-linear performance features of cones (Endeman & Kamermans, 2010; Howlett *et al.*, 2017). To ensure that all aspects other than skewness remained equal, we first picked short stretches from the NTSCI that were positively skewed, then simply flipped these around the mean to generate negatively skewed stimuli.

To generate Stimulus set#1 we divided the NTSCI (Van Hateren *et al.*, 2002) into one-second long stretches. Then from each stretch we subtracted its minimum value, adjusted their mean light intensities to be equal and picked stretches with similar power spectra and root mean

square (r.m.s.), and median contrasts (between 0.23 and 0.25). The r.m.s. and median contrast

for each stretch was calculated respectively as the ratio between the stretch's standard deviation

and its mean, and the ratio between its deviation from the median, and its median. To ensure

an absolute similarity between positively and negatively skewed stimuli, we selected only

stretches where the maximum value was not larger than 2 times the mean. Next, we chose

stretches with skewness's of 0.9, 1.6, 2.2. The skewness was calculated with the equation (1):

174
$$S = < \frac{(I - Imean)^3}{N\sigma^3} > \quad (1)$$

where N is the number of elements in the stretch, I corresponds to the light intensity of an

element, I_{mean} and σ are mean and standard deviation within the stretch and brackets denotes

averaging over the period.

We further narrowed our selection to three stretches all with similar power spectra (data not

shown). Power spectra were calculated by Welch's averaged periodogram method (Welch,

1967). No window function was used, the length of the Fourier transform was same as the

length of each corresponding data sequence. Total stimulus power was calculated as the

integral under power spectra, the differences in the total stimulus power were no more than

10%. Finally, an additional pink noise stimulus with zero-skew and similar power spectra was

added to the set. In total, Skew Stimulus set#1 consisted of seven 1-second stimuli.

Skew Stimulus set#2. This stimulus set consisted of three 4-second long stretches with a

skewness of 2.2, 0, and -2.2. They were generated in the same way as the Skew Stimulus set#1,

but with one additional condition: the degree of skewness delivered by the stimulus remained

unchanged by the cone's temporal filtering. This was ensured by first convolving the NTSCI

stretch with the mean photocurrent impulse response function (see below) obtained from

responses to Skew Stimulus set #1. The skew of the convolution product, representing the

"effective" stimulus, was then compared with the skew of the original stimulus (Figure 3). This

was further confirmed by determining the effective skewness after convolving the stimuli with

the impulse response function of each cone measured under Skew Stimulus set #2 conditions

(Figure 3B).

Calcium current isolation

To measure I_{Ca}, we used the mean voltage response (7 cells, 69 repeats in total) of cone

photoreceptors to Stimulus set #2 as the command voltages for the voltage clamp experiments.

To isolate I_{Ca} we followed the approach described by Fahrenhoft et al. (Fahrenfort *et al.*, 1999).

Briefly, to eliminate the calcium-dependent chloride current E_{Cl} was set at -41 mV and 100 μM

of niflumic acid added to the Ringer's solution; delayed rectifying, and hyperpolarization-

activated, potassium currents were blocked by substituting 5 mM of NaCl in Ringer's solution

with 5 mM of CsCl; light-activated conductances were saturated by a 20 µm spot of bright

white light focused on the cone outer segment; linear leak currents were removed by

subtraction. The leak current was estimated by clamping cones at -70mV, stepping to potentials

between -100 and 20 mV in 5 mV steps for 100 ms, calculating the mean current between 20

and 60 ms after the step onset, then determining the linear fit of the IV-relation between -100

and -60 mV (Vroman *et al.*, 2014; Kamar *et al.*, 2019).

Data analysis

For each cell, the skewness of its mean response to each stimulus was determined using

equation (1). In Figures 2, 3D and 8A, data was fitted using build-in Matlab least square

methods. All data analysis was performed in Matlab and Python.

Parallel cascade identification is the most rigorous method to describe the signal processing

properties of cone responses to naturalistic stimuli (Korenberg, 1991) . However, for practical

reasons our analysis only focuses on the estimation of the first parallel cascade, which is

effectively a linear filter followed by a static non-linearity. Apart from the computational and

descriptive simplicity, this approach is also justifiable as it accurately describes cone responses,

accounting for over 95% of the variance (Figure 8A).

The linear temporal filtering properties of a cone was described by its impulse response

function. Impulse response functions were estimated as the inverse Fourier transform of the

ratio between stimulus-response cross-power and stimulus power spectrum (Wiener, 1964;

Kim & Rieke, 2001). The spectra were calculated using Welch averaged periodogram method

(Welch, 1967). Stimuli and responses were detrended, divided into 500 ms long stretches, with

50% overlap, and windowed with a hamming function. The length of the Fourier transform

was 1024 ms. Cone integration time was calculated as the integral of the impulse response

function's initial polarization lobe (Daly & Normann, 1985).

To estimate the effective" stimuli perceived by each cone photoreceptors during each stimulus,

we convolved the cone's impulse response function during that condition with the original light

stimulus. Skews of these "effective" stimuli were calculated with equation (1). Discrepancies

between the skewness of the "effective" stimuli and the skewness of the cone's responses, were

considered a result of non-linear cone properties.

For "effective" Weber contrast steps (Figures 7&8A), light stimuli were first converted into

Weber contrasts steps (Figure 7) with equation (2):

233
$$C = \frac{(I - Imean)}{Imean} \ (\mathbf{2})$$

These Weber contrast steps were then convolved with a mean impulse response function to

obtain the "effective'' Weber contrast steps. The mean impulse response function used here

was the averaged voltage-response derived impulse response function of all 16 cones measured

in current clamp, which was subsequently scaled such that the integral under its curve yielded

one (Figure 7; Howlett et al., 2017). Even though a cone's impulse response function is affected

by the skewness of the stimulus (Figure 8B&C), we used this mean impulse response function

for the following reasons.

Firstly, observed changes to the impulse response function shape had little effect on the cone

response dynamics. We convolved Skew Stimulus Set #2 with impulse response functions

derived from responses to positive and negative stimulus of each of the cones (n=7) and found

that the resulting convolution products were highly correlated (r≈0.98 ± 0.01). Secondly,

averaging across different stimulus skew conditions was crucial to account for biases in the

estimate of the amplitude of the impulse response function arising from skewness of the light

stimuli (Chichilnisky, 2001; Simoncelli *et al.*, 2004; Bonin *et al.*, 2006; Tkačik *et al.*, 2014).

Finally, the goal of this analysis was to illustrate cone photoreceptor gain asymmetries rather

than to provide a veridical description of the gain dependence on stimulus contrast.

To estimate the cone's non-linear gain function parameters we fitted the relationship between

all the mean cone voltage responses and the "effective" Weber contrast steps (Figure 8A, 19000

data points) as the power function of input contrast (Van Hateren & Snippe, 2006) with

equation 3:

$$\Delta V = a * (C + b)^d + e \ \textbf{(3)}$$

Here ΔV denotes voltage response, C is the "effective" Weber contrast, a, b, d, and e are fit

parameters. At the biophysical level b) corresponds to the baseline rate of phosphodiesterase

activity (PDE), d) describes the inter-dependence between the PDE activity rate and the voltage

response, e) is proportional to the baseline concentration of cyclic guanosine monophosphate

and a) is a scaling factor (Van Hateren & Snippe, 2006). The quality of the fit was quantified

with the adjusted coefficient of determination (R^2). The highest R^2 (0.95) was obtained using

the following parameter values (value ± 95 % confidence interval): a=0.05138 ± 0.03506,

b=1.166 ± 0.072, d= -0.1251 ± 0.0959, e=-0.05046 ± 0.03552. Our estimation of the power

function (d) was close to that obtained by Van Hateren and Snippe (2006) in their theoretical

study.

Model

Photoreceptor responses were modelled in matlab using Van Hateren's model of vertebrate

photoreceptors (van Hateren & Snippe, 2007), which was shown to be remarkably precise in

capturing the processing steps involved in generating a cone's signal. Apart from the activation

of hyperpolarization-activated current (I_h; Howlett et al., 2017; Kamermans et al., 2017), the

model closely simulates all the cone's biophysical processing steps from the photon-initiated

activation of conopsins to the cGMP-regulated changes in the photocurrent, followed by the

generation of the voltage response. The model simulates cone photoreceptors as a cascade of

low-pass filters, a static (instantaneous and memoryless) non-linearity, and two divisive

feedback loops (van Hateren, 2005; van Hateren & Snippe, 2007). The low-pass filters

correspond to the kinetics of the different biophysical processing steps. The non-linearity

describes the inverse proportional dependence between light intensity and changes in the

cGMP concentration. The first feedback loop describes the regulation of the rate of cGMP

production by calcium influx through cGMP-gated channels. The second feedback loop

corresponds to the regulation of the membrane voltage by voltage-sensitive channels in the

cone inner segment. The cone's non-linear gain (Figure 8A) originates from the interplay

between the hydrolysis of the cGMP by PDE and calcium-regulated (feedback loop) production

of   the   cGMP   by   guanylyl   cyclase   (GC)

Figure 2

Example of the performance of the Van Hateren model for goldfish cones. Left. Voltage responses as Z-scores to

Skew Stimulus set #2 for a representative recorded cone (black line) and for the simulated cone (orange line). The

coefficient of determination (R^2) between these two traces was 0.98. Right. Impulse response functions obtained

from the simulated responses to negatively (red) and positively (blue) skewed stimuli. Parameters for the

simulation are listed in the Table 1.

We verified that the Van Hateren model could capture responses to skewed stimuli. For this

we fitted the model to the voltage responses of 7 goldfish cones recorded under Skew Stimulus

Set#2 conditions. The model parameters were modulated within the ranges determined by

Endeman and Kamermans (2010) and are shown in the Table 1. For all 7 cells, the correlation

coefficient between modelled and recorded voltage responses were no less than 0.97 (adjusted

$R^2 \geq 0.94$ Figure 2). Moreover, the impulse response functions estimated from the simulated

responses to positively and negatively skewed stimuli retained features of the impulse response

functions derived from the recorded voltage responses. For example, for both recorded and

simulated cone voltage responses, impulse response functions peaked 3 ms (8.5±1%,

p=0.00013) later under the negatively skewed stimulus compared to the positively skewed

condition but showed no statistically significant difference in their full width at half maximum

(FWHM), or in integration time (Figure 2). Thus, Van Hateren's model reproduces accurately

cone responses to skewed stimuli.

Next, we used the Van Hateren model to estimate the "effective" stimuli perceived by

salamander and cat cones in the studies by Tkacik et al. (2014) and Bonin et al. (2006),

respectively. To model salamander cones we adjusted the parameters of Van Hateren's model

such that the time course of the impulse response functions of simulated cones resembled those

of salamander cones reported by Rieke (2001) and Baccus and Meister (2002). The exact

simulation parameters are reported in the Table 1. Similarly, to model cat cones the Van

Hateren model parameters were adjusted so that the impulse response function time course

resembled the estimates made by Donner and Hemila (1996). For the cat, exact parameters of

the simulation are reported in the Table 1.

"Light" stimuli mimicking those used by Bonin et al. and Tkacik et al. (2014) where

respectively used to study the responses of the modelled cat (Figure 9) and salamander (Figure

10) cones to changes in skewness. The only difference was that for illustrative ease the

positively and negatively skewed stimuli were mirror copies of each other. Cat stimuli had a

r.m.s. contrast of 0.7, skews of ±0.4 and a flat power spectrum bandlimited to 124 Hz.

Salamander stimuli had a r.m.s. contrast of 0.2, skews of ±2 and a flat power spectrum

bandlimited to 30 Hz.

Statistics

320 All data are presented as mean ± SEM unless otherwise stated. Differences between groups

321 were tested using two-tailed t-test (Student, 1908).

Results

Cone responses vary with skewness

Direct light responses of photoreceptors are not affected by the spatial structure of stimuli and instead only depend on the intensity of light falling on the photoreceptor outer segment (see Material and Methods-Light stimuli). Hence, to assess the photoreceptor's contribution to skew discrimination, we exposed goldfish cones to a series of modified naturalistic time series of chromatic intensities (NTSCI) from the Van Hateren library (Van Hateren et al., 2002; Skew Stimulus set#1 in Material and Methods) and recorded their photocurrent and voltage responses (Figure 3A). Stimuli were equal in terms of mean intensity, root mean square contrast and median contrast, and had similar power spectra, while their skewness varied from -2.2 to +2.2. Positively and negatively skewed stimuli were mirror copies of each other, therefore any asymmetries between corresponding responses would reflect an asymmetry in the cone's processing.

To determine whether cones process negatively and positively skewed light stimuli differently, we plotted the skews of the photocurrent (Figure 3B) and voltage responses (Figure 3C) against the skews of the light stimuli. If there is no difference in processing, the skewness of the response will be equal to the light stimulus skewness and thus the data points will fall along a straight slope. However, if there is an asymmetry in the processing of positive and negative contrasts it would necessarily lead to the deviation of the data points from the grey line. Figures 3B&C shows that for positively skewed stimuli, the photocurrent and voltage responses are skewed to a lesser degree than are the light stimuli, whereas for the negatively skewed stimuli they are almost as equally skewed as the light stimuli. Note that the signal sign-inversion of the voltage response also sign-inverts its skewness. Figures 3B&C indicate an asymmetry in the processing of negatively and positively skewed stimuli by cone photoreceptors.

Figure 3

Cone processing affects stimulus skewness. **A.** Skew Stimulus set#1 (left) together with examples of voltage (middle), and photocurrent (right), responses recorded from cone photoreceptors. **B.** Photocurrent skewness as a function of stimulus skewness for Skew Stimulus set #1 (n=8). Each black square represents the skewness of a cell's photocurrent response to a skewed stimulus condition. The grey line depicts the situation where the response skewness is equal to the stimulus skewness. The panel shows cone responses to positive skewed stimuli were less skewed than their stimulus, but to stimuli with zero or negative skews cone responses were more skewed than their stimulus. **C.** Voltage response skewness as a function of stimulus skewness (n=9). Note that the voltage responses and stimuli skews have different sign due to the signal sign-inversion. The grey line describes the situation where response and stimulus skews are equal in magnitude. As was the case in B), cone responses to positive skewed stimuli were less skewed than their stimulus. To stimuli with zero or negative skews, the skewness of the cone's response was greater than that of its stimulus. **D.** Skewness of the voltage response as a function of the photocurrent. For each stimuli condition, the skews of each corresponding voltage response, and photocurrent, were averaged and the resulting means (±SEM) plotted as a voltage response verses photocurrent function. Note that as the cone's voltage

response is sign-inverted relative to its photocurrent their skews are also sign inverted. The situation where the

voltage response and photocurrent have equal magnitude of skewness is described by the grey line, which fits the

data with adjusted R^2 of ≈ 0.98. This indicates that the asymmetric processing of positively and negatively skewed

inputs originates in the phototransduction cascade. **E.** "Effective" skewness perceived by the cones as the function

of the skewness of the light stimuli. "Effective" skews were estimated from the convolution product of the light

stimuli with the cone's impulse response function (Figures 4A&B). The grey line depicts the situation, where

"effective" and response skews are equal. Note that for illustrative convenience the "effective" skewness

estimated from voltage responses were multiplied by -1. Figure 3E indicates that even though naturalistic stimuli

were used, some aspects of the stimuli were still unavailable to drive cone responses on account of the cone's

temporal filtering properties. This in turn reduced the range of "effective" skews by almost 30% relative to the

original -2.2 to +2.2 range of stimulus skews.

Figure 4

A. Representative examples of the cone impulse response functions obtained using the photocurrent (left) and

voltage responses (right) to Skew Stimulus set #1. **B.** Left. The mean cone impulse response function obtained as

the averaged photocurrent impulse response functions to all of the stimuli in Skew stimulus set#1 in all of the

recorded cells (n=8) (Figure 4A left). This mean photocurrent impulse response function was used, via

convolution, to identify segments of the NTSCI where the skewness of the original and "effective" stimuli

remained equal, a subsection of which formed Skew stimulus set#2. Right. Skewness of the "effective" stimuli as

a function of the skewness of the original stimuli for Skew Stimulus set #2. Violet squares correspond to the

"effective" skewness obtained by the convolution of the light stimuli with the mean impulse response function

shown on the left. Black squares depict the "effective" skewness 'perceived by each cone under Skew Stimulus

set #2 conditions. This was estimated by convolving each Skew Stimulus set #2 stimulus with the cone's impulse

400 response function obtained for corresponding stimulus. The grey line describes the condition where temporal

401 filtering does not affect stimulus skewness. Since all squares are aligned with the grey line, the "effective"

402 skewness is approximately equal to the original light stimulus skewness. Hence, for Skew Stimulus set #2 cone

403 temporal filtering does not change the skewness delivered by the stimuli.

The processing asymmetry originates exclusively within the phototransduction cascade

407 What are the cellular mechanisms leading to the differences in the processing of negatively and

408 positively skewed stimuli? To tease apart the relative contributions of the phototransduction

409 cascade and the voltage activated membrane conductances, we plotted the skewness of the

410 voltage responses and photocurrent against each other in Figure 3D. The grey line depicts the

411 condition where photocurrent and voltage response skews are equal in magnitude. All data

412 points fall on this line ($R^2 = 0.98$), meaning that the skewness of the photocurrent accounts for

413 98% of the skewness of the voltage responses. This means that the phototransduction cascade

414 is the primary source of the asymmetric processing of the positively and negatively skewed

415 stimuli.

Temporal filtering affects stimulus skewness

417 A cone's finite kinetics may act to temporally filter our light stimulus and thus affect the

418 skewness of the "effective" stimulus perceived by a cone. To determine the "effective" stimuli

419 skews we first estimated the temporal filters of the cone's photocurrent responses to each of

420 the skewed stimuli stretches following the Wiener approach (Wiener, 1964; Rieke, 2001;

421 Figures 4A&B). Next, we convolved the estimated filters with their corresponding skewed

422 stimuli to obtain the "effective" stimuli perceived by cone photoreceptors. Then we calculated

423 the skews of the "effective" stimuli and plotted them against the skews of the original light

424 stimuli in Figure 3E, where the grey line describes the situation where the "effective" skew is

equal to the original skew. Data points for the positively skewed light stimuli are lower than

the grey line and higher for the negatively skewed light stimuli. Consequently, temporal

filtering reduced the "effective" skewness perceived by the cones.

Asymmetry in the responses to "effective" stimuli

How do goldfish cones process these "effective" stimuli? Figure 3E shows that linear temporal

filtering reduces "effective" skewness and decreases the dynamic range over which responses

to skewed stimuli were measured by almost 30% (from ±2.2 to ±1.6). Therefore, we first

completed our data set by recording the cone's voltage responses to stimuli with "effective"

skews of ±2.2 (Figures 4C&5A). Next, we plotted the skews of the responses against the

"effective" stimulus skews (Figure 5B) and found that goldfish cones decrease the magnitude

of skewness when the stimuli are skewed positively and increase the magnitude of skewness

when the stimuli are skewed negatively.

Asymmetry in the processing of negatively and positively skewed stimuli is also reflected in

the amplitudes of the corresponding responses: standard deviations of the responses to

negatively skewed stimuli were up to 50% larger than the standard deviations of the responses

to positively skewed stimuli (Figure 5C).

Figure 5

Asymmetries in cone responses to positively and negatively skewed stimuli. **A**. Top. Depiction of Skew Stimulus

set #2. The key property of this stimulus set is that the light stimulus skewness is not affected by the temporal

filtering properties of cones (Figure 3B). Bottom. An example of a recorded voltage response to Skew Stimulus

set #2. **B**. Skews voltage and photocurrent responses against the "effective" stimuli skews. This figure illustrates

the relation between "effective" stimulus skew (see Figure 3E, Figure 4B) and the skewness for the photocurrent

(n=8) and voltage responses (n=9) of cones stimulated with Skew Stimulus set #1, and the voltage responses of

cones (n=7) recorded with Skew Stimulus set #2. This data indicates the cone response skewness differ from the

skewness of the "effective" stimulus. Cone responses are less skewed than the stimulus when it delivers higher levels of positive "effective" skew. The opposite occurs when the stimulus delivers higher levels of negative "effective" skew, the cone responses are more skewed than the stimulus. The grey line depicts the situation where the cone response and "effective" stimulus are equally skewed. For illustrative convenience, the voltage responses skews were multiplied by -1. **C**. Differences in cone response amplitudes (photocurrent – grey, voltage response – red) to negatively and positively skewed stimuli for each "effective" skew stimuli pair. When the "effective" skew magnitude was low (± 0.2) the photocurrent or voltage responses amplitudes were unaffected by the skew direction. However, at higher "effective" skew-magnitudes cone response amplitudes to negatively skewed stimuli were larger than for positively skewed stimuli (photocurrent difference: ± 0.7, 15.8% $\pm 4.52\%$ p=0.01; ± 1.6, 15.7 $\pm 4.33\%$ p=0.0083; voltage response difference: ± 0.7, 13.0 $\pm 4.67\%$ p=0.024; ± 1.6, 13.3 $\pm 2.64\%$ p=0.0.00097; ± 2.2, 45.2 ± 4.20 % p=0.00004). Changes in response amplitude were assessed as the ratio of standard deviations of a cell's response to corresponding negatively and positivity skewed stimuli. The "effective" skew values shown were estimated by the mean impulse response (Figure 7), obtained using the impulse response functions of all cells under all stimuli conditions. Data shown as mean $\pm$ SEM.

Asymmetry in the cone's output

To be perceived by the downstream neurons, asymmetries in the cone's responses to positively

and negatively skewed stimuli (Figures 5B&C) should be reflected in the synaptic release. In

photoreceptors, glutamate release is directly proportional to the calcium current (Schmitz &

Witkovsky, 1997; Thoreson *et al.*, 2004). Consequently, one can estimate changes in cone

glutamate release by recording its calcium current (I_{Ca}).

Figure 6

Asymmetrical processing of skewed stimuli at the cone I_{Ca} level. **A.** Top panel depicts the mean cone voltage

response to Skew Stimulus set #2, which was used as the stimulus for I_{Ca} measurements. An example I_{Ca}

measurement, when using this stimulus, is shown in the bottom panel. **B.** Skewness of I_{Ca} as a function of stimulus

skewness. The measurements of I_{Ca} were performed at three different potentials along its activation curve: -30mV

(cyan), -40mV (green) and -50mV (magenta). Note that as a decrease in voltage causes an increase in I_{Ca}, skews

for the stimulus and response have opposite signs. The grey line denotes the situation where I_{Ca} and its stimulus

are equally skewed. Regardless of the holding potential, data points are aligned with grey line, indicating that the

cone's I_{Ca} maintains any skewness present in its voltage response. Since cone-photoreceptor glutamate-release is

directly proportional to I_{Ca} (Schmitz & Witkovsky, 1997; Thoreson *et al.*, 2004), it is highly likely cone output

retains any I_{Ca} skewness.

We measured skew-dependent modulation of I_{Ca} by using recorded voltage responses to light

stimuli with "effective" skews of ±2.2 and -0.35 as the command voltages at three different

potentials (-30, -40, -50 mV) along the I_{Ca} activation curve (Figure 6A). To isolate I_{Ca}

responses, we blocked all other active conductance and subtracted the leak current. We then

plotted the skews of the I_{Ca} signals against the skews of the voltage responses (Figure 6B).

Note, that since depolarization produces an inward I_{Ca}, the skews of the voltage response and

I_{Ca} response have opposite signs.

Regardless of the clamping potential, all the data points in Figure 6B approximately fall on the

grey line, indicating that the skewness of the cone's signal is largely unaffected by the

transformation from membrane potential to the I_{Ca}. Consistent with this, the amplitudes of I_{Ca}

during the +2.2 and -2.2 skew conditions differed to the same degree (from $50 \pm 1.2\%$ to $54 \pm$

1% depending on the holding potential) as those of the voltage responses (Figure 5C). Thus,

for negatively and positively skewed stimuli, the asymmetries present at the cone's earlier

processing stages are preserved and even somewhat enhanced in the cone's output.

Figure 7

"Effective" Weber contrast steps. Left. Mean impulse response function used to obtain "effective" Weber contrast.

The mean impulse response was estimated by averaging the individual voltage-responses impulse-response

functions of all cells for all stimuli (n=16) (Figure 4A right, Figure 8B). This mean was scaled such that the

integral under its curve was 1. Right. Upper. Grey line: The original light stimuli of Skew Stimulus set #1 and

set#2 converted to Weber contrast steps. Black line: The "effective" Weber contrast steps obtained by the

convolution of the original Weber contrast steps with the mean impulse response function shown on the left.

Bottom. The averaged cone voltage response to each stimuli. For both upper and lower panels, the green dashed

lines separate the different stimuli stretches during which the effective skews were (from left to the right): -1.6, -

0.2, 0.2, -0.7, 0.06, 1.6, 0.7, -2.2, -0.35, 2.2.

Asymmetrical gain of the cone photoreceptors

What type of non-linear gain leads to the skew-dependent changes in the amplitude of the cone

responses? To determine how the voltage response amplitude depends on the Weber contrast

step we first converted the "effective" stimuli intensities into Weber contrast steps (Figure 7).

Next, we plotted baseline subtracted mean voltage responses (Figure 7) as a function of the

"effective" Weber contrast steps (Figure 8A). Figure 8A shows that cone responses are larger

for Weber contrast steps below -0.4 than they are for Weber contrast steps above 0.4. Hence,

the response gain of cones is greater for high negative, than for high positive, contrasts.

What kind of input-output relation supports the asymmetric gain of cones? We fitted the

relation between voltage responses and "effective" Weber contrast steps with equation (3)

(Figure 8A orange) and found that the cone voltage response is proportional to Weber contrast

step with an exponent of -0.1251 ($R^2 \approx$ 0.95, 95% confidence interval $\pm$ 0.0959). This

dependence is close to the one determined in the theoretical study by Van Hateren and Snippe

(2006), who suggested that voltage responses of the vertebrate cone is proportional to Weber

contrast with an exponent of -0.12.

Figure 8

The cone signal transfer properties. **A.** Asymmetric gain of cone photoreceptors. To estimate cone photoreceptor gain, we plotted the voltage response of cones as a function of "effective" Weber contrast. This relationship was well described (orange line, adjusted $R^2 = 0.95$) by the Weber contrast power function given in equation (3) and clearly indicates that cone gain is higher for negative contrasts than for the positive contrasts. The differences become prominent from Weber contrast steps around ±0.4 and are especially vivid for contrast steps beyond ±0.6. "Effective" Weber contrast was obtained by converting the light intensities into Weber contrast steps with equation (2), which were then convoluted with the mean cone impulse response function scaled such that integral under its curve yielded 1 (see methods, Figure 7). The voltage response shown is the baseline-subtracted average of all cells for each stimulus conditions. In total, there are 19000 data points in this figure. **B.** Stimulus skewness changes the shape of the cone's impulse response function. The figure depicts two representative examples of a cone's impulse response function in conditions with -2.2 (red) and +2.2 (blue) "effective'' stimulus skewness. **C.**

537   The cone impulse response functions shown in **B**, normalized by the amplitude of their initial lobe. On average,

538   impulse response functions peaked 3.6 ms, or 9 ± 1.0 %, later for negatively skewed stimulus than for the

539   positively skewed condition (p=0.0001, n=7) whereas the impulse response function FWHM (Δ = 1± 1.35 %,

540   p=0.37, n=7) and the cone integration time (Δ =1.1 ± 1.61 % ,p = 0.51, n=7) were unchanged.

Stimulus skewness affects the shape of the cone impulse response function

The processing time of cones is inversely proportional to light intensity such that responses to

steps of strong positive contrast peak earlier than responses to strong negative contrasts

(Nikonov *et al.*, 2000; Lee *et al.*, 2003; van Hateren, 2005; Van Hateren & Snippe, 2006;

Angueyra *et al.*, 2021). Therefore, one might expect that such a dependency would lead to a

difference in the time courses of the responses to positively and negatively skewed stimuli.

To test whether stimulus skewness has an effect on the cone kinetics, we compared impulse

response functions derived from the voltage responses to the -2.2 and the +2.2 "effective" skew

stimuli (Figure 8B). To better visualize the differences in kinetics we normalized these impulse

response functions by the amplitude of their initial lobe (Figure 8C). Interestingly, while on

average the cone impulse response functions peaked 3.6 ms (or 9 ± 1.0 %) later for the

negatively skewed stimulus than for the positively skewed stimulus (Figure 8C, p=0.0001,

n=7), there were no statistically significant differences neither in its full width at the half

maximum (FWHM) (Δ = 1.0 ± 1.35 %; p=0.37; n=7), nor in the cone's integration time (Δ=

1.1 ± 1.61 %; p=0.51; n=7). For the FWHM and integration time to remain unchanged while

the time to peak shifts suggests that during negatively skewed stimuli the reduced rise time of

the impulse response function's initial flank is largely offset by the peak falling back to baseline

at a faster rate.

Discussion

We studied responses of cone photoreceptors to differently skewed stimuli and found cone response amplitudes to negatively skewed stimuli are up to 50% greater than to positively skewed stimuli (Figures 2, 5A&C, 6A, 7). This amplitude difference originates from the asymmetrical weighting of positive and negative contrasts by the phototransduction cascade. Its gain is inversely proportional to Weber contrast steps raised to the power of -0.125 (Van Hateren and Snippe, 2006; Figure 8A) and may serve as the basis for the Blackshot mechanism proposed by Chubb et al. (1994, 2004). Additionally, we observed stimulus skewness changes the cone's impulse response function shape. For the normalized impulse response function, the rising flank was faster and the falling flank slower for positively compared to negatively skewed stimuli (Figure 8C).

The Blackshot mechanism

Psychophysical studies reported that humans can discriminate visual stimuli based on skewness (Chubb *et al.*, 1994, 2004; Graham *et al.*, 2016). Our results suggest that this discrimination starts as early as the phototransduction cascade. Chubb et al. (1994, 2004) described the sensitivity to skewness with the so-called Blackshot mechanism, which has a disproportionally strong response to high negative contrasts. Our data indicates that differences in the response to positively and negatively skewed stimuli originates in the phototransduction's asymmetric gain function, which leads to higher response amplitudes to negative contrasts than to positive contrasts (Figure 5C, Figure 8A). Moreover, in full accordance with psychophysical studies, the difference in cone response amplitudes to positive and negative contrasts becomes more prominent with larger contrast steps (Figure 8A). For the ±1.6 "effective" skew stimuli pair, where the maximal "effective" Weber contrast step was 0.5, the difference in the response was

about 15% while for the ±2.2 "effective" skew pair, where the maximal "effective" Weber

contrast step was 0.8, the difference was almost 50% (Figure 5C).

Asymmetries in responses to positive and negative contrasts are reported throughout the entire

visual system in various species (Laughlin, 1981; Van Hateren, 1997; Lee *et al.*, 2003;

Zaghloul *et al.*, 2003; Jin *et al.*, 2008; Yeh *et al.*, 2009; Endeman & Kamermans, 2010; Baden

et al., 2013; Kremkow *et al.*, 2014; Cooper & Norcia, 2015), including the human visual cortex

(Zemon *et al.*, 1988; Kremkow *et al.*, 2014). Although it was shown that differences in response

amplitudes to positive and negative contrasts are additionally amplified by the visual cortex

(Kremkow *et al.*, 2014), our data clearly indicates that the primary origin of this asymmetry is

within the cone's phototransduction cascade.

The phototransduction's asymmetric gain function enables cones to efficiently encode the

entire range of contrasts present in natural scenes. Photoreceptors encode changes in their input

with a graded output. Information theory states that such a system encodes a signal efficiently

only when the statistical distribution of its output is Gaussian, which implies a symmetrical

engagement of the system's dynamic range (Shannon, 1948; Van Hateren, 1997). On the other

hand, from a given mean, the light intensity cannot decrease by more than 100%, but can easily

increase by many orders of magnitude. This means that the dynamic range of positive contrasts

is wider than that of negative. Thus, although some visual scenes can be skewed negatively

(Tkačik *et al.*, 2014), the total distribution of contrasts at any given intensity is skewed

positively with negative contrasts being smaller in amplitude, but more frequent than positive

contrasts (Laughlin, 1983; Ruderman, 1994; Van Hateren, 1997; Ruderman *et al.*, 1998;

Cooper & Norcia, 2015). Consequently, to encode signals efficiently and to provide

symmetrical outputs, cones compensate for this asymmetry in their input by weighting high

positive contrasts with lower gain (Figure 8A), such that when stimulated with the entire range

of contrasts in natural scenes, cones provide a Gaussian output (Laughlin, 1983; Van Hateren,

1997; Endeman & Kamermans, 2010). We therefore suggest that the Blackshot mechanism is

simply a consequence of the more fundamental necessity to efficiently encode the range of

contrasts present in natural scenes.

It is also important to note that although when it comes to dealing with wide dynamic ranges

of light intensities the non-linear gain of photoreceptors is well-acknowledged, its influence on

perception is often ignored such that responses of the downstream neurons are often modelled

with an implicit assumption of the photoreceptor linearity (Chander & Chichilnisky, 2001; Kim

& Rieke, 2001; Rieke, 2001; Carandini *et al.*, 2005; Mante *et al.*, 2005; Manookin & Demb,

2006; Ozuysal & Baccus, 2012; Pitkow & Meister, 2012; Karamanlis & Gollisch, 2021;

Schreyer & Gollisch, 2021). However, there are a number of occasions where this non-linearity

is important to account for perceptual features (Kremkow *et al.*, 2014; Angueyra *et al.*, 2021).

For instance, higher spatial resolution for darker than for brighter patches is a consequence of

the cone photoreceptor non-linear gain (Kremkow *et al.*, 2014). Our data presents another

example of the far-reaching consequences of the early visual non-linearity and highlights

importance of accounting for the photoreceptor non-linearity when studying the visual system.

Shape of the impulse response function

We found the cone's impulse response function peaks ≈ 3.6 ms later for negatively skewed

stimuli whereas the cone's integration time is unaffected by stimulus skewness (Figure 8B&C).

The rising and falling flanks of the cone's impulse response are governed by different

biophysical mechanisms. The former is heavily influenced by the phosphodiesterase (PDE)

hydrolysis of cGMP. The time constant of this process is inversely proportional to the light

intensity. Hence, it decreases upon positive, and increases upon negative contrast steps

(Nikonov *et al.*, 2000; van Hateren, 2005; Endeman & Kamermans, 2010). The terminating

flank is largely regulated by the guanylyl cyclase (GC) mediated production of cGMP which

is modulated by the Ca^{2+} influx through the CNG channels. The time constant of this process

is not light-dependent but its gain is inversely proportional to the 4th power of light intensity

(Burns et al., 2002; van Hateren, 2005; van Hateren and Snippe, 2007). The interplay between

these two underlying processes is thought to account for the cone's impulse response function

shape, the asymmetric rising and falling response phases to sinusoidal stimuli, and for light

adaptation to decreases in light intensity being slower than for light intensity increases (Baylor

& Hodgkin, 1973; Lankheet *et al.*, 1991; Nikonov *et al.*, 2000; Lee *et al.*, 2003; van Hateren,

2005; Endeman & Kamermans, 2010; Angueyra *et al.*, 2021).

For our stimuli, the PDE hydrolysis of cGMP time-constant will have been shorter during

positively skewed stimuli as all large changes in light intensity were associated with positive

contrasts. This in turn manifest as a faster rate of change in the impulse response functions'

initial flank and hence an earlier time to peak. For negatively skewed stimuli, as all large

changes in light intensity were associated with negative contrast, GC mediated production of

cGMP was pronounced. The highly non-linear light-dependent gain function of this process,

and the ensuing Ca^{2+} influx when cones depolarized, increased the rate the cone CNG channels

reopened. This resulted in an increased decay rate for the terminating flank of the impulse

response function. Hence, the initial flank's faster onset rate during the positively skewed

stimulus is largely offset by the terminating flank's faster decay rate during the negatively

skewed stimulus. Consequently, the cone impulse response function integration time to

positive and negative skewed stimuli does not differ while it's time to peak does.

Relation to previous studies using skewed stimuli

Why did previous studies find stimulus skewness had little to no effect on RGC (Tkačik *et al.*,

2014) and LGN neurons (Bonin *et al.*, 2006)? We suggest a methodological factor. In both

studies, a large proportion of the stimulus power spectrum was outside the cone's temporal

frequency bandwidth. Indeed, Bonin et al. (2006) used white-noise stimuli bandlimited to 124

Hz to study cat LGN neurons responses, while the cat visual system barely responds to

frequencies above 32 Hz (Shapley & Victor, 1978; Mante *et al.*, 2005). Similarly, Tkacik et al.

(2014) studied salamander RGC responses with white-noise bandlimited to 30 Hz, whereas the

salamander retina hardly reacts to frequencies above 10 Hz (Kim & Rieke, 2001). Thus, in both

these studies a large part of their stimuli were 'filtered out' and the remaining "effective"

stimuli were only able to elicit marginal skew dependent effects.

To illustrate this point, we estimated the "effective" stimuli delivered by Bonin et al. (2006)

and Tkacik et al. (2014) using Van Hateren's cone photoreceptor model (van Hateren &

Snippe, 2007). Simulations of the cat cone indicate that while a wide range of "effective"

Weber contrasts were present (Figure 9A) the "effective" skewness was approximately half

that of the original stimuli employed, reducing from a range of ±0.4 to approximately ±0.2

(Figure 9B). Hence, both stimuli delivered largely similar distributions of "effective" Weber

contrasts. As a result, the simulated voltage responses to the positive and negatively skewed

stimuli only differed in amplitude by approximately 4% (Figure 9 C, D), which is within the

range of standard error estimates for the amplitude differences we find here (Figure 5C).

Simulations of the salamander cone reveal a different situation that none the less leads to the

same outcome. The "effective" skewness range remained relatively large despite being less

than half that of the original stimuli employed (±0.8 vs ±2, Figure 10B), but the range of

"effective" contrasts reduced to just ±0.2 Weber unit (Figure 10A). Over this limited range of

contrasts the cone photoreceptor gain is mostly symmetrical (Figure 8A) and as such the

voltage response amplitude to the negatively skewed stimulus was only 2% higher than for the

positively skewed stimulus (Figure 10C&D). Hence, even though the stimuli had substantially

different distributions of "effective" Weber contrasts, the range of contrast values they

delivered were too narrow to generate a notable effect.

To conclude, our results show that to study visual processes under varying skewness

conditions, the stimuli must be able to deliver sufficient levels of "effective" skewness over

sufficient ranges of "effective" Weber contrasts.

Figure 9

Simulated responses of cat cone photoreceptors. **A.** Light blue and light red lines depict the positively and

negatively skewed stimuli used by Bonin et al. (2006) and here for the simulation. Blue and red lines depict the

"effective" positively and negatively skewed stimuli, obtained by the convolution of the "original" stimuli with

the impulse response functions shown in **D**. **B.** Comparison of the skews of the "original" and "effective"

positively (blue) and "negatively" (red) skewed stimuli. Due to the temporal filtering the "effective" stimuli are

almost symmetrical around the mean, their skew range having decreased from ±0.4 to ≈ ±0.2. **C.** Simulated cone

voltage responses to the positively (blue) and negatively (red) skewed stimuli. The response amplitude to the

negatively skewed stimulus was 4% higher than the response amplitude to the positively skewed stimulus. The

simulated voltage response amplitudes were quantified by their standard deviations, as for Figure 5C. Parameters

for the simulation are listed in the Table 1. **D.** Impulse response functions derived from the simulated cat cone

voltage responses. These impulse response functions were used to estimate the "effective" stimuli shown in **A.**

Figure 10

Simulated responses of salamander cone photoreceptors. **A.** Light blue and light red lines depict the positively and negatively skewed stimuli used in the simulation. These stimuli had the same properties as those used by Tkacik et al.(2014). Blue and red lines depict the "effective" positively and negatively skewed stimuli, obtained by the convolution of the "original" stimuli with the impulse response functions shown in **D**. The temporal filtering leads to small Weber contrast ranges of the "effective" stimuli. **B.** Comparison of the skews of the "original" and "effective" positively (blue) and "negatively" (red) skewed stimuli. The salamander cone's temporal filtering reduced the stimulus skewness from ±2 to ±0.8. **C.** Simulated cone voltage responses to the positively (blue) and negatively (red) skewed stimuli. The response amplitude to the negatively skewed stimulus was 2% higher than the response amplitude to the positively skewed stimulus. The simulated voltage response amplitudes were quantified by their standard deviations, as for Figure 5C. Parameters for the simulation are listed in the Table 1. **D.** Impulse response functions derived from the simulated salamander cone voltage responses. These impulse response functions were used to estimate the "effective" stimuli shown in **A.**

References

Angueyra JM, Baudin J, Schwartz GW & Rieke F (2021). Multiple time scales of adaptation allow cones to encode the inputs created by visual exploration of natural scenes.

*bioRxiv*2021.02.13.431101.

Baccus SA & Meister M (2002). Fast and slow contrast adaptation in retinal circuitry. *Neuron*

36, 909–919.

Baden T, Schubert T, Chang L, Wei T, Zaichuk M, Wissinger B & Euler T (2013). A tale of

two retinal domains: Near-Optimal sampling of achromatic contrasts in natural scenes

through asymmetric photoreceptor distribution. *Neuron* **80,** 1206–1217.

Baylor DA & Hodgkin AL (1973). Detection and resolution of visual stimuli by turtle

photoreceptors. *J Physiol* **234,** 163–198.

Bonin V, Mante V & Carandini M (2006). The statistical computation underlying contrast gain

control. *J Neurosci* **26,** 6346–6353.

Burns ME, Mendez A, Chen J & Baylor DA (2002). Dynamics of cyclic GMP synthesis in

retinal rods. *Neuron* **36,** 81–91.

Carandini M, Demb JB, Mante V, Tolhurst DJ, Dan Y, Olshausen BA, Gallant JL & Rust NC

(2005). Do we know what the early visual system does? *J Neurosci* **25,** 10577–10597.

Chander D & Chichilnisky EJ (2001). Adaptation to Temporal Contrast in Primate and

Salamander Retina. *J Neurosci* **21,** 9904–9916.

Chichilnisky EJ (2001). A simple white noise analysis of neuronal light responses. *Netw*

Comput Neural Syst **12,** 199–213.

Chubb C, Econopouly J & Landy MS (1994). Histogram contrast analysis and the visual

segregation of IID textures. *J Opt Soc Am A* **11,** 2350.

Chubb C, Landy MS & Econopouly J (2004). A visual mechanism tuned to black. *Vision Res*

44, 3223–3232.

Cooper EA & Norcia AM (2015). Predicting Cortical Dark/Bright Asymmetries from Natural

Image Statistics and Early Visual Transforms. *PLoS Comput Biol* **11,** 1–25.

Daly SJA & Normann R (1985). Temporal Information Processing In Cones: Effects of Light

Adaptation on Temporal Summation and Modulation. *Vision Res* **25,** 1197–1206.

Donner K & Hemila S (1996). Modelling the spatio-temporal modulation response of ganglion

cells with difference-of-Gaussians receptive fields: Relation to photoreceptor response

kinetics. *Vis Neurosci* **13,** 173–186.

Endeman D & Kamermans M (2010). Cones perform a non-linear transformation on natural

stimuli. *J Physiol* **588,** 435–446.

Fahrenfort I, Habets RL, Spekreijse H & Kamermans M (1999). Intrinsic cone adaptation

modulates feedback efficiency from horizontal cells to cones. *J Gen Physiol* **114,** 511–

524.

Frazor RA & Geisler WS (2006). Local luminance and contrast in natural images. *Vision Res*

46, 1585–1598.

Graham D, Schwarz B, Chatterjee A & Leder H (2016). Preference for luminance histogram

regularities in natural scenes. *Vision Res* **120,** 11–21.

van Hateren H (2005). A cellular and molecular model of response kinetics and adaptation in

primate cones and horizontal cells. *J Vis* **5,** 331–347.

Van Hateren JH (1997). Processing of natural time series of intensities by the visual system of

the blowfly. *Vision Res* **37,** 3407–3416.

Van Hateren JH, Rüttiger L, Sun H & Lee BB (2002). Processing of natural temporal stimuli

by macaque retinal ganglion cells. *J Neurosci* **22,** 9945–9960.

van Hateren JH & Snippe HP (2007). Simulating human cones from mid-mesopic up to high-

photopic luminances. *J Vis* **7,** 1.

Van Hateren JH & Snippe HP (2006). Phototransduction in primate cones and blowfly

photoreceptors: Different mechanisms, different algorithms, similar response. *J Comp*

Physiol A Neuroethol Sensory, Neural, Behav Physiol **192,** 187–197.

Howlett MHC, Smith RG & Kamermans M (2017). A novel mechanism of cone photoreceptor

adaptation. *PLoS Biol* **15,** e2001210.

Jin JZ, Weng C, Yeh CI, Gordon JA, Ruthazer ES, Stryker MP, Swadlow HA & Alonso JM

(2008). On and off domains of geniculate afferents in cat primary visual cortex. *Nat*

Neurosci **11,** 88–94.

Kamar S, Howlett MHC & Kamermans M (2019). Silent-substitution stimuli silence the light

responses of cones but not their output. *J Vis* **19,** 1–11.

Kamermans W, Ariette M, Farajnia S, Howlett M & Kamermans M (2017). A quantitative

model for adaptation of in cone photoreceptors to contrast. ; DOI:

10.6084/m9.figshare.12813191.v1. Available at:

https://figshare.com/articles/poster/A_quantitative_model_for_adaptation_of_in_cone_p

hotoreceptors_to_contrast/12813191.

Karamanlis D & Gollisch T (2021). Nonlinear spatial integration underlies the diversity of

retinal ganglion cell responses to natural stimuli. **41,** 3479–3498.

Kim KJ & Rieke F (2001). Temporal contrast adaptation in the input and output signals of

salamander retinal ganglion cells. *J Neurosci* **21,** 287–299.

Korenberg MJ (1991). Parallel cascade identification and kernel estimation for nonlinear

systems. *Ann Biomed Eng* **19,** 429–455.

Kremkow J, Jin J, Komban SJ, Wang Y, Lashgari R, Li X, Jansen M, Zaidi Q & Alonso JM
(2014). Neuronal nonlinearity explains greater visual spatial resolution for darks than
lights. *Proc Natl Acad Sci U S A* **111,** 3170–3175.

Lankheet MJM, van Wezel RJA & van de Grind WA (1991). Light adaptation and frequency
transfer properties of cat horizontal cells. *Vision Res* **31,** 1129–1142.

Laughlin S (1981). A simple coding procedure enhances a neuron's information capacity.
Zeitschrift fur Naturforsch - Sect C J Biosci **36,** 910–912.

Laughlin S (1983). Matching Coding to Scenes to Enhance Efficiency. 42–52.

Lee BB, Dacey DM, Smith VC & Pokorny J (2003). Dynamics of sensitivity regulation in
primate outer retina: The horizontal cell network. *J Vis* **3,** 513–526.

Malchow RP & Yazulla S (1986). Separation and light adaptation of rod and cone signals in
the retina of the goldfish. *Vision Res* **26,** 1655–1657.

Manookin MB & Demb JB (2006). Presynaptic Mechanism for Slow Contrast Adaptation in
Mammalian Retinal Ganglion Cells. *Neuron* **50,** 453–464.

Mante V, Frazor RA, Bonin V, Geisler WS & Carandini M (2005). Independence of luminance
and contrast in natural scenes and in the early visual system. *Nat Neurosci* **8,** 1690–1697.

Nikonov S, Lamb TD & Pugh EN (2000). The role of steady phosphodiesterase activity in the
kinetics and sensitivity of the light-adapted salamander rod photoresponse. *J Gen Physiol*
116, 795–824.

Ozuysal Y & Baccus SA (2012). Linking the Computational Structure of Variance Adaptation
to Biophysical Mechanisms. *Neuron* **73,** 1002–1015.

Pitkow X & Meister M (2012). Decorrelation and efficient coding by retinal ganglion cells.

Nat Publ Gr **15,** 628–635.

Rieke F (2001). Temporal contrast adaptation in salamander bipolar cells. *J Neurosci* **21,** 9445–

9454.

Ruderman DL (1994). The statistics of natural images. *Netw Comput Neural Syst* **5,** 517–548.

Ruderman DL, Cronin TW & Chiao C-C (1998). Statistics of cone responses to natural images:

implications for visual coding. *J Opt Soc Am A* **15,** 2036.

Schmitz Y & Witkovsky P (1997). Dependence of photoreceptor glutamate release on a

dihydropyridine- sensitive calcium channel. *Neuroscience* **78,** 1209–1216.

Schreyer HM & Gollisch T (2021). Nonlinear spatial integration in retinal bipolar cells shapes

the encoding of artificial and natural stimuli. *Neuron* **109,** 1692-1706.e8.

Shannon CE (1948). A Mathematical Theory of Communication. **27,** 379–423.

Shapley RM & Victor JD (1978). The effect of contrast on the transfer properties of cat retinal

ganglion cells. *J Physiol*275–298.

Simoncelli EP, Pillow JW, Paninski L & Schwartz O (2004). Characterization of Neural

Responses with Stochastic Stimuli.

Student (1908). The probable Error of a Mean. *Biometrika* **6,** 1–25.

Thoreson WB, Rabl K, Townes-Anderson E & Heidelberger R (2004). A highly Ca2+-

sensitive pool of vesicles contributes to linearity at the rod photoreceptor ribbon synapse.

Neuron **42,** 595–605.

Tkačik G, Ghosh A, Schneidman E & Segev R (2014). Adaptation to changes in higher-order

stimulus statistics in the salamander retina. *PLoS One*; DOI:

10.1371/journal.pone.0085841.

Vroman R, Klaassen LJ, Howlett MHC, Cenedese V, Klooster J, Sjoerdsma T & Kamermans

M (2014). Extracellular ATP Hydrolysis Inhibits Synaptic Transmission by Increasing pH

Buffering in the Synaptic Cleft. *PLOS Biol* **12,** e1001864.

Welch PD (1967). Welchs Periodogram.pdf. *IEEE Trans Audio Electroacoust* **15,** 70–73.

Wiener N (1964). *Extrapolation, Interpolation, and Smoothing of Stationary Time Series*. The

MIT Press.

Yeh CI, Xing D & Shapley RM (2009). "Black" responses dominate macaque primary visual

cortex V1. *J Neurosci* **29,** 11753–11760.

Zaghloul KA, Boahen K & Demb JB (2003). Different circuits for ON and OFF retinal

ganglion cells cause different contrast sensitivities. *J Neurosci* **23,** 2645–2654.

Zemon V, Gordon J & Welch J (1988). Asymmetries in ON and OFF visual pathways of

humans revealed using contrast-evoked cortical potentials. *Vis Neurosci* **1,** 145–150.

Table 1

Parameter	Goldfish	Cat	Salamander
Lifetime of activated conopsin	8 – 31 ms	8 ms	88 ms
Lifetime of activated transducin	16 – 30 ms	12 ms	101 ms
Dark PDE activity	0.003 ms^{-1}	0.0028 ms^{-1}	0.003 ms^{-1}
Constant of the dependence of PDE activity on transducin activation	0.00004 - 0.00023	0.00016	0.0002
Apparent Hill coefficient of CNG channels	1	1	1
Hill coefficient of GC activation	4	4	4
Time constant of calcium extrusion	12 -28 ms	9 ms	24 ms
GC activation constant	0.1	0.09	0.1
Capacitive membrane time constant	15 ms	6 ms	15 ms
Parameter of membrane non-linearity	0.7 – 1.1	0.8	0.85
Constant of membrane nonlinearity	0.03 – 0.07	0.07	0.085
Time constant of membrane non-linearity	300 ms	120 ms	300 ms

Parameters of the Van Hateren model used to simulate the responses of goldfish, salamander

and cat cones. For the goldfish, model parameters were obtained by fitting cones responses

(n=7) to Skew Stimulus Set #2 while constraining the range the parameters varied to within

that determined by Endeman and Kamermans (2010). For the cat and salamander, parameters

were adjusted such that the simulated-cone's impulse response function time-to-peak

approximately matched that estimated, respectively, by Donner and Hemila (1996) and by

Rieke (2001) and Baccus and Meister (2002).