Title: Environmental filtering in marine fish communities is weakened during severe

hypoxia season

Abstract

Compositional variation among local communities reflects the differences in abiotic

environments, though the extent of the influence varies in natural systems. Generally, the

environmental filtering is expected to be strong if the environmental gradient encompasses

severe habitats where species sorting is highly likely to work. However, this hypothesis has

rarely been tested in dynamic systems, where the higher dispersal ability of species allows

them to easily respond to stressful environments. Here, with the dynamics of fish

communities in a Japanese bay revealed by environmental DNA analyses as a model case,

we examined how harmful seasonal hypoxia (low concentration of oxygen in bottom waters

in summer) affected the strength of environmental filtering. We found that, in summer,

dissolved oxygen (DO) concentration was significantly low and fish species richness

decreased in the bottom water compared to the surface, suggesting that the organisms were

adversely affected by the hypoxia. At the same time, the between-depths heterogeneity in

DO concentration was larger, but species composition was less divergent and the influence

of DO on species composition appeared weaker during summer. These results imply that

environmental filtering is weakened when the bottom water was characterized by extremely

severe environments. Furthermore, there was a shift in the species occurrence from bottom

to surface waters in summer that was consistent across species, suggesting that the

extremely severe hypoxia adversely affected fish species irrespective of their identity.

These results collectively suggest that environmental filtering is weaker during summer

despite more severity and heterogeneity in environments, most likely because individual

movements to avoid unpreferable environments in the bottom waters occurred

quasi-neutrally. By providing evidence against the prevailing understanding that

environmental filtering strongly works in severe environments, these findings invoke

further investigation on how the filtering acts in various conditions.

Keywords:

Community assembly, eDNA, environmental filtering, fish, hypoxia

Introduction

Understanding the processes structuring local communities has been a central challenge in ecology (Hutchinson 1959, HilleRisLambers et al. 2012). One of the primary determinants of local communities is the abiotic environment; species composition is thought to deterministically align with environmental gradients as determined by each species' requirements for some habitats or abiotic conditions (Grinnell 1917, Tilman 1982), a concept commonly referred to as 'environmental filtering'. However, it has also been acknowledged that more stochastic processes such as random movements of individuals among localities influence the community structure (Sale 1978, Hubbell 2001). Therefore, species composition can vary even if the localities are similar in their environmental conditions. Acknowledging that both deterministic and stochastic processes act in concert in nature, much research on community assembly has been devoted to understanding to what extent and in what situation the deterministic (or stochastic) community assembly matters (Cottenie 2005, Chase and Myers 2011).

Growing body of literature suggests that deterministic assembly plays an important role in more abiotically severe habitats (Chase 2007, Lepori and Malmqvist 2009, Guo et al. 2014), because the stressful environment filters the potential colonizers. The reduced potential species pool would hamper the importance of stochastic assembly by reducing the among-habitats variation in species composition and the probability of alternative states (Chase 2007). Therefore, environmental filtering is expected to be more influential when the environmental gradient encompasses severe conditions that strongly sort species. However, there exist a couple of studies providing opposing results that stochasticity is amplified in stressful environments (Lepori and Malmqvist 2009, Kim 2019). For example,

although the deterministic control is weak in streams without disturbance, it is also so in

highly disturbed habitats because the disturbance can induce extinction of individuals

irrespective of species traits (Lepori and Malmqvist 2009). Therefore, whether

environmental severity intensifies or dampens the deterministic control of environments is

still an open question in community ecology. Furthermore, as previous studies have focused

on organisms with limited or passive dispersal abilities (Chase 2007, Guo et al. 2014),

knowledge is particularly scarce in dynamic communities characterized by higher dispersal

ability of species. If species can perceive and easily respond to environmental severity, the

filtering may act more precisely. Otherwise, it is also probable that species actively avoid

unpreferable, though tolerable, environments, which dampens the species sorting by

environments.

In marine ecosystems, one of the environmental variables that most drastically

influence community structures is hypoxia or the decline in dissolved oxygen (DO; Diaz

and Rosenberg 1995, 2008, Breitburg et al. 2018). One of the typical forms of hypoxia is

seasonal: In summer, water warming forms the stratification of water columns, which

reduce the DO concentration in the bottom water to a critical level. As oxygen is essential

to organisms, the decreased DO adversely affects demersal organisms' performance (Levin

2003, Hrycik et al. 2017). Therefore, the bottom hypoxia is expected to serve as a strong

environmental filter by sorting out the species that are vulnerable to the lower DO

conditions (Levin 2003, Kodama and Horiguchi 2011), which aligns with the idea that

severe abiotic conditions intensify the deterministic assembly (Chase 2007, Lepori and

Malmqvist 2009, Guo et al. 2014). On the contrary, if the bottom hypoxia was so intense

that all species suffer from or avoid it to a similar extent, species composition would no

longer be correlated with the surface-bottom gradient in DO, making environmental

filtering weak.

The objective of this study is to determine how extremely severe conditions alter

the strength of environmental filtering in a dynamic system. We focused on the influence of

bottom hypoxia in summer on fish communities in Tokyo Bay, one of the largest bay areas

in Japan, as a model case. In this study system, it is already known that the water

stratification in summer causes hypoxia (Yasui et al. 2016) which severely affects

organisms in the bottom waters (Kodama and Horiguchi 2011). We have two alternative

hypotheses: (1) filtering by DO is stronger when the bottom hypoxia is more intense (in

summer) as fish species are different in their ability to tolerate lower oxygen concentration.

This predicts that explanatory power of DO for fish species composition is high in summer

because the species composition is clearly separated between oxygen-limited bottom and

oxygen-rich surface. Alternatively, (2) the filtering is weak in summer because most of the

fish species similarly suffer from or avoid the lower DO in the bottom. This alternative

hypothesis predicts that the explanatory power of DO is weak and surface-bottom

compositional differentiation is less evident in summer because the frequency of

occurrence in the bottom water is decreased irrespective of species identity. To this end, we

took advantage of the availability of data of seasonal dynamics of fish communities in

Tokyo Bay revealed by environmental DNA (eDNA) analyses (Hongo et al. 2021) which

has emerged as a powerful and efficient tool for monitoring biodiversity especially in

aquatic ecosystems (Bohmann et al. 2014; Deiner et al. 2017). By extracting genetic

materials shed from organisms from environmental samples (e.g., water and soils), this

technique allows for the non-invasive assessment of biodiversity, with a detection ability

that is comparable to that of traditional sampling methods (Thomsen et al. 2012; Sigsgaard

et al. 2017; Yamamoto et al. 2017). Importantly, owing to the short persistence (from a few

hours [Murakami et al. 2019] to several days [Thomsen et al. 2012]) and low diffusion rate

(less than 100 m; Port et al. 2016; Murakami et al. 2019) of eDNA in seawater, this

technique enables the estimation of contemporary and local species composition

(Yamamoto et al. 2016). Taking advantage of this technique allowed us to sample fish in

both bottom and surface waters, at a high frequency throughout a single year.

Materials and methods

Water sampling and eDNA sequencing

Water sampling was conducted monthly at 14 sites across Tokyo Bay, Japan (approximately 1380 km^2, 35°30' N, 139°50' E, Fig. S1) from January to December 2019. At each sampling location, 1 L of water was collected from the surface and bottom waters. Bottom depth varied among sites from approximately 6 to 70 m. Water samples were filtered and frozen onboard a ship. The frozen filters were then transferred to and stored in the laboratory. The extraction of eDNA and the construction of amplicon sequencing variants (ASVs) were performed as described by Hongo et al. (2021) using MiFish universal primers (Miya et al. 2015). The constructed ASVs were assigned to fish species using Blastn against the MitoFish database (Sato et al. 2018). As the correlation between eDNA concentration and abundance of target fish species is variable under natural conditions (Yates et al. 2019), the count data of ASVs was replaced to presence/absence data. The eDNA sequence data are available at the DNA Data Bank of Japan Sequence Read Archive under the accession number DRA010909.

Concentration of DO was measured at the same time as the water sampling. As expected, we found that DO of the bottom water was lower and the value was more divergent between surface and bottom during summer (Fig. 1). Specifically, from June to September, the minimum value of DO concentration among samples was lower than 2 mg/L, a common threshold of hypoxia in previous studies (Pihl et al. 1991, Hrycik et al. 2017, Breitburg et al. 2018). Following the established threshold in the literature, we determined these four months as "hypoxia season" and other months as "normoxia" (normal levels of oxygen). We note that even in the hypoxia season, the DO values

considerably varied among samples, with many of them being higher than the threshold

(Fig. 1). Nevertheless, we classified all samples of the months when the minimum value

falls below the threshold as the hypoxia season. This is because our hypothesis was that the

explanatory power of DO for fish species composition is strong (or weak) when the

environmental gradient encompasses the severe conditions, and we therefore needed to

analyze the samples obtained in the same environmental gradient as one unit.

Data analysis

To keep the dataset free from unexpected biases such as sequencing errors or

contamination from rivers or nearby fish markets, we filtered out the species that did not

meet the following two criteria from further analyses: (1) those whose names were not

matched during the search in Fishbase, a global database of fish (Froese and Pauly 2019;

https://www.fishbase.se/), and (2) those whose habitats were known to be out of Tokyo Bay

based on information from a range of studies (Nakabo 2002, Kouno et al. 2011). The first

criterion mainly filtered out hybrid species, while the second one, which was based on each

fish species' distribution in Japan and observation records in Tokyo Bay, removed fish

species known to be freshwater or endemic in other regions.

All statistical analyses were performed using R version 4.0.2 (R Core Team 2020)

and the figures were illustrated using the *ggplot2* package (Wickham 2016). First, to

examine the overall effects of lower DO on fish communities, we constructed generalized

linear models (GLMs) that modelled the fish species richness as a function of DO with

Poisson distribution (log link). The models were built for the bottom and surface samples

separately. Second, the explanatory power of DO for the fish species composition was

evaluated using distance-based redundancy analyses (db-RDA, Legendre and Anderson

1999) that takes the Jaccard dissimilarity matrix (based on presence-absence data) as a

response and DO value as an explanatory variable. The analyses were conducted separately

for each month, and the explanatory power was evaluated as the R^2 value that was adjusted

for the number of variables (Peres-Neto et al. 2006) using the *RsquareAdj* function in the

vegan package (Oksanen et al. 2019). The R^2 values were compared between hypoxia and

normoxia months by Wilcoxon's rank sum exact test. Third, to look at the surface-bottom

divergence of species composition, we conducted nonmetric multidimensional scaling

(NMDS) on the whole species composition data (n=281) using the *metaMDS* function. This

allows the compositional dissimilarity to be interpreted in a low-dimensional space by

positioning similar data nearby. For the NMDS analysis, we used the Jaccard dissimilarity

index on the presence-absence dataset and three dimensions (k=3) to facilitate convergence.

For the interpretation, we plotted values of the first two MDS axes in the figure and

illustrated ellipses of 80% confidence level for multivariate normal distributions for each

season (hypoxia or normoxia) and depth (surface or bottom). Difference in the species

composition among seasons and depths was tested using permutational multivariate

analysis of variance (PERMANOVA; Anderson 2001, the *adonis* function). In addition, we

included the interaction term between season and depth in the factorial design of the

PERMANOVA to test the difference in the degree of surface-bottom divergence of species

composition between seasons. Since the number of sampling sites was small (n=2) due to

logistical problems, we did not conduct the variation partitioning analysis and

PERMANOVA for the October data.

Finally, to examine whether the responses to bottom hypoxia are species-specific,

we compared the frequency of occurrence of fish species among different seasons (hypoxia

or normoxia) and depths (surface or bottom), using generalized linear mixed models

(GLMMs) with species-specific random intercept and coefficients. The models were

implemented using the *glmmTMB* function in the *glmmTMB* package (Brooks et al. 2017).

The response variable was the presence probability of species i in a sample s ($p_{i,s}$), and the

explanatory variables were seasons, depths, and their interaction:

$$logit(p_{i,s}) = \beta_{0,i} + \beta_{1,i} \times Season_s + \beta_{2,i} \times Depth_s + \beta_{3,i} \times Season_s \times Depth_s.$$

Note that the season and depth were transformed into dummy variables here (hypoxia: 0,

normoxia: 1, and surface: 0, bottom: 1). To assure sufficient statistical power, we restricted

this analysis to the frequently observed species, that is, top 10% most frequently observed

species (17 out of 168 species). Therefore, the dataset used in this GLMM consisted of

3842 presence or absence records (17 species x 226 samples). The species-specific

intercept ($\beta_{0,i}$) and coefficients ($\beta_{1,i}$, $\beta_{2,i}$, $\beta_{3,i}$) were modelled to follow normal

distributions, reflecting that the baseline of occurrence probability, the effects of seasons,

depth and their interaction on the probability may differ among species. Significance of the

estimated fixed coefficients suggests that the effects of the explanatory variables were

consistent while accounting for the difference among species. In light of our hypothesis, the

significantly positive coefficient of the fixed effect of the interaction term implies that the

relative occurrence in the bottom over surface water in normoxia was higher on average

across species, suggesting that species generally suffered from bottom hypoxia. Conversely,

if the coefficient of the interaction was not significant, the response to bottom hypoxia is

not evident at the average level among species. To further test the hypothesis on

species-specific responses to bottom hypoxia, we additionally constructed a GLMM that

was similar to the previous one except for that the coefficient of the interaction term ($\beta_{3,i}$)

was modelled as a fixed effect (β_3). We compared the fittings of the two GLMMs by the

likelihood ratio test using the *anova* function, which informs us whether modelling the

species-specific responses to bottom hypoxia improves the model. If the test indicated no

significant differences between them, we conclude that the effect of season-depth

interaction can be modelled as common across species.

Results

We obtained 308 water samples for the eDNA extraction, and gene amplification were successful for 281 samples. Among these, corresponding DO data were available for 225 samples (n = 22, 22, 22, 21, 20, 20, 18, 21, 20, 2, 15, and 22 from January to December, respectively). A total of 220 fish species were detected in the 225 samples. Of these, our first criterion filtered out 12 species, and the second criterion eliminated 40 species, resulting in 168 species retained in the subsequent analyses (Table S1).

The average species richness per sample was higher in the bottom layer (mean: 9.73, standard deviation: 4.78) than in the surface layer (mean: 6.89, standard deviation: 3.20) (Fig. S2). The GLMs that modelled fish species richness as functions of DO indicate that the species richness in the bottom water was positively correlated with DO (GLM: coefficient = 0.037; $p < 0.001$, Fig. 2), while that in the surface water decreased with DO (coefficient = -0.085; $p < 0.001$).

The variation partitioning analyses suggested that DO value explained 6.4% of the variation of fish species composition on average across months (n = 11). Note that the analysis was not conducted here for the October data because of small sample size. The explanatory power was 4.2% on average in hypoxia (n = 4), whereas that was 7.7% in normoxia (n = 7). Wilcoxon's rank sum exact test indicated that the explanatory power was significantly higher in normoxia (W = 3; $p = 0.040$, Fig. S3).

The PERMANOVA suggested that the fish species composition significantly differed between the depths (surface vs. bottom, $F_{1, 225} = 9.21$; $R^2 = 0.039$; $p < 0.001$, Fig. 3), and the seasons (hypoxia vs. normoxia, $F_{1, 225} = 5.37$; $R^2 = 0.022$; $p < 0.001$). Furthermore, the interactive term between the depth and season was also significant ($F_{1, 225}$

$= 2.58$; $R^2 = 0.011$; $p < 0.001$), suggesting that the degree of surface-bottom divergence

differed between the two seasons. The degree of surface-bottom divergence was smaller in

hypoxia, as the distance in the three-MDS dimensions between the centroids of the surface

and bottom samples was 0.27 for hypoxia, and 0.50 for normoxia (Fig. 3).

Relative frequency of occurrence of fish species in the surface and bottom waters

differed between hypoxia and normoxia. For example, *Konosirus punctatus*, the second

most frequently observed species (216 out of 281 samples), occurred in the bottom water

(73.3%, Fig. 4a) slightly more frequently than in the surface (72.2%) in normoxia. However,

this tendency reversed in hypoxia, and it more frequently occurred in the surface (90.0%)

than in the bottom (78.4%). Among the top 10% most frequently observed species

(*Engraulis japonicus*, *Konosirus punctatus*, *Lateolabrax japonicus*, *Hemitrygon akajei*,

Mugil cephalus, *Acanthopagrus schlegelii*, *Pennahia argentata*, *Callionymus valenciennei*,

Sardinella lemuru, *Scomber japonicus*, *Sardinops sagax*, *Scomberomorus niphonius*,

Seriola quinqueradiata, *Trachurus japonicus*, *Mustelus manazo*, *Branchiostegus japonicus*

and *Trichiurus lepturus* , in order of observed frequency), such an interactive effect of

depth and season on the frequency of occurrence (i.e. an increase of the relative occurrence

in the surface in hypoxia) was observed for 14 out of 17 species, excepting for Japanese

anchovy (*Engraulis japonicus*), Japanese sea bass (*Lateolabrax japonicus*) and Chub

mackerel (*Scomber japonicus*) (Fig. 4b). The GLMM, where the occurrence of each fish

species was modelled as a function of water depth, season, and their interaction with

species-specific random intercept and coefficients, showed that depth and season did not

consistently influence the occurrence probability (depth: coefficient $= 0.228$; standard error

$= 0.225$; $p = 0.311$, season: coefficient $= -0.409$; standard error $= 0.216$, $p = 0.058$, Table

1a). On the other hand, their fixed interactive term was significant (coefficient = 0.643;

standard error = 0.228; p = 0.005) with smaller random variance (0.328) than the other two

coefficients of marginal effects (depth: 0.489, season: 0.462). Indeed, the species-specific

coefficients of the interaction term estimated by random effects were positive for all the 17

species (Fig. S4). Furthermore, the likelihood test that compared the GLMM with another

one, where the random effect was not modelled for the interaction term (Table 1b), showed

no significant differences between the models (χ^2 = 2.56, df = 1, p = 0.109). These results

implies that the hypoxia at bottom has a negative effect on the occurrence probability

consistently across species.

Discussion

Previous studies have argued that environmental filtering works more in abiotically more severe habitats (Chase 2007, Lepori and Malmqvist 2009, Guo et al. 2014). However, this hypothesis has rarely been tested in dynamic systems, where higher dispersal ability of species allows them to easily respond to stressful environments. In the present study system, extremely low (< 2 mg/L) DO values in the sea water were observed during summer (from June to September, Fig, 1), which likely adversely affected fish species performance and decreased the species richness in the bottom waters (Fig. 2). Based on the discussion in previous studies (Chase 2007, Lepori and Malmqvist 2009, Guo et al. 2014), the lower DO in hypoxia season was expected to strongly sort specific sets of species from the regional species pool, as tolerating such extreme oxygen conditions would require certain morphological and physiological traits (Bickler and Buck 2007, Mandic et al. 2009). Against the expectation, the results showed that filtering by DO was weak (Fig. S3) and consequently the species composition of surface and bottom waters was less distinctive (Fig. 3) during the hypoxia.

Why weak environmental filtering during hypoxia?

These results may appear to be somewhat paradoxical, as filtering by DO was weaker during the hypoxia when the environmental conditions were more severe in the bottom and more heterogeneous between the surface and bottom waters (Fig. 1). Nevertheless, a series of evidence can explain this counterintuitive result. For example, Pihl et al. (1991) found that three demersal fish species differed in their response to bottom hypoxia in summer in Chesapeake Bay, USA. However, when the oxygen level fell below

the threshold (2 mg/L), all three species similarly showed shifts in their distribution from

deep to shallow waters (Pihl et al. 1991). Therefore, it is likely that, although the decrease

in DO in the bottom does serve as a filter for species composition, once it exceeds a certain

level of environmental severity, fish behaviors (e.g., survival, foraging, growth) will be

adversely affected regardless of species identity, at which point the filtering by DO no

longer plays a role in species sorting. Indeed, the GLMM in this study indicated that the

random variation in the coefficient of the interaction term between water depth and hypoxia

is smaller than that of depth and hypoxia (Table 1a), and the species-specific coefficients

were in the same direction for all the investigated species (Fig. S4). In addition, the two

GLMMs (with and without the random variation in the coefficient of the interaction term)

showed similar fittings. These results collectively imply that even though a large variation

in the occurrence probability in different depths and seasons exists among species (as

shown by the estimated large variance of random effects, Table 1), responses to the bottom

hypoxia (i.e., less frequent occurrence in the bottom waters in hypoxia) do not vary much

across species. Partially supporting this conclusion that the effect of bottom hypoxia is

non-species-specific, a meta-analysis revealed that effects of hypoxia on 29 fish species

were not well explained by species ecological characteristics (Hrycik et al. 2017).

Moreover, a recent study showed that the effect of hypoxia is variable among individuals of

the same species (Steube et al. 2021), suggesting that the filtering by DO may not be

precisely species-specific. Our results and other literature collectively support our second

hypothesis that the filtering is weak during hypoxia because extremely low DO

concentration influences the behavior of fish species in a similar way irrespective of species

identity.

The observed shifts in the relative occurrence frequency from the bottom to

surface waters during hypoxia can be a result of upward movements of individuals or

higher mortality in the bottom (Levin 2003, Diaz and Rosenberg 2008). Here, we speculate

that the former is responsible for the observed pattern. As fish are highly mobile, they can

respond sharply to the environmental changes by avoiding unfavorable conditions.

Therefore, when the bottom water was hypoxic in summer, they could move upward to

escape from the stressful habitat (Pihl et al. 1991). This would be particularly true for

free-swimming or floating species such as those frequently observed in this study.

Furthermore, we observed increases in species richness per sample in the surface water

during hypoxia, as well as the decreases in the bottom (Fig. S2), which supports this

explanation. On the other hand, we did not obtain strong evidence for the case that higher

mortality in the bottom is responsible for the observed shifts in the species occurrences. If

this was the case, species richness per sample should have only decreased in the bottom and

not increased in the surface, which was not in agreement with our findings (Fig. S2).

Although there may indeed be higher mortality rates in the bottom water during hypoxia, it

would be most influential for demersal species whose habitats are restricted to the bottom

environment. However, our dataset mainly consisted of free-swimming species. Among the

17 most frequently species, only six species (*Hemitrygon akajei*, *Acanthopagrus schlegelii*,

Callionymus valenciennei, *Seriola quinqueradiata*, *Mustelus manazo*, *Branchiostegus*

japonicus) are classified as demersal based on the description in Fishbase (Froese and

Pauly 2019). Though being beyond the scope of this paper, comparing the responses to

bottom hypoxia for between demersal and floating fish species would be one of the fruitful

investigations of future studies.

Implications for community assembly studies

Our results have general implication in the context of community assembly.

Previous studies have shown that deterministic assembly plays a more important role in

stressful habitats because the severe environment filters potential colonizers (Chase 2007,

Lepori and Malmqvist 2009, Guo et al. 2014). Our results are inconsistent with this in that

deterministic control by environments was weaker when the environment was more severe.

Nevertheless, a limited number of studies have suggested that this may be the case. Lepori

and Malmqvist (2009) have found that the deterministic control is strongest at the

intermediate level of disturbance. They agreed with the previous discussion that more

frequent disturbance intensified species sorting by selecting disturbance-tolerant species,

but they concluded that the disturbance that was too severe induced extinction of

individuals irrespective of species traits (Lepori and Malmqvist 2009), which would

hamper the environmental filtering. Similarly, Kim (2019) pointed to a possibility that

species living in extremely severe environments share similar tolerance traits. If this was

the case, environmental filtering may no longer play a role in such "neutral" communities

(Kim 2019). Our findings add to this series of evidence, by showing that weaker

environmental filtering was acting in more stressful environments.

Furthermore, the results showing weaker filtering in summer when the

surface-bottom gradient of DO was more distinct (Fig. 1) challenge the pervasive view that

stronger filtering is expected when the environmental gradient was more heterogeneous

(Kraft et al. 2015). Based on our results, we speculate that although increasing

environmental heterogeneity may indeed lead to stronger filtering, once the environmental

356 gradient becomes wide enough to encompass extremely severe conditions, the filtering

357 would become weak.

359 *Caveats and utility of eDNA analyses*

360  Although the ability of eDNA to detect species from environmental samples has

361 been established (Thomsen et al. 2012; Sigsgaard et al. 2017; Yamamoto et al. 2017), this

362 technique has rarely been applied to the evaluation of community assembly processes. In

363 this study, the eDNA analysis successfully reveals the seasonal shifts in the strength of

364 environmental filtering according to hypoxia, which illustrates its capability to be applied

365 to community assembly studies. However, the dataset we used contained only

366 presence–absence information, which would have missed some important information.

367 Estimating species abundance data from eDNA samples is challenging at the present time

368 as the correlation between eDNA concentration and abundance of target fish species has

369 not been established under natural conditions (Yates et al. 2019). To avoid potential biases,

370 we did not use the read number of DNA as an abundance index. Moreover, the detection of

371 eDNA is influenced not only by the presence of the species but also by the degradation and

372 accumulation of the eDNA itself ("ecology of eDNA" sensu Barnes and Turner 2016).

373 For example, it is known that the degradation of eDNA after release into water is faster at

374 higher temperatures (Tsuji et al. 2017, Jo et al. 2019), and it has been assumed that eDNA

375 accumulates in surface waters (Eichmiller et al. 2014, Moyer et al. 2014). We consider that

376 these biases are not the case here or at least do not change our main finding, because

377 species richness in surface water was high in summer where the degradation is expected to

378   be faster (Fig. S2), and the richness was generally higher in the bottom where the

379   accumulation is expected to be reduced (Fig. S2).

Conclusion

In ecosystems, species assemblages are determined by abiotic environmental

conditions, but to a variable extent. We found that the extremely severe environment, that is

the seasonal bottom hypoxia in coastal ecosystems, made the filtering weaker likely

because its adverse effect was non-species-specific. This result may seem to be

contradictory to the current pervasive view that deterministic control by abiotic conditions

is stronger in more stressful habitats. However, we speculate that these two contrasting

conclusions are compatible: While the increasing environmental severity surely strengthens

the filtering on species composition by sorting out vulnerable species, once it exceeds a

certain level, the filtering would no longer work because environments that are too severe

harm all the species in a similar way. Therefore, the deterministic control is expected to be

strongest at the intermediate level of environmental severity, a hypothesis to be further

confirmed in the future studies. Furthermore, by applying the eDNA analyses for detecting

community assembly processes for the first time to our best knowledge, this study

highlights the potential applications of this promising technique across a wide range of

disciplines of ecology.

References

Anderson, M. J. M. 2001. A new method for non-parametric multivariate analysis of

400   variance. Austral Ecology 26:32–46.

401   Barnes, M. A., and C. R. Turner. 2016. The ecology of environmental DNA and

402   implications for conservation genetics. Conservation Genetics 17:1–17.

403   Bickler, P. E., and L. T. Buck. 2007. Hypoxia tolerance in reptiles, amphibians, and fishes:

404   Life with variable oxygen availability. Annual Review of Physiology.

405   Bohmann, K., A. Evans, M. T. P. Gilbert, G. R. Carvalho, S. Creer, M. Knapp, D. W. Yu,

406   et al. 2014. Environmental DNA for wildlife biology and biodiversity monitoring.

407   Trends in Ecology and Evolution 29:358–367.

408   Breitburg, D., L. A. Levin, A. Oschlies, M. Grégoire, F. P. Chavez, D. J. Conley, V.

409   Garçon, et al. 2018. Declining oxygen in the global ocean and coastal waters.

410   Science 359.

411   Brooks, M. E., K. Kristensen, K. J. van Benthem, A. Magnusson, C. W. Berg, A. Nielsen,

412   H. J. Skaug, et al. 2017. glmmTMB Balances Speed and Flexibility Among

413   Packages for Zero-inflated Generalized Linear Mixed Modeling. The R Journal,

414   9(2), 378–400. https://journal.r-project.org/archive/2017/RJ-2017-066/index.html.

415   Chase, J. M. 2007. Drought mediates the importance of stochastic community assembly.

416   Proceedings of the National Academy of Sciences of the United States of America

417   104:17430–17434.

418   Chase, J. M., and J. A. Myers. 2011. Disentangling the importance of ecological niches

419   from stochastic processes across scales. Philosophical Transactions of the Royal

420   Society B: Biological Sciences 366:2351–2363.

Cottenie, K. 2005. Integrating environmental and spatial processes in ecological

community dynamics. Ecology Letters 8:1175–1182.

Deiner, K., H. M. Bik, E. Mächler, M. Seymour, A. Lacoursière-Roussel, F. Altermatt, S.

Creer, et al. 2017. Environmental DNA metabarcoding: Transforming how we

survey animal and plant communities. Molecular Ecology 26:5872–5895.

Diaz, R. J., and R. Rosenberg. 1995. Marine benthic hypoxia: A review of its ecological

effects and the behavioural response of benthic macrofauna MARINE.

Oceanography and Morine Biology: an Annual Review 33:245–303.

Diaz, R. J., and R. Rosenberg. 2008. Spreading Dead Zones and Consequences for Marine

Ecosystems. Science 321:926–929.

Eichmiller, J. J., P. G. Bajer, and P. W. Sorensen. 2014. The relationship between the

distribution of common carp and their environmental DNA in a small lake. PLoS

ONE 9:1–8.

Froese, R., and D. Pauly. (Eds). 2019. FishBase. World Wide Web electronic publication.

www.fishbase.org, version (12/2019).

Grinnell, J. 1917. The Niche-Relationships of the California Thrasher. The Auk

34:427–433.

Guo, H., K. Wieski, Z. Lan, and S. C. Pennings. 2014. Relative influence of deterministic

processes on structuring marsh plant communities varies across an abiotic gradient.

Oikos 123:173–178.

HilleRisLambers, J., P. B. Adler, W. S. Harpole, J. M. Levine, and M. M. Mayfield. 2012.

Rethinking Community Assembly through the Lens of Coexistence Theory. Annual

Review of Ecology, Evolution, and Systematics 43:227–248.

Hongo, Y., S. Nishijima, Y. Kanamori, S. Sawayama, K. Yokouchi, N. Kanda, S. Oori, et

al. 2021. Fish environmental DNA in Tokyo Bay: A feasibility study on the

availability of environmental DNA for fisheries. Regional Studies in Marine

Science 47:101950.

Hrycik, A. R., L. Z. Almeida, and T. O. Höök. 2017. Sub-lethal effects on fish provide

insight into a biologically-relevant threshold of hypoxia. Oikos 126:307–317.

Hubbell, S. P. 2001. The Unified Neutral Theory of Biodiversity and Biogeography.

Hutchinson, G. E. 1959. Homage to Santa Rosalia or Why Are There So Many Kinds of

Animals? The American Naturalist 93:145–159.

Jo, T., H. Murakami, S. Yamamoto, R. Masuda, and T. Minamoto. 2019. Effect of water

temperature and fish biomass on environmental DNA shedding, degradation, and

size distribution. Ecology and Evolution 9:1135–1146.

Kim, D. 2019. Selection of scale can reverse the importance of stochastic controls on

community assembly. Physical Geography 40:111–126.

Kodama, K., and T. Horiguchi. 2011. Effects of hypoxia on benthic organisms in Tokyo

Bay, Japan: A review. Marine Pollution Bulletin 63:215–220.

Kouno, H., H. Kanou, and T. Yokoo. (Eds.) 2011. A Photographic Guide to the Fishes in

Tokyo Bay. Tokyo, Heibon-sha. [in Japanese]

Kraft, N. J. B., P. B. Adler, O. Godoy, E. C. James, S. Fuller, and J. M. Levine. 2015.

Community assembly, coexistence and the environmental filtering metaphor.

Functional Ecology 29:592–599.

Legendre, P., and M. J. Andersson. 1999. Distance-based redundancy analysis: Testing

multispecies responses in multifactorial ecological experiments. Ecological

Monographs 69:1–24.

Lepori, F., and B. Malmqvist. 2009. Deterministic control on community assembly peaks at

intermediate levels of disturbance. Oikos 118:471–479.

Levin, L. A. 2003. OXYGEN MINIMUM ZONE BENTHOS: ADAPTATION AND

COMMUNITY RESPONSE TO HYPOXIA. Oceanography and Marine Biology:

an Annual Review 41:1–45.

Mandic, M., A. E. Todgham, and J. G. Richards. 2009. Mechanisms and evolution of

hypoxia tolerance in fish. Proceedings of the Royal Society B: Biological Sciences

276:735–744.

Miya, M., Y. Sato, T. Fukunaga, T. Sado, J. Y. Poulsen, K. Sato, T. Minamoto, et al. 2015.

MiFish, a set of universal PCR primers for metabarcoding environmental DNA from

fishes: Detection of more than 230 subtropical marine species. Royal Society Open

Science 2.

Moyer, G. R., E. Díaz-Ferguson, J. E. Hill, and C. Shea. 2014. Assessing environmental

DNA detection in controlled lentic systems. PLoS ONE 9.

Murakami, H., S. Yoon, A. Kasai, T. Minamoto, S. Yamamoto, M. K. Sakata, T. Horiuchi,

et al. 2019. Dispersion and degradation of environmental DNA from caged fish in a

marine environment. Fisheries Science 85:327–337.

Nakabo, T. (Ed.) 2002. Fishes of Japan with Pictorial Keys to the Species. Kanagawa,

Tokai University Press. [in Japanese]

Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R.

488   Minchin, R. B. O'Hara, G. L. Simpson, P. Solymos, et al. 2019. vegan: Community

489   Ecology Package. R package version 2.5-6.

490   https://CRAN.R-project.org/package=vegan

491   Peres-Neto, P. R., P. Legendre, S. Dray, and D. Borcard. 2006. Variation partitioning of

492   species data matrices: Estimation and comparison of fractions. Ecology

493   87:2614–2625.

494   Pihl, L., S. P. Baden, and R. J. Diaz. 1991. Effects of periodic hypoxia on distribution of

495   demersal fish and crustaceans. Marine Biology 108:349–360.

496   Port, J. A., J. L. O'Donnell, O. C. Romero-Maraccini, P. R. Leary, S. Y. Litvin, K. J.

497   Nickols, K. M. Yamahara, et al. 2016. Assessing vertebrate biodiversity in a kelp

498   forest ecosystem using environmental DNA. Molecular Ecology 25:527–541.

499   R Core Team. 2020. R: A language and environment for statistical computing. R

500   Foundation for Statistical Computing, Vienna, Austria. URL

501   https://www.R-project.org/

502   Sale, P. F. 1978. Coexistence of coral reef fishes - a lottery for living space. Environmental

503   Biology of Fishes 3:85–102.

504   Sato, Y., M. Miya, T. Fukunaga, T. Sado, and W. Iwasaki. 2018. MitoFish and mifish

505   pipeline: A mitochondrial genome database of fish with an analysis pipeline for

506   environmental DNA metabarcoding. Molecular Biology and Evolution

507   35:1553–1555.

508 Sigsgaard, E. E., I. B. Nielsen, H. Carl, M. A. Krag, S. W. Knudsen, Y. Xing, T. H.

509   Holm-Hansen, et al. 2017. Seawater environmental DNA reflects seasonality of a

510   coastal fish community. Marine Biology 164:1–15.

511 Steube, T. R., M. E. Altenritter, and B. D. Walther. 2021. Distributive stress: individually

512   variable responses to hypoxia expand trophic niches in fish. Ecology 102.

513 Thomsen, P. F., J. Kielgast, L. L. Iversen, P. R. Møller, M. Rasmussen, and E. Willerslev.

514   2012. Detection of a Diverse Marine Fish Fauna Using Environmental DNA from

515   Seawater Samples. PLoS ONE 7:1–9.

516 Tilman, D., S. S. Kilham, and P. Kilham. 1982. Phytoplankton community ecology: the role

517   of limiting nutrients. Annual review of ecology and systematics. Volume 13

518   13:349–372.

519 Tsuji, S., M. Ushio, S. Sakurai, T. Minamoto, and H. Yamanaka. 2017. Water

520   temperature-dependent degradation of environmental DNA and its relation to

521   bacterial abundance. PLoS ONE 12:1–13.

522 Wickham H 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New

523   York. ISBN 978-3-319-24277-4, https://ggplot2.tidyverse.org.

524 Yamamoto, S., R. Masuda, Y. Sato, T. Sado, H. Araki, M. Kondoh, T. Minamoto, et al.

525   2017. Environmental DNA metabarcoding reveals local fish communities in a

526   species-rich coastal sea. Scientific Reports 7:1–12.

527 Yamamoto, S., K. Minami, K. Fukaya, K. Takahashi, H. Sawada, H. Murakami, S. Tsuji, et

528   al. 2016. Environmental DNA as a "snapshot" of fish distribution: A case study of

529   Japanese jack mackerel in Maizuru Bay, Sea of Japan. PLoS ONE 11:1–18.

Yasui, S., J. Kanda, T. Usui, and H. Ogawa. 2016. Seasonal variations of dissolved organic

matter and nutrients in sediment pore water in the inner part of Tokyo Bay. Journal

of Oceanography 72:851–866.

Yates, M. C., D. J. Fraser, and A. M. Derry. 2019. Meta□analysis supports further

refinement of eDNA for monitoring aquatic species□specific abundance in nature.

Environmental DNA 1:5–13.

Figures and tables

Figure 1 Temporal trend of concentration of dissolved oxygen (DO) of each sample.

Open and filled circles represent samples in surface and bottom waters,

respectively. The minimum value of DO concentration was lower than the

common threshold (2 mg/L) from June to September, the period we determined

as hypoxia season (coloured with orange).

Figure 2   Relationship between dissolved oxygen (DO) concentration and fish species

547   richness. Open and filled circles represent samples in surface and bottom waters,

548   respectively. The vertical broken line indicates the threshold for the hypoxia (2

549   mg/L). The lines were drawn based on the generalized linear models separately

550   for the depths, after confirming the statistical significance ($p < 0.05$) of the

551   slopes (see the main text).

555   Figure 3   Nonmetric multidimensional scaling (NMDS) results for species composition

556   based on Jaccard dissimilarity (dimension = 3, stress = 0.190). Each ellipse

557   represents the 80% confidence level of the samples of each category (depth and

558   season) in the first two MDS axes. The colour and line type of the ellipses

559   correspond to season (orange: hypoxia, black: normoxia) and depth (solid:

560   surface, broken: bottom), respectively. Background points similarly depict the

561   value of the first two MDS axes of samples, with the colour and point type

562   corresponding to season and depth (open: surface, filled: bottom), respectively.

566   Figure 4   Observed relative occurrence probability in the surface and bottom waters of

567   fish species in hypoxia and normoxia. (a) An example for *Konosirus punctatus*,

568   the second most frequently observed species. The relative occurrence in surface

569   compared to bottom is higher in hypoxia. (b) The difference in the occurrence

570   probability in the surface and bottom waters in different seasons for the top

571   10% most frequently observed species (17 species). The points representing the

572   same species are linked with lines, and a bold line corresponds to the example

573   in (a). The increase of the relative occurrence in the surface in hypoxia similar

574   to (a) was observed for 14 species.

Table 1 Results of the two GLMMs that modelled the presence of species as a function of

water depth (surface or bottom), season (hypoxia or normoxia), and their

interaction with different modelling for the random effects: (a) Species-specific

random effects were modelled for the intercept, and coefficients of the three

explanatory variables. (b) Species-specific random effects were modelled for the

intercept and coefficients of depth and season only (not for the interaction).

Estimated coefficients, standard errors (SE), p-values of the fixed effects,

estimated variance of the random effect are shown. Significant estimates are

shown with bold. The likelihood ratio test indicates that the fits of these models

were not significantly different ($Chi^2 = 2.56$, df = 1, $p = 0.109$).

(a)	Fixed effects			Random effects
	Estimate	SE	p	Variance
Intercept	**-0.838**	**0.422**	**0.047**	2.81
Depth (Bottom)	0.228	0.225	0.311	0.489
Season (Normoxia)	-0.409	0.216	0.058	0.462
Depth (Bottom): Season (Normoxia)	**0.643**	**0.228**	**0.005**	0.328

(b)	Fixed effects			Random effects
	Estimate	SE	p	Variance
Intercept	-0.857	0.449	0.056	3.20
Depth (Bottom)	0.245	0.248	0.323	0.671
Season (Normoxia)	-0.387	0.244	0.112	0.676
Depth (Bottom): Season (Normoxia)	**0.642**	**0.179**	**< 0.001**	–