Natural tolerance to transposition is associated with double-strand break repair and germ-cell differentiation

Jyoti Lama[1], Satyam Srivastav[1], Sadia Tasnim[1], Donald Hubbard[1] , Savana Hadjipanteli[1] & Erin S. Kelleher[1*]

[1] Department of Biology and Biochemistry, University of Houston

[*] To whom correspondence should be addressed: eskelleher@uh.edu

Abstract

Transposable elements (TE) are mobile genetic parasites whose unregulated activity in the germline causes DNA damage and sterility. While the regulation of TE mobilization by hosts is studied extensively, little is known about mechanisms that could allow germline cells to persist in the face of genotoxic stress imposed by active transposition. Such tolerance mechanisms are predicted to be beneficial when new TEs invade and host repression has not yet evolved. Here we use hybrid dysgenesis—a sterility syndrome of *Drosophila* caused by transposition of invading DNA transposons—to uncover genetic variants that confer tolerance to transposition. Using a panel of highly recombinant inbred lines of *Drosophila melanogaste*r, we identified two linked quantitative trait loci (QTL), that determine tolerance in young and old females, respectively. Through transcriptomic and phenotypic comparisons, we provide evidence that young tolerant females exhibit enhanced repair of double-stranded breaks, explaining their ability to withstand high germline transposition rates. We furthermore identify the germline differentiation factor *brat* as an independent tolerance factor, whose activity may promote germline maintenance in aging dysgenic females. Together, our work reveals the diversity of potential tolerance mechanisms across development, as well as tolerant variants that may be beneficial in the context of *P*-element transposition.

INTRODUCTION

Transposable elements (TE) are mobile DNA sequences that spread through host genomes by replicating in germline cells. Although individual TE insertions are sometimes beneficial, genomic TEs are foremost genetic parasites [reviewed in 1]. Unrestricted transposition not only produces deleterious mutations, but also double-stranded breaks (DSBs) that lead to genotoxic stress in developing gametes. Generally, hosts avoid the fitness costs of invading parasites, pathogens and herbivores by two distinct mechanisms: resistance and tolerance [2–4]. Resistance reduces parasite proliferation, whereas tolerant individuals experience reduced fitness costs from parasitism. With respect to TEs, host resistance has been the focus of extensive research, and occurs through production of regulatory small RNAs that transcriptionally and post-transcriptionally silence TEs in the germline [5–7]. By contrast, tolerance mechanisms that could ameliorate the fitness costs of transposition during gametogenesis remain largely unstudied.

The lack of research on tolerance in part reflects the ubiquity of resistance, since in the absence of high transposition rates, tolerance will not be beneficial or apparent. For example, in *Drosophila melanogaster* all actively-transposing TE families are silenced in developing gametes by the Piwi-interacting RNA (piRNA) pathway [5]. However, host genomes are frequently invaded by new TE families, against which they lack piRNA mediated resistance. Following these invasions, tolerant genetic variants may be critical for maintaining host fertility until resistance evolves. A classic example of this occured with the *P*-element DNA transposon, which invaded natural populations of *D. melanogaster* around 1950 [8–10]. When males bearing genomic *P*-elements (P-strain) are mated to naive females lacking *P*-elements and corresponding piRNAs (M-strain), they produce dysgenic offspring that do not regulate *P*-elements in germline cells [11]. A range of fertility effects result from unregulated *P*-element transposition, including the complete loss of germline cells and sterility [12]. Interestingly, naive M genotypes differ in their propensity to produce dysgenic progeny when crossed to reference *P*-strain males, suggesting the presence of tolerant variants [8,10,13,14].

In dysgenic offspring, *P*-element transposition occurs in germline cells throughout the life cycle of the fly, providing multiple opportunities for tolerant phenotypes to emerge. Starting at the second-instar larval stage dysgenic females exhibit reduced primordial germ cells (PGCs), suggesting an early onset of *P*-element transposition [15–17]. Dysgenic PGC loss is partially suppressed by overexpression of *myc*, which encodes a transcription factor that promotes stem cell maintenance [17]. PGC loss may also be suppressed by mutations in *checkpoint kinase 2* (*chk2*), a key factor in germline response to DSBs [18,19]. Tolerance of PGCs to *P*-element transposition could therefore arise through increased signaling for stem cell maintenance, or increased DNA repair in damaged PGCs.

Similar to larvae, mechanisms that reduce accumulated DNA damage, such as DNA repair, could also confer tolerance in adult females. In mature dysgenic ovaries, differentiating pre-meiotic cells undergo *chk2*-dependent cell-death at an elevated rate [15,20]. However,

unlike their larval precursors the PGCs, adult germline stem cells (GSCs) are not lost at a high
rate due to *P*-element transposition [15,20]. Rather, the more dramatic phenotype is that *P*-
element transposition causes delay in differentiation of cytoblasts (CBs), the immediate progeny
of GSCs, which results in a temporary block to oogenesis [15,20,21]. Therefore, tolerance could
also emerge in adult females through mechanisms that facilitate the escape of CBs from
arrested differentiation.
7
Through QTL mapping in a panel of highly recombinant inbred lines from the *Drosophila*
Synthetic Population Resource (DSPR Population A RILs, [22]), we recently uncovered a
natural tolerance allele that is associated with reduced expression of *bruno*, a female germline
differentiation factor [14]. Here we present results from a second QTL mapping study in an
independent panel of DSPR RILs (Population B, [22]). We describe two natural alleles that
determine germline tolerance to *P*-element activity in young and aged females, respectively. We
further interrogated the tolerance phenotype by contrasting RNA expression, small RNA
expression, and radiation sensitivity between tolerant and sensitive genotypes, as well as by
performing mutational analysis of the candidate tolerance factor *brat*. Our results suggest that
young tolerant females enjoy enhanced DSB repair when compared to sensitive genotypes,
allowing them to minimize dysgenic PGC loss. In contrast, we uncover the germline
differentiation factor *brat* as a candidate tolerance factor in aged females. Together our results
reveal the complexity of natural variation in TE tolerance, and highlight potential targets of
positive selection following *P*-element invasion in natural populations of *D. melanogaster*.

RESULTS

23 1. QTL mapping of 2nd chromosome centromere:

24
The DSPR RILs are all *P*-element free M-strains, which were isolated from natural
populations before the *P*-element invasion [22]. We therefore screened for tolerant alleles
among the panel B RIL genomes by crossing RIL females to males from the reference P-strain
Harwich, and examining the morphology of the F1 ovaries (**Figure 1a**). Atrophied ovaries are
indicative of germline loss resulting from *P*-element activity, while non-atrophied ovaries are
indicative of tolerance [14,23]. Since dysgenic females differ across development [15], and
some females exhibit age-dependent recovery from *P*-element hybrid dysgenesis [24], we
phenotyped F1 females at two developmental time points: 3 days and 21 days post-eclosion.

Similar to our observations with the Population A RILs [14], we found continuous
variation in the frequency of ovarian atrophy among dysgenic offspring of different RIL mothers,
indicating genetic variation in tolerance (**Supplemental table S1 and 2**). Based on a combined
linear model of F1 atrophy among 3 and 21 day old females, we estimated the broad-sense
heritability of tolerance in our experiment to be ~42.5%. However, the effect of age on the
proportion of F1 atrophy was significant but minimal (X^2= 7.03, df = 1, p-value = 0.008) with 3-

day-old females showing only 0.7% increase in atrophy as compared to 21-day-old females.

Therefore, age-dependent recovery from dysgenic sterility is not common among the genotypes

we sampled.

To identify the genomic regions associated with genetic variation in germline tolerance,

we performed QTL mapping using the published RIL genotypes [22]. We found a large QTL

peak near the 2nd chromosome centromere in both 3 and 21 day-old F1 females **(Figure 1b,**

Table 1; Supplemental table S3 and S4). However, the genomic intervals within which the

causative change separating sensitive and tolerant most likely resides are non-overlapping

between the 3 and 21 day-old data sets **(Figure 1c, Table 1)**. The major QTL in 21 day-old

females (hereafter, QTL-21d) resides in the euchromatic region and is quite small (990 kb)

compared to the major QTL in 3 day-old females (hereafter QTL-3d), which spans the

centromere and pericentromeric regions (9.6 Mb, **Figure 1d**). Therefore, there are likely at least

two polymorphisms that influence tolerance near the 2nd chromosome centromere, one of which

is more important in young 3-day old females, and the other of which is more important in 21

day-old females.

We further evaluated the age-specific effect of two linked QTL through haplotype

analysis. We modeled residual F1 ovarian atrophy as a function of QTL haplotype for the 3 day

and 21 day peaks, thereby disentangling synergistic (e.g. sensitive 3d, sensitive 21d) from

opposing (e.g. sensitive 3d, tolerant 21d) allelic combinations (**Supplemental figure S4**). We

observed that the 3 day old QTL is solely-determinant of tolerance in the 3 day old offspring.

However, in 21-day-old offspring only the genotypes containing tolerant alleles at both QTL

differ from sensitive. This suggests QTL-3d may determine germ cell maintenance in the larval,

pupal and early adult stages, but QTL-21d may be additionally required to maintain tolerance in

aging females. The presence of two tolerance QTL is further supported by the phenotypic

classes we detected among founder alleles (B1-B8) for each of the QTL peaks (**Figure 1e**). For

QTL-21d, both B2 and B6 founder alleles are sensitive and greatly increase dysgenic ovarian

atrophy, while all other founder alleles are tolerant. By contrast for QTL-3d, only the B6 founder

allele is associated with increased sensitivity.

We next sought to determine whether reduced ovarian atrophy in tolerant alleles truly

increases fitness by restoring fertility, or merely allows for the production of inviable gametes.

To this end, we generated isogenic lines that carry either sensitive (B6) or tolerant (B8) alleles

at both QTL loci in an otherwise identical genetic background (**Supplemental figure S5**).

Consistent with our QTL mapping, tolerant alleles display less F1 ovarian atrophy (24-31%) than

sensitive strains when crossed with Harwich males (**Figure 1f, Supplemental table S17**).

Furthermore, fertility rates are higher than sensitive alleles (13-29)%, suggesting they are

beneficial in dysgenic females (**Figure 1g, Supplemental table S18**). Finally, while tolerant

females produce few offspring, offspring counts were significantly higher for tolerant females

from one isogenic stock when compared to sensitive (**Figure 1h).**

Figure 1: QTL mapping of variation in *P*-element tolerance. **a)** Crossing scheme to phenotype the variation in tolerance to *P*-elements among the RILs by screening for ovarian atrophy in 3 and 21 day-old dysgenic F1 females. Representative images of atrophied and non-atrophied ovaires are from Kelleher *et al.* [14] **b)** The log of odds (LOD) plot for QTL mapping of germline tolerance using 3 day-old (orange) and 21 day-old (blue) F1 females. The dotted line is the LOD threshold and x-axis represents the chromosomal positions. **c)** Zoomed-in figure of QTL mapping from 3 days (orange) and 21 days (blue). The colored boxes show the genomic interval that likely contains the causative genetic variant of each QTL, based on a Δ2LOD drop from the peak position [25]. The pairs of dotted lines indicate the peak Δ2LOD scores that determines the interval. The solid horizontal line is the LOD significance threshold based on 1,000 permutations of the phenotype data. **d)** Cytological map depicting the interval of the two QTL peaks [26,27]. **e)** Graph showing F1 atrophy (y-axis) associated with each of the eight founder alleles (x-axis) at the QTL peaks. All the QTL peaks show 2 phenotypic classes: sensitive (light green) and tolerant (dark green). **(f-g)** Percentage of **(f)** ovarian atrophy and **(g)**

sterility among dysgenic female offspring from crosses between Harwich males and isogenic females carrying sensitive (B6) and tolerant (B8) alleles. Tolerant strains show significant reduction in F1 atrophy (Tolerant_B8[1] vs. Sensitive_B6[1]: χ^2= 37.05, df = 1, *p*-value = 1.15e-09; Tolerant_B8[1] vs. Sensitive_B6[2]: χ^2= 13.7, df = 1, *p*-value = 0.0002; Tolerant_B8[2] vs. Sensitive_B6[1]: X-squared = 37.85, df = 1, *p*-value = 7.63e-10; Tolerant_B8[2] vs. Sensitive_B6[2]: χ^2 = 14.14, df = 1, *p*-value = 0.0001) as well as F1 sterility (Tolerant_B8[1] vs. Sensitive_B6[3]: χ^2= 10.55, df = 1, *p*-value = 0.001; Tolerant_B8[1] vs. Sensitive_B6[2]: χ^2= 8.41, df = 1, *p*-value = 0.003; Tolerant_B8[2] vs. Sensitive_B6[3]: χ^2 = 4.4, df = 1, *p*-value = 0.03; Tolerant_B8[2] vs. Sensitive_B6[2]: χ^2 = 3.4, df = 1, *p*-value = 0.06) compared to the sensitive strains. Subscripts [1,2] and [3] denote isogenic lines that were independently generated. **h)** Number of F2 offspring produced by individual dysgenic F1 females from crosses between Harwich males and isogenic tolerant and sensitive females. The horizontal line indicates the mean. Tolerant_B8[1] strains show a significantly higher number of F2 offspring (Tolerant_B8[1] vs. Sensitive_B6[3]: Z = 2.83, *p*-value = 0.004; Tolerant_B8[1] vs. Sensitive_B6[2]: Z = 2.52, *p*-value = 0.012). Error bars in **e, f** and **g** represent the standard error. The data used to generate plot in panel **b**,**c**, and **e** are provided in **Supplemental table S3 and S4** and that used for plot in panel **f**, **g and h** are provided in **Supplemental table Supplemental figure S17** and **S18** respectively.

Analysis	LOD Score	Peak Position	Δ2LOD CI	BCI	% variation
3-day	15.2	2R:6,192,495	2L:20,710,000-2R:7,272,495	2L:20,820,000-2R:6,942,495	11.13
21-day	10.13	2L:19,420,000	2L:19,170,000-20,080,000	2L:19,010,000-20,000,000	9.78

Table 1: QTL positions for tolerance in 3 and 21-day old females. The peak position, Δ2LOD drop confidence interval (Δ2LOD CI), and the Bayesian Credible Interval (BCI) in dm6 [28] are provided for each analysis. The data used to identify the LOD peaks and intervals for 3 and 21-day old females can be found in **Supplemental table S3** and **S4**, respectively.

2. Sensitive and tolerant alleles may differ in DSB repair and heterochromatin formation.

Both the QTL regions contain large numbers of protein coding and non-coding RNA genes, piRNA clusters, and repeats, which could influence tolerance (**Figure 1d**). To better understand the differences between tolerant and sensitive genotypes, we compared their ovarian gene expression profiles by stranded total RNA-seq. To avoid the confounding effects of germline loss under dysgenic conditions, we focused on 3-5 day old RIL females, rather than their dysgenic offspring. To account for potential background effects, we examined three pairs of RILs that carried either a sensitive (B6) or tolerant (B4) QTL haplotype across the QTL region (dm6 2L:19,010,000-2R:7,272,495) in otherwise similar genetic backgrounds (shared 44-47% of

founder alleles outside the QTL). Principal component analysis (PCA) of read counts reveals two independent axes that resolve sensitive and tolerant gene expression profiles, which together account for 40% and 16% of variation (**Figure 2a, Supplemental table S14**). One biological replicate of RIL 21188 (tolerant) was an outlier, which we excluded from our downstream analysis of differentially expressed genes.

We found a total of 530 genes differentially expressed between sensitive and tolerant genotypes (Benjamini-Hochberg adjusted *p*-value <=0.05, fold-change > 1.5; **Supplemental table S5**). The most significantly enriched GO term among genes upregulated in tolerant ovaries is chorion assembly (Bonferroni corrected *P* value <0.01, **Figure 2b, Supplemental table S7: full report**). Indeed, all of the major chorion genes are significantly upregulated in the tolerant ovaries (**Figure 2c**, [29,30]). It is unlikely that chorion synthesis promotes tolerance because chorion synthesis occurs in late-stage oocytes [stages 10B-14, 31], whereas atrophy results from the loss of larval PGCs and pre-meiotic adult cysts (GSCs) [15–17,19]. However, chorion genes reside in clusters that undergo multiple rounds of gene amplification [32,33], generating abundant DSBs at the boundaries of the amplified region that need to be repaired to permit transcription [34]. Therefore, upregulation of chorion genes in tolerant genotypes could indicate more efficient DSB repair.

Genes upregulated in the sensitive genotypes are enriched for functions in chromatin assembly and transcription, cell division, and translation. However, a careful inspection of genes underlying these enriched terms reveals that with the exception of translation, they are majorly explained by the increased expression of replication-dependent (RD) histone gene copies (**Figure 2d**). Notably, the expression of both histone and chorion genes are increased in late oogenesis [35–38], meaning that their inverted differential expression between sensitive and tolerant genotypes cannot be explained by differential abundance of late stage oocytes. Furthermore, histone upregulation may reduce tolerance to *P*-element activity, since overexpression of RD histones is associated with increased sensitivity to DNA damage [39–43], and excess Histones are reported to compete with DNA repair proteins for binding to damage sites [40].

The *D. melanogaster* histone gene cluster is located in the pericentromeric region of QTL-3d and consists of ~100 copies of a 5-kb cluster containing each of the 5 RD histones (*his1, his2A, his2B, his3 and his4)*. However, the differential regulation of histones is unlikely to reflect the presence of a *cis*-regulatory variant within the QTL, since the histone gene cluster exhibits coordinated and dosage compensated regulation in a unique nuclear body called the histone locus body (HLB, [44]). We therefore postulate that sensitive and tolerant alleles may differ in heterochromatin formation, since many negative regulators of histone gene transcription are also suppressors of position effect variegation [43,45]. In support of this model, sensitive (B6) genotypes exhibit increased expression of pericentromeric genes, as well as genes on the heterochromatic 4th chromosome (**Figure 2e**). We also discovered increased expression of

pericentromeric genes associated with the B6 haplotype in a previously published microarray

dataset from head tissue ([46] **Supplemental figure S1**), suggesting B6 is unusual among the

founder alleles in exhibiting reduced heterochromatin formation.

Figure 2: Tolerance is associated with increased chorion gene expression, whereas

sensitivity is associated with increased expression of replication-dependent histones. a)

PCA analysis of gene expression data for pairs of S/sensitive (B6) and T/ tolerant (B4) RILs.

Members of the same RIL pair with otherwise similar genetic backgrounds are represented by

the same shape. **b)** GO terms enriched among genes upregulated in tolerant and sensitive

genotypes. **c)** Log2 fold increase in expression in tolerant genotypes for chorion genes residing

in the four amplicons (*Drosophila* Amplicons in Follicle Cells, DAFCs) as well as outside

amplicons [29,30]. **d)** Log2 fold increase in RD histone expression in sensitive genotypes. **e)**.

Probability density plot of log2 fold change values for all euchromatic (blue), pericentromeric

(red), telomeric (green) genes and 4th chromosome (gray) between strains carrying sensitive

and tolerant alleles. The mean of each distribution is represented by a dotted line. Sensitive

genotypes display significantly higher expression of pericentromeric genes (two-sample t-test,

t_{141}= -9.32, *p*-value = 2.335e-16) and 4th chromosome genes (two-sample t-test, t_{53} = -4.56, *p*-
value = 3.014e-05) when compared to euchromatic genes. For **e**) the x-axis boundaries were
confined from (-1.5 to 2) for a better visualization. The pericentromere-euchromatin boundaries
were drawn from [28,47] and subtelomeric-euchromatin boundary coordinates from [48–50]. The
data represented in panel **a** is provided in **Supplemental table S14** and plot in panel **c, d,** and **e**
in **Supplemental table S5**).

3. Sensitive alleles are associated with radiation sensitivity

Our gene expression data suggest that sensitive and tolerant alleles may differ in their
capacity to repair DSBs. Mutations in repair genes are widely known to cause radiation
sensitivity [51–55]. We therefore compared the sensitivity of the tolerant and sensitive larvae
from isogenic lines to X-ray radiation.
After exploring a range of radiation doses, we found that doses above 10 Gy showed
high lethality, making it difficult to detect differences in radiation sensitivity between the
genotypes **(Supplemental table S19)**. Therefore, we compared the response of sensitive and
tolerant larvae to radiation doses of 0 Gy, 5 Gy and 10 Gy. We observed that tolerant genotypes
had significantly higher survival (53-58%) than the sensitive genotypes (25-30%) at 10 Gy
(**Figure 3**). These results are consistent with differences between sensitive and tolerant alleles
in DSB repair.
19

Figure 3. Tolerance is associated with enhanced DNA damage repair. Bar graph showing
the percentage of mock treated and irradiated (5 Gy and 10 Gy) larvae that survived to
adulthood for the tolerant, sensitive and the control genotypes. CS refers to Canton-S and
marker refers to the multiply marked stock *b cn* (#44229), which was used to generate isogenic
lines. The X-axis represents the different strains with the colors representing the type of
genotype. The Y-axis is the percentage of irradiated larvae that survived to adulthood. The
numbers in the brackets refer to the sample size. The number of larvae that survived and died
were compared between tolerance and sensitive genotypes . For 5 Gray irradiation,
Tolerant_B8[1] vs. Sensitive_B6[3]: χ^2= 15.66, df=1, *p*-value =0.0008; Tolerant_B8[2] vs.

3. piRNA clusters in QTL-3d exhibit differential activity that does not translate to TE deregulation.

Although the RIL mothers do not produce or transmit *P*-element-derived piRNAs (**Supplementary table S8**), the *D. melanogaster* genome harbors >100 resident TE families [56,57] that are also regulated by piRNAs [5]. Transposition of resident TEs could add to genotoxic stress triggered by *P*-element activity, thereby reducing tolerance. Furthermore, transposition rates of resident (non *P*-element) TEs differ between wild-type strains [58–60]. Two features of our data suggest potential differences in piRNA cluster activity between sensitive and tolerant alleles. First, QTL-3d contains numerous piRNA clusters, including major ovarian piRNA cluster *42AB,* which could differ in activity between sensitive and tolerant alleles (**Figure 1d)**. Second, differential heterochromatin formation between sensitive and tolerant genotypes could impact piRNA cluster expression, which is dependent upon the heterochromatic histone modification, histone 3 lysine 9 trimethylation (H3K9me3) [61,62]. We therefore evaluated whether tolerant and sensitive alleles differ in the activity of piRNA clusters by performing small RNA-seq on the same ovarian samples used for total RNA-seq.

A PCA of piRNA cluster expression reveals that sensitive and tolerant genotypes differ in the activity of some piRNA clusters, and are resolved by the second principal component, accounting for 22% variation in expression (**Figure 4a, Supplemental table S15**). However, the major piRNA clusters—including *42AB*—are not differentially expressed between sensitive and tolerant alleles, suggesting that the proposed reduction in heterochromatin formation in sensitive genotypes does not globally inhibit piRNA biogenesis (**Figure 4b, Supplemental Table S8**). Nevertheless, we discovered two small pericentromeric piRNA clusters located within QTL-3d that were active in tolerant genotypes but largely quiescent in sensitive genotypes (**Figure 4b, c and d; Supplemental figure S2 and S3; Supplemental table S16).** These piRNA clusters are largely composed of TE fragments that are relatively divergent from the consensus (65 to 95% sequence similarity; **Supplemental table S9**), or are most similar to a consensus TE from other (non-*melanogaster*) *Drosophila* species. Given that transpositionally active TEs are generally highly similar to the consensus sequence [63], and piRNA silencing is disrupted by mismatches between the piRNA and its target [64], this suggests that the differential activity of these two piRNA clusters is unlikely to impact the expression of transpositionally active TEs.

To directly address if differences in tolerance are related to resident TE regulation, we compared genome-wide resident TE expression between sensitive and tolerant genotypes in our RNA-seq data. None of the TE families represented in the QTL-3d piRNA clusters were upregulated in sensitive genotypes (**Figure 4e, Supplemental table S10**). Furthermore, while some TE families are differentially expressed, there is no systematic increase in TE activity in

1   the sensitive genotypes. Rather, more TE families are upregulated in tolerant genotypes (13
2   TEs) when compared to sensitive (4 TEs) genotypes. Therefore, despite the conspicuous
3   position of QTL-3d surrounding piRNA producing-regions, as well as evidence for differential
4   heterochromatin formation that could impact piRNA biogenesis (**Figure 2b and e**), we find no
5   evidence that tolerance is determined by resident TE silencing.

6
7   **Figure 4: Tolerance is not determined by differential activity of piRNA cluster or TE**
8   **deregulation. a**) PCA analysis for piRNA cluster expression data of sensitive (S) and tolerant
9   (T) genotypes. Members of the same RIL pair are represented by the same shapes. **b)** Heat
10   map showing the expression of seven major piRNA clusters [5] and the two differentially

1. expressed QTL clusters in QTL-3d. RIL pairs are plotted adjacent to each other. **c and d)**
2. Uniquely mapping piRNAs within two differentially active QTL-3d piRNA clusters are compared
3. between sensitive (21183) and tolerant (21213) genotypes. Positive value indicates piRNAs
4. mapped to the sense strand of the reference genome and negative value indicates those from
5. the antisense strand. TE insertions in each cluster are presented according to family by different
6. colors; TE-others indicate the insertion was most similar to a consensus TE from a sibling
7. species of *D. melanogaster*. See **Supplemental figure S2-3** for cluster expression in the
8. remaining RIL pairs. For **b, c and d,** piRNA cluster expression levels are estimated by log2
9. scale transformed of reads per million mapped reads [log2(RPM+1)]. **e)** Genome-wide
10. differences in TE family expression between sensitive and tolerant genotypes (fold change =
11. 1.5, base mean >= 100, adjusted *p*-value <= 0.05), based on alignment to consensus
12. sequences. The data used to plot panel **a** is provided in **Supplemental table S15,** for panel **b** in
13. **Supplemental table S8**, for panel **c** and **d** in **Supplemental table S16 and S9**, and for panel **e**
14. in **Supplemental table S10**)
15.

4. Identifying candidate tolerance genes

17.   We next sought to identify candidate genes that explain the tolerance differences using
18. three criteria: 1) location within a QTL, 2) differential expression and 3) the presence of "in-
19. phase" single nucleotide polymorphisms (SNPs) (**Supplemental table S11, S12, and S13**). In-
20. phase SNPs are those where the genotypic differences between the founder alleles are
21. consistent with their tolerance phenotype class [65]**re 5a,** [65]) . Of 530 differentially expressed
22. genes (**Figure 5b**), 43 are within the QTL region, representing an approximately five-fold
23. enrichment in the QTL regions compared to the rest of the genome (*X-squared* = 255.54, *df* = 1,
24. *p*-value < 2.2e-16, **Figure 5b**). Ultimately, we identified 14 and 5 differentially expressed genes
25. that also carry in-phase SNPs within the QTL-3d and 21d, respectively (**Figure 5c and d**;
26. **Supplemental table 12**). Furthermore, we identified 37 genes in QTL-3d and 4 genes in QTL-
27. 21d containing in-phase non-synonymous SNPs, which may affect the function of the encoded
28. protein (Supplemental table S13). These genes represent the strongest candidates to contain
29. tolerant variants.

30.   We next scoured our list of candidate genes for those with known functions in
31. heterochromatin formation and DSB repair, whose differential function or regulation are
32. plausibly related to phenotypic differences associated with sensitive and tolerant alleles. Within
33. QTL-3d, *Nipped-A*—which contains a non-synonymous in-phase SNP—stood out as a member
34. of the Tat interacting protein 60 kD (TIP60) complex. The TIP60 complex has functions in DSB
35. repair and heterochromatin formation [66–70]: providing a clear connection to our gene
36. expression and radiation assays. The non-synonymous SNP that separates sensitive and
37. tolerant alleles of this gene are located in the HEAT2 domain, which is predicted to be essential
38. for protein-protein interaction [71–73]. Furthermore, two additional members/interactors of
39. TIP60 complex residing within QTL-3d (*yeti* and *dRSF-1*) and three members outside QTL

(*dom, E(Pc)* & *DMAP1;* **Supplemental table S6**) are differentially expressed between tolerant
and sensitive genotypes [67,74,75].
Within QTL-21d, we did not find any genes with function in heterochromatin formation or
DSB repair. However, the germline differentiation factor *brat* was exceptional in containing 14
in-phase SNPs in introns and downstream regions, and is upregulated in the tolerant genotypes
(**Supplemental table S5** and **S11**). In adult ovaries, Brat is excluded from GSCs, but is
expressed in CBs and promotes differentiation [76]. Because DNA damage blocks cystoblast
differentiation by suppressing *bam* translation [21]*, brat* could confer tolerance in older females
by helping cytoblasts escape arrest.
10
11

12
13 **Figure 5: Differential expression and in-phase SNPs identify candidate tolerance genes.**
14 **a)** Hypothetical in-phase and out of phase SNPs are shown. Sequences of each of the B
15 founder B strains are colored based on their phenotypic classification, either tolerant or sensitive

(**Figure 1e**). Bold letters indicate SNPs. **b)** The proportion of genes differentially expressed
(DEG) is compared inside and outside the QTL. The dotted line is the genome wide average. **c**
and d) Venn diagrams showing the overlap of differentially expressed genes (DEG) and genes
carrying in-phase SNPs for QTL-21d (c) and QTL-3d (d). The number outside the bracket
indicates all genes with in-phase SNPs, whereas the number within the brackets indicates the
genes carrying non-synonymous in-phase SNPs only. The data for differential expression of
genes for tolerant and sensitive genotypes is provided in **Supplemental table S5**. The data on
in-phase polymorphisms for each QTL peak are provided in **Supplemental table S11**. List of
candidate genes that have both in-phase polymorphisms and are differentially expressed, and
those having non-synonymous in-phase polymorphisms are provided in **Supplemental table**
S12 and **S13**, respectively.
12

6. Investigating the role of *brat* in tolerance.

To determine the impact of *brat* on tolerance, we examined the tolerance phenotypes of
a *brat* loss-of-function mutation *(brat^1)* and multiple deficiencies overlapping *brat*. The candidate
causative variants in *brat* that are proposed to influence tolerance are most likely heterozygous
in dysgenic hybrid offspring. We therefore evaluated the heterozygous effect of *brat^1* and
overlapping deficiencies by comparing the incidence of ovarian atrophy between mutant or
deficiency offspring to balancer siblings from dysgenic crosses (*brat/CyO* x Harwich).
In absence of dysgenesis, *brat* loss of function alleles impact oogenesis recessively [76].
However, we found that the *brat^1* heterozygotes showed a significantly higher frequency of
ovarian atrophy (68.6%) than their balancer control siblings (37.5%) (**Figure 6, Supplemental**
table S20). Furthermore, two out of three deficiency stocks with deletions overlapping *brat*
increased ovarian atrophy similarly to the *brat^1* mutant, suggesting that this phenotype is not an
effect of the 2nd chromosome of the *brat^1* mutant line (**Figure 6, Supplemental table S20**). The
deficiency line (Df(2L)*brat* [ED1231]) that shows no change in the incidences of ovarian atrophy
may carry deletions in genes with opposing function to that of *brat,* or suppressors elsewhere in
the genome. Our results suggest that *brat* activity increases fertility in dysgenic females, which
is consistent with our observation that tolerant alleles exhibit increased *brat* expression
(**Supplemental table S5)**. Notably, the fertility effects of *brat* were observed in 3 day-old
offspring, as attempts to look at older females (21 day-olds) were unsuccessful due to a high
mortality rate.
33

1
2
Figure 6. **Loss-of-function mutation of *brat* increases severity of hybrid dysgenesis.** The
percentage of F1 ovarian atrophy is compared between control balancer siblings CyO/+,
heterozygous *brat¹* mutants and heterozygous deficiency lines Df(2L)*brat*. *brat¹* mutant: χ^2=
13.55, df=1, *p*-value =0.0002. Df(2L)*brat* [Exel8040]: χ^2= 14.78, df=1, *p*-value =0.0001.
Df(2L)*brat* [ED1231]: χ^2= 0.06, df=1, *p*-value =0.8. Df(2L)*brat* [ED1200]: χ^2= 3.66, df=1, *p*-value
=0.05. The underlying data are provided in **Supplemental table S20**.
9
10

1
2 **Figure 7. Hypothesized mechanisms of TE tolerance in young and old females.** a) We
3 propose that in larval and young-adult females, germline tolerance to *P*-elements may be
4 determined by enhanced DSB repair through increased TIP60 activity. b) In aging dysgenic
5 females, *brat* may determine tolerance by promoting differentiation of arrested cytoblasts, thus
6 aiding in their escape from the cell-cycle arrest imposed by *P*-element mediated DNA damage.

Discussion

Although small RNA mediated TE regulation is widely studied, little is known about
cellular and molecular mechanisms that confer tolerance to transposition. Here we uncovered
natural variation in tolerance to *P*-element DNA transposons, which is associated with two or
more loci proximal to the second chromosome centromere in *D. melanogaster*. We further
showed that tolerant and sensitive genotypes may differ in their ability to enact DSB repair,
potentially explaining their differential responses to *P*-element transposition. Finally, we
identified candidate genes in each QTL that potentially determine the phenotypic differences
between tolerant and sensitive alleles. Within QTL-3d, *Nipped-A* has a non-synonymous in-
phase SNP that could alter the activity of encoded protein. By contrast, *brat,* located in QTL-
21d, has in-phase SNPs in its intronic and downstream regions, and is upregulated in tolerant
genotypes.
19
Differences in DSB repair and TIP60 activity

We propose that in young females, tolerance is determined by the ability to repair DSBs
resulting from *P*-element activity in larval PGCs. In our ovarian RNA-seq data, we saw two
circumstantial indicators of differences in DSB repair in tolerant genotypes: increased chorion
gene expression and decreased histone gene expression (**Figure 2b-d**). Chorion gene
amplification is dependent upon DSB repair; thus, while we did not directly assay amplification,
their increased expression may indicate more efficient repair [34]. Conversely, in yeast, excess
histones inhibit DSB repair, potentially by competing with repair complexes for access to DNA
[40,77]. Increased histone expression in sensitive genotypes may therefore inhibit DSB repair.
Consistent with both of these observations, we observed that tolerant genotypes are
significantly more resilient to X-ray radiation (**Figure 3**), which is widely associated with
increased activity of DNA repair genes [51–55].
Enhanced repair in tolerant genotypes may be explained by increased activity of the
TIP60 complex: a conserved chromatin remodeling complex with functions in DSB repair [75,78]
and heterochromatin formation [66–69]. While two additional TIP60 components reside within
QTL-3d and are differentially expressed between sensitive and tolerant genotypes (*yeti* and
dRSF-1), *Nipped-A* is unique in containing a non-synonymous in-phase SNP. Consistent with a
deleterious effect, the amino acid change carried by the sensitive allele is quite rare in recently
sampled natural populations worldwide (collected after *P*-element invasion), occurring in only
four of 645 sequenced strains [79,80]. Interestingly, one of these strains (RAL799) was recently
examined for radiation sensitivity and found to be highly sensitive [81].
21   While the functional consequences of the non-synonymous SNP that separates tolerant
and sensitive *Nipped-A* alleles is not clear, the upregulation of four other TIP60 members in
tolerant genotypes (*dRSF-1, dom, E(Pc)* & *DMAP1*), together with evidence of enhanced
heterochromatin formation, suggests increased TIP60 activity (**Supplemental table S6**).
Increased TIP60 could directly facilitate DSB repair through its function in the exchange of
phosphorylated Histone 2AV at DSBs [75]. However, enhanced heterochromatin formation
resulting from TIP60 function could also facilitate DSB repair indirectly by reducing the
expression of histones. The latter is more speculative, because although the histone locus body
has a specialized chromatin state determined by multiple suppressors of variegation [43,45],
there is limited evidence that TIP60 regulates the histone locus body [82].
31
Germ cell differentiation and tolerance in adult females.
We identified *brat* as a promising candidate to explain natural variation in tolerance of
aging (21 day) females. *brat* resides in QTL-21d, contains 14 in-phase SNPs and was
upregulated in tolerant genotypes. Consistent with *brat* function promoting tolerance, we
observed that a *brat* loss-of-function mutation and multiple *brat* deficiencies are dominant
enhancers of dysgenic ovarian atrophy (**Figure 6b**), while their effects on oogenesis in non-
dysgenic germlines are recessive [83].
We propose that in aging adult females, *brat* could confer tolerance by promoting
cystoblast (CB) differentiation, thereby opposing the arrested differentiation that results from
DSBs [Figure 7b; 21]. Indeed, CB accumulation is observed when hybrid dysgenesis is induced
by temperature shift in adult females [15,20]. CB differentiation is delayed following DNA
damage by repressing the translation of *bam*, a key differentiation factor [21]. Brat acts to
promote the translation of *bam* by repressing the translation of Mad and Myc [83]. Interestingly

in the larval gonad, Myc activity is associated with retention of PGCs in dysgenic germlines [17],
further highlighting how tolerance mechanisms may differ over the course of development.
Our demonstration that *brat* promotes tolerance also presents an intriguing contrast to
our previous report that another germline differentiation factor, *bruno,* reduces tolerance [14].
Unlike *brat* alleles, *bruno* alleles and deficiencies are dominant suppressors of hybrid
dysgenesis. Similar to *brat, bruno* encodes an mRNA binding protein that promotes
differentiation of pre-meiotic cysts in the female germline, albeit at a later 4-cell stage [84,85].
bruno further differs from *brat* in acting independently of the Bam/Bgcn pathway that is
repressed in CBs after DSBs [21,86]. Collectively therefore our data speak to a careful
orchestration of germ cell differentiation that can facilitate germline persistence in the face of
DNA damage.
12
Conclusion:
Our work reveals that natural tolerance to transposition can arise throughout the lifecycle
of the fly, ensuring the maintenance of germline cells during development and the production of
gametes in adults. This contrasts our previous study of natural variation in the population A RILs
of the DSPR, which uncovered a single major effect QTL of differentiation factor, *bruno*, on
tolerance in both young and old females [14]. Furthermore, while DNA damage signaling is a
clear determinant of dysgenic germ cell loss [15,19,20], we for the first time provide evidence of
natural variation in DNA repair offsetting the damaging effects of transposition. Our observations
therefore point to multiple new mechanisms through which germlines could withstand the
genotoxic effects of unregulated transposition, which may respond to natural selection after new
TEs invade.
24
25
26

Methods

28
***Drosophila* Strains and Husbandry.** The recombinant inbred lines (RILs) were generously
provided by Stuart Macdonald. Harwich (#4264), *b cn* (#44229), *brat[1]* (#3988),
Df(2L)*brat*[Exel8040] (#7847), Df(2L)*brat* [ED1231] (#9174) and Df(2L)*brat* [ED1200] (#9173)
were obtained from the Bloomington *Drosophila* stock center. Canton-S was obtained from
Brigitte Dauwalder. All flies were maintained in standard cornmeal media.
Alleles of the second chromosome centromeric region, containing both QTL, were
extracted from three recombinant inbred lines carrying B6 QTL allele (#21076, #21218, #21156)
and two RILs carrying B8 QTL allele (#21077, #21154) into a common background by crossing
them to multiply marked stocks *b cn* (#44229). After 7 rounds of backcrossing followed by
inbreeding, the final isogenic lines (Sensitive_B6[1] , Sensitive_B6[2] , Sensitive_B6[3] and
Tolerant_B8[1] , Tolerant_B8[2]) were generated. The lines were made homozygous for the 2[nd]
chromosome by inbreeding and selecting for wild type phenotype. The genotype of the isogenic
lines were verified through PCR using five different primers within the two QTL.
chr2L:19383155-19383970: AACCCTTTTTCGCTGACAATAACA, ATTATCAGCAGGAGCCGGAAACTT;

chr2L:21333500-21334300: AAGTGAAGCTAACAACGTGACAAC,CGTTTGACCATCGCTTACAACTAA;

chr2R:2392800-2393600: AACAGGAGGTCGAAAGCCAAATA, ATGCAGAGTCATATTCTGGGTTGG;

chr2R:6203290-6204284: AATGGAGACCGTTGATTTTGGTAA,CTTTTCTGCGGCATCAGGTG;

chr2R:6058000-6059000: TGGCAATTGCAATCCTTTTGGTAT, ATAACACGAACTACGACCTTTCCA

Phenotyping. Phenotyping of ovarian atrophy was performed as described previously in

Kelleher *et al* [14]. Briefly, crosses between virgin RIL females and Harwich males were

transferred to fresh food every 3-5 days. Since crosses reared at a restrictive temperature (29

°C) result in complete gonadal atrophy in F1 offspring, we reared our crosses at a lower

permissive temperature (25 °C), which produces an intermediate phenotype that better reveals

the variation in severity of dysgenesis [12,14,15,87]. F1 offspring were maintained for 3 days or

21 days, at which point their ovaries were examined using a squash prep [87]. 21 day- old

females were transferred onto new food every 5 days as they aged to avoid bacterial growth.

Females who produced 1 or more chorionated egg chambers were scored as having non-

atrophied ovaries, and females producing 0 egg chambers were scored as having atrophied

ovaries.

Crosses and phenotyping were performed for 673 RILs across 22 experimental blocks

for 3 day-old F1 females, and 552 RILs across 18 experimental blocks for 21 day-old F1

females. If fewer than 21 F1 offspring were phenotyped for the same cross, it was discarded

and repeated if possible. In total, we phenotyped >20 3-day old and 21 day-old F1 female

offspring for 595 RILs and 456 RILs, respectively.

QTL mapping. QTL mapping was performed as described in Kelleher *et al.* [14]. Briefly, for

each developmental time point, we modeled the arcsine transformed proportion of F1 ovarian

atrophy as a function of two random effects: experimental block and undergraduate

experimenter. Regression models were fit using the lmer function from the lme4 package [88].

We then used the residuals as a response for QTL mapping with the DSPRqtl package [22] in R

3.02 [89]. The LOD significance threshold was determined from 1,000 permutations of the

observed data, and the confidence interval around each LOD peak was identified by a

difference of -2 from the LOD peak position (Δ2-LOD) [25], or from the Bayes Confidence

Interval [90]. For Δ2-LOD intervals, we took the conservative approach of determining the

longest contiguous interval where the LOD score was within 2 of the peak value. We further

calculated the broad sense heritability of ovarian atrophy as in Kelleher *et al. [14]*.

Estimation of Founder Phenotypes and QTL phasing. To estimate the phenotypic effect

associated with each founder allele at the QTL peak, we considered the distribution of

phenotypes from all RILs carrying the founder haplotype at the LOD peak position (genotype

probability >0.95%) [22]. QTL were then phased into allelic classes by identifying the minimal

number of partitions of founder haplotypes that describes phenotypic variation associated with

the QTL peak, as described previously [14,22].

Fertility Assays. Virgin female offspring from dysgenic crosses between isogenic lines carrying

tolerant_$B8_1$/$B8_2$(21077, 21154) and tolerant_$B8_2$/$B8_3$ (21218, 21156) alleles and Harwich

males were collected daily and individually placed in a vial containing two Canton-S males.

Females were allowed to mate for 5 days and were transferred to a new vial for another 5 days
after which the parents were discarded. The presence and total number of F2 individuals were
counted from the two vials.
4
5 **Identification of in-phase polymorphisms.** The SNP data of B founders that used to infer in-
6 phase SNPs is based on dm3 [22]. To identify in-phase SNPs we looked for alternate SNP
7 alleles that match the predicted phenotypic class for each of the QTL peaks. For QTL-21d we
8 used the criteria: sensitive class (B2, B6) and the tolerant class (B1, B3, B4, B7, B8), whereas
9 for QTL-3d: sensitive class (B6) and the tolerant class (B1, B2, B3, B4, B7, B8).
10
Selection of paired RILs with alternate QTL alleles. We identified background matched RILs
containing either the B6 ("sensitive") or B4 ("tolerant") haplotypes from the start position of the
QTL-21d confidence interval (2L: 19,010,000) to the end position of QTL-3d confidence interval
(2R: 6,942,495) ($P > 0.9$), based on their published HMM genotypes [22]. For all possible RIL
pairs (B6 and B4), we then calculated the number of 10 Kb genomic windows in which they
carried the same RIL haplotype ($P > 0.9$). We selected three pairs of RILs, which carry the
same founder genotype for 47% (21213 & 21183), 46% (21147 & 21346) and 44% (21291 &
21188) of genomic windows outside of the QTL.
19
Small RNA-seq and total RNA-seq. RILs were maintained at 25°C, and three biological
replicates of 20 ovaries were dissected from 3-5 day old females. Ovaries were homogenized
in TRIzol and stored at -80°C until RNA extraction. 50 μg of total RNA from each of 18 biological
samples (3 biological replicates x 3 pairs) was size fractionated in a 15% denaturing
polyacrylamide gel and the 18-30 nt band was excised. 2S-depleted small RNA libraries for
Illumina sequencing were then constructed according to the method of Wickersheim and
Blumenstiel [91]. Ovarian small RNA libraries were published previously [SRP160954, 92].
Ribodepleted and stranded total RNA libraries were generated from the same ovarian samples
using NuGen total RNA kit (TECAN). All 18 small RNA and total RNA libraries were sequenced
on an Illumina Nextseq 500 at the University of Houston Seq-N-Edit Core, and are deposited in
the NCBI BioProject PRJNA490147.
31
Small-RNA analysis. Sequenced small RNAs were separated based on size into
miRNAs/siRNAs (18-22nt) and piRNAs (23-30nt) [11]. Reads corresponding to contaminating
rRNAs, including 2S-rRNA, were removed from each library by aligning to annotated transcripts
from flybase [93]. To determine the piRNA cluster activity we first uniquely aligned the piRNAs
to reference genome (dm6 [28]) using Bowtie1 (-v 1 -m 1) [94]. We then used a customized perl
script (https://github.com/JLama75/piRNA-cluster-Coverage-script) to count reads that
mapped to a set of previously annotated piRNA clusters from the same genotypes (497 piRNA
clusters, [95]). Read counts normalized to total mapped microRNAs for each library were used
to infer differential expression using DESeq2 [96]. Sliding window estimates of piRNA
abundance (**Figure 2c and d)** were calculated using bedtools genomecov [97], normalizing the
read counts to total mapped miRNA reads.
43

Total RNA analysis. Residual ribosomal RNAs (rRNAs) were identified in ribodepleted libraries based on alignment to annotated rRNAs from flybase [93], and excluded from further analysis. Retained reads aligned to the library of consensus satellite and TE sequences from repbase [98], plus additional satellite consensus sequences from Larracuente [99]. For TE expression, the total reads mapped to TE sequences were counted using unix commands (uniq -c). Remaining reads that failed to map were aligned to *D. melanogaster* transcriptome (dm6/BDGP6) using Kallisto with default parameters [100]. Differentially expressed TEs and genes were identified from a combined analysis in DESeq2 [96]. Genes and TEs with base mean >= 100, Adjusted *P*-value <= 0.05 and whose expression pattern differed (fold change >= 1.5) were considered differentially expressed between the B6 and B4 QTL haplotype.

Radiation Sensitivity. Third instar larvae were either mock treated or irradiated in a Rad Source RS 1800 X-ray machine set at 12.5 mA and 160 kV. To obtain 3rd instar larvae, embryos were collected for 24 hr and aged for 5 days at 25 degree Celsius. The food vials containing larvae were then X-ray irradiated at doses from 5-80 Gray after which an optimal dose that clearly depicts the phenotypic difference was selected. Survival to adulthood was determined by scoring the number of empty and full pupal cases at 10 days after radiation.

References

1. Chuong EB, Elde NC, Feschotte C. Regulatory activities of transposable elements: from conflicts to benefits. Nat Rev Genet. 2017;18: 71–86.

2. Roy BA, Kirchner JW. Evolutionary dynamics of pathogen resistance and tolerance. Evolution. 2000;54: 51–63.

3. Mauricio R. Natural selection and the joint evolution of toleranceand resistance as plant defenses. Evolutionary Ecology. 2000. pp. 491–507. doi:10.1023/a:1010909829269

4. Råberg L. How to live with the enemy: understanding tolerance to parasites. PLoS Biol. 2014;12: e1001989.

5. Brennecke J, Aravin AA, Stark A, Dus M, Kellis M, Sachidanandam R, et al. Discrete small RNA-generating loci as master regulators of transposon activity in Drosophila. Cell. 2007;128: 1089–1103.

6. Nishida KM, Saito K, Mori T, Kawamura Y, Nagami-Okada T, Inagaki S, et al. Gene silencing mechanisms mediated by Aubergine piRNA complexes in Drosophila male gonad. RNA. 2007;13: 1911–1922.

7. Malone CD, Hannon GJ. Small RNAs as guardians of the genome. Cell. 2009;136: 656–668.

8. Kidwell, M.G., Frydryck, T., and Novy, J.B. The hybrid dysgenesis potential of Drosophila melanogaster strains of diverse temporal and geographical natural origins. Drosophila Information Service. 1983;59: 59–63.

9. Kidwell MG. Evolution of hybrid dysgenesis determinants in Drosophila melanogaster. Proc Natl Acad Sci U S A. 1983;80: 1655–1659.

10. Anxolabéhère D, Kidwell MG, Periquet G. Molecular characteristics of diverse populations are consistent with the hypothesis of a recent invasion of Drosophila melanogaster by mobile P elements. Mol Biol Evol. 1988;5: 252–269.

11. Brennecke J, Malone CD, Aravin AA, Sachidanandam R, Stark A, Hannon GJ. An epigenetic role for maternally inherited piRNAs in transposon silencing. Science. 2008;322: 1387–1392.

12. Kidwell MG, Kidwell JF, Sved JA. Hybrid Dysgenesis in DROSOPHILA MELANOGASTER: A Syndrome of Aberrant Traits Including Mutation, Sterility and Male Recombination. Genetics. 1977;86: 813–833.

13. Ignatenko OM, Zakharenko LP, Dorogova NV, Fedorova SA. P elements and the determinants of hybrid dysgenesis have different dynamics of propagation in

Drosophila melanogaster populations. Genetica. 2015;143: 751–759.

14. Kelleher ES, Jaweria J, Akoma U, Ortega L, Tang W. QTL mapping of natural variation reveals that the developmental regulator bruno reduces tolerance to P-element transposition in the Drosophila female germline. PLoS Biol. 2018;16: e2006040.

15. Dorogova NV, Bolobolova EU, Zakharenko LP. Cellular aspects of gonadal atrophy in Drosophila P-M hybrid dysgenesis. Dev Biol. 2017;424: 105–112.

16. Teixeira FK, Okuniewska M, Malone CD, Coux R-X, Rio DC, Lehmann R. piRNA-mediated regulation of transposon alternative splicing in the soma and germ line. Nature. 2017;552: 268–272.

17. Ota R, Kobayashi S. Myc plays an important role in Drosophila P-M hybrid dysgenesis to eliminate germline cells with genetic damage. Commun Biol. 2020;3: 185.

18. Shim HJ, Lee E-M, Nguyen LD, Shim J, Song Y-H. High-dose irradiation induces cell cycle arrest, apoptosis, and developmental defects during Drosophila oogenesis. PLoS One. 2014;9: e89009.

19. Tasnim S, Kelleher ES. p53 is required for female germline stem cell maintenance in P-element hybrid dysgenesis. Dev Biol. 2018;434: 215–220.

20. Moon S, Cassani M, Lin YA, Wang L, Dou K, Zhang ZZ. A Robust Transposon-Endogenizing Response from Germline Stem Cells. Dev Cell. 2018;47: 660–671.e3.

21. Ma X, Han Y, Song X, Do T, Yang Z, Ni J, et al. DNA damage-induced Lok/CHK2 activation compromises germline stem cell self-renewal and lineage differentiation. Development. 2016;143: 4312–4323.

22. King EG, Merkes CM, McNeil CL, Hoofer SR, Sen S, Broman KW, et al. Genetic dissection of a model complex trait using the Drosophila Synthetic Population Resource. Genome Res. 2012;22: 1558–1566.

23. Schaefer RE, Kidwell MG, Fausto-Sterling A. Hybrid Dysgenesis in DROSOPHILA MELANOGASTER: Morphological and Cytological Studies of Ovarian Dysgenesis. Genetics. 1979;92: 1141–1152.

24. Khurana JS, Wang J, Xu J, Koppetsch BS, Thomson TC, Nowosielska A, et al. Adaptation to P element transposon invasion in Drosophila melanogaster. Cell. 2011;147: 1551–1563.

25. Lander ES, Botstein D. Mapping mendelian factors underlying quantitative traits using RFLP linkage maps. Genetics. 1989;121: 185–199.

26. Bridges CB. SALIVARY CHROMOSOME MAPS: With a Key to the Banding of the Chromosomes of Drosophila Melanogaster. J Hered. 1935;26: 60–64.

27. Bridges PN. A NEW MAP OF THE SALIVARY GLAND 2L-CHROMOSOME: of Drosophila Melanogaster. J Hered. 1942;33: 403–408.

28. Hoskins RA, Carlson JW, Wan KH, Park S, Mendez I, Galle SE, et al. The Release 6 reference sequence of the Drosophila melanogaster genome. Genome Res. 2015;25: 445.

29. Tootle TL, Williams D, Hubb A, Frederick R, Spradling A. Drosophila Eggshell Production: Identification of New Genes and Coordination by Pxt. PLoS One. 2011;6: e19943.

30. Kim JC, Nordman J, Xie F, Kashevsky H, Eng T, Li S, et al. Integrative analysis of gene amplification in Drosophila follicle cells: parameters of origin activation and repression. Genes Dev. 2011;25: 1384.

31. Waring GL. Morphogenesis of the eggshell in Drosophila. Int Rev Cytol. 2000;198. doi:10.1016/s0074-7696(00)98003-3

32. Spradling AC. The organization and amplification of two chromosomal domains containing drosophila chorion genes. Cell. 1981;27: 193–201.

33. Claycomb JM, Benasutti M, Bosco G, Fenger DD, Orr-Weaver TL. Gene Amplification as a Developmental Strategy: Isolation of Two Developmental Amplicons in Drosophila. Dev Cell. 2004;6: 145–155.

34. Alexander JL, Barrasa MI, Orr-Weaver TL. Replication Fork Progression during Re-replication Requires the DNA Damage Checkpoint and Double-Strand Break Repair. Curr Biol. 2015;25: 1654–1660.

35. Potter-Birriel J, Gonsalvez GB, Marzluff WF. A region of Drosophila SLBP distinct from the histone pre-mRNA binding and processing domains is essential for deposition of histone mRNA in the oocyte. Cold Spring Harbor Laboratory. 2020. p. 2020.04.16.030577. doi:10.1101/2020.04.16.030577

36. Ruddell A, Jacobs-Lorena M. Biphasic pattern of histone gene expression during Drosophila oogenesis. Proc Natl Acad Sci U S A. 1985;82: 3316–3319.

37. Ambrosio L, Schedl P. Two discrete modes of histone gene expression during oogenesis in Drosophila melanogaster. Dev Biol. 1985;111: 220–231.

38. Waring GL. Morphogenesis of the eggshells in Drosophila. International Review of Cytology. Academic Press; 2000. pp. 67–108.

39. Gunjan A, Verreault A. A Rad53 kinase-dependent surveillance mechanism that regulates histone protein levels in S. cerevisiae. Cell. 2003;115. doi:10.1016/s0092-

8674(03)00896-1

40. Liang D, Burkhart SL, Singh RK, Kabbaj M-HM, Gunjan A. Histone dosage
regulates DNA damage sensitivity in a checkpoint-independent manner by the
homologous recombination pathway. Nucleic Acids Res. 2012;40: 9604.

41. Murga M, Jaco I, Fan Y, Soria R, Martinez-Pastor B, Cuadrado M, et al. Global
chromatin compaction limits the strength of the DNA damage response. J Cell Biol.
2007;178: 1101.

42. Landais S, D'Alterio C, Jones DL. Persistent replicative stress alters polycomb
phenotypes and tissue homeostasis in Drosophila melanogaster. Cell Rep. 2014;7.
doi:10.1016/j.celrep.2014.03.042

43. Ozawa N, Furuhashi H, Masuko K, Numao E, Makino T, Yano T, et al. Organ
identity specification factor WGE localizes to the histone locus body and regulates
histone expression to ensure genomic stability in Drosophila. Genes Cells.
2016;21. doi:10.1111/gtc.12354

44. McKay DJ, Klusza S, Penke TJR, Meers MP, Curry KP, McDaniel SL, et al.
Interrogating the function of metazoan histones using engineered gene clusters.
Dev Cell. 2015;32: 373–386.

45. Ner SS, Harrington MJ, Grigliatti TA. A Role for the Drosophila SU(VAR)3-9 Protein
in Chromatin Organization at the Histone Gene Cluster and in Suppression of
Position-Effect Variegation. Genetics. 2002;162: 1763–1774.

46. King EG, Sanderson BJ, McNeil CL, Long AD, Macdonald SJ. Genetic dissection of
the Drosophila melanogaster female head transcriptome reveals widespread allelic
heterogeneity. PLoS Genet. 2014;10: e1004322.

47. Riddle NC, Minoda A, Kharchenko PV, Alekseyenko AA, Schwartz YB, Tolstorukov
MY, et al. Plasticity in patterns of histone modifications and chromosomal proteins
in Drosophila heterochromatin. Genome Res. 2011;21: 147.

48. Karpen GH, Spradling AC. Analysis of Subtelomeric Heterochromatin in the
Drosophila Minichromosome Dp1187 by Single P Element Insertional Mutagenesis.
Genetics. 1992;132: 737.

49. Yin H, Lin H. An epigenetic activation role of Piwi and a Piwi-associated piRNA in
Drosophila melanogaster. Nature. 2007;450: 304–308.

50. Walter MF, Jang C, Kasravi B, Donath J, Mechler BM, Mason JM, et al. DNA
organization and polymorphism of a wild-type Drosophila telomere region.
Chromosoma. 1995;104: 229–241.

51. Koval L, Proshkina E, Shaposhnikov M, Moskalev A. The role of DNA repair genes
in radiation-induced adaptive response in Drosophila melanogaster is differential

and conditional. Biogerontology. 2019;21: 45–56.

52. Uri A, Martha K, Veronika B-I, Anna B, Trudi S. An essential role for Drosophila
hus1 in somatic and meiotic DNA damage responses. J Cell Sci. 2007;120: 1042.

53. Ruike T, Takeuchi R, Takata K-I, Oshige M, Kasai N, Shimanouchi K, et al.
Characterization of a second proliferating cell nuclear antigen (PCNA2) from
Drosophila melanogaster. FEBS J. 2006;273: 5062–5073.

54. Staeva-Vieira E, Yoo S, Lehmann R. An essential role of DmRad51/SpnA in DNA
repair and meiotic checkpoint control. EMBO J. 2003;22: 5863–5874.

55. Sterpone S, Cozzi R. Influence of XRCC1 Genetic Polymorphisms on Ionizing
Radiation-Induced DNA Damage and Repair. J Nucleic Acids. 2010;2010.

56. Kaminker JS, Bergman CM, Kronmiller B, Carlson J, Svirskas R, Patel S, et al. The
transposable elements of the Drosophila melanogaster euchromatin: a genomics
perspective. Genome Biol. 2002;3: 1–20.

57. Quesneville H, Bergman CM, Andrieu O, Autard D, Nouaud D, Ashburner M, et al.
Combined evidence annotation of transposable elements in genome sequences.
PLoS Comput Biol. 2005;1: 166–175.

58. Nuzhdin SV, Mackay TF. Direct determination of retrotransposon transposition
rates in Drosophila melanogaster. Genet Res. 1994;63: 139–144.

59. Nuzhdin SV, Mackay TF. The genomic rate of transposable element movement in
Drosophila melanogaster. Mol Biol Evol. 1995;12: 180–181.

60. Nuzhdin SV, Pasyukova EG, Morozova EA, Flavell AJ. Quantitative genetic
analysis of copia retrotransposon activity in inbred Drosophila melanogaster lines.
Genetics. 1998;150: 755–766.

61. Mohn F, Sienski G, Handler D, Brennecke J. The rhino-deadlock-cutoff complex
licenses noncanonical transcription of dual-strand piRNA clusters in Drosophila.
Cell. 2014;157: 1364–1379.

62. Le Thomas A, Stuwe E, Li S, Du J, Marinov G, Rozhkov N, et al.
Transgenerationally inherited piRNAs trigger piRNA biogenesis by changing the
chromatin of piRNA clusters and inducing precursor processing. Genes Dev.
2014;28: 1667–1680.

63. Bergman CM, Bensasson D. Recent LTR retrotransposon insertion contrasts with
waves of non-LTR insertion since speciation in Drosophila melanogaster. Proc Natl
Acad Sci U S A. 2007;104: 11340–11345.

64. Post C, Clark JP, Sytnikova YA, Chirn G-W, Lau NC. The capacity of target
silencing byDrosophilaPIWI and piRNAs. RNA. 2014. pp. 1977–1986.

doi:10.1261/rna.046300.114

65. Long AD, Macdonald SJ, King EG. Dissecting Complex Traits Using the Drosophila Synthetic Population Resource. Trends Genet. 2014;30: 488.

66. Qi D, Jin H, Lilja T, Mannervik M. Drosophila Reptin and Other TIP60 Complex Components Promote Generation of Silent Chromatin. Genetics. 2006;174: 241.

67. Hanai K, Furuhashi H, Yamamoto T, Akasaka K, Hirose S. RSF Governs Silent Chromatin Formation via Histone H2Av Replacement. PLoS Genet. 2008;4. doi:10.1371/journal.pgen.1000011

68. Sinclair DA, Clegg NJ, Antonchuk J, Milne TA, Stankunas K, Ruse C, et al. Enhancer of Polycomb is a suppressor of position-effect variegation in Drosophila melanogaster. Genetics. 1998;148: 211–220.

69. Ruhf ML, Braun A, Papoulas O, Tamkun JW, Randsholt N, Meister M. The domino gene of Drosophila encodes novel members of the SWI2/SNF2 family of DNA-dependent ATPases, which contribute to the silencing of homeotic genes. Development. 2001;128: 1429–1441.

70. Sun Y, Jiang X, Chen S, Fernandes N, Price BD. A role for the Tip60 histone acetyltransferase in the acetylation and activation of ATM. Proc Natl Acad Sci U S A. 2005;102: 13182.

71. Knutson BA, Hahn S. Domains of Tra1 important for activator recruitment and transcription coactivator functions of SAGA and NuA4 complexes. Mol Cell Biol. 2011;31: 818–831.

72. Perry P, Sauer S, Billon N, Richardson WD, Spivakov M, Warnes G, et al. A dynamic switch in the replication timing of key regulator genes in embryonic stem cells upon neural induction. Cell Cycle. 2004;3: 1645–1650.

73. Murr R, Vaissière T, Sawan C, Shukla V, Herceg Z. Orchestration of chromatin-based processes: mind the TRRAP. Oncogene. 2007;26: 5358–5372.

74. Messina G, Damia E, Fanti L, Atterrato MT, Celauro E, Mariotti FR, et al. Yeti, an essential Drosophila melanogaster gene, encodes a protein required for chromatin organization. J Cell Sci. 2014;127: 2577–2588.

75. Kusch T, Florens L, Macdonald WH, Swanson SK, Glaser RL, Yates JR, et al. Acetylation by Tip60 is required for selective histone variant exchange at DNA lesions. Science. 2004;306. doi:10.1126/science.1103455

76. Harris RE, Pargett M, Sutcliffe C, Umulis D, Ashe HL. Brat Promotes Stem Cell Differentiation via Control of a Bistable Switch that Restricts BMP Signaling. Developmental Cell. 2011. pp. 72–83. doi:10.1016/j.devcel.2010.11.019

77. Kumar K, Moirangthem R, Kaur R. Histone H4 dosage modulates DNA damage response in the pathogenic yeast Candida glabrata via homologous recombination pathway. PLoS Genet. 2020;16: e1008620.

78. Sun Y, Jiang X, Xu Y, Ayrapetov MK, Moreau LA, Whetstine JR, et al. Histone H3 methylation links DNA damage detection to activation of the tumour suppressor Tip60. Nat Cell Biol. 2009;11: 1376–1382.

79. Lack JB, Cardeno CM, Crepeau MW, Taylor W, Corbett-Detig RB, Stevens KA, et al. The Drosophila genome nexus: a population genomic resource of 623 Drosophila melanogaster genomes, including 197 from a single ancestral range population. Genetics. 2015;199. doi:10.1534/genetics.115.174664

80. Lack JB, Lange JD, Tang AD, Corbett-Detig RB, Pool JE. A Thousand Fly Genomes: An Expanded Drosophila Genome Nexus. Mol Biol Evol. 2016;33: 3308.

81. Vaisnav M, Xing C, Ku H-C, Hwang D, Stojadinovic S, Pertsemlidis A, et al. Genome-Wide Association Analysis of Radiation Resistance in Drosophila melanogaster. PLoS One. 2014;9. doi:10.1371/journal.pone.0104858

82. Messina G, Damia E, Fanti L, Atterrato MT, Celauro E, Mariotti FR, et al. Yeti, a Drosophila melanogaster essential gene, encodes a protein required for chromatin organization. J Cell Sci. 2014. Available: https://www.academia.edu/download/46330185/jcs.150243.full.pdf

83. Harris RE, Pargett M, Sutcliffe C, Umulis D, Ashe HL. Brat promotes stem cell differentiation via control of a bistable switch that restricts BMP signaling. Dev Cell. 2011;20: 72–83.

84. Parisi MJ, Deng W, Wang Z, Lin H. The arrest gene is required for germline cyst formation during Drosophila oogenesis. Genesis. 2001;29: 196–209.

85. Wang Z, Lin H. Sex-lethal is a target of Bruno-mediated translational repression in promoting the differentiation of stem cell progeny during Drosophila oogenesis. Dev Biol. 2007;302: 160–168.

86. Xin T, Xuan T, Tan J, Li M, Zhao G, Li M. The Drosophila putative histone acetyltransferase Enok maintains female germline stem cells through regulating Bruno and the niche. Dev Biol. 2013;384: 1–12.

87. Srivastav SP, Kelleher ES. Paternal Induction of Hybrid Dysgenesis in *Drosophila melanogaster* Is Weakly Correlated with Both *P* -Element and *hobo* Element Dosage. G3: Genes|Genomes|Genetics. 2017; g3.117.040634.

88. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4. J Stat Softw. 2015;67: 1–48.

89. Team, TRDC. The R Project for Statistical Computing. http://www.r-project.org.

2008 [cited 3 Mar 2021]. Available: https://ci.nii.ac.jp/naid/10027310073

90. Manichaikul A, Dupuis J, Sen S, Broman KW. Poor performance of bootstrap
confidence intervals for the location of a quantitative trait locus. Genetics.
2006;174: 481–489.

91. Wickersheim ML, Blumenstiel JP. Terminator oligo blocking efficiently eliminates
rRNA from *Drosophila* small RNA sequencing libraries. Biotechniques. 2013;55:
269–272.

92. Zhang S, Kelleher ES. piRNA-mediated silencing of an invading transposable
element evolves rapidly through abundant beneficial de novo mutations. bioRxiv.
2019. p. 611350. doi:10.1101/611350

93. Gramates LS, Marygold SJ, Santos G dos, Urbano J-M, Antonazzo G, Matthews
BB, et al. FlyBase at 25: looking to the future. Nucleic Acids Res. 2017;45: D663–
D671.

94. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. Genome Biol. 2009;10:
R25.

95. Zhang S, Pointer B, Kelleher ES. Rapid evolution of piRNA-mediated silencing of
an invading transposable element was driven by abundant de novo mutations.
Genome Res. 2020;30: 566–575.

96. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. 2014;15: 550.

97. Quinlan AR. BEDTools: The Swiss-Army Tool for Genome Feature Analysis. Curr
Protoc Bioinformatics. 2014;47: 11.12.1–34.

98. Bao W, Kojima KK, Kohany O. Repbase Update, a database of repetitive elements
in eukaryotic genomes. Mob DNA. 2015;6: 11.

99. Larracuente AM. The organization and evolution of the Responder satellite in
species of the Drosophila melanogaster group: dynamic evolution of a target of
meiotic drive. BMC Evol Biol. 2014;14: 233.

100.   Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq
quantification. Nat Biotechnol. 2016;34: 525–527.

Figure S1) Sensitivity is associated with increased expression of pericentromeric genes in the head. a) Mean expression of genes located in the pericentromere, euchromatin, telomere and the fourth chromosome from RILs carrying each of the eight B founder genotypes at the QTL-3d region. Error bars represent the standard deviation among mean expression levels of different genes. The sensitive/B6 (light green) shows high pericentromeric gene expression compared to the tolerant strains (dark green) (Anova; $F_{6,494}$=7.775, P<5.24e-08). The letters indicate significantly different expression levels based on Tukey-HSD comparisons between RILs with different founder alleles.

Figure S2) Expression profile of QTL piRNA clusters in a sensitive and tolerant NIL pair. The piRNA expression between sensitive and tolerant genotypes from 21188-21291 NIL pairs along the two QTL piRNA clusters: 2L:23,328,000-23,337,026 and 2L:23,222,004-23,246,024, respectively. Only uniquely mapping piRNAs are considered. The TE families at the top of each panel are represented by different colors. TE-others represent the repeat families coming from sibling species of D. melanogaster. Positive value indicates piRNAs mapped to the sense strand of the reference genome and negative value indicates those from the antisense strand. The piRNA cluster expression levels are estimated by log2 scale transformed of reads per million mapped reads [log2(RPM+1)].

Figure S3) Expression profile of QTL piRNA clusters in a sensitive and tolerant NIL pair. The piRNA expression between sensitive and tolerant genotypes from 21346-21147 NIL pairs along the two QTL piRNA clusters: 2L:23,328,000-23,337,026 and 2L:23,222,004-23,246,024, respectively. Only uniquely mapping piRNAs are considered. The TE families at the top of each figure are represented by different colors. TE-others represent the repeat families coming from sibling species of D. melanogaster. Positive value indicates piRNAs mapped to the sense strand of the reference genome and negative value indicates those from the antisense strand. The piRNA cluster expression levels are estimated by log2 scale transformed of reads per million mapped reads [log2(RPM+1)].

Figure S4. Tolerance among 3 and 21 day females based on QTL haplotype. Four haplotypes are compared, which comprise all possible combinations of tolerance alleles at 2 QTL. The allele at the 3 day QTL is indicated first and is represented by the color of the violin plot (light green = sensitive, dark green = tolerant). The allele at the 21 day QTL is indicated second and represented by the color of the points on the scatter plot. Y-axis is residual variation in F1 atrophy after accounting for student experimenter and block. Among 3 day old females, haplotypes containing different alleles for the 3 day old QTL are significantly different from each other (Tukey HSD P=0.016-0). However, haplotypes containing alternative QTL for the 21d only do not differ from each other (Tukey HSD P>0.74). This suggests phenotypic variation in 3 day old females is not influenced by their genotype at the 21 day QTL. In contrast, among 21 day old females tolerant alleles in both QTL loci are required to significantly increase tolerance above sensitive allele containing haplotypes (Tukey HSD P = 0.01-0).

marker

b   cn

b   cn

black body
bright red eyes

tolerant/sensitive

b+   cn+

b+   cn+

wild type phenotype

X

F1

recombination

b   cn

b+   cn+

X

marker

b   cn

b   cn

7 rounds of backcrossing

b+   cn

b   cn

bright red eye

b   cn+

b   cn

black body

b+   cn+

b   cn

wild type phenotype

b   cn

b   cn

black body
bright red eyes

Inbreeding

b+   cn+

Isogenic sensitive/tolerant lines