**A non-coding A-to-T Kozak site change related to the transmissibility of Alpha,**

**Delta and Omicron VOCs**

Jianing Yang[1,†], Guoqing Zhang[1,†], Dalang Yu[1,†], Ruifang Cao[1,†], Xiaoxian Wu[2],

Yunchao Ling[1], Yi-Hsuan Pan[4], Chunyan Yi[3], Xiaoyu Sun[3], Bing Sun[3], Yu Zhang[2],

Guo-Ping Zhao[1,2,5,*], Yixue Li[1,6,*], Haipeng Li[1,*]

[1]Bio-Med Big Data Center, CAS Key Laboratory of Computational Biology,

Shanghai Institute of Nutrition and Health, University of Chinese Academy of

Sciences, Chinese Academy of Sciences, Shanghai 200031, China.

[2]Key Laboratory of Synthetic Biology, CAS Center for Excellence in Molecular Plant

Sciences, Institute of Plant Physiology and Ecology, Chinese Academy of Sciences,

Shanghai 200032, China.

[3]Laboratory of Cell Biology, Shanghai Institute of Biochemistry and Cell Biology,

Center for Excellence in Molecular Cell Science, Chinese Academy of Sciences,

Shanghai 200031, China.

[4]Key Laboratory of Brain Functional Genomics of Ministry of Education, School of

Life Science, East China Normal University, Shanghai 200062, China.

[5]School of Life and Health Sciences, Hangzhou Institute for Advanced Study,

University of Chinese Academy of Sciences, Hangzhou, China.

[6]Bioland Laboratory (Guangzhou Regenerative Medicine and Health Guangdong

Laboratory), Guangzhou 510005, China.

[†]These authors contributed equally.

[*]Corresponding authors: gpzhao@sibs.ac.cn; yxli@sibs.ac.cn; lihaipeng@picb.ac.cn

**Abstract**

Three prevalent SARS-CoV-2 Variants of Concern (VOCs) were emerged and

caused epidemic waves. It is essential to uncover the key genetic changes that cause

the high transmissibility of VOCs. However, different viral mutations are generally

tightly linked so traditional population genetic methods may not reliably detect

beneficial mutation. In this study, we proposed a new pandemic-scale phylogenomic

approach to detect mutations crucial to transmissibility. We analyzed 3,646,973

high-quality SARS-CoV-2 genomic sequences and the epidemiology metadata. Based

on the sequential occurrence order of mutations and the instantaneously accelerated

furcation rate, the analysis revealed that two non-coding mutations at the position of

28271 (g.a28271-/t) might be crucial for the high transmissibility of Alpha, Delta and

Omicron VOCs. Both two mutations cause an A-to-T change at the core Kozak site of

the *N* gene. The analysis also revealed that the non-coding mutations (g.a28271-/t)

alone are unlikely to cause high viral transmissibility, indicating epistasis or

multilocus interaction in viral transmissibility. A convergent evolutionary analysis

revealed that g.a28271-/t, S:P681H/R and N:R203K/M occur independently in the

three-VOC lineages, suggesting a potential interaction among these mutations.

Therefore, this study unveils that non-synonymous and non-coding mutations could

affect the transmissibility synergistically.

**Keywords:** SARS-CoV-2; VOCs; non-coding mutation; Kozak sequence;

transmissibility; a28271-; a28271t

## Introduction

Three prevalent SARS-CoV-2 Variants of Concern (VOCs) were emerged in the last two years and draw the most attention for the epidemic waves that they caused (Fig. 1A). These VOCs were named as the Alpha, Delta and Omicron VOCs. The Alpha VOC, also known as SARS-CoV-2 lineage B.1.1.7, is a variant first detected in the UK [1] and has higher transmissibility than the preexisting variants in September 2020 [2]. Its high transmissibility remains similar across different age, sex and socioeconomic strata [3]. The Delta VOC was first identified in India in October 2020 [4] and has higher transmissibility [5-7]. The Omicron VOC was first identified in Southern Africa in late November 2021 [8]. Then the variant spreads more rapidly than the Delta variant [9] and has been detected in 76 countries globally in less than a month [10].

It is essential to uncover the key genetic changes that cause the high transmissibility of VOCs [11,12]. All the three VOCs carry a large number of mutations. Comparing with the reference genomic sequence of SARS-CoV-2 (GenBank accession number: NC_045512.2) [13], the sequences of the Alpha, Delta, Omicron BA.1 and Omicron BA.2 variants have 21, 27, 52 and 51 amino-acid change mutations, respectively [4] while two, three, one and two non-coding mutations can be found [14]. Moreover, different viral mutations are generally tightly linked since the virus lacks recombination occurred during the stage of meiosis in sexual species. Therefore, traditional population genetic methods, such as selective sweeps, may not reliably detect beneficial mutations [15-19] in SARS-CoV-2, and a novel approach is urgently needed.

In this study, we proposed a new pandemic-scale phylogenomic approach based

on the sequential occurrence order of mutations and the instantaneously accelerated

furcation rate to detect mutations crucial to transmissibility. Let us assume that the

high transmissibility of a VOC is due to the emergence of a beneficial haplotype with

multiple crucial mutations. Thus, the last (*i.e.*, the most recent) crucial mutation is

likely to increase the furcation rate in phylogenetic tree (Fig. 1B) because the VOC

with a beneficial haplotype emerges. To identify the crucial coding and non-coding

mutations for the three VOCs, we used the Coronavirus GenBrowser (CGB) [14] to

analyze a tip-dated tree with 3,646,973 high-quality SARS-CoV-2 genomic sequences

and the associated epidemiology metadata. The CGB is a free platform offering a

panoramic vision of the transmission and evolution of SARS-CoV-2. The analysis

revealed that two non-coding mutations at the position of 28,271 (g.a28271-/t) may be

crucial for the high transmissibility of the three VOCs. Both two mutations cause an

A-to-T change at the core Kozak site of the *N* gene. The analysis also revealed that the

non-coding mutations (g.a28271-/t) alone are not associated with high viral

transmissibility, indicating that the non-coding mutations may interact with other

coding mutations to increase the viral transmissibility.

**Methods**

**Data sources**

The annotated evolutionary tree and evolutionary network data were obtained

from the Coronavirus GenBrowser (CGB) [14] and VENAS [20]. All sequence data of

SARS-CoV-2 were obtained from the 2019nCoVR database [21,22], which is an integrated

resource based on Global Initiative on Sharing All Influenza Data (GISAID) [23,24],

National Center for Biotechnology Information (NCBI) GenBank [25], China National

GeneBank DataBase (CNGBdb) [26], the Genome Warehouse (GWH) [27], and the

National Microbiology Data Center (NMDC, https://nmdc.cn/). Two data versions of

CGB ("data.2021-03-06" and "data.2022-04-14") were used in this study, which

contains 400,051 and 3,777,753 genomic sequences, respectively.

**Furcation rate to evaluate virus transmissibility**

To evaluate virus transmissibility, we proposed to calculate furcation rate (FR)

of nodes in a considered phylogenetic clade (Fig. 1B). Furcation rate of the $i$-th node

($FR_i$) can be calculated using the following equation,

$FR_i = C_i/l_i,$

where $C_i$ is the number of child nodes and $l_i$ the branch length of the node with

time in units. Since $FR_i$ represents the furcation rate during the spanning time period

of the branch, the furcation rate at time $t$ ($FR_t$) can be calculated by averaging the

$FR_i$ of branches that cover the time point in the considered phylogenetic clade. The

estimated curve of $FR_t$ can be smoothed to reduce the fluctuation. In this study, it

was done by applying a sliding window with seven days in size.

**The effectiveness of mutations in improving viral transmissibility**

To evaluate the effectiveness of a mutation in improving transmissibility, we

tested whether two sibling clades with or without the mutation have similar viral

transmissibility (Fig. 1B). The null hypothesis is that the two sibling clades have the

same viral transmissibility. The genetic background of the two sibling clades has only

two-mutation difference in the presented example (Fig. 1B). The alternative

hypothesis is that the clade with the mutation (red) has higher viral transmissibility

than the sibling clade without the mutation. The binomial probability was applied to

test the null hypothesis and the test was one-tailed. Because the three VOCs occurred

at different time points, the analysis for Alpha was based on the data version

"data.2021-03-06" ($n = 400,051$) and the analysis of Delta and Omicron was based

on the data version "data.2022-04-14" ($n = 3,777,753$), where $n$ is the number of

viral strains. The two datasets were obtained from the CGB [14].

**Calculation of *Rt***

Considering that the sequenced strains were sampled from the infected patients,

the frequency of each variant at time $t$ can be estimated from the CGB datasets and

the number of new infections of a variant at time $t$ ($I_{V,t}$) can be approximated using

the following equation,

$I_{V,t} = f_{V,t} * I_{T,t}$ ,

where $I_{T,t}$ represents the total number of new infections at time $t$ and $f_{V,t}$ is the

frequency of the variant at time $t$.

To reduce the fluctuation caused by the small frequency of the Alpha-like

variant, we used a window size of 35 days to smooth its frequency curve before using

it to calculate $Rt$. The total number of new infections ($I_{T,t}$) in the United Kingdom

was downloaded from the UK Coronavirus Dashboard

(https://coronavirus.data.gov.uk/). Then the $Rt$ was calculated for the Alpha and

Alpha-like variants using the EpiEstim [28] with a mean ($\pm$SD) serial interval of 7.5$\pm$

3.4 days [29].

## Reappearance of g.a28271- and g.a28271t in the evolutionary tree

To examine the reappearance of mutations, we searched the non-coding

mutations in the CGB evolutionary tree [14]. We used the string "A28271-" to search

g.a28271- and "A28271T" to search g.a28271t. To present the reappearance patterns

of mutations, the data version "data.2021-04-14" ($n = 3,777,753$) of the CGB [14] was

used. This data was also used to examine the frequency trajectory of g.a28271-/t when

the Alpha, Delta and Omicron lineages were excluded.

## Identification of new canonical Alpha genomic sequence

The CGB was employed to identify a new canonical Alpha genomic sequence [30]

that carries the deletion g.a28271- and all other Alpha characteristic mutations (Fig.

2A, Supplementary Table S1). We first selected all the strains in the Alpha

(CGB84017.91425) clade that carries all the Alpha characteristic mutations including

g.a28271-. Then we filtered the strains by date and only kept the strains collected

before 1 November, 2020. Viral strains with extra mutations were ignored. Then the

sequence with accession EPI_ISL_629703, as the suggested new canonical Alpha

genomic sequence, is the first collected high-quality sequence without any extra

mutations after the deletion g.a28271- occurred (Supplementary Fig. S1).

**Results**

**Instantaneously accelerated furcation rate and high transmissibility**

It has been recognized that the phylogenetic tree is related to the transmission

tree and different transmission patterns can be reflected in phylogenetic tree shapes [31].

Thus furcation, as the process of lineage splitting [32], is associated with transmission.

A variant with high transmissibility could cause more infections and has a greater

chance to form new lineages in the examined evolutionary tree. These short and

multifurcation internal nodes can be measured by the accelerated rate of furcation (see

the details in Methods). Super-spreader transmission can also result in a

multifurcation node, while it only affects the furcation rate of a single node. Thus, it

can be distinguished from cases with high transmissibility.

Let us consider that a SARS-CoV-2 VOC has a beneficial haplotype with

multiple crucial mutations. To explain the sudden appearance of its high

transmissibility, we assume an instantaneous increase of fitness advantage as soon as

the last crucial mutation occurs (Fig. 1B). In this model, we call variant with a subset

of the beneficial mutations as pre-high-transmission variant, the furcation rate of

which may remain nearly unchanged. However, as soon as the last crucial mutation

occurs, the beneficial variant emerges and its transmissibility increases suddenly and

causes an instantaneously accelerated furcation rate in phylogenetic tree.

## A crucial non-coding deletion in the Alpha VOC

The Alpha lineage was examined using the Coronavirus GenBrowser (CGB) [14]. The sequential occurrence of the Alpha characteristic mutations was shown (Fig. 2A). To confirm the evolutionary path of the Alpha VOC, we also applied the VENAS [20] to obtain an evolution network of SARS-CoV-2 major haplotypes (Supplementary Fig. S2, Supplementary Table S1). The VENAS results are consistent with the CGB evolutionary tree.

The CGB evolutionary tree shows that many nodes remain to be bifurcating where the Alpha characteristic amino acid mutations occur (Fig. 2A). Multi-furcating nodes are frequently observed after a non-coding deletion (g.a28271-) occurs in the Alpha clade. The rate of furcation was calculated (Fig. 2B). The rate of furcation in the Alpha clade is higher than that in the Alpha-like clade. The Alpha-like clade indicates the viral strains lacking the non-coding deletion but carrying all other characteristic mutations of the Alpha VOC (Fig. 2A, Supplementary Table S1) [30], including all Alpha spike mutations [33]. The Alpha clade carries all those characteristic mutations and the non-coding deletion. It was also found that the higher furcation rate of the Alpha clade lasted for five months, indicating that it is not due to super-spreader transmission.

The frequencies of Alpha and Alpha-like strains were calculated (Fig. 2C). It was found that, within six months, the global frequency of Alpha increases rapidly to over 80% while that of Alpha-like is no more than 1%. Moreover, the reproduction

numbers (Rt) of Alpha and Alpha-like in the United Kingdom were estimated using

the EpiEstim [28]. The Rt of the Alpha lineage was generally larger than that of the

Alpha-like lineage (Fig. 2D). The larger Rt estimates of the Alpha lineage indicates

that the Alpha lineage has a transmission advantage over the Alpha-like. Therefore,

the non-coding deletion g.a28271- may effectively increase the viral transmissibility.

The higher transmissibility of the Alpha lineage causes that the number of

descendants in the Alpha lineage is highly significantly different with that in the

Alpha-like lineage ($n = $ 92,688 $vs$ 259, $P$-value$< 4.9 \times 10^{-324}$). Since pooling data

of viral sequences from different countries is likely to be biased due to complex

differences in sampling with respect to either viral genome sequencing capacities or

anti-contagion policies on the pandemic among the targeted countries [34]. To address

this problem, the numbers of descendants in the Alpha-like and Alpha lineages were

pairwise compared for individual countries and continents, i.e., England (27 $vs$

76,871), Spain (30 $vs$ 712), Switzerland (8 $vs$ 1,332), Germany (2 $vs$ 570), USA (8 $vs$

1,028), Australia (1 $vs$ 58), South America (1 $vs$ 22), Africa (1 $vs$ 86), and Asia (3 $vs$

642). The transmissibility of strains without or with the non-coding deletion is

significantly unequal (Supplementary Table S2, $P$-value$\leq 2.74 \times 10^{-6}$). The same

highly significant statistics was observed in 10 more countries, such as India and Italy.

Moreover, the same conclusion holds when considering different gender and age

groups (Supplementary Tables S3, S4). Therefore, the significantly difference in

transmissibility between the Alpha and Alpha-like clades has been observed which is

likely due to the non-coding deletion g.a28271-.

### The crucial non-coding deletion in the Delta VOC

The non-coding deletion (g.a28271-) was also found in the Delta VOC (Fig. 3A).

We used a recent data version (data.2022-04-14) to analyze this VOC since it became

prevalent globally after the Alpha VOC. It was observed that the number of furcation

increases as soon as the non-coding deletion (g.a28271-) occurs in the Delta lineage

(Fig. 3A), similar with the case in the Alpha lineage. The rate of furcation in the Delta

lineage is generally higher than that in the Delta-like lineage (Fig. 3C). The Delta-like

lineage represents the viral strains lacking the non-coding deletion but carrying all

other characteristic mutations of the Delta VOC (Fig. 3A).

The global frequencies of Delta and Delta-like strains were calculated (Fig. 3B).

It reveals that the global frequency of Delta increases rapidly to over 90% within six

months while that of Delta-like remains to be around 1%. The higher transmissibility

of the Delta lineage also causes that the number of its descendants is highly

significantly different with that in the Delta-like lineage ($n = $ 1,514,270 $vs$ 18,977,

$P$-value$< 4.9 \times 10^{-324}$). All these results support the modulation of non-coding

deletion g.a28271- on viral transmissibility.

### A crucial non-coding mutation in the Omicron VOC

The Omicron VOC first reported in South Africa in November 2021 [8] that

evolved into multiple sub-lineages, such as BA.1 and BA.2. The CGB evolutionary

tree shows that the most Omicron samples also carry a mutation at position 28271

(g.a28271t) (Fig. 4A). The mutation occurred independently in the two sub-lineages

(BA.1 and BA.2). Therefore, to examine the effects of the mutation on the viral

transmissibility, the furcation rate was calculated for the lineages with and without the

mutation. To keep the consistence, the term of Omicron stands for strains with the

mutation g.a28271t while Omicron-like indicates strains without the mutation in the

considered lineage (Fig. 4A). It was found that the Omicron lineage appears to have a

higher furcation rate than the Omicron-like lineage, indicating that the mutation

g.a28271t plays a key role in the viral transmissibility. Different from the situation in

Alpha and Delta, a delayed increase of furcation rate was observed (Fig. 4), which

suggests that g.a28271t may interact with other mutations to form a beneficial

haplotype.

**The non-coding mutations changes the core Kozak site of *N* gene**

The base 28,271 is located at the third base upstream of the start codon of the *N*

gene. It is the core Kozak site of the gene (Supplementary Fig. S3). The Kozak

sequence is a short sequence around the start codon and functions as the protein

translation initiation site in higher eukaryotes [35]. Nucleotides in each site of Kozak

sequence influence the translational efficiency, especially the positions -3 and +4. The

translational efficiency of the gene will be higher with A or G in position -3 and G in

position +4 [35]. The position 28,271 is the core position -3 of Kozak sequence of the *N*

gene. The g.a28271- deletion makes t28,270 to slip one base and changes the Kozak

context of gene *N* from a suboptimal one (A at -3, T at +4) to an undesirable one (T at

-3, T at +4) (Supplementary Fig. S3) [35]. It was expected that the mutation g.a28271t

may produce a similar effect by making the Kozak context of gene *N* change to an

undesirable one (T at -3, T at +4). When the homological site of the SARS-CoV

genome was mutated to another undesirable one (C at -3, T at +4), the expression of N

protein was reduced and the translation of ORF9b protein increased [36]. The ORF9b

protein was encoded by an alternative reading frame within the ORF of gene *N* and

the protein was found to be translated via a leaky ribosomal scanning mechanism [36]. It

has been found that SARS-CoV-2 ORF9b has an interferon (IFN) antagonistic activity

and suppresses type I interferon (IFN-I) responses [37,38]. Therefore, the non-coding

mutations may strengthen the interference of ORF9b on IFN-1 responses by changing

the core Kozak site of *N* gene.

### The non-coding mutations alone do not increase viral transmissibility

The non-coding mutations g.a28271- and g.a28271t were found to be crucial for

the high transmissibility of SARS-CoV-2. It was investigated whether these two

mutations function alone or co-act with other crucial mutations by considering

recurrent g.a28271- and g.a28271t mutations (including mutations due to

recombination). The non-coding mutations have appeared multiple times in the

evolution of SARS-CoV-2 (Supplementary Fig. S4), the frequency trajectories of the

two mutations were checked when the Alpha, Delta and Omicron lineages were

excluded. The first sample that carries one of the mutations appeared on June 16,

2020 (EPI_ISL_732537, with g.a28271-) and the frequency of the two mutations

generally remains much less than 10% in the next six months (Supplementary Fig. S5).

Therefore, the two non-coding mutations require epistasis or multilocus interaction

with other mutations to increase the viral transmissibility, consistent with the

observations in the Omicron VOC.

**The convergent evolution of the Alpha, Delta and Omicron VOCs**

To identify mutations that interact with g.a28271-/t, the convergent evolution of

the Alpha, Delta and Omicron VOCs may provide important clues. The three VOCs

diverged in January, 2020, and evolved independently in the background of the

D614G substitution (Fig. 5A). Thus the mutations of the three VOCs were screened in

the phylogenetic tree. The two noncoding mutations (g.a28271- and g.a28271t)

evolved convergently in the three VOCs (Fig. 5B). Moreover, another two

non-synonymous mutations (S:P681H/R and N:R203K/M) were found to occur

independently in the lineages.

Position 681 in the S protein is immediately upstream of the furin cleavage site

[39]. The mutation S:P681H occurred independently in the Alpha and Omicron VOCs

(Fig. 5B). It has been found that the mutated spike (681H) can be partially cleaved

when a protease inhibitor is added, while the un-mutated spike (P681) is nearly

uncleaved. This result suggests that the P681H mutation may increase furin cleavage

[40]. The mutation S:P681R, occurred in Sep, 2020 in the Delta VOC, also has the

similar effects [41]. Therefore, both the mutations S:P681H/R may be important for the

furin cleavage and viral entry.

The N protein is required for replication and packaging. Position 203 is in the

linker region of the N protein. The mutation N:R203K occurred on the common

ancestor of Alpha and Omicron VOCs in Feb, 2020 (Fig. 5B). Its adjacent mutation

N:G204R also occurs on the two VOCs (Fig. 5A). It has been found that the

double-mutated strain (203K/204R) has a replication advantage over the wild-type

(R203/G204) [42]. Similarly, the mutation N:R203M occurred in Sep, 2020, before the

Delta VOC emerged (Fig. 5B). The mutation N:R203M can enhance replication in

lung epithelial cells [43]. Therefore, both the mutations N:R203K/M may be crucial for

replication.

**DISCUSSION**

It is generally difficult to map causal mutations with fitness advantage along the

SARS-CoV-2 genome using traditional population genetic methods, such as selective

sweep [15,18,19], because of the extreme rarity of recombination in asexual population.

All mutations will be tightly linked and the methods to detect selective sweep actually

are not developed to be applied to viral genomes. For the same reason,

machine-learning based approaches [16] may also face the same challenge. Therefore,

we analyze the sequential occurrence order of mutations and the change of furcation

rate in the pandemic-scale phylogenetic tree of SARS-CoV-2 to map viral mutations

with fitness advantage. We find that the non-coding g.a28271- mutation may play a

crucial role in the high transmissibility of the Alpha and Delta VOCs, and g.a28271t

for the Omicron VOC. Both the mutations cause the A-to-T change at the core Kozak

site of the gene *N*. The Alpha-like and the Alpha strains encode the same spike protein; however, these strains appear to have different viral transmissibility. The Alpha and the Alpha-like spikes improve the angiotensin-converting enzyme 2 (ACE2) affinity for about 5-fold, comparing with the D614G spike [33]. However, our analysis shows that the increase of ACE2 affinity alone may not cause the high transmissibility of the Alpha VOC when the non-coding deletion g.a28271- is lack.

Sequence with accession EPI_ISL_601443 was previously recommended to be the canonical Alpha genomic sequence [30]. However, it does not carry the crucial non-coding deletion g.a28271-. Therefore, to investigate the viral transmissibility, we suggest using the sequence with accession EPI_ISL_629703 as the canonical Alpha genomic sequence (collected 21 October, 2020, in the UK) (Supplementary Fig. S1).

Genomic mutations related to the transmissibility of a pandemic etiological pathogen such as SARS-CoV-2 is complex and difficult to be revealed merely *via* genetic analysis with limited and incomplete supporting data of epidemiology. However, this study unveils that non-synonymous and non-coding mutations could affect the transmissibility synergistically as a beneficial haplotype. Overall, our analyses indicate that the non-coding A-to-T kozak site change of the gene *N* could be crucial for the high transmissibility of SARS-CoV-2 and may interact with mutations on the spike S1/S2 cleavage site and mutations that influence virus replication.

**Acknowledgments**

We thank Wolfgang Stephan for discussing on the issue of selective sweep and

the researchers who generated and deposited sequence data of SARS-CoV-2 in

GISAID, GenBank, CNGBdb, GWH, and NMDC. This work was supported by grants

from the Strategic Priority Research Program of the Chinese Academy of Sciences

(Grant No. XDB38030100), the National Key Research and Development Project

(Grant Nos. 2020YFC0847000, 2021YFC0863300, and 2020YFC0845900), the

National Natural Science Foundation of China (Grant No. 91531306), and the

Shandong Academician Workstation Program #170401 (to G.P.Z.).

## Data Availability

All the SARS-CoV-2 data can be obtained from the Coronavirus GenBrowser and

VENAS. Raw data were deposited on Mendeley at

https://data.mendeley.com/datasets/tbbxjy3gyr/draft?a=1be355b0-e347-4599-89dd-19

c950051f02. Other data are available from the corresponding authors upon request.

## Authors' contributions

Conceptualization, J.Y., G.Z., D.Y., Y.H.P., B.S., Y.Z., G.P.Z., Y.L., H.L.; Data

Analysis, J.Y., G.Z., D.Y., R.C., X.W., Y.L., C.Y., X.S.; Writing, J.Y., G.Z., D.Y., R.C.,

Y.L., Y.H.P., C.Y., X.S., Y.Z., G.P.Z., H.L.; Supervision & Funding Acquisition, G.Z.,

Y.Z., G.P.Z., Y.L., H.L.

## References

1        Rambaut, A. *et al. Preliminary genomic characterisation of an emergent SARS-CoV-2
         lineage in the UK defined by a novel set of spike mutations*,

<https://virological.org/t/preliminary-genomic-characterisation-of-an-emergent-sars-
cov-2-lineage-in-the-uk-defined-by-a-novel-set-of-spike-mutations/563> (2020).

2   Volz, E. *et al.* Assessing transmissibility of SARS-CoV-2 lineage B.1.1.7 in England. *Nature*
**593**, 266-269, doi:10.1038/s41586-021-03470-x (2021).

3   Davies, N. G. *et al.* Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7
in England. *Science* **372**, eabg3055 (2021).

4   WHO. Tracking SARS-CoV-2 variants.
*https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/* (2021).

5   Dhar, M. S. *et al.* Genomic characterization and epidemiology of an emerging
SARS-CoV-2 variant in Delhi, India. *Science* **374**, 995-999, doi:10.1126/science.abj9932
(2021).

6   Campbell, F. *et al.* Increased transmissibility and global spread of SARS-CoV-2 variants of
concern as at June 2021. *Euro Surveill* **26**, 2100509 (2021).

7   Meng, Z. *et al.* Transmission dynamics of an outbreak of the COVID-19 Delta variant
B.1.617.2 — Guangdong Province, China, May–June 2021. *China CDC Weekly* **3**, 584-586,
doi:10.46234/ccdcw2021.148 (2021).

8   Viana, R. *et al.* Rapid epidemic expansion of the SARS-CoV-2 Omicron variant in
southern Africa. *Nature* **603**, 679-686, doi:10.1038/s41586-022-04411-y (2022).

9   Suzuki, R. *et al.* Attenuated fusogenicity and pathogenicity of SARS-CoV-2 Omicron
variant. *Nature* **603**, 700-705, doi:10.1038/s41586-022-04462-1 (2022).

10  Hui, K. P. Y. *et al.* SARS-CoV-2 Omicron variant replication in human bronchus and lung
ex vivo. *Nature* **603**, 715-720, doi:10.1038/s41586-022-04479-6 (2022).

11  Kang, L. *et al.* A selective sweep in the Spike gene has driven SARS-CoV-2 human
adaptation. *Cell* **184**, 4392-4400, doi:10.1016/j.cell.2021.07.007 (2021).

12  Obermeyer, F. *et al.* Analysis of 6.4 million SARS-CoV-2 genomes identifies mutations
associated with fitness. *Science* **376**, 1327-1332 (2022).

13  Wu, F. *et al.* A new coronavirus associated with human respiratory disease in China.
*Nature* **579**, 265-269 (2020).

14  Yu, D. *et al.* Coronavirus GenBrowser for monitoring the transmission and evolution of
SARS-CoV-2. *Brief Bioinform*, bbab583 (2022).

15  Li, H. & Stephan, W. Maximum likelihood methods for detecting recent positive selection
and localizing the selected site in the genome. *Genetics* **171**, 377-384 (2005).

16  Lin, K., Li, H. P., Schlotterer, C. & Futschik, A. Distinguishing positive selection from neutral
evolution: boosting the performance of summary statistics. *Genetics* **187**, 229-244,
doi:10.1534/genetics.110.122614 (2011).

17  Sabeti, P. C. *et al.* Detecting recent positive selection in the human genome from
haplotype structure. *Nature* **419**, 832-837, doi:10.1038/nature01140 (2002).

18  Kim, Y. & Stephan, W. Detecting a local signature of genetic hitchhiking along a
recombining chromosome. *Genetics* **160**, 765-777 (2002).

19  Stephan, W. Selective sweeps. *Genetics* **211**, 5-13, doi:10.1534/genetics.118.301319
(2019).

20  Ling, Y. C. *et al.* An interactive viral genome evolution network analysis system enabling
rapid large-scale molecular tracing of SARS-CoV-2. *Sci Bull* **67**, 665-669,
doi:10.1016/j.scib.2022.01.001 (2022).

21    Zhao, W.-M. *et al.* The 2019 novel coronavirus resource. *Hereditas (Beijing)* **42**, 212-221
(2020).

22    Gong, Z. *et al.* An online coronavirus analysis platform from the National Genomics Data
Center. *Zool Res* **41**, 705-708 (2020).

23    Elbe, S. & Buckland-Merrett, G. Data, disease and diplomacy: GISAID's innovative
contribution to global health. *Glob Chall* **1**, 33-46 (2017).

24    Shu, Y. L. & McCauley, J. GISAID: Global initiative on sharing all influenza data - from
vision to reality. *Euro Surveill* **22**, 30494 (2017).

25    Sayers, E. W. *et al.* Database resources of the National Center for Biotechnology
Information. *Nucleic Acids Res* **49**, D10-D17 (2021).

26    Chen, F. *et al.* CNGBdb: China National GeneBank DataBase. *Hereditas (Beijing)* **42**,
799-809 (2020).

27    Zhang, Z. *et al.* Database resources of the National Genomics Data Center in 2020.
*Nucleic Acids Res* **48**, D24-D33 (2020).

28    Cori, A., Ferguson, N. M., Fraser, C. & Cauchemez, S. A New Framework and Software to
Estimate Time-Varying Reproduction Numbers During Epidemics. *Am J Epidemiol* **178**,
1505-1512, doi:10.1093/aje/kwt133 (2013).

29    Li, Q. *et al.* Early transmission dynamics in Wuhan, China, of novel coronavirus–infected
pneumonia. *N Engl J Med* **382**, 1199-1207 (2020).

30    Chand, M. *et al.* Investigation of novel SARS-CoV-2 variant of concern 202012/01.
*https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachme*
*nt_data/file/959438/Technical_Briefing_VOC_SH_NJL2_SH2.pdf* (2020).

31    Colijn, C. & Gardy, J. Phylogenetic tree shapes resolve disease transmission patterns. *Evol*
*Med Public Health* **2014**, 96-108, doi:10.1093/emph/eou018 (2014).

32    Oakley, T. H., Plachetzki, D. C. & Rivera, A. S. Furcation, field-splitting, and the
evolutionary origins of novelty in arthropod photoreceptors. *Arthropod Struct Dev* **36**,
386-400, doi:10.1016/j.asd.2007.08.002 (2007).

33    Gobeil, S. M.-C. *et al.* Effect of natural mutations of SARS-CoV-2 on spike structure,
conformation, and antigenicity. *Science* **373**, eabi6226 (2021).

34    Hsiang, S. *et al.* The effect of large-scale anti-contagion policies on the COVID-19
pandemic. *Nature* **584**, 262-267 (2020).

35    Kozak, M. At least six nucleotides preceding the AUG initiator codon enhance translation
in mammalian cells. *J Mol Biol* **196**, 947-950, doi:10.1016/0022-2836(87)90418-9 (1987).

36    Xu, K. *et al.* Severe acute respiratory syndrome coronavirus accessory protein 9b is a
virion-associated protein. *Virology* **388**, 279-285, doi:10.1016/j.virol.2009.03.032 (2009).

37    Jiang, H. W. *et al.* SARS-CoV-2 Orf9b suppresses type I interferon responses by targeting
TOM70. *Cell Mol Immunol* **17**, 998-1000, doi:10.1038/s41423-020-0514-8 (2020).

38    Wu, J. *et al.* SARS-CoV-2 ORF9b inhibits RIG-I-MAVS antiviral signaling by interrupting
K63-linked ubiquitination of NEMO. *Cell Rep* **34**, 108761 (2021).

39    Huang, Y., Yang, C., Xu, X. F., Xu, W. & Liu, S. W. Structural and functional properties of
SARS-CoV-2 spike protein: potential antivirus drug development for COVID-19. *Acta*
*Pharmacol Sin* **41**, 1141-1149, doi:10.1038/s41401-020-0485-4 (2020).

40    Lubinski, B. *et al.* Functional evaluation of the P681H mutation on the proteolytic
activation of the SARS-CoV-2 variant B.1.1.7 (Alpha) spike. *iScience* **25**, 103589,

464   doi:10.1016/j.isci.2021.103589 (2022).

465 41 Lubinski, B. *et al.* Spike protein cleavage-activation mediated by the SARS-CoV-2 P681R

466   mutation: a case-study from its first appearance in variant of interest (VOI) A.23.1

467   identified in Uganda. *bioRxiv*, doi:10.1101/2021.06.30.450632 (2022).

468 42 Wu, H. *et al.* Nucleocapsid mutations R203K/G204R increase the infectivity, fitness, and

469   virulence of SARS-CoV-2. *Cell Host Microbe* **29**, 1788-1801 (2021).

470 43 Syed Abdullah, M. *et al.* Rapid assessment of SARS-CoV-2–evolved variants using

471   virus-like particles. *Science* **374**, 1626-1632, doi:10.1126/science.abl6184 (2021).

**Figure 1. COVID-19 waves and a sudden appearance of high transmissibility**

**model.**

(A) Three COVID-19 waves caused by the Alpha, Delta and Omicron VOCs.

(B) Schematic diagram of a sudden appearance of high transmissibility model. The

triangles represent collapsed clades and the size of the triangle represents the size

of the clade. Each notch of branches represents a mutation. The last crucial

mutation is indicated in red.

**Figure 2. High transmissibility of Alpha compared with Alpha-like.**

(A) CGB evolutionary tree of SARS-CoV-2 lineage Alpha. The analysis was

performed on 400,051 high-quality SARS-CoV-2 genomic sequences (data

version "data.2021-03-06") using the Coronavirus GenBrowser [14]. The searchable

CGB ID of the internal node with g.a28271- is CGB84017.91425, assigned by the

CGB binary nomenclature system. The triangles represent collapsed clades. The

size of collapsed clades was labeled after the collapse triangle. The mutations on

the highlighted branches were labeled and the mutations on the Alpha and

Alpha-like clades were marked by notches. The number of Alpha-like strains is

259, and the number of Alpha strains is 92,688.

(B) The furcation rate of Alpha and Alpha-like.

(C) The frequency trajectory of Alpha and Alpha-like in United Kingdom.

(D) The Rt of Alpha and Alpha-like in United Kingdom. The grey area represents the

95% confidence interval.

**Figure 3. High transmissibility of Delta compared with Delta-like.**

(A) CGB evolutionary tree of SARS-CoV-2 lineage Delta. The analysis was

performed on 3,777,753 high-quality SARS-CoV-2 genomic sequences (data

version "data.2022-04-14") using the Coronavirus GenBrowser [14]. The searchable

CGB ID of the internal node with g.a28271- is CGB531065.736525, assigned by

the CGB binary nomenclature system. The triangles represent collapsed clades.

The size of collapsed clades was labeled after the collapse triangle. The mutations

on the highlighted branches were labeled and the mutations on the Delta and

Delta-like clades were marked by notches. The number of Delta-like strains is

18,977, and the number of Delta strains is 1,514,270.

(B) The frequency trajectory of Delta and Delta-like.

(C) The furcation rate of Delta and Delta-like.

**Figure 4. High transmissibility of Omicron compared with Omicron-like**

(A) CGB evolutionary tree of SARS-CoV-2 lineage Omicron. The analysis was

performed on 3,777,753 high-quality SARS-CoV-2 genomic sequences (data

version "data.2022-04-14") using the Coronavirus GenBrowser [14]. The CGB ID

of the internal nodes with g.a28271t are CGB3145480.3251694 and

CGB2437061.2958631 for BA.1, CGB3053091.3265861 for BA.2. The triangles

represent collapsed clades. The size of collapsed clades was labeled after the

collapse triangle. The mutations were marked by notches.

(B) The furcation rate of Omicron and Omicron-like.

**Figure 5. The convergent evolution of the Alpha, Delta and Omicron VOCs**

(A) Heatmap showing the presence of mutations in Alpha, Delta and Omicron (BA.1,

BA.2) VOCs. In the heatmap, four background mutations are in grey, mutated

alleles are in red and ancestral alleles in blue.

(B) Schematic diagram of convergent evolution of Alpha, Delta and Omicron VOCs.

Mutations under convergent evolution are labeled, the occurring time of

mutations are inferred by the CGB [14].