1

Title: Action at a distance: Defects in division plane positioning in the root meristematic zone affect cell organization in the differentiation zone

Authors: Alison M. Mills[1]**,** Carolyn G Rasmussen[1,2,*]

[1]Graduate Group in Biochemistry and Molecular Biology
[2]Department of Botany and Plant Sciences, Center for Plant Cell Biology, Institute of Integrative Genome Biology, University of California, Riverside.

[*]Correspondence to Carolyn Rasmussen: carolyn.rasmussen@ucr.edu

ORCID CGR 0000-0002-4354-6295
ORCID AMM 0000-0002-7391-1409

Keywords: Mitosis, phragmoplast, root twisting, cell file rotation, division plane orientation, Arabidopsis

Summary Statement:
Expression of *TAN1* in the root meristematic zone rescues cell file rotation defects in *tan1 air9* mutants, suggesting defects that occur in mitosis may influence organization of nondividing cells.

23
24
25
Abstract

Cell division plane orientation is critical for plant and animal development and growth. TANGLED1

(TAN1) and AUXIN-INDUCED-IN-ROOT-CULTURES9 (AIR9) are division-site localized

microtubule-binding proteins required for division plane positioning. *tan1* and *air9 Arabidopsis*

thaliana single mutants have minor or no noticeable phenotypes but the *tan1 air9* double mutant

has synthetic phenotypes including stunted growth, misoriented divisions, and aberrant cell-file

rotation in the root differentiation zone. These data suggest that TAN1 plays a role in nondividing

cells. To determine whether TAN1 is required in elongating and differentiating cells in the *tan1*

air9 double mutant, we limited its expression to actively dividing cells using the G2/M-specific

promoter of the syntaxin *KNOLLE* (*pKN:TAN1-YFP*). Unexpectedly, in addition to rescuing

division plane defects, *pKN:TAN1-YFP* rescued root growth and the root differentiation zone cell

file rotation defects in the *tan1 air9* double mutant. This suggests that defects that occur in the

meristematic zone later affect the organization of elongating and differentiating cells.

39
40
Introduction

Correct division plane orientation is key for patterning and growth across kingdoms. Because

plant cells are confined by cell walls, division positioning is tightly regulated (Facette et al., 2018;

Livanos and Müller, 2019; Rasmussen and Bellinger, 2018; Wu et al., 2018). Division plane

determination begins during S or G2, when the nucleus is repositioned within the cell (Facette et

al., 2018; Frey et al., 2010; Wada, 2018; Yi and Goshima, 2020). Polarity is often established and

maintained by nuclear repositioning and polar localization of proteins during asymmetric division

(Facette et al., 2018; Guo et al., 2021; Kimata et al., 2016; Muroyama and Bergmann, 2019; Shao

and Dong, 2016; Wada, 2018). Next, land-plant cells typically form a structure around the nucleus

at the cell cortex called the preprophase band (PPB). The PPB is a ring of microtubules,

microfilaments and associated proteins that marks the future position of the new cell wall, called

the division site (Li et al., 2015; Pickett-Heaps et al., 1999; Rasmussen and Bellinger, 2018;

Smertenko et al., 2017; Van Damme, 2009). Nuclear and PPB positioning often match division

predictions based on cell geometry (Martinez et al., 2018; Moukhtar et al., 2019). PPB

disassembly upon nuclear envelope breakdown precedes spindle formation (Dixit and Cyr, 2002).

After chromosome separation, the phragmoplast forms from the anaphase spindle to direct new

cell wall synthesis. The phragmoplast is an antiparallel array of microtubules with plus-ends facing

the cell center (Ho et al., 2012; McMichael and Bednarek, 2013; Müller and Jürgens, 2016).

Kinesins transport vesicles to form the cell plate (Lee and Liu, 2013; Smertenko et al., 2018). New

microtubule nucleation expands the phragmoplast outwards until the cell plate contacts the

division site (Gu and Rasmussen, 2022; Murata et al., 2013; van Oostende-Triplet et al., 2017).

62
Division-site localized proteins including TANGLED1 (TAN1), PHRAGMOPLAST ORIENTING

KINESIN1 (POK1), POK2, MICROTUBULE-ASSOCIATED PROTEIN 65-4 (MAP65-4), RAN

GTPASE ACTIVATING PROTEIN (RAN-GAP), MYOSIN VIII and KINESIN-LIKE CALMODULIN

BINDING PROTEIN (KCBP) remain at the cell cortex at the division site throughout cell division

(Buschmann et al., 2015; Herrmann et al., 2018; Li et al., 2017; Lipka et al., 2014; Morgan et al.,

2008; Walker et al., 2007; Wu and Bezanilla, 2014). Many of these proteins are important for

division plane positioning, often during telophase. TAN1 is a division-site-localized protein

required for phragmoplast guidance to the division site in maize (Martinez et al., 2017; Smith et

al., 2001; Walker et al., 2007). TAN1 organizes microtubules at the cell cortex called cortical

telophase microtubules which are incorporated into the phragmoplast to direct its movement at

the cell cortex (Bellinger et al.). TAN1 binds and bundles microtubules in vitro (Martinez et al.,

2020). Although the *tan1* maize mutant has misplaced divisions and stunted growth, *tan1*

Arabidopsis thaliana (Arabidopsis) mutants grow as well as wild-type plants and have minor

division placement defects (Walker et al., 2007). Another division-site-localized protein, AIR9,

also binds microtubules. AIR9 localizes to interphase cortical microtubule arrays, as well as co-

localizing with the PPB, the phragmoplast, and localizing to the division site during late telophase

(Buschmann et al., 2006). Similar to *tan1* single mutants, *air9* single mutants resemble wild-type

plants (Buschmann et al., 2015). Due to their similar division-site localization, *tan1 air9* double

mutants were generated in Arabidopsis. Combining mutations in both *tan1* and *air9* results in

division-plane-positioning defects, stunted growth, and root twisting in the differentiation zone (Mir

et al., 2018). While PPBs and phragmoplasts were both frequently misoriented in *tan1 air9*

mutants, improper phragmoplast guidance is the primary defect (Mir et al. 2018). Transforming

the *tan1 air9* double mutant with *TAN1-YFP* driven by the constitutive viral Cauliflower mosaic

CaMV35S promoter rescues root growth, misoriented divisions, and cell-file-rotation defects (Mir

et al., 2018).

We hypothesized that TAN1 may also have a role in organizing interphase microtubules in

elongating and differentiated cells, because *tan1 air9* mutants had aberrant cell-file rotation in the

root differentiation zone, minor defects in interphase microtubule organization, and root growth

defects that were enhanced by the microtubule-depolymerizing-drug propyzamide (Mir et al.,

2018). Cell file rotation phenotypes are often caused by mutations in microtubule-associated

proteins or tubulin that alter the organization or stability of the interphase cortical microtubule

array (Abe et al., 2004; Buschmann and Borchers, 2020; Buschmann et al., 2004; Hashimoto,

2015; Ishida et al., 2007; Nakajima et al., 2004; Sakai et al., 2008; Sedbrook et al., 2004; Shoji et

al., 2004). For example, in several *alpha-tubulin* mutants, cell file rotation occurred in hypocotyls

and root differentiation zones and in isolated cultured mutant cells (Abe et al., 2004; Buschmann

et al., 2009; Ishida et al., 2007; Thitamadee et al., 2002). Cell file twisting also occurs when cell

elongation differs between epidermal and cortical cells. Arabidopsis treated with compounds that

affect microtubule stability, such as oryzalin or propyzamide, have helical cell files due to cortical

cell swelling and reduced longitudinal cell expansion (Furutani et al., 2000; Hashimoto, 2002).

Therefore, defects in organ twisting are sometimes due to interphase microtubule disruption and

likely independent of division-plane defects. However, several examples suggest that division-

plane-orientation defects may lead to cell-file-rotation defects (Cnops et al., 2000; Wasteneys and

Collings, 2009). Double mutants in two related receptor-like kinases have defects in division plane

orientation near the quiescent center and in the endodermis and also have abnormal root skewing

(Goff and Van Norman, 2021). Therefore, it is possible that either mitotic or non-mitotic defects

lead to aberrant growth and root twisting defects.

To determine whether mitotic *TAN1* expression was sufficient to rescue root twisting in the

differentiation zone of *tan1 air9* double mutants, we drove *TAN1* expression using the G2/M-

phase-specific *KNOLLE* promoter (Lukowitz et al., 1996; Menges et al., 2005). KNOLLE is a

syntaxin/Qa-SNARE required for cell-plate vesicle fusion (Strompen et al., 2002; Völker et al.,

2001). The *KNOLLE* promoter drove *TAN1* expression in mitotic cells which rescued root growth

and cell-file-rotation defects in the *tan1 air9* double mutant. Our results suggest that cell-file-

rotation defects in the *tan1 air9* double mutant are likely due to defects that occur in actively

dividing meristematic cells, and not due to a lack of TAN1 in nondividing cells.

Results & Discussion

We generated two independent native-promoter TAN1 fluorescent protein fusions to determine

whether *TAN1-YFP or CFP-TAN1* expressed by its native promoter would rescue the *tan1 air9*

double mutant. Both constructs rescued the *tan1 air9* mutant. Previous studies showed that *35S-*

driven *TAN1* expression rescued *tan1 air9* mutants (Mir et al., 2018). We drove expression of

CFP-TAN1 and *TAN1-YFP* using 1263 bp upstream of the start codon, *pTAN:CFP-TAN1* and

pTAN:TAN1-YFP, and transformed or crossed them into the *tan1 air9* double mutant. Cell shape

in the root tip (Figure 1A) and cell file rotation in the differentiation zone of *pTAN:CFP-TAN1 tan1*

air9 plants was restored to *air9* single mutant levels (Figure 1B, 1C). Single *air9* mutants are

indistinguishable from wild-type plants (Buschmann et al., 2015; Mir et al., 2018). Root cell

division primarily occurs at the root tip (the meristematic zone). Above that, non-dividing cells

elongate in the elongation zone. Root hairs mark the differentiation zone, where root cells mature

and differentiate into different cell types (Wachsman et al. 2015). *tan1 air9* mutant roots tend to

twist left with variable transverse cell-wall angle values that skew above 90° (Mir et al., 2018).

pTAN:CFP-TAN1 rescued *tan1 air9* root growth, with *pTAN:CFP-TAN1* expressing plants growing

slightly longer than *air9* single mutants (Figure 1D). *pTAN:TAN1-YFP* also fully rescued *tan1 air9*

root growth and restored normal root tip patterning (Supplementary Figure 1). Measuring PPB

and phragmoplast angles is a metric for division plane orientation. PPB and phragmoplast angles

were measured relative to the left-hand cell wall. *pTAN:CFP-TAN1* fully rescued PPB and

phragmoplast positioning defects in *tan1 air9* mutants, restoring angle variances close to 90°

(Figure 1E). This shows that mitotic expression of *TAN1* by its native promoter and fluorescent

protein fusion at either end of the TAN1 protein is sufficient for normal plant growth, including the

expansion and patterning of nondividing cells in the *tan1 air9* double mutant.

Previous fluorescence measurements of TAN1-YFP in wild-type lines expressing *pTAN:TAN1-*

YFP demonstrated that fluorescent signal above background was limited to the meristematic zone

(Mir et al., 2018). We hypothesized that TAN1 accumulated at low but undetectable levels in

interphase cells when driven by its native promoter. To test whether *TAN1* expression limited to

mitotic cells influenced root growth and suppressed root twisting in the *tan1 air9* double mutant,

we fused the *KNOLLE* promoter to *TAN1-YFP* (*pKN:TAN1-YFP*) and transformed it into the *tan1*

air9 double mutant. The *KNOLLE* promoter is specifically expressed in G2/M and is contingent

on the MYB (myeloblastosis) transcription factors MYB3R1 and MYB3R4 which promote mitosis-

specific gene expression (Haga et al., 2011; Yang et al., 2021). Our prediction was that

pKN:TAN1-YFP would fully rescue mitotic defects but not restore root growth or suppress

aberrant cell file rotation within the root differentiation zone in the *tan1 air9* mutant.

pKN:TAN1-YFP fully rescued the defects in *tan1 air9* mutants (Figure 2, other independent lines

in Supplementary Figure 2). This includes rescuing cell patterning and cell-file-rotation defects

(Figure 2A-C), root growth (Figure 2D), and PPB and phragmoplast positioning (Figure 2E). In

addition, *pKN*-drivenTAN1-YFP localized to the division site during mitotic stages similar to

pTAN1-driven CFP-TAN1 (Supplementary Figure 3). We compared phenotypes of *pKN:TAN1-*

YFP to the *35S:TAN1-YFP* lines which rescue the *tan1 air9* mutant (Mir et al. 2018). Both

35S:TAN1-YFP and *pKN:TAN1-YFP* significantly rescued the *tan1 air9* double mutant (Figure 3A

& 3B). Root growth and PPB and phragmoplast angles were equivalent in *tan1 air9* plants

expressing *pKN:TAN1-YFP* or *35S:TAN1-YFP* (Figure 3D & 3E). However, *pKN:TAN1-YFP*

reduced cell-file-rotation variability slightly more than *35S:TAN1-YFP* (Figure 3C). This suggests

that expressing *TAN1* in dividing cells is sufficient to fully rescue the *tan1 air9* double mutant. To

determine why rescue with the *KNOLLE* promoter resulted in less cell-file-rotation variance, we

measured TAN1-YFP fluorescence intensities in the *35S:TAN1-YFP* and *pKN:TAN1-YFP* lines.

pKN:TAN1-YFP was expressed strongly in the meristematic zone of root tips (Figure 4B), often

showing TAN1-YFP fluorescence in recently divided cells, similar to native promoter driven

accumulation (Figure 1, Supplementary Figure 1, (Mir et al., 2018)). Indeed, TAN1-YFP

accumulated at higher levels in the meristematic zone when expression was driven by the

KNOLLE promoter (Figure 4G). However, unlike TAN1-YFP from *p35S:TAN1-YFP*, TAN1-YFP

did not accumulate above background levels in the elongation and differentiation zone of roots

expressing *pKN:TAN1-YFP* (Figure 4F & 4G). Lack of TAN1-YFP outside the meristematic zone

and more complete rescue of *tan1 air9* cell file rotation by *pKN:TAN1-YFP* suggests that TAN1 is

not required in elongating and differentiating cells. In other words, cell-file-rotation defects may

be a consequence of defects that occur within the root meristematic zone either during mitosis or

shortly afterwards.

Another example of defects in mitotic expression and division plane positioning affecting

nondividing cell organization occurs in the MYB activated GRAS-type (GIBBERELLIC-ACID

INSENSITIVE, REPRESSOR of GAI and SCARECROW-type) transcription factor *scarecrow-like*

28-3 (*scl28-3)* mutant. G2/M specific gene expression controlled by SCL28 is important for mitotic

progression and division-plane positioning. *scl28-3* mutants have both misoriented divisions and

root twisting (Goldy et al., 2021).

How do defects that occur within the meristematic zone influence the patterning or shape of

nondividing, differentiating root cells and root growth? Our hypothesis is that misshapen cells and

improper division plane orientation in *tan1 air9* double mutants cause the uneven distribution of

mechanical stresses across the root, which then triggers cell wall integrity responses that limit

growth and alter root organization. Cell wall stress patterns depend on cell geometry and the

mechanical properties of cell walls (Cosgrove, 2018; Hamant and Haswell, 2017; Whitewoods

and Coen, 2017; Schopfer, 2006).

Division-plane positioning is a way plants may respond to mechanical stress (Chakrabortty et al.,

2018; Louveaux et al., 2016). Cell division relieves mechanical stress by creating smaller cells

with less surface; further, divisions along maximal tensile stress promote growth homogeneity

(Alim et al., 2012; Sapala et al., 2018). Microtubules often align parallel to maximal tensile stress

(Hamant et al., 2008; Heisler et al., 2010; Sampathkumar et al., 2014; Uyttewaal et al., 2012) and

cortical-microtubule alignment often influences PPB placement (Louveaux et al., 2016;

Rasmussen et al., 2013; Wick and Duniec, 1983). However, division plane positioning is disrupted

in mutants with division plane orientation defects. Although *tan1 air9* cells may perceive

mechanical stress, phragmoplast guidance defects prevent construction of new cell walls in an

orientation that minimizes mechanical stress. Abnormal stresses are perceived by receptor-like

kinases involved in the cell-wall-integrity response. Cell-wall-integrity responses trigger slow

growth, upregulation of stress responses, and changes in cell morphogenesis, (Buschmann and

Borchers, 2020; Caño-Delgado et al., 2003; Gonneau et al., 2018; Hématy et al., 2007; Wolf et

al., 2014), which may contribute to the stunted growth and twisted cell files observed in the *tan1*

air9 double mutant.

Materials and methods

<u>Plasmid Construction</u>

pKN:TAN1-YFP was generated by amplifying 2152 bp of the 5' *KNOLLE* (AT1G08560) promoter

from Columbia with primers pKN-5'SacI Fw and pKN-5'EcoRI Rw. EcoRI and StuI double

digestion was used to introduce the *KNOLLE* promoter into pEZT-NL containing the TAN1 coding

sequence. Primers 35SpKN5' Fw and YFP XhoI Rw were used to amplify *pKN:TAN1-YFP* then

XhoI and StuI double digestion was used to clone *pKN:TAN1-YFP* into pEGAD, a gift from

Professor Sean Cutler (University of California, Riverside).

pTAN:CFP-TAN1 was created by overlapping PCR. The 1263bp 5' sequence upstream of

genomic *TAN1* was amplified using *Np:AtTAN-YFP (Walker et al., 2007)* as a template with the

primers NpTANSacIFor and NpTANceruleanRev. Cerulean fluorescent protein (CFP) was

amplified using Cerulean CDS in pDONR221P4r/P3r, a kind gift from Professor Anne Sylvester

(University of Wyoming), as template with the primers NpTANceruleanFor and

CeruleanpEarleyRev. TAN1 CDS was amplified using *35S:YFP-TAN1* in pEarley104 as a

template with the primers CeruleanpEarleyFor and pEarleyOCSPstIRev. The 1263bp TAN1

native promoter, CFP, and TAN1 CDS were combined to create *pTAN:CFP-TAN1* by overlapping

PCR with the primers NpTANSacI and pEarleyOCSPstIRev. SacI and PstI double digest was

used to subclone *pTAN:CFP-TAN1* into pJHA212G, a kind gift from Professor Meng Chen

(University of California, Riverside).

<u>Generation of Transgenic Lines</u>

Transgenic Arabidopsis lines were generated using *Agrobacterium tumefaciens*-mediated floral

dip transformation as described (Clough and Bent, 1999). Previously described *tan1 air9* mutants

(Mir et al., 2018), *csh-tan* (*TAN1,* AT3G05330 (Walker et al., 2007)) and *air9-31* (*AIR9*,

AT2G34680 (Buschmann et al., 2015)), were used for floral dip transformation of *pKN:TAN1-YFP*

and T1 transgenic plants were subsequently selected on 1/2 MS plates containing 15 μg/mL

glufosinate (Finale; Bayer). TAN1-YFP signal in T1 plants was confirmed by confocal microscopy

before being transferred to soil and selfed. The genotypes of *csh-tan1 air9-31* transformants was

confirmed using the primers AIR9_cDNA 2230 F and AIR9 gnm7511 R (to identify *AIR9* wild-

type), AIR9 gnm7511 R and Ds5-4 (to identify T-DNA insertion in *AIR9*), ATLP and AtTAN 733-

CDS Rw (to identify TAN1 wild-type), and AtTAN 733-CDS Rw and Ds5-4 (to identify T-DNA

insertion in *TAN1*). The microtubule marker *CFP-TUBULIN* (Kirik et al., 2007), a kind gift from

Professor David Erhardht (Stanford University) was crossed into *pKN:TAN1-YFP tan1 air9* plants

using *tan1 air9 CFP-TUBULIN* plants (Mir et al., 2018).

air9-5 tan-mad Columbia/Wassilewskija double mutants (Mir et al., 2018) expressing the

microtubule marker *UBQ10:mScarlet-MAP4* (Pan et al., 2020)*, a* kind gift from Professor

Zhenbiao Yang (University of California, Riverside), was used for floral dip transformation of

pTAN1:CFP-TAN1 and selected on 1/2 MS plates containing 100 μg/mL gentamicin (Fisher

Scientific). T1 seedlings were screened for mScarlet and CFP signal and then transferred to soil

to self.

<u>Growth conditions and root length measurements</u>

Plates containing ½ strength Murashige and Skoog (MS) media (MP Biomedicals; Murashige and

Skoog, 1962) containing 0.5 g/L MES (Fisher Scientific), pH 5.7, and solidified with 0.8% agar

(Fisher Scientific) were used to grow Arabidopsis seedlings. *tan1 air9* transgenic lines expressing

T3 *p35S:TAN1-YFP,* T2 *pKN:TAN1-YFP,* and T2 *pTAN:CFP-TAN1* were used for root length

experiments. At least 3 biological replicates were used for each root growth assay. 5-7 1/2 MS

plates were used for each replicate. 12-15 seeds were sown in a single level line on each plate

with untransformed *tan1 air9* double mutants and *air9* single mutants sown on plates alongside

double mutants expressing *TAN1* constructs. Seeds were stratified on plates in the dark at 4°C

for 2 to 5 days. After stratifying, plates were positioned vertically in a growth chamber (Percival)

with a 16/8-h light/dark cycle and temperature set to 22°C. Each biological replicate was placed

in the growth chamber on different days. 8 days after stratification, plates were scanned (Epson)

and root lengths were measured using FIJI (ImageJ, http:/ /fiji.sc/). Transgenic seedlings were

screened for fluorescence by confocal microscopy to identify seedlings expressing YFP, CFP and

mScarlet translational fusion transgenes when present. Each root growth experiment had a

minimum of 3 biological replicates. Statistical analysis of root length was determined using

Welch's t-test with Prism (GraphPad) and replicates were checked for discrepancies in statistical

significance before pooling replicates for analysis. Root length plots were created using Prism

(GraphPad).

To assess the ability of *TAN1* driven by its native promoter to rescue the *tan1 air9* double mutant,

Np:AtTAN-YFP (Walker et al., 2007) was crossed to *tan-mad air9-5* double mutants. The progeny

of *pTAN1:TAN1-YFP tan-mad*/+ *air9-5*/+ plants were sown on 1/2 MS media and grown as

described above. The seedlings were screened by confocal microscopy for the presence of TAN1-

YFP and then collected for genotyping. Seedlings were genotyped with primers AtExon1_1For

and At255AfterStopRev (to identify TAN wild-type), JL202 and ATLP (to identify T-DNA insertion

in *TAN1*), AIR9-5RP and AIR9-5LP (to identify wild-type *AIR9*), and LBb1.3 and AIR9RP (to

identify T-DNA insertion in *AIR9*) (Supplementary Table 1). The length of *tan1 air9* double mutants

expressing *pTAN1:TAN1-YFP* was compared to *tan1 air9* double mutants and *air9* single mutant

siblings lacking *pTAN1:TAN1-YFP*. *air9* single mutants used for root length analysis included

air9/air9 TAN1/TAN1 and *air9/air9 TAN1/tan1* plants.

<u>Confocal Microscopy</u>

Imaging and screening was performed using Micromanager software (micromanager.org) running

an inverted Ti Eclipse (Nikon) with motorized stage (ASI Piezo) and spinning-disk confocal

microscope (Yokogawa W1) built by Solamere Technology. Solid-state lasers (Obis) were used

with standard emission filters (Chroma Technology). Excitation 445, emission 480/40 (for CFP-

translational fusions); excitation 514, emission 540/30 (for YFP-translational fusions); and

excitation 561, emission 620/60 (for propidium iodide and mScarlet-MAP4) were used. The 20x

objective has a 0.75 numerical aperture. The 60x objective was used with perfluorocarbon

immersion liquid (RIAAA-6788, Cargille) and has a 1.2 numerical aperture objective.

<u>Measurements of PPB and phragmoplast angles and cell file rotation</u>

All angle data was gathered from at least 3 biological replicates. Each replicate consisted of 5-7

1/2 MS plates with 12-15 seeds sown on each plate. 4-5 seeds of each genotype were sown on

each plate to ensure growing conditions were identical. Each replicate was moved from stratifying

to the growth chamber on independent days. Seedlings were then imaged at 8 days after

stratification. Images were collected by confocal microscopy using the 20x objective to collect

images of the differentiation zone for cell file angles and the 60x objective to collect images of

root tips expressing a microtubule marker (*CFP-TUBULIN* or *mScarlet-MAP4*) for PPB and

phragmoplast angles. The differentiation zone was identified by the presence of root hairs. Angles

were measured using FIJI. Cell file angles were measured from the left hand side of the cell taking

the angle between the long axis of the root and the transverse cell wall using images of root cells

in the differentiation zone. PPB and phragmoplast angles were measured by taking the angle

between the left-hand cell wall and the PPB or phragmoplast. Because CFP-TUBULIN was faint

and poorly marked cell boundaries, the cell walls of *CFP-TUBULIN* expressing seedlings were

stained with 10 µM PI for 1 minute before destaining in distilled water prior to imaging. Each angle

measurement represents a single angle measured from one cell.

Statistical analyses were performed using Excel (Microsoft Office) and Prism (GraphPad). To

compare normally distributed variance of PPB and phragmoplast angles F-test was used.

Levene's test was used to compare variances of cell file angle measurements because *tan1 air9*

cell file angles are non-normally distributed due to left hand twisting of the roots. Angle variance

across biological replicates was checked before pooling data.

<u>Fluorescence Intensity Measurements</u>

air9, *35S:TAN1-YFP tan1 air9*, and *pKN:TAN1-YFP tan1 air9* plants were grown on 1/2 MS plates

as described above. 8 days after stratification, plants were imaged by confocal microscopy using

identical settings. Root tips were imaged using the 60X objective. The median fluorescence

intensity of an 116,001.5 μm^2 area was measured from multiple individual plants of each

genotype. Each fluorescence measurement represents the median fluorescence from a single

meristematic zone from one plant. Elongation zone and differentiation zone images were taken

with the 20x objective and the median fluorescence intensity of an 12,323.4 μm^2 area was

measured from multiple individual plants of each genotype. Each fluorescence measurement

represents the median fluorescence from a single elongation or differentiation zone from one

plant.

Acknowledgements

We thank Stephanie Martinez (University of California, Riverside, UCR) and Aimee Uyehara
(UCR) for improving manuscript clarity, Professors Henrik Buschmann (Osnabruck University),
Meng Chen (UCR), Zhenbio Yang (UCR), Sean Cutler (UCR), David Ehrhardt (Carnegie
Institute), Anne Sylvester (University of Wyoming) and Dr. Ricardo Mir (UCR) for materials or

facilities, and Professor Jaimie Van Norman (UCR) for cloning advice. Funding from NSF-MCB

grant 1716972, NSF-CAREER grant 1942734, and USDA CA-R-BPS-5108-H

is gratefully acknowledged.

Figure 1: The TAN1 native promoter fused to TAN1 (*pTAN:CFP-TAN1*) rescues the *tan1 air9* double mutant. A) Maximum projections of 20 1-µm Z-stacks of root tips of an *air9*, *pTAN:CFP-TAN1* (magenta) *tan1 air9*, and untransformed *tan1 air9* plants expressing microtubule marker *UBQ10:mScarlet-MAP4* (green in the top middle panel). Bars = 25 µm. B) Maximum projections of 15 1-µm Z-stacks of the differentiation zone of an *air9*, *pTAN:CFP-TAN1 tan1 air9*, and untransformed *tan1 air9* plants expressing *UBQ10:mScarlet-MAP4*. Bars = 50 µm. C) Cell-file-rotation angles of *air9*, *pTAN:CFP-TAN1 tan1 air9*, and untransformed *tan1 air9* plants, n>11 plants for each genotype and n>57 cells for angle measurements. Cell-file-angle variances were compared with Levene's test due to the non-normal distribution. D) Root-length measurements from 8 days after stratification of *air9*, *pTAN:CFP-TAN1 tan1 air9*, and untransformed *tan1 air9*

354 plants, n>25 plants for each genotype, compared by two-tailed t-test with Welch's corrections. E)
355 PPB and phragmoplast angle measurements in *air9*, *pTAN:CFP-TAN1 tan1 air9*, and
356 untransformed *tan1 air9* plants, n>9 plants for each genotype. N>41 cells for angle
357 measurements. PPB and phragmoplast angle variations compared with F-test. ns indicates not
358 significant, ** P-value <0.01, **** P-value <0.0001.

Figure 2: Full rescue of the *tan1 air9* double mutant with the G2/M-specific *KNOLLE* promoter fused to TAN1 (*pKN:TAN1-YFP*). A) Propidium iodide (PI) stained cell walls in root tips of an *air9*, *pKN:TAN1-YFP tan1 air9*, and untransformed *tan1 air9* plants. Bars = 25 µm. B) Maximum projections of 10 1-µm Z-stacks of PI-stained differentiation zone root cell walls. Bars = 50 µm. C) Cell-file-rotation angles of *air9*, *pKN:TAN1-YFP tan1 air9*, and untransformed *tan1 air9* plants, n>23 plants for each genotype. N>114 cells for angle measurements. Cell-file-rotation angle variances were compared with Levene's test due to the non-normal distribution. D) Root-length measurements from 8 days after stratification of *air9*, *pKN:TAN1-YFP tan1 air9*, and untransformed *tan1 air9* plants, n>25 plants for each genotype, compared by two-tailed t-test with Welch's corrections. E) PPB and phragmoplast angle measurements in *air9*, *pKN:TAN1-YFP*

372   *tan1 air9*, and untransformed *tan1 air9* plants, n>20 plants for each genotype. N>34 cells for
373   angle measurements. PPB and phragmoplast angle variations compared with F-test. ns indicates
374   not significant, **** P-value <0.0001.
375

376
377

378
379

Figure 3: Comparison between *KNOLLE* promoter driven (*pKN:TAN1-YFP)* and *35S* driven
TAN1 (*p35S:TAN1-YFP*) rescue of the *tan1 air9* double mutants A) Propidium iodide (PI)
stained root tips of *tan1 air9* mutants expressing *p35S:TAN1-YFP*, *pKN:TAN1-YFP*, and
untransformed plants. Bars = 25 μm. B) Maximum projections of 10 1-μm Z-stacks of PI-stained
differentiation-zone root cell walls of *tan1 air9* mutants expressing *p35S:TAN1-YFP*, *pKN:TAN1-*
YFP, and untransformed plants. Bars = 50 μm. C) Cell-file-rotation angles of *tan1 air9* mutants

expressing *p35S:TAN1-YFP*, *pKN:TAN1-YFP*, and untransformed plants, n>13 plants for each
genotype. N>64 cells for angle measurements. Angle variances were compared with Levene's
test. D) Root-length measurements from 8 days after stratification of *tan1 air9* mutants expressing
p35S:TAN1-YFP, *pKN:TAN1-YFP*, and untransformed plants, n>17 plants for each genotype,
compared by two-tailed t-test with Welch's corrections. E) PPB and phragmoplast angle
measurements in *tan1 air9* double mutants expressing *p35S:TAN1-YFP*, *pKN:TAN1-YFP*, and
untransformed plants, n>12 plants for each genotype. N>39 cells for angle measurements. PPB
and phragmoplast angle variations compared with F-test. ns indicates not significant, * P-value
<0.05, **** P-value <0.0001.

Figure 4: Comparison of TAN1-YFP fluorescence intensity when driven by the constitutive
***35S* promoter (*p35S:TAN1-YFP)* and G2/M-specific *KNOLLE* promoter (*pKN:TAN1-YFP)* in**
***tan1 air9* roots.** (A and B) Micrographs of the meristematic zone, (C and D) maximum projections
of 3 1-µm Z-stacks of the elongation zone, and (E and F) maximum projections of 10 1-µm Z-
stacks of the differentiation zone of *tan1 air9* mutants expressing (A, C, E) *p35S:TAN1-YFP* or
(B, D, F) *pKN:TAN1-YFP*. Cell walls were stained with propidium iodide. Root tip and elongation
zone, bars = 50µm. Differentiation zone, bars = 200µm. G) TAN1-YFP fluorescence-intensity
measurements (arbitrary units, a.u.) from the meristematic zone, elongation zone, and
differentiation zone of *tan1 air9* mutants expressing *p35S:TAN1-YFP*, *pKN:TAN1-YFP*, and *air9*

407   mutants, n=8 plants for each genotype, fluorescence compared with Mann-Whitney U test. *** P-
408   value <0.001. ns indicates not significant.

425   **Supplementary Figures 1-3 and Supplementary Table**
426

427
428
Supplementary Figure 1. *TAN1-YFP* expressed by its native promoter (*pTAN:TAN1-YFP*)
rescues *tan1 air9* double mutant root growth. Confocal images of propidium iodide-stained
roots of *tan1 air9* plants. A) A *tan1 air9* plant expressing *pTAN:TAN1-YFP*. B) A negative sibling
tan1 air9 plant. Bars = 50 µm. C) Root length measurements from 8 days after stratification of
air9 single mutants (left), *pTAN1:TAN1-YFP tan1 air9* double mutants (middle), and *tan1 air9*
double mutants (right). n > 10 plants for each genotype, compared by two-tailed t-test with Welch's
correction. ns indicates not significant, **** P-value <0.0001.
436
437
438

439
Supplementary Figure 2. *pKN:TAN1-YFP tan1 air9* **lines show significant rescue compared to untransformed** *tan1 air9.* A) Cell file rotation angles of *air9* single mutants (left), two transgenic lines expressing pKN*:TAN1-YFP* in the *tan1 air9* double mutant designated as line 4 (center left) and line 5 (center right) and untransformed plants (right), n >17 plants for each genotype. N > 146 cells for angle measurements. Angle variances were compared with Levene's test. B) Root length measurements from 8 days after stratification of *air9* single mutants (left), two transgenic lines expressing pKN*:TAN1-YFP* in the *tan1 air9* double mutant (middle), and untransformed plants (right), n > 21 plants for each genotype, compared by two-tailed t-test with Welch's correction. C) PPB and phragmoplast angle measurements in dividing root cells of *air9* single mutants (left), two transgenic lines expressing pKN*:TAN1-YFP* in the *tan1 air9* double mutant (middle), and untransformed plants (right), n > 15 plants for each genotype. N > 69 cells

451 for angle measurements. Angle variations compared with F-test. ns indicates not significant, * P-
452 value < 0.05, **** P-value <0.0001.

453
454

455
456
457

Supplementary Figure 3. Division site localization of TAN1-YFP driven by the *KNOLLE*
promoter (*pKN:TAN1-YFP*) and CFP-TAN1 driven by the *TAN1* promoter (*pTAN:CFP-TAN1*)
in *tan1 air9* double mutants. A-D) Confocal images of propidium iodide-stained (Cell Wall) roots
of *tan1 air9* plants expressing *pKN:TAN1-YFP* and *CFP-TUBULIN* (MTs) in dividing root tip cells.
E-F) Maximum projections of 3 1-μm Z-stacks of *tan1 air9* plants expressing *pTAN:CFP-TAN1*
and the microtubule (MTs) marker *UBQ10:mScarlet-MAP4* in dividing root tip cells.
Representative images of cells with (A&E) broad early PPBs, (B&F) late narrow PPBs, (C&G)
metaphase spindles, and (D&H) phragmoplasts. Bars = 10 μm.

466
467

468 **Supplementary Table 1. Primers used for cloning and genotyping.**

469
470

Primer Name	Sequence
ATRP	ATCTCTTAGGAACCAAAACCGGACGCTGT
ATLP	GATCCGTTACGAAAGTGAACACCTTTATC
JL202	CATTTTATAATAACGCTGCGGACATCTAC
AIR9-5RP	TGGATCAGCTGCAACATTATTC
AIR9-5LP	ATTAACATTTTGCAACGCAGG
LBb1.3	ATTTTGCCGATTTCGGAAC

Ds5-4	TACGATAACGGTCGGTACGG
AtTAN 733-CDS Rw	AAATAGAGGGGTTCGGAAAAAGAACC
AIR9 gnm7511 R	CCTCCAGTATATGAAGCAACAAAGC
AIR9_cDNA 2230 F	GATGAGGAATATATGTTATCTTTAGATG
pKN-5'SacI Fw	GAGGAGCTCCAGAAGAAAAAGAAAAAGTTCTC
pKN-5'EcoRI Rw	TAAGCGGAATTCCTTTTTCACCTGAAA
35SpKN5' Fw	ACCCACAGATGGTTAGAGagg
YFP XhoI Rw	ATAATGCTCGAGAGAGTCGCG
NpTANSacIFor	GTATGAGCTCCGGTAGAGTTGAACCAG
NpTANceruleanRev	CCTCGCCCTTGCTCACCATCTTCTATATATATTTTCTTTA
NpTANceruleanFor	TAAAGAAAATATATATAGAAGATGGTGAGCAAGGGCGAGG
CeruleanpEarleyRev	GGCCCGCGGTACCGTCCTTGTACAGCTCGTCCATGC
CeruleanpEarleyFor	GCATGGACGAGCTGTACAAGGACGGTACCGCGGGCC
pEarleyOCSPstIRev	CCATCTGCAGCTGCTGAGCCTCGACAT
AtExon1_1For	CTCAACTCAGATCTTCTCAAGGAAACG
At255AfterStopRev	GCATAGTGGTACCCTCAAATTACACC

471
472

References:

References:

Abe, T., Thitamadee, S. and Hashimoto, T. (2004). Microtubule defects and cell
morphogenesis in the lefty1lefty2 tubulin mutant of Arabidopsis thaliana. *Plant Cell Physiol.*
45, 211–220.

Alim, K., Hamant, O. and Boudaoud, A. (2012). Regulatory role of cell division rules on tissue
growth heterogeneity. *Front. Plant Sci.* **3**, 174.

Bellinger, M. A., Uyehara, A. N., Martinez, P., McCarthy, M. C. and Rasmussen, C. G. Cell
cortex microtubules contribute to division plane positioning during telophase in maize.

Buschmann, H. and Borchers, A. (2020). Handedness in plant cell expansion: a mutant
perspective on helical growth. *New Phytol.* **225**, 53–69.

Buschmann, H., Fabri, C. O., Hauptmann, M., Hutzler, P., Laux, T., Lloyd, C. W. and
Schäffner, A. R. (2004). Helical Growth of the Arabidopsis Mutant tortifolia1 Reveals a
Plant-Specific Microtubule-Associated Protein. *Curr. Biol.* **14**, 1515–1521.

Buschmann, H., Chan, J., Sanchez-Pulido, L., Andrade-Navarro, M. A., Doonan, J. H. and
Lloyd, C. W. (2006). Microtubule-associated AIR9 recognizes the cortical division site at
preprophase and cell-plate insertion. *Curr. Biol.* **16**, 1938–1943.

Buschmann, H., Hauptmann, M., Niessing, D., Lloyd, C. W. and Scha, A. R. (2009). Helical
Growth of the Arabidopsis Mutant tortifolia2 Does Not Depend on Cell Division Patterns but
Involves Handed Twisting of Isolated Cells. **21**, 2090–2106.

Buschmann, H., Dols, J., Kopischke, S., Pen, E. J., Andrade-navarro, M. A., Heinlein, M.,
Szymanski, D. B., Zachgo, S., Doonan, J. H. and Lloyd, C. W. (2015). Arabidopsis
KCBP interacts with AIR9 but stays in the cortical division zone throughout mitosis via its
MyTH4-FERM domain. 2033–2046.

Caño-Delgado, A., Penfield, S., Smith, C., Catley, M. and Bevan, M. (2003). Reduced
cellulose synthesis invokes lignification and defense responses in Arabidopsis thaliana.
Plant J. **34**, 351–362.

Chakrabortty, B., Willemsen, V., de Zeeuw, T., Liao, C.-Y., Weijers, D., Mulder, B. and
Scheres, B. (2018). A Plausible Microtubule-Based Mechanism for Cell Division
Orientation in Plant Embryogenesis. *Curr. Biol.* **28**, 3031–3043.e2.

Clough, S. J. and Bent, A. F. (1999). Floral dip : a simplified method for Agrobacterium-
mediated transformation of Arabidopsis thaliana. **16**, 735–743.

Cnops, G., Wang, X., Linstead, P., Van Montagu, M., Van Lijsebettens, M. and Dolan, L.
(2000). Tornado1 and tornado2 are required for the specification of radial and
circumferential pattern in the Arabidopsis root. *Development* **127**, 3385–3394.

Cosgrove, D. J. (2005). Growth of the plant cell wall. *Nat. Rev. Mol. Cell Biol.* **6**, 850–861.

Cosgrove, D. J. (2018). Diffuse Growth of Plant Cell Walls. *Plant Physiol.* **176**, 16–27.

Dixit, R. and Cyr, R. J. (2002). Spatio-temporal relationship between nuclear-envelope
breakdown and preprophase band disappearance in cultured tobacco cells. *Protoplasma*

219, 116–121.

Facette, M. R., Rasmussen, C. G. and Van Norman, J. M. (2018). A plane choice:
coordinating timing and orientation of cell division during plant development. *Curr. Opin.*
Plant Biol. **47**, 47–55.

Fache, V., Gaillard, J., Van Damme, D., Geelen, D., Neumann, E., Stoppin-Mellet, V. and
Vantard, M. (2010). Arabidopsis kinetochore fiber-associated MAP65-4 cross-links
microtubules and promotes microtubule bundle elongation. *Plant Cell* **22**, 3804–3815.

Frey, N., Klotz, J. and Nick, P. (2010). A kinesin with calponin-homology domain is involved in
premitotic nuclear migration. *J. Exp. Bot.* **61**, 3423–3437.

Furutani, I., Watanabe, Y., Prieto, R., Masukawa, M., Suzuki, K. and Naoi, K. (2000). The
SPIRAL genes are required for directional control of cell elongation in Arabidopsis thaliana.
4453, 4443–4453.

Goff, J. and Van Norman, J. M. (2021). Polarly localized receptor-like kinases PXC2 and IRK
act redundantly during Arabidopsis root development in the radial axis. *bioRxiv*
2021.02.11.429611.

Goldy, C., Pedroza-Garcia, J.-A., Breakfield, N., Cools, T., Vena, R., Benfey, P. N., De
Veylder, L., Palatnik, J. and Rodriguez, R. E. (2021). The Arabidopsis GRAS-type SCL28
transcription factor controls the mitotic cell cycle and division plane orientation. *Proc. Natl.*
Acad. Sci. U. S. A. **118**,.

Gonneau, M., Desprez, T., Martin, M., Doblas, V. G., Bacete, L., Miart, F., Sormani, R.,
Hématy, K., Renou, J., Landrein, B., et al. (2018). Receptor Kinase THESEUS1 Is a
Rapid Alkalinization Factor 34 Receptor in Arabidopsis. *Curr. Biol.* **28**, 2452–2458.e4.

Gu, Y. and Rasmussen, C. G. (2022). Cell biology of primary cell wall synthesis in plants. *Plant*
Cell **34**, 103–128.

Guo, X., Wang, L. and Dong, J. (2021). Establishing asymmetry: stomatal division and
differentiation in plants. *New Phytol.* **232**, 60–67.

Haga, N., Kobayashi, K., Suzuki, T., Maeo, K., Kubo, M., Ohtani, M., Mitsuda, N., Demura,
T., Nakamura, K., Jürgens, G., et al. (2011). Mutations in MYB3R1 and MYB3R4 cause
pleiotropic developmental defects and preferential down-regulation of multiple G2/M-
specific genes in Arabidopsis. *Plant Physiol.* **157**, 706–717.

Hamant, O. and Haswell, E. S. (2017). Life behind the wall: sensing mechanical cues in plants.
BMC Biol. **15**, 59.

Hamant, O., Heisler, M. G., Jönsson, H., Krupinski, P., Uyttewaal, M., Bokov, P., Corson,
F., Sahlin, P., Boudaoud, A., Meyerowitz, E. M., et al. (2008). Developmental patterning
by mechanical signals in Arabidopsis. *Science* **322**, 1650–1655.

Hashimoto, T. (2002). Molecular genetic analysis of left-right handedness in plants. *Philos.*
Trans. R. Soc. Lond. B Biol. Sci. **357**, 799–808.

Hashimoto, T. (2015). Microtubules in plants. *Arabidopsis Book* **13**, e0179.

Heisler, M. G., Hamant, O., Krupinski, P., Uyttewaal, M., Ohno, C., Jönsson, H., Traas, J.
and Meyerowitz, E. M. (2010). Alignment between PIN1 polarity and microtubule
orientation in the shoot apical meristem reveals a tight coupling between morphogenesis
and auxin transport. *PLoS Biol.* **8**, e1000516.

Hématy, K., Sado, P.-E., Van Tuinen, A., Rochange, S., Desnos, T., Balzergue, S., Pelletier,
S., Renou, J.-P. and Höfte, H. (2007). A Receptor-like Kinase Mediates the Response of
Arabidopsis Cells to the Inhibition of Cellulose Synthesis. *Current Biology* **17**, 922–931.

Herrmann, A., Livanos, P., Lipka, E., Gadeyne, A., Hauser, M.-T., Van Damme, D. and
Müller, S. (2018). Dual localized kinesin-12 POK2 plays multiple roles during cell division
and interacts with MAP65-3. *EMBO Rep.* **19**, e46085.

Ho, C.-M. K., Lee, Y.-R. J., Kiyama, L. D., Dinesh-Kumar, S. P. and Liu, B. (2012).
Arabidopsis microtubule-associated protein MAP65-3 cross-links antiparallel microtubules
toward their plus ends in the phragmoplast via its distinct C-terminal microtubule binding
domain. *Plant Cell* **24**, 2071–2085.

Ishida, T., Kaneko, Y., Iwano, M. and Hashimoto, T. (2007). Helical microtubule arrays in a
collection of twisting tubulin mutants of Arabidopsis thaliana. *Proc. Natl. Acad. Sci. U. S. A.*
104, 8544–8549.

Kimata, Y., Higaki, T., Kawashima, T., Kurihara, D., Sato, Y. and Yamada, T. (2016).
Cytoskeleton dynamics control the first asymmetric cell division in Arabidopsis zygote.

Kirik, V., Herrmann, U., Parupalli, C., Sedbrook, J. C., Ehrhardt, D. W. and Hülskamp, M.
(2007). CLASP localizes in two discrete patterns on cortical microtubules and is required for
cell morphogenesis and cell division in Arabidopsis. *J. Cell Sci.* **120**, 4416–4425.

Lee, Y.-R. J. and Liu, B. (2013). The rise and fall of the phragmoplast microtubule array. *Curr.*
Opin. Plant Biol. **16**, 757–763.

Li, S., Sun, T. and Ren, H. (2015). The functions of the cytoskeleton and associated proteins
during mitosis and cytokinesis in plant cells. *Front. Plant Sci.* **6**, 282.

Li, H., Sun, B., Sasabe, M., Deng, X., Machida, Y., Lin, H., Lee, Y. J. and Liu, B. (2017).
Arabidopsis MAP 65-4 plays a role in phragmoplast microtubule organization and marks the
cortical cell division site. *New Phytologist* **215**, 187–201.

Lipka, E., Gadeyne, A., Stöckle, D., Zimmermann, S., De Jaeger, G., Ehrhardt, D. W., Kirik,
V., Van Damme, D. and Müller, S. (2014). The Phragmoplast-Orienting Kinesin-12 Class
Proteins Translate the Positional Information of the Preprophase Band to Establish the
Cortical Division Zone in Arabidopsis thaliana. *Plant Cell* **26**, 2617–2632.

Livanos, P. and Müller, S. (2019). Division Plane Establishment and Cytokinesis. *Annu. Rev.*
Plant Biol.

Louveaux, M., Julien, J.-D., Mirabet, V., Boudaoud, A. and Hamant, O. (2016). Cell division
plane orientation based on tensile stress in Arabidopsis thaliana.

Lukowitz, W., Mayer, U. and Jürgens, G. (1996). Cytokinesis in the Arabidopsis embryo
involves the syntaxin-related KNOLLE gene product. *Cell* **84**, 61–71.

Martinez, P., Luo, A., Sylvester, A. and Rasmussen, C. G. (2017). Proper division plane
orientation and mitotic progression together allow normal growth of maize. *Proc. Natl. Acad.*
Sci. U. S. A. **114**, 2759–2764.

Martinez, P., Allsman, L. A., Brakke, K. A., Hoyt, C., Hayes, J., Liang, H., Neher, W., Rui,
Y., Roberts, A. M., Moradifam, A., et al. (2018). Predicting Division Planes of Three-
Dimensional Cells by Soap-Film Minimization. *Plant Cell* **30**, 2255–2266.

Martinez, P., Dixit, R., Balkunde, R. S., Zhang, A., O'Leary, S. E., Brakke, K. A. and
Rasmussen, C. G. (2020). TANGLED1 mediates microtubule interactions that may
promote division plane positioning in maize. *J. Cell Biol.* **219**,.

McMichael, C. M. and Bednarek, S. Y. (2013). Cytoskeletal and membrane dynamics during
higher plant cytokinesis. *New Phytol.* **197**, 1039–1057.

Menges, M., de Jager, S. M., Gruissem, W. and Murray, J. A. H. (2005). Global analysis of
the core cell cycle regulators of Arabidopsis identifies novel genes, reveals multiple and
highly specific profiles of expression and provides a coherent model for plant cell cycle
control. *Plant J.* **41**, 546–566.

Mir, R., Morris, V. H., Buschmann, H. and Rasmussen, C. G. (2018). Division Plane
Orientation Defects Revealed by a Synthetic Double Mutant Phenotype. *Plant Physiol.* **176**,
418–431.

Morgan, X., Zhao, Q., Rodrigo-peiris, T., Brkljacic, J., Sylvia, C., Mu, S. and Meier, I.
(2008). RanGAP1 is a continuous marker of the Arabidopsis cell division plane. **105**,
18637–18642.

Moukhtar, J., Trubuil, A., Belcram, K., Legland, D., Khadir, Z., Urbain, A., Palauqui, J.-C.
and Andrey, P. (2019). Cell geometry determines symmetric and asymmetric division
plane selection in Arabidopsis early embryos. *PLoS Comput. Biol.* **15**, e1006771.

Müller, S. and Jürgens, G. (2016). Plant cytokinesis—No ring, no constriction but centrifugal
construction of the partitioning membrane. *Semin. Cell Dev. Biol.* **53**, 10–18.

Murata, T., Sano, T., Sasabe, M., Nonaka, S., Higashiyama, T., Hasezawa, S., Machida, Y.
and Hasebe, M. (2013). Mechanism of microtubule array expansion in the cytokinetic
phragmoplast. *Nat. Commun.* **4**, 1967.

Muroyama, A. and Bergmann, D. (2019). Plant Cell Polarity: Creating Diversity from Inside the
Box. *Annu. Rev. Cell Dev. Biol.* **35**, 309–336.

Nakajima, K., Furutani, I., Tachimoto, H., Matsubara, H. and Hashimoto, T. (2004).
SPIRAL1 encodes a plant-specific microtubule-localized protein required for directional
control of rapidly expanding Arabidopsis cells. *Plant Cell* **16**, 1178–1190.

Pan, X., Fang, L., Liu, J., Senay-Aras, B., Lin, W., Zheng, S., Zhang, T., Guo, J., Manor, U.,
Van Norman, J., et al. (2020). Auxin-induced signaling protein nanoclustering contributes
to cell polarity formation. *Nat. Commun.* **11**, 3914.

Pickett-Heaps, J. D., Gunning, B. E. S., Brown, R. C., Lemmon, B. E. and Cleary, A. L.
(1999). The cytoplast concept in dividing plant cells: cytoplasmic domains and the evolution

of spatially organized cell division. *Am. J. Bot.* **86**, 153–172.

Rasmussen, C. G. and Bellinger, M. (2018). An overview of plant division-plane orientation.
New Phytol.

Rasmussen, C. G., Wright, A. J. and Müller, S. (2013). The role of the cytoskeleton and
associated proteins in determination of the plant cell division plane. *Plant J.* **75**, 258–269.

Sakai, T., Honing, H. van der, Nishioka, M., Uehara, Y., Takahashi, M., Fujisawa, N., Saji,
K., Seki, M., Shinozaki, K., Jones, M. A., et al. (2008). Armadillo repeat-containing
kinesins and a NIMA-related kinase are required for epidermal-cell morphogenesis in
Arabidopsis. *Plant J.* **53**, 157–171.

Sampathkumar, A., Krupinski, P., Wightman, R., Milani, P., Berquand, A., Boudaoud, A.,
Hamant, O., Jönsson, H. and Meyerowitz, E. M. (2014). Subcellular and supracellular
mechanical stress prescribes cytoskeleton behavior in Arabidopsis cotyledon pavement
cells. *Elife* **3**, e01967.

Sapala, A., Runions, A., Routier-Kierzkowska, A.-L., Das Gupta, M., Hong, L., Hofhuis, H.,
Verger, S., Mosca, G., Li, C.-B., Hay, A., et al. (2018). Why plants make puzzle cells, and
how their shape emerges. *Elife* **7**,.

Schopfer, P. (2006). Biomechanics of plant growth. *Am. J. Bot.* **93**, 1415–1425.

Sedbrook, J. C., Ehrhardt, D. W., Fisher, S. E., Scheible, W.-R. and Somerville, C. R.
(2004). The Arabidopsis SKU6/SPIRAL1 gene encodes a plus end--localized microtubule-
interacting protein involved in directional cell expansion. *Plant Cell* **16**, 1506–1520.

Shao, W. and Dong, J. (2016). Polarity in plant asymmetric cell division: Division orientation
and cell fate differentiation. *Dev. Biol.* **419**, 121–131.

Shoji, T., Narita, N. N., Hayashi, K., Hayashi, K., Asada, J., Hamada, T., Sonobe, S.,
Nakajima, K. and Hashimoto, T. (2004). Plant-specific microtubule-associated protein
SPIRAL2 is required for anisotropic growth in Arabidopsis. *Plant Physiol.* **136**, 3933–3944.

Smertenko, A. P., Kaloriti, D., Chang, H.-Y., Fiserova, J., Opatrny, Z. and Hussey, P. J.
(2008). The C-terminal variable region specifies the dynamic properties of Arabidopsis
microtubule-associated protein MAP65 isotypes. *Plant Cell* **20**, 3346–3358.

Smertenko, A., Assaad, F., Baluška, F., Bezanilla, M., Buschmann, H., Drakakaki, G.,
Hauser, M.-T., Janson, M., Mineyuki, Y., Moore, I., et al. (2017). Plant Cytokinesis:
Terminology for Structures and Processes. *Trends Cell Biol.* **27**, 885–894.

Smertenko, A., Hewitt, S. L., Jacques, C. N., Kacprzyk, R., Liu, Y., Marcec, M. J., Moyo, L.,
Ogden, A., Oung, H. M., Schmidt, S., et al. (2018). Phragmoplast microtubule dynamics -
a game of zones. *J. Cell Sci.* **131**, jcs203331.

Smith, L. G., Gerttula, S. M., Han, S. and Levy, J. (2001). Tangled1: a microtubule binding
protein required for the spatial control of cytokinesis in maize. *J. Cell Biol.* **152**, 231–236.

Song, H., Golovkin, M., Reddy, A. S. and Endow, S. A. (1997). In vitro motility of AtKCBP, a
calmodulin-binding kinesin protein of Arabidopsis. *Proc. Natl. Acad. Sci. U. S. A.* **94**, 322–
327.

Strompen, G., El Kasmi, F., Richter, S., Lukowitz, W., Assaad, F. F., Jürgens, G. and
Mayer, U. (2002). The Arabidopsis HINKEL gene encodes a kinesin-related protein
involved in cytokinesis and is expressed in a cell cycle-dependent manner. *Curr. Biol.* **12**,
153–158.

Thitamadee, S., Tuchihara, K. and Hashimoto, T. (2002). Microtubule basis for left-handed
helical growth in Arabidopsis. *Nature* **417**, 193–196.

Uyttewaal, M., Burian, A., Alim, K., Landrein, B., Borowska-Wykręt, D., Dedieu, A.,
Peaucelle, A., Ludynia, M., Traas, J., Boudaoud, A., et al. (2012). Mechanical stress
acts via katanin to amplify differences in growth rate between adjacent cells in Arabidopsis.
Cell **149**, 439–451.

Van Damme, D. (2009). Division plane determination during plant somatic cytokinesis. *Curr.*
Opin. Plant Biol. **12**, 745–751.

van Oostende-Triplet, C., Guillet, D., Triplet, T., Pandzic, E., Wiseman, P. W. and
Geitmann, A. (2017). Vesicle Dynamics during Plant Cell Cytokinesis Reveals Distinct
Developmental Phases. *Plant Physiol.* **174**, 1544–1558.

Völker, A., Stierhof, Y. D. and Jürgens, G. (2001). Cell cycle-independent expression of the
Arabidopsis cytokinesis-specific syntaxin KNOLLE results in mistargeting to the plasma
membrane and is not sufficient for cytokinesis. *J. Cell Sci.* **114**, 3001–3012.

Wachsman, G., Sparks, E. E. and Benfey, P. N. (2015). Genes and networks regulating root
anatomy and architecture. *New Phytol.* **208**, 26–38.

Wada, M. (2018). Nuclear movement and positioning in plant cells. *Semin. Cell Dev. Biol.* **82**,
17–24.

Walker, K. L., Müller, S., Moss, D., Ehrhardt, D. W. and Smith, L. G. (2007). Arabidopsis
TANGLED identifies the division plane throughout mitosis and cytokinesis. *Curr. Biol.* **17**,
1827–1836.

Wasteneys, G. O. and Collings, D. A. (2009). 3 Expanding beyond the great divide: the
cytoskeleton and axial growth. *Differentiation and Development* **10**, 83.

Whitewoods, C. D. and Coen, E. (2017). Growth and Development of Three-Dimensional Plant
Form. *Curr. Biol.* **27**, R910–R918.

Wick, S. M. and Duniec, J. (1983). Immunofluorescence microscopy of tubulin and microtubule
arrays in plant cells. I. Preprophase band development and concomitant appearance of
nuclear envelope-associated tubulin. *J. Cell Biol.* **97**, 235–243.

Wolf, S., van der Does, D., Ladwig, F., Sticht, C., Kolbeck, A., Schürholz, A.-K., Augustin,
S., Keinath, N., Rausch, T., Greiner, S., et al. (2014). A receptor-like protein mediates the
response to pectin modification by activating brassinosteroid signaling. *Proc. Natl. Acad.*
Sci. U. S. A. **111**, 15261–15266.

Wu, S.-Z. and Bezanilla, M. (2014). Myosin VIII associates with microtubule ends and together
with actin plays a role in guiding plant cell division. *Elife* **3**,.

Wu, S.-Z., Yamada, M., Mallett, D. R. and Bezanilla, M. (2018). Cytoskeletal discoveries in the

plant lineage using the moss Physcomitrella patens. *Biophys. Rev.*

Yang, W., Cortijo, S., Korsbo, N., Roszak, P., Schiessl, K., Gurzadyan, A., Wightman, R.,
Jönsson, H. and Meyerowitz, E. (2021). Molecular mechanism of cytokinin-activated cell
division in Arabidopsis. *Science* **371**, 1350–1355.

Yi, P. and Goshima, G. (2020). Rho of Plants GTPases and Cytoskeletal Elements Control
Nuclear Positioning and Asymmetric Cell Division during Physcomitrella patens Branching.
Curr. Biol. **30**, 2860–2868.e3.

Yi, P. and Goshima, G. (2022). Division site determination during asymmetric cell division in
plants. *Plant Cell*.

714