Minimal phrase composition revealed by intracranial recordings

Authors:

Elliot Murphy[1,2], Oscar Woolnough[1,2], Patrick S. Rollo[1,2], Zachary Roccaforte[1], Katrien Segaert[3,4], Peter Hagoort[4,5], Nitin Tandon[1,2,6,*]

1. Vivian L. Smith Department of Neurosurgery, McGovern Medical School, University of Texas Health Science Center at Houston, Houston, TX, USA

2. Texas Institute for Restorative Neurotechnologies, University of Texas Health Science Center at Houston, Houston, TX, USA

3. School of Psychology & Centre for Human Brain Health, University of Birmingham, Birmingham, UK

4. Max Planck Institute for Psycholinguistics, Nijmegen, The Netherlands

5. Donders Institute for Brain, Cognition and Behaviour, Nijmegen, The Netherlands

6. Memorial Hermann Hospital, Texas Medical Center, Houston, Texas, USA

*Corresponding author: Nitin Tandon (nitin.tandon@uth.tmc.edu)

Number of figures: 6 (+4 suppl.)

Abstract

The ability to comprehend phrases is an essential component of language. Here we evaluate
the neural processes that enable the transition from single word processing to a minimal
compositional scheme using intracranial recordings. 19 patients implanted with penetrating
depth or surface subdural intracranial electrodes heard auditory recordings of adjective-noun,
pseudoword-noun and adjective-pseudoword phrases and judged whether the phrase
matched a picture. Stimulus-dependent alterations in broadband gamma power activity (BGA),
low frequency power or phase consistency, and phase-locking values across the
compositional network were analyzed. The posterior superior temporal sulcus (pSTS) and
temporo-occipital junction (TOJ) revealed a fine-structured cortical mosaic, with closely
neighboring tissue displaying exclusive sensitivity to either lexicality or phrase structure, but
not both. During phrase composition, greater functional connectivity was seen between pSTS-
TOJ and both pars triangularis and temporal pole. These two regions also encode anticipation
of composition in broadband low frequencies. These results implicate pSTS-TOJ as a crucial
hub for the retrieval and computation of minimal phrases. Overall, this work reveals an
interface of sparsely interwoven coding for lower and higher level linguistic features, coupled
with large-scale network organization, with direct relevance to our understanding of cognitive
networks in the human brain.

Keywords: composition, electrocorticography, semantics, posterior temporal lobe,
anticipation, human, language

Introduction

How the brain integrates individual word meanings to comprehend the meaning of multi-word utterances is an issue that has vexed the cognitive neuroscience of language for decades. How the brain computes 'complex meaning' (Hagoort, 2020) from combinations of multiple lexical items is still largely a mystery. This linguistic compositional process – the combination of words into larger structures with new and complex meaning – is also referred to as "Merge" (Chomsky, 1995; Chomsky et al., 2019) or "Unification" (Hagoort, 2013).

The simplest form of studying complex meaning is by using minimal phrases such as in the *red-boat* paradigm, which focuses on simple combinations of two words (Bemis & Pylkkänen, 2011, 2013a; Bozic et al., 2015; Brennan & Pylkkänen, 2012; Flick et al., 2018; Flick & Pylkkänen, 2020; Pylkkänen, 2020; Westerlund et al., 2015; Westerlund & Pylkkänen, 2014), avoiding a range of confounds associated with more complex linguistic stimuli (Berwick, & Stabler, 2019). A "red boat" is interpreted as a boat which is red, and not a red object which hosts boat-related properties, with phrases delivering novel syntactic and conceptual formats (Leivada & Murphy, 2021; Murphy, 2015a; Pietroski, 2018). Experimental work using the *red-boat* paradigm implicates the left anterior temporal lobe (ATL), specifically the temporal pole (Antonucci et al., 2008; Bemis & Pylkkänen, 2011; Lambon Ralph et al., 2012; Pobric et al., 2010; Westerlund & Pylkkänen, 2014; Wilson et al., 2014; Zhang & Pylkkänen, 2018), and posterior temporal regions (Flick & Pylkkänen, 2020; Matchin & Hickok, 2020) as crucial nodes for phrase composition. *Red-boat* experiments isolate semantic composition, which in turn encompasses syntactic composition. These studies report responses that vary in timing (between 180–350ms post-composition) and duration (50–100ms). The left posterior temporal lobe and angular gyrus (AG) also show greater activity for sentences than word lists, and for phrases than words (Brennan et al., 2016; Matchin & Hickok, 2020), making them candidate regions for the retrieval of phrasal templates (Hagoort, 2003, 2017),

Phrase composition is a rapid process that is likely dependent on finely organized sets of distributed cortical substrates. Previous work in the *red-boat* paradigm has been limited by spatiotemporal resolution. Using recordings from intracranial electroencephalography (iEEG) with depth electrodes penetrating grey matter or electrodes located on the cortical surface, we conducted a study of minimal phrase composition with auditory presentations of adjective-noun phrases across a large patient cohort (n = 19).

We utilized grammatical noun phrases ("red boat") and pseudoword phrases with variable positions of the pseudoword ("bleeg boat", "red fulg") to isolate semantic compositional processing. An adjective-pseudoword condition ("red fulg") was included to isolate semantic compositionality and reduce predictability. Given the large number of patients in our study, we

were able to analyze not only regions of interest delineated by previous studies, but also a
range of temporal, frontal and parietal cortices not previously implicated in models of phrase
composition.

Studies using scalp electroencephalography and magnetoencephalography (MEG) were used
to constrain our analytic space. iEEG provides uniquely high spatiotemporal resolution
recordings and is less susceptible to artifacts (e.g., muscle movements) (Arya, 2019; Flinker
et al., 2011). High frequency gamma changes (Conner et al., 2019; Forseth et al., 2018;
Johnson et al., 2020; Leszczyński et al., 2020; Towle et al., 2008; Yi et al., 2019) typically
index local cortical processing and are implicated in a range of cognitive processes (Buzsáki
& Watson, 2012; Hovsepyan et al., 2020; Jensen et al., 2019; Marko et al., 2019; Packard et
al., 2020; Prystauka & Lewis, 2019). An early anterior negative deflection immediately
preceding the critical noun in combinatorial contexts (-50–100ms) (Neufeld et al., 2016) likely
indexes syntactic prediction (Lau et al., 2006). Low frequency power increases have also been
noted during the anticipatory window (Bastiaansen & Hagoort, 2015; Lewis et al., 2016;
Segaert et al., 2018), during phrase composition (Segaert et al., 2018) and in a variety of
auditory phrase and sentence processing paradigms (Ding et al., 2016; Keitel et al., 2018;
Keitel et al., 2017; Mai et al., 2016). Therefore, we also evaluated the role of low frequencies
during anticipatory composition. Lastly, our task required patients to determine whether the
words they heard matched a subsequent image, enabling analyses related to phrase-matched
and phrase-contrasted contexts.

Results

Patients were presented with auditory recordings grouped randomly into three conditions:
Adjective-Noun, Pseudoword-Noun, Adjective-Pseudoword. A subsequent colored image was
presented, and patients were tasked with responding, with a button press, to whether the
image matched the phrase or not (Fig. 1A). Across the cohort, we had good coverage over
lateral and medial temporal lobe, inferior parietal lobe and inferior frontal regions, with some
coverage reaching into other portions of fronto-parietal regions (Fig. 1B). We saw activation
in response to the auditory stimuli most prominently in superior and middle temporal regions
(Fig. 1C). Below we report results pertaining to lexicality; phrase anticipation; phrase
composition; and linguistic-visual unification.

<u>**Figure 1**</u>: **Experimental paradigm with coverage and activation maps.** (A) Experimental

design. Average stimuli length: adjectives (420 ± 39 ms; mean ± SD), nouns (450 ± 75 ms),

pseudowords (430 ± 38 ms). (B) Group coverage map of included, left hemisphere electrodes,

plotted on a semi-inflated standardised N27 surface. (C) Broadband gamma activity (BGA)

increases from baseline (-500 to -100ms prior to first word) for all conditions from 100 to 400ms

after onset of the first word (threshold: %BGA > 5%, t > 1.96, patient coverage ≥ 3; *p* < 0.01

corrected). Black surfaces fell below patient coverage threshold.

Behavioral performance

Performance in the image matching task was highly accurate 97 ± 3% (311 ± 11/320 trials)

with an average response time of 1599 ± 539 ms. Only correct trials were analyzed further.

Effects of lexicality

To disentangle single word semantic effects from those of combinatorial semantics, we probed

the difference in representation of words vs. pseudowords. We generated a surface-based,

population-level map of cortical activity using a surface-based mixed-effects multi-level

analysis (SB-MEMA) (Conner et al., 2011; Fischl et al., 1999; Kadipasaoglu et al., 2014;

Kadipasaoglu et al., 2015), a method specifically designed to account for sampling variations

in iEEG and minimize effects of outliers. An SB-MEMA contrasting adjectives vs. pseudowords

in word position 1 and nouns vs. pseudowords in word position 2 (Fig. 2A) revealed

significantly greater broadband gamma activity (BGA; 70–150 Hz) for pseudowords than

words in posterior superior temporal gyrus (pSTG), posterior superior temporal sulcus (pSTS),

temporo-occipital junction (TOJ) and pars triangularis 300–700ms after word onset, indexing

the additional processing of unfamiliar word sounds in an attempt to derive potential lexicality.

The greater BGA increase for pseudowords than words at position 2 than position 1 around

pars triangularis and surrounding frontal areas (Fig. 2C), suggests involvement of this region
in effortful lexical processing (perhaps in phonological space) to facilitate semantic unification
(Hagoort, 2005, 2013; Hagoort & Indefrey, 2014).

Figure 2: **Grouped analysis for lexicality.** (A) SB-MEMA comparing words vs pseudowords,
red colors indexes greater BGA (70–150 Hz) for pseudowords and blue colors for words
(threshold: %BGA > 5%, t > 1.96, patient coverage ≥ 3; $p < 0.01$ corrected). Top row: word
position 1; bottom row: word position 2. (B) Exemplar electrodes for the words vs pseudowords
analysis (red: pseudowords; blue: words). Error bars (colored shading) set at one standard
error. Sylvian fissure is marked with a yellow line for reference on each surface. (C) SB-MEMA
indicating BGA increases for pseudowords at the second word position relative to
pseudowords at the first word position. (D) Electrodes with significant differences in inter-trial
phase coherence (ITC) for words vs pseudowords, in posterior temporal cortex (9 patients).
(E) ITC in alpha (8–12 Hz) at all electrodes in (D) for words (blue) and pseudowords (red) on
the left. T0 = first word onset.

Additionally, to broaden our assessment of the contributions of lower frequencies, we
computed inter-trial phase coherence (ITC) across regions and low frequencies. We found
significantly greater ITC values for words relative to pseudowords in and around pSTG (Fig.
2D, E; Suppl. Fig. 1) in the alpha band (8–12 Hz). No other ROIs (ATL, pars triangularis, pars
orbitalis, MTG, supramarginal gyrus: all FDR-corrected q-values >0.05) displayed such effects
for this, or for phrase composition. Alpha phase coherence in pSTG may therefore encode
information relevant to lexical interpretation.

Compositional anticipation

We next contrasted Adjective-Noun and Pseudoword-Noun conditions at the onset of second
word presentation, with only the former condition licensing any phrasal anticipation. During
the anticipatory window for phrase formation (from -200ms to 0ms prior to the second word
onset), low frequency power (8–30 Hz) exhibited an anticipatory effect (Fig. 3). This
anticipatory effect was found in IFG, pSTG and ATL, which all appear to be involved in
syntactic-semantic prediction generation. In addition, these anticipatory effects were unrelated
to BGA, and such widespread, regional effects were not found in delta or theta. For
comparison, we also plot the anticipatory window in BGA for -100 to 0ms (Fig. 3B), which
exhibited no relation with the lower frequency effects (compare also Fig. 3D with Fig. 3E).

Figure 3: **Syntactic-semantic compositional anticipation.** (A) SB-MEMA for the anticipatory time window centred around second word onset (-200–0ms) for low frequency power. Pink: Greater power for composition anticipation. Green: Greater for no anticipation (i.e. after having heard a Pseudoword at first word position). (B) SB-MEMA for BGA for the same contrast for -100–0ms prior to second word onset. (C)-(E) Exemplar electrodes with location (C), low frequency power traces and (D) BGA traces in dashed green boxes (E), across three patients sorted by row.

Phrase composition

The full combinatorial contrast (Adjective-Noun vs [Adjective-Pseudoword + Pseudoword-Noun]), revealed greater BGA for portions of pSTS-TOJ during phrase composition than non-composition (Fig. 4A). The specific onset of this effect was around 210ms after noun onset, with peak BGA at around 300ms (Suppl. Fig. 3). The same region exhibited greater BGA for non-composition at later time points, implicating it in effortful phrase structure derivations.

Certain portions of pSTS-TOJ across patients displayed exclusive sensitivity to phrase
composition, and not lexicality (Fig. 4B). Other regions – pSTG and inferior frontal gyrus (IFG)
– did not show any significant BGA differences between these conditions (Suppl. Fig. 2). Since
previous combinatorial effects in phrasal and sentential processing have also localized to
anterior temporal pole, we also focused on this region (Suppl. Fig. 2, 4). These portions of left
anterior temporal lobe (temporal pole) exhibited increased broadband low frequency activity
in response to phrases from approximately 380–460ms and 700–900ms after noun
presentation. This activity was unrelated to BGA, indicating that low frequency activity in ATL
is an independent component of phrase structure comprehension; likely late-stage semantic
processing. The ERPs were dissociable from BGA across the ROIs plotted (Suppl. Fig. 2): A
late effect for Non-Composition was found in Broca's area, likely due to greater attempted
lexical access for the pseudowords, while a late signature was detected in temporal pole for
Composition, potentially related to late-stage conceptual access.

<u>Figure 4</u>: **Grouped analysis for phrase composition.** (A) SB-MEMAs for the phrase
composition analysis for BGA. (B) Exemplar electrodes with FDR-corrected (one-tailed t-tests,
q<0.05) significance bars in purple plotted in native patient space. Includes compositional

contrast and lexicality contrast, dissociating neighbouring portions of pSTS-TOJ that
responded exclusively for phrase composition and not lexicality (orange dots = composition
effect; green dots = lexicality effect). Error bars set at one standard error. Numbers denote
distinct patients. (C) Electrodes exhibiting a significant BGA contrast between Compositional
and Non-Compositional trials. 39 electrodes (orange) from 12 patients exhibited an FDR-
significant contrast (one-tailed t-tests, q<0.05) for compositionality at some point between 0-
1000ms after the second word; 97 electrodes across 17 patients for lexicality; 3 electrodes for
both across 3 patients.

Next, we isolated regions of interest to derive cortical functional connectivity during phrase
composition. These were based either on results from our main analysis (pSTS-TOJ, temporal
pole) or on composition effects described in the literature (inferior frontal regions). To
characterize functional connectivity between these regions during phrase composition, we
computed phase-locking values (PLV) for electrode pairs situated within pSTS-TOJ with either
pars triangularis or anterior temporal lobe. We computed the generalized phase of the
wideband filtered (3–50 Hz) signal that has previously been shown to be more effective than
the use of narrowband alpha or theta filters (Davis et al., 2020).

Among patients with concurrent coverage in pSTS-TOJ and pars triangularis (n = 8, electrode
pairs = 231), the majority (n = 5) exhibited significantly greater PLVs for Compositional than
for Non-Compositional trials during the 0–500ms period after second word onset, averaging
across PLV values for each pair. In patients with joint coverage in pSTS-TOJ and anterior
temporal lobe (temporal pole) (n = 8, electrode pairs = 274), the majority (n = 6) showed
greater PLVs for the same contrast during the same time window. We plot the magnitude of
phase-locking values across these regions and in two exemplar patients (Fig. 5).

Figure 5: Phase-locking between semantic composition regions of interest. (A) Left: Average generalized phase-locking values (gPLV) for 5 patients showing greater gPLV for phrase composition relative to non-composition between pSTS-TOJ (HCP index: TPOJ1) and pars triangularis (HCP index: 45, IFSa, IFSp, 47l). Right: The same plot but for the 6 patients showing greater gPLV between pSTS-TOJ and temporal pole (HCP index: TGd). (B) Posterior temporal lobe (pSTS-TOJ) gPLVs with inferior frontal gyrus (specifically, pars triangularis) (Top) and anterior temporal lobe (specifically, temporal pole) (Bottom). Left plots show the localization in native space of electrodes significantly involved (q < 0.05) in inter-regional phase-locking (3–50 Hz). Right plots show average time courses (mean ± SEM) of phase-locking value changes from baseline in phrase composition (brown) and non-composition (blue) trials.

Integration of linguistic and visual information

Following each phrase, patients were presented with a matching or non-matching visual representation of the phrase. Comparing the Adjective-Noun trials which contained a matching picture, or a non-matching picture, during the 250–500ms post-picture window, SB-MEMAs revealed two notable effects (Fig. 6): Anterior insula exhibited greater BGA for phrase-picture

matches, while more dorsal frontal regions, centered around inferior frontal sulcus, exhibited greater BGA for phrase-picture non-matches. Thus, these regions appear to be implicated in linguistic-visual integration and monitoring, likely anchored around the semantic content denoted by both phrases and images and their successful association via some conjunction operator (i.e. [*red ∧ boat*], rather than a monadic adjectival or nominal representation). These two regions flank Broca's area, subserving distinct composition-related goals. Since activity of opercular regions can be misattributed to the insula (Woolnough et al., 2019), we ensured that these effects across patients specifically came from electrodes in insula proper, rather than operculum or Broca's area.

Figure 6: **Grouped analysis for linguistic-visual integration.** (A) SB-MEMA in BGA for phrase-picture match (orange) and phrase-picture contrast (turqoise) increases, 250–500ms after picture onset. (B) Exemplar insula electrode.

Discussion

We localized the neural signatures of combinatorial phrase structure comprehension using minimal adjective-noun phrases in a large patient cohort, with both penetrating depth and surface grid intracranial electrodes. We identified a broad portion of posterior temporal cortex as being sensitive to lexicality, and then isolated closely neighboring portions of pSTS-TOJ within patients that responded exclusively to phrase composition, and not lexicality, and vice versa. This computational contrast within a narrow strip of pSTS-TOJ provides a finely

organized cortical mosaic for the language system (Amunts & Zilles, 2015), going beyond
more traditional, broader structure-function mappings for higher-order syntax-semantics. Our
results can contribute to the development of a micro-map of pSTS-TOJ with respect to the
demands of computational complexity; a necessary step towards understanding the
organizational principles of the brain at distinct spatial scales, including ultimately cellular
components.

Phrase composition was indexed by two main components: Greater BGA in portions of pSTS-
TOJ at an early window (around 210-300ms); greater low frequency power in temporal pole
at 380–460ms and 700–900ms after the presentation of a phrase-licensing nominal. Greater
functional connectivity for phrase composition was found between these regions (pSTS-TOJ
and temporal pole) and also between pSTS-TOJ and pars triangularis. pSTS-TOJ is initially
active for phrase composition and subsequently shows greater local cortical activity for non-
compositional stimuli, indicating its involvement in phrase-related processing. In addition,
pSTS-TOJ and pSTG encode local processing of pseudowords, with pSTG likely coding
speech signals into articulatory representations (Martin et al., 2019). Anterior portions of IFG
exhibit greater BGA for pseudowords at the second word position, possibly indexing increased
unification demands with the preceding adjective, compared to a real word (Hagoort, 2005,
2013; Hagoort & Indefrey, 2014). This effect may hence index the role of this region in the
syntactic and semantic unification of more or less expected continuations, in agreement with
earlier results in a sentence context (Hagoort, 2004).

In addition, STG, ATL and IFG index composition anticipation in low frequency power,
consistent with a predictive neural mechanism for higher-order linguistic information
(Bastiaansen & Hagoort, 2015; Benítez-Burraco & Murphy, 2019; Lewis et al., 2016; Murphy,
2015b; Prystauka & Lewis, 2019). These anticipatory effects were found in the context of our
stimuli being controled for predictability (the addition of the Adjective-Pseudoword condition is
the simplest way to achieve this). We additionally showed that alpha phase encodes lexical
information exclusively in pSTG via ITC (and not phrase-relevant information).

Lastly, anterior insula and inferior frontal sulcus (IFS) subserve the integration of language
input with the visual referent. Both of these regions have been argued to be the convergence
zones of the ventral (anterior insula) and dorsal (IFS) attentional networks (Cazzoli et al.,
2021), while task-evoked pupillary dynamics have been linked to the insula (Kucyi & Parvizi,
2020); given current task demands, these dynamics may also underscore our findings.

Lexicality

Our results replicate prior studies in which BGA localized to p/mSTG and pSTS tracks lexicality
(Canolty et al., 2007; Humphries et al., 2006; Tanji et al., 2005). In addition, we show

recruitment of anterior IFG in this process. Further, pSTG alpha phase coherence may index
the coordination of smaller-scale ensembles which may bundle sub-lexical features to yield
coherent word-level interpretations of auditory stimuli, in accord with a recent
neurocomputational model for lexical feature-set construction (Murphy, 2020).

Anticipatory response

According to a range of parsing models (Berwick et al., 2019), adjective-noun syntax is
constructed predictively, in anticipation of the noun. Neural response differences in the period
preceding the noun in combinatorial trials would likely reflect the generation of this structure.
For instance, particular anticipatory amplitude increases have been isolated to the alpha (8–
12 Hz) and beta (15–30 Hz) bands in fronto-central and fronto-temporal scalp EEG electrodes,
and have been related to the pre-activation of linguistic structure/information (Gastaldon et al.,
2020; Segaert et al., 2018). Based on our results, low frequency power in IFG and ATL may
be preparing a syntactic slot to be filled by predicted upcoming nominal content, or initiating
top-down local coordination of subsequent composition-indexing BGA in pSTS-TOJ. These
low frequency dynamics might not even pertain to lexico-semantic prediction specifically, but
rather anticipatory stages pertaining to phrasal initiation (Terporten et al., 2019). Our findings
are also potentially in line with the notion that beta oscillations can index the construction and
maintenance of sentence-level meaning (scaling up from minimal phrases) (Lewis et al.,
2016), and the claims that beta can index aspects of syntactic anticipation and phrasal
category generation (Benítez-Burraco & Murphy, 2019; Murphy, 2020).

Our results may relate to increased recruitment (Bonnefond & Jensen, 2013) of pSTS in late
stage lexical processing effort (-200–0ms before the noun). The portion of temporal pole and
MTG exhibiting low frequency increases during the anticipatory window could be also involved
in late-stage lexico-conceptual processing of adjectival representations, although given the
relatively late post-adjective window it is more likely to be related to anticipation. Given
increasing evidence for IFG involvement in linear morpho-syntactic processing (Matchin &
Hickok, 2020), it is also possible that effects in IFG index a form of lexical anticipation.
Predictable word processing specifically pinpoints anterior IFG and ATL in preceding cortical
activations (Pulvermüller & Grisoni, 2020). During basic semantic composition, anterior IFG
activation has been shown to increase with the amount of lexical information being integrated
into the structure (Schell et al., 2017), again pointing to a role for this region as a lexical
memory buffer and/or the locus of higher-order predictions.

Minimal phrase composition

We found that neighbouring portions of pSTS-TOJ across patients can exclusively code either
for lexicality or phrase composition, helping to isolate the core compositional operation in

phrase structure generation. While a large area of posterior and superior temporal cortex was sensitive to lexicality, a narrower portion was recruited exclusively to code for phrase structure, generating from simple lexical meanings the more computationally demanding structures of 'complex meaning' (Hagoort, 2020; Pietroski, 2018).

One of the main candidates for phrase composition from previous research, ATL, was implicated via low frequency activity (specifically, temporal pole activity). This may indicate that temporal pole is involved in compositional processing within a noun phrase, with low frequency activity possibly coordinating broader areas in the service of constructing a coherent semantic referent (Wang et al., 2020). Despite ATL being the "most consistent integrative node" (Pylkkänen & Brennan, 2020) in the syntax-semantics combinatorial network, the high spatiotemporal resolution of iEEG presented another candidate for composition: pSTS-TOJ. Increasing morpho-orthographic complexity in a similar minimal phrasal paradigm does not implicate ATL in early processing windows, but seems to delay ATL activity until after intra-word morphological composition (Flick et al., 2018). This suggests that there must be regions independent of semantic and orthographic processing that are common to all forms of (early) phrasal composition. Our findings suggest that pSTS-TOJ might be one such region, although since we only presented auditory stimuli we cannot make any stronger claims. Comparing the discovered composition effect in BGA (Fig. 4A) with the auditory stimuli activation map (Fig. 1C) suggests a clear overlap between sites of initial sensitivity to auditory stimuli and composition-sensitive sites. On the other hand, while there is indeed overlap, there is also a portion of posterior TOJ which seems uniquely active for composition but which is not prominently sensitive to audition, which may point to the involvement of a more modality-independent compositional process. Since MEG cannot detect high frequency activity deep in sulci (Irimia et al., 2012), our iEEG investigation was able to provide new insights here.

With respect to activation timings, we found greater BGA for phrase composition in pSTS-TOJ at 210ms, diminishing around 300ms, after which the majority of responses were greater for non-compositional trials. It is possible that composition-specific computations initiated earlier than this, and that it was only during these time windows that condition differences arose. The later, greater responses for non-composition could be due to more efficient encoding of phrases (hence lower BGA), or, more likely given the latency, late-stage compositional effort/evaluation. It is also possible that the low frequency responses we found to composition in ATL are responses to post-syntactic (pSTS-TOJ) processing, leading to conceptual integration. The timing of these late wrap-up effects might be due to the slow nature of these frequencies and could also be due to some form of traveling oscillatory component (Muller et al., 2018) from posterior temporal areas.

Our findings are in line with recent models implicating these posterior portions of temporal
lobe in phrase structure building (Flick & Pylkkänen, 2020; Law & Pylkkänen, 2021; Matar et
al., 2021; Matchin & Hickok, 2020; Murphy, 2018, 2020); a recent meta-analysis of sign
language implicating TOJ and pMTG/pSTS in comprehension (Trettenbrein et al., 2020); and
the finding that cross-linguistic (French and Chinese) reading competence is linked to pSTS
(Feng et al., 2020), contributing to the neural localization of complex composition (Chomsky
et al., 2019; Pietroski, 2018). In particular, evidence from neuroimaging and lesion-deficit
mapping indicates a role for pMTG in syntactic comprehension and production (Matchin &
Hickok, 2020), while pSTS has been implicated via iEEG in hierarchical syntactic
computations (Nelson et al., 2017). Data from brain stimulation implicate postero-middle
temporal areas in syntactic processing (Duffau et al., 2014), and a recent *red-boat* study in
fMRI implicated a broad portion of pMTG in meaningful phrase composition (Graessner et al.,
2021). Impairments in grammaticality judgments have primarily been linked to posterior
temporal lobe damage (Wilson & Saygin, 2004). Lastly, widespread monotonic fronto-
temporal language network gamma power increases over the course of sentence reading
have been documented in iEEG (Fedorenko et al., 2016); our findings isolate a more specific
site in the network responsible for the core compositional operator underlying sentence
comprehension.

We also discovered greater functional connectivity between pSTS-TOJ and both temporal
pole and pars triangularis after the presentation of the phrase-licensing nominal (Fig. 5). White
matter atlas consultation reveals that the subcortical distribution of responses to semantic
stimuli reflects the continuous and homogenous ventral course of functional information in the
deep white matter of the temporal lobe, from TOJ, ascending to white matter underneath the
posterior two-thirds of IFG (Sarubbo et al., 2020). These ventral workflow pathways subserve
our regions of interest (temporal pole; TOJ; IFG). In addition, arcuate fasciculus is classically
implicated in connectivity between pSTS-TOJ and pars triangularis (Figley et al., 2017;
Glasser & Rilling, 2008). In relation to current findings and frameworks, it is possible that the
formation of phrase structure in pSTS-TOJ feeds categorial information (e.g. phrase type) to
interpretation interfaces in ATL and memory/control interfaces in pars triangularis (a form of
"spell-out"). One such piece of information encoded via PLV would be the phrase
category/label (Adger, 2012; Murphy, 2015a; Murphy & Shim, 2020). These results may point
to a degree of top-down control from pars triangularis on initial phrase construction in pSTS-
TOJ (Hagoort et al., 2009). The more precise computational role of such connectivity metrics
requires further empirical, and conceptual, advances.

Inhibition of a portion of left posterior temporal lobe (pSTS, pSTG and temporo-parietal
junction) selectively impairs the ability to produce appropriate functional morphemes for

particular hierarchical relations, but does not effect any other sub-linguistic process such as semantic or lexical retrieval (Lee et al., 2018). Our data are compatible with the idea that in connection with other areas, such as IFG (Baggio & Hagoort, 2011; Zaccarella et al., 2021), pSTS-TOJ contributes to both morphological and phrasal unification (Jackendoff & Audring, 2020; Rizzi, 2016). Our results are concordant with a division between sensitivity to non-local dependency structure (in left anterior temporal pole and IFG) and sensitivity to phrase structure (in left pSTG) (Lopopolo et al., 2021). Data from sEEG comparing noun phrases and acoustically-matched, identical words acting as verb phrases (with the latter demanding richer syntactic structure) reveals greater high gamma activity in MTG and STG in the coding of complex syntax (Artoni et al., 2020). This points to partial spatial overlap with our composition effects, in compliance with the idea that the re-application of recursive hierarchical structure generation calls upon the same basic compositional operator (i.e., that the computation implicated in forming verb phrases is the same operation that forms "red boat").

While we did not find composition effects in inferior parietal lobule (in particular, angular gyrus), and had good electrode coverage here, this may be due to this region's role in higher-order semantic processing (e.g. thematic role assignment, discourse structure) and event structure (Boylan et al., 2015). For instance, there is a range of evidence, including from intracranial recordings, that taxonomic conceptual relationships are encoded in ATL, while thematic relationships are encoded in angular gyrus (Davis & Yee, 2019; Thye et al., 2021).

Given the nature of our stimuli, we can make direct claims about semantic composition, and we suspect that portions of pSTS-TOJ are also implicated in minimal syntax, however the dynamics of the syntax network may differ from that of the semantic compositional network. Overall, our results contribute to isolating the neural signatures of basic phrase structure composition, an elementary computational procedure of natural language and potentially a species-defining property (Friederici et al., 2017; Murphy, 2019; Murphy et al., 2021), and we hope that future intracranial research can expand on our findings.

Materials and Methods

Participants: 19 patients (10 male, 18-41 years, IQ 97 ± 12, 2 left-handed) participated in the experiment after written informed consent was obtained. All were native English speakers. All experimental procedures were reviewed and approved by the Committee for the Protection of Human Subjects (CPHS) of the University of Texas Health Science Center at Houston as Protocol Number HSC-MS-06-0385.

Electrode Implantation and Data Recording: Data were acquired from either subdural grid electrodes (SDEs; 6 patients) or stereotactically placed depth electrodes (sEEGs; 12 patients) (Fig. 1B). SDEs were subdural platinum-iridium electrodes embedded in a silicone elastomer sheet (PMT Corporation; top-hat design; 3mm diameter cortical contact), and were surgically implanted via a craniotomy (Conner et al., 2011; Pieters et al., 2013; Tandon, 2012; Tong et al., 2020). sEEGs were implanted using a Robotic Surgical Assistant (ROSA; Medtech, Montpellier, France) (Rollo et al., 2020; Tandon et al., 2019). Each sEEG probe (PMT corporation, Chanhassen, Minnesota) was 0.8mm in diameter and had 8-16 electrode contacts. Each contact was a platinum-iridium cylinder, 2.0mm in length and separated from the adjacent contact by 1.5-2.43mm. Each patient had 12-20 such probes implanted. Following surgical implantation, electrodes were localized by co-registration of pre-operative anatomical 3T MRI and post-operative CT scans in AFNI (Cox, 1996). Electrode positions were projected onto a cortical surface model generated in FreeSurfer (Dale et al., 1999), and displayed on the cortical surface model for visualization (Pieters et al., 2013). Intracranial data were collected during research experiments starting on the first day after electrode implantation for sEEGs and two days after implantation for SDEs. Data were digitized at 2 kHz using the NeuroPort recording system (Blackrock Microsystems, Salt Lake City, Utah), imported into Matlab, initially referenced to the white matter channel used as a reference for the clinical acquisition system and visually inspected for line noise, artifacts and epileptic activity. Electrodes with excessive line noise or localized to sites of seizure onset were excluded. Each electrode was re-referenced to the common average of the remaining channels. Trials contaminated by inter-ictal epileptic spikes, saccade artefacts and trials in which participants responded incorrectly were discarded.

Stimuli and Experimental Design: Participants were presented with two-word auditory phrases, grouped by three conditions: Adjective-Noun ("red boat"), Adjective-Pseudoword ("red neub"), Pseudoword-Noun ("zuik boat"). Since these pseudoword phrases include phonologically viable nonwords, differences in the second position of the phrase between these items and the grammatical noun phrases are likely a result of compositional processing. While most previous studies have presented a licensable noun, our inclusion of the Adjective-Pseudoword condition further isolates composition and reduces predictability. To ensure attention was maintained, after each trial participants were shown a colored drawing and asked to press a button indicating whether the picture matched the phrase they had just heard. Participants were told to respond positively only when the picture fully matched the phrase (e.g. the phrase "red boat" followed by a picture of a red car, or a blue boat, would be negative). Auditory and visual stimuli were recruited from previous *red-boat* experiments (Bemis & Pylkkänen, 2013b).

A fixation cross was presented in the centre of the screen for 700ms followed by the first word
and 800ms later the second word was presented. 1600ms after the onset of the second word
the picture was presented, and 1400ms after picture presentation participants were prompted
to respond (Fig. 1A). Following their response, a blank screen was shown for 1500ms. Stimuli
were presented in a pseudorandom order, with no repetition amongst items. The number of
trials per block across the full experiment was as follows: Adjective-Noun (80), Pseudoword-
Noun (40), Adjective-Pseudoword (40). All patients undertook 2 blocks. Half of the Adjective-
Noun trials matched the picture presented (i.e. "red boat" was heard by the patient, and a
picture of a red boat was then presented), and the other half did not match. 6 adjectives were
used: black, blue, brown, green, pink, red (length M: 4.3, SD: 0.7; SUBTLEXus log-frequency
3.64). 20 nouns were used: bag, bell, boat, bone, cane, cross, cup, disc, flag, fork, hand, heart,
house, key, lamp, leaf, lock, plane, shoe, star (length M: 4.0, SD: 0.6; log-frequency 3.38)
(Brysbaert, New, & Keuleers, 2012). 6 pseudowords were used: beeg, cresp, kleg, nar, neub,
zuik (length M: 4.0, SD: 0.6). Average stimuli length: Adjectives (420 ms), Nouns (450 ms),
Pseudowords (430ms).

Stimuli were presented using Psychtoolbox (Kleiner et al., 2007) on a 15.4" LCD screen
positioned at eye-level, 2-3' from the patient. Auditory stimuli were presented using stereo
speakers (44.1 kHz, MacBook Pro 2015).

Signal Analysis: A total of 3458 electrode contacts were implanted in these patients; 2135 of
these were included for analysis after excluding channels proximal to the seizure onset zone
or exhibiting excessive inter-ictal spikes or line noise. Analyses were performed by first
bandpass filtering the raw data of each electrode into broadband gamma activity (BGA; 70–
150 Hz) following removal of line noise and its harmonics (zero-phase 2nd order Butterworth
band-stop filters). Electrodes were also visually inspected for saccade artefacts. A frequency
domain bandpass Hilbert transform (paired sigmoid flanks with half-width 1.5 Hz) was applied
and the analytic amplitude was smoothed (Savitzky-Golay FIR, 3rd order, frame length of
251ms; Matlab 2019b, Mathworks, Natick, MA). BGA was defined as percentage change from
baseline level; -500 to -100ms before the presentation of the first word in each trial. Periods
of significant activation were tested using a one-tailed t-test at each time point and were
corrected for multiple comparisons with a Benjamini-Hochberg false detection rate (FDR)
threshold of q<0.05. For the grouped analysis, all electrodes were averaged within each
subject and then the between-subject averages were used.

To provide statistically robust and topologically precise estimates of BGA, and to account for
variations in sampling density, population-level representations were created using surface-
based mixed-effects multilevel analysis (SB-MEMA) (Conner et al., 2011; Fischl et al., 1999;

Kadipasaoglu et al., 2014; Kadipasaoglu et al., 2015). This method accounts for sparse
sampling, outlier inferences, as well as intra- and inter-subject variability to produce population
maps of cortical activity. A geodesic Gaussian smoothing filter (3mm full-width at half-
maximum) was applied. Significance levels were computed at a corrected alpha-level of 0.01
using family-wise error rate corrections for multiple comparisons. The minimum criterion for
the family-wise error rate was determined by white-noise clustering analysis (Monte Carlo
simulations, 5000 iterations) of data with the same dimension and smoothness as that
analyzed (Kadipasaoglu et al., 2014). Results were further restricted to regions with at least
three patients contributing to coverage and BGA percent change exceeding 5%.

Anatomical groups of electrodes were delineated, firstly, through indexing electrodes to the
closest node on the standardized cortical surface (Saad & Reynolds, 2012), and secondly,
through grouping channels into parcellations determined by Human Connectome Project
(HCP) space (Glasser et al., 2016). Parametric statistics were used since HCP regions of
interest contained >30 electrodes. When contrasting experimental conditions, two-sided
paired t-tests were evaluated at each time point for each region and significance levels were
computed at q<0.01 using an FDR correction for multiple comparisons.

To generate event-related potentials (ERPs), the raw data were band pass filtered (0.1–50
Hz). Trials were averaged together and the resultant waveform was smoothed (Savitzky-Golay
FIR, third-order, frame length of 251ms). To account for differences in polarity between
electrodes, ERPs were converted to root mean square (RMS), using a 50ms sliding window.
All electrodes were averaged within each subject, within ROI, and then the between subject
averages were used.

There is increasing evidence that the phase of activation across distributed neural populations
correlates with distinct underlying neural processes from the amplitude (Buzsáki, 2019; Fries,
2015; Sauseng & Klimesch, 2008). We investigated the influence of lexicality and phrase
construction on inter-trial phase coherence (ITC). Phase information was extracted from the
band-pass filtered data (zero-phase 3rd order Butterworth band-pass filter) using a Hilbert
transform. ITC was calculated as the circular mean (absolute vector length) of instantaneous
phase of stimulus-aligned trials computed as the median value of 10 iterations of a random
50% of trials.

To explore the functional connectivity between regions of interest, we used a generalized
phase-locking analysis to estimate the dominant spatio-temporal distributions of field activity,
and the strength of the coupling between them. Phase information was extracted from the
down-sampled (200 Hz) and wide band-pass filtered data (3–50 Hz; zero-phase 8th order
Butterworth band-pass filter) using the 'generalized phase' method (Davis et al., 2020) with a

single-sided Fourier transform approach (Marple, 1999). This method captures the phase of the predominant fluctuations in the wideband signal and minimizes filter-related distortion of the waveform. PLV was calculated as the circular mean (absolute vector length) of the instantaneous phase difference between each electrode pair at each time point and baselined to the period -500 to -100 ms before onset of the first word. Statistics were calculated using the mean PLV of correctly answered trials between 0 to 500ms after second word onset, comparing against a null distribution generated by randomly re-pairing trial recordings across the electrode pairs 500 times. Significant PLV was accepted at a threshold of $p < 0.05$.

Acknowledgements

We express our gratitude to all the patients who participated in this study; the neurologists at the Texas Comprehensive Epilepsy Program who participated in the care of these patients; and the nurses and technicians in the Epilepsy Monitoring Unit at Memorial Hermann Hospital who helped make this research possible. This work was supported by the National Institute of Neurological Disorders and Stroke NS098981. Human subjects: Patients participated in the experiments after written informed consent was obtained. All experimental procedures were reviewed and approved by the Committee for the Protection of Human Subjects (CPHS) of the University of Texas Health Science Center at Houston as Protocol Number: HSC-MS-06-0385.

References

Adger, D. (2012). *A Syntax of Substance*. https://doi.org/10.7551/mitpress/9780262018616.001.0001

Amunts, K., & Zilles, K. (2015). Architectonic Mapping of the Human Brain beyond Brodmann. *Neuron*, *88*(6), 1086–1107. https://doi.org/10.1016/j.neuron.2015.12.001

Antonucci, S., Beeson, P., Labiner, D., & Rapcsak, S. (2008). Lexical retrieval and semantic knowledge in patients with left inferior temporal lobe lesions. *Aphasiology*, *22*(3), 281–304. https://doi.org/10.1080/02687030701294491

Artoni, F., D'Orio, P., Catricalà, E., Conca, F., Bottoni, F., Pelliccia, V., … Moro, A. (2020). High gamma response tracks different syntactic structures in homophonous phrases. *Scientific Reports*, *10*(1), 7537. https://doi.org/10.1038/s41598-020-64375-9

Arya, R. (2019). Similarity of spatiotemporal dynamics of language-related ECoG high-gamma modulation in Japanese and English speakers. *Clinical Neurophysiology*, *130*(8), 1403–1404. https://doi.org/10.1016/j.clinph.2019.05.006

Baggio, G., & Hagoort, P. (2011). The balance between memory and unification in semantics: A dynamic account of the N400. *Language and Cognitive Processes*, *26*(9), 1338–1367. https://doi.org/10.1080/01690965.2010.542671

Bastiaansen, M., & Hagoort, P. (2015). Frequency-based Segregation of Syntactic and Semantic Unification during Online Sentence Level Language Comprehension. *Journal of Cognitive Neuroscience*, *27*(11), 2095–2107. https://doi.org/10.1162/jocn_a_00829

Bemis, D. K., & Pylkkänen, L. (2013). Basic linguistic composition recruits the left anterior temporal lobe and left angular gyrus during both listening and reading. *Cerebral Cortex*. https://doi.org/10.1093/cercor/bhs170

Bemis, Douglas K., & Pylkkänen, L. (2011). Simple composition: A magnetoencephalography investigation into the comprehension of minimal linguistic phrases. *Journal of Neuroscience*, *31*(8), 2801–2814. https://doi.org/10.1523/JNEUROSCI.5003-10.2011

Bemis, Douglas K., & Pylkkänen, L. (2013). Combination across domains: An MEG investigation into the relationship between mathematical, pictorial, and linguistic processing. *Frontiers in Psychology, 3*, 583. https://doi.org/10.3389/fpsyg.2012.00583

Benítez-Burraco, A., & Murphy, E. (2019). Why Brain Oscillations Are Improving Our Understanding of Language. *Frontiers in Behavioral Neuroscience*. https://doi.org/10.3389/fnbeh.2019.00190

Berwick, R. C., Stabler, E. P., Berwick, R. C., & Stabler, E. P. (2019). Minimalist parsing. In *Minimalist Parsing*.

https://doi.org/10.1093/oso/9780198795087.003.0001

Boylan, C., Trueswell, J. C., & Thompson-Schill, S. L. (2015). Compositionality and the angular gyrus: A multi-

voxel similarity analysis of the semantic composition of nouns and verbs. *Neuropsychologia*, *78*, 130–141.

https://doi.org/10.1016/j.neuropsychologia.2015.10.007

Bozic, M., Fonteneau, E., Su, L., & Marslen-Wilson, W. D. (2015). Grammatical analysis as a distributed

neurobiological function. *Human Brain Mapping*, *36*(3), 1190–1201. https://doi.org/10.1002/hbm.22696

Brennan, J., & Pylkkänen, L. (2012). The time-course and spatial distribution of brain activity associated with

sentence processing. *NeuroImage*, *60*(2), 1139–1148. https://doi.org/10.1016/j.neuroimage.2012.01.030

Brennan, J. R., Stabler, E. P., Van Wagenen, S. E., Luh, W. M., & Hale, J. T. (2016). Abstract linguistic structure

correlates with temporal activity during naturalistic comprehension. *Brain and Language*, *157–158*, 81–94.

https://doi.org/10.1016/j.bandl.2016.04.008

Brysbaert, M., New, B., & Keuleers, E. (2012). Adding part-of-speech information to the SUBTLEX-US word

frequencies. *Behavior Research Methods*, *44*(4), 991–997. https://doi.org/10.3758/s13428-012-0190-4

Buzsáki, G. (2019). *The Brain from Inside Out*. https://doi.org/10.1093/oso/9780190905385.001.0001

Buzsáki, G., & Watson, B. O. (2012). Brain rhythms and neural syntax: implications for efficient coding of

cognitive content and neuropsychiatric disease. *Dialogues in Clinical Neuroscience*, *14*(4), 345–367.

https://doi.org/10.31887/DCNS.2012.14.4/gbuzsaki

Canolty, R. T., Soltani, M., Dalal, S. S., Edwards, E., Dronkers, N. F., Nagarajan, S. S., … Knight, R. T. (2007).

Spatiotemporal Dynamics of Word Processing in the Human Brain. *Frontiers in Neuroscience*, *1*(1), 185–

196. https://doi.org/10.3389/neuro.01.1.1.014.2007

Cazzoli, D., Kaufmann, B. C., Paladini, R. E., Müri, R. M., Nef, T., & Nyffeler, T. (2021). Anterior insula and

inferior frontal gyrus: where ventral and dorsal visual attention systems meet. *Brain Communications*.

https://doi.org/10.1093/braincomms/fcaa220

Chomsky, N. (1995). *The Minimalist Program*. Cambridge, MA: MIT Press.

Chomsky, N., Gallego, A. J., & Ott, D. (2019). Generative grammar and the faculty of language: insights,

questions, and challenges. *Catalan Journal of Linguistics Special Issue*, 226–261.

Conner, C. R., Ellmore, T. M., Pieters, T. A., di Sano, M. A., & Tandon, N. (2011). Variability of the relationship

between electrophysiology and BOLD-fMRI across cortical regions in humans. *Journal of Neuroscience*,

31, 12855–12865. https://doi.org/10.1523/JNEUROSCI.1457-11.2011

Conner, C. R., Kadipasaoglu, C. M., Shouval, H. Z., Hickok, G., & Tandon, N. (2019). Network dynamics of

Broca's area during word selection. *PLoS ONE*, *14*(12), 1–30. https://doi.org/10.1371/journal.pone.0225756

Cox, R. W. (1996). AFNI: Software for Analysis and Visualization of Functional Magnetic Resonance

Neuroimages. *Computers and Biomedical Research*, *29*(3), 162–173.

https://doi.org/10.1006/cbmr.1996.0014

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical Surface-Based Analysis: I. Segmentation and Surface

Reconstruction. *NeuroImage*, *9*(2), 179–194. https://doi.org/10.1006/nimg.1998.0395

Davis, C. P., & Yee, E. (2019). Features, labels, space, and time: factors supporting taxonomic relationships in

the anterior temporal lobe and thematic relationships in the angular gyrus. *Language, Cognition and*

Neuroscience, *34*(10), 1347–1357.

Davis, Z. W., Muller, L., Martinez-Trujillo, J., Sejnowski, T., & Reynolds, J. H. (2020). Spontaneous travelling

cortical waves gate perception in behaving primates. *Nature*, *587*(7834), 432–436.

https://doi.org/10.1038/s41586-020-2802-y

Ding, N., Melloni, L., Zhang, H., Tian, X., & Poeppel, D. (2016). Cortical tracking of hierarchical linguistic

structures in connected speech. *Nature Neuroscience*, *19*(1), 158–164. https://doi.org/10.1038/nn.4186

Duffau, H., Moritz-Gasser, S., & Mandonnet, E. (2014). A re-examination of neural basis of language processing:

Proposal of a dynamic hodotopical model from data provided by brain stimulation mapping during picture

naming. *Brain and Language*, *131*, 1–10. https://doi.org/10.1016/j.bandl.2013.05.011

Fedorenko, E., Scott, T. L., Brunner, P., Coon, W. G., Pritchett, B., Schalk, G., & Kanwisher, N. (2016). Neural

correlate of the construction of sentence meaning. *Proceedings of the National Academy of Sciences of the*

United States of America, *113*(41), E6256–E6262. https://doi.org/10.1073/pnas.1612132113

Feng, X., Altarelli, I., Monzalvo, K., Ding, GuoshengRamus, F., Shu, H., Dehaene, S., … Dehaene-Lambertz, G.

(2020). A universal reading network and its modulation by writing system and reading ability in French and

Chinese children. *ELife*, *9*, e54591.

Figley, T. D., Mortazavi Moghadam, B., Bhullar, N., Kornelsen, J., Courtney, S. M., & Figley, C. R. (2017).

Probabilistic white matter atlases of human auditory, basal ganglia, language, precuneus, sensorimotor,

visual and visuospatial networks. *Frontiers in Human Neuroscience*, *11*(June), 1–12.

https://doi.org/10.3389/fnhum.2017.00306

Fischl, B., Sereno, M. I., & Dale, A. M. (1999). Cortical surface-based analysis: II. Inflation, flattening, and a

surface-based coordinate system. *NeuroImage*, *9*, 195–207. https://doi.org/10.1006/nimg.1998.0396

Flick, G., Oseki, Y., Kaczmarek, A. R., Al Kaabi, M., Marantz, A., & Pylkkänen, L. (2018). Building words and

phrases in the left temporal lobe. *Cortex*, *106*, 213–236. https://doi.org/10.1016/j.cortex.2018.06.004

Flick, G., & Pylkkänen, L. (2020). Isolating syntax in natural language: MEG evidence for an early contribution of

left posterior temporal cortex. *Cortex*, *127*, 42–57. https://doi.org/10.1016/j.cortex.2020.01.025

Flinker, A., Chang, E. F., Barbaro, N. M., Berger, M. S., & Knight, R. T. (2011). Sub-centimeter language

organization in the human temporal lobe. *Brain and Language*, *117*(3), 103–109.

https://doi.org/10.1016/j.bandl.2010.09.009

Forseth, K. J., Kadipasaoglu, C. M., Conner, C. R., Hickok, G., Knight, R. T., & Tandon, N. (2018). A lexical

semantic hub for heteromodal naming in middle fusiform gyrus. *Brain*, *141*(7), 2112–2126.

https://doi.org/10.1093/brain/awy120

Friederici, A. D., Chomsky, N., Berwick, R. C., Moro, A., & Bolhuis, J. J. (2017). Language, mind and brain.

Nature Human Behaviour, *1*(10), 713–722. https://doi.org/10.1038/s41562-017-0184-4

Fries, P. (2015). Rhythms for Cognition: Communication through Coherence. *Neuron*, *88*, 220–235.

https://doi.org/10.1016/j.neuron.2015.09.034

Gastaldon, S., Arcara, G., Navarrete, E., & Peressotti, F. (2020). Commonalities in alpha and beta neural

desynchronizations during prediction in language comprehension and production. *Cortex*, *133*, 328–345.

https://doi.org/10.1016/j.cortex.2020.09.026

Glasser, M. F., Coalson, T. S., Robinson, E. C., Hacker, C. D., Harwell, J., Yacoub, E., … Van Essen, D. C.

(2016). A multi-modal parcellation of human cerebral cortex. *Nature*, *536*(7615), 171–178.

https://doi.org/10.1038/nature18933

Glasser, M. F., & Rilling, J. K. (2008). DTI Tractography of the Human Brain's Language Pathways. *Cerebral*

Cortex, *18*(11), 2471–2482. https://doi.org/10.1093/cercor/bhn011

Graessner, A., Zaccarella, E., & Hartwigsen, G. (2021). Differential contributions of left-hemispheric language

regions to basic semantic composition. *Brain Structure and Function*. https://doi.org/10.1007/s00429-020-

02196-2

Hagoort, P. (2003). How the brain solves the binding problem for language: A neurocomputational model of

syntactic processing. *NeuroImage*, *20*, S18–S29. https://doi.org/10.1016/j.neuroimage.2003.09.013

Hagoort, P. (2004). Integration of Word Meaning and World Knowledge in Language Comprehension. *Science*,

304(5669), 438–441. https://doi.org/10.1126/science.1095455

Hagoort, P. (2005). On Broca, brain, and binding: A new framework. *Trends in Cognitive Sciences*.

https://doi.org/10.1016/j.tics.2005.07.004

Hagoort, P. (2013). MUC (Memory, Unification, Control) and beyond. *Frontiers in Psychology*, *4*.

https://doi.org/10.3389/fpsyg.2013.00416

Hagoort, P. (2017). The core and beyond in the language-ready brain. *Neuroscience & Biobehavioral Reviews*,

81, 194–204. https://doi.org/10.1016/j.neubiorev.2017.01.048

Hagoort, P. (2020). The meaning-making mechanism(s) behind the eyes and between the ears. *Philosophical*

Transactions of the Royal Society of London. Series B, Biological Sciences, *375*(1791), 20190301.

https://doi.org/10.1098/rstb.2019.0301

Hagoort, P., Baggio, G., & Willems, R. M. (2009). Semantic unification. In M. S. Gazzaniga (Ed.), *The Cognitive*

Neurosciences (4th ed., pp. 819–836). MIT Press.

Hagoort, P., & Indefrey, P. (2014). The Neurobiology of Language Beyond Single Words. *Annual Review of*

Neuroscience, *37*(1), 347–362. https://doi.org/10.1146/annurev-neuro-071013-013847

Hovsepyan, S., Olasagasti, I., & Giraud, A.-L. (2020). Combining predictive coding and neural oscillations

enables online syllable recognition in natural speech. *Nature Communications*, *11*(1), 3117.

https://doi.org/10.1038/s41467-020-16956-5

Humphries, C., Binder, J. R., Medler, D. A., & Liebenthal, E. (2006). Syntactic and semantic modulation of neural

activity during auditory sentence comprehension. *Journal of Cognitive Neuroscience*.

https://doi.org/10.1162/jocn.2006.18.4.665

Irimia, A., Van Horn, J. D., & Halgren, E. (2012). Source cancellation profiles of electroencephalography and

magnetoencephalography. *NeuroImage*, *59*(3), 2464–2474.

https://doi.org/10.1016/j.neuroimage.2011.08.104

Jackendoff, R., & Audring, J. (2020). *The Texture of the Lexicon: Relational Morphology and the Parallel*

Architecture. Oxford: Oxford University Press.

Jensen, O., Spaak, E., & Zumer, J. M. (2019). Human brain oscillations: From physiological mechanisms to

analysis and cognition. In *Magnetoencephalography: From Signals to Dynamic Cortical Networks: Second*

Edition. https://doi.org/10.1007/978-3-030-00087-5_17

Johnson, E. L., Kam, J. W. Y., Tzovara, A., & Knight, R. T. (2020). Insights into human cognition from intracranial

EEG: A review of audition, memory, internal cognition, and causality. *Journal of Neural Engineering*, *17*(5),

051001. https://doi.org/10.1088/1741-2552/abb7a5

Kadipasaoglu, C. M., Baboyan, V. G., Conner, C. R., Chen, G., Saad, Z. S., & Tandon, N. (2014). Surface-based

mixed effects multilevel analysis of grouped human electrocorticography. *NeuroImage*, *101*, 215–224.

https://doi.org/10.1016/j.neuroimage.2014.07.006

Kadipasaoglu, Cihan M., Forseth, K., Whaley, M., Conner, C. R., Rollo, M. J., Baboyan, V. G., & Tandon, N.

(2015). Development of grouped icEEG for the study of cognitive processing. *Frontiers in Psychology*,

6(1008). https://doi.org/10.3389/fpsyg.2015.01008

Keitel, A., Gross, J., & Kayser, C. (2018). Perceptually relevant speech tracking in auditory and motor cortex

reflects distinct linguistic features. *PLOS Biology*, *16*(3), e2004473.

https://doi.org/10.1371/journal.pbio.2004473

Keitel, A., Ince, R. A. A., Gross, J., & Kayser, C. (2017). Auditory cortical delta-entrainment interacts with

oscillatory power in multiple fronto-parietal networks. *NeuroImage*, *147*, 32–42.

https://doi.org/10.1016/j.neuroimage.2016.11.062

Kleiner, M., Brainard, D., Pelli, D., Ingling, A., Murray, R., & Broussard, C. (2007). What's new in psychtoolbox-3.

Perception, *36*(14), 1–16.

Kucyi, A., & Parvizi, J. (2020). Pupillary Dynamics Link Spontaneous and Task-Evoked Activations Recorded

Directly from Human Insula. *The Journal of Neuroscience*, *40*(32), 6207–6218.

https://doi.org/10.1523/JNEUROSCI.0435-20.2020

Lambon Ralph, M. A., Ehsan, S., Baker, G. A., & Rogers, T. T. (2012). Semantic memory is impaired in patients

with unilateral anterior temporal lobe resection for temporal lobe epilepsy. *Brain*, *135*(1), 242–258.

https://doi.org/10.1093/brain/awr325

Lau, E., Stroud, C., Plesch, S., & Phillips, C. (2006). The role of structural prediction in rapid syntactic analysis.

Brain and Language, *98*(1), 74–88. https://doi.org/10.1016/j.bandl.2006.02.003

Law, R., & Pylkkänen, L. (2021). Lists with and without Syntax: A New Approach to Measuring the Neural

Processing of Syntax. *The Journal of Neuroscience*, *41*(10), 2186–2196.

https://doi.org/10.1523/JNEUROSCI.1179-20.2021

Lee, D. K., Fedorenko, E., Simon, M. V., Curry, W. T., Nahed, B. V., Cahill, D. P., & Williams, Z. M. (2018).

Neural encoding and production of functional morphemes in the posterior temporal lobe. *Nature*

Communications, *9*, 1877. https://doi.org/10.1038/s41467-018-04235-3

Leivada, E., & Murphy, E. (2021). Mind the (terminological) gap: 10 misused, ambiguous, or polysemous terms in linguistics. *Ampersand*, *8*, 100073. https://doi.org/10.1016/j.amper.2021.100073

Leszczyński, M., Barczak, A., Kajikawa, Y., Ulbert, I., Falchier, A. Y., Tal, I., … Schroeder, C. E. (2020). Dissociation of broadband high-frequency activity and neuronal firing in the neocortex. *Science Advances*, *6*(33), eabb0977. https://doi.org/10.1126/sciadv.abb0977

Lewis, A. G., Schoffelen, J.-M., Schriefers, H., & Bastiaansen, M. (2016). A Predictive Coding Perspective on Beta Oscillations during Sentence-Level Language Comprehension. *Frontiers in Human Neuroscience*, *10*, 83. https://doi.org/10.3389/fnhum.2016.00085

Lopopolo, A., van den Bosch, A., Petersson, K.-M., & Willems, R. M. (2021). Distinguishing Syntactic Operations in the Brain: Dependency and Phrase-Structure Parsing. *Neurobiology of Language*, *2*(1), 152–175. https://doi.org/10.1162/nol_a_00029

Mai, G., Minett, J. W., & Wang, W. S.-Y. (2016). Delta, theta, beta, and gamma brain oscillations index levels of auditory sentence processing. *NeuroImage*, *133*, 516–528. https://doi.org/10.1016/j.neuroimage.2016.02.064

Marko, M., Cimrová, B., & Riečanský, I. (2019). Neural theta oscillations support semantic memory retrieval. *Scientific Reports*, *9*(1), 17667. https://doi.org/10.1038/s41598-019-53813-y

Marple, L. (1999). Computing the discrete-time "analytic" signal via FFT. *IEEE Transactions on Signal Processing*, *47*(9), 2600–2603.

Martin, S., Millán, J. del R., Knight, R. T., & Pasley, B. N. (2019). The use of intracranial recordings to decode human language: Challenges and opportunities. *Brain and Language*, *193*, 73–83. https://doi.org/10.1016/j.bandl.2016.06.003

Matar, S., Dirani, J., Marantz, A., & Pylkkänen, L. (2021). Left posterior temporal cortex is sensitive to syntax within conceptually matched Arabic expressions. *Scientific Reports*, *11*(7181).

Matchin, W., & Hickok, G. (2020). The Cortical Organization of Syntax. *Cerebral Cortex*, *30*(3), 1481–1498. https://doi.org/10.1093/cercor/bhz180

Muller, L., Chavane, F., Reynolds, J., & Sejnowski, T. J. (2018). Cortical travelling waves: mechanisms and computational principles. *Nature Reviews Neuroscience*, *19*(5), 255–268. https://doi.org/10.1038/nrn.2018.20

Murphy, E. (2015a). Labels, cognomes, and cyclic computation: An ethological perspective. *Frontiers in*
Psychology, *6*(715). https://doi.org/10.3389/fpsyg.2015.00715

Murphy, E. (2015b). The brain dynamics of linguistic computation. *Frontiers in Psychology*, *6*(1515).
https://doi.org/10.3389/fpsyg.2015.01515

Murphy, Elliot. (2015). Labels, cognomes, and cyclic computation: An ethological perspective. *Frontiers in*
Psychology, *6*(715). https://doi.org/10.3389/fpsyg.2015.00715

Murphy, Elliot. (2018). Interfaces (travelling oscillations) + recursion (delta-theta code) = language. In E. Luef &
M. Manuela (Eds.), *The Talking Species: Perspectives on the Evolutionary, Neuronal and Cultural*
Foundations of Language (pp. 251–269). Graz.

Murphy, Elliot. (2019). No Country for Oldowan Men: Emerging Factors in Language Evolution. *Frontiers in*
Psychology, *10*. https://doi.org/10.3389/fpsyg.2019.01448

Murphy, Elliot. (2020). *The Oscillatory Nature of Language*. Cambridge: Cambridge University Press.

Murphy, Elliot, Holmes, E., & Friston, K. (2021). Natural language syntax complies with the free-energy principle.
PsyArXiv. https://doi.org/doi.org/10.31234/osf.io/r3fcx

Murphy, Elliot, & Shim, J.-Y. (2020). Copy invisibility and (non-)categorial labeling. *Linguistic Research*, *37*(2),
187–215. https://doi.org/10.17250/khisli.37.2.202006.002

Nelson, M. J., El Karoui, I., Giber, K., Yang, X., Cohen, L., Koopman, H., … Dehaene, S. (2017).
Neurophysiological dynamics of phrase-structure building during sentence processing. *Proceedings of the*
National Academy of Sciences of the United States of America, *114*(18), E3669–E3678.
https://doi.org/10.1073/pnas.1701590114

Neufeld, C., Kramer, S. E., Lapinskaya, N., Heffner, C. C., Malko, A., & Lau, E. F. (2016). The electrophysiology
of basic phrase building. *PLoS ONE*, *11*(10), 1–22. https://doi.org/10.1371/journal.pone.0158446

Packard, P. A., Steiger, T. K., Fuentemilla, L., & Bunzeck, N. (2020). Neural oscillations and event-related
potentials reveal how semantic congruence drives long-term memory in both young and older humans.
Scientific Reports. https://doi.org/10.1038/s41598-020-65872-7

Pieters, T. A., Conner, C. R., & Tandon, N. (2013). Recursive grid partitioning on a cortical surface model: an
optimized technique for the localization of implanted subdural electrodes. *Journal of Neurosurgery*, *118*(5),
1086–1097. https://doi.org/10.3171/2013.2.JNS121450

Pietroski, P. M. (2018). Conjoining meanings: Semantics without truth values. In *Conjoining Meanings: Semantics*

Without Truth Values. https://doi.org/10.1093/oso/9780198812722.001.0001

Pobric, G., Jefferies, E., & Lambon Ralph, M. A. (2010). Category-Specific versus Category-General Semantic Impairment Induced by Transcranial Magnetic Stimulation. *Current Biology*, *20*(10), 964–968. https://doi.org/10.1016/j.cub.2010.03.070

Prystauka, Y., & Lewis, A. G. (2019). The power of neural oscillations to inform sentence comprehension: A linguistic perspective. *Language and Linguistics Compass*, *13*(9). https://doi.org/10.1111/lnc3.12347

Pulvermüller, F., & Grisoni, L. (2020). Semantic prediction in brain and mind. *Trends in Cognitive Sciences*, *24*(10), 781–784.

Pylkkänen, L. (2020). Neural basis of basic composition: what we have learned from the red-boat studies and their extensions. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *375*(1791), 20190299. https://doi.org/10.1098/rstb.2019.0299

Pylkkänen, L., & Brennan, J. (2020). Composition: the neurobiology of syntactic and semantic phrase structure building. In D. Poeppel, G. . Mangun, & M. . Gazzaniga (Eds.), *The Cognitive Neurosciences* (6th ed.). Cambridge, MA: MIT Press.

Rizzi, L. (2016). Monkey morpho-syntax and merge-based systems. *Theoretical Linguistics*, *42*(1–2), 139–145. https://doi.org/10.1515/tl-2016-0006

Rollo, P. S., Rollo, M. J., Zhu, P., Woolnough, O., & Tandon, N. (2020). Oblique trajectory angles in robotic stereo-electroencephalography. *Journal of Neurosurgery*, 1–10. https://doi.org/10.3171/2020.5.JNS20975

Saad, Z. S., & Reynolds, R. C. (2012). SUMA. *NeuroImage*, *62*, 768–773. https://doi.org/10.1016/j.neuroimage.2011.09.016

Sarubbo, S., Tate, M., De Benedictis, A., Merler, S., Moritz-Gasser, S., Herbet, G., & Duffau, H. (2020). Mapping critical cortical hubs and white matter pathways by direct electrical stimulation: an original functional atlas of the human brain. *NeuroImage*, *205*(September 2019), 116237. https://doi.org/10.1016/j.neuroimage.2019.116237

Sauseng, P., & Klimesch, W. (2008). What does phase information of oscillatory brain activity tell us about cognitive processes? *Neuroscience and Biobehavioral Reviews*, *32*, 1001–1013. https://doi.org/10.1016/j.neubiorev.2008.03.014

Schell, M., Zaccarella, E., & Friederici, A. D. (2017). Differential cortical contribution of syntax and semantics: An fMRI study on two-word phrasal processing. *Cortex*, *96*, 105–120.

https://doi.org/10.1016/j.cortex.2017.09.002

Segaert, K., Mazaheri, A., & Hagoort, P. (2018). Binding language: structuring sentences through precisely timed

oscillatory mechanisms. *European Journal of Neuroscience*, *48*(7), 2651–2662.

https://doi.org/10.1111/ejn.13816

Tandon, N. (2012). Mapping of human language. In D. Yoshor & E. Mizrahi (Eds.), *Clinical Brain Mapping* (pp.

203–218). McGraw Hill Education.

Tandon, N., Tong, B. A., Friedman, E. R., Johnson, J. A., Von Allmen, G., Thomas, M. S., … Thompson, S. A.

(2019). Analysis of Morbidity and Outcomes Associated with Use of Subdural Grids vs

Stereoelectroencephalography in Patients with Intractable Epilepsy. *JAMA Neurology*, *76*, 672–681.

https://doi.org/10.1001/jamaneurol.2019.0098

Tanji, K., Suzuki, K., Delorme, A., Shamoto, H., & Nakasato, N. (2005). High-frequency γ-band activity in the

basal temporal cortex during picture-naming and lexical-decision tasks. *Journal of Neuroscience*, *25*(13),

3287–3293. https://doi.org/10.1523/JNEUROSCI.4948-04.2005

Terporten, R., Schoffelen, J. M., Dai, B., Hagoort, P., & Kösem, A. (2019). The Relation between Alpha/Beta

Oscillations and the Encoding of Sentence induced Contextual Information. *Scientific Reports*.

https://doi.org/10.1038/s41598-019-56600-x

Thye, M., Geller, J., Szaflarski, J. P., & Mirman, D. (2021). Intracranial EEG evidence of functional specialization

for taxonomic and thematic relations. *Cortex*. https://doi.org/10.1016/j.cortex.2021.03.018

Tong, B. A., Esquenazi, Y., Johnson, J., Zhu, P., & Tandon, N. (2020). The Brain is Not Flat: Conformal Electrode

Arrays Diminish Complications of Subdural Electrode Implantation, A Series of 117 Cases. *World*

Neurosurgery, *144*, e734–e742. https://doi.org/10.1016/j.wneu.2020.09.063

Towle, V. L., Yoon, H.-A., Castelle, M., Edgar, J. C., Biassou, N. M., Frim, D. M., … Kohrman, M. H. (2008).

ECoG gamma activity during a language task: differentiating expressive and receptive speech areas. *Brain*,

131(8), 2013–2027. https://doi.org/10.1093/brain/awn147

Trettenbrein, P., Papitto, G., Friederici, A. D., & Zaccarella, E. (2020). Functional neuroanatomy of language

without speech: An ALE meta-analysis of sign language. *Human Brain Mapping*, *doi.org/10*.

Wang, X., Men, W., Gao, J., Caramazza, A., & Bi, Y. (2020). Two Forms of Knowledge Representations in the

Human Brain. *Neuron*, *107*(2), 383–393. https://doi.org/10.1016/j.neuron.2020.04.010

Westerlund, M., Kastner, I., Al Kaabi, M., & Pylkkänen, L. (2015). The LATL as locus of composition: MEG

evidence from English and Arabic. *Brain and Language*, *141*, 124–134.

https://doi.org/10.1016/j.bandl.2014.12.003

Westerlund, M., & Pylkkänen, L. (2014). The role of the left anterior temporal lobe in semantic composition vs.

semantic memory. *Neuropsychologia*, *57*(1), 59–70.

https://doi.org/10.1016/j.neuropsychologia.2014.03.001

Wilson, S. M., DeMarco, A. T., Henry, M. L., Gesierich, B., Babiak, M., Mandelli, M. L., … Gorno-Tempini, M. L.

(2014). What role does the anterior temporal lobe play in sentence-level processing? neural correlates of

syntactic processing in semantic variant primary progressive aphasia. *Journal of Cognitive Neuroscience*,

26(5), 970–985. https://doi.org/10.1162/jocn_a_00550

Wilson, S. M., & Saygin, A. P. (2004). Grammaticality Judgment in Aphasia: Deficits Are Not Specific to Syntactic

Structures, Aphasic Syndromes, or Lesion Sites. *Journal of Cognitive Neuroscience*, *16*, 238–252.

https://doi.org/10.1162/089892904322984535

Woolnough, O., Forseth, K. J., Rollo, P. S., & Tandon, N. (2019). Uncovering the functional anatomy of the

human insula during speech. *ELife*, *8*, 1–14. https://doi.org/10.7554/eLife.53086

Yi, H. G., Leonard, M. K., & Chang, E. F. (2019). The Encoding of Speech Sounds in the Superior Temporal

Gyrus. *Neuron*, *102*(6), 1096–1110. https://doi.org/10.1016/j.neuron.2019.04.023

Zaccarella, E., Papitto, G., & Friederici, A. D. (2021). Language and action in Broca's area: Computational

differentiation and cortical segregation. *Brain and Cognition*, *147*, 105651.

https://doi.org/10.1016/j.bandc.2020.105651

Zhang, L., & Pylkkänen, L. (2018). Semantic composition of sentences word by word: MEG evidence for shared

processing of conceptual and logical elements. *Neuropsychologia*, *119*(April), 392–404.

https://doi.org/10.1016/j.neuropsychologia.2018.08.016

 Supplementary Figures

<u>**Supplementary Figure 1**</u>: **Inter-trial phase coherence and lexicality.**

(A) Electrodes with significant differences in ITC for words vs pseudowords, in a portion of
posterior temporal cortex with a high concentration of electrodes exhibiting significant
differences (9 patients); HCP index: area STV (Glasser et al., 2016). (B) Gamma traces (green
dashed box) for electrodes in (A) for words vs. pseudowords and compositional vs. non-
compositional trials (C) ITC in alpha (8-12Hz) for all electrodes in (A) for words (blue) and
pseudowords (red) on the left, with alpha (orange dashed box) amplitude traces, and for
phrase composition (purple) and non-composition (black) on the right isolated around the
onset of the second word (no significant effect for composition contrast). T0 = first word onset.

Supplementary Figure 2: **Regions of interest and their BGA/ERP signatures.** Regional broadband gamma amplitude (middle column) with event-related potentials (right column). Red: Compositional. Black: Non-Compositional. HCP index from top-down: posterior superior temporal gyrus (pSTG) = [A4]; temporal pole = [TGd]; inferior frontal gyrus (IFG) = [FOP4, 44, 45, IFSp, p47r, IFSa, 47l]; posterior middle temporal gyrus (pMTG) = [PHT, TE1p]. BGA traces are thresholded by p<0.05 significantly active from baseline (first word onset, -500 to -100ms) with a minimum of 10% BGA amplitude increase during the 100-400ms post-second word onset period. ERPs were computed across the four ROIs plotted on the left column, with significant condition differences being computed across 0-1000ms in the same (FDR-corrected) manner as the BGA plots (see Methods).

Supplementary Figure 3: **pSTS-TOJ regional spectrograms.** Spectrograms and electrodes
in pSTS-TOJ for all active channels (HCP index: TPOJ1; electrodes = 23, patients = 10).
Second word onset was at 800ms.

Supplementary Figure 4: **Temporal pole effects.** Low frequency (2–15Hz) temporal pole
trace with FDR-significant bars for the Compositional (brown) vs Non-Compositional (purple)
contrast (HCP index: TGd) with spectrograms, thresholded at 10% increased activity from
baseline; T0 = second word onset. For BGA, see Suppl. Fig. 2.