Minimal phrase composition revealed by intracranial recordings

Authors:

Elliot Murphy[1,2], Oscar Woolnough[1,2], Patrick S. Rollo[1,2], Zachary Roccaforte[1], Katrien Segaert[3,4], Peter Hagoort[4,5], Nitin Tandon[1,2,6,*]

1. Vivian L. Smith Department of Neurosurgery, McGovern Medical School, University of Texas Health Science Center at Houston, Houston, TX 77030

2. Texas Institute for Restorative Neurotechnologies, University of Texas Health Science Center at Houston, Houston, TX 77030

3. School of Psychology & Centre for Human Brain Health, University of Birmingham, Birmingham, UK, B15 2TT

4. Max Planck Institute for Psycholinguistics, Nijmegen, The Netherlands, 6525 XD Nijmegen

5. Donders Institute for Brain, Cognition and Behaviour, Nijmegen, The Netherlands, 6525 HR Nijmegen

6. Memorial Hermann Hospital, Texas Medical Center, Houston, TX 77030

*Corresponding author: Nitin Tandon (nitin.tandon@uth.tmc.edu), +1 713-500-4472, Vivian L. Smith Department of Neurosurgery, McGovern Medical School, University of Texas Health Science Center at Houston, Houston, TX 77030

Running title: Minimal phrase composition revealed by intracranial recordings

Number of pages: 30

Number of figures: 6

Word length: 242 (Abstract), 115 (Significance), 603 (Introduction), 1496 (Discussion)

Conflict of interest: The authors declare no competing financial interests.

Acknowledgements: We express our gratitude to all the patients who participated in this study; the neurologists at the Texas Comprehensive Epilepsy Program who participated in the care of these patients; and the nurses and technicians in the Epilepsy Monitoring Unit at Memorial Hermann Hospital who helped make this research possible. This work was supported by the National Institute of Neurological Disorders and Stroke NS098981. Human subjects: Patients participated in the experiments after written informed consent was obtained. All experimental procedures were reviewed and approved by the Committee for the Protection of Human Subjects (CPHS) of the University of Texas Health Science Center at Houston as Protocol Number: HSC-MS-06-0385.

Author Contributions: N.T., K.S., P.H. designed research; E.M., P.S.R., Z.R. performed research; E.M. analyzed data; E.M. wrote the first draft of the paper; E.M., O.W., K.S., P.H., and N.T. edited the paper; N.T. conducted funding acquisition and neurosurgical procedures.

Abstract

The ability to comprehend phrases is an essential integrative property of the brain. Here we
evaluate the neural processes that enable the transition from single word processing to a
minimal compositional scheme. Previous research has reported conflicting timing effects of
composition, and disagreement persists with respect to inferior frontal and posterior temporal
contributions. To address these issues, 19 patients (10 male, 19 female) implanted with
penetrating depth or surface subdural intracranial electrodes heard auditory recordings of
adjective-noun, pseudoword-noun and adjective-pseudoword phrases and judged whether
the phrase matched a picture. Stimulus-dependent alterations in broadband gamma activity,
low frequency power and phase-locking values across the language-dominant left hemisphere
were derived. This revealed a mosaic located in the posterior superior temporal sulcus (pSTS),
in which closely neighboring cortical sites displayed exclusive sensitivity to either lexicality or
phrase structure, but not both. Distinct timings were found for effects of phrase composition
(210–300 ms) and pseudoword processing (approximately 300–700 ms), and these were
localized to neighboring electrodes in pSTS. The pars triangularis and temporal pole encode
anticipation of composition in broadband low frequencies, and both regions exhibit greater
functional connectivity with pSTS during phrase composition. Our results suggest that the
pSTS is a highly specialized region comprised of sparsely interwoven heterogeneous
constituents that encodes both lower and higher level linguistic features. This hub in pSTS for
minimal phrase processing may form the neural basis for the human-specific computational
capacity for forming hierarchically organized linguistic structures.

Keywords: semantics, human, phrase structure, auditory, language

Significance

Linguists have claimed that the integration of multiple words into a phrase demands a computational procedure distinct from single word processing. Here, we provide intracranial recordings from a large patient cohort, with high spatiotemporal resolution, to track the cortical dynamics of phrase composition. Epileptic patients volunteered to participate in a task in which they listened to phrases ("red boat"), word-pseudoword or pseudoword-word pairs (e.g., "red fulg"). At the onset of the second word in phrases, greater broadband high gamma activity was found in posterior superior temporal sulcus in electrodes that exclusively indexed phrasal meaning, and not lexical meaning. These results provide direct, high-resolution signatures of minimal phrase composition in humans, a potentially species-specific computational capacity.

Introduction

How the brain integrates individual word meanings to comprehend multi-word utterances is an issue that has vexed the cognitive neuroscience of language for decades (Hagoort, 2020). This linguistic compositional process – the combination of words into larger structures with new and complex meaning – has been referred to as "Merge" (Chomsky et al., 2019) or "Unification" (Hagoort, 2013). A simple paradigm for studying complex meaning is the use of minimal phrases, such as in the *red-boat* paradigm, which focuses on simple combinations of two words and avoids confounds associated with more complex linguistic stimuli (Bemis and Pylkkänen, 2011, 2013; Brennan and Pylkkänen, 2012; Bozic et al., 2015; Flick et al., 2018; Flick and Pylkkänen, 2020; Pylkkänen, 2020). A "red boat" is interpreted as a boat which is red, and not a red object which hosts boat-related properties, with phrases delivering novel syntactic and conceptual formats (Murphy, 2015). *Red-boat* experiments isolate semantic composition, which in turn encompasses syntactic composition.

Functional neuroimaging studies using the *red-boat* paradigm implicate the left anterior temporal lobe (ATL), specifically the temporal pole (Antonucci et al., 2008; Lambon Ralph et al., 2012; Wilson et al., 2014; Zhang and Pylkkänen, 2018), inferior frontal regions (Graessner et al., 2021b, 2021a) and posterior temporal regions (Flick and Pylkkänen, 2020; Matchin and Hickok, 2020) as crucial nodes for phrase composition with variations in the timing (180–350ms post-composition) and duration (50–100ms) of their engagement (Kochari et al., 2021). Posterior temporal cortex is more broadly implicated in syntactic comprehension and production (Duffau et al., 2014; Artoni et al., 2020; Matchin and Hickok, 2020; Graessner et al., 2021b; Lopopolo et al., 2021). The left posterior temporal lobe and angular gyrus (AG) show greater activity for sentences than word lists, and for phrases than words, making them candidate regions for the retrieval of phrasal templates (Hagoort, 2003, 2017; Brennan et al., 2016; Matchin and Hickok, 2020). According to a range of parsing models, adjective-noun syntax is also constructed *predictively* (Berwick et al., 2019), and anticipatory amplitude increases have been isolated to the alpha/beta bands (Gastaldon et al., 2020).

Phrase composition is a rapid process that is likely dependent on finely organized sets of distributed cortical substrates. Previous work using the *red-boat* paradigm has been limited by spatiotemporal resolution. Using recordings from intracranial electroencephalography (iEEG) with depth electrodes penetrating grey matter or electrodes located on the cortical surface, we conducted a study of minimal phrase composition with auditory presentations of the red boat paradigm, building directly and using stimuli from an established design (Bemis and Pylkkänen, 2013). Given our large cohort, we were able to perform a data driven analysis of iEEG data from the whole brain acquired at unprecedented spatiotemporal resolution.

Most evidence that left ATL and posterior temporal regions are involved in basic composition
is from magnetoencephalography (MEG), while the evidence supporting a role for IFG mostly
comes from functional magnetic resonance imaging (fMRI). Signal loss in fMRI may partly
explain the lack of ATL effects in this portion of the imaging literature (due to proximity to
sinuses) (Olman et al., 2009; Bonner and Price, 2013), yet even within the MEG literature
there is variation in the timing of composition effects. This suggests that intracranial recordings
can contribute to resolving the spatiotemporal dynamics of phrase processing. Our central
research questions concerned: (i) the precise localization of composition effects, addressing
a current tension in the literature with respect to frontal, temporal and parietal contributions to
minimal phrase processing (Schell et al., 2017; Flick and Pylkkänen, 2020; Graessner et al.,
2021a); (ii) the precise timing and duration of composition effects, addressing the above noted
variations documented in the literature.

Materials and Methods

Participants: 19 patients (10 male, 18-41 years, IQ 97 ± 12, 2 left-handed) participated in the
experiment after written informed consent was obtained. All were native English speakers. All
experimental procedures were reviewed and approved by the Committee for the Protection of
Human Subjects (CPHS) of the University of Texas Health Science Center at Houston as
Protocol Number HSC-MS-06-0385.

Electrode Implantation and Data Recording: Data were acquired from either subdural grid
electrodes (SDEs; 6 patients) or stereotactically placed depth electrodes (sEEGs; 13 patients)
(Fig. 1B). SDEs were subdural platinum-iridium electrodes embedded in a silicone elastomer
sheet (PMT Corporation; top-hat design; 3mm diameter cortical contact), and were surgically
implanted via a craniotomy (Conner et al., 2011; Pieters et al., 2013; Tong et al., 2020;
Kohlhase et al., 2021). sEEGs were implanted using a Robotic Surgical Assistant (ROSA;
Medtech, Montpellier, France) (Rollo et al., 2020; McCarty et al., 2021). Each sEEG probe
(PMT corporation, Chanhassen, Minnesota) was 0.8mm in diameter and had 8-16 electrode
contacts. Each contact was a platinum-iridium cylinder, 2.0mm in length and separated from
the adjacent contact by 1.5-2.43mm. SDE patients had a number of cortical arrays implanted
(mean ± SD: 9 ± 1) with a mean of 146.5 electrodes (SD ± 53.5). sEEG patients had
penetrating depth probes (mean 15.1 ± 2.3) with a mean of 199.9 electrodes (SD ± 30.1).
Typical coverage was fronto-temporal, dictated by location of the epilepsy in the antero-mesial
temporal lobe in the majority, with parietal and occipital coverage in a number of patients.
Following surgical implantation, electrodes were localized by co-registration of pre-operative
anatomical 3T MRI and post-operative CT scans in AFNI (Cox, 1996). Electrode positions

were projected onto a cortical surface model generated in FreeSurfer (Dale et al., 1999), and
displayed on the cortical surface model for visualization (Pieters et al., 2013). Intracranial data
were collected during research experiments starting on the first day after electrode
implantation for sEEGs and two days after implantation for SDEs. Data were digitized at 2 kHz
using the NeuroPort recording system (Blackrock Microsystems, Salt Lake City, Utah),
imported into Matlab, initially referenced to the white matter channel used as a reference for
the clinical acquisition system and visually inspected for line noise, artifacts and epileptic
activity. iEEG provides uniquely high spatiotemporal resolution recordings and is less
susceptible to artifacts (e.g., muscle movements) (Flinker et al., 2011; Arya, 2019). Electrodes
with excessive line noise or localized to sites of seizure onset were excluded. Each electrode
was re-referenced to the common average of the remaining channels. Trials contaminated by
inter-ictal epileptic spikes, saccade artefacts and trials in which participants responded
incorrectly were discarded. Electrodes contributing to regions of interest were taken from both
SDE and sEEG patients; for instance, we follow previous intracranial work that has included
subdural contacts in monitoring activity from posterior temporal sulcus (Uno et al., 2015).

Experimental Design and Statistical Analysis: Grammatical noun phrases ("red boat") and
pseudoword phrases with variable positions of the pseudoword ("bleeg boat", "red fulg") were
used to isolate semantic compositional processing. An adjective-pseudoword condition ("red
fulg") allowed for the isolation of semantic compositionality and reduced predictability. The
inclusion of pseudowords in our stimulus set builds upon a previous study, from which we
abridged our list of words and graphic images (Bemis and Pylkkänen, 2013). We focused on
high frequency gamma changes (Forseth et al., 2018; Conner et al., 2019; Johnson et al.,
2020; Leszczyński et al., 2020) that typically index local cortical processing and are implicated
in a range of cognitive processes (Buzsáki and Watson, 2012; Hovsepyan et al., 2020;
Packard et al., 2020). An early anterior negative deflection immediately preceding the critical
noun in combinatorial contexts (-50–100ms) likely indexes syntactic prediction (Neufeld et al.,
2016). Low frequency power increases have also been noted during the anticipatory window
and during phrase composition (Bastiaansen and Hagoort, 2015; Lewis et al., 2016; Segaert
et al., 2018), and in a variety of auditory phrase and sentence processing paradigms (Ding et
al., 2016; Mai et al., 2016; Keitel et al., 2017, 2018; Leivada and Murphy, 2021). Therefore,
we also evaluated the role of low frequencies during anticipatory composition, focusing on the
alpha/beta bands given the joint involvement of these in the literature and the similar
anticipatory dynamics attributed to them. Lastly, we asked patients to determine whether the
words they heard matched a subsequent image, enabling validation of their attention as well
as analyses related to phrase-matched and phrase-contrasted contexts. Our analyses were
restricted to language-dominant left hemisphere electrode coverage.

Participants were presented with two-word auditory phrases, grouped by three conditions: Adjective-Noun ("red boat"), Adjective-Pseudoword ("red neub"), Pseudoword-Noun ("zuik boat"). Since these pseudoword phrases include phonologically viable nonwords, differences in the second position of the phrase between these items and the grammatical noun phrases are likely a result of compositional processing. While most previous studies have presented a licensable noun, our inclusion of the Adjective-Pseudoword condition further isolates composition and reduces predictability. To ensure attention was maintained, after each trial participants were shown a colored drawing and asked to press a button indicating whether the picture matched the phrase they had just heard. Participants were told to respond positively only when the picture fully matched the phrase. Auditory and visual stimuli were adapted from a previous *red-boat* experiment (Bemis and Pylkkänen, 2013), from which we obtained our list of real words and graphic images.

A fixation cross was presented in the centre of the screen for 700 ms followed by the first word, and 800 ms later the second word was presented. 1600 ms after the onset of the second word the picture was presented, and 1400 ms after picture presentation participants were prompted to respond (Fig. 1A). Following their response, a blank screen was shown for 1500 ms. Stimuli were presented in a pseudorandom order, with no repetition amongst items. The number of trials per block across the full experiment was as follows: Adjective-Noun (80), Pseudoword-Noun (40), Adjective-Pseudoword (40). All patients undertook 2 blocks. Half of the Adjective-Noun trials matched the picture presented (i.e., "red boat" was heard by the patient, and a picture of a red boat was then presented), and the other half did not match. 6 adjectives were used: black, blue, brown, green, pink, red (length M: 4.3, SD: 0.7; SUBTLEXus log-frequency 3.64). 20 nouns were used: bag, bell, boat, bone, cane, cross, cup, disc, flag, fork, hand, heart, house, key, lamp, leaf, lock, plane, shoe, star (length M: 4.0, SD: 0.6; log-frequency 3.38) (Brysbaert et al., 2012). 6 pseudowords were used: beeg, cresp, kleg, nar, neub, zuik (length M: 4.0, SD: 0.6). Average stimuli length: Adjectives (420 ms), Nouns (450 ms), Pseudowords (430ms). Stimuli were presented using Psychtoolbox (Kleiner et al., 2007) on a 15.4" LCD screen positioned at eye-level, 2-3' from the patient. Auditory stimuli were presented using stereo speakers (44.1 kHz, MacBook Pro 2015).

A total of 3458 electrode contacts were implanted in patients; 2135 of these were included for analysis after excluding channels proximal to the seizure onset zone or exhibiting excessive inter-ictal spikes or line noise. Analyses were performed by first bandpass filtering the raw data of each electrode into broadband gamma activity (BGA; 70–150 Hz) following removal of line noise and its harmonics (zero-phase 2nd order Butterworth band-stop filters). Electrodes were also visually inspected for saccade artefacts. A frequency domain bandpass Hilbert transform (paired sigmoid flanks with half-width 1.5 Hz) was applied and the analytic amplitude

was smoothed (Savitzky-Golay FIR, 3rd order, frame length of 251ms; Matlab 2019b, Mathworks, Natick, MA). BGA was defined as percentage change from baseline level; -500 to -100ms before the presentation of the first word in each trial. Periods of significant activation were tested using a one-tailed t-test at each time point and were corrected for multiple comparisons with a Benjamini-Hochberg false detection rate (FDR) threshold of $q<0.05$, where the q-value denotes the standard term given to adjusted p-values for FDR significance corrected for multiple comparisons (Benjamini and Hochberg, 1995; Storey, 2011). For the grouped analysis, all electrodes were averaged within each subject and then the between-subject averages were used.

To provide statistically robust and topologically precise estimates of BGA, and to account for variations in sampling density, population-level representations were created using surface-based mixed-effects multilevel analysis (SB-MEMA) (Fischl et al., 1999; Conner et al., 2011; Kadipasaoglu et al., 2014, 2015). This method accounts for sparse sampling, outlier inferences, as well as intra- and inter-subject variability to produce population maps of cortical activity. A geodesic Gaussian smoothing filter (3mm full-width at half-maximum) was applied. Significance levels were computed at a corrected alpha-level of 0.01 using family-wise error rate corrections for multiple comparisons. The minimum criterion for the family-wise error rate was determined by white-noise clustering analysis (Monte Carlo simulations, 5000 iterations) of data with the same dimension and smoothness as that analyzed (Kadipasaoglu et al., 2014). Results were further restricted to regions with at least three patients contributing to coverage and BGA percent change exceeding 5%.

Anatomical groups of electrodes were delineated, firstly, through indexing electrodes to the closest node on the standardized cortical surface (Saad and Reynolds, 2012), and secondly, through grouping channels into parcellations determined by Human Connectome Project (HCP) space (Glasser et al., 2016). Parametric statistics were used since HCP regions of interest contained over 30 electrodes. When contrasting experimental conditions, two-sided paired t-tests were evaluated at each time point for each region and significance levels were computed at $q < 0.01$ using an FDR correction for multiple comparisons.

To generate event-related potentials (ERPs), the raw data were band pass filtered (0.1–50 Hz). Trials were averaged together and the resultant waveform was smoothed (Savitzky-Golay FIR, third-order, frame length of 251 ms). To account for differences in polarity between electrodes, ERPs were converted to root mean square (RMS), using a 50 ms sliding window. All electrodes were averaged within each subject, within ROI, and then the between-subject averages were used.

To explore the functional connectivity between ROIs, we used a generalized phase-locking
analysis to estimate the dominant spatiotemporal distributions of field activity, and the strength
of the coupling between them. Phase information was extracted from the down-sampled (200
Hz) and wide band-pass filtered data (3–50 Hz; zero-phase 8th order Butterworth band-pass
filter) using the 'generalized phase' method (Davis et al., 2020) with a single-sided Fourier
transform approach. This method captures the phase of the predominant fluctuations in the
wideband signal and minimizes filter-related distortion of the waveform. PLV was calculated
as the circular mean (absolute vector length) of the instantaneous phase difference between
each electrode pair at each time point and baselined to the pre-stimulus period -500 to -100
ms before onset of the first word. Statistics were calculated using the mean PLV of correctly
answered trials between 0 to 500 ms after second word onset, comparing against a null
distribution generated by randomly re-pairing trial recordings across the electrode pairs 500
times. Significant PLV from pre-stimulus baseline was accepted at a threshold of $p < 0.05$.
When computing conditional differences, significance was accepted at $q < 0.05$ using an FDR
correction for multiple comparisons.

Data Accessibility: The datasets generated from this research are not publicly available due
to their containing information non-compliant with HIPAA, and the human participants from
whom the data were collected have not consented to their public release. However, they are
available on request from the corresponding author.

Code Accessibility: The custom code that supports the findings of this study is available from
the corresponding author on request.

Results

Patients were presented with auditory recordings grouped randomly into three conditions:
Adjective-Noun, Pseudoword-Noun, Adjective-Pseudoword. A subsequent colored image was
presented, and patients were tasked with responding, with a button press, to whether the
image matched the phrase or not (Fig. 1A). Across the cohort, we had good coverage over
lateral and medial temporal lobe, inferior parietal lobe and inferior frontal regions, with some
coverage reaching into other portions of fronto-parietal cortex (Fig. 1B). We saw activation in
response to the auditory stimuli most prominently in superior and middle temporal regions
(Fig. 1C). Below we report results pertaining to lexicality; phrase anticipation; phrase
composition; and linguistic-visual unification.

Behavioral performance

Performance in the image matching task was highly accurate at 97 ± 3% (311 ± 11/320 trials),

with an average response time of 1599 ± 539 ms. Only correct trials were analyzed further.

Effects of lexicality

To disentangle single word semantic effects from those of combinatorial semantics, we probed

the difference in representation between words and pseudowords. We generated a surface-

based, population-level map of cortical activity using a surface-based mixed-effects multi-level

analysis (SB-MEMA) (Fischl et al., 1999; Conner et al., 2011; Kadipasaoglu et al., 2014, 2015),

a method specifically designed to account for sampling variations in iEEG and minimize effects

of outliers. An SB-MEMA contrasting adjectives and pseudowords in word position 1, and

nouns and pseudowords in word position 2 (Fig. 1D), revealed significantly greater broadband

gamma activity (BGA; 70–150 Hz) for pseudowords than words in posterior superior temporal

gyrus (pSTG) (word 1, 300–500 ms: $\beta = 0.88$; $p = .006$), posterior superior temporal sulcus

(pSTS) (word 1, 300–700 ms: $\beta = 0.30$; $p <.001$; word 2, 300–700 ms: $\beta = 0.23$; $p <.001$) and

pars triangularis (word 2, 300–700 ms: $\beta = 0.08$; $p = .004$). We found greater BGA increases

for pseudowords than words at position 2 than position 1 around pars triangularis and

surrounding frontal areas (Fig. 1F) (pars triangularis, 300–700 ms: $\beta = 0.09$; $p <.001$; pSTS,

300–500 ms: $\beta = 0.09$; $p = .008$). A Word 2–1 subtraction for both non-compositional trials

(Adjective-Pseudoword, Pseudoword-Adjective) revealed no such effects in posterior

temporal regions, and a BGA increase for nouns over frontal opercular sites (300–500 ms: β

$= 0.10$; $p = .002$) (Fig. 1G).

Figure 1: **Patient coverage map and grouped analysis for lexicality.** (A) Experimental design. Average stimuli length: adjectives (420 ± 39 ms; mean ± SD), nouns (450 ± 75 ms), pseudowords (430 ± 38 ms). (B) Group coverage map of left hemisphere electrodes included in analyses, plotted on a semi-inflated standardised N27 surface. (C) Broadband gamma activity (BGA) increases from pre-stimulus baseline (-500 to -100ms prior to first word) for all conditions from 100 to 400 ms after first word onset (threshold: %BGA > 5%, t > 1.96, patient coverage ≥ 3; $p < 0.01$ corrected). Black surfaces fell below patient coverage threshold. (D) SB-MEMA comparing words vs pseudowords. Red coloration indexes greater BGA (70–150 Hz) for pseudowords, and blue for words (same thresholds as in (C)). Top row: word position 1; bottom row: word position 2. (E) Exemplar electrodes for the words vs pseudowords analysis. Error bars (colored shading) set at one standard error. Sylvian fissure is marked with a yellow line for reference on each surface. Time 0 ms = word 1 onset. (F) SB-MEMA indicating BGA increases for pseudowords at the second word position relative to pseudowords at the first word position (time windows collapsed for both word 1 and word 2 positions). (G) SB-MEMA contrast for real words from both non-compositional conditions across the 300–500 ms window (Adjective-Pseudoword and Pseudoword-Noun). Colors indicate nouns (red) and adjectives (blue).

Compositional anticipation

We next contrasted Adjective-Noun and Pseudoword-Noun conditions at the onset of second word presentation, with only the former condition licensing any phrasal anticipation. Given that both traditional alpha (8–12 Hz) and beta (12–30 Hz) bands have been regularly implicated in

linguistic prediction and anticipatory composition (Lewis et al., 2016; Segaert et al., 2018; Hardy et al., 2021), with previous research in this domain (Gisladottir et al., 2018; Terporten et al., 2019) and neighboring domains (Piai et al., 2020) collapsing these bands, we analyzed activity across the 8–30 Hz range. During the anticipatory window for phrase formation (from -200ms to 0 ms prior to the second word onset), low frequency power (8–30 Hz) exhibited a significant conditional difference (Fig. 2). Greater alpha/beta power for anticipatory trials was found in pars triangularis ($β$ = 0.18; p = .009), ATL ($β$ = 0.15; p = .008) and STG ($β$ = 0.13; p = .008). These effects were unrelated to BGA. For comparison, we also plot the anticipatory window in BGA for -100 to 0 ms (Fig. 2B), which exhibited no clear relation with the lower frequency effects.

Figure 2: **Syntactic-semantic compositional anticipation.** (A) SB-MEMA for the anticipatory time window centred around second word onset (-200–0 ms) for low frequency alpha/beta power. Pink: Greater power for composition anticipation. Green: Greater for no anticipation (i.e., after having heard a Pseudoword at word 1 position). The same SB-MEMA thresholds as in Fig. 1C were applied. (B) SB-MEMA for BGA for -100–0 ms prior to word 2 onset. (C)-(E) Exemplar electrodes with location (C), low frequency power traces (D) and BGA traces (E) across three patients sorted by row.

 Phrase composition

The combinatorial contrast (Adjective-Noun vs [Adjective-Pseudoword + Pseudoword-Noun])

revealed greater BGA for portions of pSTS during phrase composition than non-composition

(100–300 ms: β = 0.10; p = .003) (Fig. 3A). The specific onset of this effect was around 210ms

after noun onset, with peak BGA at around 300ms (Fig. 4B). The same region exhibited

greater BGA for non-compositional trials at later time points (300–700 ms: β = 0.10; p = .001).

Certain portions of pSTS across patients displayed exclusive sensitivity to phrase

composition, and not lexicality (Fig. 3B, 3C). Other regions – pSTG and inferior frontal gyrus

(IFG) – did not show any significant BGA differences between these conditions (Fig. 4A). ERP

responses were dissociable from BGA across the ROIs plotted: A late effect for non-

composition was found in Broca's area, and a late signature was detected in temporal pole for

composition.

Figure 3: **Grouped analysis for phrase composition.** (A) SB-MEMAs for the phrase composition analysis for

BGA (70–150 Hz). Orange indexes greater BGA for phrase composition and purple indexes greater BGA for non-

composition. The same SB-MEMA thresholds as in Fig. 1C were applied. Time 0 ms = word 2 onset. (B) Exemplar

electrodes with FDR-corrected (one-tailed t-tests, $q < 0.05$) significance bars in purple plotted in native patient space. Includes compositional contrast and lexicality contrast, dissociating neighboring portions of pSTS that responded exclusively for phrase composition and not lexicality (orange dots = composition effect; green dots = lexicality effect; red dots = effect for both composition and lexicality). Electrodes that showed greater, later BGA increases for non-composition were spatially distinct from those sensitive to composition in early windows. Error bars set at one standard error. Numbers denote distinct patients (B1–B3). Time 0ms = word 1 onset; word 2 onset at 800ms. (C) Electrodes exhibiting a significant BGA contrast for compositional vs non-compositional trials, and words vs pseudowords. 39 electrodes (orange) from 12 patients exhibited an FDR-significant contrast (one-tailed t-tests, $q < 0.05$) for compositionality at some point between 0-1000 ms after the second word; 97 electrodes (green) across 17 patients for lexicality; 3 electrodes (red) for effects of both lexicality and compositionality across 3 patients.

Figure 4: **Regions of interest and their broadband high gamma signatures.**

(A) Regions of interest (left column) with representations of their broadband gamma activity (middle column) and event-related potentials (right column). Red: Compositional. Black: Non-Compositional. HCP index from top-down: posterior superior temporal gyrus (pSTG) = [A4]; temporal pole = [TGd]; inferior frontal gyrus (IFG) = [FOP4, 44, 45, IFSp, p47r, IFSa, 47l]; posterior middle temporal gyrus (pMTG) = [PHT, TE1p]. BGA traces are thresholded by $p < 0.05$ significantly active from pre-stimulus baseline (-500 to -100 ms) with a minimum of 10% BGA amplitude increase during the 100-400ms window after word 2 onset. ERPs were calculated across the four ROIs plotted on the left column, with significant condition differences being calculated across 0-1000 ms in the same (FDR-corrected) manner as the BGA plots. (B) Spectrograms and electrodes in pSTS for all active channels (HCP index:

TPOJ1; electrodes = 23, patients = 10). pSTS electrode coverage per patient: 1.9 ± 1.4 (mean ± SD). Word 2
conset was at 800 ms.

Next, we isolated regions of interest to derive cortical functional connectivity during phrase
composition. These were based either on results from our main analysis (pSTS) or on
composition effects described in the literature (inferior frontal regions, temporal pole)
(Graessner et al., 2021b). To characterize functional connectivity between these regions
during phrase composition, we computed phase-locking values (PLV) for electrode pairs
situated within pSTS with either pars triangularis or anterior temporal lobe. We computed the
generalized phase of the wideband filtered (3–50 Hz) signal that has previously been shown
to be more effective than the use of narrowband alpha or theta filters (Davis et al., 2020).

Among patients with concurrent coverage in pSTS and pars triangularis (n = 8, electrode pairs
= 231), the majority (n = 5) exhibited significantly greater PLVs for compositional than for non-
compositional trials during the 0–500 ms period after second word onset, averaging across
PLV values for each pair. In patients with joint coverage in pSTS and temporal pole (n = 8,
electrode pairs = 274), the majority (n = 6) showed greater PLVs for the same contrast during
the same time window (Fig. 5A, B). We also contrasted PLVs for compositional electrodes and
non-compositional electrodes in pSTS, for the subset of patients that had such electrodes.
This revealed that compositional electrodes in pSTS exhibited significantly greater PLVs with
their paired electrodes in temporal pole and pars triangularis than non-compositional
electrodes, with compositional trials also yielding greater peak PLV values. An exemplar
patient for pSTS-pars triangularis connectivity is plotted in Fig. 5C.

<u>Figure 5</u>: **Phase-locking between semantic composition regions of interest**. (A) Left: Average generalized phase-locking values (gPLV) for 5 patients showing greater gPLV for phrase composition relative to non-composition between pSTS (HCP index: TPOJ1) and pars triangularis (HCP index: 45, IFSa, IFSp, 47l). Right: Average gPLVs for the 6 patients showing greater phase-locking between pSTS and temporal pole (HCP index: TGd). Purple lines indicate points of significant conditional differences in gPLV values (FDR-corrected for multiple comparisons). gPLV values are plotted from pre-stimulus baseline (-500 to -100ms before first word onset). (B) Posterior temporal lobe (pSTS) gPLVs with inferior frontal gyrus (specifically, pars triangularis) (Top) and anterior temporal lobe (specifically, temporal pole) (Bottom). Left plots show the localization in native space of electrodes significantly involved (q < 0.05) in inter-regional phase-locking (3–50 Hz). Right plots show average time courses (mean ± SEM) of phase-locking value changes from baseline in phrase composition (brown) and non-composition (blue) trials. (C) Phase-locking values between pSTS and pars triangularis in an exemplar patient, contrasting PLVs for electrodes in pSTS that showed an effect of phrase composition (top plot, orange electrodes) with those in

369 pSTS that did not show an effect of composition (bottom plot, black electrodes). Since pSTS is a small ROI, only a
370 subset of our patients satisfied the criteria for this analysis: (1) exhibiting joint coverage across pSTS and pars
371 triangularis; (2) having electrodes in pSTS that did show an effect of composition, and other electrodes that did not.

Integration of linguistic and visual information

Comparing the Adjective-Noun trials which contained a matching versus non-matching
picture, during the 250–500 ms post-picture window SB-MEMAs revealed two notable effects
(Fig. 6): The anterior insula ($\beta = 0.16$; $p <.001$) and pars triangularis ($\beta = 0.09$; $p = .002$)
exhibited greater BGA for phrase-picture matches, while more dorsal frontal regions, centered
around inferior frontal sulcus ($\beta = 0.10$; $p <.001$), exhibited greater BGA for phrase-picture
non-matches. Since activity of opercular regions can be misattributed to the insula (Naidich et
al., 2004; Woolnough et al., 2019), we ensured that these effects across patients specifically
came from electrodes in insula proper by manually checking MRI reconstructions of electrode
localizations.

Figure 6: **Grouped analysis for linguistic-visual integration.**

(A) SB-MEMA in BGA for phrase-picture match (orange) and phrase-picture contrast (turqoise) increases, 250–
500 ms after picture onset (threshold: %BGA > 5%, t > 1.96, patient coverage ≥ 3; $p < 0.01$ corrected). (B) Exemplar
insula electrode.

Discussion

We localized the neural signatures of phrase comprehension using minimal adjective-noun phrases in a large patient cohort. We identified a broad portion of posterior temporal cortex as being sensitive to lexicality, and identified neighboring portions in pSTS that respond exclusively to phrase composition. This finely organized heterogeneity in responses implies a cortical topography that takes us beyond traditional structure-function mappings for higher-order syntax-semantics (Naidich et al., 2004). This mosaic architecture has been ascribed to sensory cortices, and to the best of our knowledge has not been shown for higher-level processing in associative cortices (Fox et al., 2020; Tsao, 2020). We speculate that such organization may be a foundational principle underpinning human language. In this section, we discuss these results in the context of previous findings, and evaluate alternative explanations.

Anterior portions of IFG exhibited greater BGA for pseudowords at word 2 position relative to word 1, possibly indexing increased unification demands with the preceding adjective (Hagoort, 2005, 2013). This implies a role for IFG in the unification of more or less expected continuations, in agreement with earlier results (Hagoort, 2004).

Our results address current concerns in the literature with respect to the spatiotemporal dynamics of phrase composition: While IFG activity indexes some aspect of phrase anticipation and phrase-picture matching, it is pSTS that appears to encode the earliest responsiveness to meaningful phrases. These results are supported by lesion-behavior mapping research implicating anterior IFG in executive control for decisions on semantic composition, and broad portions of MTG in representations of individual phrase constituents (Graessner et al., 2021a).

Lastly, anterior insula, pars triangularis and IFS subserve the integration of linguistic input with visual referents. Anterior insula and the IFS have been argued to be the convergence zones of the ventral and dorsal attentional networks (Cazzoli et al., 2021), and our results align with previous models (Willems et al., 2008).

Lexicality

Our results replicate prior studies in which BGA localized to p/mSTG and pSTS tracks lexicality (Tanji et al., 2005; Humphries et al., 2006; Canolty et al., 2007). This activity may index the bundling of sub-lexical features to yield coherent word-level interpretations of auditory stimuli. Our discovery of greater BGA increases for pseudowords than words at position 2 than position 1 around pars triangularis and surrounding frontal areas (Fig. 1F) suggests involvement of this region in effortful lexical processing to facilitate semantic unification (Hagoort, 2005, 2013). Greater BGA in posterior temporal regions for pseudowords in position 1 may speak to an effect of auditory repetition suppression, or it may indicate greater

424 processing effort as a function of pseudoword phrase position: Greater *lexical access effort* at
425 position 1 (pSTS), and greater *unification effort* (IFG) in position 2.

Anticipatory response

Based on our results, we suggest that low frequency power in IFG and ATL indexes the
preparation of a syntactic slot to be filled by predicted upcoming nominal content. Our findings
are in line with the notion that beta oscillations can index the construction and maintenance of
sentence-level meaning (Lewis et al., 2016), and the claim that alpha/beta increases can index
aspects of syntactic anticipation and category generation (Benítez-Burraco and Murphy, 2019;
Murphy, 2020).

Minimal phrase composition

We found that neighboring portions of pSTS can exclusively code either for lexicality or phrase
composition. While a large area of posterior and superior temporal cortex was sensitive to
lexicality, a narrower portion was recruited exclusively to code for phrase structure (210–300
ms), generating from simple lexical meanings representations of 'complex meaning' (Hagoort,
2020). The same portion of pSTS exhibited greater BGA for non-compositional trials (i.e.,
pseudowords) at later time points (300-700 ms), implicating it in effortful phrase structure
derivations or late-stage lexical search effort/reanalysis. In addition, the greater BGA for
phrase composition in pSTS was crucially found across different electrodes to those sites that
showed greater BGA for late-stage pseudoword processing, which we believe contributes to
a cortical mosaic. ERP responses were also dissociable from BGA: A late effect for non-
composition was found in Broca's area, likely due to greater attempted lexical access for
pseudowords, while a late signature was detected in temporal pole for composition, potentially
related to late-stage conceptual access.

Phrase composition was also marked via increased functional connectivity between pSTS
electrodes implicated in composition and both inferior frontal regions and temporal pole. This
appears to reflect network-level interactions seemingly required for basic phrase formation
(Baggio and Hagoort, 2011; Schoffelen et al., 2017). While the duration of composition effects
lasted approximately 500 ms, there were nevertheless also late-stage periods of brief (~80
ms), reversed effects. As such, future intracranial research involving either more dense or fine-
grained coverage could provide a clearer picture.

Anatomical connectivity has been elaborated by white matter pathways that connect pSTS
with IFG and ATL (Glasser and Rilling, 2008; Figley et al., 2017; Sarubbo et al., 2020). We
theorize that the formation of phrase structure in pSTS feeds categorial information to

conceptual interfaces in ATL and memory/control interfaces in pars triangularis. One such piece of information would be the phrase category/label.

Previous intracranial research found that the lower bank of the STS is involved in lexico-semantic processing (Nourski et al., 2021), and our results appear to suggest greater involvement of the lower bank of the pSTS in semantic composition than the cytoarchitechtonically distinct upper bank (Zheng et al., 2010). Future work utilizing even higher scale recording techniques, such as single-unit recordings (Bitterman et al., 2008), will more comprehensively address this issue.

One of the main candidates for phrase composition from previous research, ATL, was implicated via late-stage ERPs. ATL activity has been found to be delayed until after intra-lexical morphological composition (Flick et al., 2018). This implies the existence of composition-related activity independent of semantic and orthographic processing. Our findings suggest that pSTS might be one such region, although since we only presented auditory stimuli we cannot make any stronger claims.

Our findings are concordant with recent results and models implicating the pSTS in phrase composition (Nelson et al., 2017; Flick and Pylkkänen, 2020; Matchin and Hickok, 2020; Murphy, 2020; Law and Pylkkänen, 2021; Matar et al., 2021); producing hierarchically organized functional morphemes (Lee et al., 2018); sign language comprehension (Trettenbrein et al., 2021); cross-linguistic reading competence (Feng et al., 2020).

Existing models have posited distinct neural sites across gross portions of the cortical mantle for lexicality and basic composition. Typically, this is a large scale "posterior vs anterior temporal" (Friederici, 2012) or "dorsal vs ventral stream" distinction (Bornkessel-Schlesewsky and Schlesewsky, 2013), or a distinction between ATL (conceptual-semantic) and pMTG (hierarchical structure) (Matchin & Hickok, 2020). In contrast, we report overlapping functionality at a much smaller scale in the pSTS.

We have claimed that pSTS is "tiled" in a mosaic-like structure coding for lexicality and phrasal meaning. However, there is an alternative account invoking pseudoword and task-related processing effort. We believe that it is likely that our late-stage (300-700 ms) signature of greater BGA for non-compositional trials in pSTS can be explained by general difficulties with pseudowords. Yet, the effort-related account seems unrelated to our early (100-300 ms) BGA increases for compositional trials; we found no clear posterior temporal differences for real words in non-compositional trials, which the effort-related account would predict (due to nouns being presented after pseudowords); patients that showed early BGA increases for composition also showed distinct signatures of non-composition processing effort in neighboring (but distinct) sites in pSTS and surrounding cortex (Fig. 3). A comparison of real

word processing in the non-compositional trials (Fig. 1G) revealed no clear effects in posterior temporal regions, only a BGA increase for nouns over frontal opercular sites, suggesting any degree of task-related effort was isolated to late-stage pseudoword processing.

We have provided a more spatiotemporally reliable profile of basic semantic composition than previous non-invasive methodologies, resolving some conflicts in the literature concerning the variety of effect timings documented. In addition, we believe that our results point to the need to move beyond simpler hypotheses about pSTS either being involved in phrase composition or not: Instead, it appears to be a question of *when* and *over which specific portions of tissue* (e.g., on the order of 10mm). We believe that recordings with higher spatial resolution may reveal greater insights into the operations of this cortical mosaic that we have documented.

While our paradigm only allows us to make direct claims about semantic composition, we suspect that portions of pSTS are also implicated more narrowly in minimal syntax, however the functional connectome of the syntax network may differ from that of the semantic compositional network. Given that recent research has revealed substantial overlap in the distribution of the fronto-temporal language network for speakers of 45 languages across 11 language families (Ayyash et al., 2021), and given that the computational process our paradigm isolated was generic, we also expect that pSTS activity codes for meaningful phrase structures across languages other than English. The specific neural signatures of basic phrase structure composition isolated here represent an elementary computation in comprehending natural language that can be probed further in future intracranial research.

References

Amunts K, Zilles K (2015) Architectonic Mapping of the Human Brain beyond Brodmann. Neuron 88:1086–1107.

Antonucci S, Beeson P, Labiner D, Rapcsak S (2008) Lexical retrieval and semantic knowledge in patients with left inferior temporal lobe lesions. Aphasiology 22:281–304.

Artoni F, D'Orio P, Catricalà E, Conca F, Bottoni F, Pelliccia V, Sartori I, Russo G Lo, Cappa SF, Micera S, Moro A (2020) High gamma response tracks different syntactic structures in homophonous phrases. Scientific Reports 10:7537.

Arya R (2019) Similarity of spatiotemporal dynamics of language-related ECoG high-gamma modulation in Japanese and English speakers. Clinical Neurophysiology 130:1403–1404.

Ayyash D, Malik-Moraleda S, Gallée J, Affourtit J, Hoffman M, Mineroff Z, Jouravlev O, Fedorenko E (2021) The universal language network: A cross-linguistic investigation spanning 45 languages and 11 language families. Neuroscience. Available at: http://biorxiv.org/lookup/doi/10.1101/2021.07.28.454040 [Accessed October 12, 2021].

Baggio G, Hagoort P (2011) The balance between memory and unification in semantics: A dynamic account of the N400. Language and Cognitive Processes 26:1338–1367.

Bastiaansen M, Hagoort P (2015) Frequency-based Segregation of Syntactic and Semantic Unification during Online Sentence Level Language Comprehension. Journal of Cognitive Neuroscience 27:2095–2107.

Bemis DK, Pylkkänen L (2011) Simple composition: A magnetoencephalography investigation into the comprehension of minimal linguistic phrases. Journal of Neuroscience 31:2801–2814.

Bemis DK, Pylkkänen L (2013) Basic linguistic composition recruits the left anterior temporal lobe and left angular gyrus during both listening and reading. Cerebral Cortex.

Benítez-Burraco A, Murphy E (2019) Why Brain Oscillations Are Improving Our Understanding of Language. Front Behav Neurosci 13:190.

Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B (Methodological) 57:289–300.

Berwick RC, Stabler EP, Berwick RC, Stabler EP (2019) Minimalist parsing. Oxford: Oxford University Press.

Bitterman Y, Mukamel R, Malach R, Fried I, Nelken I (2008) Ultra-fine frequency tuning revealed in single neurons of human auditory cortex. Nature 451:197–201.

Bonner MF, Price AR (2013) Where Is the Anterior Temporal Lobe and What Does It Do? Journal of Neuroscience 33:4213–4215.

Bornkessel-Schlesewsky I, Schlesewsky M (2013) Reconciling time, space and function: A
new dorsal–ventral stream model of sentence comprehension. Brain and Language
125:60–76.

Bozic M, Fonteneau E, Su L, Marslen-Wilson WD (2015) Grammatical analysis as a
distributed neurobiological function. Human Brain Mapping 36:1190–1201.

Brennan J, Pylkkänen L (2012) The time-course and spatial distribution of brain activity
associated with sentence processing. NeuroImage 60:1139–1148.

Brennan JR, Stabler EP, Van Wagenen SE, Luh WM, Hale JT (2016) Abstract linguistic
structure correlates with temporal activity during naturalistic comprehension. Brain
and Language 157–158:81–94.

Brysbaert M, New B, Keuleers E (2012) Adding part-of-speech information to the SUBTLEX-
US word frequencies. Behavior Research Methods 44:991–997.

Buzsáki G, Watson BO (2012) Brain rhythms and neural syntax: implications for efficient
coding of cognitive content and neuropsychiatric disease. Dialogues in Clinical
Neuroscience 14:345–367.

Canolty RT, Soltani M, Dalal SS, Edwards E, Dronkers NF, Nagarajan SS, Kirsch HE,
Barbaro NM, Knight RT (2007) Spatiotemporal Dynamics of Word Processing in the
Human Brain. Frontiers in Neuroscience 1:185–196.

Cazzoli D, Kaufmann BC, Paladini RE, Müri RM, Nef T, Nyffeler T (2021) Anterior insula and
inferior frontal gyrus: where ventral and dorsal visual attention systems meet. Brain
Communications.

Chomsky N, Gallego AJ, Ott D (2019) Generative grammar and the faculty of language:
insights, questions, and challenges. Catalan Journal of Linguistics Special
Issue:226–261.

Conner CR, Ellmore TM, Pieters TA, di Sano MA, Tandon N (2011) Variability of the
relationship between electrophysiology and BOLD-fMRI across cortical regions in
humans. Journal of Neuroscience 31:12855–12865.

Conner CR, Kadipasaoglu CM, Shouval HZ, Hickok G, Tandon N (2019) Network dynamics
of Broca's area during word selection Hinojosa JA, ed. PLOS ONE 14:e0225756.

Cox RW (1996) AFNI: Software for Analysis and Visualization of Functional Magnetic
Resonance Neuroimages. Computers and Biomedical Research 29:162–173.

Dale AM, Fischl B, Sereno MI (1999) Cortical Surface-Based Analysis: I. Segmentation and
Surface Reconstruction. NeuroImage 9:179–194.

Davis ZW, Muller L, Martinez-Trujillo J, Sejnowski T, Reynolds JH (2020) Spontaneous
travelling cortical waves gate perception in behaving primates. Nature 587:432–436.

Ding N, Melloni L, Zhang H, Tian X, Poeppel D (2016) Cortical tracking of hierarchical
linguistic structures in connected speech. Nat Neurosci 19:158–164.

Duffau H, Moritz-Gasser S, Mandonnet E (2014) A re-examination of neural basis of
language processing: Proposal of a dynamic hodotopical model from data provided
by brain stimulation mapping during picture naming. Brain and Language 131:1–10.

Feng X, Altarelli I, Monzalvo K, Ding, GuoshengRamus F, Shu H, Dehaene S, Meng X,
Dehaene-Lambertz G (2020) A universal reading network and its modulation by
writing system and reading ability in French and Chinese children. eLife 9:e54591.

Figley TD, Mortazavi Moghadam B, Bhullar N, Kornelsen J, Courtney SM, Figley CR (2017)
Probabilistic white matter atlases of human auditory, basal ganglia, language,
precuneus, sensorimotor, visual and visuospatial networks. Frontiers in Human
Neuroscience 11:1–12.

Fischl B, Sereno MI, Dale AM (1999) Cortical surface-based analysis: II. Inflation, flattening,
and a surface-based coordinate system. NeuroImage 9:195–207.

Flick G, Oseki Y, Kaczmarek AR, Al Kaabi M, Marantz A, Pylkkänen L (2018) Building words
and phrases in the left temporal lobe. Cortex 106:213–236.

Flick G, Pylkkänen L (2020) Isolating syntax in natural language: MEG evidence for an early
contribution of left posterior temporal cortex. Cortex 127:42–57.

Flinker A, Chang EF, Barbaro NM, Berger MS, Knight RT (2011) Sub-centimeter language
organization in the human temporal lobe. Brain and Language 117:103–109.

Forseth KJ, Kadipasaoglu CM, Conner CR, Hickok G, Knight RT, Tandon N (2018) A lexical
semantic hub for heteromodal naming in middle fusiform gyrus. Brain 141:2112–
2126.

Fox NP, Leonard M, Sjerps MJ, Chang EF (2020) Transformation of a temporal speech cue
to a spatial neural code in human auditory cortex. eLife 9.

Friederici AD (2012) The cortical language circuit: from auditory perception to sentence
comprehension. Trends in Cognitive Sciences 16:262–268.

Gastaldon S, Arcara G, Navarrete E, Peressotti F (2020) Commonalities in alpha and beta
neural desynchronizations during prediction in language comprehension and
production. Cortex 133:328–345.

Gisladottir RS, Bögels S, Levinson SC (2018) Oscillatory Brain Responses Reflect
Anticipation during Comprehension of Speech Acts in Spoken Dialog. Front Hum
Neurosci 12:34.

Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E, Ugurbil K,
Andersson J, Beckmann CF, Jenkinson M, Smith SM, Van Essen DC (2016) A multi-
modal parcellation of human cerebral cortex. Nature 536:171–178.

Glasser MF, Rilling JK (2008) DTI Tractography of the Human Brain's Language Pathways.
Cerebral Cortex 18:2471–2482.

Graessner A, Zaccarella E, Friederici AD, Obrig H, Hartwigsen G (2021a) Dissociable
contributions of frontal and temporal brain regions to basic semantic composition.
Brain Communications 3:fcab090.

Graessner A, Zaccarella E, Hartwigsen G (2021b) Differential contributions of left-
hemispheric language regions to basic semantic composition. Brain Structure and
Function.

Hagoort P (2003) How the brain solves the binding problem for language: A
neurocomputational model of syntactic processing. NeuroImage 20:S18–S29.

Hagoort P (2004) Integration of Word Meaning and World Knowledge in Language
Comprehension. Science 304:438–441.

Hagoort P (2005) On Broca, brain, and binding: a new framework. Trends in Cognitive
Sciences 9:416–423.

Hagoort P (2013) MUC (Memory, Unification, Control) and beyond. Frontiers in Psychology
4.

Hagoort P (2017) The core and beyond in the language-ready brain. Neuroscience &
Biobehavioral Reviews 81:194–204.

Hagoort P (2020) The meaning-making mechanism(s) behind the eyes and between the
ears. Philosophical transactions of the Royal Society of London Series B, Biological
sciences 375:20190301.

Hardy SM, Jensen O, Wheeldon L, Mazaheri A, Segaert K (2021) Modulation in Alpha Band
Activity Reflects Syntax Composition: An MEG Study of Minimal Syntactic Binding.
Neuroscience. Available at: http://biorxiv.org/lookup/doi/10.1101/2021.07.09.451797
[Accessed October 15, 2021].

Hovsepyan S, Olasagasti I, Giraud A-L (2020) Combining predictive coding and neural
oscillations enables online syllable recognition in natural speech. Nature
Communications 11:3117.

Humphries C, Binder JR, Medler DA, Liebenthal E (2006) Syntactic and semantic
modulation of neural activity during auditory sentence comprehension. Journal of
Cognitive Neuroscience 18:665–679.

Johnson EL, Kam JWY, Tzovara A, Knight RT (2020) Insights into human cognition from
intracranial EEG: A review of audition, memory, internal cognition, and causality.
Journal of Neural Engineering 17:051001.

Kadipasaoglu CM, Baboyan VG, Conner CR, Chen G, Saad ZS, Tandon N (2014) Surface-
based mixed effects multilevel analysis of grouped human electrocorticography.
NeuroImage 101:215–224.

Kadipasaoglu CM, Forseth K, Whaley M, Conner CR, Rollo MJ, Baboyan VG, Tandon N
(2015) Development of grouped icEEG for the study of cognitive processing.
Frontiers in Psychology 6.

Keitel A, Gross J, Kayser C (2018) Perceptually relevant speech tracking in auditory and
motor cortex reflects distinct linguistic features Bizley J, ed. PLoS Biol 16:e2004473.

Keitel A, Ince RAA, Gross J, Kayser C (2017) Auditory cortical delta-entrainment interacts
with oscillatory power in multiple fronto-parietal networks. NeuroImage 147:32–42.

Kleiner M, Brainard D, Pelli D, Ingling A, Murray R, Broussard C (2007) What's new in
psychtoolbox-3. Perception 36:1–16.

Kochari AR, Lewis AG, Schoffelen J-M, Schriefers H (2021) Semantic and syntactic
composition of minimal adjective-noun phrases in Dutch: An MEG study.
Neuropsychologia 155:107754.

Kohlhase K, Zöllner JP, Tandon N, Strzelczyk A, Rosenow F (2021) Comparison of
minimally invasive and traditional surgical approaches for refractory mesial temporal
lobe epilepsy: A systematic review and meta-analysis of outcomes. Epilepsia
62:831–845.

Lambon Ralph MA, Ehsan S, Baker GA, Rogers TT (2012) Semantic memory is impaired in
patients with unilateral anterior temporal lobe resection for temporal lobe epilepsy.
Brain 135:242–258.

Law R, Pylkkänen L (2021) Lists with and without Syntax: A New Approach to Measuring the
Neural Processing of Syntax. The Journal of Neuroscience 41:2186–2196.

Lee DK, Fedorenko E, Simon M V., Curry WT, Nahed B V., Cahill DP, Williams ZM (2018)
Neural encoding and production of functional morphemes in the posterior temporal
lobe. Nature Communications 9:1877.

Leivada E, Murphy E (2021) Mind the (terminological) gap: 10 misused, ambiguous, or
polysemous terms in linguistics. Ampersand 8:100073.

Leszczyński M, Barczak A, Kajikawa Y, Ulbert I, Falchier AY, Tal I, Haegens S, Melloni L,
Knight RT, Schroeder CE (2020) Dissociation of broadband high-frequency activity
and neuronal firing in the neocortex. Science Advances 6:eabb0977.

Lewis AG, Schoffelen J-M, Schriefers H, Bastiaansen M (2016) A Predictive Coding
Perspective on Beta Oscillations during Sentence-Level Language Comprehension.
Frontiers in Human Neuroscience 10:83.

Lopopolo A, van den Bosch A, Petersson K-M, Willems RM (2021) Distinguishing Syntactic
Operations in the Brain: Dependency and Phrase-Structure Parsing. Neurobiology of
Language 2:152–175.

Mai G, Minett JW, Wang WS-Y (2016) Delta, theta, beta, and gamma brain oscillations index
levels of auditory sentence processing. NeuroImage 133:516–528.

Matar S, Dirani J, Marantz A, Pylkkänen L (2021) Left posterior temporal cortex is sensitive
to syntax within conceptually matched Arabic expressions. Scientific Reports 11.

Matchin W, Hickok G (2020) The Cortical Organization of Syntax. Cerebral Cortex 30:1481–
1498.

McCarty MJ, Woolnough O, Mosher JC, Seymour J, Tandon N (2021) The Listening Zone of
Human Electrocorticographic Field Potential Recordings. Neuroscience. Available at:
http://biorxiv.org/lookup/doi/10.1101/2021.10.22.465519 [Accessed November 7,
2021].

Murphy E (2015) Labels, cognomes, and cyclic computation: An ethological perspective.
Frontiers in Psychology 6.

Murphy E (2020) The Oscillatory Nature of Language. Cambridge: Cambridge University
Press.

Naidich TP, Kang E, Fatterpekar GM, Delman BN, Gultekin SH, Wolfe D, Ortiz O, Yousry I,
Weismann M, Yousry TA (2004) The insula: anatomic study and MR imaging display
at 1.5 T. AJNR Am J Neuroradiol 25:222–232.

Nelson MJ, El Karoui I, Giber K, Yang X, Cohen L, Koopman H, Cash SS, Naccache L, Hale
JT, Pallier C, Dehaene S (2017) Neurophysiological dynamics of phrase-structure
building during sentence processing. Proceedings of the National Academy of
Sciences of the United States of America 114:E3669–E3678.

Neufeld C, Kramer SE, Lapinskaya N, Heffner CC, Malko A, Lau EF (2016) The
electrophysiology of basic phrase building. PLoS ONE 11:1–22.

Nourski KV, Steinschneider M, Rhone AE, Kovach CK, Banks MI, Krause BM, Kawasaki H,
Howard MA (2021) Electrophysiology of the Human Superior Temporal Sulcus during
Speech Processing. Cerebral Cortex 31:1131–1148.

Olman CA, Davachi L, Inati S (2009) Distortion and Signal Loss in Medial Temporal Lobe
García AV, ed. PLoS ONE 4:e8160.

Packard PA, Steiger TK, Fuentemilla L, Bunzeck N (2020) Neural oscillations and event-
related potentials reveal how semantic congruence drives long-term memory in both
young and older humans. Scientific Reports.

Piai V, Klaus J, Rossetto E (2020) The lexical nature of alpha-beta oscillations in context-
driven word production. Journal of Neurolinguistics 55:100905.

Pieters TA, Conner CR, Tandon N (2013) Recursive grid partitioning on a cortical surface
model: an optimized technique for the localization of implanted subdural electrodes.
Journal of Neurosurgery 118:1086–1097.

Pylkkänen L (2020) Neural basis of basic composition: what we have learned from the red-
boat studies and their extensions. Philosophical transactions of the Royal Society of
London Series B, Biological sciences 375:20190299.

Rollo PS, Rollo MJ, Zhu P, Woolnough O, Tandon N (2020) Oblique trajectory angles in
robotic stereo-electroencephalography. Journal of Neurosurgery:1–10.

Saad ZS, Reynolds RC (2012) SUMA. NeuroImage 62:768–773.

Sarubbo S, Tate M, De Benedictis A, Merler S, Moritz-Gasser S, Herbet G, Duffau H (2020)
Mapping critical cortical hubs and white matter pathways by direct electrical
stimulation: an original functional atlas of the human brain. NeuroImage 205:116237.

Schell M, Zaccarella E, Friederici AD (2017) Differential cortical contribution of syntax and
semantics: An fMRI study on two-word phrasal processing. Cortex 96:105–120.

Schoffelen J-M, Hultén A, Lam N, Marquand AF, Uddén J, Hagoort P (2017) Frequency-
specific directed interactions in the human brain network for language. Proceedings
of the National Academy of Sciences 114:8083–8088.

Segaert K, Mazaheri A, Hagoort P (2018) Binding language: structuring sentences through
precisely timed oscillatory mechanisms. European Journal of Neuroscience 48:2651–
2662.

Storey JD (2011) False Discovery Rate. In: International Encyclopedia of Statistical Science
(Lovric M, ed), pp 504–508. Berlin, Heidelberg: Springer Berlin Heidelberg. Available
at: http://link.springer.com/10.1007/978-3-642-04898-2_248 [Accessed October 14,
2021].

Tanji K, Suzuki K, Delorme A, Shamoto H, Nakasato N (2005) High-frequency γ-band
activity in the basal temporal cortex during picture-naming and lexical-decision tasks.
Journal of Neuroscience 25:3287–3293.

Terporten R, Schoffelen JM, Dai B, Hagoort P, Kösem A (2019) The Relation between
Alpha/Beta Oscillations and the Encoding of Sentence induced Contextual
Information. Scientific Reports 9.

Tong BA, Esquenazi Y, Johnson J, Zhu P, Tandon N (2020) The Brain is Not Flat:
Conformal Electrode Arrays Diminish Complications of Subdural Electrode
Implantation, A Series of 117 Cases. World Neurosurgery 144:e734–e742.

Trettenbrein PC, Papitto G, Friederici AD, Zaccarella E (2021) Functional neuroanatomy of
language without speech: An ALE meta-analysis of sign language. Human Brain
Mapping 42:699–712.

Tsao DY (2020) Does finding a face cell tell us anything much at all? Progress in
Neurobiology 195:101925.

Uno T, Kawai K, Sakai K, Wakebe T, Ibaraki T, Kunii N, Matsuo T, Saito N (2015)
Dissociated Roles of the Inferior Frontal Gyrus and Superior Temporal Sulcus in
Audiovisual Processing: Top-Down and Bottom-Up Mismatch Detection Pouratian
NN, ed. PLoS ONE 10:e0122580.

Willems RM, Özyürek A, Hagoort P (2008) Seeing and Hearing Meaning: ERP and fMRI
Evidence of Word versus Picture Integration into a Sentence Context. Journal of
Cognitive Neuroscience 20:1235–1249.

Wilson SM, DeMarco AT, Henry ML, Gesierich B, Babiak M, Mandelli ML, Miller BL, Gorno-
Tempini ML (2014) What role does the anterior temporal lobe play in sentence-level
processing? neural correlates of syntactic processing in semantic variant primary
progressive aphasia. Journal of Cognitive Neuroscience 26:970–985.

Woolnough O, Forseth KJ, Rollo PS, Tandon N (2019) Uncovering the functional anatomy of
the human insula during speech. eLife 8:1–14.

Zhang L, Pylkkänen L (2018) Semantic composition of sentences word by word: MEG
evidence for shared processing of conceptual and logical elements.
Neuropsychologia 119:392–404.

Zheng ZZ, Wild C, Trang HP (2010) Spatial Organization of Neurons in the Superior
Temporal Sulcus. Journal of Neuroscience 30:1201–1203.