1

7   EEG biomarkers of free recall

9   Katerman, B. S., Li, Y., Pazdera, J. K., Keane, C., and Kahana, M. J.

10   University of Pennsylvania

23   Author Note

24   The authors gratefully acknowledge support from National Institutes of Health

25   grant MH55687. Correspondence concerning this should be addressed to Michael

26   J. Kahana (kahana@psych.upenn.edu).

28   Abstract

Brain activity in the moments leading up to successful verbal recall provide a

window into the cognitive processes underlying memory retrieval. But these same

recordings also subsume neural signals unrelated to mnemonic retrieval, such as

response-related motor activity. Here we examined spectral EEG biomarkers of

successful recall under an extreme manipulation of mnemonic demands: subjects

either recalled items after a few seconds or several days. This manipulation

isolated EEG components specifically related to episodic retrieval.   Theta and

gamma power (4-8 Hz and 40-128 Hz respectively) increased immediately prior to

long-delay recall, whereas 8-20 Hz power decreased.   A direct comparison of long-

delay and immediate recall revealed a nearly identical pattern, indicating that these

spectral biomarkers of successful retrieval reflect memory-specific processes.

Ruling out a confound of motor related activity, these results identify theta and

gamma activity as biomarkers of successful episodic memory retrieval.

Keywords: **theta, gamma, alpha, retrieval, reinstatement**

45   EEG biomarkers of free recall

47   **Introduction**

48   Our ability to recall past experiences is the hallmark of episodic memory. In laboratory

49   studies, lists of discrete nameable items, such as words, serve as sets of mini-experiences, and the act

50   of recall is the motor output (vocalization or typing) of the remembered items. The magic of memory

51   retrieval happens during the moments leading up to the motor output, but this period also includes

52   other non-mnemonic signals, such as the planning of a motor response and the activation of semantic

53   and perceptual representations associated with the remembered item.

54   Prior research using intracranial EEG recordings has uncovered a network of brain regions in

which increased high frequency activity and concomitant decreases in lower frequency activity mark

the moments leading up to spontaneous verbal recall as compared with matched periods of silence

during the recall period (Burke, Sharan, et al., 2014; Greenberg, Burke, Haque, Kahana, & Zaghloul,

2015; Solomon et al., 2017). In some cortical sub-regions and for some retrieval contrasts, low-

frequency (3-8 Hz) theta activity increases during the putative retrieval period (see Herweg et al., 2016,

for a review). Researchers would ideally like to use these recall biomarkers to probe the dynamics of

memory retrieval in the absence of motor output.   Such signals could, for example, be used to study

the hypothesized role of covert retrieval in memory consolidation and other learning processes.

However, the interpretation of these recall biomarkers is problematized by the concomitant presence

of non-mnemonic neural processes that accompany recall even under minimal memory demands.

65   Comparisons of correct and incorrect recalls (intrusions) offers a potential solution (Long et

al., 2017) These studies have shown, for example, that hippocampal high-frequency activity increases

moreso prior to correct recalls than prior to intrusion errors. As incorrect recalls presumably involve

identical, or at least very similar, motor planning activity, differences between these conditions more

likely reflect mnemonic retrieval.   However, intrusions may show similar signals to those of successful

recall as the subject is actively engaged in recall of a word they believe to have been previously

presented. As such, these contrasts may mask important neural correlates of memory retrieval that

appear similarly for both true and false memories.

73   The present study sought to elucidate the biomarkers of episodic memory retrieval by

74   examining recall of words under an extreme manipulation of retrieval demands: In an immediate recall

75   condition, subjects recalled a single just-presented word after a brief delay.   In a long-delayed recall

76   test, subjects attempted to recall items learned across multiple days prior to the test day (i.e., a

77   minimum of 16 hours prior to the recall period). Over the course of ten experimental sessions

78   administered on different days, subjects contributed data in each of these conditions.

80   **Method**

81   **Subjects**

82   Fifty-seven young adults (ages 18-35), recruited among the students and staff at the

University of Pennsylvania and neighboring institutions, each contributed 10 sessions of immediate

word recall data and five sessions of delayed recall data. All subjects provided informed consent to

participate in our study, which was approved by the Institutional-Review Board at the University of

Pennsylvania. Subjects were excluded (n=17) if they did not complete at least 7 sessions or if they had

an average rate of recall lower than 70% during the immediate recall task. The data reported here came

from Experiment 5 of the Penn Electrophysiology of Encoding and Retrieval Study (PEERS). This is the

first paper to report data from PEERS-Experiment 5.

Data Availability

92   All PEERS data, including the full dataset reported and analyzed in the present manuscript,

93   may be freely downloaded from our public repository http://memory.psych.upenn.edu/data. Analysis

94   code for this manuscript is also available at the same URL.

96   **Experimental Task and Behavioral Analyses**

97   Each of the first five sessions (henceforth, *Phase I*) consisted of one block of 10 practice trials,

followed by 24 blocks of 24 trials each. Each block began with a 10-second countdown. After the

countdown was complete, the first trial of the block began. On each trial, a black screen was shown for

a jittered 1000--1600 ms (uniformly distributed), after which a single word appeared onscreen in white

text for 1200--1800 ms (uniformly distributed). Following presentation, the screen went blank again

and subjects were instructed to pause briefly, and then vocalize the word they had just seen. If they

began speaking within 1.0 seconds of word offset, the message "Too fast." appeared on the screen in

red text. By avoiding these messages subjects could increase the size of their bonus payment.   After

the subject finished speaking, a tone sounded, marking the end of the current trial. Speech was

detected using a volume amplitude threshold. In addition to the 10-second countdown between blocks,

two 2-minute mid-session breaks were administered after block eight and block 16. Phase II of the

experiment began on the day of the sixth session and continued to the final session of the experiment.

In Phase II the practice block and 24 experimental blocks were preceded by a 10-minute initial

externalized free recall period. Subjects were instructed to recall as many words as possible from the

previous sessions in any order, while also vocalizing any additional words that come to mind in their

attempt to recall these items (e.g., Kahana, Dolan, Sauder, & Wingfield, 2005; Lohnas, Polyn, & Kahana,

2015; Zaromb et al., 2006). Our lists comprised the same 576-word pool as used in the PEERS4 study

(Aka, Phan, & Kahana, 2020; Kahana, Aggarwal, & Phan, 2018; Weidemann & Kahana, 2020). Subjects

saw the same 576 words in each of their 10 sessions, but the ordering of these words was randomized

for each session.

EEG Post-Processing and Spectral Decomposition

119   We recorded electroencephalographic (EEG) data using a 128-channel Biosemi system with a

2048 Hz sampling rate. We applied the following preprocessing steps to the data from each session.

First, we applied a 0.5 Hz highpass filter to eliminate baseline drift. We then partitioned the recording

into thirds by splitting at the end of each mid-session break. For each partition of the data, we identified

bad channels as those with extremely high or low ($|z|>3$) log-variance, or which had an extremely high

($z>3$) Hurst exponent relative to other channels. We identified bad channels separately for each

partition, as problematic electrodes were often corrected during mid-session breaks.[1] We then dropped

the bad channels from their respective partitions and applied a common average reference scheme.

We next performed independent component analysis (ICA) on each of the three partitions to

decompose the data into 128-(n+1) components, where n is the number of bad channels that were

dropped, and used the localized component filtering method of DelPozo-Banos and Weidemann (2017)

to filter artifactual time points from the components. Data points were identified as artifacts if they

exceeded that component's interquartile range by three times the magnitude of that range. We then

reconstructed the original channels from the cleaned components, interpolated bad channels using

spherical splines, and applied a fourth-order Butterworth notch filter at 60 Hz to eliminate electrical line

noise.

135   We used a multitaper method (MNE-Python software package; Gramfort et al., 2013,2014)

136   to estimate spectral power at each electrode over 4--128 Hz and log transformed the resulting signal.

---

[1] Recordings during break periods were frequently noisy as a result of these adjustments and participant
movement, and were therefore excluded when calculating variances and Hurst exponents, as well as when
performing ICA and calculating artifact thresholds for localized component filtering.

Electrodes were grouped into regions of interest (see Figure 2) and the corresponding powers averaged

for each frequency. We spaced frequencies every 2 Hz in the range of 4--26 Hz, and every 6 Hz within

26--128 Hz, resulting in 29 frequencies of interest. We avoided the Morlet wavelet method, as

convolving low frequency wavelets using buffer periods may allow speech artifacts to intrude in the

power estimates of intervals just prior to subjects' vocal responses. We used a 500 ms moving window

centered at multiple time-points relative to the start of recall events (i.e., vocalization of a recalled item),

with a 50 ms step size. To minimize potential artifacts from pre-motor activity, we extracted spectral

patterns in a 500 ms window ending 250 ms prior to the annotated time of speech onset unless

otherwise noted. We excluded from our analysis recall events that occurred within 1500 ms of the onset

of the prior recalled item. In addition to the interval preceding vocalization, we identified 1000 ms

"deliberation" periods of silence during the delayed recall test that did not overlap with a preceding

vocalization (i.e., within 500 ms of vocalization onset) or a subsequent retrieval interval of interest.

Deliberation periods for both immediate and delayed recall contrasts were matched to recall events

using linear sum optimization, which minimizes the total time difference between recall events and

matched deliberation periods, and were constrained to events between the first and last delayed recalls

within a session.

During delayed recall, we defined successful memory events as the memory search intervals

immediately preceding recalls of list items (correct recalls). Unsuccessful memory search periods

preceded recall errors (i.e., intrusions from outside of the word pool: extra-list intrusions). To identify

spectral features specific to successful recall on the delayed recall test, we performed contrasts against

successful immediate recall and successful delayed recall across eight regions of interest (ROI). Only

immediate recall events from the first ten minutes of the first five sessions were included in these

contrasts; this time period is matched to the delayed recall that occurs in later sessions and only covers

immediate recall events where the subject is unaware of future delayed recall tests. We also performed

a contrast between immediate recall events from the first five sessions and deliberation periods from

the long-delayed recall test using the same matching procedure. We aggregated desired events from

each session within subjects and performed a t-test to produce a variance-normalized difference score

for every frequency-by-ROI pair within subjects. We subsequently used two-tailed one sample t-tests

(with FDR correction) to assess group-level differences.

170   **Results**

171   Figure 1A illustrates the basic structure of the experiment. In each of 10 experimental

sessions, subjects performed a simple immediate-recall task.   As each of 576 words appeared

individually, subjects read each word silently, and then, after a one second delay, recited the word

aloud. At the start of the 6th session (phase II), subjects were given a surprise long-delay recall task:

We asked them to recall as many of the 576 words as they could remember in any order as well as any

words that come to mind that may have not been presented in previous sessions. We then had subjects

perform the same immediate recall task as on previous sessions. Sessions 7-10 replicated the methods

of Session 6 except that the initial recall test could no longer be deemed a surprise.

179   As expected, subjects recalled items with very high accuracy (above 96%) in the immediate

recall condition. Figure 1B illustrates accuracy of immediate recall across the 24 test blocks separately

for phases one and two. Accuracy fell slightly across blocks but recovered after each of the two-minute

breaks (see Figure 1A), possibly due to build-up and release from proactive interference (e.g., Lohnas et

al., 2015; Underwood, 1957). Figure 1C illustrates accuracy of long-delay free recall. On the first session

of phase II, subjects recalled an average of 29 words (5% of the total pool of 576 words).   Their

performance on subsequent sessions increased dramatically, reaching an average recall rate of 103

items by the 10th session. This rate of increase (about 25 items per session) is much higher than the

implied rate of learning across sessions 1-5, but this likely reflects the difference between incidental and

intentional encoding.

Figure 1. **Experimental Paradigm and Behavioral Data. A.** During Sessions 1-5, subjects performed an immediate recall task for each of 576 words, silently reading a word which they verbally recalled after waiting approximately 1 second. On average, subjects responded 1.53 s after word onset. Each of sessions 6-10 began with subjects attempting to freely recall the 576 words that they had seen on each of the preceding sessions along with any other words that come to mind during a 10 minute retrieval interval. Subjects then performed the same immediate recall task as on earlier sessions. **B.** Subjects exhibited very high levels of immediate recall across all sessions, with performance dropping modestly across blocks and recovering following breaks. **C.** Subjects exhibited modest levels of delayed recall on the first (surprise) recall test given on Session 6, but performance rose sharply across subsequent sessions, hitting an average of 103 correct recalls by the final session. **D.** Intrusions similarly rose across sessions, but much less quickly than successful recalls. **E.** Inter-response times (IRTs) increased with output position; on sessions where subjects recalled a larger proportion of items, IRTs were generally faster throughout the retrieval period but rose sharply during the last few correct recalls.

Figure 1D illustrates the number of extra-list intrusions committed by subjects across the five sessions of phase II. Here we see that subjects committed a much larger percentage of intrusions than seen in standard within-session immediate and delayed recall tasks (Zaromb et al., 2006). However, the overall fraction of intrusions decreased over sessions, dropping from 37% on Session 6 to 23% on Session 10. Given the unusually-long delay in our free recall condition, and the very high rates of recall achieved by the 10th session, we asked whether inter-response times (IRTs) across the recall period exhibited the same pattern of growth as documented in more standard free-recall

214   paradigms (Rohrer & Wixted, 1994; Murdock & Okada, 1970). Figure 1E shows average IRTs based on

215   the total number of recalled words during delayed-recall sessions. This analysis demonstrates when

216   subjects recall many items, they exhibit much faster IRT overall. However, subjects still display a sharp

217   increase in their IRTs as they approach their final recalled item.

218   Our primary question concerned how neural activity, as measured through spectral analysis

of EEG recordings, signaled the process of spontaneous retrieval of previously experienced items. We

first addressed this question by comparing the 500 ms pre-vocalization period in the long-delay recall

condition with matched periods of silence separated by at least 0.5 sec from prior and subsequent

recalls (we refer to these as deliberation periods; see *Methods*). This comparison of retrieval and

deliberation periods revealed increased high-frequency activity and decreased alpha-band power across

most regions of interest (See Figure 2A; black outlined regions indicate statistically-significant

frequency-region pairs, FDR-corrected p<0.05), extending previous intracranial-recording studies that

identified similar retrieval biomarkers in both cued recall and free recall tasks (Burke, Ramayya, &

Kahana, 2015; Burke, Sharan, et al., 2014; Greenberg et al., 2015). Whereas those earlier studies

examined recall that took place within minutes of item encoding, the present study asked subjects to

recall items that had not been seen (experimentally) for at least 16 hours.

230   At anterior electrodes we observed increased theta-band activity during the pre-vocalization

memory-retrieval period. Although previous studies have frequently reported theta increases during

successful recognition memory (Addante, Watrous, Yonelinas, Ekstrom, & Ranganath, 2011; Guderian

& Düzel, 2005; Herweg et al., 2016; Osipova et al., 2006) most (standard) free recall studies find

retrieval-related *decreases* in theta and alpha-band power (Burke, Sharan, et al., 2014; Kragel, Koban,

Barrett, & Wager, 2018; Solomon et al., 2017; Solomon, Stein, et al., 2019). However, several free and

cued recall studies have found mixed results, with positive effects in specific brain regions, such as right

anterior temporal pole (Burke, Long, et al., 2014) and for specific contrasts, such as semantically-

clustered vs. non-clustered recall transitions (Solomon, Lega, Sperling, & Kahana, 2019). Our study

demonstrates retrieval-related increases in low theta power coupled with alpha-band decreases in a

long-delay recall task, conditions likely to place stronger demands on associative retrieval processes

(Herweg, Solomon, & Kahana, 2020).

Figure 2. **Statistical maps illustrating relative increases (red) and decreases (blue) in spectral power**

across key memory contrasts for eight regions of interest. A. Delayed Recall vs. Deliberation. **B.**

Delayed Recall vs. Immediate Recall. **C.** Deliberation vs. Immediate Recall. Panels **A** and **B** use matched

time periods from the first five sessions of Immediate Recall and the five sessions of Delayed Recall

following those first five sessions. Black-bordered regions indicated significant FDR-corrected t-tests on

within subject difference scores (p < .05). Electrodes locations corresponding to each region of interest

appear on a schematic view of an electrode net (LAI,RAI: left/right anterior inferior; LSA,RAS: left/right

anterior superior; LPI,RPI: left/right posterior inferior; LPS,RPS: left/right posterior superior).

Comparisons of pre-vocalization and deliberation intervals do not, however, uniquely identify

the process of memory retrieval. This is because prevocalization periods also differ from deliberation

in the presence of premotor activity related to the vocalization of recalled items. Switching to another

recall modality, e.g., typing, would simply replace this confound with a different one. We therefore

employed an immediate recall task as a control for motor activity that may confound our comparison

between pre-vocalization and deliberation intervals.

Using the pre-vocalization period in our immediate-recall task as a control, we found that

increased theta, decreased alpha, and increased HFA mark successful long-delay recall (Figure 2B; black

outlines indicate statistically-significant frequency-region pairs, that met an FDR-corrected p<0.05

threshold). This comparison recapitulates the spectral pattern observed in our comparison between

long-delay recall and deliberation, but without the confound of vocalization present in Figure 2A and

earlier studies. One difference, however, is that in our tighter contrast between immediate recall and

long-delay recall, the retrieval related theta effect appeared to be restricted to anterior-inferior ROIs.

Given the far greater demands on episodic memory retrieval in the long-delay condition, but the

matched vocalization in both conditions, we interpret these biomarkers as reflecting neural correlates

of context-dependent memory retrieval. Comparisons between immediate recall and deliberation

(Figure 2C) further support this interpretation. Here we see that trying to retrieve after a long delay (in

the deliberation periods) exhibits higher HFA and low alpha-band power than during the period

immediately preceding a motor response in the immediate recall condition (black outlines indicate

statistically-significant frequency-region pairs that met an FDR corrected *p*<0.05 threshold).

Figure 3. **Time course of high-frequency activity leading up to correct recalls.** Delayed recall, immediate

recall, and delayed recall baseline log high-frequency activity (98-110 Hz power) at posterior electrodes

in the 1.0 second leading up to vocalization of recalled items, averaged for all correctly recalled items

and across subjects. Results are shown separately for phases one and two. Error bands reflect 95%

confidence computed by the method of Loftus and Masson (1994).

281   Because high-frequency signals provide excellent temporal resolution, we sought to examine

the timing of the increased retrieval-related HFA seen prominently in posterior ROIs. Figure 3 shows

the time course of high-frequency activity (HFA) leading up to retrieval in the long delay and immediate

recall conditions. We illustrate this time course separately for immediate recalls contributed during

phases one and two. Error bands indicate the 95% confidence interval based upon between-subject

variability in high-frequency activity. For all three conditions, HFA rose in the moments leading up to

recall, but the prevocalization HFA was highest for delayed recall, lower for phase-two immediate recall,

and lowest for phase-one immediate recall. For comparison, we indicate the baseline gamma power in

the long delay deliberation periods. This ordering of conditions aligns with the hypothesized episodic

memory retrieval demands across these conditions (such demands being highest in delayed recall, and

lowest in immediate recall when a memory test is not expected). Given that the expectation of a

subsequent test would lead to better memory encoding during phase II, as shown in Figure 1C,

mnemonic processes likely exerted a greater influence on immediate recall in phase II than in phase

one.

296   **Discussion**

297   Humans possess a remarkable ability to search their memory for previously experienced items

learned in a given context.   When asked to recall without the aid of specific cues, subjects generate

their own retrieval cues, based upon the context at the time of test as well as the contextual

representations evoked by recently remembered items (Kahana, 2020).   Here we examined the

electrophysiological (EEG) correlates of spontaneous memory retrieval under conditions designed to

vary subjects' reliance on contextual retrieval between two extremes: In a long-delay recall condition,

we asked subjects to freely recall items not seen in at least 16 hours but encoded on previous sessions;

in an immediate recall condition, we asked subjects to read a word, pause for >1 seconds, and then

speak the word aloud. We chose these extreme contrasts to help distinguish pre-motor activity related

to vocalization from context-dependent memory-retrieval processes required in the long-delay task.

307   Contrasting immediate recall of a single word with long-delayed free recall of the entire 576-

word pool identified a spectral signature of successful episodic memory retrieval. Increased theta- and

gamma-band power (4-8 Hz and >40 Hz respectively), coupled with reduced alpha- and beta-band

power, marked periods of successful memory retrieval. This spectral signature appears to identify the

neural activity related to context-dependent memory retrieval. Our contrast of immediate recall and

long-delay recall allowed us to disentangle the premotor and electromyographic signals from episodic-

memory retrieval processes. These findings help to establish theta and gamma activity in the periods

preceding recall as a spectral signature of episodic memory retrieval.

315   Most prior EEG investigations of human memory have either compared encoding of

subsequently remembered and forgotten items (Staudigl & Hanslmayr, 2013; Klimesch, Doppelmayr,

Russegger, & Pachinger, 1996; Burke et al., 2015; Solomon et al., 2017; Hanslmayr, Spitzer, & Bäuml,

2008) or have compared successful and unsuccessful discrimination between targets and lures in a

recognition task (Jacobs, Hwang, Curran, & Kahana, 2006; van Vugt, Sekuler, Wilson, & Kahana, 2013;

Guderian & Düzel, 2005; Hsieh & Ranganath, 2014; Osipova et al., 2006). These, and other prior studies

reviewed by (Herweg et al., 2020) and (Hanslmayr & Staudigl, 2014), report both positive and negative

correlations between theta-band power and successful mnemonic processes. In the present study we

focused on the EEG correlates of successful free recall, a process rarely studied using non-invasive

recording methods due to the presence of large electromyographic (EMG) artifacts caused by

vocalization during recall. By contrasting long-delayed recall with immediate recall, we find that

increased theta and decreased alpha power mark periods of successful episodic memory retrieval.

Pastötter and Bäumel (2014) report a similar theta-alpha pattern in a cued recall task.

328   Our finding that memory-related theta increases appeared at the lower bound of the theta

range aligns with several earlier reports using intracranial recordings to examine memory encoding

(Miller et al., 2018; Lega, Jacobs, & Kahana, 2012; Lin et al., 2017) as well as scalp EEG and MEG studies

of successful recognition memory (Herweg et al., 2016; Gruber, Tsivilis, Giabbiconi, & Müller, 2008).

Studies reporting negative correlations between theta and successful memory encoding or retrieval

either did not use contrasts that selected for the associative or contextual aspects of memory retrieval,

or they averaged across a broader frequency range that would include both low-frequency increases

and alpha-band decreases (see Herweg et al., 2020, for a review).

336   Simultaneous recordings of local field potentials and single neuron activity have implicated

high-frequency activity (HFA) as a correlate of neuronal firing rates (Manning, Jacobs, Fried, & Kahana,

2009). Studying neurosurgical patients with indwelling electrodes, Burke et al (2015), Solomon et al

(2017), and Long et al. (2018) have all found greater HFA immediately preceding recall than during

matched deliberation intervals. Although invasive recordings minimize the influence of premotor

artifacts, it is still likely that some non-mnemonic factors contributed to those reported results. Our

finding that high-frequency activity increases in the moments leading up to successful recall when

comparing long-delay with immediate recall (Figure 2B) directly implicates HFA increases in the

cognitive processes involved in episodic memory retrieval.

345   The present study identifies a pattern of electrophysiological biomarkers of successful

episodic recall: increases in frontal slow-theta (+T), and decreases in alpha-band power (-A) and increased

high-frequency, or gamma, activity with a posterior distribution (+G). These biomarkers, tag (+T-A+G) the

process of retrieval from episodic memory. They also add to an emerging body of evidence demonstrating

the utility of non-invasive methods for decoding cognitive states (Chakravarty, Chen, & Caplan, 2020; Noh,

Liao, Mollison, Curran, & de Sa, 2018; Weidemann & Kahana, 2019, 2020). Given the ease of collecting

non-invasive EEG data from human research subjects, and human's potentially unique ability to

spontaneously recall verbal items, the +T-A+G biomarker of episodic recall can serve as a basis for future

studies that investigate the role of retrieval cues, such as the temporal, semantic and spatial contexts

surrounding experienced items.

356   References

357   Addante, R. J., Watrous, A. J., Yonelinas, A. P., Ekstrom, A. D., & Ranganath, C. (2011). Prestimulus theta
358   activity predicts correct source memory retrieval. *Proceedings of the National Academy of*
359   *Sciences, USA*, *108*(26), 10702-10707.

360   Aka, A., Phan, T., & Kahana, M. J. (2020). Predicting recall of words and lists. *Journal of Experimental*
361   *Psychology: Learning, Memory, and Cognition*.

362   Burke, J. F., Long, N. M., Zaghloul, K. A., Sharan, A. D., Sperling, M. R., & Kahana, M. J. (2014). Human
363   intracranial high-frequency activity maps episodic memory formation in space and time.
364   *NeuroImage*, *85* , 834–843. doi: 10.1016/j.neuroimage.2013.06.067

365   Burke, J. F., Ramayya, A. G., & Kahana, M. J. (2015). Human intracranial high-frequency activity during
366   memory processing: Neural oscillations or stochastic volatility? *Current Opinion in Neurobiology*,
367   *31*, 104–110.

368   Burke, J. F., Sharan, A. D., Sperling, M. R., Ramayya, A. G., Evans, J. J., Healey, M. K., . . . Kahana, M. J.
369   (2014). Theta and high–frequency activity mark spontaneous recall of episodic memories.
370   *Journal of Neuroscience*, *34*(34), 11355–11365. doi: 10.1523/JNEUROSCI.2654-13.2014

371   Chakravarty, S., Chen, Y., & Caplan, J. (2020). Predicting memory from study-related brain activity.
372   *Journal of Neurophysiology*, *124*, 2060-2075.

373   DelPozo-Banos, M., & Weidemann, C. T. (2017). Localized component filtering for electroencephalogram
374   artifact rejection. *Psychophysiology*, *54*(4), 608-619.

375   Gramfort, A., Luessi, M., Larson, E., Engemann, D. A., Strohmeier, D., Brodbeck, C., . . . Hämäläinen, M.
376   (2013). MEG and EEG data analysis with MNE-Python. *Frontiers in Neuroscience*, *7*(267). doi:
377   10.3389/fnins.2013.00267

378   Greenberg, J. A., Burke, J. F., Haque, R., Kahana, M. J., & Zaghloul, K. A. (2015, July). Decreases in theta
379   and increases in high frequency activity underlie associative memory encoding. *NeuroImage*,
380   *114*, 257–263. doi: 10.1016/j.neuroimage.2015.03.077

381   Gruber, T., Tsivilis, D., Giabbiconi, C.-M., & Müller, M. (2008). Induced electroencephalo gram oscillations
382   during source memory: familiarity is reflected in the gamma band, recollection in the theta band.
383   *Journal of Cognitive Neuroscience*, *20*(6), 1043-53.

384   Guderian, S., & Düzel, E. (2005). Induced theta oscillations mediate large-scale synchrony with
385   mediotemporal areas during recollection in humans. *Hippocampus*, *15*(7), 901–912.

386   Hanslmayr, S., Spitzer, B., & Bäuml, K.-H. T. (2008). Brain oscillations dissociate between

387   semantic and nonsemantic encoding of episodic memories. *Cerebral Cortex*, *19*(7),

388   1631-1640. doi: 10.1093/cercor/bhn197.

389   Herweg, N. A., Apitz, T., Leicht, G., Mulert, C., Fuentemilla, L., & Bunzech, N. (2016). Theta-alpha

390   oscillations bind the hippocampus, prefrontal cortex, and striatum during recollection: Evidence

391   from simultaneous EEG–fMRI. *Journal of Neuroscience*, *36* (12), 3579–3587.

392   Herweg, N. A., Solomon, E., & Kahana, M. J. (2020). Theta oscillations in human memory. *Trends in

393   Cognitive Science*, *24*(3), 208-227.

394   Hsieh, L.-T., & Ranganath, C. (2014). Frontal midline theta oscillations during working memory

395   maintenance and episodic encoding and retrieval. *NeuroImage*, *85*, 721-729.

396   Jacobs, J., Hwang, G., Curran, T., & Kahana, M. J. (2006). EEG oscillations and recognition memory: Theta

397   correlates of memory retrieval and decision making. *NeuroImage*, *15*(2), 978–87.

398   Kahana, M. J. (2020, January). Computational models of memory search. *Annual Review of Psychology,

399   71* (1), 107–138. Retrieved from https://doi.org/10.1146/annurev-psych-010418-103358 doi:

400   10.1146/annurev- psych-010418-103358

401   Kahana, M. J., Aggarwal, E. V., & Phan, T. D. (2018). The variability puzzle in human memory. *Journal of

402   Experimental Psychology: Learning, Memory, and Cognition*, *44*(12), 1857–1863. doi:

403   10.1037/xlm0000553

404   Kahana, M. J., Dolan, E. D., Sauder, C. L., & Wingfield, A. (2005). Intrusions in episodic recall: Age

405   differences in editing of overt responses. *Journal of Gerontology: Psychological Sciences*, *60*(2),

406   92–97. doi: 10.1093/geronb/60.2.P92

407   Klimesch, W., Doppelmayr, M., Russegger, H., & Pachinger, T. (1996). Theta band power in the human

408   scalp EEG and the encoding of new information. *NeuroReport*, *7*, 1235–1240.

409   Kragel, P. A., Koban, L., Barrett, L. F., & Wager, T. D. (2018). Representation, pattern information, and

410   brain signatures: From neurons to neuroimaging. *Neuron*, *99*(2), 257–273.

411   Lega, B., Jacobs, J., & Kahana, M. (2012). Human hippocampal theta oscillations and the formation of

412   episodic memories. *Hippocampus*, *22*(4), 748–761.

413   Lin, J. J., Rugg, M. D., Das, S., Stein, J., Rizzuto, D. S., Kahana, M. J., & Lega, B. C. (2017). Theta band power

414   increases in the posterior hippocampus predict successful episodic memory encoding in humans.

415   *Hippocampus*, *27*(10), 1040-1053. doi: 10.1002/hipo.22751

416   Loftus, G. R., & Masson, M. E. J. (1994). Using confidence intervals in within-subject designs. *Psychonomic*
417   *Bulletin & Review*, *1*(4), 476–490. doi: 10.3758/BF03210951

418   Lohnas, L. J., Polyn, S. M., & Kahana, M. (2015). Expanding the scope of memory search: Modeling
419   intralist and interlist effects in free recall. *Psychological Review*, *122*(2), 337–363. doi:
420   10.1037/a0039036

421   Long, N. M., Sperling, M. R., Worrell, G. A., Davis, K. A., Gross, R. E., Lega, B. C., . . . Kahana, M. J. (2017).
422   Contextually mediated spontaneous retrieval is specific to the hippocampus. *Current Biology*,
423   *27*(7), 1074–1079. doi: 10.1016/j.cub.2017.02.054

424   Manning, J. R., Jacobs, J., Fried, I., & Kahana, M. J. (2009). Broadband shifts in local field potential power
425   spectra are correlated with single-neuron spiking in humans. *Journal of Neuroscience*, *29*(43),
426   13613–13620. doi: 10.1523/JNEUROSCI.2041-09.2009

427   Miller, J. F., Watrous, A. J., Tsitsiklis, M., Lee, S. A., Sheth, S. A., Schevon, C. A., . . . Jacobs, J. (2018).
428   Lateralized hippocampal oscillations underlie distinct aspects of human spatial memory and
429   navigation. *Nature communications*, *9*(1), 2423.

430   Murdock, B. B., & Okada, R. (1970). Interresponse times in single-trial free recall. *Journal of Verbal*
431   *Learning and Verbal Behavior*, *86*, 263-267.

432   Long, N. M., Kuhl, B. A., & Chun, M. M. (2018). Memory and attention. In *Stevens' handbook of*
433   *experimental psychology and cognitive neuroscience*, Fourth Edition, Wixted, J. T. (pp. 1-37). doi:
434   10.1002/9781119170174.epcn109

435   Noh, E., Liao, K., Mollison, M., Curran, T., & de Sa, V. R. (2018). Single-trial eeg analysis predicts memory
436   retrieval and reveals source-dependent differences. *Frontiers in Human Neuroscience*, *12*, 258.

437   Osipova, D., Takashima, A., Oostenveld, R., Fernández, G., Maris, E., & Jensen, O. (2006). Theta and
438   gamma oscillations predict encoding and retrieval of declarative memory. *Journal of*
439   *Neuroscience*, *26*(28), 7523–7531. doi: 10.1523/JNEUROSCI.1948- 06.2006

440   Pastötter, B., & Bäumel, K. H. T. (2014). Distinct slow and fast cortical theta dynamics in episodic memory
441   retrieval. *Neuroimage*, *94*, 155-161.

442   Rohrer, D., & Wixted, J. T. (1994). An analysis of latency and interresponse time in free recall. *Memory &*
443   *Cognition*, *22*, 511- 524.

Solomon, E. A., Kragel, J., Sperling, M., Sharan, A., Worrell, G., Kucewicz, M., . . . Kahana, M. (2017).

Widespread theta synchrony and high-frequency desynchronization underlies enhanced

cognition. *Nature Communications*, *8*(1), 1704. doi: 10.1038/s41467- 017-01763-2

Solomon, E. A., Lega, B. C., Sperling, M. R., & Kahana, M. J. (2019). Hippocampal theta codes for distances

in semantic and temporal spaces. *Proceedings of the National Academy of Sciences*, *116*(48),

24343-24352.

Solomon, E. A., Stein, J. M., Das, S., Gorniak, R., Sperling, M. R., Worrell, G., . . . Kahana, M. J. (2019).

Dynamic theta networks within the human medial temporal lobe support episodic encoding and

retrieval. *Current Biology*, *29*(7), 1100-1111. doi: 10.1016/j.cub.2019.02.020

Staudigl, T., & Hanslmayr, S. (2013). Theta oscillations at encoding mediate the context-dependent

nature of human episodic memory. *Current Biology*, *23*(12), 1101-1106.

Underwood, B. J. (1957). Interference and forgetting. *Psychological Review*, *64*, 49-60. doi:

10.1037/h0044616

van Vugt, M. K., Sekuler, R., Wilson, H. R., & Kahana, M. J. (2013). An electrophysiological signature of

summed similarity in visual working memory. *Journal of Experimental Psychology: General*,

142(2), 412–425.

Weidemann, C. T., & Kahana, M. J. (2019). Dynamics of brain activity reveal a unitary recognition signal.

Journal of Experimental Psychology: Learning, Memory, and Cognition, *45*(3), 440–451. doi:

http://dx.doi.org/10.1037/xlm0000593

Weidemann, C. T., & Kahana, M. J. (2020). Neural measures of subsequent memory reflect endogenous

variability in cognitive function. *Journal of Experimental Psychology: Learning, Memory, and*

Cognition.

Zaromb, F. M., Howard, M. W., Dolan, E. D., Sirotin, Y. B., Tully, M., Wingfield, A., & Kahana, M. J. (2006).

Temporal associations and prior-list intrusions in free recall. *Journal of Experimental Psychology:*

Learning, Memory, and Cognition, *32*(4), 792– 804. doi: 10.1037/0278-7393.32.4.792