TITLE PAGE

Title: Modularity and connectivity of nest structure scale with colony size

Running head: Nest architecture scales with colony size

Authors: Julie S. Miller[1], Emma Wan[1], Sean O'Fallon[1] & Noa Pinter-Wollman[1]

Author Affiliations
1. Department of Ecology and Evolutionary Biology, University of California Los Angeles,
612 Charles E. Young Dr. East, Los Angeles, CA 90095-7246.

Corresponding Author: julie.serena@gmail.com; +1-310-766-3581; 612 Charles E. Young Dr.
East, Los Angeles, CA 90095-7246.

Author contributions: JSM conceived of the study, analyzed the data and wrote the first draft of
the paper. JSM and EW collected the data. SO collected data on ant body lengths. JSM and NPW
designed the study, and all authors helped write the manuscript.

Acknowledgements: We are thankful to Doug Booher for sharing his nest castings and to
Walter Tschinkel for additional images of his published castings. We thank Dana Williams,
Felipe Zapata, Kevin Loope, Tyler McCraney and the UCLA IDRE statistical consulting unit for
advice on the statistical analysis and to Saray Shai for advice on generating reference models.
We thank Gabby Najm, James Lichtenstein, Natalie Lemanski, Andrea Perna, and two
anonymous reviewers for helpful comments on the manuscript. The authors declare no conflicts
of interest.

Funding: JSM was supported by NSF Postdoctoral Research Fellowship in Biology DBI
1812292 and NSF grants IOS 1456010/1708455 to NPW.

Data Accessibility Statement: Data available on Dryad, DOI https://doi.org/10.5068/D18D7B;
Code available on Zenodo, DOI https://doi.org/10.5281/zenodo.5636719

TITLE: Modularity and connectivity of nest structure scale with colony size

ABSTRACT

Large body sizes have evolved structures to facilitate resource transport. Like unitary organisms,

social insect colonies must transport information and resources. Colonies with more individuals

may experience transport challenges similar to large-bodied organisms. In ant colonies, transport

occurs in the nest, which may consist of structures that facilitate movement. We examine three

attributes of nests that might have evolved to mitigate transport challenges related to colony size:

(1) subdivision - nests of species with large colonies are more subdivided to reduce crowd

viscosity; (2) branching - nest tunnels increase branching in species with large colonies to reduce

travel distances; and (3) short-cuts – nests of species with large colonies have cross-linking

tunnels to connect distant parts of the nest and create alternative routes. We test these hypotheses

by comparing nest structures of species with different colony sizes in phylogenetically controlled

meta-analyses. Our findings support the hypothesis that nest subdivision and branching evolved

to mitigate transport challenges related to colony size. Nests of species with large colonies

contain more chambers and branching tunnels. The similarity in how ant nests and bodies of

unitary organisms have evolved in response to increasing size suggests common solutions across

taxa and levels of biological organization.

KEYWORDS: nest, morphological evolution, network, colony size, ants, allometry

INTRODUCTION

The consequences of body size for the evolution of phenotypic diversity are far reaching,

influencing morphological (Schmidt-Nielson 1984), physiological (Kleiber 1947), and

behavioral traits (Calder 1984; Dial et al. 2008) in systematic ways (Peters 1983). Evolutionary

allometry (Thompson 1917; Huxley 1932), the study of how body size relates to trait size across

lineages (Gould 1971; Cheverud 1982), has offered a window into the underlying mechanisms

that drive trait diversity (e.g. McMahon 1975; Fairbairn 1997; Frankino et al. 2005). An

important consequence of increasing body size is that it alters the basic physical conditions in an

organism by increasing the distances over which transport of nutrients, waste products, and

signals must traverse (Bonner 2004). This physical transport challenge has been resolved through

the evolution of a variety of structural features across the tree of life (McMahon and Bonner

1983), including multi-cellularity (Niklas and Newman 2016), body plans that increase surface-

area-to-volume ratios (e.g., jellyfish and sponges; Rupppert et al. 2004), branching vascular

systems in animals and plants (West et al. 1997; Banavar et al. 1999), and compartmentalization

of specialized tissues into organs (Gould 1966).

Social insect colonies are often compared to whole organisms (Wheeler 1910; Seeley

1989; Holldobler and Wilson 2009), and large colonies may experience transport challenges that

are similar to those of large bodied organisms (Gillooly et al. 2010). In larger colonies, the speed

and/or reliability with which resources, or information, reach disparate parts of the colony will

depend on transfer occurring among more individuals and across greater distances (Pacala et al.

1996; O'Donnell and Bulova 2007; Dornhaus et al. 2012). The outstanding diversity of the ant

lineage (Hölldobler and Wilson 1990), and multiple independent evolutionary increases in

colony size (Burchill and Moreau 2016), provide an ideal foundation for comparative work that

investigates how physical structures evolved to mitigate transport challenges related to size.

Transport of resources and information in social insect colonies may depend on the

layout of their nests. Nest structure affects the routes that individuals take as they move resources

and information. Therefore, nest structure influences the way in which individuals interact

(Pinter-Wollman et al. 2013; Pinter-Wollman 2015b; Tschinkel 2015; Pinter-wollman et al.

2018; Vaes et al. 2020) and behave as a collective (Burd et al. 2010; Pinter-Wollman 2015a).

Having more individuals may slow the delivery of resources and information, but the physical

structure of the nest can promote rapid movement. Here we investigate macroevolutionary trends

in how key nest traits, including subdivision, branching, and short-cuts, relate to typical colony

size, i.e., the average number of individuals in a colony, in ants (Formicidae).

Subdivision, or compartmentalization, is one structural feature that large systems evolved

to mitigate the challenges of size. For example, large multicellular organisms consist of

subdivided cells and organs (Kempes et al. 2012; Okie 2013). However, the size of the units are

constrained by the costs of expanding surface area-to-volume ratios (Bonner 1988; Lorenz et al.

2011). In social insect nests, larger nest chambers can hold more individuals, but make it harder

for individuals to contact each other (Brian 1956; Aguilar et al. 2018). Constraining the size of

nest chambers reduces the number of individuals occupying each chamber while reducing the

distances they need to travel to interact. The "subdivision hypothesis" predicts that the nests of

species with larger colony sizes will consist of more chambers compared to nests of species with

small colonies, while chamber sizes will not differ in relation to colony size.

Another way that large systems have evolved to improve transport is by increasing their

connectivity and in turn, reducing travel distances. Travel distances can be reduced by adding

paths that branch off from main routes. Many natural systems exhibit branching structures which

reduce travel distances without the addition of cross-linking paths. For example, hierarchical

branching structures are common in the vascular networks of animals (West et al. 1997; Tekin et

al. 2016) and plants (McKown et al. 2010; Blonder et al. 2018). Branching structures are

common when resources enter through a single site and need to be homogenously distributed

throughout the system (Dodds 2010). The "branching hypothesis" predicts that nests of species

with larger colonies will have tunnels with high branching frequency to reduce travel distances in

the nest.

Reducing travel distances and increasing connectivity can further be accomplished by

adding cross-linking paths, or short-cuts, that connect distant parts of a network and create

alternative access routes to maintain traffic flow (Corson 2010; brains: Bullmore and Sporns

2012; plants: Katifori et al. 2010; Blonder et al. 2018; termite nests: Valverde et al. 2009). The

"short-cut hypothesis" predicts that nests of species with larger colonies will contain more cross-

linking paths than nests of species with smaller colonies. Despite the transport benefits of adding

short-cuts, every additional connection has a building and maintenance cost, and highly

connected nests may be structurally unsound (Monaenkova et al. 2015).

Ants offer an ideal system to study how nest structure co-evolves with group size on

macroevolutionary scales because ants span a large range of colony sizes (Bourke and Franks

1995), and exhibit tremendous structural diversity in their nests (Tschinkel 2015). Early lineages

of ant species have small to medium sized colonies (Burchill and Moreau 2016) and build small

nests with a single tunnel and/or chambers (Tschinkel 2003), similar to their Apoid wasp

ancestors (Branstetter et al. 2017). The evolutionary radiation of the ants resulted in a diversity

of nest structures and forms, but their diversity has not been quantified and related to patterns of

colony size evolution. Furthermore, it is unknown how the transport challenges introduced by

large colony sizes have been mitigated in the evolutionary history of ants. Here, we focus on

ground-nesting species because, out of all ants, they have the greatest potential to control the

structure of their nests.

Prior studies of subterranean ant nest structure in relation to colony size have focused on

the ontogeny of single species, showing that ant nests grow in volume and increase chamber size

and/or number as colonies age (Tschinkel 1987, 2004, 2005, 2011, 2015). Fewer studies have

examined how tunnel connectivity changes with colony development (Buhl et al. 2004a), so

branching patterns in ant nests are not well understood. Furthermore, these studies on the

ontogeny of nest structure do not provide information on the evolution of nest structures in

relation to mature colony sizes, which we examine here in a phylogenetically controlled cross-

species comparison.

Here we investigate interspecific variation in ant nest structure and colony size within a

phylogenetic context. We consider an adaptive hypothesis for the evolution of nest structure

diversity: if ant colonies face size-associated challenges in transport and communication,

analogous to other systems, then the structure of nests should relate to colony size in ways that

ease the movement of individuals, resources, and information throughout the nest. We quantify

three attributes of nest structure, (1) subdivision, (2) branching, and (3) short-cuts, and examine

whether these traits relate to colony size in phylogenetically controlled, cross-species meta-

analyses. We further consider that variation in nest structure could be the result of non-adaptive

processes, like genetic drift or constraints imposed by evolutionary history. To determine the

extent to which phylogenetic history predicts variation in nest structure, we measure the

phylogenetic signal for the above nest traits and colony size.

METHODS

Obtaining nest and colony size data

To test our hypotheses, we gathered data on nest architecture of subterranean ant species. We identified publications with graphical representations of ant nests by searching for the terms "nest", "structure", "architecture", "casting", and "excavation/ed" on Google Scholar and Web of Science. We identified 24 papers that contained photographs or illustrations of castings or excavations of complete ant nests with distinguishable entrances, tunnels, and chambers. For the subdivision hypothesis, we included data from 3 additional papers which contain tables with measures of chamber size and number, even though nest images were not available (Table S1). We excluded 14 publications from our dataset because interior chambers and/or tunnels were obscured, nests were constructed under artificial conditions, or were from incipient colonies (Table S2). Authors were contacted for additional images when parts of the nest were obscured, accounting for approximately 10% of the images we used. Physical castings of unpublished nests were also obtained (Table S1). When more than one nest was available for a species, we took an average of all measurable nests.

To relate nest structure with colony size – i.e., the number of ants in a colony - we searched the scientific literature and the web (AntWiki.org and AntWeb.org) for information on the number of workers in a mature colony. For the majority of species, we used the average colony size from the literature. When only a range (minimum - maximum) for number of ants was available, we took the midpoint. Information about the number of ants in a colony at the time of nest excavation or casting was available in less than half of the studies we obtained nest structures from, but when available, we used the reported number of adult workers for those particular nests. When colony size was not available for a particular species, or when nest

identity at the species level was unknown, we used colony sizes representative of the order of

magnitude for the genus obtained from a sister species. This approximation was necessary for the

Forelius sp nest cast in Anza Borrego, CA (Booher, personal communication). At this site, only

two species of *Forelius* occur and they have the same reported colony size, so we used the

colony size from one of them - *F. pruinosus* (10^4). For *Dorymyrmex bureni* we used the only

species from this genus with a recorded colony size - *Dorymyrmex bicornis* (10^3) (Table S1).

We excluded *Pheidole oxyops* and *Acromyrmex subterraneus* nests from our analysis because

colony size was not available for these species, and because we could not make a reliable

approximation due to substantial variation in recorded colony sizes (Table S2).

Nests in our sample come from species belonging to 6 subfamilies, including from the

most diverse subfamilies (Myrmecinae, Formicinae, and Dolichoderinae), and the early

branching Ponerinae, which tend be characterized by smaller colony sizes and more basal traits

in general. However, we lacked nest data from other early lineages, like the Leptanillinae and the

Amblyoponinae.

Nest Subdivision Hypothesis

To test the subdivision hypothesis, we related colony size to the number and size of nest

chambers. We counted chambers in 296 nests from 43 species in 24 genera and measured their

sizes in 188 nests from 37 species in 21 genera. To quantify the size of chambers, we measured

the maximum width of each chamber using ImageJ (Rueden et al. 2017). We excluded nests

from the size analysis if images did not include a scale bar or if species identity was unknown for

a nest (e.g., *Forelius sp.*). We used chamber width as a proxy for chamber area because we relied

on 2D images of 3D structures. When we had access to physical castings, we measured chamber

width directly from the structures using a measuring-tape. We validated that chamber width was

a reliable indicator of chamber area using an independent dataset (see *Chamber Width Validation*

section of the Supplemental Materials).

To allow for cross-species comparison, we standardized chamber width to worker size.

Worker size was measured using a standard metric of ant body size, Weber's length (Brown

1953), using ant profile images containing a scale bar from AntWeb.org or other publications.

For all species, we measured the available images up to 8 specimens (1-8). For monomorphic

species and for species with continuous worker polymorphism, we took the average length from

all available images. For species with discrete worker polymorphism (i.e., majors and minors),

we used a weighted average, based on typical frequency of majors according to the literature

(Brown and Traniello 1998; Murdock and Tschinkel 2015). Standardized widths of all chambers

in each nest were averaged to obtain a single standardized chamber width per nest.

Nest Connectivity

Nests as Networks

To examine nest connectivity, we depicted each nest as a network, as in (Buhl et al. 2004b; Perna

et al. 2008; Viana et al. 2013; Pinter-Wollman 2015a; Kwapich et al. 2018), by manually

assigning a node ID to each chamber, nest entrance, tunnel junction, and ends of tunnels that did

not reach a chamber ('end nodes') and connecting these nodes with edges depicting tunnels. See

Figure S1 for an illustration of this method. When quantifying nest casts as networks we

excluded the "mushroom top" where the casting material pools around the nest entrance when it

is poured (Tschinkel 2010). We designated the nest entrance as the narrowest section of the

mushroom top. We obtained networks of 170 nests of 38 species from 21 genera, excluding

images in which the network structure could not be inferred because tunnel connections were

obscured or ambiguous (Table S1 & 2). We excluded polydomous nests because it was not

possible to calculate the network measures we used, mean travel distances and number cycles

(see below), without information about the structure of trails connecting different nests.

Branching Hypothesis

To test the if branching reduces travel distances in large colonies, i.e., the 'branching hypothesis'

we quantified the length of the travel paths in a nest as the mean distance of a network. Mean

distance of a network is the sum of the lengths of the shortest paths that connect all possible pairs

of nodes, divided by the number of all possible node pairs (Figure 1):

$$Mean\ Distance = \frac{1}{n(n-1)} \sum_{i=1}^{n} d_{ij}$$

where n is the number of nodes, and d is the shortest path between all node pairs i and j

(Newman 2018). We used the "mean_distance()" function in igraph R package (Csárdi 2020).

To determine if observed nests had more branching than expected based on their size, i.e.,

the number of chambers, or nodes, in the network, we compared the mean distance of networks

from observed nests to mean distance of reference models of networks with known branching

properties. We evaluated how mean distance relates to nest size under different assumptions

about how new nodes are added to a network (i.e., generative models (Hobson et al. 2021)). We

used three generative reference models that represent the upper and lower bounds on

connectivity for how nest networks might increase in size (Buhl et al. 2004b; Bebber et al. 2007;

Tero et al. 2010; Latty et al. 2011): (1) chain networks (Figure 1A), in which new nodes are

added to the last node in a chain – i.e., no branching; (2) triangulated networks (Figure 1C), in

which new nodes are connected to at least two other nodes such that they form a triangle while

avoiding edge overlap. We generated random triangulated networks using Delaunay

triangulation with the deldir() function in the 'deldir' R package (Turner 2020). The x,y

coordinates of the nodes to connect were randomly generated using the 'sample()' function; and

(3) minimum spanning trees (MSTs; Figure 1B), in which each new node is added to a terminal

node to form a new branch in a hierarchical structure. New nodes are added such that the total

length of the network is minimized and no cycles (Figure 1C) are formed. We generated MST

networks by pruning the triangulated graphs generated in (2) using the 'mst' function in the

'igraph' R package (Csárdi 2020). Each of the above generative processes was run for network

sizes ranging from 3 to 100 nodes, in increments of one node. We repeated this process 1000

times for each network size for both triangulated networks and MSTs, because they were

probabilistic generative processes, but not for the chain network generation because each size has

only one solution. For each generated network, we calculated the mean distance to create

reference distributions for comparison with the observed data (Hobson et al. 2021; Figure 3B).

See R code in the supplementary materials.

We further scaled the observed networks to the reference models to relate mean distance

to colony size (number of individuals) in a way that accounted for expected changes in network

size (number of nodes). To create scaled measures of mean distance, we divided the observed

mean distance by the mean value of mean distance of the reference networks (from the 1000

simulations) for each network size (3-100 nodes). The closer the scaled mean distance value is to

1, the more similar the observed network is to the reference model. Scaled values greater than 1

indicate that the observed network has a higher mean distance than expected according to the

reference model. Likewise, scaled values less than 1 indicate a smaller mean distance than

expected according to the reference model. We related these scaled values with colony size using

phylogenetically controlled comparisons as described below in Statistical Analyses (Figure 3C).

Short-Cut Hypothesis

To test if larger colonies have nests with more short-cuts, i.e., the short-cut hypothesis, we

quantified short cuts using the number of cycles in a network (Buhl et al. 2004b; Pinter-Wollman

2015a). A cycle is a path along a network that returns to its starting node after passing through at

least one other node (Newman 2018; Figure 1C). The number of cycles in a nest is:

$$Number\ of\ cycles = \ m - n + 1$$

where *n* is the number of nodes and *m* is the number of edges. Cycles are a good approximation

of path redundancy of a network, or presence of short-cuts, indicating how often there is more

than one route to any given part of the nest (Perna and Latty 2014).

To account for how network size affects the rate at which new cycles are added, we

compared the observed nests to the reference models of networks described above. Following a

similar procedure to the one described for network mean distance, we calculated the number of

cycles in triangulated networks (Figure 3E) and then scaled the number of cycles in observed

nests to triangulated networks of equivalent size. Because both chain and MST reference models

do not produce cycles, we did not include these models when we scaled the observed nests. We

related the scaled number of cycles in a nest with colony size using phylogenetically controlled

comparisons as described below (Figure 3F).

Figure 1: Examples of nest topologies with 10 chambers connected based on the three

generative reference models we tested: (A) Chain network, (B) Minimum Spanning Tree, and

(C) Fully connected Triangulated network. Above each network we provide its mean distance

(MD), which is the average of shortest paths that connect all possible pairs of nodes. As an

example, in (C), the length of the shortest path between nodes A and B is one, indicated by the

light blue edge X. The length of shortest path between nodes A and C is two, indicated by the

light blue edges X and Y. We further provide the number of cycles in each network. As an

example, in (C) the purple edges are a cycle connecting nodes D, E, and F.

Statistical Analyses

We examined the effect of colony size on the above nest measures while controlling for

phylogeny. We used a pruned evolutionary tree of the ants (Formicidae) from (Blanchard and

Moreau 2017) and calculated the mean trait value for each species. A complete species-level tree

was not available for all species in our dataset, so we selected only one species per genus and ran

our analysis using a genus-level tree (Figure 4). When there were data for more than one species

per genus, we used the species which had the most nests in our dataset (subset 1 in Tables S3-

S13). To ensure this sub-setting of species did not bias our results, we re-ran each analysis with

previously excluded species as representatives of a genus, and report the proportion of those runs

that were statistically significant. See supplementary materials for results from all runs (Tables

S3-13).

We ran phylogenetic generalized least squares (PGLS) comparing each nest trait to the

log of colony size. PGLS is a multiple regression that controls for the degree of similarity

between species based on phylogenetic distance in a co-variance matrix. We ran PGLS by

estimating the phylogenetic signal, lambda λ, through maximum likelihood in the "caper"

package v1.0.1 in R (Orme et al. 2018), with $\lambda=0$ meaning no phylogenetic signal, and $\lambda=1$

meaning the observations completely match phylogeny (Pagel 1999). Due to the small sample

size (20 - 24 genera) we could not estimate lambda with high confidence (Freckleton 2009).

Therefore, we further conducted PGLS for fixed lambda values (0, 1) using the "ape" package

v5.1 in R (Paradis et al. 2018). When our models did not meet the assumption of normality of

residuals we analyzed these relationships by calculating phylogenetic independent contrasts

(PICs) instead (Paradis et al. 2018). PIC transformed data were modeled using OLS regression,

with the intercept set to the origin (R Core Team 2019). Due to small sample sizes, we calculated

95% confidence intervals for OLS model coefficients using a bootstrap (1000 iterations).

To estimate phylogenetic signal for all our measures of nest structure and the log of

colony size, we estimated Pagel's lambda with the 'phylosig' function in the "phytools" R

package (Revell 2017). This function estimates the value of Pagel's lambda from 1000

simulations and compares it to a model of no phylogenetic signal (λ=0) using a likelihood ratio

test. Due to small sample size, we also compared the fit of our estimated lambda model of

phylogenetic signal to a Brownian Motion model (λ=1) and to a model with no phylogenetic

signal (λ=0) using AICc values calculated from the 'fitContinuous' function in the "geiger"

package in R (Harmon et al. 2020). Finally, to visualize colony size and nest trait variation

across the phylogeny, we mapped standardized trait values onto a genus level phylogeny from

species subset 1 (Figure 4) using the 'contMap' and 'phylo.heatmap' functions in the "phytools"

R package (Revell 2017).

RESULTS

Subdivision Hypothesis

Both predictions of the subdivision hypothesis, that chamber number would be larger in species

with large colony sizes and that chamber size would not relate to colony size, were supported.

We found a statistically significant positive relationship between number of chambers per nest

and colony size (Figure 2A; N= 24 genera, OLS of PIC: β = 4.024, 95% CI = 2.014, 6.292, R^2 =

0.560, $p < 0.0001$; Table S3). The effect size of this relationship was large ($3.7 < \beta < 4.3$ across

all data subsets) and statistically significant in all species subsets (19/19 datasets). We did not

detect a statistically significant relationship between the average standardized chamber width and

colony size (Figure 2B; N=21 genera, OLS of PICs: β = 0.350, 95% CI = -0.049, 0.959, R^2 =

0.140, p = 0.095; Table S4) in the majority of the species subsets (13/15 datasets).

Figure 2: Relationship between the log of colony size (number of ants in a colony) and (A)

average number of chambers per nest or (B) average standardized chamber width. Each data

point represents a mean value for a single species, whiskers indicate standard errors when

multiple nests were sampled from the species. Colors represent ant subfamilies. Lines are fits

from the PGLS (λ=1) with a statistically significant fit as a solid line and a relationship that is not

statistically significant as a dashed line.

Branching Hypothesis

The "branching hypothesis", that structural branching would reduce travel distances in nests of

species with large colonies, was supported. Raw, unscaled, mean distance of nests significantly

increased with colony size (Figure 3A; N = 21 genera, PGLS λ=MaxLL: β =1.36 ± 0.247,

λ=0.585, R^2 = 0.615, *p* = 0.0007; λ=1: β = 0.844 ± 0.227, p=0.0015; λ=0: β = 1.013 ± 0.251,

p<0.0001; Tables S5-S7). In the chain network reference models, mean distances increased

fastest and linearly with network size (i.e., number of nodes; Figure 3B). In the triangulated

reference models, mean distances were smallest and increased nonlinearly, slowing their rate of

increase with increasing network size. In the MST reference models, mean distances were

intermediate between the values expected from the chain and triangulated reference models, and

also increased nonlinearly (Figure 3B). The mean distances of empirically observed nests were

between the expected values for chain and MST reference models (Figure 3B). When we scaled

the observed mean distance with the expected mean distance based on each of the three reference

models, we found that the relationship between scaled mean distance and colony size differed

across the three models (Figure 3C). Mean distances scaled to chain networks declined with

colony size and the relationship was statistically significant in almost all data subsets (15/16

datasets; PGLS $\lambda=1$: $\beta = -0.055 \pm 0.024$, $p = 0.031$; 16/16 datasets; $\lambda=0$: $\beta = -0.061 \pm 0.024$, $p =$

0.018; Tables S8 & S9). In contrast, mean distances scaled to triangulated networks increased

significantly with colony size in almost all the data subsets (15/16 datasets; OLS of PIC: $\beta =$

0.236, 95% CI = 0.055, 0.466, $R^2 = 0.348$, $p = 0.0048$; Table S10). Mean distances scaled to

MSTs also increased significantly with colony size (15/16 species subsets), but the effect size

was smaller than for triangulated networks (OLS of PIC: $\beta = 0.11$, 95% CI = -0.022, 0.240, $R^2 =$

0.281, $p = 0.013$; Table S11).

Short-cut Hypothesis

The "short-cut hypothesis," that nests from species with larger colonies would contain more

cycles, was not supported. The relationship between the number of cycles in a nest had a positive

relationship with colony size, but this trend was not statistically significant in the majority of the

data subsets (Figure 3D; 15/16 subsets; N = 21 genera, OLS of PIC: $\beta =0.293$, 95% CI = -0.089,

0.528, $R^2 = 0.160$, $p = 0.072$; Table S12). The triangulated reference networks showed a steep

increase in the number of cycles as a function of network size, and far exceeded the number of

cycles in observed nests given their networks' size (Figure 3E). However, the number of cycles

in observed nests was similar to that expected from the chain and MST reference networks,

which do not produce cycles at any network size (Figure 3E). The measure of cycles scaled to

the triangulated reference models significantly decreased with colony size in all species subsets

(16/16 subsets; Figure 3F; OLS of PIC: β = -0.048, 95% CI = -0.102, -0.015, R^2 = 0.090, p =

0.186; Tables S13).

Figure 3: **Nest Connectivity: branching (A-C) and short-cut (D-F) hypotheses.** (A, D)

Relationship between log colony size (average number of ants in a colony) and (A) network

mean distance and (D) number of cycles in nests. Each data point represents a mean value for a

single species, whiskers indicate standard errors around the mean when multiple nests were

sampled for a given species, colors represent ant subfamily (see legend in A). Dashed line is the

regression fit for PGLS when λ=1, solid when statistically significant and dashed when not

statistically significant; (B, E) The mean distance (B) or number of cycles (E) according to

network size (number of nodes). Networks from observed nests are plotted as grey dots, sized

according to sample size. In (B) lines are mean distances as a function of number of nodes for

chain (red), MST (yellow), and triangulated (blue) reference networks, with 95% confidence

intervals from 1000 simulation runs shown as shading around the lines; in (E) Blue line is the

number of cycles in triangulated networks as a function of number of nodes. Orange line depicts

the number of cycles for both MSTs and chain networks, which contain zero cycles at all

network sizes; (C, F) Relationship between log colony size (average number of ants in a colony)

and scaled mean distance (C) and scaled number of cycles (F). Mean distance (C) is scaled to

chain (red), MST (yellow), and triangulated (blue) networks. Cycles (F) are only scaled to

triangulated networks (blue). Each point represents a mean of scaled values from the nests of

each species. Grey dashed line at y=1 represents values at which observed nests contain the same

value as expected from the reference model. Values below the dashed line indicate networks that

have lower values than expected by the reference model. Values above the dashed line indicate

networks which have values greater than expected according to the reference model. Solid

colored lines indicate statistically significant regression fits for PGLS (λ=1).

Phylogenetic Signal

We detected a significant phylogenetic signal for nest connectivity traits (cycles and mean

distance) but not for colony size or nest subdivision traits (chamber number and size). Most

traits, except colony size and number of cycles, tended to fit a Brownian Motion model of

evolution, however the differences in fit between the alternative models (no signal and estimated

Pagel's lambda) were not significant (<4 AIC units). We did not detect a significant phylogenetic

signal for colony size (Table S14; $p = 0.348$, $\lambda = 0.440$, $AIC_{\lambda} = 99.03$) in any species subsets

(0/19). Colony size had a slightly better fit to the no signal model ($AIC_0 = 97.29$) relative to the

Brownian Motion model ($AIC_{BM} = 99.35$) in all species subsets (19/19 subsets). We did not

detect a significant phylogenetic signal for chamber number (Table S15; $p = 0.091$, $\lambda = 1.07$,

$AIC_{\lambda} = 182.72$) in all species subsets (19/19). Chamber number had a slightly better fit to the

Brownian Motion model ($AIC_{BM} = 180.09$) relative to the no signal model ($AIC_0 = 182.60$) in all

species subsets (19/19 subsets). We did not detect a significant phylogenetic signal for

standardized chamber size (Table S16; $p = 0.084$, $\lambda = 0.71$, $AIC_{\lambda} = 88.79$) in the majority of

species subsets (12/15). Chamber size had a slightly better fit to the Brownian Motion model

($AIC_{BM} = 86.83$) relative to the no signal model ($AIC_0 = 89.02$) in most species subsets (13/15).

Nest mean distance normalized to a chain network had a significant phylogenetic signal (Table

S17; $p = 0.001$, $\lambda = 1.12$, $AIC_{\lambda} = -2.22$) and had a slightly better fit to the Brownian Motion

model ($AIC_{BM} = -4.97$) relative to the no signal model ($AIC_0 = -2.19$) in all species subsets

(16/16). The number of cycles had a significant phylogenetic signal (Table S18; $p = 0.005$, $\lambda =$

1.12, $AIC_{\lambda} = 74.97$), however we could not distinguish between the fit with the Brownian Motion

model ($AIC_{BM} = 72.23$) and the no signal model ($AIC_0 = 72.39$) in most species subsets (13/16).

Figure 4: Nest traits and colony size mapped to a genus-level phylogeny of the ants for all

genera on which we had data. The log of colony size is mapped onto the branches of the

phylogeny, with cooler colors indicating smaller colony sizes and warmer colors larger colony

sizes. Values for the four nest traits we analyzed are depicted as a heatmap, with color indicating

trait value standardized as a z-score. Branching frequency is a measure of how much nests

branch relative to their size; such that values are nest Mean Distance standardized to Chain

networks with values multiplied by -1 to ease interpretation, so that larger values indicate greater

branching. Data are displayed for species subset one, and slashes indicate missing data.

DISCUSSION

Our results support the adaptive hypothesis that nest architecture mitigates the challenges of

larger colony sizes, i.e., more individuals in a colony. Our comparative analysis found that nests

of species with larger colonies are more subdivided and reduce travel distances through

increased branching. Overall, we did not find a strong effect of phylogeny, but our findings lay

out a foundation for future experimental work on the evolution of mechanisms that allow ant

colonies to optimize the structures of their nests.

As predicted by the subdivision hypothesis, the number of chambers was greater in

species with larger colonies and chamber size was not related to colony size (Figure 2). Thus, ant

nests have evolved from the basal single-chambered form to accommodate larger colonies by

adding repeating modules of similar sized chambers rather than increasing chamber (module)

size. These results suggest that nests of species with large colonies are divided into more sub-

units than nests of species with small colonies. This pattern of increased spatial subdivision in

ant nests is shared across a variety of living systems, particularly those that undergo an

evolutionary increase in body size. For instance, the mammalian brain subdivides into more

cortical units in lineages with larger brain sizes (Kaas 2000, 2017). At smaller biological scales,

similar patterns of increased spatial subdivision form the basis for the evolution of multi-

cellularity that made larger body sizes possible (Bonner 1998). Even man-made networks, like

cities become more subdivided as they increase in size (Anas et al. 1998). Given our

correlational findings, it remains unclear if subdivision in nests serves a similar function as it

does in these other systems. Further data are needed on the distribution of individuals throughout

nest chambers to determine if ants utilize chambers in a way that is consistent with subdivision

(e.g., Tschinkel 1999; Murdock and Tschinkel 2015). Nest subdivision may facilitate division of

labor through spatial segregation of tasks (Tofts and Franks 1992; Richardson et al. 2011;

Mersch et al. 2013), and slow down the spread of disease (Pie et al. 2004; Stroeymeyt et al.

2014, 2018). However, the only direct test of the benefit of nest subdivision showed that

subdivision does not increase brood rearing efficiency (Cassill et al. 2002; Tschinkel 2018).

Limits on chamber size may also be imposed by physiological processes such as gas exchange or

temperature regulation (Jones and Oldroyd 2006), or by structural constraints, because large

chambers may collapse. Thus, the causes and consequences of nest subdivision remain to be

tested.

The connectivity of nests changed according to colony size, but only one of our

hypotheses about nest connectivity was supported. The nests of species with larger colonies had

reduced travel distances due to their branching structure, consistent with the branching

hypothesis. In the nests of species with small colonies (i.e., up to ~1000 workers in nests

containing up to ~10 nodes), mean distances tended to match the values expected for chain

networks, and indeed the nests of many small colony species were chains. In the nests of species

with large colonies, mean travel distances were shorter than would be expected for chain

networks but greater than expected for MSTs, and far greater than expected for triangulated

networks (Figure 3B, C). These findings indicate that the nests of species with large colonies

have increased branching (i.e., are more tree-like) relative to the nests of species with smaller

colonies. These findings show that nest branching evolves in association with colony size, and

that increased branching is not necessarily a more "derived" trait or explained by phylogenetic

relatedness alone. For example, the nests of some early branching species (e.g. *Dinoponera*)

have relatively more branching than the nests of some later branching species (e.g. leaf-cutters,

Mycetagrocious; Figures 3C & 4). By increasing the frequency of branching, large colonies may

curb rising travel distances as their nests increase in size (i.e., contain more chambers). Networks

are subject to an inherent tradeoff, in which they cannot simultaneously minimize travel

distances (i.e., be highly interconnected) and minimize their number of edges (i.e., minimize

infrastructure costs). Increased branching is a common way for biological transport networks to

balance this tradeoff (e.g., ant foraging trails: Buhl et al. 2009; slime molds: Tero et al. 2010;

neural networks: Raj and Chen 2011; plant growth forms: Conn et al. 2017), especially as body

size, or population size, increases (Ringo 1991; Banavar et al. 1999; Ercsey-Ravasz et al. 2013;

Cabanes et al. 2015). While these previous studies have identified branching as a way to balance

the connectivity-infrastructure cost tradeoff in single species (i.e., over development or between

individuals), ours is the first study to show this pattern across lineages in an evolutionary

context. Branching is thus a general low-cost way for biological transport networks, including

ant nests, to maintain connectivity as they increase in size.

502   The short-cuts hypothesis, that nests from species with larger colonies will have more

cross-linking paths or cycles, was not supported. Although the number of cycles in a nest

increased with colony size, this increase was not statistically significant (Figure 3D). Cycles

were added at a slower rate than expected for a triangulated network (Figure 3E), which is not

surprising given that triangulations represent the maximum rate at which new cycles could be

added with increasing network size (for 2D networks). This finding lends further support to the

conclusion that the increase in nest connectivity with colony size is due to increased branching

and not the addition of short-cuts. While we did not find support for this adaptive function of

cycles, our results lend some support to the hypothesis that variation in cycles is due to

phylogenetic processes, like drift, evolutionary history, etc., as cycle number had a significant

phylogenetic signal. However, we caution that our analysis conservatively excludes nests that

might have the most cycles because we could not accurately infer the topology of the most

convoluted nests. Our inability to detect a significant increase in the number of cycles with

colony size could be due to this bias in our dataset. More refined imaging techniques could

provide a solution to this issue and expand future work on nest architecture in general.

517   The scarcity of cycles in the nests we measured suggest that the benefits of having

alternative access routes, which help maintain movement (Corson 2010; Katifori et al. 2010;

Blonder et al. 2018, Jyothi et al. 2016) and facilitate forager recruitment (Pinter-Wollman

2015a), might not outweigh the potential costs of additional tunnels. For instance, there are costs

to building and maintaining tunnels (da Silva Camargo et al. 2013; Monaenkova et al. 2015) and

there may be negative consequences of many cross-linking paths, including structural instability

or heightened exposure to risks, like pathogens (Scovell 1983; Viana et al. 2013; Stroeymeyt et

al. 2014, 2018), parasites (Scovell 1983), or predators (Jeanne 1975; LaPolla et al. 2002). Nests

are multi-purpose structures that have other functions, like thermoregulation, humidity control or

ventilation (Jones and Oldroyd 2006; King et al. 2015), which might suffer when nests have

many cycles. In systems where the costs of cross-linking paths are low, like the plasmodia of

slime molds, the mycelia of fungus, or the pheromone trails of ants, cycles are ephemeral and

created for exploration, yet they are pruned once resources are discovered (Bebber et al. 2007;

Buhl et al. 2009; Tero et al. 2010; Latty et al. 2011). Conversely, when the benefits to movement

are very high, cycles become more common as size increases, but this only appears to be true in

some man-made networks, like urban roads or computer routing networks (Levinson 2012;

Jyothi et al. 2016). Future investigation into the building costs of tunnels and the functional

consequences of interconnected nest structures will help reveal why ant nests do not contain

more short-cuts, or cycles.

While we found support for our adaptive hypotheses, we also found that phylogenetic

relatedness explains some of the variation in ant nest structural diversity. Nest traits relating to

connectivity (number of cycles and mean distance normalized to chain networks), had significant

phylogenetic signals, whereas traits relating to subdivision (chamber number and size) did not.

We caution, however, that we did not detect a strong effect of phylogeny overall. The results of

our regression analyses did not differ when we assumed a strong phylogenetic signal ($\lambda=1$) or

none at all ($\lambda=0$). Furthermore, the differences in fit between alternative models of evolution (no

signal, Brownian Motion, and estimated Pagel's lambda) were rarely large enough to robustly

distinguish between models, which may be due to lack of statistical power. Notably, we did not

detect a significant phylogenetic signal for colony size, even though prior studies of ant colony

size suggest that phylogenetic relatedness is a factor in colony size evolution (Boulay et al. 2014;

Ferguson-gow et al. 2014; Burchill and Moreau 2016). While we cannot reach robust

conclusions about the extent of phylogenetic effect on the traits we measured, the co-

evolutionary patterns we analyze here account for potential phylogenetic non-independence.

As with any meta-analysis, our dataset is constrained by the available literature. Many of the nests we found in the literature were built in sandy soils, and so our analysis has an over-representation of species adapted to the climate and ecology of sandy soils (Table S1). Fungus growers are highly-represented in our analysis (represented by the crown spanning from *Mycocepurus* to *Mycetophylax* in Figure 4), and constraints imposed by requirements to maintain healthy fungus gardens could have influenced our findings. Additionally, because we do not have data on the size or age of the colonies that built the majority of nests in our meta-analysis, we might have included immature colonies that have not reached their full size. Further work could investigate whether nest connectivity changes throughout development within a species in natural environments (e.g., Tschinkel 1987, 2004, 2005, 2011) or whether there are species-specific connectivity patterns that can be found at different ages.

In conclusion, we show that ant nests exhibit modular structures that minimize travel distancesf in species with larger colonies. The similarity in how ant nests and the bodies of unitary organisms respond to increasing size suggest that their shared architectural features, branching transportation networks and compartmentalization, provide common solutions to the challenges created by large size across the tree of life.

REFERENCES

Aguilar, J., D. Monaenkova, V. Linevich, W. Savoie, B. Dutta, H.-S. Kuan, M. D. Betterton, M. A. D. Goodisman, and D. I. Goldman. 2018. Collective clog control: Optimizing traffic flow in confined biological and robophysical excavation. Science (80-.). 361:672–677.

Anas, A., R. Arnott, and K. A. Small. 1998. Urban spatial structure. Am. Econ. Assoc. 36:1426–

1464.

Banavar, J. R., A. Maritan, and A. Rinaldo. 1999. Size andform in efficient transportation

networks. Nature 399:130–132.

Bebber, D. P., J. Hynes, P. R. Darrah, L. Boddy, and M. D. Fricker. 2007. Biological solutions to

transport network design. Proc. R. Soc. B Biol. Sci. 274:2307–2315.

Blanchard, B. D., and C. S. Moreau. 2017. Defensive traits exhibit an evolutionary trade-off and

drive diversification in ants. Evolution (N. Y). 71:315–328.

Blonder, B., N. Salinas, L. P. Bentley, A. Shenkin, P. O. Chambi Porroa, Y. Valdez Tejeira, T. E.

Boza Espinoza, G. R. Goldsmith, L. Enrico, R. Martin, G. P. Asner, S. Díaz, B. J. Enquist,

and Y. Malhi. 2018. Structural and defensive roles of angiosperm leaf venation network

reticulation across an Andes–Amazon elevation gradient. J. Ecol. 106:1683–1699.

Bonner, J. T. 2004. Perspective: The Size-Complexity Rule. Evolution (N. Y). 58:1883–1890.

Bonner, J. T. 1988. The Evolution of Complexity by Means of Natural Selection. Princeton

University Press, Princeton, NJ.

Bonner, J. T. 1998. The origins of multicellularity. Integr. Biol. 1:27–36.

Boulay, R., X. Arnan, X. Cerdá, and J. Retana. 2014. The ecological benefits of larger colony

size may promote polygyny in ants. J. Evol. Biol. 27:2856–2863.

Bourke, A. F., and N. R. Franks. 1995. Social Evolution in Ants. Princeton University Press,

Princeton, NJ.

Branstetter, M. G., B. N. Danforth, J. P. Pitts, B. C. Faircloth, P. S. Ward, M. L. Buffington, M.

W. Gates, R. R. Kula, and S. G. Brady. 2017. Phylogenomic Insights into the Evolution of

Stinging Wasps and the Origins of Ants and Bees. Curr. Biol. 27:1019–1025.

Brian, M. V. 1956. Group Form and Causes of Working Inefficiency in the Ant Myrmica rubra

L. Physiol. Zool. 29:173–194.

Brown, J. J., and J. F. A. Traniello. 1998. Regulation of Brood-Care Behavior in the Dimorphic

Castes of the Ant Pheidole morrisi (Hymenoptera : Formicidae): Effects of Caste Ratio ,

Colony Size , and Colony Needs. J. Insect Behav. 11:209–219.

Brown, W. L. 1953. Revisionary Studies in the Ant Tribe Dacetini. Am. Midl. Nat. 50:1.

Buhl, J., J. Gautrais, J.-L. Deneubourg, and G. Theraulaz. 2004a. Nest excavation in ants :

group size effects on the size and structure of tunneling networks. Naturwissenschaften

91:602–606.

Buhl, J., J. Gautrais, R. V Sole, P. Kuntz, S. Valverde, J.-L. Deneubourg, and G. Theraulaz.

2004b. Efficiency and robustness of ant networks of galleries. Eur. Phys. J. B 129:123–129.

Buhl, J., K. Hicks, E. R. Miller, S. Persey, O. Alinvi, and D. J. T. Sumpter. 2009. Shape and

efficiency of wood ant foraging networks. Behav. Ecol. Sociobiol. 63:451–460.

Bullmore, E., and O. Sporns. 2012. The economy of brain network organization. Nat. Rev.

Neurosci. 13:336–349.

Burchill, A. T., and C. S. Moreau. 2016. Colony size evolution in ants: macroevolutionary

trends. Insectes Soc. 62:291–298.

Burd, M., N. Shiwakoti, M. Sarvi, and G. Rose. 2010. Nest architecture and traffic flow : large

potential effects from small structural features. Ecol. Entomol. 35:464–468.

Cabanes, G., E. Van Wilgenburg, M. Beekman, and T. Latty. 2015. Ants build transportation

networks that optimize cost and efficiency at the expense of robustness. Behav. Ecol.

26:223–231.

Calder, W. A. 1984. Size, Function, and Life History. Harvard University Press, Cambridge,

MA.

Cassill, D., W. R. Tschinkel, and S. B. Vinson. 2002. Nest complexity , group size and brood

rearing in the fire ant , Solenopsis invicta. Insectes Soc. 49:158–163.

Cheverud, J. M. 1982. Relationships among ontogenetic, static, and evolutionary allometry. Am.

J. Phys. Anthropol. 59:139–149.

Conn, A., U. V. Pedmale, J. Chory, and S. Navlakha. 2017. High-Resolution Laser Scanning

Reveals Plant Architectures that Reflect Universal Network Design Principles. Cell Syst.

5:53-62.e3. Elsevier Inc.

Corson, F. 2010. Fluctuations and Redundancy in Optimal Transport Networks. 048703:1–4.

Csárdi, G. 2020. Package ' igraph ': Network Analysis and Visualization.

da Silva Camargo, R., J. F. S. Lopes, and L. C. Forti. 2013. Energetic cost of digging behavior in

workers of the leaf-cutting ant Atta sexdens (Fabricius). Rev. Bras. Entomol. 57:401–404.

Dial, K. P., E. Greene, and D. J. Irschick. 2008. Allometry of behavior. Trends Ecol. Evol.

23:394–401.

Dodds, P. S. 2010. Optimal form of branching supply and collection networks. Phys. Rev. Lett.

104:1–4.

Dornhaus, A., S. Powell, and S. Bengston. 2012. Group size and its effects on collective

organization. Annu. Rev. Entomol. 57:123–41.

Ercsey-Ravasz, M., N. T. Markov, C. Lamy, D. C. VanEssen, K. Knoblauch, Z. Toroczkai, and

H. Kennedy. 2013. A Predictive Network Model of Cerebral Cortical Connectivity Based

on a Distance Rule. Neuron 80:184–197.

Fairbairn, D. J. 1997. Allometry for sexual size dimorphism: Pattern and process in the

coevolution of body size in males and females. Annu. Rev. Ecol. Syst. 28:659–687.

Ferguson-gow, H., S. Sumner, A. F. G. Bourke, and K. E. Jones. 2014. Colony size predicts

division of labour in attine ants. Proc. R. Soc. B Biol. Sci. 281:20141411.

Frankino, W. A., B. J. Zwaan, D. L. Stern, and P. M. Brakefield. 2005. Natural selection and

developmental constraints in the evolution of allometries. Science (80-.). 307:718–720.

Freckleton, R. P. 2009. The seven deadly sins of comparative analysis. J. Evol. Biol. 22:1367–

1375.

Gillooly, J. F., C. Hou, and M. Kaspari. 2010. Eusocial insects as superorganisms: Insights from

metabolic theory. Commun. Integr. Biol. 3:360–362.

Gould, S. J. 1966. Allometry and size in ontogeny and phylogeny. Biol. Rev. Camb. Philos. Soc.

41:587–640.

Gould, S. J. 1971. Geometric Similarity in Allometric Growth: A Contribution to the Problem of

Scaling in the Evolution of Size. Am. Nat. 105:113–136.

Harmon, A. L., M. Pennell, C. Brock, J. Brown, W. Challenger, J. Eastman, R. Fitzjohn, R. Glor,

G. Hunt, L. Revell, G. Slater, J. Uyeda, and J. Weir. 2020. Package ' geiger ': Analysis of

Evolutionary Diversification.

Hobson, E. A., M. J. Silk, N. H. Fefferman, D. B. Larremore, P. Rombach, S. Shai, and N.

Pinter-Wollman. 2021. A guide to choosing and implementing reference models for social

network analysis. Biol. Rev., doi: 10.1111/brv.12775.

Holldobler, B., and E. O. Wilson. 2009. The superorganism: the beauty, elegance, and

strangeness of insect societies Title. W.W. Norton, New York.

Hölldobler, B., and E. O. Wilson. 1990. The Ants. Belknap Press, Cambridge, MA.

Huxley, J. 1932. Problems of Relative Growth. Methuen and Co., Ltd., London.

Jeanne, R. L. 1975. The Adaptiveness of Social Wasp Nest Architecture. Q. Rev. Biol. 50:267–

287.

Jones, J. C., and B. P. Oldroyd. 2006. Nest Thermoregulation in Social Insects. Adv. In Insect

Phys. 33:153–191.

Jyothi, S. A., A. Singla, P. B. Godfrey, and A. Kolla. 2016. Measuring and Understanding

Throughput of Network Topologies. Int. Conf. High Perform. Comput. Networking, Storage

Anal. SC 0:761–772.

Kaas, J. H. 2017. Evolution of the Brain, Cognition, and Emotion in Vertebrates. Evol. Brain,

Cogn. Emot. Vertebr. 59–80.

Kaas, J. H. 2000. Why is Brain Size so Important:Design Problems and Solutions as Neocortex

Gets Biggeror Smaller. Brain Mind 1:7–23.

Katifori, E., G. J. Szöllosi, and M. O. Magnasco. 2010. Damage and fluctuations induce loops in

optimal transport networks. Phys. Rev. Lett. 104:1–4.

Kempes, C. P., S. Dutkiewicz, and M. J. Follows. 2012. Growth , metabolic partitioning , and the

size of microorganisms. Proc. Natl. Acad. Sci. 109:495–500.

677 King, H., S. A. Ocko, and L. Mahadevan. 2015. Termite mounds harness diurnal temperature

678   oscillations for ventilation. Proc. Natl. Acad. Sci. 112:11589–11593.

679 Kleiber, M. 1947. Body size and metabolic rate. Physiol. Rev. 27:511–541.

680 Kwapich, C. L., G. Valentini, and B. Hölldobler. 2018. The non-additive effects of body size on

681   nest architecture in a polymorphic ant. Philos. Trans. R. Soc. B 373:20170235.

682 LaPolla, J. S., U. G. Mueller, M. Seid, and S. P. Cover. 2002. Predation by the army ant

683   Neivamyrmex rugulosus on the fungus-growing ant Trachymyrmex arizonensis. Insectes

684   Soc. 49:251–256.

685 Latty, T., K. Ramsch, K. Ito, T. Nakagaki, D. J. T. Sumpter, M. Middendorf, and M. Beekman.

686   2011. Structure and formation of ant transportation networks. J. R. Soc. Interface 8:1298–

687   1306.

688 Levinson, D. 2012. Network structure and city size. PLoS One 7:e29721.

689 Lorenz, D. M., A. Jeng, and M. W. Deem. 2011. The emergence of modularity in biological

690   systems. Phys Life Rev. 8:129–160.

691 McKown, A. D., H. Cochard, and L. Sack. 2010. Decoding leaf hydraulics with a spatially

692   explicit model: Principles of venation architecture and implications for its evolution. Am.

693   Nat. 175:447–460.

694 McMahon, T. A. 1975. Using body size to understand the structural design of animals:

695   quadrupedal locomotion. J. Appl. Physiol. 39:619–627.

696 McMahon, T. A., and J. T. Bonner. 1983. On Size and Life. Scientific American Books, Inc,

697   New York.

Mersch, D. P., A. Crespi, and L. Keller. 2013. Tracking individuals shows spatial fidelity is a

key regulator of ant social organization. Science (80-.). 340:1090–3.

Monaenkova, D., N. Gravish, G. Rodriguez, R. Kutner, M. A. D. Goodisman, and D. I.

Goldman. 2015. Behavioral and mechanical determinants of collective subsurface nest

excavation. J. Exp. Biol. 218:1295–1305.

Murdock, T. C., and W. R. Tschinkel. 2015. The life history and seasonal cycle of the ant,

Pheidole morrisi Forel, as revealed by wax casting. Insectes Soc. 62:265–280.

Newman, M. E. J. 2018. Networks: An Introduction. 2nd Editio. Oxford University Press,

Oxford.

Niklas, K. J., and S. A. Newman (eds). 2016. Multicellularity: Origins and Evolution. MIT Press,

Cambridge, MA.

O'Donnell, S., and S. J. Bulova. 2007. Worker connectivity: A review of the design of worker

communication systems and their effects on task performance in insect societies. Insectes

Soc. 54:203–210.

Okie, J. G. 2013. General models for the spectra of surface area scaling strategies of cells and

organisms: Fractality, geometric dissimilitude, and internalization. Am. Nat. 181:421–439.

Orme, A. D., R. Freckleton, G. Thomas, T. Petzoldt, N. Isaac, W. Pearse, and M. D. Orme. 2018.

Package ' caper '□: Comparative Analyses of Phylogenetics and Evolution in R.

Pacala, S. W., D. M. Gordon, and H. C. J. Godfray. 1996. Effects of social group size on

information transfer and task allocation. Evol. Ecol. 10:127–165.

Pagel, M. 1999. Inferring historical patterns of biological evolution. Nature 401:877–884.

Paradis, E., S. Blomberg, B. Bolker, J. Brown, J. Claude, H. S. Cuong, R. Desper, G. Didier, B.

Durand, J. Dutheil, R. Ewing, O. Gascuel, T. Guillerme, C. Heibl, A. Ives, B. Jones, F.

Krah, D. Lawson, V. Lefort, P. Legendre, J. Lemon, E. Marcon, R. McCloskey, J.

Nylander, R. Opgen-Rhein, A.-A. Popescu, M. Royer-Carenzi, K. Schliep, K. Strimmer,

and D. de Vienne. 2018. Package 'ape': Analyses of Phylogenetics and Evolution.

Perna, A., C. Jost, E. Couturier, S. Valverde, S. Douady, and G. Theraulaz. 2008. The structure

of gallery networks in the nests of termite Cubitermes spp. revealed by X-ray tomography.

Naturwissenschaften 95:877–884.

Perna, A., and T. Latty. 2014. Animal transportation networks. J. R. Soc. Interface 11.

Peters, R. H. 1983. The Ecological Implications of Body Size. Cambridge University Press.

Pie, M. R., R. B. Rosengaus, and J. F. A. Traniello. 2004. Nest architecture , activity pattern ,

worker density and the dynamics of disease transmission in social insects. J. Theor. Biol.

226:45–51.

Pinter-Wollman, N. 2015a. Nest architecture shapes the collective behaviour of harvester ants.

Biol. Lett. 11:20150695-.

Pinter-Wollman, N. 2015b. Persistent variation in spatial behavior affects the structure and

function of interaction networks. Curr. Zool. 61:98–106.

Pinter-Wollman, N., A. Bala, A. Merrell, J. Queirolo, M. C. Stumpe, S. Holmes, and D. M.

Gordon. 2013. Harvester ants use interactions to regulate forager activation and availability.

Anim. Behav. 86:197–207. Elsevier Ltd.

Pinter-wollman, N., A. Penn, G. Theraulaz, and S. M. Fiore. 2018. Interdisciplinary approaches

for uncovering the impacts of architecture on collective behaviour. Philos. Trans. R. Soc. B 373:20170232.

R Core Team. 2019. R: A language environment for statistical computing.

Raj, A., and Y. H. Chen. 2011. The wiring economy principle: Connectivity determines anatomy in the human brain. PLoS One 6.

Revell, M. L. J. 2017. Package ' phytools ':Phylogenetic Tools for Comparative Biology (and Other Things).

Richardson, T. O., K. Christensen, N. R. Franks, H. J. Jensen, and A. B. Sendova-franks. 2011. Ants in a Labyrinth: A Statistical Mechanics Approach to the Division of Labour. PLoS One 6:e18416.

Ringo, J. L. 1991. Neuronal interconnection as a function of brain size. Brain. Behav. Evol. 38:1–6.

Rueden, C. T., J. Schindelin, M. C. Hiner, B. E. Dezonia, A. E. Walter, E. T. Arena, and K. W. Eliceiri. 2017. ImageJ2: ImageJ for the next generation of scientific image data. BMC Bioinformatics 18:1–26.

Ruppert, E. E., R. S. Fox, and R. D. Barnes. 2004. Invertebrate Zoology: A Functional Evolutionary Approach. Thomson-Brooks/Cole.

Schmidt-Nielson, K. 1984. Scaling: Why is Animal Size So Important? Cambridge University Press, New York.

Scovell, E. 1983. Dominance and reproductive success among slave-making worker ants. Nature 304:724–725.

Seeley, T. D. 1989. The honey bee colony as a superorganism. Am. Sci. 77:546–553.

Stroeymeyt, N., B. Casillas-Pérez, and S. Cremer. 2014. Organisational immunity in social

insects. Curr. Opin. Insect Sci. 5:1–15.

Stroeymeyt, N., A. V Grasse, A. Crespi, D. P. Mersch, S. Cremer, and L. Keller. 2018. Social

network plasticity decreases disease transmission in a eusocial insect. Science (80-.).

362:941–945.

Tekin, E., D. Hunt, M. G. Newberry, and V. M. Savage. 2016. Do Vascular Networks Branch

Optimally or Randomly across Spatial Scales? PLoS Comput. Biol. 12:e1005223.

Tero, A., S. Takagi, T. Saigusa, K. Ito, D. P. Bebber, M. D. Fricker, K. Yumiki, R. Kobayashi,

and T. Nakagaki. 2010. Rules for biologically inspired adaptive network design. Science

(80-.). 327:439–442.

Thompson, D. W. 1917. On Growth and Form. Cambridge University Press, Cambridge, MA.

Tofts, C., and N. R. Franks. 1992. Doing the right thing: Ants, honeybees and naked mole-rats.

Trends Ecol. Evol. 7:346–349.

Tschinkel, W. R. 2010. Methods for casting subterranean ant nests. J. Insect Sci. 88:1–17.

Tschinkel, W. R. 1987. Seasonal life history and nest architecture of a winter-active ant,

Prenolepis imparis. Insectes Soc. 34:143–164.

Tschinkel, W. R. 1999. Sociometry and sociogenesis of colonies of the harvester ant,

Pogonomyrmex badius: distribution of workers, brood and seeds within the nest in relation

to colony size and season. Ecol. Entomol. 24:222–237.

Tschinkel, W. R. 2003. Subterranean ant nests: trace fossils past and future? Palaeogeogr.

Palaeoclimatol. Palaeoecol. 192:321–333.

Tschinkel, W. R. 2018. Testing the effect of a nest architectural feature in the fire ant Solenopsis

invicta (Hymenoptera: Formicidae). Myrmecological News 27:1–5.

Tschinkel, W. R. 2015. The architecture of subterranean ant nests: beauty and mystery underfoot.

J. Bioeconomics 17:271–291.

Tschinkel, W. R. 2005. The nest architecture of the, ant Camponotus socius. J. Insect Sci. 5:1–

18.

Tschinkel, W. R. 2004. The nest architecture of the Florida harvester ant , Pogonomyrmex

badius. J. Insect Sci. 21:1–19.

Tschinkel, W. R. 2011. The nest architecture of three species of north Florida Aphaenogaster

ants. J. insect Sci. 11:1–30.

Turner, R. 2020. Package ' deldir ': Delaunay Triangulation and Dirichlet (Voronoi) Tessellation

Author.

Vaes, O., A. Perna, and C. Detrain. 2020. The effect of nest topology on spatial organization and

recruitment in the red ant Myrmica rubra. Sci. Nat. 107. The Science of Nature.

Viana, M. P., V. Fourcassié, A. Perna, F. Costa, and C. Jost. 2013. Accessibility in networks :

A useful measure for understanding social insect nest architecture. Chaos, Solitons and

Fractals 46:38–45. Elsevier Ltd.

West, G. B., J. H. Brown, and B. J. Enquist. 1997. A General Model for the Origin of Allometric

Scaling Laws in Biology. Science (80-.). 276:122–126.

Wheeler, W. M. 1910. Ants: Their structure, Development and Behavior. Columbia University

Press, New York.

804