TITLE PAGE

Title: Modularity and connectivity of nest structure scale with colony size

Authors: Julie S. Miller[1], Emma Wan[1] & Noa Pinter-Wollman[1]

Author Affiliations
1. Department of Ecology and Evolutionary Biology, University of California Los Angeles, 612 Charles E. Young Dr. East, Los Angeles, CA 90095-7246.

ABSTRACT

Large body sizes have evolved structures to facilitate resource transport. Like unitary organisms,

social insect colonies must transport information and resources, and colonies with more

individuals may experience transport challenges similar to large-bodied organisms. In ant

colonies, transport occurs in the nest, which may consist of structures that facilitate movement.

We examine three attributes of nest structures that could mitigate transport challenges related to

colony size: (1) subdivision - nests of species with large colonies are more subdivided to reduce

viscosity of large crowds; (2) branching - nest tunnels increase branching in species with large

colonies to reduce travel distances; and (3) short-cuts – nests of species with large colonies have

cross-linking tunnels to connect distant parts of the nest and create alternative routes. We test

these hypotheses by comparing nest structures of species with different colony sizes in

phylogenetically controlled meta-analyses. Our findings support the subdivision and branching

hypotheses. The nests of species with large colonies contain more, but not larger, chambers and

reduce travel distances by increasing branching frequency. The similarity in how ant nests and

the bodies of unitary organisms respond to increasing size suggests structural solutions that are

common across levels of biological organization.

INTRODUCTION

Organisms with large body sizes are faced with the challenge of transporting nutrients, waste

products, and signals across greater distances compared to organisms with small bodies (Bonner

2004). This physical challenge has been resolved through a variety of structural features across

the tree of life (McMahon and Bonner 1983), including multi-cellularity (Niklas and Newman

2016), body plans that increase surface-area-to-volume ratios (e.g., jellyfish and sponges)

(Rupppert et al. 2004), branching vascular systems in animals and plants (West G. B. et al. 1997;

Banavar et al. 1999), and compartmentalization of specialized tissues into organs (Gould 1966).

Social insect colonies have frequently been compared to whole organisms (Wheeler 1910; Seeley

1989; Holldobler and Wilson 2009), and large colonies may experience transport challenges that

are similar to those of large bodied organisms (Gillooly et al. 2010). In larger colonies, the speed

and/or reliability with which resources or information reach disparate parts of the colony will

depend on transfer occurring among more individuals and across greater distances (Pacala et al.

1996; O'Donnell and Bulova 2007; Dornhaus et al. 2012).

Transport of resources and information in social insect colonies may depend on the

layout of their nests. Nest structure affects the routes that individuals take as they move resources

and information and therefore the way in which individuals interact (Pinter-Wollman et al. 2013;

Pinter-Wollman 2015b; Tschinkel 2015; Pinter-wollman et al. 2018; Vaes et al. 2020) and

behave as a collective (Burd et al. 2010; Pinter-Wollman 2015a). Having more individuals may

slow the delivery of resources and information, but the physical structure of the nest can promote

rapid or efficient movement. Here we investigate how features of the nest, including subdivision,

branching, and short-cuts, relate to typical colony size, i.e., the average number of individuals in

a colony, of different ant species.

Subdivision, or compartmentalization, is one way in which large systems mitigate the

challenges of size. For example, large cities subdivide into smaller units, such as neighborhoods

(Anas et al. 1998) and large multicellular organisms consist of subdivided cells and organs

(Kempes et al. 2012; Okie 2013). The size of the units are constrained by the costs of expanding

surface area-to-volume ratios (Bonner 1988; Lorenz et al. 2011). In social insect nests, larger

chambers can hold more individuals, but make it harder for individuals to contact each other

(Brian 1956; Aguilar et al. 2018). Constraining the size of nest chambers reduces the number of

individuals occupying each chamber while reducing the distances they need to travel to interact.

The "subdivision hypothesis" predicts that the nests of species with larger colony sizes will

consist of more chambers compared to nests of species with small colonies, while chamber sizes

will not differ in relation to colony size.

Another way that large systems improve transport is by increasing their connectivity and

in turn, reducing travel distances. Travel distances can be reduced by branching paths. Many

natural systems exhibit branching structures which reduce travel distances without the addition

of cross-linking paths. For example, hierarchical branching structures are common in the

vascular networks of animals (West G. B. et al. 1997; Tekin et al. 2016) and plants (McKown et

al. 2010; Blonder et al. 2018). Branching structures are also common when resources enter

through a single site and need to be homogenously distributed throughout the system (Dodds

2010). The "branching hypothesis" predicts that nests of species with larger colonies will have

tunnels with high branching frequency to reduce travel distances in the nest.

Reducing travel distances and increasing connectivity can further be accomplished by

adding cross-linking paths, or short-cuts, that connect distant parts of a network and create

alternative access routes to maintain traffic flow (Corson 2010; Katifori et al. 2010; Bullmore

and Sporns 2012; Blonder et al. 2018). Indeed, the frequency of cross-linking paths often

increases with population size and traffic flow (e.g,. roads: Levinson 2012; and computer routing

networks: Jyothi et al. 2016). The "short-cut hypothesis" predicts that nests of species with larger

colonies will contain more cross-linking paths than nests of species with smaller colonies.

Despite the transport benefits of adding short-cuts, every additional connection has a building

and maintenance cost, and a highly connected nest may be structurally unsound (Monaenkova et

al. 2015).

Ants offer an ideal system to study the interaction between group size and structural

adjustments to size due to their large range of colony sizes (Bourke and Franks 1995) and the

structural diversity of their nests (Tschinkel 2015). We focus our study on ground-nesting

species because of all ants, they have the greatest potential to control the structure of their nests.

Prior studies of subterranean ant nest structure in relation to colony size have focused on single

species, showing that ant nests grow in volume and increase chamber size and/or number over

time (Tschinkel 1987, 2004, 2005, 2011, 2015). Fewer studies have examined how tunnel

connectivity changes with colony development (but see Buhl et al. 2004a), so branching patterns

in ant nests are not well understood. These studies on the ontogeny of nest structure do not

provide information on the evolution of nest structures in relation to their mature colony size,

which we examine here in a phylogenetically controlled cross-species comparison.

We hypothesize that if ant colonies face size-associated challenges in transport and

communication, analogous to other systems, then the structure of nests should relate to colony

size in ways that ease the movement of individuals, resources, and information throughout the

nest. We examine the relationship between three attributes of nest structure: (1) subdivision, (2)

branching and (3) and short-cuts, with colony size by comparing different ant species in

phylogenetically controlled meta-analyses.

METHODS

Obtaining nest and colony size data

To test our hypotheses, we gathered data on nest architecture of subterranean ant species. We

identified publications with graphical representations of ant nests by searching for the terms

"nest", "structure", "architecture", "casting", and "excavation/ed" on Google Scholar and Web

of Science. We identified 24 papers that contained photographs or illustrations of castings or

excavations of complete ant nests with distinguishable entrances, tunnels, and chambers. For the

subdivision hypothesis, we included data from an additional 3 papers containing tables with

measures of chamber size and number, even though nest images were not available (Table S1).

We excluded 14 publications from our dataset because interior chambers and/or tunnels were

obscured, nests were constructed under artificial conditions, or were from incipient colonies

(Table S2). Authors were contacted for additional images when parts of the nest were obscured,

accounting for approximately 10% of the images we used. Physical castings of unpublished nests

were also obtained (Table S1). When more than one nest was available for a species, we took an

average of all measurable nests.

To relate nest structure with colony size – i.e., the number of ants in a colony - we

searched the scientific literature and the web (AntWiki.org and AntWeb.org) for information on

the number of workers in a mature colony. For the majority of species, we used the average

colony size from the literature. When only a range (minimum - maximum) for number of ants

was available, we took the midpoint. Information about the number of ants in a colony at the

time of nest excavation or casting was available in less than half of the studies we obtained nest

structures from, but when available, we used the reported number of adult workers for those

particular nests. When colony size was not available for a particular species, or when nest

identity at the species level was unknown, we used colony sizes representative of the order of

magnitude for the genus obtained from a sister species. This approximation was necessary for

Forelius sp - in which colony size from *F. pruinosus* (10^4) was used, and for *Dorymyrmex bureni*

- in which colony size from *Dorymyrmex bicornis* (10^3) was used (Table S1). We excluded

Pheidole oxyops and *Acromyrmex subterraneus* nests from our analysis because colony size was

not available for these species, and because we could not make a reliable approximation due to

substantial variation in their colony size (Table S2).

Nest Subdivision Hypothesis

To test the subdivision hypothesis, we related colony size to the number and size of nest

chambers. We counted chambers in 296 nests from 43 species in 24 genera and measured their

sizes in 188 nests from 37 species in 21 genera. To quantify the size of chambers, we measured

the maximum width of each chamber using ImageJ (Rueden et al. 2017). We excluded nests

from the size analysis if images did not include a scale bar or if species identity was unknown for

a nest (e.g., *Forelius sp.*). We used chamber width as a proxy for chamber area because we relied

on 2D images of 3D structures. When we had access to physical castings, we measured chamber

width directly from the structures using a measuring-tape. We validated that chamber width was

a reliable indicator of chamber area using an independent dataset (see Supplemental Materials).

To allow for cross-species comparison, we standardized chamber width to worker size.

Worker size was measured using a standard metric of ant body size, Weber's length (Brown

1953), using ant profile images containing a scale bar from AntWeb.org or other publications. In

monomorphic species, one ant length measurement was made per species. In species with

continuous worker polymorphism, we took the average length from all available images. In

species with discrete worker polymorphism (i.e., majors and minors), we used a weighted

average, based on typical frequency of majors according to the literature (Brown and Traniello

1998; Murdock and Tschinkel 2015). Standardized widths of all chambers in each nest were

averaged to obtain a single standardized chamber width per nest.

Nest Connectivity

Nests as Networks

To examine nest connectivity, we depicted each nest as a network, as in (Buhl et al. 2004b; Perna

et al. 2008; Viana et al. 2013; Pinter-Wollman 2015a; Kwapich et al. 2018), by manually

assigning a node ID to each chamber, nest entrance, tunnel junction, and ends of tunnels that did

not reach a chamber ('end nodes') and connecting these nodes with edges depicting tunnels.

When quantifying nest casts as networks we excluded the "mushroom top" where the casting

material pools around the nest entrance when it is poured (Tschinkel 2010). We designated the

nest entrance as the narrowest section of the mushroom top. We obtained networks of 170 nests

of 38 species from 21 genera, excluding images in which the network structure could not be

inferred because tunnel connections were obscured or ambiguous (Table S1 & 2). We excluded

polydomous nests because it was not possible to calculate the network measures we used, mean

travel distances and number cycles (see below), without information about the structure of trails

connecting different nests.

Branching Hypothesis

To test the if branching reduces travel distances in large colonies, i.e., the 'branching hypothesis'

we quantified the length of the travel paths in a nest as the mean distance of a network. Mean

distance of a network is the sum of the lengths of the shortest paths that connect all possible pairs

of nodes, divided by the number of all possible node pairs:

$$Mean\ Distance = \frac{1}{n(n-1)} \sum_{i=1}^{n} d_{ij}$$

where n is the number of nodes, and d is the shortest path between all node pairs i and j

(Newman 2018). We used the "mean_distance()" function in igraph R package (Csárdi 2020).

To determine if observed nests had more branching than expected based on their size,

i.e., the number of nodes in the network, we compared the mean distance of networks from

observed nests to mean distance of reference models of networks with known branching

properties. We evaluated how mean distance relates to nest size under different assumptions

about how new nodes are added to a network (i.e., generative models (Hobson et al 2021)). We

used three generative reference models that represent the upper and lower bounds on

connectivity for how nest networks might increase in size (Buhl et al. 2004b; Bebber et al. 2007;

Tero et al. 2010; Latty et al. 2011): (1) chain networks, in which new nodes are added to the last

node in a chain – i.e., no branching; (2) triangulated networks, in which new nodes are connected

to at least two other nodes such that they form a triangle while avoiding edge overlap. We

generated random triangulated networks using Delaunay triangulation with the deldir() function

in the 'deldir' R package (Turner 2020). The x,y coordinates of the nodes to connect were

randomly generated using the sample() function; and (3) minimum spanning trees (MSTs), in

which each new node is added to a terminal node to form a new branch in a hierarchical

structure. New nodes are added such that the total length of the network is minimized and no

cycles are formed. We generated MST networks by pruning the triangulated graphs generated in

the prior step using the 'mst' function in the 'igraph' R package (Csárdi 2020). Each of the above

generative processes was run for network sizes ranging from 3 to 100 nodes, in increments of

one node. We repeated this process 1000 times for each network size for both triangulated

networks and MSTs because they were probabilistic generative processes, but not for the chain

network generation because it has only one solution. For each generated network, we calculated

the mean distance to create reference distributions for comparison with the observed data

(Hobson et al. 2020; Figure 2B). See R code in the supplementary materials.

We further scaled the observed networks to the reference models to relate mean distance

to colony size (number of individuals) in a way that accounted for expected changes in network

size (number of nodes). To create scaled measures of mean distance, we divided the observed

mean distance by the mean value of mean distance of the reference networks (from the 1000

simulations) for each network size (3-100 nodes). The closer the scaled mean distance value is to

1, the more similar the observed network is to the reference model. Scaled values greater than 1

indicate that the observed network has a higher mean distance than expected according to the

reference model. Likewise, scaled values less than 1 indicate a smaller mean distance than

expected according to the reference model. We related these scaled values with colony size using

phylogenetically controlled comparisons as described below in Statistical Analyses (Figure 2C).

Short-Cut Hypothesis

To test if larger colonies have nests with more short-cuts, i.e., the short-cut hypothesis, we

quantified short cuts using the number of cycles in a network (Buhl et al. 2004b; Pinter-Wollman

2015a). A cycle is a path along a network that returns to its starting node after passing through at

least one other node (Newman 2018). The number of cycles in a nest is:

$$Number\ of\ cycles = \ m - n + 1$$

where n is the number of nodes and m is the number of edges. Cycles are a good approximation

of the path redundancy of a network, or presence of short-cuts, indicating how often there is

more than one route to any given part of the nest (Perna and Latty 2014).

To account for how network size affects the rate at which new cycles are added, we

compared the observed nests to the reference models of networks described above. Following a

similar procedure to the one described for network mean distance, we calculated the number of

cycles in triangulated networks (Figure 2E) and then scaled the number of cycles in observed

nests to triangulated networks of equivalent size. Because both chain and MST reference models

do not produce cycles, we did not include these models when we scaled the observed nests. We

related the scaled number of cycles in a nest with colony size using phylogenetically controlled

comparisons as described below (Figure 2F).

Statistical Analyses

We examined the effect of colony size on the above nest measures while controlling for

phylogeny. We used a pruned evolutionary tree of the ants (Formicidae) from (Blanchard and

Moreau 2017) and calculated the mean trait value for each species. A complete species-level tree

was not available for all species in our dataset, so we selected only one species per genus and ran

our analysis using a genus-level tree. When there were data for more than one species per genus,

we used the species which had the most nests in our dataset (subset 1 in Tables S3-S13). To

ensure this sub-setting of species did not bias our results, we re-ran each analysis with previously

excluded species as representatives of a genus, and report the proportion of those runs that were

statistically significant. See supplementary materials for results from all runs (Tables S3-13).

We ran phylogenetic generalized least squares (PGLS) comparing each nest trait to the

log of colony size. PGLS is a multiple regression that controls for the degree of similarity

between species based on phylogenetic distance in a co-variance matrix. We ran PGLS by

estimating the phylogenetic signal, lambda λ, through maximum likelihood in the "caper"

package v1.0.1 in R (Orme et al. 2018), with λ=0 meaning no phylogenetic signal, and λ=1

meaning the observations completely match phylogeny (Pagel 1999). Due to the small sample

size (20 - 24 genera) we could not estimate lambda with high confidence (Freckleton 2009).

Therefore, we further conducted PGLS for fixed lambda values (0, 1) using the "ape" package

v5.1 in R (Paradis et al. 2018). When our models did not meet the assumption of normality of

residuals we analyzed these relationships by calculating phylogenetic independent contrasts

(PICs) instead (Paradis et al. 2018). PIC transformed data were modeled using OLS regression,

with the intercept set to the origin (R Core Team 2019). Due to small sample sizes, we calculated

95% confidence intervals for OLS model coefficients using a bootstrap (1000 iterations).

RESULTS

Subdivision Hypothesis

Both predictions of the subdivision hypothesis, that chamber number would be larger in species

with large colony sizes and that chamber size would not relate to colony size, were supported.

We found a statistically significant positive relationship between number of chambers per nest

and colony size (Figure 1A; N= 24 genera, OLS of PIC: $\beta = 4.024$, 95% CI = 2.014, 6.292, R^2 =

0.560, $p < 0.0001$; Table S3). The effect size of this relationship was large ($3.7 < \beta < 4.3$ across

all data subsets) and statistically significant in all species subsets (19/19 datasets). We did not

find a statistically significant relationship between the average standardized chamber width and

colony size (Figure 1B; N=21 genera, OLS of PICs: β = 0.325, 95% CI = -0.086, 0.932, R^2 =

0.102, p = 0.15; Table S4) in the majority of the species subsets (14/15 datasets). This lack of

statistical significance could be attributed to low statistical power (power < 0.4 from a post-hoc

power analysis based on N=21 genera, effect size = 0.1, and alpha = 0.05).

Figure 1: Relationship between the log of colony size (number of ants in a colony) and (A)

average number of chambers per nest or (B) average standardized chamber width. Each data

point represents a mean value for a single species, whiskers indicate standard errors when

multiple nests were sampled. Colors represent ant subfamilies. Lines are fits from the PGLS

(λ=1) with a statistically significant fit as a solid line and a relationship that is not statistically

significant is a dashed line.

Branching Hypothesis

The "branching hypothesis", that branching would reduce travel distances in nests of species

with large colonies, was supported. Raw, unscaled, mean distance of nests significantly increased

with colony size (Figure 2A; N = 21 genera, PGLS λ=MaxLL: β =1.36 ± 0.247, λ=0.585, R^2 =

0.615, p = 0.0007; λ=1: β = 0.844 ± 0.227, p=0.0015; λ=0: β = 1.013 ± 0.251, p<0.0001; Tables

S5-S7). In the chain network reference models, mean distance increased fastest and linearly with

network size (i.e., number of nodes; Figure 2B). In the triangulated reference models, mean

distances were smallest and increased nonlinearly, slowing their rate of increase with increasing

network size. In the MST reference models, mean distances were intermediate between the

values expected from the chain and triangulated reference models, and also increased nonlinearly

(Figure 2B). The mean distances of empirically observed nests were between the expected

values for chain and MST reference models (Figure 2B). When we scaled the observed mean

distance with the expected mean distance based on each of the three reference models, we found

that the relationship between scaled mean distance and colony size differed across the three

models (Figure 2C). Mean distances scaled to chain networks declined with colony size and the

relationship was statistically significant in almost all data subsets (15/16 datasets; PGLS λ=1: β

= -0.055± 0.024, p = 0.031; 16/16 datasets; λ=0: β = -0.061 ± 0.024, p = 0.018; Tables S8 & S9).

In contrast, mean distances scaled to triangulated networks increased significantly with colony

size in almost all the data subsets (15/16 datasets; OLS of PIC: β = 0.236, 95% CI = 0.055,

0.466, R^2 = 0.348, p = 0.0048; Table S10). Mean distances scaled to MSTs also increased

significantly with colony size (15/16 species subsets), but the effect size was smaller than for

triangulated networks (OLS of PIC: β = 0.11, 95% CI = -0.022, 0.240, R^2 = 0.281, p = 0.013;

Table S11).

Short-cut Hypothesis

The "short-cut hypothesis," that nests from species with larger colonies would contain more cycles, was not supported. The relationship between the number of cycles in a nest had a positive relationship with colony size, but this trend was not statistically significant in the majority of the data subsets (Figure 2D; 15/16 subsets; N = 21 genera, OLS of PIC: β =0.293, 95% CI = -0.089, 0.528, R^2 = 0.160, p = 0.072; Table S12). The triangulated reference networks showed a steep increase in the number of cycles as a function of network size, and far exceeded the number of cycles in observed nests given their networks' size (Figure 2E). However, the number of cycles in observed nests was similar to that expected from the chain and MST reference networks, which do not produce cycles at any network size (Figure 2E). The measure of cycles scaled to the triangulated reference models significantly decreased with colony size in all species subsets (16/16 subsets; Figure 2F; OLS of PIC: β = -0.048, 95% CI = -0.102, -0.015, R^2 = 0.090, p = 0.186; Tables S13).

Figure 2: **Nest Connectivity: branching (A-C) and short-cut (D-F) hypotheses.** (A, D)

Relationship between colony size (average number of ants in a colony) and (A) network mean

distance and (D) mean number of cycles in nests. Each data point represents a mean value for a

single species, whiskers indicate standard errors around the mean when multiple nests were

sampled for a given species, colors represent ant subfamily (see legend in A). Dashed line is the

regression fit for PGLS when λ=1, solid when statistically significant and dashed when not

statistically significant; (B, E) The mean distance (B) or number of cycles (E) according to

network size (number of nodes). Networks from observed nests are plotted as grey dots, sized

according to sample size. In (B) lines are mean distances as a function of number of nodes for

chain (red), MST (yellow) and triangulated (blue) reference networks, with 95% confidence

intervals from 1000 simulation runs shown as shading around the lines; in (E) Blue line is the

number of cycles in triangulated networks as a function of number of nodes. Orange line depicts

the number of cycles for both MSTs and chain networks, which contain zero cycles at all

network sizes; (C, F) Relationship between colony size (average number of ants in a colony) and

scaled mean distance (C) and scaled number of cycles (F). Mean distance (C) is scaled to chain

(red), MST (yellow) and triangulated (blue) networks. Cycles (F) are only scaled to triangulated

networks (blue). Each point represents a mean of scaled values from the nests of each species.

Grey dashed line at y=1 represents values at which observed nests contain the same value as

expected from the reference model. Values below the dashed line indicate networks that have

lower values than expected by the reference model. Values above the dashed line indicate

networks which have values greater than expected according to the reference model. Solid

colored lines indicate statistically significant regression fits for PGLS (λ=1).

DISCUSSION

Our results support the hypothesis that nest architecture mitigates the challenges of larger colony

sizes, i.e., more individuals in a colony. Our comparative analysis found that nests of species

with larger colonies are more subdivided and reduce travel distances through increased

branching. Overall, we did not find a strong effect of phylogeny.

As predicted by the subdivision hypothesis, the number of chambers was greater in species

with larger colonies and chamber size was not related to colony size (Figure 1). These results

suggest that nests of species with large colonies are divided into more sub-units than nests of

species with small colonies, and that the size of those sub-units does not increase proportionately

to colony size. Similar patterns of increased spatial subdivision with body size are the

cornerstone of multi-cellularity (Bonner 1998), and also characterize subdivision as the brain

(Kaas 2000) and cities (Anas et al. 1998) increase in size. Given our correlational findings, it

remains unclear if subdivision in nests serves a similar function as it does in these other systems.

Further data are needed on the distribution of individuals throughout nest chambers to determine

if ants utilize chambers in a way that is consistent with subdivision (e.g. Tschinkel 1999;

Murdock and Tschinkel 2015). Nest subdivision may facilitate division of labor through spatial

segregation of tasks (Tofts and Franks 1992; Richardson et al. 2011; Mersch et al. 2013), and

slow down the spread of disease (Pie et al. 2004; Stroeymeyt et al. 2014, 2018). However, the

only direct test of the benefit of nest subdivision showed that subdivision does not increase brood

rearing efficiency (Cassill et al. 2002; Tschinkel 2018). Limits on chamber size may also be

imposed by physiological processes, such as gas exchange or temperature regulation (Jones and

Oldroyd 2006), or by structural constraints, because large chambers may collapse. Thus, the

causes and consequences of nest sub-division remain to be tested.

The connectivity of nests changed according to colony size, but only one of our hypotheses

about nest connectivity was supported. The nests of species with larger colonies had reduced

travel distances due to their branching structure, consistent with the branching hypothesis. In the

nests of species with small colonies (i.e., up to ~1000 workers in nests containing up to ~10

nodes), mean distances tended to match the values expected for chain networks, and indeed the

nests of many small colony species were chains. In the nests of species with large colonies, mean

travel distances were shorter than would be expected for chain networks but greater than

expected for MSTs, and far greater than expected for triangulated networks (Figure 2B, C).

These findings indicate that the nests of species with large colonies have increased branching

(i.e., are more tree-like) relative to the nests of species with smaller colonies. By increasing the

frequency of branching large colonies may curb rising travel distances as their nests increase in

size (i.e., contain more chambers). Networks cannot simultaneously minimize travel distances

(i.e., be highly interconnected) and minimize their number of edges (i.e., minimize infrastructure

costs). Increased branching is a common way for biological transport networks to balance this

trade-off (e.g., ant foraging trails: Buhl et al. 2009; slime molds: Tero et al. 2010; neural

networks: Raj and Chen 2011; plant growth forms: Conn et al. 2017), especially as body size, or

population size, increases (Ringo 1991; Banavar et al. 1999; Ercsey-Ravasz et al. 2013; Cabanes

et al. 2015). Branching is thus a low-cost way for biological transport networks, including ant

nests, to maintain connectivity as they increase in size.

The short-cuts hypothesis, that nests from species with larger colonies will have more cross-

linking paths or cycles, was not supported. Although the number of cycles in a nest increased

with colony size, this increase was not statistically significant (Figure 2D). Cycles were added at

a slower rate than expected for a triangulated network (Figure 2E), which is not surprising given

that triangulations represent the maximum rate at which new cycles could be added with

increasing network size (for 2D networks). This finding lends further support to the conclusion

that the increase in connectivity with colony size is due to increased branching and not the

addition of short-cuts. However, we caution that our analysis conservatively excludes nests that

might have the most cycles because we could not accurately infer the topology of the most

convoluted nests. Our inability to detect a significant increase in the number of cycles with

colony size could be due to this bias in our dataset. More refined imaging techniques could

provide a solution to this issue and expand future work on nest architecture in general.

The scarcity of cycles in the nests we measured suggest that the benefits of having

alternative access routes, which help maintain movement (Corson 2010; Katifori et al. 2010;

Blonder et al. 2018, Jyothi et al. 2016) and facilitate forager recruitment (Pinter-Wollman 2015),

might not outweigh the potential costs of additional tunnels. For instance, there are costs to

building and maintaining tunnels (da Silva Camargo et al. 2013; Monaenkova et al. 2015) and

there may be negative consequences of many cross-linking paths, including structural instability

or heightened exposure to risks, like pathogens (Scovell 1983; Viana et al. 2013; Stroeymeyt et

al. 2014, 2018), parasites (Scovell 1983), or predators (Jeanne 1975; LaPolla et al. 2002). Nests

are multi-purpose structures that have other functions, like thermoregulation, humidity control or

ventilation (Jones and Oldroyd 2006; King et al. 2015), which might suffer when nests have

many cycles. In systems where the costs of cross-linking paths are low, like the plasmodia of

slime molds, the mycelia of fungus, or the pheromone trails of ants, cycles are ephemeral and

created for exploration, yet they are pruned once resources are discovered (Bebber et al. 2007;

Buhl et al. 2009; Tero et al. 2010; Latty et al. 2011). Conversely, when the benefits to movement

are very high, as in some man-made networks like urban roads or computer routing networks,

cycles become more common as size increases (Levinson 2012; Jyothi et al. 2016). Future

investigation into the building costs of tunnels and the functional consequences of interconnected

nest structures will help reveal why ant nests do not contain more short-cuts, or cycles.

As with any meta-analysis, our dataset is constrained by the available literature. Many of

the nests we found in the literature were built in sandy soils, and so our analysis has an over-

representation of species adapted to the climate and ecology of sandy soils (Table S1). Fungus

growers are highly-represented in our analysis, and constraints imposed by requirements to

maintain healthy fungus gardens could have influenced our findings. Additionally, because we

do not have data on the size or age of the colonies that built the majority of nests in our meta-

analysis, we might have included immature colonies that have not reached their full size. Further

work could investigate whether nest connectivity changes throughout development within a

species in natural environments (e.g., Tschinkel 1987, 2004, 2005, 2011) or whether there are

species-specific connectivity patterns that can be found at different ages.

In conclusion, we show that ant nests exhibit modular and efficient structures in species

with larger colonies. The similarity in how ant nests and the bodies of unitary organisms respond

to increasing size suggest that their shared architectural features, branching transportation

networks and compartmentalization, provide common solutions to the challenges created by

large size across the tree of life.

Acknowledgements
We are thankful to Doug Booher for sharing his nest castings and to Walter Tschinkel for
additional images of his published castings. We thank Dana Williams, Felipe Zapata, Kevin
Loope, Tyler McCraney and the UCLA IDRE statistical consulting unit for advice on the
statistical analysis and to Saray Shai for advice on generating reference models. We thank Gabby
Najm, James Lichtenstein, Sean O'Fallon, Natalie Lemanski and two anonymous reviewers for
helpful comments on the manuscript.
427
428
429

REFERENCES

Aguilar, J., D. Monaenkova, V. Linevich, W. Savoie, B. Dutta, H.-S. Kuan, M. D. Betterton, M.

A. D. Goodisman, and D. I. Goldman. 2018. Collective clog control: Optimizing traffic

flow in confined biological and robophysical excavation. Science (80-.). 361:672–677.

Anas, A., R. Arnott, and K. A. Small. 1998. Urban spatial structure. Am. Econ. Assoc. 36:1426–

1464.

Banavar, J. R., A. Maritan, and A. Rinaldo. 1999. Size and form in efficient transportation

networks. Nature 399:130–132.

Bebber, D. P., J. Hynes, P. R. Darrah, L. Boddy, and M. D. Fricker. 2007. Biological solutions to

transport network design. Proc. R. Soc. B Biol. Sci. 274:2307–2315.

Blanchard, B. D., and C. S. Moreau. 2017. Defensive traits exhibit an evolutionary trade-off and

drive diversification in ants. Evolution (N. Y). 71:315–328.

Blonder, B., N. Salinas, L. P. Bentley, A. Shenkin, P. O. Chambi Porroa, Y. Valdez Tejeira, T. E.

Boza Espinoza, G. R. Goldsmith, L. Enrico, R. Martin, G. P. Asner, S. Díaz, B. J. Enquist,

and Y. Malhi. 2018. Structural and defensive roles of angiosperm leaf venation network

reticulation across an Andes–Amazon elevation gradient. J. Ecol. 106:1683–1699.

Bonner, J. T. 2004. Perspective: The Size-Complexity Rule. Evolution (N. Y). 58:1883–1890.

Bonner, J. T. 1988. The Evolution of Complexity by Means of Natural Selection. Princeton

University Press, Princeton, NJ.

Bonner, J. T. 1998. The origins of multicellularity. Integr. Biol. 1:27–36.

Bourke, A. F., and N. R. Franks. 1995. Social Evolution in Ants. Princeton University Press,

Princeton, NJ.

Brian, M. V. 1956. Group Form and Causes of Working Inefficiency in the Ant Myrmica rubra

L. Physiol. Zool. 29:173–194.

Brown, J. J., and J. F. A. Traniello. 1998. Regulation of Brood-Care Behavior in the Dimorphic

Castes of the Ant Pheidole morrisi (Hymenoptera□: Formicidae): Effects of Caste Ratio ,

Colony Size , and Colony Needs. J. Insect Behav. 11:209–219.

Brown, W. L. 1953. Revisionary Studies in the Ant Tribe Dacetini. Am. Midl. Nat. 50:1.

Buhl, J., J. Gautrais, J.-L. Deneubourg, and G. Theraulaz. 2004a. Nest excavation in ants□:

group size effects on the size and structure of tunneling networks. Naturwissenschaften

91:602–606.

Buhl, J., J. Gautrais, R. V Sole, P. Kuntz, S. Valverde, J.-L. Deneubourg, and G. Theraulaz.

2004b. Efficiency and robustness of ant networks of galleries. Eur. Phys. J. B 129:123–129.

Buhl, J., K. Hicks, E. R. Miller, S. Persey, O. Alinvi, and D. J. T. Sumpter. 2009. Shape and

efficiency of wood ant foraging networks. Behav. Ecol. Sociobiol. 63:451–460.

Bullmore, E., and O. Sporns. 2012. The economy of brain network organization. Nat. Rev.

Neurosci. 13:336–349.

Burd, M., N. Shiwakoti, M. Sarvi, and G. Rose. 2010. Nest architecture and traffic flow □: large

potential effects from small structural features. Ecol. Entomol. 35:464–468.

Cabanes, G., E. Van Wilgenburg, M. Beekman, and T. Latty. 2015. Ants build transportation

networks that optimize cost and efficiency at the expense of robustness. Behav. Ecol.

26:223–231.

Cassill, D., W. R. Tschinkel, and S. B. Vinson. 2002. Nest complexity , group size and brood

rearing in the fire ant , Solenopsis invicta. Insectes Soc. 49:158–163.

Conn, A., U. V. Pedmale, J. Chory, and S. Navlakha. 2017. High-Resolution Laser Scanning

Reveals Plant Architectures that Reflect Universal Network Design Principles. Cell Syst.

5:53-62.e3. Elsevier Inc.

Corson, F. 2010. Fluctuations and Redundancy in Optimal Transport Networks. 048703:1–4.

Csárdi, G. 2020. Package ' igraph ': Network Analysis and Visualization.

da Silva Camargo, R., J. F. S. Lopes, and L. C. Forti. 2013. Energetic cost of digging behavior in

workers of the leaf-cutting ant Atta sexdens (Fabricius). Rev. Bras. Entomol. 57:401–404.

Dodds, P. S. 2010. Optimal form of branching supply and collection networks. Phys. Rev. Lett.

482  104:1–4.

483 Dornhaus, A., S. Powell, and S. Bengston. 2012. Group size and its effects on collective

484  organization. Annu. Rev. Entomol. 57:123–41.

485 Ercsey-Ravasz, M., N. T. Markov, C. Lamy, D. C. VanEssen, K. Knoblauch, Z. Toroczkai, and

486  H. Kennedy. 2013. A Predictive Network Model of Cerebral Cortical Connectivity Based

487  on a Distance Rule. Neuron 80:184–197.

488 Freckleton, R. P. 2009. The seven deadly sins of comparative analysis. J. Evol. Biol. 22:1367–

489  1375.

490 Gillooly, J. F., C. Hou, and M. Kaspari. 2010. Eusocial insects as superorganisms: Insights from

491  metabolic theory. Commun. Integr. Biol. 3:360–362.

492 Gould, S. J. 1966. Allometry and size in ontogeny and phylogeny. Biol. Rev. Camb. Philos. Soc.

493  41:587–640.

494 Hobson, E. A., M. J. Silk, N. H. Fefferman, D. B. Larremore, P. Rombach, S. Shai, and N.

495  Pinter-Wollman. 2020. A guide to choosing and implementing reference models for social

496  network analysis. arXiv.

497 Holldobler, B., and E. O. Wilson. 2009. The superorganism: the beauty, elegance, and

498  strangeness of insect societies Title. W.W. Norton, New York.

499 Jeanne, R. L. 1975. The Adaptiveness of Social Wasp Nest Architecture. Q. Rev. Biol. 50:267–

500  287.

501 Jones, J. C., and B. P. Oldroyd. 2006. Nest Thermoregulation in Social Insects. Adv. In Insect

502  Phys. 33:153–191.

Jyothi, S. A., A. Singla, P. B. Godfrey, and A. Kolla. 2016. Measuring and Understanding

Throughput of Network Topologies. Int. Conf. High Perform. Comput. Networking, Storage

Anal. SC 0:761–772.

Kaas, J. H. 2000. Why is Brain Size so Important:Design Problems and Solutions as Neocortex

Gets Biggeror Smaller. Brain Mind 1:7–23.

Katifori, E., G. J. Szöllosi, and M. O. Magnasco. 2010. Damage and fluctuations induce loops in

optimal transport networks. Phys. Rev. Lett. 104:1–4.

Kempes, C. P., S. Dutkiewicz, and M. J. Follows. 2012. Growth , metabolic partitioning , and the

size of microorganisms. Proc. Natl. Acad. Sci. 109:495–500.

King, H., S. A. Ocko, and L. Mahadevan. 2015. Termite mounds harness diurnal temperature

oscillations for ventilation. Proc. Natl. Acad. Sci. 112:11589–11593.

Kwapich, C. L., G. Valentini, and B. Hölldobler. 2018. The non-additive effects of body size on

nest architecture in a polymorphic ant. Philos. Trans. R. Soc. B 373:20170235.

LaPolla, J. S., U. G. Mueller, M. Seid, and S. P. Cover. 2002. Predation by the army ant

Neivamyrmex rugulosus on the fungus-growing ant Trachymyrmex arizonensis. Insectes

Soc. 49:251–256.

Latty, T., K. Ramsch, K. Ito, T. Nakagaki, D. J. T. Sumpter, M. Middendorf, and M. Beekman.

2011. Structure and formation of ant transportation networks. J. R. Soc. Interface 8:1298–

1306.

Levinson, D. 2012. Network structure and city size. PLoS One 7:e29721.

Lorenz, D. M., A. Jeng, and M. W. Deem. 2011. The emergence of modularity in biological

systems. Phys Life Rev. 8:129–160.

McKown, A. D., H. Cochard, and L. Sack. 2010. Decoding leaf hydraulics with a spatially

explicit model: Principles of venation architecture and implications for its evolution. Am.

Nat. 175:447–460.

McMahon, T. A., and J. T. Bonner. 1983. On Size and Life. Scientific American Books, Inc,

New York.

Mersch, D. P., A. Crespi, and L. Keller. 2013. Tracking individuals shows spatial fidelity is a

key regulator of ant social organization. Science (80-.). 340:1090–3.

Monaenkova, D., N. Gravish, G. Rodriguez, R. Kutner, M. A. D. Goodisman, and D. I.

Goldman. 2015. Behavioral and mechanical determinants of collective subsurface nest

excavation. J. Exp. Biol. 218:1295–1305.

Murdock, T. C., and W. R. Tschinkel. 2015. The life history and seasonal cycle of the ant,

Pheidole morrisi Forel, as revealed by wax casting. Insectes Soc. 62:265–280.

Newman, M. E. J. 2018. Networks: An Introduction. 2nd Editio. Oxford University Press,

Oxford.

Niklas, K. J., and S. A. Newman (eds). 2016. Multicellularity: Origins and Evolution. MIT Press,

Cambridge, MA.

O'Donnell, S., and S. J. Bulova. 2007. Worker connectivity: A review of the design of worker

communication systems and their effects on task performance in insect societies. Insectes

Soc. 54:203–210.

Okie, J. G. 2013. General models for the spectra of surface area scaling strategies of cells and

organisms: Fractality, geometric dissimilitude, and internalization. Am. Nat. 181:421–439.

Orme, A. D., R. Freckleton, G. Thomas, T. Petzoldt, N. Isaac, W. Pearse, and M. D. Orme. 2018.

Package ' caper '□: Comparative Analyses of Phylogenetics and Evolution in R.

Pacala, S. W., D. M. Gordon, and H. C. J. Godfray. 1996. Effects of social group size on

information transfer and task allocation. Evol. Ecol. 10:127–165.

Pagel, M. 1999. Inferring historical patterns of biological evolution. Nature 401:877–884.

Paradis, E., S. Blomberg, B. Bolker, J. Brown, J. Claude, H. S. Cuong, R. Desper, G. Didier, B.

Durand, J. Dutheil, R. Ewing, O. Gascuel, T. Guillerme, C. Heibl, A. Ives, B. Jones, F.

Krah, D. Lawson, V. Lefort, P. Legendre, J. Lemon, E. Marcon, R. McCloskey, J.

Nylander, R. Opgen-Rhein, A.-A. Popescu, M. Royer-Carenzi, K. Schliep, K. Strimmer,

and D. de Vienne. 2018. Package 'ape': Analyses of Phylogenetics and Evolution.

Perna, A., C. Jost, E. Couturier, S. Valverde, S. Douady, and G. Theraulaz. 2008. The structure

of gallery networks in the nests of termite Cubitermes spp. revealed by X-ray tomography.

Naturwissenschaften 95:877–884.

Perna, A., and T. Latty. 2014. Animal transportation networks. J. R. Soc. Interface 11.

Pie, M. R., R. B. Rosengaus, and J. F. A. Traniello. 2004. Nest architecture , activity pattern ,

worker density and the dynamics of disease transmission in social insects. J. Theor. Biol.

226:45–51.

Pinter-Wollman, N. 2015a. Nest architecture shapes the collective behaviour of harvester ants.

Biol. Lett. 11:20150695-.

Pinter-Wollman, N. 2015b. Persistent variation in spatial behavior affects the structure and

function of interaction networks. Curr. Zool. 61:98–106.

Pinter-Wollman, N., A. Bala, A. Merrell, J. Queirolo, M. C. Stumpe, S. Holmes, and D. M. Gordon. 2013. Harvester ants use interactions to regulate forager activation and availability. Anim. Behav. 86:197–207. Elsevier Ltd.

Pinter-wollman, N., A. Penn, G. Theraulaz, and S. M. Fiore. 2018. Interdisciplinary approaches for uncovering the impacts of architecture on collective behaviour. Philos. Trans. R. Soc. B 373:20170232.

R Core Team. 2019. R: A language environment for statistical computing.

Raj, A., and Y. H. Chen. 2011. The wiring economy principle: Connectivity determines anatomy in the human brain. PLoS One 6.

Richardson, T. O., K. Christensen, N. R. Franks, H. J. Jensen, and A. B. Sendova-franks. 2011. Ants in a Labyrinth□: A Statistical Mechanics Approach to the Division of Labour. PLoS One 6:e18416.

Ringo, J. L. 1991. Neuronal interconnection as a function of brain size. Brain. Behav. Evol. 38:1–6.

Rueden, C. T., J. Schindelin, M. C. Hiner, B. E. Dezonia, A. E. Walter, E. T. Arena, and K. W. Eliceiri. 2017. ImageJ2□: ImageJ for the next generation of scientific image data. BMC Bioinformatics 18:1–26.

Rupppert, E. E., R. S. Fox, and R. D. Barnes. 2004. Invertebrate Zoology: A Functional Evolutionary Approach. Thomson-Brooks/Cole.

Scovell, E. 1983. Dominance and reproductive success among slave-making worker ants. Nature

304:724–725.

Seeley, T. D. 1989. The honey bee colony as a superorganism. Am. Sci. 77:546–553.

Stroeymeyt, N., B. Casillas-Pérez, and S. Cremer. 2014. Organisational immunity in social insects. Curr. Opin. Insect Sci. 5:1–15.

Stroeymeyt, N., A. V Grasse, A. Crespi, D. P. Mersch, S. Cremer, and L. Keller. 2018. Social network plasticity decreases disease transmission in a eusocial insect. Science (80-.). 362:941–945.

Tekin, E., D. Hunt, M. G. Newberry, and V. M. Savage. 2016. Do Vascular Networks Branch Optimally or Randomly across Spatial Scales□? PLoS Comput. Biol. 12:e1005223.

Tero, A., S. Takagi, T. Saigusa, K. Ito, D. P. Bebber, M. D. Fricker, K. Yumiki, R. Kobayashi, and T. Nakagaki. 2010. Rules for biologically inspired adaptive network design. Science (80-.). 327:439–442.

Tofts, C., and N. R. Franks. 1992. Doing the right thing: Ants, honeybees and naked mole-rats. Trends Ecol. Evol. 7:346–349.

Tschinkel, W. R. 2010. Methods for casting subterranean ant nests. J. Insect Sci. 88:1–17.

Tschinkel, W. R. 1987. Seasonal life history and nest architecture of a winter-active ant, Prenolepis imparis. Insectes Soc. 34:143–164.

Tschinkel, W. R. 1999. Sociometry and sociogenesis of colonies of the harvester ant, Pogonomyrmex badius: distribution of workers, brood and seeds within the nest in relation to colony size and season. Ecol. Entomol. 24:222–237.

Tschinkel, W. R. 2018. Testing the effect of a nest architectural feature in the fire ant Solenopsis

invicta (Hymenoptera: Formicidae). Myrmecological News 27:1–5.

Tschinkel, W. R. 2015. The architecture of subterranean ant nests: beauty and mystery underfoot.

J. Bioeconomics 17:271–291.

Tschinkel, W. R. 2005. The nest architecture of the, ant Camponotus socius. J. Insect Sci. 5:1–

18.

Tschinkel, W. R. 2004. The nest architecture of the Florida harvester ant , Pogonomyrmex

badius. J. Insect Sci. 21:1–19.

Tschinkel, W. R. 2011. The nest architecture of three species of north Florida Aphaenogaster

ants. J. insect Sci. 11:1–30.

Turner, R. 2020. Package ' deldir ': Delaunay Triangulation and Dirichlet (Voronoi) Tessellation

Author.

Vaes, O., A. Perna, and C. Detrain. 2020. The effect of nest topology on spatial organization and

recruitment in the red ant Myrmica rubra. Sci. Nat. 107. The Science of Nature.

Viana, M. P., V. Fourcassié, A. Perna, F. Costa, and C. Jost. 2013. Accessibility in networks :

A useful measure for understanding social insect nest architecture. Chaos, Solitons and

Fractals 46:38–45. Elsevier Ltd.

West G. B., Brown J. H., and Enquist B. J. 1997. A General Model for the Origin of Allometric

Scaling Laws in Biology. Science (80-.). 276:122–126.

Wheeler, W. M. 1910. Ants: Their structure, Development and Behavior. Columbia University

Press, New York.

628