1 TITLE
2
Left Hemisphere Dominance for Bilateral Kinematic Encoding in the Human Brain

AUTHORS:

Christina M. Merrick[1*], Tanner C. Dixon[2], Assaf Breska[1], Jack J. Lin[3], Edward F., Chang[4], David
King-Stephens[5], Kenneth D. Laxer[5], Peter B. Weber[5], Jose M. Carmena[2,6,7], Robert T. Knight[1,2,4,7],
Richard B. Ivry[1,2,7]

AFFILIATIONS:

[1] Department of Psychology, University of California, Berkeley, CA, USA

[2]UC Berkeley – UCSF Graduate Program in Bioengineering

[3]Department of Neurology, University of California at Irvine, Irvine, CA, USA.

[4]Department of Neurological Surgery, University of California San Francisco, San Francisco, CA,
USA.

[5]Department of Neurology and Neurosurgery, California Pacific Medical Center, San Francisco,
CA, USA.

[6]Department of Electrical Engineering and Computer Sciences, University of California, Berkeley,
CA, USA

[7]Helen Wills Neuroscience Institute, University of California, Berkeley, CA, USA

*Correspondence: cmerrick@berkeley.edu

KEYWORDS: lateralization; left hemisphere; motor control; reaching movements; ipsilateral;
praxis; electrocorticography

ABSTRACT (149 words)

Neurophysiological studies in humans and non-human primates have revealed movement
representations in both the contralateral and ipsilateral hemisphere. Inspired by clinical
observations, we ask if this bilateral representation differs for the left and right hemispheres.
Electrocorticography (ECoG) was recorded in human participants during an instructed-delay
reaching task, with movements produced with either the contralateral or ipsilateral arm. Using a
cross-validated kinematic encoding model, we found stronger bilateral encoding in the left
hemisphere, an effect that was present during preparation and was amplified during execution.
Consistent with this asymmetry, we also observed better across-arm generalization in the left
hemisphere, indicating similar neural representations for right and left arm movements. Notably,
these left hemisphere electrodes were centered over premotor and parietal regions. The more
extensive bilateral encoding in the left hemisphere adds a new perspective to the pervasive
neuropsychological finding that the left hemisphere plays a dominant role in praxis.

INTRODUCTION

A primary tenet of neurology is the contralateral organization of movement. The vast majority of the fibers from the corticospinal tract cross to the opposite side of the body (Nyberg-Hansen & Rinvik, 1963) and functionally, hemiparesis resulting from cortical stroke is manifest on the contralateral side of the body (Bourbonnais, & Noven, 1989). Although direct control of arm movements is primarily mediated through contralateral projections, unimanual arm movements elicit bilateral activity in the primary motor cortex (M1, Babiloni et al., 1999; Ghacibeh et al., 2007), indicating that neural activity in the ipsilateral hemisphere contains information relevant to ongoing movement. Correspondingly, kinematic and movement parameters of the ipsilateral limb can be decoded from ipsilateral hemisphere intracortical recordings in monkeys (Ganguly et al., 2009; Ames & Churchland, 2019) and from electrocorticography (ECoG) in humans (Bundy, Szrama, Pahwa & Leuthardt, 2018; Ganguly et al., 2009, Wisneski et al., 2008). Ipsilateral signals represent an intriguing source of neural activity, both for understanding how activity across the two hemispheres results in coordinated movement and because this information might be exploited for rehabilitative purposes.

While it is established that information about unimanual movements is contained within the ipsilateral hemisphere, there remains considerable debate about what this signal represents. Previous studies have centered on the question of whether ipsilateral representations overlap or are independent of contralateral representations, leading to mixed results. Consistent with the overlap hypothesis, neural activity for the contralateral and ipsilateral limb movements show several similarities, including shared target tuning preferences and the ability to cross predict kinematic features from a model trained on the opposite arm (Bundy, Szrama, Pahwa & Leuthardt, 2018; Cisek, Crammond & Kalaska, 2003; Steinberg et al., 2002, Willett et al., 2020). Consistent with the independence hypothesis, intracortical recordings in monkeys have revealed that the lower dimensional representations of the two arms lie in orthogonal subspaces (Ames, Churchland, 2019; Heming, Cross, Takei, Cook & Scott, 2019). These hypotheses are not mutually exclusive. For example, the degree of overlap or independence may depend on the gesture type (e.g., overlapping representations for grasping but not arm movement, Downey et a., 2020), or brain region (e.g., premotor cortex displays stronger preservation of tuning preferences across the two arms than primary motor cortex, Cisek, Crammond & Kalaska, 2003).

One factor that has received little attention in this literature is the recording hemisphere. This is surprising given the marked asymmetries between the two hemispheres in terms of praxis (Corballis, Badzakova-Trajkov & Häberling, 2012; Rothi, Ochipa & Heilman, 1997). Tracing back to the early 20[th] century, marked hemispheric asymmetries have been defined by the behavioral deficits observed following unilateral brain injury (Schaefer, Haaland & Sainburg, 2007; Liepmann 1908, cited in Renzi, & Lucchelli, 1988). Apraxia, an impairment in the production of coordinated, meaningful movement in the absence of muscle recruitment deficits, is much more common after left compared to right hemisphere insult (Haaland, Harrington & Knight, 2000; Renzi, & Lucchelli, 1988). Moreover, left hemisphere stroke will frequently result in apraxic symptoms for gestures produced with either hand, as well as impairments in action comprehension (Renzi, & Lucchelli, 1988). Hemispheric asymmetries are also evident in neuroimaging activation patterns in healthy participants, with the left hemisphere having stronger activation during ipsilateral movement than the right hemisphere, especially with increasing task difficulty (Chettouf et al., 2020, Verstynen et al. 2005; Verstynen and Ivry 2011; Schäfer et al, 2007). These patterns raise the possibility that the ipsilateral cortical representation differs between the left and right hemispheres.

In the present study, we use intracranial recordings from the cortical surface (ECoG) to examine the degree of cortical overlap for ipsilateral and contralateral upper limb movement in the left and the right hemisphere. The data were collected from six patients, three with left hemisphere implants and three with right hemisphere implants, while they engaged in an instructed-delay reaching task. We focus on predicting the temporal dynamics of high frequency activity (HFA; 70-200Hz), a surrogate for infragranular single-unit activity and supragranular dendritic potentials, (Leszczynski et al., 2020) which tracks local activation of the cortex (Muthukumaraswamy, 2010). Going beyond previous studies that use decoding models which combine multiple neural features from multiple electrodes to predict kinematics, we employed an encoding model which uses kinematic features to predict neural activity for each electrode, allowing us to retain the high spatial and temporal resolution of the ECoG signal. This approach allows us to create high-resolution topographic maps depicting encoding strength on the surface of the cortex for movements produced with the contralateral and ipsilateral arm. This is preferable to projecting the weights obtained from decoding models since these models have difficulty disambiguating between informative and uninformative electrodes (Kriegeskorte & Douglas, 2019). Moreover, our approach provides a way to map kinematics to neural activity in a time-resolved manner (rather than as single weights), allowing us to identify time ranges of representational overlap and divergence across the two arms for each electrode.

RESULTS

Behavior. We used a delayed response, out-and-back reaching task. On each trial, a cue indicating the target location was presented on a touchscreen followed, after a short delay by an imperative signal. Participants were instructed to prepare to move to the target during the delay period. The participant was free to move at their own pace, with the instructions emphasizing that the participant should focus on touching the screen near the target and then returning to the start position. Left and right arm reaches were tested in separate blocks, with the position of the non-responding hand fixed throughout the block.

Table 1 summarizes the total number of successful trials, along with the reaction time and movement time data. A trial was considered unsuccessful if the reach was initiated before the go cue or if contact with the touchscreen was outside the boundary of the target. The percentage of unsuccessful trials was low, ranging between 0% to 12.5% across individuals. The movements had roughly, bell-shaped velocity profiles for the outbound and the inbound segments (Fig 1C,1E) and the outbound reaches were, on average, faster than inbound reaches. The marked inter-individual differences in reaction time and movement time reflect the fact that the instructions emphasized accuracy and smoothness.

Patient ID	Age	Handedness	RT	Outbound reach	Inbound reach	Total trials
L1	36	Right	392 (102)	703 (118)	907 (287)	152
L2	26	Right	1574 (871)	598 (173)	1003 (406)	146
L3	55	Ambidextrous	771 (345)	946 (183)	1429 (252)	148
R1	26	Right	518 (194)	940 (211)	1027 (345)	132
R2	32	Right	335 (55)	602 (94)	590 (120)	145
R3	34	Right	534 (157)	721 (145)	1017 (318)	145

Table 1. Summary of performance measures for each participant. Reaction times are averages
over left and right arm reaches since RT differences were negligible for the two arms.

Figure 1. Task and model design. A. Task overview. Participants performed an instructed-delay reaching
task, moving to targets that appeared on a touchscreen monitor with either the left or right arm. **B.**
Target positions. The position of each target with respect to the start position of the reaching arm can be
defined on the basis of three Cartesian coordinates (X, Y, Z) and two spherical angles (Theta and Phi). **C.**
Representative kinematics. Average estimated position and velocity traces for a representative series of

trials performed with the left arm. **D. Reach accuracy**. Accuracy was quantified as the absolute distance
from the center of each target (target diameter = 2.5cm) to the touch location for all four targets with the
left (blue) and right (grey) arm. **E. Kinematic encoding model**. Time lagged estimated kinematic features
were used to the predict high frequency activity (HFA) for each electrode using ridge regression. Four
kinematic features were included in the model: Position in the Z dimension, speed in the Z dimension and
the two spherical angles phi and theta. Kinematic features were trained on a subset of the HFA data and
predictions of HFA activity were evaluated on held-out test sets.

At a more fine-grained level of spatial accuracy, we calculated the distance from the center of
each target to the touch location for each trial. On average, the mean distance from the center
of the 2.5 cm circle was 0.80 cm (*SD* = 0.10 cm) for right-handed reaches and 0.90 cm (*SD* = 0.17
cm) for left-handed reaches (Fig 1D). These values did not differ from one another [*t* = 1.538, *p* =
.222].

Stronger bilateral encoding in the left hemisphere. We examined the extent to which movement
kinematics were encoded for contralateral and ipsilateral reaches in individual electrodes. To do
this we fit a kinematic encoding model that maps continuous kinematic features to the HFA
signal (Fig 1F) for the 665 electrodes meeting our inclusion criteria. This procedure was done
separately for contralateral and ipsilateral reaches. We quantified the cross-validated model fit
by generating HFA predictions using the kinematic features from held-out trials of the same
condition and calculating prediction performance as the square of the linear correlation (R^2)
between the predicted and actual HFA signal (Fig 2B).

Figure 2. Stronger bilateral encoding in left hemisphere. Held-out prediction performance (R^2) was computed for each electrode during contralateral reaches (C) and ipsilateral reaches (I). R^2 was calculated as the squared linear correlation between the actual HFA and the predictions based on the model. **A. Prediction performance maps for individual patients.** Performance of each electrode, shown at the idiosyncratic electrode location for each participant (location based on clinical criteria). Electrodes that did not account for at least .05% of the variance ($R^2 < .05$) in either the contralateral or ipsilateral condition are shown as smaller dots. **B. Model predictions.** Representative time series of the actual HFA and model-based predictions for three electrodes during contralateral and ipsilateral reaches. **C. Summary across patients.** Scatter plot displaying R^2 values separately for patients with electrodes in either the left (upper) or right (lower) hemisphere. Individual patients can be identified in the scatterplot by the color surrounding the patient label (e.g., L1) in A. R^2 for contralateral predictions are plotted against R^2 for ipsilateral predictions. Electrodes close to the unity line encode both arms equally whereas electrodes off the unity line indicate stronger encoding of one arm. Points above the unity line indicate stronger encoding of the contralateral arm. These differences are summarized in the frequency histograms in the upper right of each panel. The histogram shows less of a shift in the left hemisphere, a signature consistent with stronger bilateral encoding. *p<0.05, **p<0.01, ***p<0.001, permutation test.

Figure 2A displays R^2 values for each electrode for the contralateral and ipsilateral condition, presented on the individual patient MRIs. Electrodes with high prediction performance were primarily located in arm areas of sensorimotor cortex. In line with previous research (Downey et a., 2020), a sizeable percentage of the electrodes were able to predict the HFA at or above our criterion of $R^2 > .05$ (examples shown in Figure 2B). This degree of prediction was observed not only when the data were restricted to contralateral movement (31% of electrodes), but also when the data were from ipsilateral movement (25%). A number of electrodes (24%) were predictive in both the contralateral and ipsilateral models. Electrodes that did not meet this criterion for either arm are represented as small dots in figure 2A and were excluded from further analysis, leaving a total of 216 predictive electrodes (32%, 141 = left hemisphere, 75 = right hemisphere).

Before comparing prediction performance across the two hemispheres, we first evaluated the distribution of the electrodes in the right and left hemispheres. Electrode placement for ECoG is determined solely for clinical purposes; as such, hemispheric asymmetries could arise from differences in coverage rather than functional differences. To evaluate coverage, we categorized the position of all electrodes based on a cortical parcellation (Desikan et. al., 2006; Figure 2- figure supplement 1). When the categorization data were pooled across the three left hemisphere and the three right hemisphere patients, the proportion of electrodes was similar across the eight parcellations that encompass premotor, sensorimotor motor, and parietal regions [$averages: premotor_{right} = 11\%,\ premotor_{left} = 10\%,\ sensorimotor_{right} = 17\%, sensorimotor_{right} = 16\%, parietal_{right} = 5\%, parietal_{left} = 3\%;\ \chi^2(7) = 0.057, p = .999$; Figure 2- figure supplement 1].

Given the similar distributions, we next asked whether prediction performance for the two arms differed across the two hemispheres. Figure 2C compares the predictive performance of each electrode for the contralateral and ipsilateral conditions for patients with left or right hemisphere grids. Values close to the unity line yield similar predictions for the two conditions; values off the unity line indicate that encoding is stronger for one arm compared to the other. To compare prediction performance at the group level, we fit a permutation-based mixed effects model with fixed factors of *Arm* and *Hemisphere* and a random factor of *Participant.* We found a main effect of *Arm* with contralateral reaches being encoded more strongly than ipsilateral

reaches [$\chi^2(1)$ = 29.34, p < .001]. We found no effect of *Hemisphere* [$\chi^2(1)$ = 0.46, p > .50], but
we found a significant interaction between *Arm* and *Hemisphere* [$\chi^2(1)$ = 12.03, p < .001].

To further explore this interaction, we calculated the difference between the R^2 values for the
contralateral and ipsilateral conditions for each electrode, using this as a proxy of an encoding
bias between the two arms (Figure 2C, upper right corner of each scatterplot). Values close to
zero indicate similar encoding across the two arms, whereas positive values correspond to
stronger contralateral encoding and negative values stronger ipsilateral encoding. The
distribution for each condition was positively skewed indicating that, overall, there was a bias for
better encoding for contralateral reaches [permutation test: p_{right} < .001, p_{left} < .001]. However,
there was a significant difference in the distributions for the two hemispheres: The bias scores
were lower in the left hemisphere compared to the right hemisphere [permutation test: p <
.001] indicating stronger bilateral encoding in the left hemisphere. We also found that the
contralateral bias becomes weaker the further the electrodes are from the primary motor
cortex, an effect observed in both hemispheres [r_{left} = -0.48, p_{left} < .001, r_{right} = -0.45, p_{right} < .001;
Figure 2 - figure supplement 3].

Opposing patterns of kinematic encoding for the left and right hemisphere during planning and
execution. As neural activity unfolds from preparation to movement, the underlying
computations may change substantially (Elsayed et al., 2016). To examine if hemispheric
asymmetries in encoding depend on task state, we repeated the mixed-effects model described
in the previous section, but now added a factor *Task Phase*, separating the data to test the held-
out predictions during the instruction and movement phases (Fig 3A). The effect of *Arm* was
again significant, with contralateral reaches more strongly encoded than ipsilateral reaches
[$\chi^2(1)$ = 16.19, p < .001]. The main effects of *Hemisphere* [$\chi^2(1)$ = 0.72, p > .40] and *Task Phase*
were not significant [$\chi^2(1)$ = 0.01, p > .90]. Importantly, there was a three-way interaction
between *Arm, Hemisphere* and *Task Phase* indicating that the level of encoding for the two arms
varied across the two hemispheres for the two task phases [$\chi^2(4)$ = 22.47, p < .001].

Figure 3. Opposing encoding patterns for left and right hemisphere across task phase. For all predictive electrodes the time series was segmented into instruction and movement epochs. R^2 was then calculated separately for each epoch. **A) Example model predictions.** Time series of a representative electrode with boxes surrounding the instruction (teal) and movement epochs (grey). **B) Prediction performance during movement and instruction.** Comparison of R^2 values for contralateral and ipsilateral predictions during the instruction epoch (top) and the movement epoch (bottom) for patients with electrodes in the left hemisphere (left) or right (right) hemisphere. Bilateral encoding was stronger in the left hemisphere, an effect that was especially pronounced during the movement phase.

To explore this interaction we again examined the distribution of difference scores for each electrode. The right hemisphere electrodes show a positive skew in both the planning and movement phase [permutation test: $p_{right\_move} < .001$, $p_{right\_planning} < .001$]. However, this pattern is only seen in the left hemisphere during the planning phase [$p_{left\_planning} < .001$]; the mean difference score was not statistically different from zero for the left hemisphere electrodes in the movement phase [$p_{left\_move} = .482$]. Analyzing simple effects within each hemisphere, we found that the difference score was smaller (i.e., more bilateral encoding) in the left hemisphere during the movement phase compared to the planning phase [$\Delta R^2_{left\_move} = 0.01$, $\Delta R^2_{left\_planning} = 0.04$, $p < .001$]. In contrast, the opposite pattern was observed in the right hemisphere, with encoding being more bilateral during the instruction phase [$\Delta R^2_{right\_move} = 0.13$, $\Delta R^2_{right\_planning} = 0.10$, $p < .001$]. Taken together, these results suggest that the left and right hemisphere may have different roles in bilateral encoding with regard to task phase.

Across arm generalization: More overlap between arms in the left hemisphere. The preceding analyses focused on an encoding analysis for within-arm prediction. We next evaluate the

overlap between the neural representations for contralateral and ipsilateral movement. To this
end, we examined across-arm prediction performance by training the kinematic encoding model
with the data from movements produced with one arm and testing prediction performance
using the data from movements produced with the other arm.

Figure 4. Stronger across-arm generalization in the left hemisphere. Across-arm predictions were created
by training the model on ipsilateral reaches and using the trained weights to predict HFA during
contralateral reaches. (Within-arm predictions generated the same as in Figs 2 and 3.) Electrodes close to
the unity line have overlapping neural representations across the two arms whereas electrodes off the
unity line indicate that the two arms are being encoding differentially. **A. Model Predictions.** Predicted
and actual HFA for two electrodes selected from the distribution of left-hemisphere electrodes during
movement, one that generalizes well across arms (E1) and one that fails to generalize (E2). Bottom row
shows within-arm performance for ipsilateral trials, demonstrating that the failure to generalize across
arms does not necessarily indicate poor ipsilateral performance. **B. Across-arm generalization across
patients.** R^2 for within-arm predictions plotted against R^2 for across-arm predictions, with the analysis
performed separately for the instruction and movement phases. Left hemisphere electrodes showed
better generalization than right hemisphere electrodes, an effect that was magnified in the movement
phase.

Figure 4A shows the traces for two representative electrodes, one that shows good
generalization across the two arms and the other that shows poor generalization. For the
electrode that shows good generalization (E1), prediction performance for held-out contralateral
reaches is comparable when the model is trained on data from either the contralateral or
ipsilateral arm. This suggests that there is overlap between the neural representations for
reaches performed with either upper limb for this electrode. In contrast, the electrode showing
poor generalization (E2) showed good prediction for contralateral reaches when trained with
contralateral data, but poor prediction when trained with ipsilateral data. Here the neural
representations for the arms do not overlap. Note that E2 showed relatively strong within-arm
ipsilateral encoding (R^2 = .25); thus, the inability of this electrode to generalize across arms is not

a result of poor encoding of the ipsilateral arm. Rather, E2 encodes movement produced by
either arm, but the manner in which they are encoded differs.

Figure 4B summarizes the comparison of within-arm prediction (y axis) against across-arm
prediction (x axis), with the data separated for the instruction and movement phases. In this
depiction, electrodes close to the unity line have overlapping neural representations during
contra- and ipsilateral movement, whereas electrodes off the unity line encode the two arms
differentially. We again used a permutation-based mixed effects model, this time with fixed
factors of *Generalization, Task Phase* and *Hemisphere* and a random factor of *Participant*. As
expected, we found a main effect of *Generalization* with within-arm predictions outperforming
across-arm predictions [$\chi^2(1)$ = 103, $p < .001$]. There was also a main effect of *Task Phase* [$\chi^2(1)$
= 30.51, $p < .001$] with better prediction accuracy during the movement phase compared to the
instruction phase. The main effect of *Hemisphere was not significant* [$\chi^2(1) < .01$, $p > .90$], but
there was a significant three-way interaction between the three factors [$\chi^2(4)$ = 19.56, $p > .001$].

To evaluate this interaction, we used a difference score, calculated by subtracting the across-arm
R^2 from the within-arm R^2 for each electrode (Figure 4B, upper right corner of each scatterplot).
As such, the larger (more positive) the score, the poorer the electrode is in predicting activity
when the movement is produced with the other arm. Overall, across-arm generalization was
better during the instruction phase compared to the movement phase [$\Delta R^2_{instruction}$ = 0.05,
$\Delta R^2_{movement}$ = 0.08; $p < .001$]. In terms of laterality differences, the left hemisphere showed
stronger across-arm generalization (lower difference values) than the right hemisphere [ΔR^2_{left} =
0.04, ΔR^2_{right} = 0.11; $p < .001$]. Analyzing simple effects within each task phase, the left
hemisphere had better across-arm generalization compared to the right hemisphere for both the
instruction and movement phase [simple effect analysis: $p_{instruct} < .001$, $p_{move} < .001$]. In addition,
better across-arm generalization was found during the instruction phase in both the left and the
right hemispheres [simple effect analysis: $p_{left} < .001$, $p_{right} < .001$]. In sum, the three-way
interaction reflects the fact that electrodes in the left hemisphere exhibit better generalization
across arms than right hemisphere electrodes, an effect observed in both the instruction and
movement phases. In addition, the difference in across arm generalization between phases is
more pronounced in the right hemisphere.

Considered together, the within-arm and across-arm encoding results indicate that left
hemisphere electrodes show stronger bilateral encoding and more similar encoding during left
and right arm movements. Interestingly, although overall encoding was stronger during
movement, across-arm generalization was stronger during the instruction phase.

Temporal and spatial topography of across-arm generalization. To examine how generalization
varied across the cortex, we categorized each electrode as showing either good across-arm
generalization (decrease of up to 20% relative to within-arm performance) or poor across-arm
generalization (decrease of more than 50%; Fig 5A). We focused on the extremes of the
generalization distribution based on the assumption that these electrodes were more likely to
share similar underlying neural profiles. This also allowed us to have similar numbers of
electrodes in each group.

Figure 5B plots ipsilateral encoding strength (R^2) for electrodes that either generalize well or
generalize poorly. Electrodes that generalize poorly have lower encoding strength than
electrodes that generalize well [t = 5.921, $p < .001$]. Despite this difference, 61% (23/38) of
electrodes that generalize poorly meet our encoding criteria of $R^2 > 0.05$.

To examine the dynamics of representational overlap and divergence, we averaged the time-
resolved HFA amplitude across electrodes, restricted to those showing good or poor across-arm
generalization. Figure 5C displays the average time series for contralateral (solid line) and
ipsilateral (dashed line) predictions for electrodes that generalize well (white) or poorly
(magenta). The temporal profile of HFA activity is similar for electrodes that generalize well,
showing a single peak in the movement phase. A cluster-based permutation test identified two
periods where the HFA amplitude differed for contralateral and ipsilateral reaches, one during
instruction and one well into the movement period. In contrast, the temporal profiles are
radically different for those that generalize poorly. The lack of correspondence likely arises, at
least in part, because of the weak modulation for these electrodes during ipsilateral reaches.

Figure 5. Spatial and temporal relationship of across-arm generalization. A. Generalization index.
Electrodes were classified as showing good across-arm generalization (white, generalization index > .80)
or poor across arm generalization (magenta; generalization index < .50). **B. Ipsilateral encoding strength.**
Ipsilateral encoding strength for electrodes that generalize well and generalize poorly. **C. Amplitude**
differences across arms. Average contralateral (solid line) and ipsilateral (dashed line) HFA for electrodes
that generalize well across arms (left) or generalize poorly (right). Significant clusters are represented
with a gray line. Inset: Same data but standardized to highlight shape of the timeseries independent of
absolute amplitude. **D. Spatial distribution of across-arm generalization.** Electrodes that generalize well
across arms (white) were primarily located in dorsal and ventral premotor regions of the left hemisphere.
Electrodes that generalize poorly (magenta) were clustered around the putative arm area of the central
sulcus in both the left and right hemispheres. Dashed rectangles indicate extent of electrode coverage
across the three patients for each hemisphere. **E. Modulation depth.** Depth of tuning was calculated
during instruction or movement with either the ipsilateral or contralateral hand. Greater modulation was
found during contralateral reaches and during movement. **F. Tuning similarity.** Across arms tuning

similarity was calculated for electrodes that generalize well (white) or poorly (magenta). Electrodes that
generalize well across arms had significantly more tuning similarity than electrodes that did not
generalize. *p<0.05, **p<0.01, ***p<0.001, cluster permutation test, Pearson's correlation.

It is possible that similarity in temporal structure is obscured in the preceding analysis by the
differences in HFA amplitude for the electrodes that showed poor generalization. To control for
this, we standardized the time series data by dividing each sample by the overall standard
deviation (Insets: Fig 5C). Using the standardized traces, we calculated the linear correlation
coefficient between the contralateral and ipsilateral traces, separately for instruction and
movement. As expected, electrodes that generalized well across arms showed strong across-arm
correlations for both task phases (Inset Fig 5C, *left*). In contrast, for electrodes that generalize
poorly across arms, the correlation between arms was negative during instruction and then rose
to a moderate positive correlation during movement (Inset Fig 5C, *right*). Thus, the poor
generalization of these electrodes is due in part to the temporal divergence of the two arms
during instruction, where the ipsilateral trace becomes inhibited compared to the contralateral
trace. Interestingly, although the ipsilateral trace remains inhibited during movement, the
temporal structure between the two arms re-emerges.

We next examined the relationship between generalization, hemisphere and spatial position on
the cortical surface. As can be seen in Figure 5D, electrodes that generalize well were
predominantly found in the left hemisphere (proportion of electrodes that generalize across
patients: L1 = 22%, L2 = 39%, L3 = 29%, R1 = 0%, R2 = 5%, R3 = 4%; $\chi^2(2)$ = 48813, p < .001). In
contrast, electrodes showing poor generalization were observed in both hemispheres
(proportion of electrodes that generalize poorly: L1 = 21%, L2 = 5%, L3 = 6%, R1 = 14%, R2 = 15%,
R3 = 37%; $\chi^2(2)$ = 0.400, p = .818). Moreover, in both hemispheres, electrodes showing poor
generalization were clustered near the dorsal portion of the central sulcus, a region
corresponding to the arm area of motor cortex. Electrodes showing strong generalization
(mostly limited to the left hemisphere) tended to be in dorsal and ventral premotor cortices
(PMd, PMv), along with a few in superior parietal cortex. This pattern was also observed when
we analyzed all electrodes, rather than restrict the analysis to those showing extreme values.
Here we used a continuous measure, correlating the amount of across-arm generalization with
the distance (absolute value) from the dorsal aspect of the central sulcus. The correlation was
significant in the left hemisphere [r_{left} = 0.46, p_{left} < .001] but did not reach significance in the
right hemisphere, although the trend was in the same direction [r_{right} = 0.22, p_{right} = .068; Figure 5
- Figure supplement 1].

Target modulation and tuning similarity across arms. To examine the extent of target modulation
for the contralateral and ipsilateral arm, we calculated the modulation depth of each electrode
during the instruction and movement phases. The modulation index reflects the amount of
variability in the signal captured by target tuning (or target specificity): A modulation index of .1
means 10% of the variance is captured by the difference between the response to the four
target locations. The modulation values overall were relatively low (Fig 5D). However, it should
be noted that the reaches were all within the fronto-parallel plane which comprise a
considerably smaller range of movement compared to studies that use a center-out reaching
task. For both electrode types (showing good or poor across arm generalization), there was a
main effect of arm, with ipsilateral modulation lower than contralateral modulation
[permutation test: $p_{Generalize\_well}$ < .001; $p_{Generalize\_poorly}$ < .005]. Both subgroups of electrodes also
displayed a main effect of task phase, with the depth of modulation greater during the

movement phase compared to the instruction phase [permutation test: $p_{Generalize\_well} < .001$;
$p_{Generalize\_poorly} < .005$]. No significant interactions were found for either group.

We also examined the representational overlap between the two arms in terms of their tuning
profiles. We computed a tuning similarity index, defined as the sum of squared errors for
average HFA predictions to the same target between the contralateral and ipsilateral arms. A
similarity index of 1 would correspond to identical tuning preferences for the arms whereas a
similarity index of 0 would indicate completely disparate tuning preferences. The similarity data
were analyzed with a mixed design permutation test, including the factors task phase and
electrode type (good vs. poor generalizers). Electrodes that generalize well across the two arms
(predominately found in the left hemisphere) showed more overlap of tuning preferences
compared to electrodes that generalized poorly [main effect of generalizability: $p < .001$]. While
there was no effect of phase [$p > .70$], the interaction was significant [$p < .005$], with electrode
types showing more comparable tuning similarity during instruction and tuning similarity
diverging during movement. Simple effects analysis revealed that for electrodes that generalize
poorly, tuning similarity was higher during the instruction phase compared to the movement
phase [permutation test: $p < .001$]. In contrast, for electrodes that generalize well, tuning
similarity was higher during movement compared to instruction [permutation test: $p < .001$].
These analyses demonstrate that a number of electrodes in the left hemisphere encode
kinematic variables for both arms, including similar tuning preferences across the two arms,
which was especially pronounced during the movement phase.

Figure 6. Relationship of kinematic features to HFA. A. Relative contribution of kinematic
features to the encoding model. Average contribution of the four kinematic features for
contralateral (top) and ipsilateral (bottom) reaches. Error bars represent the standard deviation
across patients. **B. Temporal pattern of weights across time.** Average weights for all predictive
electrodes are plotted for each kinematic feature for contralateral (top) and ipsilateral (bottom)
reaches. Error bars represent the standard error across electrodes.

HFA relationship to kinematics. In the final analysis, we examined how the kinematic features of the movements contribute to the encoding model used to predict HFA. Each of the four kinematic features includes 400 time lags and thus 400 weights that contribute to the model. To obtain a metric of the relative contribution of the features, we calculated the total contribution of each feature and normalized these values by dividing by the total contribution of the four features. The calculation was done for each patient separately and then averaged, with error bars representing the standard deviation across patients (Fig 6A). The relative contribution of the four kinematic features was similar for contralateral and ipsilateral reaches. We next examined the temporal profile of the weights (Fig 6B) and found that this was also similar for the two conditions, although the average weights for ipsilateral reaches are substantially lower, consistent with the observation of lower performance metrics for ipsilateral reaches across all predictive electrodes (Supplementary figure 3).

As can be seen in Fig 6A, speed and position, kinematic features which are associated with timing and movement initiation make a strong contribution to the encoding model (relative contribution: contra = 68%, ipsi = 63%). This is in contrast with the smaller contribution of theta and phi, features which provide information about movement direction (relative contribution: contra = 32%, ipsi = 37%). This result is similar to that observed in single-unit and population activity recorded in premotor and motor cortex of non-human primates. Kaufman et al. (2016) observed that the largest response component was associated with movement timing/initiation rather than features such as movement direction. Similarly, this direction-independent signal occurs twice during sequential movements (Zimnik and Churchland, 2021); in our data, speed has two prominent peaks, one occurring before the reach and the second occurring before the return movement. We were surprised to see the markedly differential weighting for the vertical (theta) and horizontal (phi) directional features. We assume this is likely idiosyncratic to the layout of our targets.

We also examined the correspondence between HFA and the kinematic features as a function of whether electrodes generalize well or poorly (Fig 7). For electrodes that generalize well, position most closely corresponds to HFA for both contralateral and ipsilateral movements. The maximum cross-correlation for contralateral and ipsilateral movements was found at a lag of 200 ms and 150 ms, respectively, with HFA leading hand position. For electrodes that generalize poorly, the kinematic feature that most closely corresponds to HFA for both contralateral and ipsilateral movements is speed. For these electrodes, the maximum cross-correlation for contralateral and ipsilateral movements were both at a lag of 200 ms, with HFA again leading the kinematic feature. Although the ipsilateral HFA signals are considerably lower in amplitude, the pattern between HFA and speed is quite similar for both ipsilateral and contralateral movements. The fact that the neural activity from electrodes that generalize poorly (primarily located over M1) correlates well with speed provides additional evidence that a strong component of the HFA ECoG signal is related to timing and movement initiation (Kaufman et al., 2016).

Figure 7. Relationship of kinematic features to HFA for electrodes that generalize well or poorly. A. HFA overlaid with kinematics. Each plot shows average HFA activity along with average values for position, velocity, or speed, time-locked to movement onset. **B. Cross correlation between HFA and kinematic features.** The cross-correlation between each kinematic feature and the HFA is plotted for electrodes that generalize well or poorly. For each plot, the lag (in ms) yielding the highest correlation between HFA and kinematics is represented by a dashed line with the lag time and correlation coefficient displayed in the inset at the bottom right of each plot.

Discussion

Although the most prominent feature of cortical motor pathways is their contralateral organization, unimanual movements are well represented in the ipsilateral hemisphere. Single-unit activity and local field potentials obtained from motor cortex in non-human primates (Ganguly et al., 2009; Ames & Churchland, 2019), as well as ECoG activity in humans (Bundy, Szrama, Pahwa & Leuthardt, 2018; Ganguly et al., 2009, Wisneski et al., 2008) can be decoded to predict complex kinematic variables and EMG activity during arm movements of the ipsilateral arm.

Here we report the results from ECoG data obtained from six patients, three with left hemisphere implants and three with right hemisphere implants, with each patient having at least 17 predictive electrodes. To examine how kinematic features are represented in each hemisphere, we built an electrode-wise encoding model. Such models allow prediction of the full time series for each electrode thus retaining the high spatial and temporal resolution of the intracranial signal. From these metrics we could compare kinematic encoding and across-arm generalization between the two hemispheres as well as the spatial distribution of the information-carrying electrodes within each hemisphere.

Hemispheric asymmetry in movement encoding. We observed a striking asymmetry between the two hemispheres for ipsilateral movement encoding. While contralateral movements were encoded similarly across the two hemispheres, ipsilateral encoding was much stronger in the left hemisphere, an effect that was especially pronounced during movement execution. In addition, there was greater overlap between the representation of contra- and ipsilateral movement in the left hemisphere compared to the right hemisphere.

The bilateral encoding effect size is quite substantial (Cohen's d = 1.34), exceeding the conventional criterion for a large effect (d = .80). Given the size of the hemispheric asymmetry effect, it is surprising that this asymmetry has not be described in previous reports. This may in part reflect the smaller sample size in ECoG studies. For example, in Bundy et al. (2018), three of the four patients had left hemisphere grids, leaving a hemisphere analysis dependent on the data from a single right hemisphere patient. Studies with non-human primates tend to ignore hemispheric differences, perhaps because these animals do not show consistent patterns of hand-dominance across individuals. One exception here is a study by Cisek, Crammond and Kalaska (2003) who reported no hemispheric differences in neural recordings obtained from M1 and PMd during ipsilateral and contralateral arm reaches.

In addition to examining hemispheric differences in the encoding of unimanual movement, we also asked if kinematic features were encoded differently for contra- and ipsilateral movements by testing across-arm generalization. We categorized electrodes as showing either good across-arm generalization (decrease of up to 20% relative to within-arm performance) or poor across-arm generalization (decrease of more than 50%). This categorization scheme revealed a striking anatomical division, with electrodes showing good across-arm generalization clustering in the left premotor and superior parietal regions and electrodes that generalized poorly clustering in left and right M1. This result does not appear to reflect a sampling bias. There are more left hemisphere electrodes compared to right hemisphere electrodes because two of the left hemisphere patients had high density grid implants. However, the electrodes in our left and right hemisphere samples were similarly distributed over premotor, sensorimotor and parietal regions (see Supplementary Figure 2).

Examining the temporal profile of electrodes that generalize poorly, we found that the divergence between the two arms occurs during instruction, with the ipsilateral trace becoming inhibited relative to the contralateral trace. TMS studies have shown reduced corticospinal excitability of the non-selected hand during movement preparation (e.g., Leocani, Cohen, Wassermann, Ikoma & Hallett, 2000; Liepert, Dettmers, Terborg & Weiller, 2001), an effect that has been hypothesized to reflect inhibition from prefrontal regions to facilitate response selection (Duque et al., 2012).

We further examined the spatial tuning of the electrodes. Target tuning in the HFA band was found for both contralateral and ipsilateral movement, although ipsilateral tuning was significantly shallower. Interestingly, electrodes that generalized well across-arms had similar spatial/target tuning for each arm. This suggests that for these electrodes, ipsilateral signals are not just encoding generic movement, but encoding movement direction in a similar manner to contralateral signals. A similar overlap in tuning has been observed in single unit recordings from PMd (Cisek, Crammond & Kalaska, 2003) and can be inferred from the across-arm generalization decoding results reported by Bundy et al., (2018). In contrast, electrodes that failed to

550 generalize, located primarily in M1 in either left or right hemisphere, exhibited disparate tuning
551 for contra- and ipsilateral reaches.

Functional implications of hemispheric asymmetries in movement encoding. By using a delayed
response task, we were able to segregate activity into an instruction phase during which the
participant was presented with the target location for the forthcoming movement and a
movement phase, defined at the onset of the reach. With this design, we found that the
encoding model could predict neural activity during the instruction phase based on the
kinematics of the forthcoming reach, evidence that the participants were indeed planning the
upcoming movement.

This task phase analysis also revealed robust asymmetries between the two hemispheres. There
was a main effect of hemisphere, with the left hemisphere displaying stronger bilateral encoding
overall compared to the right hemisphere. However, there was also an interaction: In the left
hemisphere bilateral encoding was stronger during the movement phase whereas in the right
hemisphere bilateral encoding was stronger during the instruction phase. Surprisingly, in the left
hemisphere the contralateral bias completely disappeared during the movement phase, with
both the contra- and ipsilateral arms being encoded to the same extent. Stronger bilateral
encoding during movement (compared to instruction) is surprising given the spatial distribution
of electrodes that encode ipsilateral movement were primarily outside of M1, regions typically
associated more with planning than execution (e.g., premotor cortices, parietal cortex).

The asymmetry observed here is in accord with the long-standing recognition of hemispheric
asymmetries in praxis. Starting with the classic observations of Liepmann at the turn of the 20[th]
century on the association of the left hemisphere and apraxia (Liepmann 1908, cited in Renzi, &
Lucchelli, 1988; see also Schaefer et al., 2007) and continuing with functional imaging studies in
neurotypical populations, a large body of evidence points to a dominant role for the left
hemisphere in skilled movement (Corballis, Badzakova-Trajkov & Häberling, 2012; Przybylski &
Króliczak, 2017). This asymmetry is most pronounced in tasks involving functional object use
(Buxbaum et al., 2006), symbolic gestures (Xu, Gannon, Emmorey, Smith & Bruan, 2009) and
intransitive pantomimes (Bohlhalter et al., 2009). While the neuropsychological and
neuroimaging work have highlighted the involvement of left premotor and parietal cortex in
praxis, corresponding asymmetries have also been noted in subcortical structures such as the
basal ganglia, cerebellum, and thalamus (Swinnen et al., 2010). We note that for our patient
population, we are limited to regions of the brain where we have sufficient electrode coverage.

Apraxia, following left-hemisphere damage can be manifest in movements produced with either
limb (Renzi, & Lucchelli, 1988), and are usually associated with lesions that encompass premotor
and parietal cortices (Haaland, Harrington & Knight, 2000). While this asymmetry may be linked
to hand dominance (Ochipa, Rothi & Heilman, 1989), functional imagining studies with relatively
large sample sizes have shown that handedness only influences the strength of the left
hemisphere bias for skilled movement but does not produce a reversal in left handers
(Vingerhoets et al., 2012; Verstynen et al., 2005, Chettouf et al., 2020, Vingerhoets et al., 2013).
Of the six patients tested in the current study, five are right-handed and the remaining patient
reported being ambidextrous with a slight preference for using the left hand. We note that the
results from this patient (L3) did not qualitatively differ from the other two left hemisphere
patients.

Ipsilateral encoding was most prominent in the premotor and parietal cortex of the left
hemisphere, overlapping with the neural regions implicated in praxis. However, two features of
our results do not map on readily to an interpretation that focuses on hemispheric asymmetries
in praxis. First, our task involved simple reaching movements, whereas praxis generally
encompasses more complex learned movements associated with tool use or symbolic gestures.
Second, ipsilateral encoding became more pronounced during movement execution.

Assuming that ipsilateral encoding is indicative of bilateral motor representations, one might
have expected, a priori, that hemispheric asymmetries in ipsilateral encoding would be more
prevalent during movement planning. We recognize that, even with delayed response tasks, it is
overly simplistic to assume that the activity cleanly separates into planning and execution
phases. This is especially true with sequential movements where planning effects are observed
both prior to and during movement execution (Ariani & Diedrichsen, 2019; Zimnick &
Churchland, 2021). While the experimental task was to reach to a cued target, the participants
made out-and-back movements, returning to the home position in a relatively smooth manner
(see Supplemental Figure 1). It is reasonable to assume that some component of the activity
during the primary outward movement was related to planning the return movement. As such,
we are hesitant to draw strong inferences about the differences between ipsilateral encoding
during action planning and movement execution.

To this point, we have focused on how ipsilateral activity may be reflective of control processes
associated with movements of the ipsilateral arm. However, it is possible that this activity is
related to postural stabilization of the body during reaching. Indeed, extrapyramidal pathways
such as the reticulospinal track have a prominent ipsilateral projection associated with postural
control (Cleland, Madhavan, 2021), and these pathways receive cortical input. We aimed to
reduce postural demands in our task by having the participants seated in an upright hospital bed
with the back fully supported; nonetheless, there are surely postural shifts associated with the
reaching movement. The clinical setting precluded the use of EMG or video, measurements that
would allow a quantitative assessment of the relationship of ipsilateral activity to postural
adjustments. We do note a few points that are at odds with a posture-based account of
ipsilateral activity. First, we are unaware of evidence suggesting a left hemisphere specialization
in postural control. In fact, the evidence suggests that postural instability is more frequently
associated with right-hemisphere lesions (Bohannon, Smith & Larkin 1986; Spinazzola, Cubelli &
Della Sala 2003). Second, postural adjustments typically precede movement onset (Belen'kii,
Gurfinkel & Pal'tsev 1967, cited in Guiard 1987), whereas ipsilateral encoding in our data
increased with movement onset. Third, and perhaps most convincing, the encoding model
shows that the ipsilateral signals predict reaching kinematics with relatively high precision.

The asymmetry of ipsilateral encoding may be reflective of a prominent role of the left
hemisphere in bimanual coordination (Jäncke et al., 2000; Toyokura, Muro, Komiya & Obara,
1999; Maki, Wong, Sugiura, Ozaki & Sadato, 2008; Serrien, Cassidy & Brown, 2003; Fujiyama et
al., 2016). For example, Schaffer et al., (2020) observed greater impairments in bimanual
coordination following left hemisphere stroke compared to right hemisphere stroke.
Interestingly, the impairment was manifest prior to peak velocity, a finding interpreted as a
disruption in predictive control. It may be that the left hemisphere makes an asymmetric
contribution to inter-limb coordination by tracking or predicting where both limbs are in space.
As such, the encoding of ipsilateral arm movement might be a form of state representation, a
means to keep track of the state of the ipsilateral arm given that many actions require the

coordinated activity of the two limbs. This hypothesis, derived from the current data, is
consistent with the increased ipsilateral encoding during the movement phase. The need to
monitor the state of the other limb should hold for unimanual gestures performed with either
limb.

An important question for future work is to examine how ipsilateral representations in the left
hemisphere are affected during more complex movements, including those that involve both
limbs. Using fMRI, Diedrichsen, Wiestler and Krakauer (2013) compared ipsilateral movement
representations during unimanual and bimanual movements. Within the primary motor cortex,
ipsilateral representations could only be discerned during unimanual movement. However,
caudal premotor and anterior parietal regions retained similar ipsilateral representation during
uni- and bimanual movement. If the left hemisphere tracks both limbs to facilitate bimanual
coordination, we would predict that ipsilateral representations in premotor cortex are retained
more strongly in the left hemisphere compared to the right hemisphere when both arms are
engaged in the task.

Conclusion: Using a kinematic encoding model, we observed a striking hemispheric asymmetry,
with the left hemisphere more strongly encoding the ipsilateral arm than the right hemisphere, a
finding that was apparent during preparation and amplified during movement. This asymmetry
was primarily driven by electrodes positioned over premotor and parietal cortices, with strong
contralateral encoding for electrodes positioned over sensorimotor cortex. We propose that
these networks monitor the state of each arm, a prerequisite for most skilled actions.

METHOD (2676 words)

Participants. Intracranial recordings were obtained from six patients (2 female; 5 right-handed;
Table 1) implanted with subdural grids as part of their treatment for intractable epilepsy. Data
were recorded at three hospitals: University of California, Irvine (UCI) Medical Center (n = 2),
University of California, San Francisco (UCSF) Medical Center (n = 2) and California Pacific
Medical Center (CPMC), San Francisco (n = 2). Electrode placement was solely determined based
on clinical considerations and all procedures were approved by the institutional review boards at
the hospitals, as well as the University of California, Berkeley. All patients provided informed
consent prior to participating in the study.

Behavioral task. Participant performed an instructed-delay reaching task while sitting upright in
their hospital bed. The participant rested their arms on a horizontal platform (71 cm x 20 cm)
that was placed over a standard hospital overbed table. The platform contained two custom-
made buttons, each connected to a microswitch. At the far end of the platform (13 cm from the
buttons, approximately 55 cm from the participant's eyes), a touchscreen monitor was attached,
oriented vertically. Visual targets could appear at one of six locations, four for each arm (Figure
1a). The two central locations were used as targets for reaches with either arm; the two
eccentric targets varied depending on the arm used. Stimulus presentation was controlled with
Matlab 2016a. A photodiode sensor was placed on the monitor to precisely track target
presentation times. The analog signals from the photodiode and the two microswitches were fed
into the ECoG recording system and were digitized into the same data file as the ECoG data with
identical sampling frequency.

Testing of the contralateral and ipsilateral arms (relative to the ECoG electrodes) was conducted
in separate experimental blocks that were counterbalanced. To start each trial, the participant

placed their left and right index fingers on two custom buttons to depress the microswitches
(this indicated they were in the correct position and ready to start the trial). If both
microswitches remained depressed for 500 ms, a fixation stimulus was presented in the middle
of the screen for 750 ms, followed by the target, a circle (1.25 cm diameter) which appeared in
one of the four locations. Another hold period of 900 ms followed in which the participant was
instructed to prepare the required movement while the target remained on the screen. If the
microswitch was actuated during this hold period, an error message appeared on the screen and
the program would advance to the next trial. If the start position was maintained, a compound
imperative stimulus was presented at the end of the hold period. This consisted of an auditory
tone and an increase in the size of the target (2.5 cm diameter). The participant was instructed
that this was the signal to initiate and complete a continuous out-and-back movement,
attempting to touch the screen at the target location before returning back to the platform. The
target disappeared when the touchscreen was contacted. The imperative was withheld on 5% of
the trials ('catch' trials) to ensure that the participant only responded after the onset of the
imperative.

Once back at the home position, the screen displayed the word 'HIT' or 'MISS' for 750 ms to
indicate if the touch had occurred within the target zone. The target zone included the 2.5
diameter circle as well as a 1cm buffer around the target. After the feedback interval, the screen
was blank for 250 ms before the reappearance of the fixation stimulus, signaling the start of the
next trial. The participants were informed to release either of the buttons at any time they
wished to take a break.

Each block consisted of 40 trials (10/target), all performed with a single limb. Blocks alternated
between contra- and ipsilateral arms (relative to the ECoG electrodes), with the order
counterbalanced across participants. Each block took approximately 5-6 minutes to complete.
All participants completed at least two blocks with each per arm (Table 1).

Movement analysis and trajectory reconstruction. We used two methods to analyze the
movements. For the first method, we recorded key events defined by the release of the
microswitch at the start position, time and location of contact with the touchscreen, and return
time to the home position, defined by the time at which they depressed the home position
microswitch. For the second method, we used the Leap Motion 3-d movement analysis system
(Weichert, et al., 2013) to record continuous hand position and the full movement trajectory
(sampling rate = 60 Hz). Although the Leap system is a lightweight video-based tracking device
that is highly mobile, the unpredictable environment of the ICU led to erratic recordings from
the Leap system. For example, patients frequently had intravenous lines in one or both hands
which obstructed the visibility of the hand and interfered with the ability of the Leap system to
track the hand using their built-in hand model. This resulted in lost samples and therefore
satisfactory kinematic data was obtained from only a subset of conditions collected from
patients using the Leap system.

Given the limitations with the Leap data, we opted to use a simple algorithm to reconstruct the
time-resolved hand trajectory in each trial, estimating it from the event-based data obtained
with the first method. We used a beta distribution to estimate the velocity profile of the forward
and return reach based on reach times and the travel distance (sampling rate = 100 Hz). We
opted to use a beta distribution because this best matched the velocity profiles of the data
obtained with the Leap system.

For conditions that had clean kinematic traces (no lost samples) from the Leap system, we
compared the estimated kinematic profiles with those obtained with the Leap system. There was
a high correlation between the two data sets (r = .98 for position in the Z dimension; r = .93 for
velocity in the Z dimension; Figure 1-figure supplement 1). We note that our method of
estimating the trajectories results in a smoothed version of the movement, one lacking any
secondary or corrective movements that are sometimes observed when reaching to a visual
target (Suway & Schwartz, 2019). We believe this is still a reasonable estimation given the high
correlation with the continuous Leap data, and the fact that participants had ample time to
prepare the movements and were instructed and observed to make ballistic movements by the
experimenter who was present for all recording sessions (CMM).

Electrode Localization. Grid and strip electrode spacing was 1 cm in four patients and 4 mm in
the two other patients. The electrode locations were visualized on a three-dimensional
reconstruction of the patient's cortical surface using a custom script that takes the post-
operative computer tomography (CT) scan and co-registers it to the pre-operative structural
magnetic resonance (MR) scan (Stolk et al., 2018).

Neural data acquisition and preprocessing. Intracranial EEG data and peripheral data
(photodiode and microswitch traces) were acquired using a Nihon Kohden recording system at
UCI (128 channel, 5000 Hz digitization) and CPMC (128 channel, 1000 Hz digitization rate), and
two Tucker Davis Technologies recording systems at UCSF (128 channel, 3052 Hz digitization
rate).

Offline preprocessing included the following steps. First, if the patient's data was not sampled at
1000 Hz (UCI and UCSF recording sites), the signal from each electrode was low-pass filtered at
500 Hz using a Butterworth filter as an anti-aliasing measure before down-sampling to 1000 Hz.
Electrodes were referenced using a common average reference. Each electrode was notch-
filtered at 60, 120 and 180 Hz to remove line noise. The signals were then visually inspected and
electrodes with sustained excessive noise were excluded from further analyses. The signals were
also inspected by a neurologist (RTK) for epileptic activity and other artifacts. Electrodes that had
pathological seizure activity were also excluded from the main analyses. Out of 752 electrodes,
82 were removed due to excessive noise and 5 were removed due to epileptic activity, resulting
in a final data set of 665 electrodes. Catch trials and unsuccessful reaches were not included in
the analyses.

From the cleaned data set, we extracted the HFA instantaneous amplitude using a Hilbert
transform. To account for the $1/f$ power drop in the spectrum, we divided the broadband signal
into five narrower bands that logarithmically increased from 70 to 200 Hz (i.e., 70-86, 86-107,
107-131, 131-162, 162-200 Hz), and applied a band-pass filter within each of these ranges. We
then took the absolute value of the Hilbert transform within each band-pass, performed a z-
score transformation, and averaged the five values. Z-scoring was performed after concatenating
all the blocks for each patient, ensuring that we did not obscure possible amplitude differences
across the two arms. As a final step, the data were down-sampled to 100 Hz to reduce
computational load (e.g., number of parameters in the encoding model, see below). HFA
amplitude fluctuations (envelope; are evident at lower frequencies (Canolty et al., 2006; Pei et
al., 2011).

Feature selection. Four estimated kinematic features were used to predict HFA (Figure 1B *left*).
The first two features were position and speed in the Z dimension. This dimension captures

variability related to movement that is relatively independent of target location (i.e., along the
axis between the participant and touchscreen). The second pair of features were spherical angles
that define the specific target locations (Figure 1A *right*). Features were selected to reduce
collinearity and redundancy in the encoding model. Because we include time lags for each
kinematic feature, derivatives can emerge from the linear model (e.g., velocity and acceleration
can be created from position); thus, velocity and acceleration were not included as additional
features. Speed is a non-linear transformation of position and is added as a separate feature.

Kinematic encoding model. The estimated kinematic features were used to predict the HFA for
each electrode (Fig 1F). We created a 4 x 400 feature matrix by generating a time series for each
feature by time-lagging the values of the selected feature relative to the neural data, with lags
extending from 2 s before movement onset to 2 s after movement onset (sampling rate at 100
Hz). This wide range of lags serves two purposes. First, it provides a way to compensate for the
anticipated asynchrony between neural data and movement kinematics. Second, it allowed us to
evaluate HFA activity during the instructed delay (beginning ~1.5 s before movement onset)
period as well as during movement. HFA at each time point [HFA(t)] was modeled as a weighted
linear combination of the kinematic features at different time-lags, resulting in a set of beta
weights, b1 …, b400 per kinematic feature. To make the beta weights scale-free, the kinematic
features and neural HFA were z-scored before being fit by the model.

Model fitting. Regularized (ridge) regression (Hoerl and Kennard, 1970) was used to estimate the
weights that map each kinematic feature (X) to the HFA signal (y) for each electrode, with λ
being the regularization hyperparameter:

790
$$\hat{\beta} = (X^T X + \lambda I)^{-1} X^T y$$

For within-arm model fitting, the total dataset consisted of all clean, successful trials performed
with either the ipsilateral or contralateral arm (each arm was fit separately). Nested five-fold
cross-validation was used to select the regularization hyperparameter on inner test sets
(validation sets) and assess prediction performance on separate, outer test sets. At the outer
level, the data was partitioned into five mutually exclusive estimation and test sets. For each test
set, the remaining data served as the estimation set. For each outer fold, we further partitioned
our estimation set into five mutually exclusive inner folds to train the model (80% of estimation
set) and predict neural responses across a range of regularization values on the validation set
(20% of estimation set). For each inner fold, the regularization parameter value was selected
that produced the best prediction as measured by the linear correlation of the predicted and
actual HFA. The average of the selected regularization parameters across the five inner folds was
computed and used to calculate the prediction of the HFA on the outer test set. This procedure
was done at the outer level five times. Our primary measure is held-out prediction performance
(R^2), which we quantified as the squared linear correlation between the model prediction and
the actual HFA time series, averaged across the five mutually exclusive test sets.

To be considered as predictive, we established a criterion that an electrode must account for at
least 5% of the variability in the HFA signal ($R^2 > .05$) for either ipsilateral or contralateral reaches
(Downey et a., 2020). Electrodes not meeting this criterion were not included in subsequent
analyses.

For across-arm model fitting, the same procedure was used except the test set was partitioned
from the total dataset of the other arm. We partitioned the data in this manner (80% estimation,

20% test) to make the fitting procedure for the across-arm model comparable to that employed
in the within-arm model.

Tuning modulation and similarity across arms. Modulation depth (MD) of target tuning was
calculated as the standard deviation of the mean HFA predictions for each of the four target
locations:

$$MD = \sum_{i=1}^{n} \frac{(x_i - \bar{x})^2}{n}$$

where $\bar{x}$ is the average HFA prediction, x_i is the average HFA prediction for each of the four
target positions. i iterates through the target locations and n is the total number of target
locations. This process was done separately for contralateral and ipsilateral HFA predictions.

To assess similarity in tuning across the two arms, we computed the SSE (sum of squared errors)
for average HFA predictions to the same target between the contralateral and ipsilateral arms.
This calculation was computed for each electrode as follows:

$$SSE_e = \sum_{i=1}^{n} (\{contra_i - ipsi_i)\}^2$$

where *contra* and *ipsi* are average HFA predictions for a given target location reached with either
the contralateral or ipsilateral arm. Note that for this calculation n is limited to the two positions
that served as target locations for both arms.

This metric was only calculated for the two central targets, the targets common to both arms
(the two eccentric target locations varied depending on the arm used). These values were scaled
from 0 to 1 based on the minimum and maximum values of SSE across all electrodes. SSE
represents a metric of dissimilarity; To calculate a similarity index (SI), we converted this to a
measure of similarity by subtracting the scaled SSE values from 1:

$$SI = 1 - \frac{SSE_e - min(SSE_e)}{max(SSE_e) - min(SSE_e)}$$

Thus, higher SI represents more similar average predictions.

Separating instruction and movement phases. The encoding model was run to predict the full
HFA time course. To compare model prediction performance during different phases of the task,
the data were epoched into instruction and movement phases, using event markers recorded in
the analog channel (i.e., cue onset and movement onset). Epochs of the same task phase were
concatenated together, and prediction performance was operationalized as the square of the
Pearson correlation between the predicted and actual HFA for each task phase.

Permutation-based linear mixed effects model. Linear mixed effects models were carried out in
RStudio using software packages lme4 and permlmer (Bates, Mächler, Bolker & Walker, 2014;
Lee & Braun, 2012). Each mixed effects model used participant as a random effect and
experimental variables (e.g., reaching arm, hemisphere) as fixed effects. The models were used
to predict performance from the kinematic encoding model for all predictive electrodes. Nested

models were created to assess the effect of the fixed factors, with the null model using patient
as a random effect to predict encoding values. For the nesting, fixed factors were added to the
model to assess if each new factor improved prediction above that obtained with the null model
using a permutation-based method (Lee & Braun, 2012). Interactions were tested by comparing
a model in which the fixed effects were restricted to have additive effects to a model that could
have both multiplicative and additive effects.
852
Assessing the contribution of kinematic features in the encoding model. Average weights for
each patient were calculated for all predictive electrodes for each kinematic feature. Because
each feature has 400 time lags, there are 400 weights per feature in the full model. To assess the
relative contribution of each feature, we calculated the sum of the weights across time lags after
taking the absolute value of each weight (since negative weights are as informative as positive
weights). The sum of each feature was then plotted as a proportion against the total sum of all
weights (after taking the absolute value) to assess the relative contribution. To capture the
temporal profile of the weights, the average was taken for all predictive electrodes for all
patents.
Assessing explainable variance in the encoding model. There are several methods to assess
prediction performance for encoding models (Lage-Castellanos, Valente, Formisano & De
Martino, 2019; Nunez-Elizalde, Huth & Gallant, 2019; Schoope et al., 2016). A drawback with the
Pearson's product-moment correlation coefficient is that it does not distinguish between
explainable variability and response variability (Hsu et al., 2004). To better capture this
distinction, we calculated CC_{norm}, a measure which normalizes by signal power to account for
response variability (Schoope et al., 2016):

$$CC_{norm} = \frac{Cov(y,\hat{y})}{\sqrt{Var(\hat{y})}}\sqrt{\frac{1}{SP}}$$

871   where y is neural activity and $\hat{y}$ are predictions from the model, and signal power (SP) is:

$$SP = \frac{Var(\sum_{n=1}^{N} R_n) - \sum_{n=1}^{N} Var(R_n)}{N(N-1)}$$

where R_n is the neural time series for the n^{th} trial and N is the total number of trials.

We plot the density estimates of all predictive electrodes for both the Pearson's correlation
coefficient (CC_{abs}) and CC_{norm} (see Figure 2 – figure supplement 2). We find similar values for the
two metrics, suggesting that the number of trials in our study and signal power is sufficient for
the encoding model to capture the majority of explainable variance. (Calculations of Pearson's
correlation coefficient were always taken from held-out trials.)

Calculating distance from dorsal central sulcus. For each patient, 30 discrete (x, y) coordinates
were manually demarcated along the central sulcus on individual MRI scans. The 30 points were
then interpolated to create a line traversing the central sulcus for each individual. The dorsal
aspect of the central sulcus was defined as all points dorsal to the midpoint of the central sulcus.

We then calculated the absolute distance between each electrode and the closest point on the
dorsal aspect of the central sulcus (our interpolated line).

ACKNOWLEDGMENTS:
887
888 We want to thank Anwar Nunez-Elizalde for his invaluable guidance building the encoding
889 model, Ian Greenhouse for building the first prototype rig to bring into the ICU and William
890 Liberti for his advice with the figures. This research was supported by the National Institute of
891 Health: NS097480.

892
893
894 **DECLARATION OF INTEREST:** The authors declare no competing interests.
895

911 Figure 1-figure supplement 1. **Actual vs. estimated kinematics.** Sample of position and velocity
912 traces along the Z dimension for one participant. Kinematics from the leap motion controller are
913 plotted in black and the estimated kinematics calculated from task parameters are in blue.
914

Figure 2- figure supplement 1. Similar distribution in left and right hemispheres of electrodes in premotor, sensorimotor and parietal regions. Each electrode was categorized based on a cortical parcellation provided by the Desikan-Killiany atlas. When pooled across the three left hemisphere and three right hemisphere patients, the proportion of electrodes in the highlighted regions was similar. 70% of all electrodes were assigned to one of these regions.

Figure 2 - figure supplement 2. Assessing explainable variance in the encoding model. A.
Distribution of electrodes using Pearson's correlation coefficient (CC_{abs}). Histograms of the
average held-out correlation for predictive electrodes during ipsilateral and contralateral reaches
for the right hemisphere (top) and left hemisphere (bottom). B. Distribution of all electrodes
using the correlation coefficient normalized by signal power (CC_{norm}). Same as A except the x-
values now indicate CC_{norm} values, a measure designed to account for response variability (see
Methods). The two metrics provide similar density estimates, suggesting that the number of
observations and signal power in our experimental data are sufficient to capture the majority of
explainable variance.

Figure 2 - figure supplement 3. Distance from central sulcus negatively correlates with bilateral encoding. Degree of unimanual encoding, calculated as difference in R^2 for contralateral and ipsilateral movement, is plotted as a function of absolute distance from the dorsal aspect of the central sulcus. Electrodes further from the central sulcus had stronger encoding of the ipsilateral arm for both the left and the right hemisphere. *$p<0.05$, **$p<0.01$, ***$p<0.001$.

Figure 5 - figure supplement 1. Distance from central sulcus positively correlates with degree of
overlap between representations for the left and right movements. Across-arm generalization
was calculated as the ratio of across-arm performance divided by within-arm performance.
Electrodes further from the dorsal aspect of the central sulcus had better across-arm
generalization in the left hemisphere. This relationship was not significant in the right
hemisphere. *p<0.05, **p<0.01, ***p<0.001.

References:

Nyberg-Hansen, R., & Rinvik, E. (1963). Some comments on the pyramidal tract, with special reference to its individual variations in man. *Acta Neurologica Scandinavica*, *39*(1), 1-30.

Bourbonnais, D., & Noven, S. V. (1989). Weakness in patients with hemiparesis. *American Journal of Occupational Therapy*, *43*(5), 313-319.

Babiloni, C., Carducci, F., Cincotti, F., Rossini, P. M., Neuper, C., Pfurtscheller, G., & Babiloni, F. (1999). Human movement-related potentials vs desynchronization of EEG alpha rhythm: a high-resolution EEG study. *Neuroimage*, *10*(6), 658-665.

Ghacibeh, G. A., Mirpuri, R., Drago, V., Jeong, Y., Heilman, K. M., & Triggs, W. J. (2007). Ipsilateral motor activation during unimanual and bimanual motor tasks. *Clinical Neurophysiology*, *118*(2), 325-332.

Ganguly, K., Secundo, L., Ranade, G., Orsborn, A., Chang, E. F., Dimitrov, D. F., ... & Carmena, J. M. (2009). Cortical representation of ipsilateral arm movements in monkey and man. *Journal of Neuroscience*, *29*(41), 12948-12956.

Ames, K. C., & Churchland, M. M. (2019). Motor cortex signals for each arm are mixed across hemispheres and neurons yet partitioned within the population response. *Elife*, *8*, e46159.

Bundy, D. T., Szrama, N., Pahwa, M., & Leuthardt, E. C. (2018). Unilateral, 3D arm movement kinematics are encoded in ipsilateral human cortex. *Journal of Neuroscience*, *38*(47), 10042-10056.

Wisneski, K. J., Anderson, N., Schalk, G., Smyth, M., Moran, D., & Leuthardt, E. C. (2008). Unique cortical physiology associated with ipsilateral hand movements and neuroprosthetic implications. *Stroke*, *39*(12), 3351-3359.

Cisek, P., Crammond, D. J., & Kalaska, J. F. (2003). Neural activity in primary motor and dorsal premotor cortex in reaching tasks with the contralateral versus ipsilateral arm. *Journal of Neurophysiology*, *89*(2), 922-942.

Duque, J., & Ivry, R. B. (2009). Role of corticospinal suppression during motor preparation. *Cerebral cortex*, *19*(9), 2013-2024.

Steinberg, O., Donchin, O., Gribova, A., De Oliveira, S. C., Bergman, H., & Vaadia, E. (2002). Neuronal populations in primary motor cortex encode bimanual arm movements. *European Journal of Neuroscience*, *15*(8), 1371-1380.

Willett, F. R., Deo, D. R., Avansino, D. T., Rezaii, P., Hochberg, L. R., Henderson, J. M., & Shenoy, K. V. (2020). Hand knob area of premotor cortex represents the whole body in a compositional way. *Cell*, *181*(2), 396-409.

Heming, E. A., Cross, K. P., Takei, T., Cook, D. J., & Scott, S. H. (2019). Independent representations of ipsilateral and contralateral limbs in primary motor cortex. *ELife*, *8*, e48190.

Downey, J. E., Quick, K. M., Schwed, N., Weiss, J. M., Wittenberg, G. F., Boninger, M. L., &
Collinger, J. L. (2020). The motor cortex has independent representations for ipsilateral and
contralateral arm movements but correlated representations for grasping. *Cerebral*
Cortex, *30*(10), 5400-5409.

Corballis, M. C., Badzakova-Trajkov, G., & Häberling, I. S. (2012). Right hand, left brain: genetic
and evolutionary bases of cerebral asymmetries for language and manual action. *Wiley*
Interdisciplinary Reviews: Cognitive Science, *3*(1), 1-17.

Rothi, L. G., Ochipa, C., & Heilman, K. M. (1997). A cognitive neuropsychological model of limb
praxis and apraxia. *Apraxia: The neuropsychology of action*, 29-49.

Schaefer, S. Y., Haaland, K. Y., & Sainburg, R. L. (2007). Ipsilesional motor deficits following stroke
reflect hemispheric specializations for movement control. *Brain*, *130*(8), 2146-2158.

Liepmann, H. (1908). *Drei aufsätze aus dem apraxiegebiet* (Vol. 2545). S. Karger.

De Renzi, E., & Lucchelli, F. (1988). Ideational apraxia. *Brain*, *111*(5), 1173-1185.

Haaland, K. Y., Harrington, D. L., & Knight, R. T. (2000). Neural representations of skilled
movement. *Brain*, *123*(11), 2306-2313.

Chettouf, S., Rueda-Delgado, L. M., de Vries, R., Ritter, P., & Daffertshofer, A. (2020). Are
unimanual movements bilateral? *Neuroscience & Biobehavioral Reviews*, *113*, 39-50.

Verstynen, T., Diedrichsen, J., Albert, N., Aparicio, P., & Ivry, R. B. (2005). Ipsilateral motor cortex
activity during unimanual hand movements relates to task complexity. *Journal of*
Neurophysiology, *93*(3), 1209-1222.

Verstynen, T., & Ivry, R. B. (2011). Network dynamics mediating ipsilateral motor cortex activity
during unimanual actions. *Journal of Cognitive Neuroscience*, *23*(9), 2468-2480.

Schäfer, K., Blankenburg, F., Kupers, R., Grüner, J. M., Law, I., Lauritzen, M., & Larsson, H. B.
(2012). Negative BOLD signal changes in ipsilateral primary somatosensory cortex are associated
with perfusion decreases and behavioral evidence for functional inhibition. *Neuroimage*, *59*(4),
3119-3127.

Leszczyński, M., Barczak, A., Kajikawa, Y., Ulbert, I., Falchier, A. Y., Tal, I., ... & Schroeder, C. E.
(2020). Dissociation of broadband high-frequency activity and neuronal firing in the
neocortex. *Science Advances*, *6*(33), eabb0977.

Muthukumaraswamy, S. D. (2010). Functional properties of human primary motor cortex gamma
oscillations. *Journal of Neurophysiology*, *104*(5), 2873-2885.

Kriegeskorte, N., & Douglas, P. K. (2019). Interpreting encoding and decoding models. *Current*
Opinion in Neurobiology, *55*, 167-179.

Weichert, F., Bachmann, D., Rudak, B., & Fisseler, D. (2013). Analysis of the accuracy and
robustness of the leap motion controller. *Sensors*, *13*(5), 6380-6393.

Suway, S. B., & Schwartz, A. B. (2019). Activity in Primary Motor Cortex Related to Visual
Feedback. *Cell Reports*, *29*(12), 3872-3884.

Stolk, A., Griffin, S., Van der Meij, R., Dewar, C., Saez, I., Lin, J. J., ... & Oostenveld, R. (2018).
Integrated analysis of anatomical and electrophysiological human intracranial data. *Nature*
Protocols, *13*(7), 1699-1723.

Canolty, R. T., Edwards, E., Dalal, S. S., Soltani, M., Nagarajan, S. S., Kirsch, H. E., ... & Knight, R. T.
(2006). High gamma power is phase-locked to theta oscillations in human
neocortex. *Science*, *313*(5793), 1626-1628.

Pei, X., Leuthardt, E. C., Gaona, C. M., Brunner, P., Wolpaw, J. R., & Schalk, G. (2011).
Spatiotemporal dynamics of electrocorticographic high gamma activity during overt and covert
word repetition. *Neuroimage*, *54*(4), 2960-2972

Hoerl, A. E., & Kennard, R. W. (1970). Ridge regression: Biased estimation for nonorthogonal
problems. *Technometrics*, *12*(1), 55-67.

Anderson, M., & Braak, C. T. (2003). Permutation tests for multi-factorial analysis of variance.
Journal of statistical computation and simulation, 73(2), 85-113.

Churchland, M. M., Cunningham, J. P., Kaufman, M. T., Foster, J. D., Nuyujukian, P., Ryu, S. I., &
Shenoy, K. V. (2012). Neural population dynamics during reaching. *Nature*, *487*(7405), 51-56.

Elsayed, G. F., Lara, A. H., Kaufman, M. T., Churchland, M. M., & Cunningham, J. P. (2016).
Reorganization between preparatory and movement population responses in motor
cortex. *Nature Communications*, *7*(1), 1-15.

Schaffer, J. E., Maenza, C., Good, D. C., Przybyla, A., & Sainburg, R. L. (2020). Left hemisphere
damage produces deficits in predictive control of bilateral coordination. *Experimental Brain*
Research, *238*(12), 2733-2744.
1063
Diedrichsen, J., Wiestler, T., & Krakauer, J. W. (2013). Two distinct ipsilateral cortical
representations for individuated finger movements. *Cerebral Cortex*, *23*(6), 1362-1377.

Przybylski, Ł., & Króliczak, G. (2017). Planning functional grasps of simple tools invokes the hand-
independent praxis representation network: an fMRI study. *Journal of the International*
Neuropsychological Society, *23*(2), 108-120.

Buxbaum, L. J., Kyle, K. M., Tang, K., & Detre, J. A. (2006). Neural substrates of knowledge of
hand postures for object grasping and functional object use: Evidence from fMRI. *Brain*
Research, *1117*(1), 175-185.

Xu J, Gannon PJ, Emmorey K, Smith JF, Braun A. Symbolic gestures and spoken language are
processed by a common neural system. Proc Natl Acad Sci U S A 2009, 106:20665–20669.

Bohlhalter, S., Hattori, N., Wheaton, L., Fridman, E., Shamim, E. A., Garraux, G., & Hallett, M.
(2009). Gesture subtype–dependent left lateralization of praxis planning: An event-related fMRI
study. *Cerebral Cortex*, *19*(6), 1256-1262.

Ochipa C, Rothi LJG, Heilman KM (1989): Ideational apraxia—A deficit in tool selection and use.
Ann Neurol 25:190–193.

Vingerhoets, G., Acke, F., Alderweireldt, A. S., Nys, J., Vandemaele, P., & Achten, E. (2012).
Cerebral lateralization of praxis in right-and left-handedness: Same pattern, different
strength. *Human Brain Mapping*, *33*(4), 763-777.

Vingerhoets, G., Alderweireldt, A. S., Vandemaele, P., Cai, Q., Van der Haegen, L., Brysbaert, M.,
& Achten, E. (2013). Praxis and language are linked: evidence from co-lateralization in individuals
with atypical language dominance. *Cortex*, *49*(1), 172-183.

Vingerhoets, G., Acke, F., Alderweireldt, A. S., Nys, J., Vandemaele, P., & Achten, E. (2012).
Cerebral lateralization of praxis in right-and left-handedness: Same pattern, different
strength. *Human Brain Mapping*, *33*(4), 763-777.

Jäncke, L., Peters, M., Himmelbach, M., Nösselt, T., Shah, J., & Steinmetz, H. (2000). fMRI study
of bimanual coordination. *Neuropsychologia*, *38*(2), 164-174.

Toyokura, M., Muro, I., Komiya, T., & Obara, M. (1999). Relation of bimanual coordination to
activation in the sensorimotor cortex and supplementary motor area: analysis using functional
magnetic resonance imaging. *Brain Research Bulletin*, *48*(2), 211-217.

Serrien, D. J., Cassidy, M. J., & Brown, P. (2003). The importance of the dominant hemisphere in
the organization of bimanual movements. *Human Brain Mapping*, *18*(4), 296-305.

Maki, Y., Wong, K. F. K., Sugiura, M., Ozaki, T., & Sadato, N. (2008). Asymmetric control
mechanisms of bimanual coordination: an application of directed connectivity analysis to
kinematic and functional MRI data. *Neuroimage*, *42*(4), 1295-1304.

Cleland, B. T., & Madhavan, S. (2021). Ipsilateral motor pathways to the lower limb after stroke:
Insights and opportunities. *Journal of Neuroscience Research*, *99*(6), 1565-1578.
1100
1101   Swinnen, S. P., Vangheluwe, S., Wagemans, J., Coxon, J. P., Goble, D. J., Van Impe, A., ... &
1102   Wenderoth, N. (2010). Shared neural resources between left and right interlimb coordination
1103   skills: the neural substrate of abstract motor representations. *Neuroimage*, *49*(3), 2570-2580.
1104
1105   Fujiyama, H., Van Soom, J., Rens, G., Cuypers, K., Heise, K. F., Levin, O., & Swinnen, S. P. (2016).
1106   Performing two different actions simultaneously: The critical role of interhemispheric
1107   interactions during the preparation of bimanual movement. *Cortex*, *77*, 141-154.
1108
1109   Bohannon, R. W., Smith, M. B., & Larkin, P. A. (1986). Relationship between independent sitting
1110   balance and side of hemiparesis. *Physical therapy*, *66*(6), 944-945.
1111
1112   Spinazzola, L., Cubelli, R., & Della Sala, S. (2003). Impairments of trunk movements following left
1113   or right hemisphere lesions: dissociation between apraxic errors and postural
1114   instability. *Brain*, *126*(12), 2656-2666.
1115
1116   Guiard, Y. (1987). Asymmetric division of labor in human skilled bimanual action : The kinematic
1117   chain as a model. *Journal of Motor Behavior, 19*, 486-517.

Bresson, F., Maury, L., Pieraut-Le Bonniec, G., & de Schonen, S. (1977). Organization and lateralization of reaching in infants: An instance of asymmetric functions in hands collaboration. *Neuropsychologia*, *15*(2), 311-320.

Kaufman, M. T., Seely, J. S., Sussillo, D., Ryu, S. I., Shenoy, K. V., & Churchland, M. M. (2016). The largest response component in the motor cortex reflects movement timing but not movement type. *Eneuro*, *3*(4).

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models using lme4. *arXiv preprint arXiv:1406.5823*.

Lee, O. E., & Braun, T. M. (2012). Permutation tests for random effects in linear mixed models. *Biometrics*, *68*(2), 486-493.

Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., ... & Killiany, R. J. (2006). An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. Neuroimage, 31(3), 968-980.

Leocani, L., Cohen, L. G., Wassermann, E. M., Ikoma, K., & Hallett, M. (2000). Human corticospinal excitability evaluated with transcranial magnetic stimulation during different reaction time paradigms. Brain, 123(6), 1161-1173.

Liepert, J., Dettmers, C., Terborg, C., & Weiller, C. (2001). Inhibition of ipsilateral motor cortex during phasic generation of low force. Clinical Neurophysiology, 112(1), 114-121.

Schoppe, O., Harper, N. S., Willmore, B. D., King, A. J., & Schnupp, J. W. (2016). Measuring the performance of neural models. Frontiers in computational neuroscience, 10, 10.

Hsu, A., Borst, A., & Theunissen, F. E. (2004). Quantifying variability in neural responses and its application for the validation of model predictions. Network: Computation in Neural Systems, 15(2), 91-109.

Lage-Castellanos, A., Valente, G., Formisano, E., & De Martino, F. (2019). Methods for computing the maximum performance of computational models of fMRI responses. PLoS computational biology, 15(3), e1006397.

Nunez-Elizalde, A. O., Huth, A. G., & Gallant, J. L. (2019). Voxelwise encoding models with non-spherical multivariate normal priors. *Neuroimage*, *197*, 482-492.

Zimnik AJ, Churchland MM. Independent generation of sequence elements by motor cortex. Nat Neurosci. 2021 Mar;24(3):412-424. doi: 10.1038/s41593-021-00798-5.