Cytosolic aggregation of mitochondrial proteins disrupts cellular homeostasis by

stimulating the aggregation of other proteins

Urszula Nowicka[1, 2, 3], Piotr Chroscicki[2, 4, 7], Karen Stroobants[5, 7], Maria Sladowska[2, 4],

Michal Turek[1, 2, 4], Barbara Uszczynska-Ratajczak[2, 6], Rishika Kundra[5], Tomasz Goral[1,

[2], Michele Perni[5], Christopher M. Dobson[5], Michele Vendruscolo[5], & Agnieszka

Chacinska[1, 2, 3,]*

[1]ReMedy International Research Agenda Unit, University of Warsaw, Warsaw, Poland

[2]Centre of New Technologies, University of Warsaw, Warsaw, Poland

[3]IMol Polish Academy of Sciences, Warsaw, Poland

[4]International Institute of Molecular and Cell Biology, Warsaw, Poland

[5]Centre for Misfolding Diseases, Department of Chemistry, University of Cambridge,

Cambridge, United Kingdom

[6]Institute of Bioorganic Chemistry, Polish Academy of Sciences, Poznan, Poland

[7]These authors contributed equally to this work.

*Correspondence should be directed to a.chacinska@imol.institute

Abstract

Mitochondria are organelles with their own genomes, but they rely on the import of nuclear-encoded proteins that are translated by cytosolic ribosomes. Therefore, it is important to understand whether failures in the mitochondrial uptake of these nuclear-encoded proteins can cause proteotoxic stress and identify response mechanisms that may counteract it. Here, we report that upon impairments in mitochondrial protein import, high-risk precursor and immature forms of mitochondrial proteins form aberrant deposits in the cytosol. These deposits then cause further cytosolic accumulation and consequently aggregation of other mitochondrial proteins and disease-related proteins, including α-synuclein and amyloid β. This aggregation triggers a cytosolic protein homeostasis imbalance that is accompanied by specific molecular chaperone responses at both the transcriptomic and protein levels. Altogether, our results provide evidence that mitochondrial dysfunction, specifically protein import defects, contributes to impairments in protein homeostasis, thus revealing a possible molecular mechanism by which mitochondria are involved in neurodegenerative diseases.

Introduction

Although over one thousand proteins are utilized by mitochondria to perform their functions, only ~1% of them are synthesized inside this organelle. The majority of mitochondrial proteins are synthesized in the cytosol and need to be actively transported to mitochondria, a process that occurs via a sophisticated system that involves protein translocases and sorting machineries (Calvo et al., 2016; Morgenstern et al., 2017; Neupert & Herrmann, 2007; Pfanner et al., 2019). The consequences of mitochondrial protein import defects on cellular proteostasis can be severe and currently some response mechanisms are identified (Boos et al., 2019; Izawa et al., 2017; Kim et al., 2016; Martensson et al., 2019; Priesnitz & Becker, 2018; Wang & Chen, 2015; Weidberg & Amon, 2018; Wrobel et al., 2015; Wu et al., 2019; Poveda-Huertes et al., 2020). Mitochondrial dysfunction is closely associated with neurodegenerative disorders, and such mitochondrial defects as aberrant Ca^{2+} handling, increases in reactive oxygen species, electron transport chain inhibition, and impairments in endoplasmic reticulum-mitochondria tethering are well described pathological markers (Cabral-Costa & Kowaltowski, 2020). Still unknown, however, is whether mitochondrial defects appear as a consequence of neurodegeneration, whether they contribute to it, or whether both processes occur. Disease-related proteins can interfere with mitochondrial import and the further processing of imported proteins within mitochondria (Cenini et al., 2016; Di Maio et al., 2016; Mossmann et al., 2014; Vicario et al., 2018). Furthermore, aggregated proteins can be imported into mitochondria where they can be either cleared or sequestered in specific deposit sites (Bruderek et al., 2018; Ruan et al., 2017; Sorrentino et al., 2017). However, the reverse aspect of the way in which mitochondrial dysfunction, including mitochondrial import defects, contributes to the progression of neurodegenerative diseases remains elusive. One possible mechanism may occur through alterations of cellular homeostasis, as mitochondrial

dysfunction can affect it through multiple mechanisms (Andreasson et al., 2019; Braun &

Westermann, 2017; Escobar-Henriques et al., 2020).

Impairments in mitochondrial protein import and mitochondrial import machinery

overload result in the accumulation of mitochondria-targeted proteins in the cytosol and

stimulation of mitoprotein-induced stress (Boos et al., 2019; Wang & Chen, 2015; Wrobel et

al., 2015). These findings raise the issue of whether the accumulation of mistargeted

mitochondrial proteins contributes to the progression of neurodegenerative diseases.

Additionally, unknown are whether mitoprotein-induced stress is a general response to

precursor proteins that globally accumulate in the cytosol and whether a subset of mitochondrial

precursor proteins pose particularly difficult challenges to the protein homeostasis system and

consequently contribute to the onset and progression of neurodegenerative disorders (Boos et

al., 2020; Mohanraj et al., 2020).

The analysis of a transcriptional signature of Alzheimer's disease supports the notion

that there is a subset of mitochondrial proteins that is more dangerous than others for the cell

(Ciryam et al., 2016; Kundra et al., 2017). These studies have shown that specific mitochondrial

proteins that are functionally related to oxidative phosphorylation are transcriptionally

downregulated in Alzheimer's disease. In the present study, we investigated why these proteins

are downregulated. We hypothesized that this need arises from the potential supersaturation of

these proteins, which makes them prone to aggregation (Ciryam et al., 2016; Kundra et al.,

2017). Our results showed that when some of these mitochondrial proteins remain in the cytosol

because of mitochondrial protein import insufficiency, they formed insoluble aggregates that

disrupted protein homeostasis. These proteins triggered a prompt specific molecular chaperone

response that aimed to minimize the consequences of protein aggregation. However, when this

rescue mechanism was insufficient, these aggregates stimulated the cytosolic aggregation of

other mitochondrial proteins and led to the downstream aggregation of non-mitochondrial

proteins. Our findings indicate that metastable mitochondrial proteins can be transcriptionally

downregulated during neurodegeneration to minimize cellular protein homeostasis imbalance

that is caused by their mistargeting.

Results

Metastable mitochondrial precursor proteins can aggregate in the cytosol

The analysis of a transcriptomic signature of Alzheimer's disease identified oxidative

phosphorylation as a pathway that is metastable and downregulated in the human central

nervous system (Ciryam et al., 2016; Kundra et al., 2017). This observation suggests that a

group of mitochondrial proteins might be dangerous for cellular protein homeostasis because

of their poor supersaturation and hence solubility at cellular concentrations. From the list of

genes that were simultaneously downregulated and metastable in Alzheimer's disease patients,

we selected all genes that encode mitochondrial proteins. Next, we identified genes that encode

proteins that have homologs in yeast (**Figure 1–figure supplement 1**). Based on the yeast

homolog sequence, we generated FLAG-tagged constructs that were expressed under control

of the copper-inducible promoter (CUP1). We then established a multi-centrifugation step assay

to assess whether these proteins exceed their critical concentrations and become supersaturated

when overproduced (Vecchi et al., 2020), thereby acquiring the ability to aggregate during their

trafficking to mitochondria (**Figure 1–figure supplement 2A**). We followed a FLAG peptide

signal to determine whether the protein was present in the soluble (S_{125k}) or insoluble (P_{125k})

fraction. We found that the β and g subunits of mitochondrial F_1F_O adenosine triphosphate

(ATP) synthase (Atp2 and Atp20, respectively) were present in the insoluble fraction, indicating

that they formed high-molecular-weight deposits (**Figure 1A** and **Figure 1–figure supplement**

2B). We made a similar observation for Rieske iron-sulfur ubiquinol-cytochrome *c* reductase

(Rip1). Rip1 and subunit VIII of cytochrome *c* oxidase complex IV (Cox8) had entirely

insoluble precursor (p) forms, whereas the mature (m) forms were partially insoluble (the p form of the protein is larger and migrates on the gel above the m form; **Figure 1A** and **Figure 1–figure supplement 2C**). Furthermore, the mature forms of subunit VIII of ubiquinol cytochrome *c* reductase complex III (Qcr8), core subunit of the ubiquinol-cytochrome *c* reductase complex (Cor1), and subunit β of the mitochondrial processing protease (MPP; Mas1) were partially insoluble. Only subunit VIb of cytochrome *c* oxidase (Cox12) and subunit 6 of ubiquinol cytochrome *c* reductase complex (Qcr6) were mainly present in the soluble fraction (**Figure 1A** and **Figure 1–figure supplement 2C**). Moreover, the tendency to aggregate and its harmful consequences correlated well with growth defects of yeast transformants that overexpressed mitochondrial proteins (**Figure 1B** and **Figure 1–figure supplement 2D**). We observed that the higher amount of an overproduced protein in the insoluble fraction correlated with a greater increase in lethality. This difference was still present under heat shock conditions of 37°C, indicating that the general molecular chaperone response that was associated with an increase in temperature did not compensate for the observed changes for Atp2, Cox8, or pRip1 and could only partially compensate for Cox12 and Atp20.

To test whether these metastable mitochondrial proteins aggregate in the cytosol as a result of an inefficient mitochondrial protein import system, we followed the cytosolic fate of non-imported mitochondrial proteins that contained a cleavable targeting sequence. Consistent with previous studies (Wrobel et al., 2015), cells that were treated with the chemical uncoupler carbonyl cyanide m-chlorophenyl hydrazine (CCCP) had a compromised mitochondrial inner membrane (IM) electrochemical potential and protein import failure (Chacinska et al., 2009; Neupert & Herrmann, 2007; van der Laan et al., 2010). To observe import defects without prompting negative consequences of CCCP (e.g., autophagy), the treatment time was limited to 30 min. Perturbations of IM potential resulted in the accumulation of p forms of Rip1 (pRip1), similar to when the full precursor form of Rip1 was overproduced (**Figure 1C and 1D** and

Figure 1–figure supplement 2E). We also extended the analysis of p forms of other proteins

for which antibodies were available and allowed p form detection, namely mitochondrial matrix

superoxide dismutase (Sod2) and mitochondrial malate dehydrogenase 1 (Mdh1) (**Figure 1C**

and **Figure 1–figure supplement 2E**). The accumulation of pRip1, the p form of Sod2 (pSod2),

and the p form of Mdh1 (pMdh1) increased as CCCP concentrations increased (**Figure 1C and**

1D). Next, an aggregation assay was performed to assess the fate of precursor mitochondrial

proteins under conditions of chemical impairments in mitochondrial import. pRip, pSod2, and

pMdh1 were present in the pellet fractions, demonstrating that even without overproduction

they formed insoluble aggregates in the cell (**Figure 1E**). We also tested conditions of defective

protein import by using a temperature-sensitive mutant of TIM23 presequence translocase,

pam16-3 (Wrobel et al., 2015). pRip1 and pSod2 accumulated (**Figure 1–figure supplement**

2F) and consequently formed insoluble aggregates under these conditions (**Figure 1F**).

Therefore, the mistargeted precursor forms of mitochondrial proteins aggregated, when their

concentrations exceeded their solubility limits due to their overproduction, when the

mitochondrial import defect was stimulated chemically, and when mutants of the presequence

translocase import pathway were used.

Mitochondrial protein aggregation stimulates a cytosolic molecular chaperone response

154   To further investigate the cellular response to the accumulation of mitochondrial

precursor forms in the cytosol, we investigated global transcriptomic changes that were

triggered by pRip1 overproduction and *pam16-3* mutation (**Figure2-Source Data 1-4, Figure2**

and Figure2–figure supplement 1 and 2). pRip1 overexpression triggered small and rather

specific transcriptomic changes, in which a total of nine genes (**Figure 2C**) were upregulated

and four were downregulated (see also **Figure 2A** and **Figure2–figure supplement 1A**). The

Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment analysis indicated that the

oxidative phosphorylation pathway was upregulated, as expected, for pRip1 overexpression

(**Figure2–figure supplement 1B**). After 4 h of pRpi1 expression, the ATP-binding cassette

(ABC) transporters appeared as the second most upregulated KEGG pathway (**Figure2–figure**

supplement 1B). For pRip1, the largest fold change was observed for the *SSA1*, *SSA4*, and

HSP82 chaperones as a likely response to pRip1 aggregation (**Figure 2A** and **2C**). Extended

pRip1 overproduction started to affect cellular performance as the following genes became

downregulated: *POT1* (a peroxisomal protein that is also localized to the mitochondrial IM

space and involved in fatty acid β-oxidation), *DLD3* (involved in lactate biosynthesis), *AGP3*

(amino acid transmembrane transporter), *PDC6* (pyruvate decarboxylase)(**Figure 2C**).

Next, since we planned to use a temperature-sensitive allele, we wanted to exclude

possible effects that arise merely from a temperature shift. For that purpose, we analyzed the

effect of heat shock individually on wild-type *pam16-3* (referred to as *WT*) and the *pam16-3*

mutant. For WT the increase in temperature resulted in a consistent drop in the amount of up-

and downregulated genes (**Figure2–figure supplement 2A**). For the *pam16-3* mutant, the

number of up- and downregulated genes was nearly constant (**Figure2–figure supplement 2B**),

suggesting an unchanging need for adaptation. To determine the function of genes whose

expression changed in response to the temperature shift, we performed KEGG enrichment

analysis for WT and the *pam16-3* mutant. In both cases, we observed the upregulation of genes

that are associated with ribosomes and ribosome biogenesis (**Figure2–figure supplement 2C**

and 2D). Unsurprisingly, the higher temperature also affected metabolic pathways with the

longer exposure of *pam16-3* yeast cells to 37°C (Wrobel et al., 2015).

Finally, we studied effects of *pam16-3* mutations alone. For each heat shock time,

pam16-3 was compared with WT at the same time of exposure to 37°C. With regard to

mitochondrial defects, the *pam16-3* mutation had a stronger effect on the transcriptome than

the effect of pRip1 (**Figure2–figure supplement 1A**). Based on the KEGG analysis and as

expected, oxidative phosphorylation-, metabolic pathway-, and citrate cycle-related genes were

downregulated (**Figure2–figure supplement 1C**). More than half of the genes that were

affected encode mitochondrial proteins (Morgenstern et al., 2017) (**Figure2–figure**

supplement 1D). We also tested whether any groups of genes had related functions but were

not enriched in the KEGG analysis. We analyzed all genes that passed a 5% false discovery rate

(FDR) cut-off and were non-mitochondrial. From this list, we selected genes for which we could

identify at least 10 that shared similar functions. Based on this analysis, we noticed that a group

of genes that encode molecular chaperones was upregulated in the *pam16-3* mutant (**Figure**

2D). We observed the upregulation of *HSP82* and *SSA4*, which were also upregulated in the

case of pRip1 overexpression. We also observed the upregulation of other molecular chaperone

genes, including *SNO4*, *HSP26*, *HSP42*, *SSE2*, *HSP12*, *HSP32*, *HSP104*, and *HSP33* (**Figure**

2B, blue, and **2D**). The effect of the *pam16-3* mutant could be both direct and indirect when

considering the various pathways in which these molecular chaperones are involved. We used

the temperature-sensitive *pam16-3* mutant; therefore, we investigated the degree to which

mitochondrial dysfunction exceeded the change that was attributable to heat shock treatment

alone for each of these molecular chaperones. We compared changes in WT samples at both

19°C and 37°C and matched them to corresponding changes that were attributable to the

pam16-3 mutation (normalized to WT at the same heat shock time) (**Figure2–figure**

supplement 1E and Supplementary file 1). Our analysis showed that the molecular chaperone

response in *pam16-3* went beyond the changes that were expected solely for the high-

temperature treatment. The most pronounced changes were observed for *SSE2*, *SIS1*, *HSP42*,

and *HSP104* at 4 h of treatment (**Figure2–figure supplement 1E,** black arrows). These findings

suggest that impairments in mitochondrial protein import that originate from defects of

translocases had to some extent similar, but not identical consequences as clogging the import

sites (Boos et al., 2019). To compare the pRip1 data with the effects of the *pam16-3* mutation,

we investigated whether ABC transporter genes would also be upregulated. The analysis of significantly altered genes showed that as the heat shock time was extended, *PDR5* gene levels changed, as well as other ABC transporter genes, including *PDR15* and *SNQ2* (**Figure2–figure supplement 1F**). *PDR5* is controlled by PDR3 transcription factor, the levels of which also increased (Gulshan et al., 2008). All of these genes were shown to be involved in the mitochondrial compromised protein import response mitoCPR response (Weidberg & Amon, 2018). When we extended our analysis to other mitoCPR-related genes, we observed their upregulation beginning at 4 h of treatment (**Figure2–figure supplement 1F**).

Effects of the Hsp42 and Hsp104 chaperone response to mitochondrial import failure at the protein level

Next, we assessed whether the observed molecular chaperone upregulation would also be observed at the protein level. We used CCCP to stimulate mitochondrial dysfunction to avoid the temperature factor at initial screening. We monitored changes in protein levels for all molecular chaperones against which we had antibodies available at the time of the studies, including Hsp104, Ssa1, Ssb1, Ssc1, Hsp60, and Edg1. No changes were observed for most of them, but a significant response was observed for Hsp104 (**Figure 3A** and **3B**). Notably, Hsp104 upregulation at the protein level was observed already after 15 min of treatment (**Figure3–figure supplement 1A**). We then tested whether Hsp104 was upregulated in response to mitochondrial import defects that were caused by the *pam16-3* mutation, along with the established *pam16-1* and *pam18-1* mutants and mitochondrial intermembrane space import and assembly (MIA) import pathway mutations *mia40-4int* and *mia40-3* (Wrobel et al., 2015). For all of the mutants under permissive conditions (19°C; i.e., when only a mild protein import induction phenotype occurs), we observed the upregulation of Hsp104 at the protein level (**Figure 3C**). This effect was still present at the restrictive temperature of 37°C, but the

differences were less evident because of the activation of heat shock responses (**Figure3–figure**

supplement 1B). We also tested whether Hsp42 would be upregulated at the protein level

because its upregulation was one of the most prominent, based on transcriptomic data for the

pam16-3 mutant. Because of the lack of an antibody against Hsp42, we used a yeast strain with

hsp42 that was tagged with green fluorescent protein (GFP) that allowed us to follow Hsp42

levels by monitoring the GFP signal. We could only test the consequences of impairments in

mitochondrial import that were caused by CCCP treatment in this experimental setup.

Consistent with the transcriptomic *pam16-3* data, we observed an increase in the abundance of

Hsp42 (**Figure 3D**). After identifying Hsp42 and Hsp104 as two molecular chaperones that

change significantly when mitochondrial import is impaired, we further examined their

expression levels when metastable mitochondrial proteins were overproduced. Hsp104 levels

were significantly upregulated upon such protein overproduction (**Figure 3E** and **3F**). Hsp42

levels were again analyzed in the background of the *hsp42-GFP* strain. Here, we also observed

an increase in the abundance of Hsp42 for all metastable proteins (**Figure 3G** and **3H**). In the

case of pRip1, this response was not observed at the transcriptome level but was significant at

the protein level (**Figure 2C vs. Figure 3G** and **3H**). Thus, the upregulation of specific

molecular chaperones in response to cytosolic mitochondrial protein aggregation could be

achieved both transcriptionally and posttranscriptionally. In the case of *pam16-3* mutant the

transcriptomic response involved a number of chaperone genes, whereas on the protein level

we observed a significant response of the aggregate-related chaperones, Hsp42 and Hsp104.

Interestingly, in the case of pRip1 overproduction, Hsp42 and Hsp104 were increased without

any transcriptional stimulation, meaning that in the case of mitochondrial precursor aggregates

the posttranscriptional response can be uncoupled from transcription.

In yeast, different types of aggregates have been identified (Sontag et al., 2017;

Tyedmers et al., 2010). Both Hsp42 and Hsp104 are molecular chaperones that are involved in

aggregate handling in inclusion bodies (Balchin et al., 2016; Mogk et al., 2015; Mogk et al.,

2019). To assess whether Hsp42 and Hsp104 upregulation is associated with a response to

protein aggregation, we tested whether they can recognize metastable mitochondrial protein

aggregates. We labelled FLAG-tagged metastable proteins with the Alexa 568 fluorophore and

followed its co-localization with Hsp42-GFP by confocal microscopy. We observed the strong

accumulation of pRip1 in the form of inclusion body-like large deposits in the cytosol (**Figure**

3I). These deposits co-localized with the GFP signals, indicating that Hsp42 sequestered pRip1

aggregates. Next, we reasoned that if these specific molecular chaperones were responsible for

preventing the aggregation of metastable proteins, then we should observe a change in the

amount of accumulated proteins when they are not present in the cell. In cells with *hsp104*

deletion, we observed an increase in the accumulation of pRip1 upon CCCP treatment (**Figure**

3J). Thus, Hsp104 function was essential for limiting aggregates of mitochondrial precursors.

Based on the aggregation assay, we detected Hsp104 in the pellet fraction at higher amounts

than without CCCP treatment, suggesting that an additional amount of Hsp104 associated with

aggregates (**Figure 3K**). The lack of Hsp42 did not result in the upregulation of pRip1, pSod2,

or pMdh1 (**Figure3–figure supplement 2A and 2B**). However, the *Δhsp42* strain exhibited the

significant upregulation of Hsp104, suggesting a compensatory mechanism that diminishes

negative consequences of Hsp42 deletion (**Figure 3L and 3M**). Next, we tested whether the

overexpression of Hsp104 and Hsp42 exerts beneficial effects when mitochondrial import

defects are stimulated. Hsp104 overproduction resulted in lower levels of pRip1, pSod2, and

pMdh1 upon CCCP treatment (**Figure 3N** and **Figure3–figure supplement 2C**). In this case,

we again observed a link between Hsp42 and Hsp104 levels, in which higher levels of Hsp104

were associated with lower levels of Hsp42 in the cell (**Figure 3N** and **Figure3–figure**

supplement 2D). The overexpression of Hsp42 did not have such an effect on pRip1, pSod2,

or pMdh1, in which levels of p forms did not change (**Figure3–figure supplement 2E and 2F**).

Finally, we tested whether higher amounts of Hsp42 or Hsp104 exert positive effects on cells

that overproduce metastable proteins. Despite the beneficial effect of Hsp104 on precursor

aggregate levels, we did not observe the rescue of growth defects that were observed for strains

that expressed Atp2, Cox8, Cox12, pRip1, or Atp20 when either Hsp104 or Hsp42 was

overproduced (**Figure3–figure supplement 2G**). Altogether, our results show that the

aggregate-related chaperone response is activated in the cell to handle consequences of

cytosolic accumulation of mitochondrial precursor proteins.

Mitochondrial protein import failure impairs cellular protein homeostasis

We next investigated whether the presence of metastable and aggregation-prone

mitochondrial precursors initiates the accumulation of p forms of other mitochondrial proteins.

We verified that this was the case for some of the metastable proteins. The overproduction of

Atp2 and Cox8 resulted in the concurrent accumulation of other mitochondrial proteins, pRip1

and pSod2, in the total cell fractions, without any other stimulation beyond just the

overexpression of metastable proteins (**Figure 4A-C** and **Figure4–figure supplement 1A**).

Atp20 overproduction also stimulated a significant increase in pSod2 but not pRip1 in the total

cell fractions (**Figure 4A-C**). We hypothesized that differences in pRip1 and pSod2

accumulation for analyzed metastable proteins may reflect the differences: in the mitochondria

import rates of these metastable proteins, in their relative cytosolic abundance, and in the

protein sequence affecting their aggregation dynamics. Nevertheless, Rip1, Sod2, and Mdh1

co-aggregated with Atp2 and Cox8 metastable proteins in the insoluble fraction, based on the

aggregation assay analysis (**Figure 4D,** and **Figure4–figure supplement 1B**). Thus, the greater

abundance of aggregation-prone mitochondrial precursors resulted in the progression of

mitochondrial protein import defect and consequently the larger cytosolic aggregation of

mitochondrial precursors. This observation may justify why Atp2, Cox8, and Atp20 exhibited

the most pronounced viability decrease, demonstrated by the drop-test results (**Figure 1B**).

Moreover, this process was partially counteracted by proteasome-mediated degradation.

MG132-mediated protein inhibition led to an increase in the accumulation and aggregation of

mitochondrial precursor proteins (**Figure 4E,** and **Figure4–figure supplement 1C**).

315   Finally, we tested whether a chain reaction may occur, wherein the cytosolic aggregation

of mitochondrial proteins first impaired the import efficiency and solubility of other

mitochondrial precursors and subsequently induced the downstream aggregation of other non-

mitochondrial proteins. We took advantage of α-synuclein (α-Syn) a model protein of

neurodegenerative disorders. α-synuclein (α-Syn) WT and two of its mutations, A30P and

A53T are implicated in Parkinson's disease. α-Syn WT and A53T, but not A30P share a similar

cellular distribution when they are expressed, which allows monitoring the formation of their

aggregates in the cell (Outeiro & Lindquist, 2003). Therefore, we used these two model systems

that were tagged with GFP in our studies. Confocal microscopy was used to monitor changes

in the average number of α-Syn WT and A53T aggregates per cell by following the signal of

the GFP tag in response to impairments in mitochondrial import. Here, mitochondrial defects

were stimulated by the *pam16-3* mutant. The average number of α-Syn aggregates per cell

increased for *pam16-3* at a permissive temperature of 19°C for both α-Syn WT-GFP and A53T-

GFP compared with WT strain (**Figure 4F** and **4G,** and **Figure4–figure supplement 1D** and

1E). The increase in the average number of aggregates continued to grow for α-Syn WT-GFP

at the restrictive temperature of 37°C (**Figure 4F**). The average number of α-Syn A53T-GFP

aggregates was higher at 37°C than at the permissive temperature and did not significantly

increase upon the mitochondrial import defect in the *pam16-3* mutant (**Figure 4G**). This

observation might be justified by the different morphology of WT and A53T α-Syn aggregates.

The A53T mutation of α-Syn resulted in larger aggregates compared with α-Syn WT (Outeiro

& Lindquist, 2003). To test this hypothesis, we investigated the size of aggregates of α-Syn WT

and A53T. We did not observe any difference in the average size of aggregates for α-Syn WT

at 19°C and 37°C for both WT and *pam16-3* mutant (**Figure4–figure supplement 1F**). In the

case of α-Syn A53T, puncta were larger and better resolved than α-Syn WT puncta (**Figure**

4G) in both strains. We observed an increase in the average α-Syn A53T aggregate size for WT

and the *pam16-3* mutant at 37°C (**Figure4–figure supplement 1G**) when compared to 19°C.

These findings suggest that for α-Syn A53T at 37°C, the average number of aggregates did not

significantly change after reaching a threshold and instead they were becoming larger as the

aggregation progressed. Finally, we observed the higher accumulation of pRip at combined

conditions of a *pam16-3*-stimulated mitochondrial import defect and α-Syn aggregation at 37°C

(**Figure 4H** and **4I,** and **Figure4–figure supplement 1D and 1E**). Thus, the protein

aggregation linked to mitochondrial import defects had a dual effect: (*i*) α-Syn aggregation was

stimulated, and (*ii*) α-Syn aggregation further deepened the aggregation of mitochondrial

precursor proteins.

Our results indicated that the mitochondrial precursor aggregation in the cytosol led to

an increase in the aggregation of other non-mitochondrial proteins, and this acceleration of

protein aggregation, which may be compared to a snowball effect, reduced cellular protein

homeostasis.

Mitochondrial dysfunction results in protein aggregation in *Caenorhabditis elegans*

To assess whether mitochondrial precursor aggregation that is caused by mitochondrial

protein import deficiency compromises protein homeostasis at the organismal level, we used

Caenorhabditis elegans as a model system. We used an RNA interference (RNAi) approach to

silence *dnj-21* (a homolog of yeast Pam18) in *C. elegans*. The depletion of DNJ-21 stimulates

the mitochondrial import defect similarly to the defect that is observed in yeast for *pam16-3*.

We first monitored the aggregation of two model proteins in cytosol, red fluorescent protein

(RFP) and GFP, in the transgenic strain that expressed RFP and GFP in the body wall muscle to assess whether the RNAi silencing of *dnj-21* in early adulthood in *C. elegans* is sufficient to stimulate their aggregation (**Figure 5A-C** and **Figure5–figure supplement 1A**). We found that the RNAi of *dnj-21* increased the cytosolic aggregation of both GFP and RFP model proteins. Therefore, we next tested whether changes in cellular homeostasis that are attributable to mitochondrial defects affect the health of *C. elegans* when α-Syn is produced. The silencing of *dnj-21*, accompanied by α-Syn expression, decreased worm fitness, manifested by a slower speed and fewer bends compared with the effect of *dnj-21* silencing alone (**Figure 5D**). Finally, we investigated whether the link between mitochondrial dysfunction is only limited to α-Syn, or whether similar effects are observed for other proteins that are linked to neurodegenerative diseases. For this reason, we analyzed if mitochondrial dysfunction enhances amyloid β (Aβ) aggregation in *C. elegans*. We used worms that carried Aβ peptides in body wall muscles and exhibited paralysis in adults when the temperature shifted to 25°C (**Figure5–figure supplement 1B**) (Sorrentino et al., 2017). The level of Aβ aggregates increased at 22°C when *dnj-21* was silenced (**Figure 5E and 5F** and **Figure5–figure supplement 1C and 1D**). This stimulated aggregation resulted from the mitochondrial import defect without any accompanying changes in the expression of Aβ peptide (**Figure5–figure supplement 1E** to **1G**). We found that Aβ aggregation was accompanied by a decrease in worm motility with *dnj-21* silencing (**Figure 5G,** and **Figure5–figure supplement 1H**). Overall, our results indicate that the mechanism of aggregation stimulation that is caused by mitochondrial dysfunction is conserved between species, as demonstrated for both α-Syn and Aβ aggregates.

Discussion

The present study showed that a group of mitochondrial proteins that are downregulated in Alzheimer's disease (i.e., Rip1, Atp2, Cox8, and Atp20) can aggregate in the cytosol and that

the overexpression of these proteins upregulates Hsp42 and Hsp104, two molecular chaperones

that are associated with inclusion bodies (Balchin et al., 2016; Mogk et al., 2015; Mogk et al.,

2019). Depending how mitochondrial precursor protein aggregation is stimulated, the Hsp42

and Hsp104 upregulation is modulated transcriptionally and posttranscriptionally, with a

possibility of posttranscriptional stimulation being a dominant component. Thus, our findings

of modulated aggregate-related chaperones upregulation further contribute to the transcriptomic

response mechanisms involving chaperones due to mitochondrial import machinery overload

(Boos et al., 2019), and mechanism of proteasomal protein degradation machinery upregulation

(Wrobel et al., 2015). We show that stress responses which are induced by mitochondrial

proteins, such as the response that was identified herein, mitigate the danger that is related to

the aberrant formation of aggregates by metastable mitochondrial precursor proteins.

Our study demonstrated that mitochondrial import defects, triggered either chemically

or by mutations of the TIM23 translocase, resulted in the accumulation and aggregation of

mitochondrial precursor proteins. Consistent with the experiments that utilized the

overproduction of metastable mitochondrial proteins, an Hsp42- and Hsp104-specific

molecular chaperone response was also triggered in the cell with mitochondrial import defects.

Additionally, the ubiquitin-proteasome system could mitigate this process to some extent

(Wrobel et al., 2015).

The unresolved aggregation of mistargeted metastable mitochondrial precursor proteins

eventually resulted in a chain reaction that led to the accumulation of deposits that were formed

by other non-mitochondrial proteins. The collapse of cellular homeostasis resulted in an

increase in pRip1 and α-Syn aggregation when mitochondrial import was impaired. More

generally, we found that an increase in the aggregation of proteins linked to age-related

degenerations under conditions of mitochondrial protein import dysfunction was a conserved

cellular mechanism that was observed in both yeast and *C. elegans*. We found that the

aggregation of α-Syn and Aβ increased because of mitochondrial dysfunction at the organismal

level, accompanied by a decrease in worm fitness.

Several stress response pathways have recently been identified that counteract defects

of mitochondrial protein import (Boos et al., 2019; Izawa et al., 2017; Kim et al., 2016;

Martensson et al., 2019; Priesnitz & Becker, 2018; Wang & Chen, 2015; Weidberg & Amon,

2018; Wrobel et al., 2015; Wu et al., 2019; Poveda-Huertes et al., 2020). Unknown, however,

is whether they act independently or whether concurrent actions of all of them are required to

secure balanced cellular protein homeostasis. The cytosolic responses, which are aiming to clear

clogged translocase of the outer membrane (TOM) and to clear the precursor proteins prior to

their import through proteasomal activity, are accompanied by the aggregate-specific molecular

chaperone response identified in the present study. In addition, recent findings suggest that

some precursor proteins can accumulate within mitochondria to form insoluble aggregates

(Poveda-Huertes et al., 2020). We postulate that when these defense mechanisms may not be

sufficient to prevent mitochondrial precursor proteins aggregation in the cytosol, this

aggregation will lead to the snowball effect, where aggregation of mitochondrial proteins

stimulates further cytosolic aggregation of other proteins. These findings also indicate that

precursor protein-induced stress responses are orchestrated to complement each other (Boos et

al., 2020; Mohanraj et al., 2020).

In conclusion, our findings suggest a model in which a cascade of events that is triggered

by the cytosolic aggregation of specific metastable mitochondrial proteins contributes to the

collapse of cellular protein homeostasis accelerates the aggregation of other proteins, including

some that are hallmarks of age-related degeneration. Consequently, defects in mitochondrial

function, which are commonly observed during ageing and in neurodegeneration, trigger a

vicious cycle of protein aggregation. This notion is essential for understanding cellular events

that contribute to the onset and progression of neurodegenerative processes. Overall, our results

illustrate an important aspect of the interdependence of mitochondrial fitness and cellular

protein homeostasis systems, with direct relevance to common neurodegenerative disorders and

mitochondrial function. Our findings may suggest new avenues for therapeutic interventions to

cure or prevent diseases that are linked to protein aggregation and mitochondrial dysfunction.

Materials and methods

Experimental Design. No statistical methods were used to predetermine sample size. The

experiments were not randomized. The investigators were not blinded to allocation during the

experiments and outcome assessment. All of the experiments were repeated at least three times.

For experiments with a larger number of biological replicates, the *n* is indicated.

Yeast strains. The *Saccharomyces cerevisiae* strains were derivatives of YPH499 (MATa,

ade2-101, his3-D200, leu2-D1, ura3-52, trp1-D63, lys2-801) (524) or BY4741 (MATa, his3D1,

leu2D0, met15D0, ura3D0) (755). The descriptions of WT yeast cells refer to the BY4741 strain

unless otherwise indicated. For the inducible expression of FLAG-tagged metastable proteins,

the amino acid sequences were amplified by polymerase chain reaction (PCR) from yeast

genomic DNA. The resulting DNA fragments were cloned in frame with the FLAG tag using

the oligonucleotides that are indicated in **Supplementary file 2** into the pESC-URA vector

(Agilent), in which the GAL10 and GAL1 promoters were replaced with the Cup1 promoter.

This procedure yielded the pPCh17 (481p), pPCh18 (482p), pPCh19 (484p), pPCh20 (485p),

pPCh21 (486p), pPCh22 (487p), pPCh23 (488p), pPCh24 (489p), pPCh26 (492p), pPCh29

(490p), pPCh30 (483p), and pPB36.1 (471p) plasmids (Kowalski et al., 2018), with a FLAG

tag at the C-terminus and expressed under control of the Cup1 promoter (**Supplementary file

3**). The plasmid that expressed Hsp104 under the endogenous promoter in pFL46L vector was

provided by Prof. Magdalena Boguta (IBB PAN) (Chacinska et al., 2000). Yeast cells were

transformed according to the standard procedure. The *Δhsp104* (1001) deletion yeast strain was

purchased from Euroscarf. The *hsp42-GFP* (1005) yeast strain was purchased from Invitrogen.

The Hsp42-expressing strain on the BY4741 (1178) background under the Gal1 promoter was

provided by the Dr. Bernd Bukau laboratory (ZMBH Heidelberg). The temperature-sensitive

pam16-1 (YPH-BG-mia1-1) (733) and *pam16-3* (YPH-BGmia1-3) (734) and corresponding

pam16-WT (736) were described previously (Frazier et al., 2004). The temperature-sensitive

pam18-1 (YPH-BG-Mdj3-66) (739) and its respective *pam18-WT* (738) were described

previously (Truscott et al., 2003), similar to *mia40-3* (YPH-BGfomp2-8) (178) and *mia40-4int*

(305) and the corresponding *mia40-WT* (398) (Chacinska et al., 2004; Stojanovski et al., 2008).

Yeast growth, treatments, and analysis. In the experiments in which metastable proteins were

overproduced, the strains were grown on minimal synthetic medium (0.67% (w/v) yeast

nitrogen base, 0.079% (w/v) complete supplement mixture (CSM) amino acid mix without

uracil, containing 2% (w/v) galactose with 0.1% glucose or 2% (w/v) sucrose as carbon source.

The yeast culture was performed at 28°C to the early logarithmic growth phase and further

induced with 100 μM $CuSO_4$ for 4 h unless otherwise indicated. For the CCCP (Sigma, catalog

no. C2759) treatment experiments, yeast strains were grown in full medium that contained 2%

(w/v) glucose or 2% (w/v) sucrose at 24°C to the early logarithmic growth phase, at which time

they were treated with 0, 5, 10, or 15 μM CCCP for 15 or 30 min at the growth temperature.

The temperature-sensitive mutants were grown at a permissive temperature of 19°C and

analyzed with and without a shift to a restrictive temperature of 37°C for the indicated time.

The temperature-sensitive *pam16-3* mutants, which were transformed with p426 α-Syn WT-

GFP or A53T-GFP plasmids with galactose induction, were grown at a permissive temperature

of 19°C to the early logarithmic growth phase. Next, they were further induced with 0.5% (w/v)

galactose for another 4 h. After induction, the samples were shifted to a restrictive temperature

of 37°C for heat shock or left at 19°C as a control for 2 h. p426 GAL α-Syn WT-GFP and

A53T-GFP were provided by Dr. Tiago Outeiro (Göttingen). For the experiments that

performed proteasome inhibition, the strains were grown on minimal synthetic medium (0.67%

(w/v) yeast nitrogen base, 0.079% (w/v) CSM amino acid mix without ammonium sulphate,

supplemented with 0.1% (w/v) proline, 0.03% (w/v) SDS, and 2% (w/v) galactose as carbon

source at a permissive temperature of 19°C to the early logarithmic growth phase. Afterward,

75 μM MG132 (Enzo, catalog no. BML-PI102-0005) was added to the cell culture and

maintained at 19°C or transferred to a restrictive temperature of 37°C for the indicated time.

Proteins were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-

PAGE) using 12% or 15% gels and then transferred to polyvinyl difluoride (PVDF) membranes.

Immunodetection was performed according to standard techniques using chemiluminescence.

The following antibodies were used in the study: FLAG (Sigma, catalog no. F1804), GFP

(Sigma, catalog no. 11814460001), Hsp104 (Enzo, catalog no. ADI-SPA-1040-F), Alexa Fluor

568 (Invitrogen, catalog no. A11031), Rip1, Sod2, Mdh1, Hsp60, Pgk1, Tom70, Ssa1, Ssc1,

Rpl17, Ccp1, Qcr8, Qcr6, Cox12, Tom20, Ssb1, Edg1, Cdc48, Rpn1, Rpl5, and DNJ-21 (these

latter antibodies were non-commercial antibodies that are available in our laboratory

depository). Hsp42 antibody was provided by the Dr. Bernd Bukau laboratory (ZMBH

Heidelberg).

Aggregation assay. For the aggregation assay of metastable proteins, galactose with 0.1%

glucose was used as the carbon source. For the experiments that were followed by MG132

treatment, 2% galactose was used as the carbon source. For all other experiments, 2% (w/v)

sucrose was used as the carbon source. Ten OD_{600} units of cells were harvested by

centrifugation at $5000 \times g$ for 5 min at 4°C and re-suspended in 400 µl of Lysis buffer (30 mM

Tris-HCl pH 7.4, 20 mM KCl, 150 mM NaCl, 5 mM ethylenediaminetetraacetic acid (EDTA),

and 0.5 mM phenylmethylsulfonyl fluoride (PMSF)). Cells were homogenized by vortexing with 200 μl of glass beads (425-600 μm, Sigma-Aldrich) using a Cell Disruptor Genie (Scientific Industries) for 10 min at 4°C. The cell lysate was solubilized with 1% (v/v) Triton X-100 and mixed gently for 20 min at 4°C. The sample was centrifuged at $4000 \times g$ for 5 min at 4°C to remove unbroken cells and detergent-resistant aggregates (P_{4k}). Next, the supernatant was transferred to two tubes at equal volumes. One was saved as the total protein fraction (T). The second was centrifuged at $125,000 \times g$ for 60 min at 4°C to separate soluble proteins (S_{125k}) from protein aggregates (P_{125k}). The total (T) and soluble (S_{125k}) fractions were precipitated by 10% trichloroacetic acid. After 30 min of incubation on ice, the samples were centrifuged at $20,000 \times g$ for 15 min at 4°C, washed with ice-cold acetone, and centrifuged again. The pellets from the T and S_{125k} samples, as well as P_{4k} and P_{125k}, were resuspended in urea sample buffer (6 M urea, 6 % [w/v] SDS, 125 Tris-HCl pH 6.8, and 0.01% (w/v) bromophenol blue), incubated for 15 min at 37°C, and analyzed by SDS-PAGE followed by Western blot.

RNA-seq: sample preparation. Total RNA was isolated from the *pam16-3* mutant and corresponding WT samples, as well as BY4741 cells that were transformed with the pESC-URA plasmid that expressed pRip1-FLAG protein or an empty vector under control of the Cup1 promoter. The empty vector and pRip1 samples were grown at 28°C to the early logarithmic growth phase and further induced with 100 μM $CuSO_4$ for 1, 4, or 8 h. The *pam16WT* and *pam16-3* mutant samples after growth to the early logarithmic phase were shifted to a restrictive temperature of 37°C for 0, 1, 4, or 8 h. A total of 2.5 OD_{600} units of cells were harvested by centrifugation at $5,000 \times g$ for 5 min at 4°C, resuspended in 125 μl of RNAlater solution (Thermo Fisher Scientific, catalog no. AM7020), and incubated for 1 h at 4°C. After re-centrifugation, the cell pellet was frozen in liquid nitrogen and stored for further processing.

The *pam16-3-* and pRip1-related samples were prepared for RNA sequencing, including

four biological replicates. Each replicate was individually generated from frozen stocks. The

defrosted samples were centrifuged at $5,000 \times g$ for 5 min at 4°C to thoroughly remove the

supernatants. The loosened pellets were resuspended in 200 µl of phosphate-buffered saline

(PBS). To this sample was added 600 µl of Buffer RLT (Qiagen, catalog no. 79216) along with

acid-washed glass beads, and the cells were shaken in a cell disruptor for 10 min at maximum

speed at 4°C. Further processing was performed with agreement of the RNA extraction protocol

using the Qiagen Purification of Total RNA Kit for Yeast with Optimal On-Column DNase

Digestion (RNeasy Mini Kit (50) (Qiagen, catalog no. 74104).

RNA-seq: analysis. Sequencing was performed using an Illumina NextSeq500 instrument with

v2 chemistry, resulting in an average of 15-20 million reads per library with a 75-bp single-end

setup. The resulting raw reads were assessed for quality, adapter content, and duplication rates

with FastQC (Andrews, 2010). Reads for each sample were aligned to Ensembl *Saccharomyces*

cerevisiae transcriptome version 91 (Saccharomyces_cerevisiae.R64-1-1.91.gtf) and quantified

using Salmon (v0.11.2) (Patro et al., 2017) with default parameters. Full lists of genes for the

pam16-3 and pRip1 samples are available in **Figure2-Source Data 1** and **Figure2-Source Data**

2, respectively. Expression matrices for the *pam16-3* and pRip1 strains are available in

Figure2-Source Data 3 and **Figure2-Source Data 4**, respectively. For each time point,

differentially expressed genes were identified using DESeq2 version 1.26.0 (Love et al., 2014).

Only genes with a minimum fold change of log2 ±1, maximum Benjamini-Hochberg corrected

p value of 0.05, and minimum combined mean of 10 reads were deemed to be significantly

differentially expressed. The WT samples were used as a reference in *pam16-3*, and samples

with the empty pESC-URA plasmid were the control for the pRip1 comparisons. Raw data were

deposited in the GEO repository (accession no. GSE147284). Raw data are available at

Identification of the molecular chaperone and mitoCPR-related genes was based on an

unbiased data analysis apart from the KEGG enrichment analysis. Among all significantly

changed genes (5% FDR), we selected genes that encoded non-mitochondrial proteins. From

this group, we selected genes for which we could identify at least 10 genes that shared similar

function. This allowed for identification of group of proteins that were changed but not enriched

in KEGG enrichment analysis.

KEGG enrichment analysis. The KEGG enrichment analysis was performed for RNA-seq

using in-house developed methods based on KEGG.db (v. 3.2.3) and org.SC.sgd.db (v 3.10.0).

Pathways and genes that were selected in each developmental stage were filtered after

Benjamini-Hochberg correction for an adjusted $p < 0.05$.

Microscopy. Yeast confocal microscopy images of GAL α-Syn WT-GFP and A53T-GFP in

the *pam16-3* mutant and its corresponding WT were acquired with a Zeiss LSM700 laser-

scanning confocal microscope using a 60× oil objective (NA 1.4). Images were recorded with

pixel dimensions of 25 nm. To investigate the presence and distribution of aggregates, we used

two photomultiplier tube (PMT) detectors with 405 nm laser excitation for DAPI

(ThermoFisher Scientific, catalog no. P36962) and 488 nm for GFP fluorescence scanned

sequentially. Yeast cells were visualized using a 1AU confocal pinhole, and typically 20-25 z-

stacks were acquired, each with an optical thickness of 0.28 μm. Cytosolic aggregates were

counted manually from the z-stack Maximum Intensity Projection merge. The metastable co-

localization experiments in the *hsp42-GFP* strain were performed as above, with the additional

detection of metastable proteins by immunofluorescence. Metastable proteins were labeled

using a standard immunofluorescence protocol, with a 1:1000 dilution of the FLAG antibody

(Sigma, catalog no. F1804) and 1:500 dilution of the Alexa Fluor 568 secondary antibody

(Invitrogen, catalog no. A11031), and mounted with DAPI (ThermoFisher Scientific, catalog

no. P36962). To analyze the number of RFP and GFP aggregates in *C. elegans*, we recorded

fluorescence images of the head region of worms that expressed wrmScarlet (RFP) and GFP in

body wall muscles (ACH87 strain). Photographs were captured with a Zeiss 700 laser-scanning

confocal microscope using a 40× oil objective (NA 1.3). To investigate the presence and

distribution of aggregates, we used 2 PMT detectors with 488 nm laser excitation for GFP

fluorescence and 555 nm excitation for RFP fluorescence, scanned sequentially. Worm head

regions were visualized using a 1AU confocal pinhole, and typically 35-45 z-stacks were

acquired, each with an optical thickness of 1 μm. Eight to fifteen animals were taken per

condition per each day of worms' adulthood, starting from L4 larvae as day 0 until day 5 of

adulthood. Next, z-stacks were merged using Maximum Intensity Projection. Finally, RFP and

GFP aggregates were extracted from the photographs and counted automatically using ImageJ

software. For all of the microscopy experiments, the exposure parameters were chosen such

that no saturation effect of the fluorescent signal was present and kept at the same level during

the entire experiment.

Co-localization analysis. Fluorescence co-localization between the Alexa568 FLAG (red

channel) and Hsp42-GFP (green channel) was calculated for individual yeast cells, with three

cells per condition (marked as individual regions of interest (ROIs)). Pearson's correlation

coefficients above the Costes threshold were calculated using the Co-localization Threshold

plugin in ImageJ. This was followed by running the Co-localization Test plugin in ImageJ to

test statistical significance of the calculations. A resulting *p* value of 1.00 indicated statistical

significance.

Aggregate size analysis. Because of the small sizes of the aggregates that were at the resolution limit of the confocal microscope and the large diversity of their intensity in each cell, an automatic particle analysis based on image thresholding was not possible. Therefore, we analyzed the average size of the aggregates using a manual approach of defining aggregate boundaries and measuring the aggregate size with ImageJ software, with $n = 10$ analyzed for each condition.

Worm maintenance and strains. Standard conditions were used for the propagation of *C. elegans* (Brenner, 1974). Briefly, the animals were synchronized by hypochlorite bleaching, hatched overnight in M9 buffer (3 g/L KH_2PO_4, 6 g/L Na_2HPO_4, 5 g/L NaCl, and 1 M $MgSO_4$), and subsequently cultured at 20°C on nematode growth medium (NGM) plates (1 mM $CaCl_2$, 1 mM $MgSO_4$, 5 μg/mL cholesterol, 25 mM KPO_4 buffer pH 6.0, 17 g/L agar, 3 g/L NaCl, and 2.5 g/L peptone) seeded with the *E. coli* HB101 or OP50 strain as a food source. The following *C. elegans* strains were used:

CL2122: dvIs15 [(pPD30.38) unc-54(vector) + (pCL26) mtl-2::GFP]. Control strain for GMC101.

GMC101: dvIs100 [punc-54::A-beta-1-42::unc-54-3'-UTR + mtl-2p::GFP]

ACH87: wacIs11 [pmyo-3::mGFP::SL2 gdp-2-wrmScarlet::unc-54-3'UTR, unc-119(+)]

OW40: zgIs15 [punc-54::αsyn::YFP]IV (van Ham et al., 2008)

OW450: rmIs126 [punc-54::YFP]V. Control strain for OW40 (van Ham et al., 2008)

Molecular biology for worm studies. An RNAi construct that targeted the *dnj-21* gene was created by PCR amplification of the gene from cDNA pools that were generated from RNA. Primers that were used to generate PCR products were designed to amplify the full cDNA

sequence of *dnj-21*. The PCR product was digested with XhoI and KpnI restriction enzymes

and cloned into the XhoI and KpnI-digested L4440 vector. Constructs for the expression of

wrmScarlet (RFP; mCherry derivative) (El Mouridi et al., 2017) and mGFP in body wall

muscles of worms were created using the SLiCE method (Zhang et al., 2012) and the Three

Fragment Gateway System (Invitrogen). Briefly, wrmScarlet from the pSEM87 plasmid was

PCR amplified using the following primers: ggaaactgcttcaacgcatcatggtcagcaagggagag and

gaagagtaattggacttacttgtagagctcgtccatt. The PCR product was used to replace the mKate2

sequence in the pCG150 vector with the pmyo-3::mGFP::SL2 gdp-2-mKate2::unc-54-3'UTR

insert. Cloned constructs were sequenced for insert verification.

Worm transformation. The *C. elegans* ACH87 transgenic strain was created using biolistic

bombardment as described previously (Praitis et al., 2001). *unc-119* rescue was used as a

selection marker.

Worm RNAi. RNAi was achieved by feeding worms *E. coli* HT115(DE3) bacteria that was

transformed with a construct that targeted the *dnj-21* gene. *E. coli* HT115(DE3) that was

transformed with the L4440 empty vector was used as a control. LB medium (10 g/L tryptone,

10 g/L NaCl, and 5 g/L yeast extract) was inoculated with transformed bacteria and cultured at

37°C at 180 rotations per minute. Bacterial culture was induced with 1 mM isopropyl β-D-1-

thiogalactopyranoside (IPTG) once the culture OD_{600} reached 0.4-0.6. After 2 h, the bacteria

were pelleted. For liquid culture, the bacterial pellet was added to S-medium and used for

further worm culture. For cultures on solid medium, the plates were seeded with bacteria and

used after the bacteria dried.

Worm total protein isolation and Western blot. GMC101 L1 larvae were cultured in liquid S-medium for 3 days, first at 20°C for 24 h and then at 20°C for 48 h or at 22°C or 25°C for 48 h to induce Aβ aggregation. Total proteins were isolated from frozen worm pellets. Samples were thawed on ice in lysis buffer (20 mM Tris pH 7.4, 200 mM NaCl, and 2 mM PMSF) and sonicated three times for 10 s. The lysate was centrifuged at $2,800 \times g$ for 5 min at 4°C. The pellet that contained debris was discarded. Protein concentrations were measured by DirectDetect. Proteins were separated by SDS-PAGE (using 15% gels) or native PAGE (12% gels) and then transferred to PVDF membranes. Immunodetection was performed according to standard techniques using chemiluminescence. The following antibodies were used: custom-made rabbit antibody against DNJ-21, purified anti-beta amyloid 1-16 antibody (Biolegend, catalog no. 803001), tubulin (Sigma, catalog no. T9026), mCherry (Abcam, catalog no. ab167453), and GFP (Sigma, catalog no. 11814460001).

Motility assay. The worms were placed in a drop of M9 buffer and allowed to recover for 30 s to avoid the observation of behavior that is associated with stress, after which the number of body bends was counted for 30 s. The experiments were conducted in triplicate. For each experiment, at least 35 worms were analyzed.

Automated body bend assay. Automated motility assessment was performed with a tracking device that was developed and described previously (Perni et al., 2018). Briefly, worms were washed off the plates with M9 buffer and spread over a 9 cm NGM plate in a final volume of 5 mL, after which their movements were recorded for 120 s. Videos were analyzed in a consistent manner to track worm motility (bends per minute) and swimming speed. The results show the mean ± SEM from two independent experiments.

Amyloid β (Aβ) quantification. Aβ aggregates were calculated by dividing the signal that was detected with anti-Aβ antibody by the protein signal that was detected with Coomassie staining. For each temperature condition, aggregate levels were normalized to the control. The data are expressed as the mean ± SD ($n = 3$). Overall differences between conditions were assessed using unpaired t-tests by assuming unequal variance.

Statistical analysis. For the statistical analysis, two-tailed, unpaired t-tests were used by assuming equal variance unless otherwise stated. Values of $p \leq 0.05$ were considered statistically significant. For the statistical analysis of the confocal microscopy co-localization data, see the co-localization analysis section in the Methods.

Data and material availability. The authors declare that data that support the findings of this study are available within the paper, supporting information, and source data.

Acknowledgements: We thank Tiago Outeiro, Bernd Bukau, Axel Mogk, Magdalena Boguta, Henrik Bringmann, Thomas Boulin, Ellen Nollen, and Agnieszka Sztyler for materials and experimental assistance and CGC (funded by the National Institutes of Health Office of Research Infrastructure Programs, P40 OD010440) for providing *C. elegans* strains.

Author contributions: U.N., P.C., K.S., M.S., M.T., T.G., and M.P. performed and analyzed the biochemical experiments. U.N., P.C., K.S., B.U.R, and R.K. performed RNA-seq and the analyses. C.M.D., M.V., and A.C. conceived the project. All authors interpreted the experiments. M.V. and A.C. analyzed and supervised the study. U.N, M.V., and A.C. wrote the manuscript with input from all authors.

Funding: Research in the M.V. laboratory was supported by the Centre for Misfolding Diseases

and a donation by the William B. Harrison Foundation.

Research in the A.C. laboratory was supported by the Ministerial funds for science within Ideas

Plus programme 000263 in 2014-2017 (P.Ch, U.N), HOMING project POIR.04.04.00-00-

3FE4/17 (UN) of the Foundation for Polish Science co-financed by the European Union under

the European Regional Development Fund, "Regenerative Mechanisms for Health" project

MAB/2017/2 (M.T, A.C) carried out within the International Research Agendas programme of

the Foundation for Polish Science co-financed by the European Union under the European

Regional Development Fund, Copernicus Award (A.C, P.Ch.) of the Foundation of Polish

Science and German Research Foundation (DFG), National Science Centre grants:

2015/18/A/NZ1/00025 (M.S, A.C), and POLONEZ Fellowships 2016/23/P/NZ3/03730

(B.U.R) and UMO-2016/21JPJNZ3/03891 (M.T). The POLONEZ programme has received

funding from the European Union's Horizon 2020 research and innovation programme under

the Marie Sklodowska-Curie grant agreement No 665778. MS was also supported by the

EMBO short-term fellowship 7124.

Competing interests: The authors declare no competing financial interests.

References

727   Andreasson, C., Ott, M., & Buttner, S. (2019). Mitochondria orchestrate proteostatic and

728   metabolic stress responses. *EMBO Rep*, *20*(10), e47865.

729   https://doi.org/10.15252/embr.201947865

730   Andrews, S. (2010). *FastQC: a quality control tool for high throughput sequence data.*

731   http://www.bioinformatics.babraham.ac.uk/projects/fastqc

Balchin, D., Hayer-Hartl, M., & Hartl, F. U. (2016). In vivo aspects of protein folding and

quality control. *Science*, *353*(6294), aac4354. https://doi.org/10.1126/science.aac4354

Boos, F., Kramer, L., Groh, C., Jung, F., Haberkant, P., Stein, F., Wollweber, F., Gackstatter,

A., Zoller, E., van der Laan, M., Savitski, M. M., Benes, V., & Herrmann, J. M. (2019).

Mitochondrial protein-induced stress triggers a global adaptive transcriptional

programme. *Nat Cell Biol*, *21*(4), 442-451. https://doi.org/10.1038/s41556-019-0294-5

Boos, F., Labbadia, J., & Herrmann, J. M. (2020). How the mitoprotein-induced stress response

safeguards the cytosol: a unified view. *Trends Cell Biol*, *30*(3), 241-254.

https://doi.org/10.1016/j.tcb.2019.12.003

Braun, R. J. & Westermann, B. (2017). With the help of MOM: mitochondrial contributions to

cellular quality control. *Trends Cell Biol*, *27*(6), 441-452.

https://doi.org/10.1016/j.tcb.2017.02.007

Brenner, S. (1974). The genetics of *Caenorhabditis elegans*. *Genetics*, *77*(1), 71-94.

https://www.ncbi.nlm.nih.gov/pubmed/4366476

Bruderek, M., Jaworek, W., Wilkening, A., Rub, C., Cenini, G., Fortsch, A., Sylvester, M., &

Voos, W. (2018). IMiQ: a novel protein quality control compartment protecting

mitochondrial functional integrity. *Mol Biol Cell*, *29*(3), 256-269.

https://doi.org/10.1091/mbc.E17-01-0027

Cabral-Costa, J. V. & Kowaltowski, A. J. (2020). Neurological disorders and mitochondria.

Mol Aspects Med, *71*, 100826. https://doi.org/10.1016/j.mam.2019.10.003

Calvo, S. E., Clauser, K. R., & Mootha, V. K. (2016). MitoCarta2.0: an updated inventory of

mammalian mitochondrial proteins. *Nucleic Acids Res*, *44*(D1), D1251-1257.

https://doi.org/10.1093/nar/gkv1003

Cenini, G., Rub, C., Bruderek, M., & Voos, W. (2016). Amyloid β-peptides interfere with

mitochondrial preprotein import competence by a coaggregation process. *Mol Biol Cell*,

27(21), 3257-3272. https://doi.org/10.1091/mbc.E16-05-0313

Chacinska, A., Boguta, M., Krzewska, J., & Rospert, S. (2000). Prion-dependent switching

between respiratory competence and deficiency in the yeast nam9-1 mutant. *Mol Cell*

Biol, *20*(19), 7220-7229. https://doi.org/10.1128/MCB.20.19.7220-7229.2000

Chacinska, A., Koehler, C. M., Milenkovic, D., Lithgow, T., & Pfanner, N. (2009). Importing

mitochondrial proteins: machineries and mechanisms. *Cell*, *138*(4), 628-644.

https://doi.org/10.1016/j.cell.2009.08.005

Chacinska, A., Pfannschmidt, S., Wiedemann, N., Kozjak, V., Sanjuan Szklarz, L. K., Schulze-

Specking, A., Truscott, K. N., Guiard, B., Meisinger, C., & Pfanner, N. (2004). Essential

role of Mia40 in import and assembly of mitochondrial intermembrane space proteins.

EMBO J, *23*(19), 3735-3746. https://doi.org/10.1038/sj.emboj.7600389

Ciryam, P., Kundra, R., Freer, R., Morimoto, R. I., Dobson, C. M., & Vendruscolo, M. (2016).

A transcriptional signature of Alzheimer's disease is associated with a metastable

subproteome at risk for aggregation. *Proc Natl Acad Sci U S A*, *113*(17), 4753-4758.

https://doi.org/10.1073/pnas.1516604113

Claros, M. G. & Vincens, P. (1996). Computational method to predict mitochondrially imported

proteins and their targeting sequences. *Eur J Biochem*, *241*(3), 779-786.

https://www.ncbi.nlm.nih.gov/pubmed/8944766

Di Maio, R., Barrett, P. J., Hoffman, E. K., Barrett, C. W., Zharikov, A., Borah, A., Hu, X.,

McCoy, J., Chu, C. T., Burton, E. A., Hastings, T. G., & Greenamyre, J. T. (2016). α-

Synuclein binds to TOM20 and inhibits mitochondrial protein import in Parkinson's

disease. *Sci Transl Med*, *8*(342), 342ra78. https://doi.org/10.1126/scitranslmed.aaf3634

El Mouridi, S., Lecroisey, C., Tardy, P., Mercier, M., Leclercq-Blondel, A., Zariohi, N., &

Boulin, T. (2017). Reliable CRISPR/Cas9 genome engineering in *Caenorhabditis*

elegans using a single efficient sgRNA and an easily recognizable phenotype. *G3*, *7*(5),

1429-1437. https://doi.org/10.1534/g3.117.040824

Escobar-Henriques, M., Altin, S., & Brave, F. D. (2020). Interplay between the ubiquitin

proteasome system and mitochondria for protein homeostasis. *Curr Issues Mol Biol*, *35*,

35-58. https://doi.org/10.21775/cimb.035.035

Frazier, A. E., Dudek, J., Guiard, B., Voos, W., Li, Y., Lind, M., Meisinger, C., Geissler, A.,

Sickmann, A., Meyer, H. E., Bilanchone, V., Cumsky, M. G., Truscott, K. N., Pfanner,

N., & Rehling, P. (2004). Pam16 has an essential role in the mitochondrial protein

import motor. *Nat Struct Mol Biol*, *11*(3), 226-233. https://doi.org/10.1038/nsmb735

Fukasawa, Y., Tsuji, J., Fu, S. C., Tomii, K., Horton, P., & Imai, K. (2015). MitoFates:

improved prediction of mitochondrial targeting sequences and their cleavage sites. *Mol*

Cell Proteomics, *14*(4), 1113-1126. https://doi.org/10.1074/mcp.M114.043083

Gulshan, K., Schmidt, J. A., Shahi, P., & Moye-Rowley, W. S. (2008). Evidence for the

bifunctional nature of mitochondrial phosphatidylserine decarboxylase: role in Pdr3-

dependent retrograde regulation of *PDR5* expression. *Mol Cell Biol*, *28*(19), 5851-5864.

https://doi.org/10.1128/MCB.00405-08

Izawa, T., Park, S. H., Zhao, L., Hartl, F. U., & Neupert, W. (2017). Cytosolic protein Vms1

links ribosome quality control to mitochondrial and cellular homeostasis. *Cell*, *171*(4),

890-903.e18. https://doi.org/10.1016/j.cell.2017.10.002

Kim, H. E., Grant, A. R., Simic, M. S., Kohnz, R. A., Nomura, D. K., Durieux, J., Riera, C. E.,

Sanchez, M., Kapernick, E., Wolff, S., & Dillin, A. (2016). Lipid biosynthesis

coordinates a mitochondrial-to-cytosolic stress response. *Cell*, *166*(6), 1539-1552.e16.

https://doi.org/10.1016/j.cell.2016.08.027

Kowalski, L., Bragoszewski, P., Khmelinskii, A., Glow, E., Knop, M., & Chacinska, A. (2018).

Determinants of the cytosolic turnover of mitochondrial intermembrane space proteins.

BMC Biol, *16*(1), 66. https://doi.org/10.1186/s12915-018-0536-1

Kundra, R., Ciryam, P., Morimoto, R. I., Dobson, C. M., & Vendruscolo, M. (2017). Protein

homeostasis of a metastable subproteome associated with Alzheimer's disease. *Proc*

Natl Acad Sci U S A, *114*(28), E5703-E5711. https://doi.org/10.1073/pnas.1618417114

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol*, *15*(12), 550.

https://doi.org/10.1186/s13059-014-0550-8

Martensson, C. U., Priesnitz, C., Song, J., Ellenrieder, L., Doan, K. N., Boos, F., Floerchinger,

A., Zufall, N., Oeljeklaus, S., Warscheid, B., & Becker, T. (2019). Mitochondrial

protein translocation-associated degradation. *Nature*, *569*(7758), 679-683.

https://doi.org/10.1038/s41586-019-1227-y

Mogk, A., Kummer, E., & Bukau, B. (2015). Cooperation of Hsp70 and Hsp100 chaperone

machines in protein disaggregation. *Front Mol Biosci*, *2*, 22.

https://doi.org/10.3389/fmolb.2015.00022

Mogk, A., Ruger-Herreros, C., & Bukau, B. (2019). Cellular functions and mechanisms of

action of small heat shock proteins. *Annu Rev Microbiol*, *73*, 89-110.

https://doi.org/10.1146/annurev-micro-020518-115515

Mohanraj, K., Nowicka, U., & Chacinska, A. (2020). Mitochondrial control of cellular protein

homeostasis. *Biochem J*, *477*(16), 3033-3054. https://doi.org/10.1042/BCJ20190654

Morgenstern, M., Stiller, S. B., Lubbert, P., Peikert, C. D., Dannenmaier, S., Drepper, F., Weill,

U., Hoss, P., Feuerstein, R., Gebert, M., Bohnert, M., van der Laan, M., Schuldiner, M.,

Schutze, C., Oeljeklaus, S., Pfanner, N., Wiedemann, N., & Warscheid, B. (2017).

Definition of a high-confidence mitochondrial proteome at quantitative scale. *Cell Rep*,

19(13), 2836-2852. https://doi.org/10.1016/j.celrep.2017.06.014

Mossmann, D., Vogtle, F. N., Taskin, A. A., Teixeira, P. F., Ring, J., Burkhart, J. M., Burger,

N., Pinho, C. M., Tadic, J., Loreth, D., Graff, C., Metzger, F., Sickmann, A., Kretz, O.,

Wiedemann, N., Zahedi, R. P., Madeo, F., Glaser, E., & Meisinger, C. (2014). Amyloid-

β peptide induces mitochondrial dysfunction by inhibition of preprotein maturation.

Cell Metab, *20*(4), 662-669. https://doi.org/10.1016/j.cmet.2014.07.024

Neupert, W. & Herrmann, J. M. (2007). Translocation of proteins into mitochondria. *Annu Rev*

Biochem, *76*, 723-749. https://doi.org/10.1146/annurev.biochem.76.052705.163409

Outeiro, T. F. & Lindquist, S. (2003). Yeast cells provide insight into alpha-synuclein biology

and pathobiology. *Science*, *302*(5651), 1772-1775.

https://doi.org/10.1126/science.1090439

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., & Kingsford, C. (2017). Salmon provides

fast and bias-aware quantification of transcript expression. *Nat Methods*, *14*(4), 417-

419. https://doi.org/10.1038/nmeth.4197

Perni, M., Challa, P. K., Kirkegaard, J. B., Limbocker, R., Koopman, M., Hardenberg, M. C.,

Sormanni, P., Muller, T., Saar, K. L., Roode, L. W. Y., Habchi, J., Vecchi, G., Fernando,

N., Casford, S., Nollen, E. A. A., Vendruscolo, M., Dobson, C. M., & Knowles, T. P. J.

(2018). Massively parallel *C. elegans* tracking provides multi-dimensional fingerprints

for phenotypic discovery. *J Neurosci Methods*, *306*, 57-67.

https://doi.org/10.1016/j.jneumeth.2018.02.005

Pfanner, N., Warscheid, B., & Wiedemann, N. (2019). Mitochondrial proteins: from biogenesis

to functional networks. *Nat Rev Mol Cell Biol*, *20*(5), 267-284.

https://doi.org/10.1038/s41580-018-0092-0

Poveda-Huertes, D., Matic, S., Marada, A., Habernig, L., Licheva, M., Myketin, L., Gilsbach,

R., Tosal-Castano, S., Papinski, D., Mulica, P., Kretz, O., Kucukkose, C., Taskin, A.

A., Hein, L., Kraft, C., Buttner, S., Meisinger, C., & Vogtle, F. N. (2020). An early

mtUPR: redistribution of the nuclear transcription factor Rox1 to mitochondria protects

against intramitochondrial proteotoxic aggregates. *Mol Cell*, *77*(1), 180-188.e9.

https://doi.org/10.1016/j.molcel.2019.09.026

Praitis, V., Casey, E., Collar, D., & Austin, J. (2001). Creation of low-copy integrated

transgenic lines in *Caenorhabditis elegans*. *Genetics*, *157*(3), 1217-1226.

https://www.ncbi.nlm.nih.gov/pubmed/11238406

Priesnitz, C. & Becker, T. (2018). Pathways to balance mitochondrial translation and protein

import. *Genes Dev*, *32*(19-20), 1285-1296. https://doi.org/10.1101/gad.316547.118

Ruan, L., Zhou, C., Jin, E., Kucharavy, A., Zhang, Y., Wen, Z., Florens, L., & Li, R. (2017).

Cytosolic proteostasis through importing of misfolded proteins into mitochondria.

Nature, *543*(7645), 443-446. https://doi.org/10.1038/nature21695

Sontag, E. M., Samant, R. S., & Frydman, J. (2017). Mechanisms and functions of spatial

protein quality control. *Annu Rev Biochem*, *86*, 97-122.

https://doi.org/10.1146/annurev-biochem-060815-014616

Sormanni, P., Aprile, F. A., & Vendruscolo, M. (2015). The CamSol method of rational design

of protein mutants with enhanced solubility. *J Mol Biol*, *427*(2), 478-490.

https://doi.org/10.1016/j.jmb.2014.09.026

Sorrentino, V., Romani, M., Mouchiroud, L., Beck, J. S., Zhang, H., D'Amico, D., Moullan,

N., Potenza, F., Schmid, A. W., Rietsch, S., Counts, S. E., & Auwerx, J. (2017).

Enhancing mitochondrial proteostasis reduces amyloid-β proteotoxicity. *Nature*,

552(7684), 187-193. https://doi.org/10.1038/nature25143

Stojanovski, D., Milenkovic, D., Muller, J. M., Gabriel, K., Schulze-Specking, A., Baker, M.

J., Ryan, M. T., Guiard, B., Pfanner, N., & Chacinska, A. (2008). Mitochondrial protein

import: precursor oxidation in a ternary complex with disulfide carrier and sulfhydryl

oxidase. *J Cell Biol*, *183*(2), 195-202. https://doi.org/10.1083/jcb.200804095

Truscott, K. N., Voos, W., Frazier, A. E., Lind, M., Li, Y., Geissler, A., Dudek, J., Muller, H.,

Sickmann, A., Meyer, H. E., Meisinger, C., Guiard, B., Rehling, P., & Pfanner, N.

(2003). A J-protein is an essential subunit of the presequence translocase-associated

protein import motor of mitochondria. *J Cell Biol*, *163*(4), 707-713.

https://doi.org/10.1083/jcb.200308004

Tyedmers, J., Mogk, A., & Bukau, B. (2010). Cellular strategies for controlling protein

aggregation. *Nat Rev Mol Cell Biol*, *11*(11), 777-788. https://doi.org/10.1038/nrm2993

van der Laan, M., Hutu, D. P., & Rehling, P. (2010). On the mechanism of preprotein import

by the mitochondrial presequence translocase. *Biochim Biophys Acta*, *1803*(6), 732-739.

https://doi.org/10.1016/j.bbamcr.2010.01.013

van Ham, T. J., Thijssen, K. L., Breitling, R., Hofstra, R. M., Plasterk, R. H., & Nollen, E. A.

(2008). *C. elegans* model identifies genetic modifiers of α-synuclein inclusion

formation during aging. *PLoS Genet*, *4*(3), e1000027.

https://doi.org/10.1371/journal.pgen.1000027

Vecchi, G., Sormanni, P., Mannini, B., Vandelli, A., Tartaglia, G. G., Dobson, C. M., Hartl, F.

U., & Vendruscolo, M. (2020). Proteome-wide observation of the phenomenon of life

on the edge of solubility. *Proc Natl Acad Sci U S A*, *117*(2), 1015-1020.

https://doi.org/10.1073/pnas.1910444117

Vicario, M., Cieri, D., Brini, M., & Cali, T. (2018). The close encounter between alpha-

synuclein and mitochondria. *Front Neurosci*, *12*, 388.

https://doi.org/10.3389/fnins.2018.00388

Wang, X. & Chen, X. J. (2015). A cytosolic network suppressing mitochondria-mediated

proteostatic stress and cell death. *Nature*, *524*(7566), 481-484.

https://doi.org/10.1038/nature14859

Weidberg, H. & Amon, A. (2018). MitoCPR-A surveillance pathway that protects mitochondria

in response to protein import stress. *Science*, *360*(6385), eaan4146.

https://doi.org/10.1126/science.aan4146

Wrobel, L., Topf, U., Bragoszewski, P., Wiese, S., Sztolsztener, M. E., Oeljeklaus, S.,

Varabyova, A., Lirski, M., Chroscicki, P., Mroczek, S., Januszewicz, E., Dziembowski,

A., Koblowska, M., Warscheid, B., & Chacinska, A. (2015). Mistargeted mitochondrial

proteins activate a proteostatic response in the cytosol. *Nature*, *524*(7566), 485-488.

https://doi.org/10.1038/nature14951

Wu, Z., Tantray, I., Lim, J., Chen, S., Li, Y., Davis, Z., Sitron, C., Dong, J., Gispert, S.,

Auburger, G., Brandman, O., Bi, X., Snyder, M., & Lu, B. (2019). MISTERMINATE

mechanistically links mitochondrial dysfunction with proteostasis failure. *Mol Cell*,

75(4), 835-848.e8. https://doi.org/10.1016/j.molcel.2019.06.031

Zhang, Y., Werling, U., & Edelmann, W. (2012). SLiCE: a novel bacterial cell extract-based

DNA cloning method. *Nucleic Acids Res*, *40*(8), e55.

https://doi.org/10.1093/nar/gkr1288

Figure 1. Supersaturated nuclear-encoded mitochondrial proteins aggregate in the cytosol. (**A**) SDS-PAGE analysis of aggregation assay fractions of WT yeast cells that overexpressed Atp2$_{FLAG}$, Atp20$_{FLAG}$, Cox8$_{FLAG}$, pRip1$_{FLAG}$, Qcr8$_{FLAG}$, Cor1$_{FLAG}$, Mas1$_{FLAG}$, Cox12$_{FLAG}$, and Qcr6$_{FLAG}$ for 3 h when 2% galactose with 0.1% glucose was used as the carbon source. Pellet fractions after 4,000 and 125,000 $\times$ g centrifugation are indicated as P$_{4k}$ and P$_{125k}$, respectively. The soluble fraction at 125,000 $\times$ g is indicated as S$_{125k}$. $n = 3$. (**B**) Ten-fold dilutions of WT cells that expressed metastable proteins or controls that were spotted on selective medium agar plates with galactose as the main carbon source at 28°C and 37°C. (**C**) Total protein cell extract from WT yeast grown at 24°C and treated with 0, 5, 10, or 15 μM CCCP for 30 min. (**D**) Quantification of pRip1, pSod2, and pMdh1 from **C**. Quantified data are shown as mean $\pm$ SEM. $n = 3$. (**E**) SDS-PAGE analysis of aggregation assay fractions of yeast cells that were treated with 15 μM CCCP for 30 min, with 2% sucrose as the carbon source. (**F**) SDS-PAGE analysis of aggregation assay fractions of WT (*pam16WT*) and *pam16-3* mutant yeast strains grown at 19°C and shifted to 37°C for 0, 3, or 5 h, with 2% sucrose as the carbon source. In **A**, **C**, **E**, and **F**, the samples were separated by SDS-PAGE and identified by Western blot with specific antisera. $n = 3$ for each experiment. *, nonspecific; p, presequence protein; i, intermediate protein; m, mature protein. *$p < 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$, ****$p \leq 0.0001$.

Figure supplement 1. Aggregation propensity characterization of mitochondrial proteins.

Figure supplement 2. Supersaturated nuclear-encoded mitochondrial protein aggregate in the cytosol and stimulate growth defects.

Figure 2. Mitochondrial protein import defects stimulate specific transcriptomic responses. (**A, B**) Volcano plot comparison of changes in expression (in terms of the logarithm of the fold change) assessed by the RNA-seq analysis of (**A**) WT yeast cells that overexpressed

pRip1 compared with the empty vector (ev) control (induction (I) was performed under control

of the CUP1promoter for 1, 4, or 8 h for both ev and pRip1) and (**B**) the *pam16-3* strain

compared with WT (*pam16WT*) (both cell types were grown at a permissive temperature of

19°C and shifted to a restrictive temperature of 37°C for 1, 4, or 8 h). HS, heat shock time (in

hours). Differentially expressed genes that encode molecular chaperones are indicated in blue.

The gene name is displayed for each molecular chaperone if it was detected as differentially

expressed (i.e., 5% FDR, log2 fold change [log2FC] ± 1). (**C**) All differentially expressed genes

in pRip1 samples after 1, 4, and 8 h of overexpression. (**D**) Analysis of changes in the expression

of genes that encode molecular chaperones using *pam16-3* samples that were treated as in **B**.

Up- and downregulated genes (5% FDR) are shown in green and pink, respectively. The

intensity of the color shades reflects the level of expression change (log2FC). Genes that were

not detected or those without statistically significant expression changes are depicted in gray.

Figure supplement 1. Characterization of gene change that was attributable to mitochondrial

protein import defects.

Figure supplement 2. Characterization of gene change that was attributable to heat shock in

WT and the *pam16-3* mutant.

Supplementary file 1. Side-by-side comparison of gene expression levels that were attributable

to heat shock and the *pam16-3* mutant.

Source Data 1. Full list of gene changes in response to pRip1 overexpression.

Source Data 2. Full list of gene changes in response to *pam16-3* overexpression.

Source Data 3. Gene expression matrices in response to pRip1 overexpression.

Source Data 4. Gene expression matrices in response to *pam16-3* mutation.

Figure 3. Cytosolic mitochondrial protein aggregation elicits a molecular chaperone response to restore cellular homeostasis. (**A, D**) Hsp104 and Hsp42 are upregulated in response to impairment in mitochondrial protein import. (**A**) Total protein cell extracts from WT yeast cells were grown at 24°C and treated with 0, 5, 10, or 15 μM CCCP for 30 min. (**B**) Quantified changes in Hsp104 protein expression from **A**. Quantified data are shown as mean ± SEM. $n = 3$. (**C**) Total protein cell extracts from WT (YPH499), *pam16-WT*, *pam16-1*, *pam16-3*, *pam18-WT*, *pam18-1*, *mia40-4WT*, *mia40-4int*, and *mia40-3* grown at a permissive temperature of 19°C and shifted to a restrictive temperature of 37°C for 3 h. (**D**) Total protein cell extracts from WT yeast cells and *hsp42-GFP* yeast grown at 24°C and treated with 0 or 15 μM CCCP for 30 min. (**E-H**) Metastable protein overexpression increases the expression levels of molecular chaperones. Total protein cell extracts that expressed the indicated metastable proteins or an empty vector control for 4 h showed higher levels of Hsp104 (**E**) and Hsp42 (**G**). (**F, H**) Quantitative analyses of Hsp104 (**F**) and Hsp42 (**H**) levels from **E** and **G**, respectively, are shown. Quantified data are shown as the mean ± SEM. $n = 3$. (**I**) Aggregated metastable proteins co-localize with Hsp42-GFP. Representative confocal microscope images show metastable proteins that were tagged with Alexa568 fluorophore in the *hsp42-GFP* yeast strain. Scale bar = 2 μm. See Methods for further details. Pearson's correlation coefficients were calculated for the indicated cells for each condition. (**J**) Total protein cell extracts from WT yeast cells and *Δhsp104* yeast grown at 24°C and treated with 0 or 15 μM CCCP for 30 min. (**K**) SDS-PAGE analysis of aggregation assay fractions of samples of WT yeast cells and *Δhsp104* yeast grown at 24°C and treated with 0 or 15 μM CCCP for 30 min, with 2% sucrose as the carbon source. (**L**) Total protein cell extracts from WT yeast cells and *Δhsp42* yeast grown at 24°C and treated with 0 or 15 μM CCCP for 30 min. (**M**) Quantified changes in Hsp104 expression from **L**. Quantified data are shown as the mean ± SEM. $n = 3$. (**N**) Total protein cell extracts from WT yeast cells and yeast cells that expressed Hsp104 grown overnight

at 24°C and treated with 0 or 15 μM CCCP for 30 min. In **A**, **C-E**, **G**, **J**, **K-L**, and **N**, the

samples were separated by SDS-PAGE and identified by Western blot with specific antisera.

Each experiment was repeated three times. p, presequence protein; m, mature protein; *,

nonspecific. Significance in **B**, **F**, **H**, and **M**: $*p < 0.05$, $**p \leq 0.01$, $***p \leq 0.001$, $****p \leq$

0.0001; ns, nonsignificant.

Figure supplement 1. Total protein cell extract changes that were attributable to mitochondrial

import defects.

Figure supplement 2. Cellular effects of Hsp42 and Hsp104 overexpression during

mitochondrial import defects.

1005

Figure 4. Mitochondrial protein import dysfunction enhances impairment in cellular

homeostasis. (**A-D**) Metastable proteins cause the accumulation and aggregation of other

mitochondrial precursor proteins. (**A**) Total protein cell extracts from *hsp42-GFP* cells that

expressed selected metastable proteins or an empty vector control overnight. Changes for pRip1

(**B**) and pSod2 (**C**) were quantified. Quantified data are shown as the mean ± SEM. $n = 3$. (**D**)

SDS-PAGE analysis of aggregation assay fractions of *hsp42-GFP* yeast cells that

overexpressed Atp2$_{FLAG}$ or an empty vector control for 3 h, with 2% sucrose as the carbon

source. insoluble, S$_{4k}$ aggregation assay fraction. (**E**) Total protein cell extract from *pam16-3*

mutant yeast strains treated with 75 μM MG132 for 1 h under permissive growth conditions

and subsequently heat shocked at 37°C for 0, 1, 2, or 4 h. (**F, G**) Representative confocal images

of α-Syn WT-GFP (**F**) and A53T-GFP (**G**) aggregates in WT (*pam16-WT*) and *pam16-3* yeast

strains. α-Syn WT-GFP and A53T-GFP were induced for 4 h at 19°C and for an additional 2 h

at 19°C for control or at 37°C for heat shock. Scale bar = 2 μm. See Methods for further details.

The bar plot shows the average number of aggregates per cell. The data are shown as the

mean ± SEM. $n = 57-83$ for α-Syn WT-GFP. $n = 154-175$ for α-Syn A53T-GFP. (**H**) Total cell

extracts of WT (*pam16-WT*) and *pam16-3* yeast strains that expressed α-Syn WT-GFP induced

for 4 h at 19°C and for an additional 2 h at 19°C for control or 37°C for heat shock. (**I**)

Quantitative analysis of pRip1 from **H**. Quantified data are shown as the mean ± SEM. $n = 3$.

In **A**, **D**, **E**, and **H**, protein samples were separated by SDS-PAGE and identified by Western

blot with specific antisera. Each experiment was repeated three times. For Western blot: p,

presequence protein; i, intermediate protein; m, mature protein; *, nonspecific. $*p < 0.05$, $**p$

≤ 0.01, $***p \leq 0.001$, $****p \leq 0.0001$; ns, nonsignificant.

Figure supplement 1. Effects of mitochondrial protein import dysfunction on cellular

homeostasis.

1030

Figure 5. Mitochondrial dysfunction results in the accumulation of Aβ aggregates in *C.*

***elegans*.** (**A-C**) Mitochondrial dysfunction stimulates the aggregation of model proteins in *C.*

elegans. (**A**) Confocal images of worms that expressed wrmScarlet and GFP in body wall

muscle [pmyo-3::wrmScarlet+pmyo::GFP]. The zoomed image is presented in the white box.

Scale bar = 20 μm. (**B**) Number of RFP aggregates at different days of adulthood of strain

[pmyo-3::wrmScarlet + pmyo-3::GFP] strain upon *dnj-21* RNAi. $n = 14$-16 worms for empty

vector. $n = 8$-16 worms for *dnj-21* RNAi. (**C**) Number of GFP aggregates present at different

days of adulthood of [pmyo-3::wrmScarlet + pmyo-3::GFP] strain upon *dnj-21* RNAi. $n = 14$-

16 worms for empty vector. $n = 8$-16 worms for *dnj-21* RNAi. (**D**) Motility of Parkinson's

disease model strain that expressed α-Syn::YFP in the body wall muscle or control strain that

expressed YFP in the body wall muscle upon the silencing of *dnj-21*. An empty vector was used

as the control. Data were obtained using an automated body bend assay. The data are shown as

the mean ± SEM, with at least $n = 700$ for each condition. (**E**) Protein aggregation under native

conditions in worms that expressed Aβ upon *dnj-21* RNAi. Worms were cultured at 20°C or

shifted to 22°C or 25°C. $n = 3$. (**F**) Aβ levels were calculated from the native aggregation data

in **E**. The data are shown as the mean ± SD. $n = 3$. (**G**) Motility in worms that expressed Aβ

and GFP upon *dnj-21* RNAi. The data are shown as the mean ± SEM. $n = 50$ worms per

condition. Overall differences between conditions were assessed by unpaired *t*-test by assuming

equal variance. *$p < 0.05$, **$p ≤ 0.01$, ***$p ≤ 0.001$, ****$p ≤ 0.0001$; ns, nonsignificant.

Figure supplement 1. Scheme and controls for experiments on mitochondrial dysfunction in

C. elegans that resulted in Aβ accumulation.

1052

Figure 1–figure supplement 1. Aggregation propensity characterization of mitochondrial

proteins. Protein sequences and information about mitochondrial targeting presequences were

acquired from the Saccharomyces Genome Database and verified using Mitofates (Fukasawa

et al., 2015) and MitoProt (Claros & Vincens, 1996) software. Protein solubility was analyzed

using CamSol (Sormanni et al., 2015) software. Proteins and sequence residues with scores < -

1 were poorly soluble and indicated as potential self-assembly hotspots (red). Scores > 1

characterize highly soluble proteins and sequence residues (blue). TMDs, transmembrane

domains; IMS, intermembrane space; IM, inner membrane.

1061

Figure 1–figure supplement 2. Supersaturated nuclear-encoded mitochondrial protein

aggregate in the cytosol and stimulate growth defects. (**A**) Schematic representation of the

aggregation assay. (**B, C**) Metastable proteins aggregate in the cell, showing SDS-PAGE

analysis of aggregation assay fractions of WT yeast cells that overexpressed $Atp2_{FLAG}$ and

$Atp20_{FLAG}$ (**B**) and $Cox8_{FLAG}$, $Qcr8_{FLAG}$, $pRip1_{FLAG}$, $iRip1_{FLAG}$, $mRip1_{FLAG}$, $Cor1_{FLAG}$,

$Mas1_{FLAG}$, $Cox12_{FLAG}$, and $Qcr6_{FLAG}$ (**C**) for 3 h at 28°C, with 2% galactose with 0.1% glucose

as the carbon source. (**D**) Metastable proteins exhibit a temperature-sensitive phenotype. Each

protein-expressing strain was subjected to consecutive 10-fold dilutions and spotted on

selective medium agar plates with glucose as the main carbon source at 28°C and 37°C. (**E**)

Cartoon representation of Rip1, Sod2, and Mdh1 mitochondrial proteins containing their

presequences. (**F**) Total protein cell extract from WT (*pam16WT*) and *pam16-3* mutant yeast

strains that were grown at a permissive temperature of 19°C and shifted to a restrictive

temperature of 37°C for 0, 1, 3, or 5 h. In **B**, **C**, and **F**, protein samples were separated by SDS-

PAGE and identified by Western blot with specific antisera. Each experiment was repeated

three times. p, presequence protein; m, mature protein; i, intermediate; *, nonspecific; aa, amino

acid.

1078

Figure 2–figure supplement 1. Characterization of gene change that was attributable to

mitochondrial protein import defects. (**A**) Bar plots of the total number of detected gene

transcripts in *pam16-3* compared with WT at the indicated time of heat shock and pRip1

compared with the empty vector (ev) at the indicated expression time by RNA-seq. The

proportions of significant genes (5% FDR) are shown in gray shades (statistically significant),

and the proportions of differentially expressed features in the *pam16-3* and pRip1 samples (5%

FDR; log2FC ± 1) are shown in green (upregulated) and pink (downregulated). The number of

genes in each group is indicated. (**B, C**) KEGG enrichment analysis of differentially expressed

genes for pRip1 *vs*. ev (**B**) and *pam 16-3 vs*. WT (**C**) samples, respectively. The results are

shown as a negative log10 *p* value after Bonferroni correction. Bars in green indicate KEGG

terms that were enriched for upregulated genes. Pink bars represent KEGG terms for

downregulated features. No KEGG terms were identified for down- and upregulated pRip1 and

pam16-3 genes, respectively. I, induction time; HS, heat shock time (in hours). (**D**) Bar plots of

the proportion of mitochondrial genes that encode the high confidence mitochondrial proteome

(HCMP) in each set. The proportions of significant genes (5% FDR) are shown in gray shades

(statistically significant). The proportions of differentially expressed features in the *pam16-3*

and pRip1 samples (5% FDR; log2FC ± 1) are shown in green (upregulated) and pink

(downregulated). The number of genes in each group is indicated. The number of genes in each

fraction is indicated in parentheses. (**E**) Expression analysis of selected molecular chaperone

genes based on RNA-seq analysis in *pam16-3* strains. Cells were grown at a permissive

temperature of 19°C and shifted to a restrictive temperature of 37°C for 1, 4, and 8 h. Dark

shades indicate the expression level in *pam16-3* compared with WT strains. Light shades show

expression changes in WT samples that were triggered by the heat shock response. To estimate

transcriptional changes that were induced by the higher temperature, WT samples that were

incubated at 37°C for 1, 4, and 8 h were compared against untreated WT samples (0h). Arrows

indicate chaperones for which the largest increase in gene levels of molecular chaperones was

observed because of the mitochondrial import defect compared with changes that were only

attributable to heat shock. (**F**) Analysis of expression changes for genes that are related to the

mitoCPR (Weidberg & Amon, 2018) response based on the RNA-seq analysis in *pam16-3*

strain samples as in **A**. Up- and downregulated genes (5% FDR) are shown in green and pink,

respectively. The intensity of the color shades reflects the level of expression change (log2FC).

1110

Figure 2–figure supplement 2. Characterization of gene change that was attributable to

heat shock in WT and the pam16-3 mutant. (**A, B**) Bar plots of the number of changed gene

transcripts in WT (**A**) and *pam16-3* (**B**) for the indicated heat shock time. Values of log2FC ±

1 are shown in green (upregulated) and pink (downregulated). The number of genes in each

group is indicated. (**C, D**) KEGG enrichment analysis for differentially expressed genes for WT

(**C**) and *pam 16-3* (**D**) samples for the indicated heat shock time, respectively. The results are

shown as a negative log10 *p* value after Bonferroni correction. Bars in green indicate enriched

KEGG terms for the upregulated genes. Pink bars represent KEGG terms for downregulated

features.

1120

Figure 3–figure supplement 1. Total protein cell extract changes that were attributable to

mitochondrial import defects. (**A**) Total protein cell extracts from WT yeast cells grown at

24°C and treated with 0, 5, 10, or 15 µM CCCP for 15 min. (**B**) Total protein cell extracts from

WT (YPH499), *pam16-WT*, *pam16-1*, *pam16-3*, *pam18-WT*, *pam18-1*, *mia40-4WT*, *mia40-*

4int, and *mia40-3* grown at 19°C and shifted to 37°C for 6 h. In **A** and **B**, protein samples were

separated by SDS-PAGE and identified by Western blot with specific antisera. Each experiment

was repeated three times.

1128

Figure 3–figure supplement 2. Cellular effects of Hsp42 and Hsp104 overexpression

during mitochondrial import defects. (**A**) Total protein cell extracts from WT yeast cells and

Δhsp42 yeast grown at 24°C and treated with 0 or 15 µM CCCP for 30 min. (**B**) Quantification

of pRip1, pSod2, and pMdh1 from **A**. Quantified data are shown as the mean ± SEM. $n = 3$. (**C**)

Quantification of pRip1, pSod2, and pMdh1 from **Figure 3N**. Quantified data are shown as the

mean ± SEM. $n = 3$. (**D**) Quantification of Hsp42 chaperone from **Figure 3N**. Quantified data

are shown as the mean ± SEM. $n = 3$. (**E**) Total protein cell extracts from WT yeast cells, yeast

cells that expressed Hsp42 under the Gal1 promoter, and *Δhsp42* yeast grown at 24°C. (**F**) Total

protein cell extracts from WT yeast cells and yeast cells that expressed Hsp42 and grown

overnight at 24°C and treated with 0 or 15 µM CCCP for 30 min. (**G**) Ten-fold dilutions of WT

cells that expressed Hsp42 or Hsp104 along with the indicated metastable protein expression

spotted on selective medium agar plates with galactose as the main carbon source at 28°C and

37°C. In **A**, **E**, and **F**, samples were separated by SDS-PAGE and identified by Western blot

with specific antisera. Each experiment was repeated three times. p, presequence protein; m,

mature protein; *, nonspecific. Significance in **B-D**: *$p < 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$, ****p

≤ 0.0001; ns, nonsignificant.

1145

Figure 4–figure supplement 1. Effects of mitochondrial protein import dysfunction on

cellular homeostasis. (**A**) The presence of metastable proteins results in the cytosolic

accumulation of other mitochondrial precursors proteins, showing the SDS-PAGE analysis of

total protein cell extracts from *hsp42-GFP* yeast cells that expressed $Atp2_{FLAG}$, $Cox8_{FLAG}$, or

an empty vector control for 3 h, corresponding to the aggregation assay from **Figure 4D** and

Figure 4–figure supplement 1B. (**B**) SDS-PAGE analysis of aggregation assay fractions of

hsp42-GFP yeast cells that overexpressed $Cox8_{FLAG}$ or an empty vector control for 3 h, with

2% sucrose as the carbon source. insoluble, S_{4k} aggregation assay fraction. (**C**) SDS-PAGE

analysis of aggregation assay fractions of the *pam16-3* mutant yeast strain grown at a permissive

temperature of 19°C to the mid-logarithmic phase. Cells were treated with 75 µM MG132 for

1 h and heat shocked at 37°C for 0, 3, or 5 h, with 2% galactose as the carbon source. (**D, E**)

WT (*pam16-WT*) and *pam16-3* yeast strains that expressed α-Syn WT-GFP (**D**) or A53T-GFP

(**E**) samples that were used for the confocal images presented in **Figure 4F** and **4G** were

subjected to the SDS-PAGE analysis of total protein cell extracts. (**F, G**) Quantification of

average aggregate size (in µm) for WT and *pam16-3* yeast strains that expressed α-Syn WT-

GFP (**F**) or A53T-GFP (**G**) from **Figure 4F** and **4G**, respectively. Quantified data are shown

as the mean ± SEM. *n* = 10. In **A-E**, protein samples were separated by SDS-PAGE and

identified by Western blot with specific antisera. Each experiment was repeated three times. **p*

< 0.05, **p ≤ 0.01, ***p ≤ 0.001, ****p ≤ 0.0001; ns, nonsignificant

1165

Figure 5–figure supplement 1. Scheme and controls for experiments on mitochondrial

dysfunction in C. elegans that resulted in Aβ accumulation. (**A**) RNAi silencing

effectiveness on protein levels upon *dnj-21* RNAi in worms strain [pmyo-3::wrmScarlet +

pmyo::GFP]. (**B**) Scheme of temperature shifts during worm culture in the experiments that are

presented in **Figure 5E-G** and **Figure 5–figure supplement 1D-F**. (**C**) Protein levels in worms

that expressed Aβ peptides upon *dnj-21* RNAi. Worms were cultured with the indicated

temperature shift. $n = 3$ biological replicates. (**D**) Coomassie staining as a loading control for

data that are presented in **Figure 5E**. $n = 3$ biological replicates. (**E**) SDS-PAGE analysis of

aggregate levels in worms that expressed Aβ peptides upon *dnj-21* RNAi. $n = 3$ biological

replicates. Worms were cultured at 20°C or with a temperature shift to 22°C or 25°C. (**F**)

Coomassie staining as a loading control for data that are presented in **E**. $n = 3$ biological

replicates. (**G**) Aβ levels calculated from the SDS-PAGE analysis of aggregates in **E**. The data

are shown as the mean ± SD. $n = 3$. Overall differences between conditions were assessed by

unpaired *t*-test by assuming unequal variance. (**H**) Motility in worms that expressed GFP upon

dnj-21 RNAi. The data are shown as the mean ± SEM. Overall differences between conditions

were assessed by unpaired *t*-test by assuming equal variance. $n = 35$ worms for culture at 20°C.

$n = 60$ worms for culture with 22°C shift. $n = 50$ worms for culture with 25°C shift. *$p < 0.05$,

**$p \leq 0.01$, ***$p \leq 0.001$, ****$p \leq 0.0001$; ns, nonsignificant.

1184

1185   **Supplementary file 2.** Nucleotides used to clone protein coding sequences and CUP1 promoter

1186   into pESC-URA.

1187   **Supplementary file 3.** List of plasmids used in this study.

1188

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
strain, (*S. cerevisiae*)	YPH499	standard laboratory strain	In-house ID: 524	MATa, ade2-101, his3-D200, leu2-D1, ura3-52, trp1-D63, lys2-801
strain, (*S. cerevisiae*)	BY4741	standard laboratory strain	In-house ID: 755	MATa, his3D1, leu2D0, met15D0, ura3D0
strain, (*S. cerevisiae*)	Δ*hsp104*	Euroscarf	In-house ID: 1001	
strain, (*S. cerevisiae*)	*hsp42-GFP*	Invitrogen	In-house ID: 1005	
strain, (*S. cerevisiae*)	*Hsp42-BY4741*	Provided by Dr. Bernd Bukau	In-house ID: 1178	-His selection; Gal1 promoter
strain, (*S. cerevisiae*)	*pam16WT*	doi.org:10.1038/nsmb735	In-house ID: 736	
strain, (*S. cerevisiae*)	*pam16-1*	doi.org:10.1038/nsmb735	In-house ID: 733	
strain, (*S. cerevisiae*)	*pam16-3*	doi.org:10.1038/nsmb735	In-house ID: 734	
strain, (*S. cerevisiae*)	*pam18WT*	doi.org:10.1083/jcb.200308004	In-house ID: 738	
strain, (*S. cerevisiae*)	*pam18-1*	doi.org:10.1083/jcb.200308004	In-house ID: 739	
strain, (*S. cerevisiae*)	*mia40WT*	doi.org:10.1038/sj.emboj.7600389	In-house ID: 398	

strain, (*S. cerevisiae*)	*mia40-3*	doi.org:10.1038/sj.emboj.7600389	In-house ID: 178	
strain, (*S. cerevisiae*)	*mia40-4int*	doi.org:10.1083/jcb.200804095	In-house ID: 739	
strain (*C. elegans*)	worms expressing GFP	Caenorhabditis Genetics Center (CGC)	CL2122 dvIs15 [(pPD30.38) unc-54(vector) + (pCL26) mtl-2::GFP]	Control strain for GM101
strain (*C. elegans*)	worms expressing Aβ and GFP	Caenorhabditis Genetics Center (CGC)	GMC101 dvIs100 [punc-54::A-beta-1-42::unc-54-3'-UTR + mtl-2p::GFP]	
strain (*C. elegans*)	α-syn::YFP	doi: 10.1371/journal.pgen.1000027	OW40 zgIs15 [punc-54::αsyn::YFP]IV	
strain (*C. elegans*)	YFP	doi: 10.1371/journal.pgen.1000027	OW450 rmIs126 [punc-54::YFP]V	Control strain for OW40
strain (*C. elegans*)	[pmyo-3::wrmScarlet+pmyo::GFP]	A strain generated in our laboratory	ACH87: wacIs11 [pmyo-3::mGFP::SL2 gdp-2-wrmScarlet::unc-54-3'UTR, unc-119(+)]	
strain (*Escherichia coli*)	HB101	Caenorhabditis Genetics Center (CGC)	HB101	*E. coli* strain used as a food source for worms
strain (*Escherichia coli*)	HT115(DE3)	Caenorhabditis Genetics Center (CGC)	HT115(DE3)	*E. coli* strain used as a food source for worms in RNAi experiments
strain (*Escherichia coli*)	OP50	Caenorhabditis Genetics Center (CGC)	OP50	*E. coli* strain used as a food source for worms
antibody	Anti-Rip1 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Sod2 (rabbit polyclonal)	Custom made		WB (1:500)

antibody	Anti-Mdh1 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Hsp60 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Pgk1 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Tom70 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Ssa1 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Ssb1 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Ssc1 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Rpl17 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Ccp1 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Qcr8 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Qcr6 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Cox12 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Tom20 (rabbit polyclonal)	Custom made		WB (1:500)

antibody	Anti-Edg1 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Cdc48 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Rpl5 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Dnj21 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Rpn1 (rabbit polyclonal)	Custom made		WB (1:500)
antibody	Anti-Hsp42 (rabbit polyclonal)	Provided by Dr. Bernd Bukau		WB (1:5000)
antibody	Anti-Hsp104 (rabbit polyclonal)	Enzo	ADI-SPA-1040-F	WB (1:1000)
antibody	Anti-GFP (mouse monoclonal)	Sigma	11814460001	WB (1:1000)
antibody	anti-3-PGDH (rabbit polyclonal)	Millipore	ABS571, RRID:AB_2783876	IF(1:1000), WB (1:1000)
antibody	Anti-beta amyloid 1-16 (mouse monoclonal)	Biolegend	803001	WB (1:1000)
antibody	Anti-Tubulin (mouse monoclonal)	Sigma	T9026	WB (1:500)
antibody	Anti-mCherry (anti-wrmScarlet/RFP) ; (rabbit polyclonal)	Abcam	ab167453	WB (1:1000)
antibody	Anti-FLAG (mouse monoclonal)	Sigma	F1804	IF (1:1000) WB (1:1000)

antibody	Alexa Fluor 568 (goat anti-mouse)	Invitrogen	A11031	IF (1:500)
Recombinant DNA reagent (plasmid)	L4440 (control for RNAi in *C. elegans*)	AddGene	#1654	
Recombinant DNA reagent (plasmid)	*dnj-21* RNAi	Recombinant DNA generated in our laboratory	pMJS1 In-house ID: 435p	Plasmid used for silencing of *dnj-21*
Recombinant DNA reagent (plasmid)	Hsp104-pFL46L	doi.org:10.1128/MCB.20.19.7220-7229.2000		Provided by Dr. Magdalena Boguta
Recombinant DNA reagent (plasmid)	α-Syn WT-GFP	doi.org:10.1126/science.1090439		Provided by Dr. Tiago Outeiro
Recombinant DNA reagent (plasmid)	α-Syn A53T-GFP	doi.org:10.1126/science.1090439		Provided by Dr. Tiago Outeiro
Recombinant DNA reagent (plasmid)	Empty vector with FLAG-tag, pCup1	Recombinant DNA generated in our laboratory	pPCh26 In-house ID: 435p	See Supplementary file 3
Recombinant DNA reagent (plasmid)	Cox8	Recombinant DNA generated in our laboratory	pPCh17 In-house ID: 481p	See Supplementary file 3
Recombinant DNA reagent (plasmid)	Atp2	Recombinant DNA generated in our laboratory	pPCh18 In-house ID: 482p	See Supplementary file 3
Recombinant DNA reagent (plasmid)	Cor1	Recombinant DNA generated in our laboratory	pPCh19 In-house ID: 484p	See Supplementary file 3
Recombinant DNA reagent (plasmid)	Mas1	Recombinant DNA generated in our laboratory	pPCh20 In-house ID: 485p	See Supplementary file 3
Recombinant DNA reagent (plasmid)	Qcr6	Recombinant DNA generated in our laboratory	pPCh21 In-house ID: 486p	See Supplementary file 3
Recombinant DNA reagent (plasmid)	Qcr8	Recombinant DNA generated in our laboratory	pPCh22 In-house ID: 487p	See Supplementary file 3

Recombinant DNA reagent (plasmid)	pRip1	Recombinant DNA generated in our laboratory	pPCh23 In-house ID: 488p	See Supplementary file 3
Recombinant DNA reagent (plasmid)	iRip1	RDNA generated in our laboratory	pPCh24 In-house ID: 489p	See Supplementary file 3
Recombinant DNA reagent (plasmid)	mRip1	Recombinant DNA generated in our laboratory	pPCh29 In-house ID: 490p	See Supplementary file 3
Recombinant DNA reagent (plasmid)	Atp20	Recombinant DNA generated in our laboratory	pPCh30 In-house ID: 483p	See Supplementary file 3
Recombinant DNA reagent (plasmid)	Cox12	Recombinant DNA generated in our laboratory doi: 10.1186/s12915-018-0536-1	pPB36.1 In-house ID: 471p	See Supplementary file 3
Sequence-based reagent	Primers	This paper	PCR primers	See Supplementary file 2
commercial assay or kit	RNeasy Mini Kit	Qiagen	74104	
chemical compound, drug	CCCP	Sigma	C2759	
chemical compound, drug	MG132	Enzo	BML-PI102-0005	
software, algorithm	ZEN	Zeiss	ZEN 2012 SP5 FP3 (black)	Confocal Microscopy Data Collection Software
software, algorithm	FastQC	www.bioinformatics.babraham.ac.uk/projects/fastqc		
software, algorithm	Salmon (v0.11.2)	doi.org:10.1038/nmeth.4197		

software, algorithm	DESeq2 version 1.26.0	doi.org:10.1186/s13059-014-0550-8		
other	ProLong Diamond antifade mountant with DAPI	ThermoFisher Scientific	P36962	

1190

Figure 1. Nowicka et al.

Human homologs	Protein name (ORF name)	Localization	Amount of TMDs	Intrinsic residue solubility along amino acid sequence	Intrinsic solubility score	Mitochondria targeting sequence
ATP5B	Atp2 (YJR121W)	peripheral IM	0		0.4783	N-terminal cleavable presequence
ATP5L	Atp20 (YPR020W)	IM	1		0.9556	N-terminal non-cleavable sequence
COX7C	Cox8 (YLR395C)	IM	1		-0.2763	N-terminal cleavable presequence
COX6B1	Cox12 (YLR038C)	IMS	0		1.2939	Cystein motifs
UQCRFS1	Rip1 (YEL024W)	IM	2		0.4783	N-terminal cleavable presequence
UQCRH	Qcr6 (YFR033C)	IMS	0		4.1091	Unknown
UQCRQ	Qcr8 (YJL166W)	IM	1		0.0079	N-terminal non-cleavable sequence
UQCRC1	Cor1 (YBL045C)	IM	0		0.3800	N-terminal cleavable presequence

Figure 1 - figure supplement 1. Nowicka et.al.

Figure 1 - figure supplement 2 Nowicka et al.

Figure 2. Nowicka et al.

Figure 2 - figure supplement 1. Nowicka et al.

Figure 2 - figure supplement 2. Nowicka et al.

Figure 3. Nowicka et al.

Figure 3 - figure supplement 1. Nowicka et al.

Figure 3 - figure supplement 2. Nowicka et al.

Figure 4. Nowicka et al.

Figure 4 - figure supplement 1. Nowicka et al.

Figure 5. Nowicka et al.

Figure 5 - figure supplement 1. Nowicka et al.