Agrochemical control of gene expression using evolved split RNA polymerase

Yuan Yuan[1], Jin Miao[2]

[1] Department of Neurophysiology and Neuropharmacology, Institute of Special Environmental

Medicine and Co-innovation Center of Neuroregeneration, Nantong University, Nantong,

226019, China;

[2] Duke Kunshan University, 8 Duke Avenue, Kunshan, Jiangsu Province, 215316, China

Corresponding Author:

Jin Miao

Duke Kunshan University, 8 Duke Avenue, Kunshan, Jiangsu Province, 215316, China

Email address: jin.miao@dukekunshan.edu.cn

Abstract

Chemical inducible gene expression systems are valuable tools for rational control of gene

expression both for basic research and biotechnology.   However, most chemical inducers are

confined to certain groups of organisms. Therefore, dissecting interaction between different

organisms could be challenging using existing chemical inducible systems. We engineered a

mandipropamid induced gene expression system (Mandi-T7) based on evolved split T7 RNAP

system. As a proof-of-principle, we induced GFP expression in *E. coli* cells grown inside plant

tissue.

Introduction

Chemical inducible gene expression systems are powerful tools that allow conditional control

over gene expression and facilitate study of toxic and essential genes. However, most chemical

inducible systems were often designed to control gene expression in a single group of

organisms in the laboratory condition. Working in complex context, like symbiosis between

plant and bacteria, requires new chemical inducers that are neutral to different groups of

organisms. Agrochemicals are attractive candidates because of their specific mode of action,

- low toxicity, and drug-like physical properties. Mandipropamid is the active ingredient of

oomyceticide REVUS® and has been repurposed to control drought tolerance of plants through

orthogonal control strategy.[1]   Mandipropamid and PYR1[MANDI] protein constitute an engineered

ligand-receptor pair, orthogonal to Abscisic acid (ABA) - PYR1 receptor pair.[1]   The

Mandipropamid - PYR1[MANDI] pair binds protein phosphatase HAB1 or ABI1 as the ABA-PYR1

pair (Figure S1).[1] RNA polymerases which are modulated by protein-protein interaction would

be desirable for coupling with Mandipropamid-sensing module to activate gene transcription.

Split T7 RNA Polymerase (RNAP) has been evolved to be a proximity-dependent biosensor

platform, which could transduce protein-protein interaction of light or small molecule sensing

modules into activation of gene expression under the control of T7 promoter.[2]   ABA inducible

CRISPR/Cas9 gene editing in mammalian cells has been demonstrated using this evolved split

T7 RNAP system.[3]   Engineering Mandipropamid induced gene expression is feasible by

modifying abscisic acid sensing module. Here, we report the repurposing of Mandipropamid to

activate gene expression using the evolved split T7 RNA polymerase (Mandi-T7) and

demonstrate the ability to regulate gene expression of bacteria grown inside plant tissue.

Materials & Methods

Plasmid construction

All genetic parts were synthesized and cloned into vectors by Genscript Biotechnology Inc.

(Nanjing, China). Plasmid pET23a(+) was used to clone the Mandipropamid-T7 RNAP

cassettes. And the backbone of compatible plasmid pCDFduet-1 was used to clone the T7pro-

sfGFP reporter cassette. Sequence of all genetic parts were described in Table S2 and Table

S3.

Mandipropamid responsive assay for evolved split T7 RNA polymerase

Both driver and reporter plasmids were co-transformed into *E. coli* strain Top10 (TIANGEN

biotech., Beijing, China). Single colonies were inoculated 2XYT medium with antibiotics at 37

°C. Overnight culture was transferred to fresh medium with antibiotics at 1:400 ratio.   At the

same time, Mandipropamid (sc-235565, Santa Cruz) or DMSO (solvent control) was added.

The culture was incubated for 6 hours, and 100 µL of each sample was added to a 96 well plate

and mixed with 190 µL 0.1% Tween-80. Both florescence (GFP, Ex: 488, Em: 510; mcherry,

Ex: 587, Em: 630) and OD_{600} was measured by Perkin Elmer Enspire^{TM} 2300 Multilabel Reader.

After the values for medium were subtracted, florescence over OD_{600} was calculated and

compared to that of the DMSO control samples.

Mandipropamid responsive assay for split GFP

Plasmids were transformed into *E. coli* strain BL21(DE3) (TRANSGEN biotech., Beijing, China).

Single colonies were inoculated 2XYT medium with antibiotics at 37 °C. Overnight culture was

transferred to fresh medium at 1:100 ratio.   When OD_{600} reached between 0.4 to 0.6, IPTG

(1mM final) was added to induce expression of GFP1-9. Mandipropamid or DMSO (solvent

control) was also added. The culture was incubated for 3 hours, and 100 µL of each sample

was transferred to a 96 well plate. Both GFP florescence (Ex: 488, Em: 510) and OD_{600} was

measured by Perkin Elmer Enspire^{TM} 2300 Multilabel Reader.   After the values for medium

were subtracted, florescence over OD_{600} was calculated and compared to that of the DMSO

control sample.

Time course fluorescence measurement

E. coli Top10 strain containing Mandi-T7 and reporter plasmids was generated as mentioned

above. Single colonies were inoculated 2XYT medium with antibiotics at 37 °C. Overnight

culture was transferred to fresh medium with antibiotics at 1:200 ratio. When OD_{600} reached

roughly 0.2, aliquots of 500 µL culture was distributed into individual wells of a deep-well plate

or into the hollow septate stems of water spinach (*Ipomoea aquatica*). Mandipropamid (sc-

235565, Santa Cruz) or DMSO (solvent control) was then added. The plate was incubated at

1000 rpm for 6 hours. Samples were harvested after 15 min, 30 min, 1 hour, 2 hour, 3 hour, 4

hour, and 6 hour. 100 µL of each sample was added to a 96 well plate. Both florescence (GFP:

Ex: 488, Em: 510; mcherry: Ex: 587, Em: 630) and OD_{600} was measured by Perkin Elmer

Enspire^{TM} 2300 Multilabel Reader.   After the values for medium were subtracted, florescence

over OD_{600} was calculated.

Western Blot analysis

E.coli culture was harvested at series of time points from deep-well plates using centrifuge.

Samples were boiled in 1X loading buffer for 5 min. Relative protein concentration was

determined by BCA Assay. Total protein was resolved using SDS-PAGE. The protein sample

was transferred to PVDF membranes (Millipore) using Mini Trans-Blot (Bio-Rad). Anti-GFP

(MF090, 1:2000 dilution) and anti-GAPDH (MF091, 1:1000 dilution) antibodies from Mei5bio

were used.

Results and discussion

To simplify circuit design and troubleshooting, we deployed the Mandi-T7 driver module and

reporter module on two compatible plasmids. For the reporter module, expression of

superfolder Green Fluorescent Protein (sfGFP) was driven by the T7 promoter. For the driver

module, we followed the reported architecture of split T7 RNAP.[3] Briefly, we fused PYR1[MANDI]

to the RNAP C-terminal half (T7 RNAPc) and the evolved RNAP N-terminal half (eRNAPn) to

the N-terminal truncated ABI1 (Figure 1A). We used relatively long flexible linker (12 aa) to fuse

PYR1[MANDI] and T7 RNAPc for achieving high-level expression upon induction.[2] We used the

catalytic inactive mutant (D143A) of ABI1 protein (ABI) to minimize unintended effects.[4] To

avoid using additional inducer molecules, we used constitutive promoters to drive the

expression of Mandi-T7. We chose standard promoter of medium strength (J23105) to drive the

expression of PYR1[MANDI]-T7 RNAPc to avoid overwhelming the metabolism.[5] To evaluate the

performance of Mandi-T7 system, we measured the response of the Mandi-T7 to different

concentration of Mandipropamid. The result showed a dose-dependent response, which

saturated at 100 µM Mandipropamid (Figure 1B and Figure S2). A fold-induction of 44X was

achieved, within the same order of magnitude as the 100X induction of the ABA-inducible RNAP

sensor at 100 µM.[3] The difference between the two systems is that ABA sensor can be further

induced to 468X fold change at 10mM.[3]

To understand kinetic characteristics of Mandi-T7 system, we measured fluorescence intensity

over the course of induction for 6 hours. Fluorescence above the background level was

detected at 15 min (Figure 1C).   The induction plateaued after 6 hours (Figure 1C). We also

detected fluorescence signal without induction (Figure 1C), which may be caused by the

residual background activity of the evolved split-T7 polymerase.[3]   We also tested mcherry as

the reporter. The dose-response curve of mcherry showed 140X induction at 100 μM (Figure

S3A). The time course expression analysis showed induction plateaued after 4 hours (Figure

S3B).   The increased fold change for mcherry is consistent with low cellular autofluorescence at

its emission region.[6]   We failed to detect toxic effect of Mandipropamid on *E.coli* growth (Figure

S4).

Figure 1, Engineering Mandi-T7 and its application. A, schematic of Mandi-T7 system and GFP
reporter under control of T7 promoter. Bent arrow: promoter; Box arrow: CDS; Large T-shape:
terminator. B, dose-response curve of Mandi-T7 in *E. coli* Top10. C, inducer response after 6
hours of induction in liquid culture. D, schematic of CP234-Mandi-T7. The original termini of
ABI were linked by a flexible linker. E, dose-response curve of CP234-Mandi-T7 in *E. coli*
Top10. F, time course after induction for 6 hours in the hollow septate stems of water spinach.
126
In a separate attempt, we tried a circular permutated version of ABI protein, which provides

alternative topology for complementation of the split T7 RNAP. We noticed that Ser 234 at the

end of helix 2 is closer to the C terminus of PYR1^{MANDI} than the original N-terminus (Figure S5).

We joined the original N and C termini of ABI by a flexible linker and generated the circular

permutant CP234, for which Ser234 of ABI serves as the new N-terminus (Figure 1D). To test

the retention of the activity to bind PYR1^{MANDI}, we used the tripartite split-GFP based system

because of its low background fluorescence and minimal effect on fusion protein.[7] We

observed concentration dependent increase in GFP florescence after Mandipropamid induction

(Figure S6). This result indicated that ABI-CP234 retained the ability to interact with PYR1^{MANDI}

in the presence of Mandipropamid. Next, we adapted ABI-CP234 to the evolved split T7 RNAP

system (CP234-Mandi-T7), replacing ABI using ABI-CP234.   The result also showed strong

sfGFP expression upon Mandipropamid induction like the Mandi-T7 system, with a fold

induction of 33 (Figure 1E and Figure S2).

To provide a proof-of-principle that Mandi-T7 is useful in complex setting, we evaluated the

ability of Mandi-T7 to induce bacterial gene expression inside the hollow septate stems of water

spinach (*Ipomoea aquatica*).   We followed reporter expression using fluorescence intensity and

Western Blot. The result showed induction after 15 min and induction level kept on increasing

over 6-hour period (Figure 1F, Figure S7). This result suggested that Mandi-T7 has the

potential to enable manipulation of gene expression of plant-associated microorganisms.

In summary, this study provides a new tool for agrochemical control gene expression. We

showed that Mandipropamid inducible gene expression can be readily built based on the

evolved split T7 RNAP system. T7 RNAP is orthogonal and widely applicable to many

prokaryotic and eukaryotic organisms. Mandi-T7 can be further integrated with other tools like

CRISPRi technology. Low toxicity, easy absorption, and cost-effectiveness of Mandipropamid

will give Mandi-T7 leverage in other complex contexts like vector-borne bacterial plant

pathogens. We also tested alternative topology (ABI-CP234) for generating ABI fusion protein

and hope this result might help expand Mandipropamid to other proximity-dependent systems.

CP234-ABI might also be applicable to ABA FRET sensor. The reporter signal in uninduced

conditions could be caused by constitutive expression of Mani-T7 module and low level self-

assembly of split T7 RNAP. Further engineering efforts like directed evolution are needed to

lower the basal expression level and increase the dynamic range. We hope the Mandi-T7

system will expand rational control over gene expression to diverse biological context.

Acknowledgements

We would like to thank Dr. Jinyue Pu and Professor Bryan Dickinson (University of Chicago) for
sharing plasmids and vector information. We want to thank Dr. Xin Zhou (Nantong University)
for technical help with Confocal imaging. We also want to thank Professor Linfeng Huang
(Duke Kunshan University) for advices 163 and generous sharing equipment and reagents.
This work was partly supported by the 164 National Natural Science Foundation of China
(Grant 31500690 to J. M.).
166

References

(1) Park, S. Y., Peterson, F. C., Mosquna, A., Yao, J., Volkman, B. F., and Cutler, S. R. (2015)
Agrochemical control of plant water use using engineered abscisic acid receptors. Nature, 520,
545-548.
(2) Pu, J., Zinkus-Boltz, J., and Dickinson, B.C. (2017) Evolution of a split RNA polymerase as
a versatile biosensor platform. Nat. Chem. Biol., 13, 432-438.
(3) Pu, J., Kentala, K., and Dickinson, B. C. (2018) Multidimensional control of cas9 by evolved
RNA polymerase-based biosensors. ACS Chem. Biol., 13, 431-437.
(4) Liang, F.S., Ho, W.Q., Crabtree, G.R. (2011) Engineering the ABA plant stress pathway for
regulation of induced proximity. Sci Signal. 4, rs2.
(5) Tan, S. I., and Ng, I. S. (2020) New insight into plasmid-driven T7 RNA polymerase in
Escherichia coli and use as a genetic amplifier for a biosensor. ACS Synth. Biol., 9, 613-622.
(6) Telford, W.G., Hawley, T., Subach, F., Verkhusha, V., Hawley, R.G. (2012) Flow cytometry
of fluorescent proteins. Methods, 57, 318-30.
(7) Cabantous, S., Nguyen, H. B., Pedelacq, J. D., Koraïchi, F., Chaudhary, A., Ganguly, K.,
Lockard, M.A., Favre, G., Terwilliger, T.C., and Waldo, G.S. (2013) A new protein-protein
interaction sensor based on tripartite split-GFP association. Sci. Rep., 3, 2854.