1 **Functional characterisation of banana (*Musa* spp.) 2-oxoglutarate-**
2 **dependent dioxygenases involved in flavonoid biosynthesis**
3
4 Short title: banana flavonoid 2-ODDs
5
6

7 Authors:
8 Mareike Busche, Christopher Acatay, Bernd Weisshaar, Ralf Stracke*
9
10 Genetics and Genomics of Plants, Faculty of Biology, Bielefeld University, 33615 Bielefeld,
11 Germany
12
13
14 *Corresponding author:
15 Ralf Stracke
16 ralf.stracke@uni-bielefeld.de

17
18
19
Keywords: banana, specialised metabolites, flavanone 3-hydroxylases, flavonol synthase,
anthocyanidin synthase

Abstract

Bananas (*Musa*) are monocotyledonous, perennial plants that are well-known for their edible

fruits. Their cultivation provides food security and employment opportunities in many countries.

Banana fruits contain high levels of minerals and phytochemicals, including flavonoids, which

are beneficial for human nutrition. To broaden the knowledge on flavonoid biosynthesis in this

major crop plant, we aimed to identify and functionally characterise selected structural genes

encoding 2-oxoglutarate-dependent dioxygenases, involved in the formation of the flavonoid

aglycon.

Musa candidates genes predicted to encode flavanone 3-hydroxylase (F3H), flavonol synthase

(FLS) and anthocyanidin synthase (ANS) were assayed. Enzymatic functionalities of the

recombinant proteins were confirmed *in vivo* using bioconversion assays. Moreover,

transgenic analyses in corresponding *Arabidopsis thaliana* mutants showed that *Musa*F3H,

*Musa*FLS and *Musa*ANS were able to complement the respective loss-of-function phenotypes,

thus verifying functionality of the enzymes *in planta*.

Knowledge gained from this work provides a new aspect for further research towards genetic

engineering of flavonoid biosynthesis in banana fruits to increase their antioxidant activity and

nutritional value.

Introduction

Banana (*Musa* spp.) plants are well-known for their edible fruit and serve as a staple food crop

in Africa, Central and South America (Arias et al., 2003). With more than 112 million tons

produced in 2016, bananas are among the most popular fruits in the world and provide many

employment opportunities (FAO, 2019). Furthermore banana fruits are rich in health promoting

minerals and phytochemicals, including flavonoids, a class of plant specialised metabolites

which contribute to the beneficial effects through their antioxidant characteristics (Forster et

al., 2003; Wall, 2006; Singh et al., 2016). Flavonoid molecules share a C6-C3-C6 aglycon core,

which can be reorganised or modified, e.g. by oxidation or glycosylation (Tanaka et al., 2008;

Le Roy et al., 2016). Modifications at the central ring structure divide flavonoids into ten major

subgroups (i.e. chalcones, aurones, flavanones, flavones, isoflavones, dihydroflavonols,

flavonols, leucoanthocyanidins, anthocyanidins, and flavan-3-ols). The diversity in chemical

structure is closely related to diverse bioactivities of flavonoids in plant biology and human

nutrition (Falcone Ferreyra et al., 2012). For example, anthocyanins are in many cases well

known to colour flowers and fruits to attract animals and thus promoting pollination and

dispersion of seeds (Ishikura and Yoshitama, 1984; Gronquist et al., 2001; Grotewold, 2006).

Flavonols can interact with anthocyanins to modify the colour of fruits (Andersen and Jordheim,

2010) and play a prominent role in protection against UV-B irradiation (Li et al., 1993) and also

in plant fertility (Mo et al., 1992).

Many researchers have attributed positive effects on human health to flavonoids: e.g.

antigenotoxic (Dauer et al., 2003), anticarcinogenic and antioxidative (Kandil et al., 2002)

effects, as well as the prevention of cardiovascular diseases have been suggested

(summarised inPerez-Vizcaino and Duarte, 2010). Additionally, Sun *et al.* (2019) suggest an

involvement of flavonoids in the plant defence against the tropical race 4 (TR4) of the *Musa*

Fusarium wilt (commonly known as 'panama disease') pathogen *Fusarium oxysporum* f. sp.

cubense (Foc), which is a threat to the global banana production. It is certainly interesting to

take a closer look at the biosynthesis of flavonoids in *Musa*.

Flavonoids are derived from the amino acid L-phenylalanine and malonyl-coenzyme A. Their

biosynthesis (Figure 1) has been analysed in many different species including the

dicotyledonous model plant *Arabidopsis thaliana* (*A. thaliana*) (Hahlbrock and Scheel, 1989;

Lepiniec et al., 2006) and the monocotyledonous crop plants *Zea mays* (*Z. mays*) (summarised

in Tohge et al., 2017) and *Oryza sativa* (*O. sativa*) (summarised in Goufo and Trindade, 2014).

The first committed step, catalysed by the enzyme chalcone synthase (CHS), is the formation

of naringenin chalcone from p-coumaroyl CoA and malonyl CoA (Kreuzaler and Hahlbrock,

1972). A heterocyclic ring is introduced during the formation of naringenin (a flavanone) from

naringenin chalcone, which can occur spontaneously or catalysed by chalcone isomerase

(CHI) (Bednar and Hadcock, 1988). The enzyme flavanone 3-hydroxylase (F3H or FHT)

converts flavanones to dihydroflavonols by hydroxylation at the C-3 position (Forkmann et al.,

1980). Alternatively, flavanones can be converted to flavones by flavone synthase I or II (FNS

I or II) (Britsch, 1990). Flavonol synthase (FLS) catalyses the conversion of dihydroflavonols

to the corresponding flavonols by introducing a double bond between C-2 and C-3 (Forkmann

et al., 1986; Holton et al., 1993). Moreover, dihydroflavonols can be converted to

leucoanthocyanidins by dihydroflavonol 4-reductase (DFR) (Heller et al., 1985), which

competes with FLS for substrates (Luo et al., 2016). Anthocyanidin synthase (ANS, also

termed leucoanthocyanidin dioxygenase, LDOX) converts leucoanthocyanidins to

anthocyanidins (Saito et al., 1999).

The flavonoid biosynthesis enzymes belong to different functional classes (summarised in

Winkel, 2006): polyketide synthases (e.g. CHS), 2-oxoglutarate-dependent dioxygenases

(2-ODD, e.g. F3H, ANS, FLS, FNS I), short-chain dehydrogenases/reductases (e.g. DFR),

aldo-keto reductases (e.g. chalcone reductase, CHR) and cytochrome P450 monooxygenases

(e.g. FNS II, F3'H). Flavonoid biosynthesis is evolutionary old in plants (Wen et al., 2020) and,

although differences exist, quite similar in dicots like *A. thaliana* and monocots like *Musa* .

In the present study, we focus on the functional class of flavonoid aglycon forming enzymes,

namely the 2-ODDs. 2-ODDs occur throughout the kingdom of life and play important roles in

many biological processes including oxygen sensing and DNA repair (Jaakkola et al., 2001;

Trewick et al., 2002). They are a class of non-heme iron-containing enzymes, which require

2-oxoglutarate, Fe^{2+} and ascorbate for substrate conversion (summarised in Prescott and

John, 1996).

Figure 1: Schematic, simplified illustration of the core flavonoid aglycon biosynthesis pathway in plants.

Chalcone synthase (CHS), the first committed enzyme in flavonoid biosynthesis, connects p-coumaroyl-CoA and malonyl-CoA, forming naringenin chalcone. Chalcone isomerase (CHI) introduces the heterocyclic ring. The resulting naringenin is converted to dihydroflavonol by flavanone 3-hydroxylase (F3H). Dihydroflavonols are converted to flavonols by flavonol synthase (FLS). Alternatively, dihydroflavonol 4-reductase (DFR) can reduce dihydroflavonols to the corresponding leucoanthocyanidin, which is converted to anthocyanidins by the activity of anthocyanidin synthase (ANS). 2-oxoglutarate-dependant oxygenases are given in bold underlined. Enzymes in the focus of this work are highlighted by grey boxes.

Plant 2-ODDs share conserved amino acid residues which coordinate ferrous iron binding

(HxDxnH) and binding of 2-oxoglutarate (RxS) (Cheng et al., 2014). They can be divided into

three distinct evolutionary classes, DOXA, DOXB and DOXC, based on amino acid similarity

(Kawai et al., 2014). The 2-ODDs involved in specialised metabolism were classified into the

DOXC class. Yet, all flavonoid biosynthesis related 2-ODDs were found in the DOXC

subclades 28 (F3H and FNS I) or 47 (FLS and ANS) (Kawai et al., 2014). In some cases, the

high amino acid similarity between the different enzymes leads to overlapping functions. For

example, FLSs from *Ginkgo biloba* (*G. biloba*) (Xu et al., 2012) and *O. sativa* (Park et al., 2019)

can accept flavanones and dihydroflavonols as substrates. Also, ANSs from *A. thaliana*

(Stracke et al., 2009) and *Malus domestica* (*M. domestica*) (Yan et al., 2005) can catalyse the

formation of flavonol glycosides.

Based on the *Musa acuminata* (*M. acuminata*) genome sequence annotation (DHont et al.,

2012), Pandey *et al.* (2016) identified 28 putative *Musa* flavonoid biosynthesis enzymes. This

includes seven 2-ODD-type enzymes, namely two F3H, four FLS and one ANS, while no

suitable FNS I candidate was identified.. While the expression of the respective genes and

correlations of expression to flavonoid metabolite accumulation have been studied, the

functionality of these enzymes has not been analysed until now.

Here, we describe the sequence-based and functional characterisation of 2-ODDs from *Musa*.

Functionalities of recombinant *Musa* enzymes were analysed using *in vivo* bioconversion

assays and by *in planta* complementation of corresponding *A. thaliana* loss-of-function

mutants. This resulted in the experimental confirmation that the *Musa* flavonoid 2-ODDs

studied have the predicted functions. The presented results contribute to the understanding of

flavonoid biosynthesis in *Musa*. They provide a strong basis for further research to enhance

the efficiency of flavonoid production in banana, in order to increase the fruits' health promoting

effects.

Materials and Methods

Plant material

Banana plants (Grand Naine, plantain) for RNA extraction were grown in the field in Lucknow,

India. *Musa* gene annotation identifiers refer to the study of Martin et al. (2016). Columbia-0

(Col-0, NASC ID N1092) and Nössen-0 (Nö-0, NASC ID N3081) were used as wildtype

controls. The *A. thaliana* mutants *tt6-2* (*f3h*, GK-292E08, Col-0 background) (Appelhagen et

al., 2014) and *ans/fls1-2* (synonym *ldox/fls1-2, ldox: SALK_028793,* Col-0 background; *fls1-2:*

RIKEN_PST16145, Nö-0 background) (Stracke et al., 2009) were used for complementation

experiments.

Phylogenetic analysis

Multiple protein sequence alignments were created with MAFFT v7 (Katoh and Standley, 2013)

using default settings. The Approximately-Maximum-Likelihood phylogenetic tree of 33 plant

2-ODDs with proven F3H, FNS I, FLS and ANS functionality (Supplementary Table S1) and

seven *Musa* 2-ODDs was constructed as described by Pucker *et al.* (2020b): MAFFT

alignments were cleaned with pxclsq (Brown et al., 2017) and the tree was constructed with

FastTree v2.1.10 using the WAG+CAT model (Price et al., 2010). The tree was visualised with

Interactive Tree Of Life (Letunic and Bork, 2019), branch lengths were neglected.

Expression analysis

Expression data for *Musa* 2-ODD genes was extracted from previous analyses (Pucker et al.,

2020a). Short Read Archive IDs can be found in Supplementary Table S2.

Total RNA extraction, cDNA synthesis and molecular cloning

Isolation of RNA from different *Musa* plant organs (leaf, pseudostem, bract, fruit peel, fruit pulp)

was performed according to a protocol from Asif et al. (2000). cDNA synthesis was performed

from 1 µg total RNA using the ProtoScript® First Strand cDNA Synthesis Kit (New England

Biolabs, NEB) with the provided random primer mix according to the suppliers' instructions.

Amplification of predicted full length CDSs (Pandey et al., 2016) were done using Q5® High-

Fidelity DNA polymerase (NEB) and gene-specific primers (Supplementary Table S3)

according to standard protocols. Creation of full-length coding sequence constructs (CDS) was

performed using the GATEWAY® Technology (Invitrogen). *MusaF3H1* (Ma02_t04650),

MusaF3H2 (Ma07_t17200), *MusaFLS1* (Ma03_t06970), *MusaFLS3* (Ma10_t25110) and

MusaANS (Ma05_t03850) CDSs were successfully amplified on different cDNA pools. The

resulting PCR products were recombined into pDONR™/Zeo (Invitrogen) with BP clonase

(Invitrogen) resulting in Entry plasmids, which were sequenced by Sanger technology (Sanger

et al., 1977) on 3730XL sequencers using BigDye terminator v3.1 chemistry (Thermo Fisher).

Entry plasmids for *AtF3H*, *AtFLS1* and *AtMYB12* were available from previous studies (Preuss

et al., 2009; Stracke et al., 2017). The full-length CDSs were introduced from the Entry

plasmids into the inducible *E. coli* expression vector pDEST17 (Invitrogen) and the binary

expression vector pLEELA (Jakoby et al., 2004) using GATEWAY LR-reaction (Invitrogen).

Heterologous expression in *E. coli*

pDEST17-based plasmids containing *proT7*-RBS-6xHis-CDS-*T7term* expression cassettes

(*proT7: T7 promoter*, RBS: ribosome binding site, 6xHis: polyhistidine tag, *T7term: T7*

transcription terminator) were transformed into BL21-AI cells (Invitrogen). Cultures were grown

in LB to an OD_{600} of about 0.4 and expression was induced with 0.2 % L-arabinose.

F3H and FLS bioconversion assay in *E. coli*

The enzyme assay was performed using 20 mL *E. coli* cultures expressing the respective constructs right after induction with L-arabinose. 100 µL Naringenin (10 mg/mL), 50 µL 2-oxoglutaric acid, 50 µL $FeSo_4$ and 50 µL 1 M sodium ascorbate were added. The cultures were incubated at 28 °C overnight. To extract flavonoids, 1 mL was removed from each culture and mixed with 200 µL ethyl acetate by vortexing for 30 s. After centrifugation for 2 min. at 14,000 g, the organic phase was transferred into a fresh reaction tube. Samples were taken after 0 h, 1 h, 2 h, 3 h, 4 h and 24 h. Flavonoid content was analysed by high-performance thin-layer chromatography (HPTLC). Naringenin (Sigma), dihydrokaempferol (Sigma) and kaempferol (Roth) were dissolved in methanol and used as standards. 3 µL of each methanolic extract were spotted on a HPTLC Silica Gel 60 plate (Merck). The mobile phase was composed of 50 % chloroform, 45 % acetic acid and 5 % water. Flavonoid compounds were detected as described previously (Stracke et al., 2007), using diphenylboric acid 2-aminoethyl ester (DPBA) and UV-light (Sheahan and Rechnitz, 1992).

Agrobacterium*-mediated transformation of *A. thaliana

T-DNA from pLEELA-based plasmids containing *2xpro35S*-driven *Musa2-ODDs*, were transformed into *A. thaliana* plants via *Agrobacterium tumefaciens* (Agrobacterium, GV101::pMP90RK, (Koncz and Schell, 1986)) mediated gene transfer using the floral dip method (Clough and Bent, 1998). Successful T-DNA integration was verified by BASTA selection and PCR-based genotyping.

Flavonoid staining of *A. thaliana* seedlings

In situ visualisation of flavonoids in norflurazon-bleached *A. thaliana* seedlings was performed according to Stracke *et al.* (2007) using DPBA/Triton X-100 and epifuorescence microscopy.

Analysis of flavonols in methanolic extracts

Flavonol glycosides were extracted and analysed as previously described (Stracke et al., 2009). *A. thaliana* rosette leaves were homogenised in 80 % methanol, incubated for 15 min. at 70 °C and centrifuged for 10 min. at 14,000 g. Supernatants were vacuum-dried. The dried pellets were dissolved in 80 % methanol and analysed by HPTLC on a Silica Gel 60 plate (Merck). Ethyl acetate, formic acid, acetic acid, water (100:26:12:12) were used as a mobile phase. Flavonoid compounds were detected as described above.

Determination of anthocyanin content

To induce anthocyanin production, *A. thaliana* seedlings were grown on 0.5 MS plates with 4 % sucrose and 16 h of light illumination per day at 22 °C. Six day old seedlings were used

224 to photometrically quantify anthocyanins as described by Mehrtens et al. (2005). All samples
225 were measured in three independent biological replicates. Error bars indicate the standard
226 error of the average anthocyanin content.
227
228
Results
Creation of cDNA constructs and sequence based characterisation of *Musa* flavonoid
2-ODDs
Previously described putative *Musa* flavonoid biosynthesis enzymes encoded in the
M. acuminata Pahang DH reference genome sequence were used. This included two F3Hs,
four FLSs and one ANS, which were classified as 2-ODDs. To functionally characterise these
Musa enzymes, we amplified the corresponding CDSs from a cDNA template collection
derived from different *Musa* organs, using primers designed on the Pahang DH reference
genome sequence. The successfully amplified cDNAs of *MusaF3H1* and *MusaF3H2* were
derived from plantain pulp, *MusaFLS1* from Grand Naine young leaf, *MusaFLS3* on Grand
Naine bract, and *MusaANS* on Grand Naine peel. Unfortunately, we were not able to amplify
MusaFLS2 and *MusaFLS4* from our template collection.
Comparison of the resulting 2-ODD cDNA sequences with the reference sequence revealed
several single-nucleotide polymorphisms (SNPs) (Supplementary File S1). The derived amino
acid sequences show close similarity to other plant 2-ODD proteins known to be involved in
flavonoid biosynthesis (Supplementary File S2). The amino acids well known to coordinate
ferrous iron binding (HxDxnH) and binding of 2-oxoglutarate (RxS) are also conserved in the
Musa 2-ODDs. Additionally, the *At*FLS1 residues which have been shown to be involved in
flavonoid substrate binding (Supplementary File S2) are conserved. F293 is conserved in all
Musa 2-ODD proteins, F134 and K202 are found in the *Musa*FLSs and *Musa*ANS and E295
is conserved in *Musa*ANS.
To analyse the evolutionary relationship between *Musa* 2-ODDs and 33 known flavonoid
biosynthesis related 2-ODDs, a phylogenetic tree was built (Figure 2). The phylogenetic tree
revealed two distinct clades, which correspond to the DOXC28 and DOXC47 classes of the
2-ODD superfamily. In the F3H and FNS I-containing DOXC28 class, *Musa*F3H1 and
*Musa*F3H2 cluster with other F3Hs from monocotyledonous plants. In the FLS and ANS-
containing DOXC47 class, the *Musa*ANS clusters with ANSs from monocotyledonous plants,
while the FLSs from monocotyledonous plants do not form a distinct group, although the
*Musa*FLSs are in proximity to *Zm*FLS and *Os*FLS.
Our analyses clearly show that the CDSs identified for *Musa*F3H1, *Musa*F3H2, *Musa*FLS1,
*Musa*FLS3 and *Musa*ANS have the potential to encode functional flavonoid 2-ODDs.
260

Figure 2: Rooted Approximately-Maximum-Likelihood (ML) phylogenetic tree of 2-ODDs involved in flavonoid biosynthesis.

Additional to the ODDs from *Musa* (marked with black dots), 33 enzymes with proven F3H, FNS I, FLS or ANS activity were included. Different grey scales indicate the F3H, FNS I, FLS and ANS clades. *At*GA3ox1 (gibberellin 3 beta-hydroxylase 1) was used as an outgroup. DOXC28 and DOXC47 classes are marked with black triangles.

Expression profiles of *Musa 2-ODD* genes

Analysis of the expression patterns of the genes studied was performed using published RNA-Seq data. We obtained normalised RNA-Seq read values for *Musa* 2-ODD genes from several organs and developmental stages (Table 1). *MusaF3H1* and *MusaF3H2* are expressed in almost all analysed organs and developmental stages. *MusaF3H1* expression is highest in early developmental stages of pulp (S1, S2), followed by intermediate developmental stages of peel (S2, S3). *MusaF3H2* shows highest expression in adult leaves.

MusaFLS2 was not expressed in any of the analysed samples. All four *MusaFLS* genes show

low or no expression in seedlings and embryonic cell suspension. *MusaFLS1* shows variance

in transcript abundance with particularly high levels in peel (S2) and in young and adult leaves.

MusaFLS3 and *MusaFLS4* transcript levels are comparatively constant and low, with highest

transcript abundance in root and peel (S1 and S3 respectively). While *MusaFLS1* transcript

abundance is highest in adult leaf, *MusaFLS3* and *MusaFLS4* lack expression in this organ.

MusaANS shows highest transcript abundance in pulp (S1, S2, S4) and peel (S3) and very

low expression in embryogenic cell suspension, seedlings and leaves.

Table 1: Expression profiles of *Musa2*-ODD genes in different organs and developmental stages based on RNA-Seq data. Expression levels are given in fragments per kilo base per million mapped reads (FPKM). Different grey scales indicate expression levels with higher levels given in darker grey. S1-S4 indicate stages of pulp and peel developments from immature to ripe (Lu et al., 2018).

gene	embryogenic	seedling	root	leaf	young leaf	adult leaf	old leaf	pulp	pulp S1	pulp S2	pulp S3	pulp S4	peel	peel S1	peel S2	peel S3	peel S4
MusaF3H1	3	9	80	44	73	27	3	170	385	191	43	60	82	46	140	104	36
MusaF3H2	21	0	23	57	39	171	7	32	96	14	3	14	19	7	27	33	9
MusaFLS1	6	0	4	189	91	643	16	2	2	1	2	2	29	3	99	7	7
MusaFLS2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
MusaFLS3	0	0	32	16	17	0	0	16	15	22	10	17	22	27	16	20	25
MusaFLS4	0	0	19	4	0	0	0	10	11	4	12	13	15	16	11	19	14
MusaANS	1	3	90	41	79	3	1	255	559	298	26	137	73	47	50	163	30

_Musa_F3H1 and _Musa_F3H2 are functional flavanone 3-hydroxylases

To confirm F3H activity *in vivo*, we used a bioconversion assay. *MusaF3H1* and *MusaF3H2*

were heterologously expressed in *E. coli* and the bacterial cultures were fed with naringenin

as substrate of F3H (Figure 1). Both recombinant proteins, *Musa*F3H1 and *Musa*F3H2, were

able to convert naringenin to dihydrokaempferol (DHK) in the presence of 2-oxoglutarate and

ferrous iron (Figure 3A). After 24 h of incubation, only small amounts of naringenin remained

un-converted. Conversion of DHK to kaempferol by *Musa*F3H1 or *Musa*F3H2 was not

observed. For further *in planta* analysis we chose a complementation assay with an *A. thaliana*

f3h single mutant *(tt6-2)* which expresses FLS and ANS but no F3H. *MusaF3H1* or *MusaF3H2*

were expressed in *tt6-2* plants under the control of the constitutive 2x35S promoter. The

accumulation of flavonol glycosides was analysed in herbizide-bleached seedlings using

DPBA-staining (Figure 3B). While Col-0 wildtype seedlings appeared orange under UV light, indicating the accumulation of flavonol glycosides, *tt6-2* seedlings showed a red fluorescence. *tt6-2* mutants expressing *MusaF3H1* or *MusaF3H2* were able to complement the mutant phenotype, showing the characteristic orange flavonol glycoside fluorescence of the wild type. These results demonstrate that *Musa*F3H1 and *Musa*F3H2 are functional enzymes with F3H activity and are able to catalyse the conversion of naringenin to dihydrokaempferol.

Figure 3: *Musa*F3H1 and *Musa*F3H2 are functional flavanone 3-hydroxylases.
(A) HPTLC analysis of F3H bioconversion assays using extracts from *E. coli* expressing recombinant *Musa*F3H1 or *Musa*F3H2. The substrate naringenin (N) and the product dihydrokaempferol (DHK) were used as standards. *At*FLS1 was used as a positive control. *At*MYB12 was used as a negative control. (B) Analysis of *tt6-2* mutant seedlings complemented with *Musa*F3H1 and *Musa*F3H2. Col-0 wildtype and *tt6-2* were used as controls. Representative pictures of DPBA-stained seedlings under UV light are given. Yellow fluorescence indicates flavonol glycoside accumulation. The different numbers indicate individual transgenic lines.

MusaFLS1 and MusaFLS3 are functional flavonol synthases

To confirm FLS activity *in vivo*, the enzymes were assayed in a coupled bioconversion experiment (Figure 4A). Two *E. coli* cultures expressing *MusaF3H1* and a *MusaFLS* were mixed and fed with naringenin in the presence of 2-oxoglutarate and ferrous irons. Here, *Musa*F3H1 converts naringenin to DHK, thus providing the substrate for the FLS enzyme to be tested. While *Musa*FLS1 was found to be able to convert DHK to kaempferol under the assay conditions, *Musa*FLS3 was not. We checked the expression of recombinant *Musa*FLS1 and *Musa*FLS3 in the *E. coli* cultures by SDS-PAGE and found both *Musa*FLS proteins being

expressed at similar levels (Supplementary File S3). To further analyse the enzymatic
properties *in planta*, *MusaFLS1* or *MusaFLS3* were expressed in the flavonol and anthocyanin
deficient *A. thaliana ans/fls1-2* double mutant. HPTLC analyses of methanolic extracts from
rosette leaves from greenhouse-grown plants (Figure 4B) showed that wildtype plants (Col-0,
Nö-0) contained flavonol glycosides, including the prominent derivatives K-3R-7R, K-3G-7R
and K-3[G-R]-7R. *ans/fls1-2* plants accumulated several dihydroflavonols but did not show
flavonol derivatives. *ans/fls1-2* mutants transformed with 2x35S::*MusaFLS1* or
2x35S::*MusaFLS3* constructs were able to form several flavonol derivatives. Nevertheless,
intensities and accumulation patterns of flavonol glycosides varied. We also analysed if
*Musa*FLS1 and *Musa*FLS3 are able to complement the anthocyanin-deficiency of the *ans/fls1-*
2 double mutant. For this, seedlings were grown on 4 % sucrose to induce anthocyanin
accumulation. As shown in Figure 4C, 6-day-old Col-0 and Nö-0 seedlings were able to
accumulate red anthocyanin pigments, while *ans/fls1-2* transformed with *Musa*FLS1 or
*Musa*FLS3 did not show visible anthocyanins. These results confirm that *Musa*FLS1 and
*Musa*FLS3 are functional proteins with FLS activity and are enzymes able to catalyse the
conversion of dihydroflavonol to flavonol.

Figure 4: *Musa*FLS1 and *Musa*FLS3 are functional flavonol synthases.

(A) HPTLC analysis of a FLS bioconversion assays using extracts from *E. coli* expressing recombinant *Musa*FLS1 or *Musa*FLS3. The F3H substrate naringenin (N), the FLS substrate dihydrokaempferol (DHK) and the product kaempferol (K) were used as standards. *At*FLS1 served as a positive control and *At*MYB12 was used as a negative control. (B) Flavonol glycoside accumulation in *Musa*FLS-complemented *ans/fls1-2* seedlings, analysed by HPTLC analysis. Col-0, Nö-0 (both wildtype) and *ans/fls1-2* were used as controls. Bright green spots belong to derivates of kaempferol, orange spots are derivates of quercetin and faint blue shows sinapate derivatives. Dark green and yellow spots indicate DHK and DHQ, respectively. G, glucose; K, kaempferol; Q, quercetin; R, rhamnose (C) Representative pictures of anthocyanin accumulation in 6-day-old *Musa*FLS-complemented *ans/fls1-2* seedlings growing on 4 % sucrose.

MusaANS is a functional anthocyanidin synthase

We analysed *Musa*ANS functionality in a complementation assay with the *ans/fls1-2* *A. thaliana* double mutant. To examine the ability of *Musa*ANS to complement the *ans/fls1-2* anthocyanin deficiency phenotype, seedlings were grown on anthocyanin-inducing media. Anthocyanin accumulation was analysed visually and quantified photometrically (Figure 5A-B). While wildtype seedlings showed red pigmentation, the *ans/fls1-2* seedlings did not. *ans/fls1-2* seedlings expressing *Musa*ANS showed accumulation of anthocyanins. The anthocyanin content in the complemented seedlings was strongly increased compared to *ans/fls1-2* knockout plants, indicating ANS activity. Furthermore, we analysed the ability of *Musa*ANS to complement the flavonol deficiency phenotype of *ans/fls1-2* plants. For this, methanolic extracts of seedlings were analysed by HPTLC followed by DPBA-staining (Figure 5C). While wildtype seedlings accumulated several kaempferol and quercetin glycosides, *ans/fls1-2* mutants expressing *Musa*ANS showed a flavonoid pattern identical to the *ans/fls1-2* mutant, accumulating dihydroflavonol derivatives, but no flavonol derivatives. These results indicate, that *Musa*ANS is a functional enzyme with ANS activity and is able to catalyse the conversion of leucoanthocyanidin to anthocyanidin.

Figure 5: *Musa*ANS is a functional anthocyanin synthase.
Analysis of *ans/fls1-2* double mutant seedlings complemented with 2x35S driven *MusaANS*
demonstrate *in planta* ANS functionality of *Musa*ANS by anthocyanin accumulation. The
different numbers indicate individual transgenic lines. (A-B) Sucrose induced anthocyanin
accumulation in 6-day-old *A. thaliana* seedlings. (A) Representative pictures of seedlings and
(B) corresponding relative anthocyanin content. Error bars indicate the standard deviation for
three independent measurements. (C) *Musa*ANS does not show *in planta* FLS activity.
Flavonol glycoside accumulation in *Musa*ANS-complemented *ans/fls1-2* seedlings, analysed
by HPTLC analysis. Col-0, Nö-0 (both wildtype) and *ans/fls1-2* were used as controls. Bright
green spots belong to derivates of kaempferol, orange spots are derivates of quercetin and
faint blue shows sinapate derivatives. Dark green and yellow spots indicate DHK and DHQ,
respectively. G, glucose; K, kaempferol; Q, quercetin; R, rhamnose

Discussion

In this study, we isolated several cDNAs of *in silico* annotated 2-ODD-type flavonoid biosynthesis enzyme coding genes from *Musa* and tested the *in vivo* functionality of the encoded proteins in *E. coli* and *A. thaliana*.

***Musa*F3H1 and *Musa*F3H2**

In vivo E. coli bioconversion assays revealed that *Musa*F3H1 and *Musa*F3H2 can covert naringenin to dihydrokaempferol. Moreover, *Musa*F3H1 and *Musa*F3H2 are able to complement the loss-of-function phenotype of *A. thaliana tt6-2* seedlings, showing *in planta* F3H-activity. Therefore we conclude that *Musa*F3H1 and *Musa*F3H2 are functional flavanone 3-hydroxylases.

Previous studies annotated *Ma02_g04650* and *Ma07_g17200* as genes encoding flavanone 3-hydroxylases (Martin et al., 2016; Pandey et al., 2016; Pucker et al., 2020b). However, the automatic approach developed by Pucker *et al*. (2020b) also considered *Ma07_g17200* as a candidate to encode a FNS I enzyme. For a parsley F3H protein it has been found that a replacement of three amino acids was sufficient to cause FNS I side activity and seven amino acid exchanges almost lead to a complete change in enzyme activity towards FNS I functionality (Gebhardt et al., 2007). These findings underline the particular high similarity between FNS I and F3H. A closer inspection of the deduced peptide sequence of *Ma07_g17200* gene revealed a lack of conservation of amino acid residues known to be relevant for FNS I function (i.e. T106M, T115I, I116V, F131I, E195D, I200V, V215L, R216K). Together with the confirmed F3H activity this indicates that the classification of *Ma07_g17200* as a FNS I encoding gene was inaccurate. Even though Ma07_p17200 is most likely no functional FNS I, flavone derivatives have been identified in *Musa* (Fu et al., 2018). In *Gerbera* and *Glycine max*, FNS II is responsible for the formation of flavones (Martens and Forkmann, 1999; Fliegmann et al., 2010). A candidate *Musa*FNS II, encoded by *Ma08_g26160*, has been identified (Pucker et al., 2020b) and is probably responsible for the accumulation of flavones in *Musa*. Since FNS II enzymes are NADPH-dependent cytochrome P450 monooxygenases (Jiang et al., 2016), the putative *Musa*FNS II (encoded by *Ma08_g26160*) was not studied in this work.

Our expression study that was based on published RNA-Seq data (Pucker et al., 2020a), revealed highest *MusaF3H1* transcript abundance in early developmental stages of pulp and intermediate stages of peel development, while highest *MusaF3H2* expression was found in adult leaves (Table 1). This is, at first glance, in contrast to the quantitative real-time PCR (qRT-PCR)-derived expression data presented by Pandey et al. (2016), reporting high transcript levels of *MusaF3H1* and *MusaF3H2* in young leaves and bract. In this study, the authors also show increased *MusaF3H1* expression in pseudostem, root and ripe pulp,

compared to *MusaF3H2*. These discrepancies are probably due to the different growth
conditions, germplasms/cultivars and sampling timepoints in the generation of the expression
data and cannot be reasonably analysed further at this point. It should be noted, however, that
MusaF3H2 expression in leaves of plantlets was found to be increased following treatment
with the phytohormone methyl jasmonate (MJ), while *MusaF3H1* expression was not (Pandey
et al., 2016). As MJ is involved in various regulatory processes, including the response against
biotic and abiotic stresses (summarised inCheong and Choi, 2003), the MJ-dependent
induction of *MusaF3H2* expression could imply that the *Musa*F3H2 enzyme plays a specialised
role in the formation of flavonoids in response to stresses. Such stress response induction of
F3H encoding genes was previously shown for a *F3H* gene from the dessert plant *Reaumuria*
soongorica, which is induced by UV-light (Liu et al., 2013) and two *F3H* genes from *Camellia*
sinensis (*C. sinensis*), which are induced by UV light and by treatment with abscisic acid (ABA)
or sucrose (Han et al., 2017). In addition, overexpression of *F3H* from *C. sinensis* (Mahajan
and Yadav, 2014) and from *Lycium chinense* (Song et al., 2016) in tobacco improved the
tolerance to salt stress and fungal pathogens in the first case and to drought stress in the latter
case. In conclusion, our results clearly show that *Musa*F3H1 and *Musa*F3H2 are functional
flavanone 3-hydroxylases and that *Musa*F3H2 might play a role in stress response.
441
_Musa_FLS1 and _Musa_FLS3
*Musa*FLS1 was found to be able to convert dihydrokaempferol to kaempferol in the *E. coli*
bioconversion assays. This ability was validated *in planta* by successful complementation of
the flavonol deficiency of *A. thaliana ans/fls1-2* double mutant seedlings, while the anthocyanin
deficiency phenotype was not restored. These observations could hint to an exclusive FLS
activity of *Musa*FLS1. FLS and DFR, the first enzymes of the flavonol and anthocyanin
branches of flavonoid biosynthesis, compete for dihydroflavonol substrates (Figure 1). This is
demonstrated by several *A. thaliana fls1* single mutants which accumulate higher levels of
anthocyanin pigments (Owens et al., 2008; Stracke et al., 2009). Falcone Ferreyra et al. (2010)
found that overexpression of *ZmFLS1* in an *A. thaliana fls1* mutant decreases the
accumulation of anthocyanins to wildtype level, again indicating that the anthocyanin and
flavonol branches of flavonoid biosynthesis can compete for dihydroflavonol substrates in the
same cell or tissue. Accordingly, the overlapping substrate usage of FLS and ANS is difficult
to analyse in a *fls1* single mutant. Here, the use of an *A. thaliana ans/fls1-2* double mutant is
a simple way to analyse the FLS and a possible ANS side activity of flavonol synthases and
vice versa *in planta*. To analyse further side activities, a *f3h fls ans* mutant would be beneficial.
A. thaliana ans/fls1-2 plants expressing *MusaFLS3* showed the accumulation of several
kaempferol derivatives. In contrast, plants complemented with *MusaFLS1* also accumulate
quercetin derivatives. This is most likely an effect of different environmental conditions in plant

growth. One set of plants was grown in the greenhouse (*Musa*FLS1), another set was grown in a growth chamber (*Musa*FLS3). In the plants used for *Musa*FLS3 experiments these conditions promoted the accumulation of dihydroquercetin, while the other growth conditions did not in the plants used for *Musa*FLS1 experiments. Accordingly, dihydroquercetin was probably not available as a substrate for *Musa*FLS1 and could not be converted to quercetin. A possible explanation for the varying dihydroquercetin amounts could be the light induced expression of F3'H, which converts dihydrokaempferol to kaempferol. The expression of *F3'H* in cultured *A. thaliana* cells can be induced by UV-light (Schoenbohm et al., 2000) and the activity of the *F3'H* promoter from *Vitis vinifera* is increased under light exposure (Sun et al., 2015). Consequently, a different light quality or higher light intensity could be responsible for increased *F3'H* expression, causing the accumulation of dihydroquercetin in the plants grown in the growth chamber. While *Musa*FLS3 was able to complement the flavonol deficiency in the *ans/fls1-2* mutant, it did not lead to an accumulation of anthocyanins. In contrast to the *in planta* results, *Musa*FLS3 did not convert dihydrokaempferol to kaempferol in the *in vivo* bioconversion assays, even though the protein was successfully expressed in the *E. coli* culture. We used the bioconversion assays as a simple and versatile tool to analyse enzymatic activities. Nevertheless, heterologous expression of eucaryotic proteins in *E. coli* is an artificial system. It can cause a divergent pattern of posttranslational modifications or production of high amounts of protein in inclusion bodies, leading to inactive protein (Sahdev et al., 2008). Preuss et al. (2009) reported that *At*FLS3 did not show FLS activity in *E. coli*, but did convert dihydroflavonols to the corresponding flavonols upon expression in yeast, indicating that the stability of *At*FLS3 was improved under the latter assay conditions. The approach carried out in this work might have similar limitations. Further limitations can be caused by different substrate preferences, as reported for *Os*FLS (Park et al., 2019). Nevertheless, the *in planta* complementation of the flavonol deficiency in the *A. thaliana ans/fls1-2* mutant and the lacking complementation of the anthocyanin deficiency phenotype in the *ans/fls1-2* mutant indicate that *Musa*FLS3 is a functional flavonol synthase, which does not exhibit significant ANS activity.

*Musa*FLS1 transcript abundance was found to be high in adult leaves and low in roots. *Musa*FLS3 and *Musa*FLS4 revealed opposed transcript levels. Such divergent expression patterns have previously been observed for *FLS1* and *FLS2* from *Freesia hybrida* and *FLS1* and *FLS2* from *Cyclamen purpurascens* (Akita et al., 2018; Shan et al., 2020). These opposed expression patterns could point to differential activity in distinct organs. Furthermore, the tandemly arranged *MusaFLS3* and *MusaFLS4* genes show very similar expression patterns (Table 1), possibly indicating functional redundancy. While we could not find expression of *Musa*FLS2 in our expression analyses, Pandey et al. (2016) observed *Musa*FLS2 expression in several organs (including bract, pseudostem and root), as well as in different developmental

stages of peel and pulp. Again, these results show that different cultivars, growth conditions,
sampling timepoints and analysed organs can have a strong influence on the resulting data.
Accordingly, data from different studies should be evaluated carefully. *MusaFLS1* and
MusaFLS2 expression in leaves of plantlets does not significantly increase after MJ treatment
(Pandey et al., 2016). However, UV radiation induces the expression of FLS from *Z. mays*
(Falcone Ferreyra et al., 2010) and *M. domestica* (Henry-Kirk et al., 2018) and the relative
expression of FLS from *Triticum aestivum* increases during drought stress (Ma et al., 2014).
Together with the knowledge that flavonols act as antioxidants (Wang et al., 2006), an
involvement of *Musa*FLSs in stress response seems feasible and should be further analysed
under a broader range of conditions. In summary, our results show that *Musa*FLS1 and
*Musa*FLS3 are functional flavonol synthase enzymes and hint at possible differential organ
specific activities of *Musa*FLS1 and *Musa*FLS3 / *Musa*FLS4.
510
_Musa_ANS
A. thaliana ans/fls1-2 seedlings expressing *Musa*ANS show a strong, red pigmentation,
revealing that *Musa*ANS can complement the anthocyanin deficiency caused by mutation of
AtANS. However, the seedlings did not display flavonol derivatives in HPTLC analyses, which
have been reported to detect flavonol glycosides at levels of 50 pMol (Stracke, 2010). These
results indicate that *Musa*ANS is a functional anthocyanidin synthase, but does not have FLS
activity. The RNA-Seq data-derived expression profiles revealed high *Musa*ANS transcript
abundance in pulp and peel. Additionally, *MusaANS* expression has been reported to be high
in bract tissue (Pandey et al., 2016), the specialised leaves surrounding the flowers and usually
coloured red or purple due to anthocyanin accumulation (Pazmiño-Durána et al., 2001). This
spatial correlation of *MusaANS* transcripts and anthocyanin metabolites in bract tissue
supports the proposed biological functionality of this enzyme, catalysing the conversion of
leucoanthocyanidins to anthocyanidins in *Musa*. *MusaANS* expression increases 24 hours
after MJ treatment and decreases following dark treatment (Pandey et al., 2016). As supposed
for *MusaF3H2*, the increased expression of *MusaANS* after MJ treatment could imply an
involvement in the formation of anthocyanins as a consequence of stress response. ABA,
salicylic acid (SA), UV-B and cold treatments have been shown to enhance *ANS* transcript
abundance in *G. biloba* (Xu et al., 2008) and overexpression of *OsANS* raises the antioxidant
potential in *O. sativa* (Reddy et al., 2007). The accumulation of anthocyanins has often been
shown to be induced by (UV-) light (Takahashi et al., 1991; Stapleton and Walbot, 1994; Meng
et al., 2004). We therefore assume that also *Musa*ANS could be involved in such stress
response.
Very recently, a R2R3-MYB-type transcription factor (MaMYB4) has been identified as a
negative regulator of anthocyanin biosynthesis in *Musa* acting as repressor on the *Musa*ANS

promoter (Deng et al., 2021). These results give a first insight into the transcriptional regulation

of *Musa*ANS expression and confirm the role of *Musa*ANS in the anthocyanin biosynthesis in

Musa. Taking all available evidence into account, *MusaANS* encodes a functional

anthocyanidin synthase with a possible involvement in the plants' stress response.

To deepen the knowledge about 2-ODDs and other enzymes involved in *Musa* flavonoid

biosynthesis, it would be beneficial to acquire even more spatially and timely highly resolved

transcriptome and in particular metabolite data. This data could serve as a starting point for

the analysis of organ-, stress- or development-specific enzyme activities and functions, as well

as possible substrate preferences. It could also be used to elucidate the regulatory network of

Musa flavonoid biosynthesis. Furthermore, knowledge about the influence of different stresses

(e.g. pathogens, light, temperature) on specific transcriptomes and metabolomes of the *Musa*

plant could help to widen the knowledge of flavonoid biosynthesis and particular the

functionality of the *Musa* 2-ODD enzymes. This could also lead to the detection of possible

restricted side activities or overlapping functionalities as described for 2-OODs in some other

plant species (Falcone Ferreyra et al., 2010; Park et al., 2019) and to further decode the cause

of these multifunctionalities in 2-ODDs.

To conclude, in this study the functionality of five 2-ODDs involved in flavonoid biosynthesis in

Musa was demonstrated *in planta*. Knowledge gained about the structural genes *MusaF3H1*,

MusaF3H2, *MusaFLS1*, *MusaFLS3* and *MusaANS* in a major crop plant provides a basis for

further research towards engineered, increased flavonoid production in banana, which could

contribute to the fruits' antioxidant activity and nutritional value, and possibly even enhance

the plants' defence against *Foc*-TR4.

Conflict of Interest

The authors declare that there is no conflict of interest.

Author Contributions

MB and RS planned the experiments. MB and CA performed the experiments and analysed

the data. RS and BW supervised the project. MB wrote the initial draft. RS and BW revised the

manuscript. All authors read and approved the final manuscript version.

Funding

This work was supported by basic funding of the chair of Genetics and Genomics of Plants

provided by Bielefeld University/Faculty of Biology.

Acknowledgements

We are grateful to Melanie Kuhlmann for her excellent assistance in the laboratory and to Andrea Voigt for her competent help in the greenhouse. We thank Anika Beckers who contributed to creation of constructs with *MusaFLS* cDNAs and Prisca Viehöver for sequencing. In addition, we thank Thomas Baier for his support with the SDS-PAGE. A special "thanks" to Stefan Martens for discussion and helpful hints to the bioconversion assay. We acknowledge support for the publication costs by the Open Access Publication Fund of Bielefeld University.

References

Akita, Y., Kitamura, S., Mikami, R., and Ishizaka, H. (2018). Identification of functional flavonol synthase genes from fragrant wild cyclamen (Cyclamen purpurascens). *Journal of Plant Biochemistry and Biotechnology* 27(2)**,** 147-155. doi: https://doi.org/10.1007/s13562-017-0423-9.

Andersen, Ø.M., and Jordheim, M. (2010). "Anthocyanins," in *Encyclopedia of Life Sciences*. (Chichester: John Wiley &Sons, Ltd).

Appelhagen, I., Thiedig, K., Nordholt, N., Schmidt, N., Huep, G., Sagasser, M., et al. (2014). Update on *transparent testa* mutants from *Arabidopsis thaliana*: characterisation of new alleles from an isogenic collection. *Planta* 240(5)**,** 955-970.

Arias, P., Dankers, C., Liu, P., and Pilkauskas, P. (2003). The World Banana Economy, 1985-2002. http://www.fao.org/3/y5102e/y5102e5100.htm.

Asif, M.H., Dhawan, P., and Nath, P. (2000). A simple procedure for the isolation of high quality RNA from ripening banana fruit. *Plant Molecular Biology Reporter* 18**,** 109-115.

Bednar, R.A., and Hadcock, J.R. (1988). Purification and Characterizationo of Chalcone Isomerase from Soybeans. *Journal of Biological Chemistry* 263**,** 9582-9588

Britsch, L. (1990). Purification and Characterization of Flavone Synthase I, a 2-Oxoglutarate-Dependent Desaturase. *Archives of Biochemistry and Biophysics* 282(1)**,** 152-160.

Brown, J.W., Walker, J.F., and Smith, S.A. (2017). Phyx: phylogenetic tools for unix. *Bioinformatics* 33(12)**,** 1886-1888. doi: 10.1093/bioinformatics/btx063.

Cheng, A.X., Han, X.J., Wu, Y.F., and Lou, H.X. (2014). The function and catalysis of 2-oxoglutarate-dependent oxygenases involved in plant flavonoid biosynthesis. *International Journal of Molecular Sciences* 15(1)**,** 1080-1095.

Cheong, J.-J., and Choi, Y.D. (2003). Methyl jasmonate as a vital substance in plants. *Trends in Genetics* 19(7)**,** 409-413. doi: 10.1016/S0168-9525(03)00138-0.

Clough, S.J., and Bent, A.F. (1998). Floral dip: a simplified method for Agrobacterium-mediated transformation of *Arabidopsis thaliana*. *The Plant Journal* 16(6)**,** 735-743. doi: 10.1046/j.1365-313x.1998.00343.x.

Dauer, A., Hensel, A., Lhoste, E., Knasmüller, S., and Mersch-Sundermann, V. (2003). Genotoxic and antigenotoxic effects of catechin and tannins from the bark of Hamamelis virginiana L. in metabolically competent, human hepatoma cells (Hep G2) using single cell gel electrophoresis. *Phytochemistry* 63(2)**,** 199-207. doi: 10.1016/S0031-9422(03)00104-3.

Deng, G.M., Zhang, S., Yang, Q.S., Gao, H.J., Sheng, O., Bi, F.C., et al. (2021). MaMYB4, an R2R3-MYB Repressor Transcription Factor, Negatively Regulates the Biosynthesis of Anthocyanin in Banana. *Frontiers in Plant Science* 11**,** 600704. doi: 10.3389/fpls.2020.600704.

DHont, A., Denoeud, F., Aury, J.M., Baurens, F.C., Carreel, F., Garsmeur, O., et al. (2012). The banana (Musa acuminata) genome and the evolution of monocotyledonous plants. *Nature* 488(7410)**,** 213-217.

Falcone Ferreyra, M.L., Rius, S., Emiliani, J., Pourcel, L., Feller, A., Morohashi, K., et al. (2010). Cloning and characterization of a UV-B-inducible maize flavonol synthase. *The Plant Journal* 62(1)**,** 77-91. doi: 10.1111/j.1365-313X.2010.04133.x.

Falcone Ferreyra, M.L., Rius, S.P., and Casati, P. (2012). Flavonoids: biosynthesis, biological functions, and biotechnological applications. *Frontiers in Plant Science* 3**,** 222. doi: 10.3389/fpls.2012.00222.

FAO (2019). "The State of Food and Agriculture 2019". (Rome, Italy: Food and Agricultural Organization (FAO)).

Fliegmann, J., Furtwangler, K., Malterer, G., Cantarello, C., Schuler, G., Ebel, J., et al. (2010). Flavone synthase II (CYP93B16) from soybean (Glycine max L.). *Phytochemistry* 71(5-6)**,** 508-514. doi: 10.1016/j.phytochem.2010.01.007.

Forkmann, G., de Vlaming, P., Spribille, R., Wiering, H., and Schram, W. (1986). Genetic and Biochemical Studies on the Conversion of Dihydroflavonols to Flavonols in Flowers of *Petunia hybrida*. *Zeitschrift für Naturforschung C* 41(1-2)**,** 179-186.

Forkmann, G., Heller, W., and Grisebach, H. (1980). Anthocyanin Biosynthesis in Flowers of *Matthiola incana* Flavanone 3- and Flavonoid 3'-Hydroxylases. *Zeitschrift für Naturforschung C* 35(9-10)**,** 691-695. doi: doi.org/10.1515/znc-1980-9-1004.

Forster, M., Rodríguez, E.R., Martín, J.D., and Romero, C.D. (2003). Distribution of Nutrients in Edible Banana Pulp. *Food Technology and Biotechnology* 41(2)**,** 167-171.

Fu, X., Cheng, S., Liao, Y., Huang, B., Du, B., Zeng, W., et al. (2018). Comparative analysis of pigments in red and yellow banana fruit. *Food Chemistry* 239**,** 1009-1018. doi: 10.1016/j.foodchem.2017.07.046.

Gebhardt, Y.H., Witte, S., Steuber, H., Matern, U., and Martens, S. (2007). Evolution of flavone synthase I from parsley flavanone 3beta-hydroxylase by site-directed mutagenesis. *Plant Physiology* 144(3)**,** 1442-1454. doi: doi: 10.1104/pp.18.00463.

Goufo, P., and Trindade, H. (2014). Rice antioxidants: phenolic acids, flavonoids, anthocyanins, proanthocyanidins, tocopherols, tocotrienols, gamma-oryzanol, and phytic acid. *Food Science & Nutrition* 2(2)**,** 75-104. doi: 10.1002/fsn3.86.

Gronquist, M., Bezzerides, A., Attygalle, A., Meinwald, J., Eisner, M., and Eisner, T. (2001). Attractive and defensive functions of the ultraviolet pigments of a flower *(Hypericum calycinum)*. *Proceedings of the National Academie of Sciences of the United States of America* 98(24)**,** 13745-13750.

Grotewold, E. (2006). *The Science of Flavonoids.* Columbus: The Ohio State University.

Hahlbrock, K., and Scheel, D. (1989). Physiology and molecular biology of phenylpropanoid metabolism. *Annual Review of Plant Physiology and Plant Molecular Biology* 40(1)**,** 347-369. doi: 10.1146/annurev.pp.40.060189.002023.

Han, Y., Huang, K., Liu, Y., Jiao, T., Ma, G., Qian, Y., et al. (2017). Functional Analysis of Two Flavanone-3-Hydroxylase Genes from *Camellia sinensis*: A Critical Role in Flavonoid Accumulation. *Genes* 8(11). doi: 10.3390/genes8110300.

Heller, W., Forkmann, G., Britsch, L., and Grisebach, H. (1985). Enzymatic reduction of (+)-dihydroflavonols to flavan-3,4-cis-diols with flower extracts from *Matthiola incana* and its role in anthocyanin biosynthesis. *Planta* 165**,** 284-287.

Henry-Kirk, R.A., Plunkett, B., Hall, M., McGhie, T., Allan, A.C., Wargent, J.J., et al. (2018). Solar UV light regulates flavonoid metabolism in apple (Malus x domestica). *Plant, Cell & Environment* 41(3)**,** 675-688. doi: 10.1111/pce.13125.

Holton, T.A., Brugliera, F., and Tanaka, Y. (1993). Cloning and expression of flavonol synthase from *Petunia hybrida*. *The Plant Journal* 4**,** 1003-1010. doi: 10.1046/j.1365-313x.1993.04061003.x.

Ishikura, N., and Yoshitama, K. (1984). Anthocyanin-Flavonol Co-pigmentation in Blue Seed Coats of Ophiopogon jaburan. *Journal of Plant Physiology* 115(2)**,** 171-175. doi: 10.1016/s0176-1617(84)80064-4.

Jaakkola, P., Mole, D.R., Tian, Y.M., Wilson, M.I., Gielbert, J., Gaskell, S.J., et al. (2001). Targeting of HIF-alpha to the von Hippel-Lindau ubiquitylation complex by O2-

regulated prolyl hydroxylation. *Science* 292(5516)**,** 468-472. doi: 10.1126/science.1059796.

Jakoby, M., Wang, H.Y., Reidt, W., Weisshaar, B., and Bauer, P. (2004). FRU (BHLH029) is required for induction of iron mobilization genes in Arabidopsis thaliana. *FEBS Letters* 577(3)**,** 528-534.

Jiang, N., Doseff, A.I., and Grotewold, E. (2016). Flavones: From Biosynthesis to Health Benefits. *Plants* 5(2). doi: 10.3390/plants5020027.

Kandil, F.E., Smith, M.A., Rogers, R.B., Pepin, M.F., Song, L.L., Pezzuto, J.M., et al. (2002). Composition of a chemopreventive proanthocyanidin-rich fraction from cranberry fruits responsible for the inhibition of 12-O-tetradecanoyl phorbol-13-acetate (TPA)-induced ornithine decarboxylase (ODC) activity. *Journal of Agricultural and Food Chemistry* 50(5)**,** 1063-1069. doi: 10.1021/jf011136z.

Katoh, K., and Standley, D.M. (2013). MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* 30(4)**,** 772-780.

Kawai, Y., Ono, E., and Mizutani, M. (2014). Evolution and diversity of the 2-oxoglutarate-dependent dioxygenase superfamily in plants. *The Plant Journal* 78(2)**,** 328-343. doi: 10.1111/tpj.12479.

Koncz, C., and Schell, J. (1986). The promoter of TL-DNA gene 5 controls the tissue-specific expression of chimaeric genes carried by a novel type of Agrobacterium binary vector. *Molecular Genetics and Genomics* 204**,** 383-396.

Kreuzaler, F., and Hahlbrock, K. (1972). Enzymatic synthesis of aromatic compounds in higher plants: formation of naringenin (5,7,4'-trihydroxyflavanone) from p-coumaroyl coenzyme A and malonyl coenzyme A *FEBS Letters* 28(1)**,** 69-72.

Le Roy, J., Huss, B., Creach, A., Hawkins, S., and Neutelings, G. (2016). Glycosylation Is a Major Regulator of Phenylpropanoid Availability and Biological Activity in Plants. *Frontiers in Plant Science* 7**,** 735.

Lepiniec, L., Debeaujon, I., Routaboul, J.M., Baudry, A., Pourcel, L., Nesi, N., et al. (2006). Genetics and biochemistry of seed flavonoids. *Annual Review of Plant Biology* 57**,** 405-430.

Letunic, I., and Bork, P. (2019). Interactive Tree Of Life (iTOL) v4: recent updates and new developments. *Nucleic Acids Research* 47(W1)**,** W256-W259. doi: 10.1093/nar/gkz239.

Li, J., Ou-Lee, T.M., Raba, R., Amundson, R.G., and Last, R.L. (1993). Arabidopsis Flavonoid Mutants Are Hypersensitive to UV-B Irradiation. *The Plant Cell* 5(2)**,** 171-179. doi: 10.1105/tpc.5.2.171.

Liu, M., Li, X., Liu, Y., and Cao, B. (2013). Regulation of flavanone 3-hydroxylase gene involved in the flavonoid biosynthesis pathway in response to UV-B radiation and drought stress in the desert plant, Reaumuria soongorica. *Plant Physiology and Biochemistry* 73**,** 161-167. doi: 10.1016/j.plaphy.2013.09.016.

Lu, P., Yu, S., Zhu, N., Chen, Y.R., Zhou, B., Pan, Y., et al. (2018). Genome encode analyses reveal the basis of convergent evolution of fleshy fruit ripening. *Nature Plants* 4(10)**,** 784-791. doi: https://doi.org/10.1038/s41477-018-0249-z.

Luo, P., Ning, G., Wang, Z., Shen, Y., Jin, H., Li, P., et al. (2016). Disequilibrium of Flavonol Synthase and Dihydroflavonol-4-Reductase Expression Associated Tightly to White vs. Red Color Flower Formation in Plants. *Frontiers in Plant Science* 6**,** 1257. doi: 10.3389/fpls.2015.01257.

Ma, D.Y., Sun, D.X., Wang, C.Y., Li, Y.G., and Guo, T.C. (2014). Expression of flavonoid biosynthesis genes and accumulation of flavonoid in wheat leaves in response to drought stress. *Plant Physiology and Biochemistry* 80**,** 60-66. doi: http://dx.doi.org/10.1016/j.plaphy.2014.03.024.

Mahajan, M., and Yadav, S.K. (2014). Overexpression of a tea flavanone 3-hydroxylase gene confers tolerance to salt stress and Alternaria solani in transgenic tobacco. *Plant Molecular Biology* 85(6)**,** 551-573. doi: 10.1007/s11103-014-0203-z.

Martens, S., and Forkmann, G. (1999). Cloning and expression of flavone synthase II from Gerbera hybrids. *The Plant Journal* 20(5)**,** 611-618.

Martin, G., Baurens, F.C., Droc, G., Rouard, M., Cenci, A., Kilian, A., et al. (2016). Improvement of the banana "Musa acuminata" reference sequence using NGS data and semi-automated bioinformatics methods. *BMC Genetics* 17(243).

Mehrtens, F., Kranz, H., Bednarek, P., and Weisshaar, B. (2005). The Arabidopsis transcription factor MYB12 is a flavonol-specific regulator of phenylpropanoid biosynthesis. *Plant Physiology* 138(2)**,** 1083-1096.

Meng, X., Xing, T., and Wang, X. (2004). The role of light in the regulation of anthocyanin accumulation in Gerbera hybrida. *Plant Growth Regulation* 44(3)**,** 243-250. doi: 10.1007/s10725-004-4454-6.

Mo, Y., Nagel, C., and Taylor, L.P. (1992). Biochemical complementation of chalcone synthase mutants defines a role for flavonols in functional pollen. *Proceedings of the National Academy of Sciences of the United States of America* 89**,** 7213-7217.

Owens, D.K., Alerding, A.B., Crosby, K.C., Bandara, A.B., Westwood, J.H., and Winkel, B.S. (2008). Functional analysis of a predicted flavonol synthase gene family in Arabidopsis. *Plant Physiology* 147(3)**,** 1046-1061. doi: 10.1104/pp.108.117457.

Pandey, A., Alok, A., Lakhwani, D., Singh, J., Asif
, M.H., and Trivedi, P.K. (2016). Genome-wide Expression Analysis and Metabolite Profiling Elucidate Transcriptional Regulation of Flavonoid Biosynthesis and Modulation under Abiotic Stresses in Banana. *Scientific Reports* 6.

Park, S., Kim, D.H., Park, B.R., Lee, J.Y., and Lim, S.H. (2019). Molecular and Functional Characterization of Oryza sativa Flavonol Synthase (OsFLS), a Bifunctional Dioxygenase. *Journal of Agricultural and Food Chemistry* 67(26)**,** 7399-7409. doi: 10.1021/acs.jafc.9b02142.

Pazmiño-Durána, E.A., Giustib, M.M., Wrolstad, R.E., and Glóriaa, M.B.A. (2001). Anthocyanins from banana bracts (Musa X paradisiaca) as potential food colorants. *Food Chemistry* 73**,** 327-332. doi: doi.org/10.1016/S0308-8146(00)00305-8.

Perez-Vizcaino, F., and Duarte, J. (2010). Flavonols and cardiovascular disease. *Molecular Aspects of Medicine* 31(6)**,** 478-494. doi: 10.1016/j.mam.2010.09.002.

Prescott, A.G., and John, P. (1996). DIOXYGENASES: Molecular Structure and Role in Plant Metabolism. *Annual Review of Plant Physiology and Plant Molecular Biology* 47**,** 245-271. doi: doi.org/10.1146/annurev.arplant.47.1.245.

Preuss, A., Stracke, R., Weisshaar, B., Hillebrecht, A., Matern, U., and Martens, S. (2009). *Arabidopsis thaliana* expresses a second functional flavonol synthase. *FEBS Letters* 583(12)**,** 1981–1986. doi: 10.1016/j.febslet.2009.05.006.

Price, M.N., Dehal, P.S., and Arkin, A.P. (2010). FastTree 2--approximately maximum-likelihood trees for large alignments. *PLoS One* 5(3)**,** e9490. doi: 10.1371/journal.pone.0009490.

Pucker, B., Pandey, A., Weisshaar, B., and Stracke, R. (2020a). The R2R3-MYB gene family in banana (Musa acuminata): Genome-wide identification, classification and expression patterns. *PLoS One* 15(10)**,** e0239275. doi: 10.1371/journal.pone.0239275.

Pucker, B., Reiher, F., and Schilbert, H.M. (2020b). Automatic Identification of Players in the Flavonoid Biosynthesis with Application on the Biomedicinal Plant Croton tiglium. *Plants* 9(9). doi: 10.3390/plants9091103.

Reddy, A.M., Reddy, V.S., Scheffler, B.E., Wienand, U., and Reddy, A.R. (2007). Novel transgenic rice overexpressing anthocyanidin synthase accumulates a mixture of flavonoids leading to an increased antioxidant potential. *Metabolic Engineering* 9(1)**,** 95-111. doi: https://doi.org/10.1016/j.ymben.2006.09.003.

Sahdev, S., Khattar, S.K., and Saini, K.S. (2008). Production of active eukaryotic proteins through bacterial expression systems: a review of the existing biotechnology strategies. *Molecular and Cellular Biochemistry* 307(1-2)**,** 249-264. doi: 10.1007/s11010-007-9603-6.

Saito, K., Kobayashi, M., Gong, Z.Z., Tanaka, Y., and Yamazaki, M. (1999). Direct evidence for anthocyanidin synthase as a 2-oxoglutarate-dependent oxygenase: molecular cloning and functional expression of cDNA from a red forma of Perilla frutescens. *The Plant Journal* 17(2)**,** 181-189.

Sanger, F., Nicklen, S., and Coulson, A.R. (1977). DNA sequencing with chain-terminating
inhibitors. *Proceedings of the National Academy of Sciences of the United States of*
America 74(12), 5463-5467.
Schoenbohm, C., Martens, S., Eder, C., Forkmann, G., and Weisshaar, B. (2000).
Identification of the *Arabidopsis thaliana* flavonoid 3'-hydroxylase gene and functional
expression of the encoded P450 enzyme. *Biological Chemistry* 381, 749-753.
Shan, X., Li, Y., Yang, S., Yang, Z., Qiu, M., Gao, R., et al. (2020). The spatio-temporal
biosynthesis of floral flavonols is controlled by differential phylogenetic MYB regulators
in Freesia hybrida. *New Phytologist*. doi: 10.1111/nph.16818.
Sheahan, J.J., and Rechnitz, G.A. (1992). Flavonoid-specific staining of Arabidopsis thaliana.
BioTechniques 13(6), 880-883.
Singh, B., Singh, J.P., Kaur, A., and Singh, N. (2016). Bioactive compounds in banana and
their associated health benefits - A review. *Food Chemistry* 206, 1-11. doi:
10.1016/j.foodchem.2016.03.033.
Song, X., Diao, J., Ji, J., Wang, G., Guan, C., Jin, C., et al. (2016). Molecular cloning and
identification of a flavanone 3-hydroxylase gene from *Lycium chinense*, and its
overexpression enhances drought stress in tobacco. *Plant Physiol Biochem* 98, 89-
100. doi: 10.1016/j.plaphy.2015.11.011.
Stapleton, A.E., and Walbot, V. (1994). Flavonoids can protect maize DNA from the induction
of ultraviolet radiation damage. *Plant Physiology* 105, 881-889.
Stracke, R., De Vos, R.C.H., Bartelniewoehner, L., Ishihara, H., Sagasser, M., Martens, S., et
al. (2009). Metabolomic and genetic analyses of flavonol synthesis in *Arabidopsis*
thaliana support the *in vivo* involvement of leucoanthocyanidin dioxygenase. *Planta*
229(2), 427-445. doi: 10.1007/s00425-008-0841-y.
Stracke, R., Ishihara, H., Huep, G., Barsch, A., Mehrtens, F., Niehaus, K., et al. (2007).
Differential regulation of closely related R2R3-MYB transcription factors controls
flavonol accumulation in different parts of the *Arabidopsis thaliana* seedling. *The Plant*
Journal 50(4), 660-677. doi: 10.1111/j.1365-313X.2007.03078.x.
Stracke, R., Turgut-Kara, N., and Weisshaar, B. (2017). The AtMYB12 activation domain maps
to a short C-terminal region of the transcription factor. *Zeitschrift für Naturforschung C*
72(7-8), 251-257. doi: 10.1515/znc-2016-0221.
Sun, J., Zhang, J., Fang, H., Peng, L., Wei, S., Li, C., et al. (2019). Comparative transcriptome
analysis reveals resistance-related genes and pathways in *Musa acuminata* banana
'Guijiao 9' in response to Fusarium wilt. *Plant Physiology and Biochemistry* 141, 83-94.
doi: 10.1016/j.plaphy.2019.05.022.
Sun, R.Z., Pan, Q.H., Duan, C.Q., and Wang, J. (2015). Light response and potential
interacting proteins of a grape flavonoid 3 '-hydroxylase gene promoter. *Plant*
Physiology and Biochemistry 97, 70-81. doi:
http://dx.doi.org/10.1016/j.plaphy.2015.09.016.
Takahashi, A., Takeda, K., and Ohnishi, T. (1991). Light-Induced Anthocyanin Reduces the
Extent of Damage to DNA in UV-Irradiated Centaurea cyanus Cells in Culture. *Plant*
Cell Physiology 32(4), 541-547. doi: 10.1093/oxfordjournals.pcp.a078113.
Tanaka, Y., Sasaki, N., and Ohmiya, A. (2008). Biosynthesis of plant pigments: anthocyanins,
betalains and carotenoids. *The Plant Journal* 54(4), 733-749. doi:
https://doi.org/10.1111/j.1365-313X.2008.03447.x.
Tohge, T., de Souza, L.P., and Fernie, A.R. (2017). Current understanding of the pathways of
flavonoid biosynthesis in model and crop plants. *Journal of Experimental Botany*
68(15), 4013-4028.
Trewick, S.C., Henshaw, T.F., Hausinger, R.P., Lindahl, T., and Sedgwick, B. (2002). Oxidative
demethylation by Escherichia coli AlkB directly reverts DNA base damage. *Nature*
419(6903), 174-178. doi: https://doi.org/10.1038/nature00908.
Wall, M.M. (2006). Ascorbic acid, vitamin A, and mineral composition of banana (Musa sp.)
and papaya (Carica papaya) cultivars grown in Hawaii. *Journal of Food Composition*
and Analysis 19(5), 434-445. doi: 10.1016/j.jfca.2006.01.002.

Wang, L., Tu, Y.-C., Lian, T.-W., Hung, J.-T., Yen, J.-H., and Wu, M.-J. (2006). Distinctive
antioxidant and antiinflammatory effects of flavonols. *Journal of Agricultural and Food*
Chemistry 54(26)**,** 9798-9804. doi: doi.org/10.1021/jf0620719.
Wen, W., Alseekh, S., and Fernie, A.R. (2020). Conservation and diversification of flavonoid
metabolism in the plant kingdom. *Current Opinion in Plant Biology* 55**,** 100-108. doi:
10.1016/j.pbi.2020.04.004.
Winkel, B.S.J. (2006). "The biosynthesis of flavonoids," in *The Science of Flavonoids,* ed. E.
Grotewold. (New York, USA: Springer Scienceand Business Media, Inc), 71-96.
Xu, F., Cheng, H., Cai, R., Li, L.L., Chang, J., Zhu, J., et al. (2008). Molecular cloning and
function analysis of an anthocyanidin synthase gene from Ginkgo biloba, and its
expression in abiotic stress responses *Molecules and Cells* 26(6)**,** 536-547.
Xu, F., Li, L., Zhang, W., Cheng, H., Sun, N., Cheng, S., et al. (2012). Isolation,
characterization, and function analysis of a flavonol synthase gene from Ginkgo biloba.
Molecular Biology Reports 39(3)**,** 2285-2296. doi: 10.1007/s11033-011-0978-9.
Yan, Y., Chemler, J., Huang, L., Martens, S., and Koffas, M.A. (2005). Metabolic engineering
of anthocyanin biosynthesis in Escherichia coli. *Applied and Environmental*
Microbiology 71(7)**,** 3617-3623.