1   Neuronal genetic rescue normalizes brain network dynamics in a lysosomal storage disorder
2   despite persistent storage accumulation
3
4

5   **Authors**
6   Rebecca C. Ahrens-Nicklas*[1,2], Luis Tecedor[3], Arron F. Hall[2], Owen Kane[2], Richard J. Chung[2], Elena
7   Lysenko[3], Eric D. Marsh[4,5], Colleen S. Stein[6], Beverly L. Davidson*[3,7]
8
9
Affiliations
[1]Department of Pediatrics, Perelman School of Medicine at the University of Pennsylvania.
[2]Division of Human Genetics, The Children's Hospital of Philadelphia.
[3]The Raymond G. Perelman Center for Cellular and Molecular Therapeutics, The Children's Hospital
of Philadelphia.
[4]Department of Neurology, Perelman School of Medicine at the University of Pennsylvania.
[5]Division of Child Neurology, The Children's Hospital of Philadelphia.
[6]Department of Internal Medicine, Fraternal Order of Eagles Diabetes Research Center, Abboud
Cardiovascular Research Center, Carver College of Medicine, University of Iowa.
[7]The Department of Pathology and Laboratory Medicine Perelman School of Medicine at the
University of Pennsylvania.
21
22
23 *Correspondence should be addressed to co-corresponding authors
24
Rebecca C. Ahrens-Nicklas, MD, PhD
Children's Hospital of Philadelphia
3615 Civic Center Blvd Abramson 1016H
Philadelphia, PA 19104,
Tel: 215-590-3412
Email: ahrensnicklasr@email.chop.edu
31
32 AND
33
Beverly L. Davidson, PhD
Children's Hospital of Philadelphia,
3501 Civic Center Boulevard, 5060 CTRB,
Philadelphia, PA 19104,
Tel: 267-426-0929,
Email: davidsonbl@chop.edu

40
41 **Short title:** Genetic rescue of network activity in CLN3 Disease
42
43
44
45

Abstract

Although neurologic symptoms occur in two-thirds of lysosomal storage disorders (LSDs), for most we

do not understand the mechanisms underlying brain dysfunction. A major unanswered question is if the

pathogenic hallmark of LSDs, storage accumulation, induces functional defects directly or is a disease

bystander. Also, for most LSDs we do not know the impact of loss-of-function in individual cell types.

Understanding these critical questions are essential to therapy development. Here, we determined the

impact of genetic rescue in distinct cell types on neural circuit dysfunction in CLN3 disease, the most

common pediatric dementia and a representative LSD. We restored *Cln3* expression via AAV-mediated

gene delivery and conditional genetic rescue in a CLN3 mouse model. Surprisingly, we found that low-

level rescue of *Cln3* expression in neurons alone normalized clinically-relevant electrophysiologic

markers of network dysfunction, despite the presence of substantial residual histopathology, in contrast to

restoring expression in astrocytes. Thus, loss of Cln3 function in neurons, not storage accumulation,

underlies neurologic dysfunction in CLN3 disease, implying that storage clearance may be an

inappropriate target for therapy development and an ineffectual biomarker.

Introduction

Lysosomal storage disorders (LSDs) are a group of approximately 50 rare metabolic diseases

characterized by progressive accumulation of storage material in lysosomes. Neurological symptoms,

including neurocognitive regression, seizures, and psychiatric symptoms, occur in more than 2/3 of LSDs

and are generally not ameliorated by current therapies [1]. For many LSDs, including CLN3 disease, the

most common cause of childhood dementia, the mechanisms underlying neurologic dysfunction are

poorly understood.

The function of CLN3 protein remains unknown, hindering mechanism-based therapy development. The

brains of CLN3 disease mouse models demonstrate progressive storage accumulation, predominantly

comprised of subunit C of the mitochondrial ATP synthase (SCMAS) [2] [3], mirroring what occurs in CLN3

disease patients[4]. Therefore, most drug development efforts rely on histopathologic markers of efficacy,

including reduction of storage material, reactive astrocytosis and microgliosis[2,4,5]. However, whether these

histopathologic changes cause disease, augment the disease process, or are simply harmless

epiphenomenon is unclear.

In addition to lack of clarity on the impact of storage, the role of various CNS cell types in driving

functional deficits in CLN3 disease is unknown. Astrocytes and microglia isolated from CLN3 disease

mouse models demonstrate altered cellular properties and physiology[6-8], and in animal models, localized

glial activation occurs in regions where neurons will later die[9] supporting a role for glia as a disease

driver. On the other hand, neurons from CLN3 animal models exhibit both histopathologic and functional

differences[2,10,11]. Ultimately, patients' symptoms such as seizures and neurologic regression arise directly

from functional changes in key neuronal circuits. Nonetheless, we do not understand what cell types

underlie these pathologic network dynamics in the brain.

Previously, we described network-level neurologic defects in both the cortex and hippocampus of two

CLN3 disease mouse models using *in vitro* voltage sensitive dye imaging and *in vivo* electrophysiology

techniques[12]. While many brain regions are affected in CLN3 disease, the cortex and hippocampus are

especially vulnerable and show early pathologic changes in mouse models and human patients[3,4,11,13].

We found that network-level changes begin early in the disease course before substantial storage

accumulation in both the hippocampus and cortex. In slice-recordings, these changes include

hypoexcitability of the dentate gyrus of the hippocampus, a region essential for learning and memory. On

in vivo electroencephalogram (EEG), CLN3 disease mice show frequent epileptiform spikes, a feature of

many epilepsy models, and a shift towards high frequency background activity, a change associated with

cognitive impairment and present in Alzheimer's disease models and patients[14,15].

To date, CLN3 disease research has been hindered by the fact that animal models have only subtle

behavioral and survival phenotypes[16,17]. Most manifest histopathologic changes including storage

accumulation and reactive astrocytosis; however, these phenotypes may not correlate with central nervous

system (CNS) dysfunction. Our network-level electrophysiologic approach provides a robust, clinically

meaningful measure of neurologic function that can probe disease mechanisms and evaluate potential

therapies.

Here, we developed a new 'conditional rescue' mouse model of CLN3 disease. The model is homozygous

for the most common human CLN3 disease mutation[2], but also carries a Cre-inducible wild-type *Cln3*

transgene, under the control of its endogenous promoter, allowing for cell-type specific rescue of

expression. We employed AAV-mediated gene replacement as a second strategy to restore *Cln3*

expression and combined these tools with network-level electrophysiology techniques to answer three

essential questions: 1) Does gene replacement correct neural circuit dysfunction in a LSD? 2) What cell

types must be corrected to improve network dynamics? and 3) Does storage accumulation underlie

deficits in CNS network dynamics in CLN3 disease?

Results

In prior work, we found that CLN3 disease mouse models demonstrate robust, progressive neuronal

network defects[12]. We first assessed if neonatal correction of most CNS cell types could prevent circuit-

level functional pathology. For this, we employed an adeno-associated virus serotype 9 (AAV9) vector

expressing wild-type murine *Cln3* under control of 1665 bp of the endogenous *Cln3* promoter, which

mimics the low levels of expression seen in the endogenous setting **(Sup Figure 1-2, Sup Methods).**

Vector was delivered at p0 via unilateral intracerebroventricular (ICV) injection into CLN3-deficient

mice to transduce both astrocytes and neurons[18] **(Sup Figure 3)**. The CLN3 disease mouse model harbors

the most common human CLN3 disease mutation, a deletion of exons 7 and 8 ($Cln3^{\Delta ex78/\Delta ex78}$) (*2*). ICV

injection of AAV9.CLN3 at a dose of $1e^{10}$ vg increased *Cln3* expression **(Figure 1A)**. One year after

injection, levels in the $Cln3^{\Delta ex78/\Delta ex78}$ hemisphere contralateral to the injection site reached 24.5% ± 4.7 of

heterozygous controls. When assessed histologically, gene replacement partially reduced storage

accumulation in both the hippocampus and motor cortex ipsilateral to the injection site **(Figure 1B, Sup**

Figure 4).

To determine the impact of p0 AAV9.CLN3 infusion on *in vivo* network dynamics, we recorded activity

throughout the brain via *in vivo* EEG. Twelve-month-old $Cln3^{\Delta ex78/\Delta ex78}$ mice have increased spike burden

and a shift in background activity towards faster frequencies, with increased power in faster frequency

bands, as compared to $Cln3^{\Delta ex78/WT}$ controls **(Figure 1C-D)**. Specifically, there is decreased power in the

slow delta frequency band (0.1-4Hz) and a trend towards increased power in the faster beta (13-25Hz)

and gamma (25-50Hz) bands in $Cln3^{\Delta ex78/\Delta ex78}$ mice, consistent with our previous report in *Cln3* knock-out

mice[12]. Remarkably, restoring gene expression at p0 normalized spiking rates **(Figure 1C)** and corrected

background activity **(Figure 1D)** in 12-month-old $Cln3^{\Delta ex78/\Delta ex78}$ AAV9.CLN3-treated animals. Correction

of EEG phenotypes was equally effective in both the hemispheres. Thus, early gene replacement can

normalize network dynamics, and this rescue does not require full inhibition of storage accumulation.

Because AAV9 delivered to neonates transduces astrocytes and neurons[18], it is unclear whether all cell

types must be corrected to improve circuit function, a question critically important to translate gene

therapies to children. AAV9 delivered into the CNS generally transduces neurons in adult mice and

juvenile and adult large animal models[19-21]. To address this, we developed a conditional rescue model of

CLN3 disease, the Flex*Cln3* mouse. This model is homozygous for the common *Cln3* exon 7-8 deletion

and also carries the Cre-inducible Flex*Cln3* allele, permitting expression of wild-type Cln3 protein in the

presence of Cre recombinase **(Figure 2, Sup Methods and Sup Figures 5-7)**. The Flex*Cln3* allele is

under control of the same 1665 bp *Cln3* promoter as used for AAV-mediated gene replacement.

We validated the model by crossing Flex*Cln3* mice with *Zp3-Cre* mice[22] to drive germline level

recombination and flip the reverse-oriented wildtype Flex*Cln3* allele into the forward orientation in all

cells. Recombination of the allele was confirmed using orientation-specific PCR primers on genomic

DNA extracted from the hippocampus **(Figure 2B, Sup Figure 7)**. In these mice *Cln3* expression was

ubiquitously restored, with expression levels dictated by the transgenic *Cln3* 1665 bp promoter. *Cln3*

expression in the hippocampus was restored to 40% of heterozygous ($Cln3^{\Delta ex78/WT}$) control levels as

determined by qPCR **(Figure 2C)**. Correction (i.e. expression of the Flex*Cln3* transgene) in all cell types

(Flex*Cln3*/ $Cln3^{\Delta ex78/\Delta ex78}$/ *Zp3*-Cre) substantially reduced SCMAS storage in the hippocampus and other

brain regions **(Figure 2D, Sup Figure 8)**.

For selective rescue of astrocytes or neurons, Flex*Cln3* mice were crossed to *Gfap*-Cre[23] or *Syn1*-Cre[24]

driver lines, respectively. Cre-mediated rearrangement of the Flex*Cln3* allele was confirmed by PCR of

genomic DNA extracted from the hippocampus using orientation specific primers **(Figure 3A)**. As Cre is

only expressed in a subset of cells (either astrocytes or neurons), unlike in the case of *Zp3*-Cre **(Figure**

2B), both the forward and reverse alleles were detected in hippocampal tissue.

Expression of forward-oriented Flex*Cln3* transcripts was confirmed in the hippocampus of the astrocyte

and neuronal-specific rescue models. As rescue did not occur in all cell types, and Cln3 protomer activity

and expression can vary between cells and be induced by disease pathology[3] it is difficult to normalize

expression levels to heterozygote controls. However, to get a relative sense of expression levels we

compared the level of Flex*Cln3* transgene expression in the astrocyte-only and neuron-only rescue

models. We found no differences in transcript levels (normalized ΔCT values of 14.1 vs. 13.4, *p*=0.53 by

unpaired t-test) between the groups **(Figure 3B)**. As there are no available antibodies that can reliably

detect endogenous levels of mouse Cln3 protein[25], we could not assess protein expression.

We next determined the impact of cell type-selective restoration of *Cln3* on histopathological readouts.

Rescue of *Cln3* expression in either astrocytes (Flex*Cln3* / *Cln3*$^{\Delta ex78/\Delta ex78}$ / *Gfap*-Cre) or neurons

(Flex*Cln3* / *Cln3*$^{\Delta ex78/\Delta ex78}$ / *Syn1*-Cre) alone did not prevent storage accumulation in the hippocampus or

motor cortex in 6 or 12-month-old mice **(Figure 3C-L, Sup Figures 8-9)**. In addition to storage

accumulation, CLN3 disease mice develop progressive, reactive astrocytosis. At twelve months of age,

both the hippocampus and motor cortex of *Cln3*$^{\Delta ex78/\Delta ex78}$ showed increased astrocytosis as compared to

wild-type controls. Rescue of *Cln3* expression in either neurons or astrocytes alone did not prevent

astrocytosis **(Sup Figure 10)**.

To investigate if the Flex*Cln3* transgene could prevent storage accumulation in the specific cell type

where it was expressed, we investigated co-localization of autofluorescent lipofuscin with a neuronal

marker, NeuN, in cortical tissue **(Figure 4)**. There is little lipofuscin accumulation in *Cln3*$^{WT/WT}$ cortex

(Figure 4, top row), as compared to *Cln3*$^{\Delta ex78/\Delta ex78}$ cortex, where there is storage in both neurons and non-

neuronal cells (**Figure 4, second row)**. Ubiquitous Flex*Cln3* allele expression (in the presence of *Zp3*-

Cre) prevented storage accumulation in all cell types **(Figure 4, third row)**. Flex*Cln3* allele expression

in neurons (in the presence of *Syn1*-Cre) did not prevent storage in either neurons or non-neuronal cells

(Figure 4, bottom row).

Next, we determined if restoring *Cln3* expression could improve electrophysiologic measures, despite

residual storage accumulation. *In vitro* voltage sensitive dye imaging of hippocampal slices from 6-

month-old animals was completed in Flex*Cln3* / *Cln3*$^{\Delta ex78/\Delta ex78}$ / *Gfap*-Cre, Flex*Cln3* / *Cln3*$^{\Delta ex78/\Delta ex78}$ /

Syn1-Cre*, and* Flex*Cln/Cln3*$^{\Delta ex78/\Delta ex78}$ /Zp3 mice. In this early-stage of disease, the internal blade of the

dentate gyrus of the *Cln3*$^{\Delta ex78/\Delta ex78}$ hippocampus was hypoexcitable in response to perforant path

stimulation as compared to heterozygous *Cln3*$^{\Delta 78/WT}$ mice **(Figure 5A-C)**, consistent with our previous

studies[12]. Specifically, 32.7% of pixels in the internal blade of the *Cln3*$^{\Delta ex78/\Delta ex78}$ dentate gyrus were

significantly hypoexcitable as compared to slices obtained from heterozygous control animals.

Interestingly, restoring expression in astrocytes did not correct hypoexcitability of the dentate gyrus

(Figure 5D-E). However, restoring *Cln3* expression in neurons was sufficient to prevent disease-induced

hypoexcitability **(Figure 5F-G)**, in fact excitability was mildly increased as compared to controls. This

rescue of dynamics occurred despite the presence of residual histopathology **(Figure 3-4 and Sup.**

Figures 8-10).

To evaluate how cell-type specific correction modulates CNS functional defects *in vivo,* we measured

network activity through long-term EEG recordings. While restoration of *Cln3* expression in either

astrocytes or neurons prevented epileptiform spikes throughout the brain (**Figure 6A**), only rescue in

neurons corrected EEG background frequency composition (i.e. frequency band power) and thereby fully

normalized network activity **(Figure 6B-C)**. Intriguingly, this correction of physiology occurred despite

the presence of substantial storage burden, revealing for the first time that loss of Cln3 protein function in

neurons, not storage burden, underlies functional defects in CLN3 disease.

Discussion

Previous cellular-level studies of CLN3 disease have demonstrated disease-associated abnormalities in both neurons and astrocytes[4,7-9,11,26-28]. Prior to this work, it was unknown how these cellular defects translate to the circuit-level dysfunction that underlies neurological symptoms such as dementia and seizures that CLN3 disease patients experience. This work highlights the fact that neurons underlie functional defects in CLN3 disease and must be targeted in future drug development efforts.

A number of novel therapies, including small molecule[5,29-33], antisense oligonucleotides[34], and gene replacement [35,36] strategies are under development for CLN3 disease. These approaches will not target all cell types equally. For example, studies in non-human primates have demonstrated that intravenous administration of AAV9 in juvenile non-human primates predominately transduces glia[21,37] while direct administration into the cerebral spinal fluid (CSF) improves delivery to neurons[38] (34). Our results suggest that maximizing neuronal correction will be essential to develop a treatment that improves functional outcomes in CLN3 disease.

In previous studies, $Cln3^{\Delta ex78/\Delta ex78}$ mice have demonstrated abnormal astrocytic-neuronal coupling[7]. Monogenic mouse models of isolated astrocyte dysfunction have epileptiform spikes and seizures on EEG because of pathologic cross-talk between astrocytes and neurons[39]. In the Flex*Cln3* mouse, while rescue of *Cln3* expression in astrocytes did not normalize EEG background, it was sufficient to prevent epileptiform spikes. The reduction in spiking likely reflects improved astrocytic-neuronal interactions when *Cln3* expression is restored in astrocytes. Collectively, our data suggests that diseased astrocytes may modulate network-level dysfunction in CLN3 disease, but diseased neurons are the primary driver of functional deficits.

Using our cell-type specific rescue model, we found that storage accumulation was prevented when *Cln3* was reintroduced to all cell types embryonically, but not when that correction was restricted to astrocytes

or neurons alone. Interestingly, prevention of storage accumulation is not required for correcting neuronal

network physiology, suggesting that storage material does not underlie functional defects and may be an

innocent bystander in the disease process. These studies highlight the importance of using clinically-

relevant functional measures, such as *in vivo* electrophysiology studies, to guide treatment development.

Histologic measures, such as storage burden, may be poor surrogate biomarkers of therapeutic efficacy in

CLN3 disease and possibly other neuronopathic LSDs.

In summary, partial restoration of Cln3 expression by AAV vectors or through Cre-based systems shows

that neuronal expression is critical for preventing deficits in CNS functional circuits in CLN3 disease

mice. This work also highlights that improvement in storage burden may be a poor readout of therapeutic

efficacy in CLN3 disease, a phenomenon that should be considered by the broader LSD community.

Materials and Methods

For additional information, see Supplemental Methods.

Study approval: All animal protocols were approved by the Institutional Animal Care and Use Committee

at the Children's Hospital of Philadelphia.

Statistical Analysis: All statistical analysis of VSDI rasters was completed using a permutation sampling

method and the VSDI toolbox in MATLAB (Mathworks) as previously described[12,40]. For summary

statistics of regions or groups, GraphPad Prism was also used. In graphs, data are expressed as mean ±

SEM. The statistical significance of the observed differences between 2 groups was assessed by two-

tailed t-test (for normally distributed data) or Mann-Whitney U test (for non-normally distributed data).

For grouped data sets that were not normally distributed a Kruskal-Wallis test followed by a multiple

comparisons test was used as detailed in each figure legend. For grouped data sets that were normally

distributed, 1-way ANOVA or 2-way ANOVA followed by multiple-comparisons test was completed as

detailed in each figure legend. Results were considered significant when $p < 0.05$. Male and female mice

were included in all groups.

Acknowledgments: NINDS R21NS084424 and NICHD HD33532 to BLD, NIH K08NS105865 to RAN,

NCL Stiftung Research Award to RAN, Children's Hospital of Philadelphia Research Institute support to

RAN and BLD.

Conflicts of interests: BLD is a founder of Spark Therapeutics and Spirovant is on the scientific advisory

boards of Intellia Therapeutics, Homology Medicines, Prevail Therapeutics, Resilience, Moment Bio,

Spirovant, Saliogen and Panorama Medicines. No other authors report conflicts of interest.

Author contributions:

Design of study: RA, LT, AH, OK, RC, EL, EM, CS, BD

Data acquisition and analysis of data: RA, LT, AH, OK, RC, EL, CS

Drafting of manuscript: RA, LT, AH, OK, CS

Critical review of data and manuscript: RA, LT, EM, CS, BD

Figures

Figure 1: Intracerebroventricular AAV9.CLN3 gene therapy given at p0 normalizes neuronal

network dynamics *in vivo* at 12 months of age. (A) p0 ICV injection of AAV9.Cln3 restored Cln3

expression in *Cln3*$^{\Delta ex78/\Delta ex78}$ mouse brain at 12 months of age, with levels in the *Cln3*$^{\Delta ex78/\Delta ex78}$ hemisphere

contralateral to the injection site reaching 24.5% ± 4.7 of heterozygous controls. **(B)** Subunit C of the

mitochondrial ATP synthase (SCMAS) accumulation is partially corrected in the hippocampus ipsilateral

to the injection site of *Cln3*$^{\Delta ex78/\Delta ex78}$ mice treated with AAV9.CLN3. Gene replacement corrects circuit

function measured on intracranial EEG including **(C)** epileptiform spiking rates throughout the cortex

(average of all cortical leads shown) and **(D)** background frequency band composition in the motor

cortex. Groups: *Cln3*$^{WT/\Delta ex78}$ (black); *Cln3*$^{\Delta ex78/\Delta ex78}$ (red); *Cln3*$^{\Delta ex78/\Delta ex78}$ + AAV9.CLN3 (gray). N=5-6

animals/condition. Results from 12 hours of recordings were averaged for each animal shown as an

individual circle. Spikes analyzed using a non-parametric Kruskal-Wallis test followed by Dunn's

multiple comparisons test. Frequency bands were analyzed by two-way ANOVA by band and genotype,

followed by Bonferonni's multiple comparisons test (significant results of multiple comparison testing

shown as $*p<0.05$, $**p<0.01$, $***p<0.001$, $****p<0.0001$). Data are shown as mean +/- s.e.m.

Figure 2: Expression of Flex*Cln3* allele in all cells reduces storage accumulation in *Cln3*^{Δex78/Δex78}

brain. (A) In the presence of Cre, the Flex*Cln3* allele undergoes inversion (Step 1) and excision (Step 2)

and is expressed under control of its endogenous promoter. **(B)** PCR of genomic DNA from hippocampus

using orientation-specific primers confirms that the Flex*Cln3* is in the forward orientation only in the

presence of Cre (see Sup Figure 7 for uncut gels). **(C)** In the presence of *Zp3*-Cre which drives ubiquitous

recombination, the Flex*Cln3* allele results in expression levels that are 40.0 +/- 0.4% of heterozygous

controls. **(D)** Subunit C of the mitochondrial ATP synthase (SCMAS) accumulates in the hippocampus of

Cln3^{Δex78/Δex78} and Flex*Cln3* / *Cln3*^{Δex78/Δex78} mice. Storage is reduced in mice with the Flex*Cln3* allele in

the forward confirmation.

Figure 3: Expression of Flex*Cln3* allele in either neurons or astrocytes alone does not prevent storage accumulation. (A) In the presence of Cre, the Flex*Cln3* allele undergoes inversion as detected by PCR with orientation-specific primers of genomic DNA from hippocampus. As both the *Gfap*-Cre and *Syn1*-Cre only drive recombination in a subset of cells, both the forward and reverse alleles are detected (see Sup Figure 7 for uncut gels). **(B)** Expression levels as measured by ΔCt values (FlexCln3-βactin) are similar in Flex*Cln3* brain in the presence of *Gfap*-Cre and *Syn1*-Cre (14.1 vs. 13.4, *p*=0.53 by unpaired t-test). **(C-G)** Expression of Flex*Cln3* in astrocytes (Flex*Cln3* / *Cln3*$^{\Delta ex78/\Delta ex78}$ / *Gfap*-Cre animals) or neurons alone (Flex*Cln3* / *Cln3*$^{\Delta ex78/\Delta ex78}$ / *Syn1*-Cre animals) does not prevent accumulation of SCMAS (green). **(H-K)** Similar findings were seen in the motor cortex. N=4-5 animals, non-parametric Kruskal-

330   Wallis test followed by Benjamini and Hochberg multiple comparisons correction to detect differences

331   from control. For all panels: *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$.

Figure 4: Expression of the Flex*Cln3* allele in neurons alone does not prevent autofluorescent lipofuscin accumulation in cortical neurons. (Top row) There is little autofluorescent lipofuscin accumulation (green) in wild-type cortical neurons (red) (blue DAPI) as compared to **(second row)** *Cln3*^{Δ78/Δ78} cortical neurons in 12 month old animals. **(Third row)** *Zp3*-Cre induced expression of the Flex*Cln3* allele in all cell types prevents autofluorescent lipofuscin accumulation in both neuronal and non-neuronal cells. **(Bottom row)** Flex*Cln3* expression in neurons (*Syn1*-Cre) did not prevent storage accumulation in neurons (arrow heads) or non-neuronal cells (arrows).

Figure 5: Restoring low level *Cln3* expression in neurons alone corrects hypoexcitability of the

dentate gyrus. Dentate gyrus excitability measured through *in vitro* voltage sensitive dye imaging of

hippocampal slices from 6-month-old animals is shown as raster plots of average fluorescence change

(ΔF/F warm colors excitation, cool colors inhibition) over time (X axis) and location within the internal

blade of the dentate gyrus (Y Axis). **(A)** Four-pulse stimulation of the perforant path in heterozygous

control (*Cln3*$^{\Delta ex78/WT}$) mice reveals robust excitation of the dentate gyrus (DG) as compared to **(B)**

Cln3$^{\Delta ex78/\Delta ex78}$ mice**. (C)** *Cln3*$^{\Delta ex78/WT}$ vs. *Cln3*$^{\Delta ex78/\Delta ex78}$ rasters were compared using a permutation

sampling method with 1000 trials. Pixels with $p > 0.05$ are masked in gray. For regions of significant

difference in excitability with $p < 0.05$, the difference in fluorescence change (ΔF/F$_{\Delta ex78/WT}$ -ΔF/F$_{\Delta ex78/\Delta ex78}$)

is shown. **(D-E)** In animals expressing *Cln3* in astrocytes (Flex*Cln3* / *Cln3*$^{\Delta ex78/\Delta ex78}$ / *Gfap*-Cre), the DG

remained hypoexcitable as compared to the *Cln3*$^{\Delta ex78/WT}$ DG. **(F-G)** Expression of *Cln3* in neurons alone

(Flex*Cln3* / *Cln3*$^{\Delta ex78/\Delta ex78}$ / *Syn1*-Cre) was sufficient to reverse hypoexcitability of the DG as compared to

Cln3$^{\Delta ex78/WT}$ animals. Group sizes (n=slices, N=mice): *Cln3*$^{\Delta ex78/WT}$ n= 23 N=4; *Cln3*$^{\Delta ex78/\Delta ex78}$ n= 30, N=6;

Flex*Cln3* / *Cln3*$^{\Delta ex78/\Delta ex78}$ / *Gfap-Cre* n=19 slices, N=4; Flex*Cln3* / *Cln3*$^{\Delta ex78/\Delta ex78}$ / *Syn1-Cre* n=17; N=4.

Figure 6: Cln3 rescue in either astrocytes or neurons reduces epileptiform spikes, but only neuronal rescue fully normalizes network dynamics as measured on EEG. (A) Correction of either astroctyes (Flex*Cln3* / *Cln3*$^{\Delta ex78/\Delta ex78}$ / *Gfap*-Cre, blue) or neurons (Flex*Cln3* / *Cln3*$^{\Delta ex78/\Delta ex78}$ / *Syn1*-Cre, green) reduced spike burden on EEG in 12 month old mice. **(B-E)** Example EEG traces from *Cln3*$^{\Delta ex78/\Delta ex78}$ (red) and Flex*Cln3* / *Cln3*$^{\Delta ex78/\Delta ex78}$ / *Gfap*-Cre motor cortex (blue) show decreased slow delta activity and increased fast activity as compared to control motor cortex (black). EEG trace from Flex*Cln3* / *Cln3*$^{\Delta ex78/\Delta ex78}$ / *Syn1*-Cre motor cortex (green) shows that rescue of *Cln3* expression in neurons normalizes EEG background frequency composition. **(F)** Power spectral analysis of EEG data recorded in the motor cortex confirms that rescue of *Cln3* expression in astrocytes (blue) does not correct disease-associated loss of delta activity, and actually exacerbates increased fast gamma activity. However, correction in

373   neurons (green) normalizes EEG frequency composition. N=4-5 animals/genotype, after passing a

374   Shapiro-Wilk test of normality, data were analyzed by two-way ANOVA, followed by Sidak's multiple

375   comparisons test. Significant differences from multiple comparisons testing are shown as *$p<0.05$,

376   **$p<0.01$, ***$p<0.001$. Frequency data in panel F in $Cln3^{WT/\Delta ex78}$ and $Cln3^{\Delta ex78/\Delta ex78}$ also shown in Fig 1D.

References

1. Platt, F.M. (2018). Emptying the stores: lysosomal diseases and therapeutic strategies.
Nat Rev Drug Discov *17*, 133-150. 10.1038/nrd.2017.214.
2. Cotman, S.L., Vrbanac, V., Lebel, L.A., Lee, R.L., Johnson, K.A., Donahue, L.R., Teed,
A.M., Antonellis, K., Bronson, R.T., Lerner, T.J., and MacDonald, M.E. (2002).
Cln3(Deltaex7/8) knock-in mice with the common JNCL mutation exhibit progressive
neurologic disease that begins before birth. Hum Mol Genet *11*, 2709-2721.
10.1093/hmg/11.22.2709.
3. Eliason, S.L., Stein, C.S., Mao, Q., Tecedor, L., Ding, S.L., Gaines, D.M., and Davidson,
B.L. (2007). A knock-in reporter model of Batten disease. J Neurosci *27*, 9826-9834.
10.1523/JNEUROSCI.1710-07.2007.
4. Tyynela, J., Cooper, J.D., Khan, M.N., Shemilts, S.J., and Haltia, M. (2004).
Hippocampal pathology in the human neuronal ceroid-lipofuscinoses: distinct patterns of
storage deposition, neurodegeneration and glial activation. Brain Pathol *14*, 349-357.
10.1111/j.1750-3639.2004.tb00077.x.
5. Tarczyluk-Wells, M.A., Salzlechner, C., Najafi, A.R., Lim, M.J., Smith, D., Platt, F.M.,
Williams, B.P., and Cooper, J.D. (2019). Combined Anti-inflammatory and
Neuroprotective Treatments Have the Potential to Impact Disease Phenotypes in Cln3 (-/-
) Mice. Front Neurol *10*, 963. 10.3389/fneur.2019.00963.
6. Xiong, J., and Kielian, T. (2013). Microglia in juvenile neuronal ceroid lipofuscinosis are
primed toward a pro-inflammatory phenotype. J Neurochem *127*, 245-258.
10.1111/jnc.12385.
7. Burkovetskaya, M., Karpuk, N., Xiong, J., Bosch, M., Boska, M.D., Takeuchi, H.,
Suzumura, A., and Kielian, T. (2014). Evidence for aberrant astrocyte hemichannel
activity in Juvenile Neuronal Ceroid Lipofuscinosis (JNCL). PLoS One *9*, e95023.
10.1371/journal.pone.0095023.
8. Parviainen, L., Dihanich, S., Anderson, G.W., Wong, A.M., Brooks, H.R., Abeti, R.,
Rezaie, P., Lalli, G., Pope, S., Heales, S.J., Mitchison, H.M., et al. (2017). Glial cells are
functionally impaired in juvenile neuronal ceroid lipofuscinosis and detrimental to
neurons. Acta Neuropathol Commun *5*, 74. 10.1186/s40478-017-0476-y.
9. Pontikis, C.C., Cella, C.V., Parihar, N., Lim, M.J., Chakrabarti, S., Mitchison, H.M.,
Mobley, W.C., Rezaie, P., Pearce, D.A., and Cooper, J.D. (2004). Late onset
neurodegeneration in the Cln3-/- mouse model of juvenile neuronal ceroid lipofuscinosis
is preceded by low level glial activation. Brain Res *1023*, 231-242.
10.1016/j.brainres.2004.07.030.
10. Brenneman, D.E., Pearce, D.A., Kovacs, A., and DeFrees, S. (2017). Pharmacological
Effects on Ceroid Lipofuscin and Neuronal Structure in Cln3 (ex7/8) Mouse Brain
Cultures. J Mol Neurosci *63*, 100-114. 10.1007/s12031-017-0962-5.
11. Grunewald, B., Lange, M.D., Werner, C., O'Leary, A., Weishaupt, A., Popp, S., Pearce,
D.A., Wiendl, H., Reif, A., Pape, H.C., Toyka, K.V., et al. (2017). Defective synaptic
transmission causes disease signs in a mouse model of juvenile neuronal ceroid
lipofuscinosis. Elife *6*. 10.7554/eLife.28685.
12. Ahrens-Nicklas, R.C., Tecedor, L., Hall, A.F., Lysenko, E., Cohen, A.S., Davidson, B.L.,
and Marsh, E.D. (2019). Neuronal network dysfunction precedes storage and
neurodegeneration in a lysosomal storage disorder. JCI Insight *4*.
10.1172/jci.insight.131961.

13.   Mitchison, H.M., Lim, M.J., and Cooper, J.D. (2004). Selectivity and types of cell death
428   in the neuronal ceroid lipofuscinoses. Brain Pathol *14*, 86-96. 10.1111/j.1750-
429   3639.2004.tb00502.x.
430   14.   Kwak, Y.T. (2006). Quantitative EEG findings in different stages of Alzheimer's disease.
431   J Clin Neurophysiol *23*, 456-461. 10.1097/01.wnp.0000223453.47663.63.
432   15.   Kent, B.A., Strittmatter, S.M., and Nygaard, H.B. (2018). Sleep and EEG Power Spectral
433   Analysis in Three Transgenic Mouse Models of Alzheimer's Disease: APP/PS1,
434   3xTgAD, and Tg2576. J Alzheimers Dis *64*, 1325-1336. 10.3233/JAD-180260.
435   16.   Faller, K.M., Gutierrez-Quintana, R., Mohammed, A., Rahim, A.A., Tuxworth, R.I.,
436   Wager, K., and Bond, M. (2015). The neuronal ceroid lipofuscinoses: Opportunities from
437   model systems. Biochim Biophys Acta *1852*, 2267-2278. 10.1016/j.bbadis.2015.04.022.
438   17.   Johnson, T.B., Langin, L.M., Zhao, J., Weimer, J.M., Pearce, D.A., and Kovacs, A.D.
439   (2019). Changes in motor behavior, neuropathology, and gut microbiota of a Batten
440   disease mouse model following administration of acidified drinking water. Sci Rep *9*,
441   14962. 10.1038/s41598-019-51488-z.
442   18.   Hammond, S.L., Leek, A.N., Richman, E.H., and Tjalkens, R.B. (2017). Cellular
443   selectivity of AAV serotypes for gene delivery in neurons and astrocytes by neonatal
444   intracerebroventricular injection. PLoS One *12*, e0188830.
445   10.1371/journal.pone.0188830.
446   19.   Hinderer, C., Bell, P., Vite, C.H., Louboutin, J.P., Grant, R., Bote, E., Yu, H., Pukenas,
447   B., Hurst, R., and Wilson, J.M. (2014). Widespread gene transfer in the central nervous
448   system of cynomolgus macaques following delivery of AAV9 into the cisterna magna.
449   Mol Ther Methods Clin Dev *1*, 14051. 10.1038/mtm.2014.51.
450   20.   Gray, S.J., Nagabhushan Kalburgi, S., McCown, T.J., and Jude Samulski, R. (2013).
451   Global CNS gene delivery and evasion of anti-AAV-neutralizing antibodies by
452   intrathecal AAV administration in non-human primates. Gene Ther *20*, 450-459.
453   10.1038/gt.2012.101.
454   21.   Samaranch, L., Salegio, E.A., San Sebastian, W., Kells, A.P., Foust, K.D., Bringas, J.R.,
455   Lamarre, C., Forsayeth, J., Kaspar, B.K., and Bankiewicz, K.S. (2012). Adeno-associated
456   virus serotype 9 transduction in the central nervous system of nonhuman primates. Hum
457   Gene Ther *23*, 382-389. 10.1089/hum.2011.200.
458   22.   Shafi, R., Iyer, S.P., Ellies, L.G., O'Donnell, N., Marek, K.W., Chui, D., Hart, G.W., and
459   Marth, J.D. (2000). The O-GlcNAc transferase gene resides on the X chromosome and is
460   essential for embryonic stem cell viability and mouse ontogeny. Proc Natl Acad Sci U S
461   A *97*, 5735-5739. 10.1073/pnas.100471497.
462   23.   Gregorian, C., Nakashima, J., Le Belle, J., Ohab, J., Kim, R., Liu, A., Smith, K.B.,
463   Groszer, M., Garcia, A.D., Sofroniew, M.V., Carmichael, S.T., et al. (2009). Pten
464   deletion in adult neural stem/progenitor cells enhances constitutive neurogenesis. J
465   Neurosci *29*, 1874-1886. 10.1523/JNEUROSCI.3095-08.2009.
466   24.   Zhu, Y., Romero, M.I., Ghosh, P., Ye, Z., Charnay, P., Rushing, E.J., Marth, J.D., and
467   Parada, L.F. (2001). Ablation of NF1 function in neurons induces abnormal development
468   of cerebral cortex and reactive gliosis in the brain. Genes Dev *15*, 859-876.
469   10.1101/gad.862101.
470   25.   Nelson, T., Pearce, D.A., and Kovacs, A.D. (2017). Lack of specificity of antibodies
471   raised against CLN3, the lysosomal/endosomal transmembrane protein mutated in
472   juvenile Batten disease. Biosci Rep *37*. 10.1042/BSR20171229.

473   26.   Pontikis, C.C., Cotman, S.L., MacDonald, M.E., and Cooper, J.D. (2005).
474   Thalamocortical neuron loss and localized astrocytosis in the Cln3Deltaex7/8 knock-in
475   mouse model of Batten disease. Neurobiol Dis *20*, 823-836. 10.1016/j.nbd.2005.05.018.
476   27.   Bosch, M.E., and Kielian, T. (2019). Astrocytes in juvenile neuronal ceroid lipofuscinosis
477   (CLN3) display metabolic and calcium signaling abnormalities. J Neurochem *148*, 612-
478   624. 10.1111/jnc.14545.
479   28.   Zhong, Y., Mohan, K., Liu, J., Al-Attar, A., Lin, P., Flight, R.M., Sun, Q., Warmoes,
480   M.O., Deshpande, R.R., Liu, H., Jung, K.S., et al. (2020). Loss of CLN3, the gene
481   mutated in juvenile neuronal ceroid lipofuscinosis, leads to metabolic impairment and
482   autophagy induction in retinal pigment epithelium. Biochim Biophys Acta Mol Basis Dis
483   *1866*, 165883. 10.1016/j.bbadis.2020.165883.
484   29.   Schultz, M.L., Tecedor, L., Lysenko, E., Ramachandran, S., Stein, C.S., and Davidson,
485   B.L. (2018). Modulating membrane fluidity corrects Batten disease phenotypes in vitro
486   and in vivo. Neurobiol Dis *115*, 182-193. 10.1016/j.nbd.2018.04.010.
487   30.   Augustine, E.F., Beck, C.A., Adams, H.R., Defendorf, S., Vierhile, A., Timm, D.,
488   Weimer, J.M., Mink, J.W., and Marshall, F.J. (2019). Short-Term Administration of
489   Mycophenolate Is Well-Tolerated in CLN3 Disease (Juvenile Neuronal Ceroid
490   Lipofuscinosis). JIMD Rep *43*, 117-124. 10.1007/8904_2018_113.
491   31.   El-Sitt, S., Soueid, J., Maalouf, K., Makhoul, N., Al Ali, J., Makoukji, J., Asser, B.,
492   Daou, D., Harati, H., and Boustany, R.M. (2019). Exogenous Galactosylceramide as
493   Potential Treatment for CLN3 Disease. Ann Neurol *86*, 729-742. 10.1002/ana.25573.
494   32.   Petcherski, A., Chandrachud, U., Butz, E.S., Klein, M.C., Zhao, W.N., Reis, S.A.,
495   Haggarty, S.J., Ruonala, M.O., and Cotman, S.L. (2019). An Autophagy Modifier Screen
496   Identifies Small Molecules Capable of Reducing Autophagosome Accumulation in a
497   Model of CLN3-Mediated Neurodegeneration. Cells *8*. 10.3390/cells8121531.
498   33.   Maalouf, K., Makoukji, J., Saab, S., Makhoul, N.J., Carmona, A.V., Kinarivala, N.,
499   Ghanem, N., Trippier, P.C., and Boustany, R.M. (2020). Exogenous Flupirtine as
500   Potential Treatment for CLN3 Disease. Cells *9*. 10.3390/cells9081872.
501   34.   Centa, J.L., Jodelka, F.M., Hinrich, A.J., Johnson, T.B., Ochaba, J., Jackson, M., Duelli,
502   D.M., Weimer, J.M., Rigo, F., and Hastings, M.L. (2020). Therapeutic efficacy of
503   antisense oligonucleotides in mouse models of CLN3 Batten disease. Nat Med *26*, 1444-
504   1451. 10.1038/s41591-020-0986-1.
505   35.   Bosch, M.E., Aldrich, A., Fallet, R., Odvody, J., Burkovetskaya, M., Schuberth, K.,
506   Fitzgerald, J.A., Foust, K.D., and Kielian, T. (2016). Self-Complementary AAV9 Gene
507   Delivery Partially Corrects Pathology Associated with Juvenile Neuronal Ceroid
508   Lipofuscinosis (CLN3). J Neurosci *36*, 9669-9682. 10.1523/JNEUROSCI.1635-16.2016.
509   36.   Kleine Holthaus, S.M., Aristorena, M., Maswood, R., Semenyuk, O., Hoke, J., Hare, A.,
510   Smith, A.J., Mole, S.E., and Ali, R.R. (2020). Gene Therapy Targeting the Inner Retina
511   Rescues the Retinal Phenotype in a Mouse Model of CLN3 Batten Disease. Hum Gene
512   Ther *31*, 709-718. 10.1089/hum.2020.038.
513   37.   Gray, S.J., Matagne, V., Bachaboina, L., Yadav, S., Ojeda, S.R., and Samulski, R.J.
514   (2011). Preclinical differences of intravascular AAV9 delivery to neurons and glia: a
515   comparative study of adult mice and nonhuman primates. Mol Ther *19*, 1058-1069.
516   10.1038/mt.2011.72.
517   38.   Bey, K., Deniaud, J., Dubreil, L., Joussemet, B., Cristini, J., Ciron, C., Hordeaux, J., Le
518   Boulc'h, M., Marche, K., Maquigneau, M., Guilbaud, M., et al. (2020). Intra-CSF AAV9

and AAVrh10 Administration in Nonhuman Primates: Promising Routes and Vectors for
Which Neurological Diseases? Mol Ther Methods Clin Dev *17*, 771-784.
10.1016/j.omtm.2020.04.001.
39. Robel, S., Buckingham, S.C., Boni, J.L., Campbell, S.L., Danbolt, N.C., Riedemann, T.,
Sutor, B., and Sontheimer, H. (2015). Reactive astrogliosis causes the development of
spontaneous seizures. J Neurosci *35*, 3330-3345. 10.1523/JNEUROSCI.1574-14.2015.
40. Bourgeois, E.B., Johnson, B.N., McCoy, A.J., Trippa, L., Cohen, A.S., and Marsh, E.D.
(2014). A toolbox for spatiotemporal analysis of voltage-sensitive dye imaging data in
brain slices. PLoS One *9*, e108686. 10.1371/journal.pone.0108686.
528