Fixed or random? On the reliability of mixed-effects models for a small number of levels in grouping variables

Authors and Affiliations:

Johannes Oberpriller[1*], Melina de Souza Leite[1,2], Maximilian Pichler[1*]

[1] Theoretical Ecology, University of Regensburg, Universitätsstraße 31, 93053 Regensburg, Germany

[2] Department of Ecology, University of São Paulo, Rua do Matão 321, Travessa 14, São Paulo, SP CEP 05508-090, Brazil

* These authors contributed equally

Keywords:

Mixed-effects models, generalized linear models, multilevel models, hierarchical models, fixed effects, random effects

Corresponding author: Johannes Oberpriller, Universitätsstraße 31, 93051 Regensburg, Germany, johannes.oberpriller@ur.de

Abstract

1. Biological data are often intrinsically hierarchical. Due to their ability to account for

such dependencies, mixed-effects models have become a common analysis

technique in ecology and evolution. While many questions around their theoretical

foundations and practical applications are solved, one fundamental question is still

highly debated: When facing a low number of levels should we model a grouping

(blocking, clustering) variable as a random or fixed effect? In such situation, the

variance of the random effect is imprecise, but whether this affects the statistical

properties of the population effect is unclear.

2. Here, we analyzed the consequences of including a grouping variable as fixed or

random effect in the correctly specified and other possible miss-specified models (too

simple or too complex models) for data with small number of levels (2 - 8). For all

these options, we calculated type I error rates and power. Moreover, we show how

these statistical properties change with the study design.

3. We found that the model choice does not influence the statistical properties of the

population effect when the effect is the same at all levels However, if an ecological

effect differs among levels, using a random slope and intercept model, and switching

to a fixed-effect model only in case of a singular fit, avoids overconfidence in the

results. Additionally, power and type I error are strongly influenced by the number of

and difference between levels.

4. We conclude that inferring the correct random effect structure is of high importance

to get correct statistical properties. When in doubt, we recommend starting with the

simpler model and using model diagnostics to identify missing components. When

having identified the correct structure, we encourage to start with a mixed-effects

model independent of the number of levels and switch to a fixed-effect model only in

case of a singular fit. With these recommendations, we allow for more informative

choices about study design and data analysis and thus make ecological inference

with mixed-effects models more robust for small number of levels.

Introduction

Many biological data from experimental or observational studies have a hierarchical

grouping (or blocking or clustering) structure that introduces dependencies among

observations (McMahon & Diez 2007; Bolker *et al.* 2009; Zuur *et al.* 2009; Harrison *et al.*

2018). When conducting statistical analyses, models have to reflect these dependencies in

their assumptions for correct statistical properties (Arnqvist 2020), a task for which linear

and generalized mixed-effects models (LMMs or GLMMs) were designed (Laird & Ware

1982; Chen & Dunson 2003). Mixed-effects models have also replaced ANOVAs as the

common tool for variance analysis (Wainwright *et al.* 2007; Boisgontier & Cheval 2016)

because they allow simultaneous analysis of variance at different hierarchical levels

(Krueger & Tian 2004; Boisgontier & Cheval 2016), handle unbalanced study designs better

(Swallow & Monahan 1984; Lindstrom & Bates 1988; Pinheiro & Bates 1995; Littell 2002),

and have better statistical properties for missing data (Baayen *et al.* 2008). Therefore,

mixed-effects models have become one of the most popular methods in ecology and

evolution (Bolker *et al.* 2009) and other fields (e.g., psychology, see Meteyard & Davies

2020).

The ability of mixed-effects models to adapt to different data structures (i.e. their flexibility to

handle many different hierarchical structures, see Box 1), which makes them in the first

place so powerful (Wainwright *et al.* 2007), raises discussions about their proper application

(Nakagawa & Schielzeth 2013; Dixon 2016). These issues include data-related problems

such as overdispersion (Harrison 2014, 2015), robustness to wrong distributional

assumptions (Schielzeth *et al.* 2020), and their correct evaluation (Nakagawa & Schielzeth

2013). Despite these rather technical challenges, there are also application-oriented issues

(Harrison *et al.* 2017; Meteyard & Davies 2020) such as the question whether to use the

most complex random effect structure (Barr *et al.* 2013; but see Matuschek *et al.* 2017), their

correct interpretation (e.g. Dixon 2016), or in which situations we should model a grouping

variable as random or fixed effect (Harrison *et al.* 2018).

A priori, modeling a grouping (or blocking) variable as fixed or random effect are equally well

suited for multilevel analysis (Townsend *et al.* 2013; Kadane 2020) and strict rules don't

exist because the best strategy generally depends on the goal of the analysis (Gelman &

Hill 2007, see Box 2). For instance, for unbalanced study designs, random effect estimates

incorporate between and within group information whereas the corresponding fixed-effect

model (grouping variable is specified as a fixed effect) only within group information(McLean

et al. 1991; Dixon 2016; Shaver 2019; but see Giesselmann & Schmidt-Catran 2020). This

is important when one is interested in the actual level effects themselves (also called narrow-

sense inference analysis), but rather negligible when only interested in the population effect

(broad-sense inference analysis), where the individual levels of the grouping variable are

only nuisance parameters.

These inferential properties stem from different assumptions underlying these two options

(specifying the grouping variable as fixed or random) (Millar & Anderson 2004). The most

prominent difference is that modeling a grouping variable as random effect implicitly

assumes that the individual levels of the grouping variable are realizations of a common

distribution, usually a normal distribution, for which the variance and the mean (the

population effect) are unknown and need to be estimated (e.g. DerSimonian & Laird 1986).

This assumption shrinks the estimates of each random effect level to the mean of the

underlying distribution. In contrast, modeling a grouping variable as a fixed effect makes no

distributional assumptions about the individual level estimates (i.e., making them

independent of each other and thus no between level information is used to estimate the

effects). Thus, compared to when modeling the grouping variable as a fixed effect, the

random effect model has to estimate a lower effective number of parameters (e.g. Gelman

and Hill, 2005), which could lead to higher power to detect significant effects of population

effects at the cost of a small bias towards the mean of the random effect estimates (e.g.

Johnson et al. 2014).

Based on these properties, and if the grouping variable is only a nuisance parameter (broad-

sense inference), random-effect modeling seems preferable over fixed-effect modeling. It is,

however, unclear if these advantages remain when the number of levels is small (cf. also

Harrison *et al.* 2018), because this might cause an imprecise and biased random effects'

variance estimate (Harrison, 2018), which then could influence the population effect

estimate of the mixed-effects model (Hox *et al.* 2017).

The ecological literature suggests, as a rule of thumb, to only fit a grouping variable as

random effect if it has a minimum of five, sometimes eight, levels to precisely estimate the

random effect' variance (Bolker 2015; Harrison 2015; Harrison *et al.* 2018). With four or

fewer levels in the blocking variable, the preferred alternative is fitting it as a fixed effect

(Gelman & Hill 2007; Bolker *et al.* 2009; Bolker 2015). In other disciplines other thresholds

are proposed, e.g. 10-20 in psychology (McNeish & Stapleton, 2016) or 30-50 in sociology

(Maas & Hox, 2005). To our knowledge, however, none of these values were based on a

systematic analysis of how the choice of modeling the grouping variable affects the statistical

estimates (i.e., type I error rate and power) of the population effects (i.e., the average slope

or intercept over a grouping variable, henceforth called the ecological or population effect).

Here, we analyze a situation where a researcher wants to make inference about the

population effect and draw conclusions for the population of levels that were sampled and

thus has decided to use a mixed-effect model, but is confronted with a low number of levels.

We simulated a hypothetical unbalanced study design on reproductive success and height

of a plant on a temperature gradient (altitudinal gradient among different mountains, see

Box 1) to compare empirical statistical properties with a varying number of levels, blocks, or

experimental units (two to eight mountains, the nuisance parameter) and two different data-

generating processes. To represent the challenge of an analyst to correctly specify the

model structure and the consequences if not, we additionally tested miss-specified models

(too complex or too simple versions of the fixed and mixed-effects models). To investigate

the consequences of these modeling choices on the population effect, we compared: type I

error rates (how often a non-existing temperature effect would be interpreted as existing)

and statistical power (the rate at which a truly existing temperature effect would be

significant). Based on our results and in the context of broad-sense inference, we give

practical recommendations on when to include grouping variables as random effects and in

which situations it might be more beneficial to fall back to specifying them as fixed effects.

Box 1: Example of an ecological study design with grouping/blocking variables

Sampling design. We want to understand the population effect of temperature on the reproductive success of a plant that grows in mountains. We hypothesize that: (H1) increasing temperature (lower altitude) increases the probability of flowering (reproduction), and (H2) the height of flowering plants. To do so, we establish altitudinal transects in many mountains (nuisance variable) and collect information from a certain number of plants.

Sampling design

Problem. The transects are not in the same geographical alignment, the type of soil varies in each mountain, and the plants are genetically very distinct among populations. All these factors introduce differences among populations that are not exactly of our interest (given our hypotheses), but statistically, plants of the same mountain are non-independent observations. The mountains can be considered as grouping, blocking, control, or nuisance parameters.

Modeling options. We may use a mixed-effects model with a random intercept and slope (Box 2) for mountain to account for the differences among populations (colored lines in H1 and H2), while still modeling the relationship of interest as fixed effects (red lines). An alternative may be to use a fixed-effects model, i.e., to include mountain as a categorical predictor (Box 2).

Hypothesis 1 - The reproductive success (flowering or not) of plants increases with temperature.

Hypothesis 2 - The height of flowering plants increases with temperature.

Box 2: Differences between modeling a grouping variable as random or fixed effect

Fixed or random effects. The question of whether to include a hierarchical/blocking variable as random or fixed effect in the analysis depends on several factors. Fixed effects are usually used when the analysts are interested in the individual level estimates of a grouping variable (Bolker *et al.* 2009) and these are independent, mutually exclusive, and completely observed (e.g. control and treatment in experiments, male and female when analyzing differences between sex) (e.g. Hedges & Vevea 1998; Gunasekara *et al.* 2014). Random effects, in comparison, are modeling choices when the variance between the different levels (Bolker *et al.* 2009) and not the exact estimates of the different levels are of interest (e.g. DerSimonian & Laird 1986). Additionally, random effects can be used when not every realization of the underlying mechanism can be observed (e.g. species across a number of observational sites in different geographic areas) but the analysts want to control for its influence (i.e. pseudo-replication, see Arnqvist 2020). The two options differ in their interpretation, mixed-effect models use between- and within-group information whereas fixed-effect models use only within-group information. This subtle difference is important when for instance treatment or group differences are the goal of the analysis. Another important difference is that when modeling the categorical variable as fixed-effect

Figure B 1: Fixed- and mixed-effects models fit to simulated data with random intercept (a,b) and random intercept and slope (c,d) for each mountain in the example from H2 Box 1. Dots represent the data points, lines the individual estimates for each mountain. The red line and shaded red intervals are the mean and its estimated standard deviation for mixed-effects models.

conclusions apply to the levels used in the study, while when modeling as random-effect conclusions apply to the population of levels from where the studied levels were randomly sampled. However, in our example (Box 1), we are mainly interested in the population effect and not in the group differences which makes the inferential distinction negligible. See Gelman (2005) or Gelman & Hill 2007 for more decision criteria for whether an effect is random or fixed.

Technical differences between random and fixed effects. When specifying a grouping variable as fixed effect, the model with default contrast in R estimates the effect of one reference level (see Schielzeth 2010) differences between the reference level and possible linear combinations of other levels (Fig. B1a,c). Thus, fixed-effect models with default contrast don't estimate mean effect over groups (i.e., the population effect), but it can be calculated using e.g. bootstrapping (see Supporting Information S1), with sum-to-zero contrasts, or follow-on packages such as emmeans (Lenth 2021). Mixed-effects models estimate the population effect and its standard deviation and from a Bayesian perspective each individual level effect or from a frequentist perspective predict future realizations of the individual random effect levels – Best Unbiased Linear Predictor (Fig. B1b, d). Blocking variables may not only imply different intercepts (Fig B1 a, b), but also different slopes (Fig B1 c, d - the temperature "ecological" effect). In fixed-effects models, this is done by introducing an interaction between the ecological effect and the grouping variable, which implicitly allows for correlations between slopes and intercepts. With mixed-effects models correlations between random slopes and random intercepts can be disabled. The choice of modeling different slopes and their correlation to intercepts for each group is related to the study design and may have impact on modeling structure and inference.

Models in R. Here are the formula R syntax[1] of fixed and mixed-effects models fitted in Figure B1

Scenario A random intercept only

    a)   Height ~ 0 + Temperature + Mountain (fixed-effect model [2])
    b)   Height ~ Temperature + (1| Mountain) (mixed-effects model)

Scenario B random intercept and slope

c) Height ~ 0 + Mountain + Temperature:Mountain   (fixed-effect model[2])
d) Height ~ Temperature + (1|Mountain) + (0 + Temperature| Mountain) (mixed-effects model, uncorrelated random effects)

e) Height ~ Temperature + (Temperature| Mountain).   (mixed-effects model, correlated random effects)

1 From lm() or glm() base R functions, lme4 (Bates et al. 2015) and glmmTMB (Brooks et al. 2017) packages.
2 This formula parametrization estimates a separate intercept and slope (only in c) for each group independently. In c, the most common parametrization is *Mountain + Temperature + Mountain:Temperature*, which is the same as *Mountain*Temperature*, that uses one group as a reference intercept and slope and the other parameters estimate differences to the reference group. For purposes of prediction as in Fig B1 c, these parametrizations don't change anything.

Methods

Simulated example

To compare random- and fixed-effect modeling of a grouping variable with small number of groups, we simulated data based on our hypothetical example from Box 1. We hypothesized first that higher temperatures improve the reproductive success (either yes or no) of a plant species (H1), and second, that higher temperatures also increase the average height of the reproductive plants (H2). To test these hypotheses, we simulated an unbalanced study design – a common scenario in ecology and evolution (Schielzeth et al. 2020) – with two to eight mountains from a varying number of plants for each mountain (H1: expected range between 40-360 plants per mountain, H2: expected range between 10-90 plants per mountain) while keeping the overall number of plants constant (H1: on average 200 plants per mountain, H2: on average 50 plants per mountain) along altitudinal transects.

We varied the number of mountains (groups) from two to eight and simulated 5,000 datasets for each case. Additionally, we performed a simulation, which varies components of the study design, to quantify their influence on the inference (see section Quantifying the influence of study design on type I error and power). We investigated performance for models with binomial (GLMM and GLM, H1) and normal distribution (LMM and LM, H2). The different choices of observations per group were made to meet the higher data request of binomial models compared to linear models.

Scenarios of data complexity and model fitting

Table 1: Data-generating and tested models for each scenario: Scenario A random intercept for each mountain and B random intercept and slope for each mountain. For the fixed-effects models, we used R syntax for model formula in *lm()*

155 function and for the mixed-effects models we used syntax from *lmer()* or *glmmTMB()* functions from *lme4* and *glmmTMB*

156 packages. Response variables are reproductive success (RS) for the binomial distribution models (H1, Box 1) and height

157 of flowering plants for the normal distribution models (H2, Box 1). T is the temperature effect on both response variables.

		Scenario A Random intercept only		Scenario B Random intercept and slope	Description
Data- generating model	(1)	Height/RS ~ T + (1|mountain)	(6)	Height/RS~ T + (1|mountain) + (0+T|mountain)	Effect of intercept (and slope in B) vary across mountains
Tested models					
Fixed- effect models	(2)	Height/RS ~ T	(7)	Height/RS ~ T	Temperature only main effect – too simple model
	(3)	Height/RS ~ 0 + T + mountain	(8)	Height/RS ~ 0 + mountain + T:mountain	Main effects of temperature and mountain (and interaction in B) – slightly more complex model
Mixed- effects models			(9)	Height/RS ~ T + (1|mountain)	Temperature and mountain both vary – too simple models
	(4)	Height/RS ~ T + (1|mountain)	(10)	Height/RS ~ T + (1|mountain) + (0+T|mountain)	Effect of intercept (and uncorrelated slope temperature in B) vary across mountains – correctly specified models
	(5)	Height/RS ~ T + (1|mountain) + (0+T|mountain)	(11)	Height/RS ~ T + (T|mountain)	Effect of intercept and slope temperature (correlated effects in B) across mountains – too complex models

Scenario A - random intercepts per mountain

In scenario A, we assumed mountains only differ in their intercepts (mean

height/reproductive success, Table 1, Eq. 1). When interested in the temperature effect, this

is an example where the ecological effect is the same for each level of the grouping variable.

For this scenario we tested two different mixed-effects model structures: a correctly specified

model which corresponds to the data generating model (Table 1, Eq. 4) and a too complex

model (Table 1, Eq. 5) with an additional random slope for each mountain. Since in real

studies the true underlying data generating process is unknown, it is useful to understand if

a too complex model can correctly estimate a zero (or nearly zero) random slope and, thus,

approximate the true model structure (Table1, Eq. 1).

As fixed-effect alternatives, we tested the correctly specified model with mountain as fixed

intercept together with temperature as slope (Table1, Eq. 3), and a too simple model omitting

mountain at all (Table 1, Eq. 2). This last model corresponds to a mixed-effects model that

estimates the random intercepts to zero.

Scenario B - random intercepts and random slopes per mountain

In scenario B, we assumed a random intercept and a random slope (without correlation

among them) for each mountain as data-generating process (Table 1, Eq. 6). When

interested in the temperature effect, this is an example where the ecological effect differs

among levels of the grouping variable. We tested three different mixed-effects model

structures: a correctly specified model corresponding to the data generating model (Table

1, Eq. 10), a too complex model with a correlation term for the random intercept and random

slope (Table 1, Eq. 11), and a too simple model with only a random intercept for each

mountain (Table 1, Eq. 9). We used the too simple model to test the effect of not accounting

for important contributions to the data-generating process. Note, however, only in case of

balanced designs and linear models the population effect estimate from the too simple model

is consistent with the full model, because of different weighting schemes (for unbalanced

designs) and the fact that the mean of a non-linear transformation (here link function) of

estimates is not the same as the non-linear transformation of the mean of these estimates.

Specifying a correlation between the different random effects within a grouping-variable is

the default in most mixed-effect model implementations, for a good reason, because omitting

the correlation can lead to higher type I error rates (Matuschek *et al.* 2017). However, we

disagree with the view of Matuschek *et al.* 2017 that trading a small increase in type I error

rate for higher power is favorable, but it could still be an interesting solution with the often-

small number of observations in ecological studies when the increase in power prevails the

increase in type I error rate.

As fixed-effect alternatives, we tested the correctly specified model with the main effects of

temperature and mountain and their interaction (Table 1, Eq. 8), and the too simple model

dropping mountain as predictor (Table 1, Eq. 7). We tested the last model since mixed-

effects models, with standard deviation estimates of zero for both random effects,

correspond to fixed-effects models omitting the grouping variable.

Model fitting

We fitted mixed-effects models to our simulated data with two of the most commonly used

packages in R: *lme4* (Bates *et al.* 2015) and *glmmTMB* (Brooks *et al.* 2017). We present

here *lme4* results because it reports singular fits, i.e., when some of the parameters of the

variance-covariance Cholesky decomposition are exactly zero. Results with *glmmTMB* can

be found in Supporting Information S1. To obtain p-values for mixed-effects models we

used the *lmerTest* (Kuznetsova *et al.* 2017) package. For fixed-effects models in scenario

A we extracted p-values from the summary and for scenario B we used the fitted variance-

covariance matrix and the individual level effects to bootstrap the population effect and its

standard error (see Supporting Information S1).

For LMMs we used the restricted maximum likelihood estimator (REML). For GLMMs we

used the maximum likelihood estimator (MLE) as REML is not supported in *lme4* for GLMMs

(for a comparison of REML and MLE see Supporting Information S1). All results of mixed-

effect models presented in scenario A and B are for the datasets without singular fits (see

section *Variances of random effects and singular fits*).

Statistical properties and simulation setup

We compared the modeling options for both data generating scenarios based on important

statistical properties of the temperature effect (ecological effect) in our simulated data: type

I error rate and statistical power. Type I error rate is the probability to identify a temperature

effect as statistically significant although the effect is zero. Statistical power is the probability

to detect the temperature effect as significant if the effect is truly greater than zero. For a

correctly calibrated statistical test, the type I error is expected to be equal to the alpha-level

(in our case 5%), and the higher the statistical power the better the test. We additionally

calculated coverage for part of the simulations, i.e., how often the true effect of temperature

falls into the 95% confidence interval of the model, see Supporting Information S1 for results.

To investigate type I error rates for the slope/intercept, we simulated data with no

slope/intercept effect, i.e., the effect of temperature/mountain on plant reproduction (average

reproductive success of 0.4) and on height (average 0.4 cm) is zero. To additionally

investigate statistical power and coverage, we simulated an example with a weak effect, i.e.,

an effect that is barely statistically significant. In our example, a weak effect corresponds to

an average increase in size per unit step of the standardized temperature (linear scale) of

0.4 cm (Hypothesis 2) and 0.4 gain in reproductive success (Hypothesis 1) at scale of the

linear predictors.

For scenario A and scenario B, the individual effects for each mountain were drawn from a

normal distribution with standard deviation of 0.1 around the average effects: 0.4 cm average

height (intercept), and 0.4 cm average increase in size or 0.4 (logit link scale) gain in

reproductive success with temperature (slope).

Variances of random effects and singular fits

To understand how the number of levels affected the random effect variance estimates, we

recorded and compared them for random intercepts and slopes from the correctly specified

mixed-effects model in scenario B (Table 1, Eq. 10). We also compared optimization routines

(REML and MLE) in terms of estimating zero variances (singular fits, see below) for both

LMMs and GLMMs (see Supporting Information S1). For bounded operations, which most

R packages apply for the variance, one would expect a null distribution to be a combination

of a point mass at zero and a chi-squared distribution (Stram & Lee 1994). For the sampling

distribution with a true variance unequal to zero there are no proofs, but one would expect

a similar distribution.

Handling of singular fits

Technically, singular fits occur when at least one of the variances (diagonal elements) in the

Cholesky decomposition of the variance-covariance matrix are exactly zero or correlations

between different random effects are estimated as -1 or 1. However, how to deal with

singular fits is being discussed. While Matuschek *et al.* 2017 suggests to think *a priori* about

using simpler models, Barr *et al.* 2013 states to start with the maximal model and simplify

the model in case of convergence issues and singular fits. While singular fits don't signal a

convergence issue, the common sense is that they should be prevented. When facing a

singular fit, a practitioner usually changes the model structure and neglects the previous fit.

Following this perspective, we decided to use only non-singular fits for calculating the above-

mentioned statistical properties for the mixed-effect models. For fixed-effects models, we

used estimates from non-singular and singular fits combined.

But using only non-singular fits for calculating the statistical properties impacts the statistical

properties (e.g., type I error) because they are conditional on this selection and thus likely

not to be at the nominal level (e.g., 5% for type I error rate). We calculated type I error rates

for non-singular and singular fits combined which can be found in the Supporting Information

S1 (Fig. S8, S10). For scenario B (random intercepts and random slopes) *lme4* gives only a

generic singular fit warning but does not specify for which random effect the singular fit

occurred (random intercept or random slope), which furthermore complicates the handling

of singular fits (random intercept singular fit should be no problem for the population slope

effect). But as our main intention is to report the type I error rates from the point of the

practitioners who may adjust the model structure to dispose of the singular fit, our reported

rates represent empirical type I error rates.

Quantifying the influences of study design on power and type I error

The statistical properties of the ecological effect may depend not only on the number of

levels (mountains) but also on the random effect' variance, the overall number of

observations and the balance of observations among levels. To further quantify their impacts

on the statistical properties of the ecological effect, we additionally ran 1,000 iterations (each

with 1,000 non-singular model fits) with the data generating model from scenario B for

hypothesis 2 (height of plants, uncorrelated random slope and random intercept). Thereby,

we sampled uniformly the number of mountains from 2 to 20, the random effects' variance

from 10^{-4} to 4, the overall number of observations from 10*number of mountains to

500*number of mountains. Additionally, to create different degrees of unbalancedness in

data, we sampled for each mountain the average share of total observations from 0.1 to 0.9.

Thereby, we ensured that there are at least 3 observations per mountain. We used the

difference between the largest and the lowest proportion as proxy for the degree of

unbalancedness.

We fitted the correctly specified LMMs and LMs from scenario B (Table 1, Eq 8) and

calculated type I error rate and statistical power of the temperature effect. We then fitted a

quantile regression using the *qgam* R-package (Fasiolo *et al.* 2020), with the statistical

property (power and type I error rate) as response and variance, number of levels, total

number of observations and the unbalancedness proxy as splines.

Results

Scenario A - random intercepts per mountain

When the ecological effect was the same among mountains, irrespectively of the number of

levels (mountains), all models except for the too complex model (random intercept and

slope) showed a type I error rate of 5% (Fig. 1a, c). Power increased (Fig. 1b) with the

number of mountains for H2 from 90% (2 mountains) to 100% (5 to 8 mountains), and for

H1 (Fig. 1d) from 35% (2 mountains) to 95% (8 mountains). Note that the model omitting the

grouping variable presented similar properties as the other models. However, when

increasing the standard deviation of the random intercept in the simulation, this model

showed much lower power (Fig. S12).

For the too complex model for LMMs, we found on average a lower type I error rate of around

1-2% (Fig. 1a), and lower statistical power to detect the temperature effect for a small

number of mountains (Fig. 1b). Binomial datasets with small number of observations per

mountain (25, 50, 100) presented similar results regarding type I error but as expected, very

low power among all models (Figure S14). The results for the intercept for the different

models (see Fig. S9) are similar to the results for the slope in scenario B (see below).

Figure 1: Type I error rates and power for linear fixed and mixed effect models (a, b) and generalized linear fixed and

mixed-effects models (c, d), fitted to simulated data with 2-8 mountains (random intercept for each mountain - Scenario A)

and 50 observations per mountain. For each scenario, 5,000 simulations and models were tested. Results for mixed-effects

models are only from datasets in which mixed-effects models converged without presenting singular fit problems, while

results for fixed-effects model are from all datasets.

Scenario B - random intercepts and slopes per mountain

In scenario B, where the ecological effect differed among levels, the modeling decision had

a more severe influence on the statistical properties (Fig. 2). We found that type I error rate

of the correctly specified mixed-effects model (Table 1, Eq. 10) slightly increased (Fig. 2a,

c) with the number of levels towards the nominal value (0.05) (Fig. 2a, c). The too complex

model with correlated random intercept and random slope (Table 1, Eq. 11) presented

similar properties, but with slightly increased type I error and decreased power (Fig. 2). For

the correctly specified fixed-effect model, type I error ($\approx 2\%$) stayed constant with the number

of levels (Fig. 2a, c) and power increased with the number of mountains. For H1 (normal

model), however, the mixed-effects model showed higher power than the fixed-effects model

irrespective of the number of mountains (Fig. 2 d). For both normal and binomial models,

the too simple model omitting the grouping variable resulted in a higher type I error rate and

higher power than the other models (Fig. 2). The results for the intercept are similar as the

results for the slope (see Fig. S11).

324

Number of mountains

Figure 2: Type I error rates and power for linear (mixed effect) models (a, b) and generalized linear (mixed-effects) models (c, d), fitted to simulated data with 2-8 mountains for scenario B (random intercept and random slope for each mountain range). For each scenario, 5,000 simulations and models were tested. Results for mixed-effects models are only from datasets in which mixed-effects models converged without presenting singular fit problems, while results for fixed-effects model are from all datasets.

Variance estimates of random effects and singular fits

We found for LMMs (singular and non-singular fit results combined) in Scenario B (random intercept and slope) that random effects' variance estimates of the correctly specified model

(Table 1, Eq. 10) showed bimodal distributions with a chi-squared distribution around the

correct value (0.01) and a point mass at zero (Fig. 3a, b median is near to zero). The point

mass at zero decreased in height with increasing levels, i.e., less models estimated a

variance of zero with an increasing number of mountains (Fig. 3a, b, see also Table S1).

There was smaller bias for the random intercept variance estimates than for the random

slope variance estimates, which were still biased for eight levels. When looking at models

without singular fits, the variance estimates were chi-square distributed (Fig. 3c, d). The bias

towards higher values was stronger compared to estimates with singular fits, especially for

the random slope estimates (Fig. 3d). Results for *glmmTMB* were consistent with *lme4* when

using a threshold of 10^{-3} to classify a variance as zero (singular fit) (Fig. S2).

By comparing the fitting algorithms, we found that using MLE led to more zero-variance

estimates, i.e., singular fits, (Fig. S3, S4) than REML. Additionally, using MLE, also non-

singular variance estimates were strongly biased (Fig. S3, S4), but the bias decreases with

increasing number of levels. For both optimization routines, increasing the number of levels

reduced the number of singular fits (Table S1).

Figure 3: Variance estimates of random intercepts (a, c) and random slopes (b, d) for linear mixed-effects models (LMM, Table 1. eq. 10) in Scenario B, fitted with *lme4* using REML to simulated data with 2-8 mountains. Figures (a) and (b) show the results for all models (singular and non-singular fits) and figures (c) and (d) show the results for only non-singular fits. For each scenario, 5.000 simulations and models were tested. The blue dotted lines represent the true variance used in the simulation (0.01) and the red lines the average variance estimates.

We found that singular fits had a high influence on type I error and power (Fig. 4) of mixed-

and fixed-effects models. For singular fits, the type I error rate of the correctly specified

mixed-effects model was constant around 10% (similar to the model omitting the grouping

variable), while with non-singular fits it was 1% for two levels and increased towards 3% with

eight levels (Fig. 4a). In comparison, the fixed-effects model had similar type I error rates

singular and non-singular fits, both increasing from 0% (two levels) towards 1% (eight levels)

(Fig. 4c).

We also found differences in power for the models between singular and non-singular fits

(Fig. 4b, d). The power of the mixed-effects model with correct structure is higher for singular

than non-singular fits especially for a low number of mountains (Fig. 4b). For the fixed-effects

model, this difference is less strong, however the model has still higher power for singular

than for non-singular fits (Fig. 4d).

Figure 4: Type I error rate and power of the correctly specified linear fixed and mixed-effects models in scenario B. We separated the datasets based on if when fitted with lme4 they presented a singular fit (red lines) or non-singular fit (blue lines). Figure (a) and (b) are results for the linear mixed-effects models, and (c) and (d) for the linear fixed-effects models. For comparisons, we show also results for the fixed-effects model that omits the grouping variable (mountain).

Quantifying the influences of study design on power and type I error

We found that the average type I error of mixed-effects models is slightly nearer to the nominal value its fixed-effects counterpart (Fig. 5a). Additionally, we found that the number

of levels most strongly influences the type I error rate for mixed- as well as fixed-effects

model (Fig. 5c). With five or more levels, however, the influence is almost zero. Differences

between the mixed- and fixed-effects models arose for the variance and the total number of

observations. Here, the mixed-effects model was less influenced by a small random effects'

variance and a low number of total observations than the fixed-effects model (Fig. 5 b, d).

Balance did not influence both models (Fig. 5e).

For power we found no difference between fixed- and mixed-effects model (Fig. 5 f-j). For

both models, an increase in variance decreased the power, while increasing the number of

levels increased the power (Fig. 5 g, i). The total number of observations and the balance

between groups had less influence (Fig. 5 h, j).

Figure 5: Comparing the influence of study design factors on the type I error rate (b - e) and power (g - j) of linear mixed- (blue lines) and fixed-effects models (red lines) with their respective average values (a, f). We found that the variance of

395 the random effects and the number of levels (*number of mountains*) are the most important values to get correct statistical

396 properties and improve the chance to find a significant effect. For this analysis, we used the plant height example

397 (Hypothesis 2, Box 1, normal distribution models) for Scenario B (random intercept and random slope). Results for mixed-

398 effects models are only from datasets in which mixed-effects models converged without presenting singular fit problems,

399 while results for fixed-effects model are from all datasets.

Discussion

Ecological data collections or experiments often produce data with grouping/blocking

structures and mixed-effects models can account for these dependencies. When such data,

however, has a small number of levels, should the analysists stick to the mixed-effects model

or fall back to a fixed-effects model? How does this decision influence the statistical

properties of the population effect? Here, we showed with simulations that mixed-effects

models with small number of levels in the grouping variable are technically more robust than

previously assumed (Fig. 2) and that the decision between random and fixed effect matters

most when the ecological effect differs among levels (Fig. 2).

When the ecological effect of interest is the same for each level of the grouping variable

(scenario A, varying intercept, random intercept model), almost all models independent of

the number of levels presented the same statistical properties for the ecological effect

(temperature, slope) (Fig. 1, see also Gomes 2021). The only exception was the too complex

model that presented too low type I errors (close to zero, Fig. 1) and lower power. We

speculate this is caused by the higher number of parameters that need to be estimated by

the model and that the model was unable to correctly estimate the random slopes to zero.

Notably, for scenario A, the too simple model omitting the grouping variable presented

correct statistical properties (Fig. 1). However, this is illusive because power decreased

strongly with increasing variance in the random effect (Fig. S12) confirming the importance

of including grouping variables to correctly partition the variance among the different

predictors to get well calibrated statistical properties (Gelman 2005; Gelman & Hill 2007;

Bell *et al.* 2019). Also including the grouping variable is mandatory if one is interested in the

average intercept, otherwise it would cause inflated type I error rates (see Fig. S9, S11; see

the following section).

When the ecological effect of interest differs for each level of the grouping variable (scenario

B; varying intercept and slope; random intercept and random slope model), the model choice

influenced the statistical properties more strongly. The mixed-effects model had a better

type I error than the fixed-effects model, especially for a larger number of mountains (Fig.

2). Power was comparable for LMMs, but for GLMMs the mixed-effects model had higher

power which may be linked to the distributional assumption of the random effects (see

Gelman & Hill 2007): the regularization of the random effects, which stems from their

distributional assumption, reduces the "effective" number of parameters as individual

estimates are constrained to the mean. In summary, mixed-effects models have better type

I error rates and higher power than fixed-effects models for linear and generalized models.

The different patterns between the linear and generalized models are mainly due to our

bootstrapping approach to calculate the average effect and its standard error. For

generalized models, the bootstrapping approach showed on average a slightly deflationary

distribution of p-values (Fig. 18) leading to more conservative type I error rates. However,

the properties of alternatives such as the zero-sum contrasts were even worse for different

simulation scenarios (Fig. 18, Fig. S20).

Too complex mixed-effects models presented in both scenarios slightly lower type I error

and power compared to the correctly parameterized mixed-effects model (Fig. 1, 2). This

trade-off between type I error and power is in line with Matuschek *et al.* (2017) for different

model complexities. Overall, the more complex (higher parametrized) models are more

conservative but have less power than the simplified models. We think these more

conservative estimates are preferable over anti-conservative estimates because

researchers unfortunately tend to try a variety of analyses and only report significant ones

(Simmons *et al.* 2011) and more conservative type I error rates counteract this procedure.

However, dropping the correlation structure between random effects should be still carefully

considered. Correlations between the random effects are very common in empirical data

and the true change in type I error rate might be higher than shown here because we

assumed no correlation in our simulations. Group-mean centering of the ecological effect

may mitigate the requirement of assuming a correlation, but it also changes the interpretation

of the model.

In scenario B, too simple models exhibited inflated type I errors (in line with Schielzeth &

Forstmeier 2009; Barr *et al.* 2013; Bell *et al.* 2019), but very high power (Fig. 2). We

speculate that additional variance coming from the difference between levels in the grouping

variable, which is not accounted, is attributed to the ecological effect and causes

overconfident estimates.

Variances of random effects and singular fits

Singular fits occurred more often for small number of levels, creating a point mass at zero

in the distribution of the variance estimates (Fig. 3, Table S1), because a singular fit in our

simulations corresponded to a zero-variance estimate as we excluded the other possibility

(we set the correlations between random slope and intercept to zero, see *Methods*). The

resulting distribution consisted of a towards higher values biased chi-squared distribution

and the point mass at zero as expected (see Stram & Lee 1994). For non-singular and

singular fits combined, the variance estimates of the random effects were biased and

imprecise with a small number of levels, but the bias decreased with the number of levels

towards zero (McNeish 2017). Removing the singular fits led to even more bias in the

variance estimates (Fig. 3c, d).

The biased variance estimates are caused by ensuring positive variances in the optimization

routines (Bates *et al.* 2015; Brooks *et al.* 2017). In case of a singular fit, the correctly

specified mixed-effects model had similar statistical properties as a fixed-effect model

dropping the grouping variable (Fig. 4): no difference between the levels, which corresponds

to a fixed-effect model without the grouping variable. However, the models still differed in

their number of parameters (and degrees of freedom) which might explain the slight

differences in statistical properties (Fig. 4).When switching to fixed-effects models for

singular fits in the random effect, the type I error rate and power were similar to the random

effect model with non-singular fits (Fig. 4).

Connection to study design

Earlier studies reported mixed recommendations about important study design factors.

While some studies only stressed the importance of the total number of observations (Martin

et al. 2011; Pol 2012), we found, in accordance with Aarts *et al. (*2014), that the number of

levels and the variance between levels have a strong influence on type I error rates and

power. Due to our simulation design, which automatically increases the number of

observations when increasing the number of levels, we however, cannot perfectly separate

the effects of number of observations and levels from each other.

The influence of the variance on the statistical properties is mixed. On the one hand,

increasing the variance had a positive effect on the type I error for both models but the fixed-

effects model was more strongly affected (Fig. 5). The different distributional assumptions

might explain this different behavior: the mixed-effects model assumes the levels to be

normally distributed and estimates flexibly the variance of the levels, whereas the fixed-

effects model makes implicitly the assumption of an infinite variance. On the other hand,

increasing the variance over a certain value (Fig. 5g) decreased the power of both models

because more variance is explained by the difference between levels, and this increases

the uncertainty of the slope effect estimate.

Given the strong influence of the number of mountains on type I error rates, we encourage

to design a study with at least 8 levels because with more than 8 levels, the type I error rate

was approximately not affected by the number of mountains (Fig. 5c). In our scenarios, the

influence of the unbalanced number of observations between levels was small (Fig. 5)

suggesting that unbalanced designs are not the main concern for mixed-effects models

given their robustness to unbalanced data (Swallow & Monahan 1984; Pinheiro & Bates

1995; Schielzeth et al. 2020) as long as one is not interested in treatment but population

effects. Moreover, the impact of study design on type I error and power stresses the

importance of pre-experiments and power analyses (e.g. Johnson *et al.* 2015; Green &

MacLeod 2016; Brysbaert & Stevens 2018) to maximize the meaningfulness and efficiency

of a study.

Practical suggestion

Before giving practical advice, we must recall the exact situation in which this manuscript

acts. We assume that a researcher is interested in a population effect, and they have already

decided to use a mixed-effects model (broad-sense analysis, not interested in the individual

levels effects), but now faces a low number of levels. Recommendations only apply to such

situations.

We found no harsh threshold for the minimum number of levels in a grouping variable

necessary to model it as a random effect (Fig. 2), but rather found that a singular fit in the

mixed-effects model indicates switching to the fixed-effects model is beneficial (more

conservative type I error rates, Fig. 2). Acknowledging that most singular fits occur with a

small number of levels (Table S1), this might also explain the common rule of thumb to do

not fit a grouping variable as random effect if it has fewer than 5 levels (Gelman & Hill 2007;

Bolker et al. 2009; Bolker 2015). Apart from the results of our study, checking the data for

possible issues causing zero variance estimates is desired. Additionally, one should

investigate whether a zero-variance estimate is ecologically reasonable.

We therefore recommend starting with the mixed-effects model, regardless of the number

of levels, and switching to a fixed-effects model only in case of a singular fit (more

conservative type I error rate). When the ecological effect differs among levels, we

recommend starting with correlated random slope and intercept (following Barr et al. 2013).

When obtaining a singular fit, switch to uncorrelated random effects (following Matuschek *et*

al. 2017) and in case of another singular fit, switch to a fixed-effects model. These

recommendations are summarized in Fig. 6.

Our previous recommendations assumed that we know whether the ecological effect differs

across levels. With empirical data, however, this is usually unknown. Here, a too complex

model leads to too low type I error rates and less power, while a too simple model has

inflated type I error rates. Thus, to prevent being overly confident, it is essential to include

random slopes if they are present (Schielzeth & Forstmeier 2009). But because of the power

issue, we doubt that its generally advisable to always start with the maximally complex

random effect structure (as suggested in Bell *et al.* 2019), and it requires a decision criterion

for the complexity of the random effect structure.

Figure 6: Consequences and recommendations for mixed-effects models with a small number of levels in the random

effect. When the ecological effect does not differ between different levels of the grouping variable (left side) all modeling

options, which include the grouping variable, lead to the same results and thus only a singular fit requires a change to a

fixed-effects model. If the ecological effect differs among levels (middle to right side), starting with the mixed-effects model

and only changing to the fixed-effects model in case of a singular fit is recommended.

Unfortunately, model selection on random effects is complicated, because the exact degrees

of freedom, that are used by a random effect, are unclear (Kuznetsova *et al.* 2017).

Approximations exist and for low number of levels a Kenward-Rogers correction is preferred

(McNeish 2017). Although this prevents a naïve use of AIC or likelihood ratio tests (LRT),

other methods such as simulated (restricted) LRTs (Wiencierz *et al.* 2011) can be used to

decide if adding a random slope is justified. Thus, we recommend starting with the simpler

structure (typically random intercepts) and then use residual checks (e.g. Hartig 2019) or

appropriate model selection criteria (e.g. Matuschek *et al.* 2017) to decide if a random slope

should be added.

Another, somewhat perpendicular option is to modify the distributional assumption about the

random effects (normal distribution), which is mainly due to mathematical convenience

(Beck & Katz 2007) and corresponds to a L2 (RIDGE) regularization. A L2 regularization

biases random effect estimates towards zero but is unable to produce exactly zero estimates

(Zou & Hastie 2005). In contrast, L1 (LASSO) regularization, which corresponds to a Laplace

distribution (Park & Casella 2008; Hans 2009; Tibshirani 2011), is able to estimate zero

random effect estimates, when there is no effect. Because mixed-effects models are

relatively robust to distributional assumptions (Bell *et al.* 2019; Schielzeth *et al.* 2020),

switching distributions to LASSO or a combination of L1 and L2 regularization (elastic net,

see Zou & Hastie 2005) could potentially allow starting with the maximal complex random

effect structure (see also McNeish & Bauer 2020). Such options should be investigated in

further studies.

Conclusion

In conclusion, we showed that mixed-effects models are more robust than previously thought, despite the biased variance estimates for low number of levels in the grouping variable. We found that the statistical properties of the population effect are robust against the model choice when the ecological effect is the same among the levels of the grouping variable, however, the model matters when the ecological effect differs among levels. When in doubt about the data-generating process, we encourage starting with a simplified model (random intercept only) and consult model diagnostics and simulated LRTs to check for evidence of random slope effects. When finding evidence for random slopes in these tests, we recommend starting with the mixed-effects model and switching only to a fixed-effects model in case of a singular fit problem. With this work, we provide a practical guideline, which helps ecologists in the study design, the data analysis and thus making ecological inference more informative and robust.

Data availability statement:

No empirical data was used in this study. Code to run and analyze the experiments can be

found at https://github.com/JohannesOberpriller/RandomEffect_Groups.

Author Statement

Acknowledgement

The idea of the manuscript originated from a discussion in the Theoretical Ecology seminar

and was further developed in the Coding Club at University of Regensburg. We thank Rainer

Spang, Carsten Dormann, Magdalena Mair, Björn Reineking, Sean McMahon, Andreas

Ettner and Florian Hartig for comments and discussions on earlier versions of the

manuscript. We also thank two anonymous reviewers for their valuable comments and

suggestions. JO was funded by the Bavarian Ministry of Science and the Arts in the context

of Bavarian Climate Research Network (bayklif). MSL was funded by the Smithsonian

Predoctoral Fellowship.

References

Aarts, E., Verhage, M., Veenvliet, J.V., Dolan, C.V. & van der Sluis, S. (2014). A solution to dependency: using multilevel analysis to accommodate nested data. *Nature Neuroscience*, 17, 491–496.

Arnqvist, G. (2020). Mixed Models Offer No Freedom from Degrees of Freedom. *Trends in Ecology & Evolution*, 35, 329–335.

Baayen, R.H., Davidson, D.J. & Bates, D.M. (2008). Mixed-effects modeling with crossed random effects for subjects and items. *Journal of Memory and Language*, Special Issue: Emerging Data Analysis, 59, 390–412.

Barr, D.J., Levy, R., Scheepers, C. & Tily, H.J. (2013). Random effects structure for confirmatory hypothesis testing: Keep it maximal. *Journal of Memory and Language*, 68, 255–278.

Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software, Articles*, 67, 1–48.

Beck, N. & Katz, J.N. (2007). Random Coefficient Models for Time-Series—Cross-Section Data: Monte Carlo Experiments. *Political Analysis*, 15, 182–195.

Bell, A., Fairbrother, M. & Jones, K. (2019). Fixed and random effects models: making an informed choice. *Quality & Quantity*, 53, 1051–1074.

Boisgontier, M.P. & Cheval, B. (2016). The anova to mixed model transition. *Neuroscience & Biobehavioral Reviews*, 68, 1004–1005.

Bolker, B.M. (2015). Linear and generalized linear mixed models. *Ecological Statistics: Contemporary theory and application*, 309–333.

Bolker, B.M., Brooks, M.E., Clark, C.J., Geange, S.W., Poulsen, J.R., Stevens, M.H.H., *et al.* (2009). Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in ecology & evolution*, 24, 127–35.

Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A., *et al.* (2017). glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *The R journal*, 9, 378–400.

Brysbaert, M. & Stevens, M. (2018). Power Analysis and Effect Size in Mixed Effects Models: A Tutorial. *Journal of Cognition*, 1.

Chen, Z. & Dunson, D.B. (2003). Random Effects Selection in Linear Mixed Models. *Biometrics*, 59, 762–769.

628 DerSimonian, R. & Laird, N. (1986). Meta-Analysis in Clinical Trials. *Controlled Clinical*
629    *Trials*, 7, 177–188.

630 Dixon, P. (2016). Should blocks be fixed or random? *2016 Conference on Applied*
631    *Statistics in Agriculture Proceedings*, 23–39.

632 Fasiolo, M., Wood, S.N., Zaffran, M., Nedellec, R. & Goude, Y. (2020). qgam: Bayesian
633    non-parametric quantile regression modelling in R. *arXiv preprint arXiv:2007.03303*.

634 Gelman, A. (2005). Analysis of variance—why it is more important than ever. *The Annals*
635    *of Statistics*, 33, 1–53.

636 Gelman, A. & Hill, J. (2007). *Data analysis using regression and multilevel/hierarchical*
637    *models*. Cambridge university press.

638 Giesselmann, M. & Schmidt-Catran, A.W. (2020). Interactions in Fixed Effects Regression
639    Models. *Sociological Methods & Research*, 0049124120914934.

640 Gomes, D.G.E. (2021). Including random effects in statistical models in ecology: fewer
641    than five levels? *bioRxiv*, 2021.04.11.439357.

642 Green, P. & MacLeod, C.J. (2016). SIMR: an R package for power analysis of generalized
643    linear mixed models by simulation. *Methods in Ecology and Evolution*, 7, 493–498.

644 Gunasekara, F.I., Richardson, K., Carter, K. & Blakely, T. (2014). Fixed effects analysis of
645    repeated measures data. *International Journal of Epidemiology*, 43, 264–269.

646 Hans, C. (2009). Bayesian lasso regression. *Biometrika*, 96, 835–845.

647 Harrison, X.A. (2014). Using observation-level random effects to model overdispersion in
648    count data in ecology and evolution. *PeerJ*, 2, e616.

649 Harrison, X.A. (2015). A comparison of observation-level random effect and Beta-Binomial
650    models for modelling overdispersion in Binomial data in ecology & evolution. *PeerJ*,
651    3, e1114.

652 Harrison, X.A., Donaldson, L., Correa-Cano, M.E., Evans, J., Fisher, D.N., Goodwin, C., *et*
653    *al.* (2017). *Best practice in mixed effects modelling and multi-model inference in*
654    *ecology* (No. e3113v1). PeerJ Inc.

655 Harrison, X.A., Donaldson, L., Correa-Cano, M.E., Evans, J., Fisher, D.N., Goodwin,
656    C.E.D., *et al.* (2018). A brief introduction to mixed effects modelling and multi-model
657    inference in ecology. *PeerJ*, 6, e4794.

658 Hartig, F. (2019). DHARMa: residual diagnostics for hierarchical (multi-level/mixed)
659    regression models. *R package version 0.2,* 4.

Hedges, L.V. & Vevea, J.L. (1998). Fixed- and random-effects models in meta-analysis.
Psychological Methods, 3, 486–504.

Hox, J.J., Moerbeek, M. & Van de Schoot, R. (2017). *Multilevel analysis: Techniques and*
applications. Routledge.

Johnson, P.C.D., Barry, S.J.E., Ferguson, H.M. & Müller, P. (2015). Power analysis for
generalized linear mixed models in ecology and evolution. *Methods in Ecology and*
Evolution, 6, 133–142.

Kadane, J.B. (2020). *Principles of uncertainty*. Chapman and Hall/CRC.

Krueger, C. & Tian, L. (2004). A Comparison of the General Linear Mixed Model and
Repeated Measures ANOVA Using a Dataset with Multiple Missing Data Points.
Biological Research For Nursing, 6, 151–157.

Kuznetsova, A., Brockhoff, P. & Christensen, R. (2017). lmerTest Package: Tests in Linear
Mixed Effects Models. *Journal of Statistical Software, Articles*, 82.

Laird, N.M. & Ware, J.H. (1982). Random-Effects Models for Longitudinal Data.
Biometrics, 38, 963–974.

Lenth, R.V. (2021). *emmeans: Estimated Marginal Means, aka Least-Squares Means*.

Lindstrom, M.J. & Bates, D.M. (1988). Newton—Raphson and EM Algorithms for Linear
Mixed-Effects Models for Repeated-Measures Data. *Journal of the American*
Statistical Association, 83, 1014–1022.

Littell, R.C. (2002). Analysis of unbalanced mixed model data: A case study comparison of
ANOVA versus REML/GLS. *JABES*, 7, 472–490.

Maas, C.J.M. & Hox, J.J. (2005). Sufficient Sample Sizes for Multilevel Modeling.
Methodology, 1, 86–92.

Martin, J.G.A., Nussey, D.H., Wilson, A.J. & Réale, D. (2011). Measuring individual
differences in reaction norms in field and experimental studies: a power analysis of
random regression models. *Methods in Ecology and Evolution*, 2, 362–374.

Matuschek, H., Kliegl, R., Vasishth, S., Baayen, H. & Bates, D. (2017). Balancing Type I
error and power in linear mixed models. *Journal of Memory and Language*, 94,
305–315.

McLean, R.A., Sanders, W.L. & Stroup, W.W. (1991). A Unified Approach to Mixed Linear
Models. *The American Statistician*, 45, 54–64.

McMahon, S.M. & Diez, J.M. (2007). Scales of association: hierarchical linear models and
the measurement of ecological systems. *Ecology Letters*, 10, 437–452.

McNeish, D. (2017). Small Sample Methods for Multilevel Modeling: A Colloquial
Elucidation of REML and the Kenward-Roger Correction. *Multivariate Behavioral*
Research, 52, 661–670.

McNeish, D. & Bauer, D.J. (2020). Reducing Incidence of Nonpositive Definite Covariance
Matrices in Mixed Effect Models. *Multivariate Behavioral Research*, 0, 1–23.

McNeish, D.M. & Stapleton, L.M. (2016). The Effect of Small Sample Size on Two-Level
Model Estimates: A Review and Illustration. *Educ Psychol Rev*, 28, 295–314.

Meteyard, L. & Davies, R.A.I. (2020). Best practice guidance for linear mixed-effects
models in psychological science. *Journal of Memory and Language*, 112, 104092.

Millar, R.B. & Anderson, M.J. (2004). Remedies for pseudoreplication. *Fisheries Research*,
70, 397–407.

Nakagawa, S. & Schielzeth, H. (2013). A general and simple method for obtaining R2 from
generalized linear mixed-effects models. *Methods in Ecology and Evolution*, 4, 133–
142.

Park, T. & Casella, G. (2008). The Bayesian Lasso. *null*, 103, 681–686.

Pinheiro, J.C. & Bates, D.M. (1995). Approximations to the Log-Likelihood Function in the
Nonlinear Mixed-Effects Model. *Journal of Computational and Graphical Statistics*,
4, 12–35.

Pol, M. van de. (2012). Quantifying individual variation in reaction norms: how study
design affects the accuracy, precision and power of random regression models.
Methods in Ecology and Evolution, 3, 268–280.

Schelldorfer, J., Bühlmann, P. & Geer, S.V.D. (2011). Estimation for High-Dimensional
Linear Mixed-Effects Models Using ℓ1-Penalization. *Scandinavian Journal of*
Statistics, 38, 197–214.

Schielzeth, H. (2010). Simple means to improve the interpretability of regression
coefficients. *Methods in Ecology and Evolution*, 1, 103–113.

Schielzeth, H., Dingemanse, N.J., Nakagawa, S., Westneat, D.F., Allegue, H., Teplitsky,
C., *et al.* (2020). Robustness of linear mixed-effects models to violations of
distributional assumptions. *Methods in Ecology and Evolution*, 11.

Schielzeth, H. & Forstmeier, W. (2009). Conclusions beyond support: overconfident
estimates in mixed models. *Behavioral Ecology*, 20, 416–420.

Shaver, J.M. (2019). Interpreting Interactions in Linear Fixed-Effect Regression Models:
When Fixed-Effect Estimates Are No Longer Within-Effects. *Strategy Science*, 4,
25–40.

Simmons, J.P., Nelson, L.D. & Simonsohn, U. (2011). False-Positive Psychology:
Undisclosed Flexibility in Data Collection and Analysis Allows Presenting Anything
as Significant. *Psychol Sci*, 22, 1359–1366.

Stram, D.O. & Lee, J.W. (1994). Variance Components Testing in the Longitudinal Mixed
Effects Model. *Biometrics*, 50, 1171–1177.

Swallow, W.H. & Monahan, J.F. (1984). Monte Carlo Comparison of ANOVA, MIVQUE,
REML, and ML Estimators of Variance Components. *Technometrics*, 26, 47–57.

Tibshirani, R. (2011). Regression shrinkage and selection via the lasso: a retrospective.
Journal of the Royal Statistical Society: Series B (Statistical Methodology), 73, 273–
282.

Townsend, Z., Buckley, J., Harada, M. & Scott, M. (2013). The Choice between Fixed and
Random Effects. In: *The SAGE Handbook of Multilevel Modeling*. SAGE
Publications Ltd, 1 Oliver's Yard, 55 City Road, London EC1Y 1SP United Kingdom,
pp. 73–88.

Wainwright, P.E., Leatherdale, S.T. & Dubin, J.A. (2007). Advantages of mixed effects
models over traditional ANOVA models in developmental studies: A worked
example in a mouse model of fetal alcohol syndrome. *Developmental
Psychobiology*, 49, 664–674.

Wiencierz, A., Greven, S. & Küchenhoff, H. (2011). Restricted likelihood ratio testing in
linear mixed models with general error covariance structure. *Electronic Journal of
Statistics*, 5, 1718–1734.

Zou, H. & Hastie, T. (2005). Regularization and variable selection via the elastic net.
Journal of the royal statistical society: series B (statistical methodology), 67, 301–
320.

Zuur, A., Ieno, E., Walker, N., Saveliev, A. & Smith, G. (2009). *Mixed effects models and
extensions in ecology with R*.
753