Biological complexity facilitates tuning of the neuronal parameter space

*Marius Schneider[a,b,c,d], Alexander D Bird[a,b,c] Albert Gidon[e], Jochen Triesch[a,d,f], Peter Jedlicka[a,c,g,1], *Hermann Cuntz[a,b,1]

[a]Frankfurt Institute for Advanced Studies, 60438 Frankfurt am Main, Germany
[b]Ernst Strüngmann Institute (ESI) for Neuroscience in cooperation with the Max Planck Society, 60528 Frankfurt am Main, Germany
[c]ICAR3R – Interdisciplinary Centre for 3Rs in Animal Research, Justus Liebig University Giessen, 35390 Giessen, Germany
[d]Faculty of Physics, Goethe University, Frankfurt/Main, 60438 Frankfurt am Main, Germany
[e]Institute for Biology, Humboldt-Universität zu Berlin, Berlin, Germany
[f]Faculty of Computer Science and Mathematics, Goethe University, 60438 Frankfurt am Main, Germany
[g]Institute of Clinical Neuroanatomy, Neuroscience Center, Goethe University, 60528 Frankfurt am Main, Germany

[1]Joint senior authors

*marius.schneider@esi-frankfurt.de, cuntz@fias.uni-frankfurt.de

Keywords

Dendrite function, Degeneracy, Electrophysiology, Compartmental modelling, Evolutionary principles, Law of large numbers

In brief

Studying ion channel diversity in neuronal models we show how robust biological systems may evolve not despite but because of their complexity.

Highlights

- 15 channel model of hippocampal granule cells (GCs) reduces to 5 ion channels without loss of spiking behaviour.

- But knocking out ion channels can be compensated only in the full model.

- Random sampling leads to $\sim 6\%$ solutions in full but only $\sim 1\%$ in reduced model.

- Law of large numbers generalises our observations to other complex biological systems.

Abstract

The electrical and computational properties of neurons in our brains are determined by a rich repertoire of membrane-spanning ion channels and elaborate dendritic trees. However, the precise reason for this inherent complexity remains unknown. Here, we generated large stochastic populations of biophysically realistic hippocampal granule cell models comparing those with all 15 ion channels to their reduced but functional counterparts containing only 5 ion channels. Strikingly, valid parameter combinations in the full models were more frequent and more stable in the face of perturbations to channel expression levels. Scaling up the numbers of ion channels artificially in the reduced models recovered these advantages confirming the key contribution of the actual number of ion channel types. We conclude that the diversity of ion channels gives a neuron greater flexibility and robustness to achieve target excitability.

Significance statement

Over the course of billions of years, evolution has led to a wide variety of biological systems. The emergence of the more complex among these seems surprising in the light of the high demands of searching for viable solutions in a correspondingly high-dimensional parameter space. In realistic neuron models with their inherently complex ion channel composition, we find a surprisingly large number of viable solutions when selecting parameters randomly. This effect is strongly reduced in models with fewer ion channel types but is recovered when inserting additional artificial ion channels. Because concepts from probability theory provide a plausible explanation for this improved distribution of valid model parameters, we propose that this may generalise to evolutionary selection in other complex biological systems.

Introduction

Throughout evolution, biological cells have emerged with increasing diversity and complexity. Optimising for multiple objectives while keeping an ever larger number of cell parameters within a viable range seems a daunting task for evolutionary processes; and it remains unclear how such a multi-objective optimisation can be achieved in the corresponding high dimensional parameter space. Here we explore the counter-intuitive hypothesis that increasing the number of mechanisms – i.e. increasing the biological complexity – potentially helps systems to evolve more quickly, easily and efficiently towards satisfying a large number of objectives.

Neurons are a good example of complex cells, typically exhibiting a great diversity in the expression of ion channels as products of such evolutionary optimisation. The channel parameters must be tuned to cooperatively generate multiple features of neuronal spiking behaviour. A palette of such spiking features has been successfully used in computational biophysical neuron models for multi-objective optimisation (MOO) using genetic algorithms (Druckmann, 2007). Mammalian neurons contain a large variety of ion channels types in their membrane (Coetzee *et al.*, 2006) producing a wide range of possible spiking mechanisms with varying temporal dynamics and excitability (Connors and Gutnick, 1990). Interestingly, a number of these ion channel variants exhibit overlapping functional properties (Coetzee *et al.*, 2006; Rudy, 1988; Herrera-Valdez *et al.*, 2013; Marder and Goaillard, 2006; Olypher and Calabrese, 2007; Hille, 2001; Goaillard and Marder, 2021). A large body of literature has explored the reason for this high diversity (Marder, 2011; Prinz *et al.*, 2004; Golowasch *et al.*, 2002; O'Leary *et al.*, 2013). However, it remains unclear what role exactly the diversity of ion channel types plays regarding evolution and its contribution to functional mechanisms that impact neuronal computations.

Neuronal computation relies on the morphology as well as on the diversity and distribution of ion channels in the membrane of the dendritic tree, the soma, and

the axon initial segment. Even small changes in the distribution of ion channels can   54
change the activity in neurons drastically (Achard and De Schutter, 2006). Large   55
differences in experimental measurements have been observed from cell to cell,   56
day to day and animal to animal in data from the same classes of cells (Marder and   57
Goaillard, 2006; Golowasch *et al.*, 2002; Golowasch and Marder, 1992; MacLean *et*   58
al., 2003; Swensen, 2005; Schulz *et al.*, 2006, 2007). The expression levels of these ion   59
channel types can vary several-fold across neurons of a defined type (Marder and   60
Goaillard, 2006; Prinz *et al.*, 2004; Golowasch *et al.*, 2002; Golowasch and Marder,   61
1992; MacLean *et al.*, 2003; Schulz *et al.*, 2006). However, many detailed biophysical   62
models of single cells ignore this variability in electrophysiological data and search   63
for a fixed set of parameters that replicates an average behaviour of a particular   64
cell type (Golowasch *et al.*, 2002).   65

How can neurons manage to achieve a functional target activity with such a wide   66
ion channel diversity? Using a spike-feature-based multi-objective approach, we   67
generated large population parameter sets of dentate granule cell (GC) models   68
with different numbers of ion channel types in order to investigate the potential   69
advantages of ion channel diversity. We then tested to which degree the different   70
models could compensate for pathological channel loss. Furthermore, we inves-   71
tigated differences in functional parameter sets, taking into account stochastic   72
fluctuations in channel-coding gene expression. Finally, we studied the stability of   73
the different models against ion channel alterations due to e.g. protein turnover.   74
We found that in all cases the complete GC model with all ion channel types was   75
more robust, stable and had more valid parameter combinations than its reduced   76
counterparts.   77

Figure 1. Simplified as well as realistic complex ion conductance-based models capture multiple spiking features of real granule cells (GCs)

A, (*Top*) 3D-reconstructed mouse GC morphology used for our simulations (Schmidt-Hieber *et al.*, 2007). (*Bottom*) Spike features used to calculate the multi-objective fitness of the GC model. **B**, Membrane potential during $200ms$ lasting current clamp of $90pA$. The coloured curves show the relative contribution of all implemented ion channels to the total inward (See next page)

Figure 1. (continued) and outward current at each time step (during the second and third spike) as a percentage of the total current. The black filled curves illustrate the total inward and outward currents on a logarithmic scale. This plot was inspired by Alonso and Marder (2019).**C**, Contribution of currents to total inward and outward current in reduced models and models that compensate for the knock out of the BK (*Left*) and Cav22 (*Right*) channel. Similar visualisation and current injection procedure as in **B**.

Results

We used a recently established multi-compartmental model comprising the 15 79
different voltage or calcium-dependent ion channels that were described in mouse 80
GCs (Beining *et al.*, 2017). The model was specifically designed to reproduce the 81
results not of a single experiment but of a large series of experiments and was 82
based on raw electrophysiology traces. Its parameters were fitted to reproduce 83
the experimental data for a number of different reconstructed (see example in **Fig-** 84
ure 1A, *Top*, from Schmidt-Hieber *et al.*, 2007) and synthetic neuronal morphologies 85
making the model robust within the GC morphological space. Furthermore, the 86
resulting model readily generalised to rat GCs as well as to adult born mouse GCs 87
(i.e. GCs from adult mouse neurogenesis) after incorporating the known changes in 88
morphology and ion channel composition. The model can therefore be considered 89
to be robust and comprehensive. This makes it an experimentally validated tool 90
to study the impact of complex ion channel compositions on robustness of the 91
spiking output. To this end, we employed a population (also called "ensemble" 92
or "database") modelling approach, which allowed us to explore the multidimen- 93
sional parameter space in large populations of stochastically generated models 94
(Prinz *et al.*, 2003; Gunay *et al.*, 2008; Britton *et al.*, 2013; Sekulic *et al.*, 2014; Rathour 95
and Narayanan, 2019; Jedlicka *et al.*, 2022). 96

The GC model cost function 97

First, we developed a cost function for an automated evaluation of the validity of 98
diverse models, which differed in their ion channel combinations and densities. 99
Since no quantitative data exists on the particular expression of the various ion 100
channels in individual GCs, some form of fitting procedure of channel densities 101
was required in the construction of the GC model. The model consists of 27 conduc- 102
tance parameters, which precludes a comprehensive grid scan for parameter fitting 103
due to the long computing time in a 27 dimensional parameter space. The model 104
has therefore previously been largely tuned manually with expert knowledge 105
from GC biology. To assess the quality of any individual set of parameters more 106
automatically, we designed a fitness function that quantified the distance to experi- 107
mental spiking data and was inspired by approaches used previously (Druckmann, 108
2007; Beining *et al.*, 2017, see Methods, **Figure S1**). A number of different methods 109
have been proposed to quantify the quality of a set of parameters in relation to 110
neuronal activity (Achard and De Schutter, 2006; Bahl *et al.*, 2012; Keren *et al.*, 2005; 111
Vanier and Bower, 1999). While most studies focus on reproducing an average 112
electrophysiological activity pattern, we wanted to focus on the distribution of 113
valid parameter combinations in the GC model taking into account the variability 114
present in experimental data. 115

We therefore used a multi-objective fitness function based on spike features, which 116
allowed us to search for optimal trade-offs between different firing properties 117
(Druckmann, 2007). We extracted 9 different spiking features from raw electrophys- 118
iology traces during a $200ms$ current clamp injection with 50 and $90pA$ at the soma 119
(**Figure 1A**, *Bottom*, see Methods). We then compared the values for these features 120
between the model and the experimental data. To generate a population of GC 121
model instances that reflected the full range of firing properties, we calculated the 122
deviation from the experimental mean in units of experimental standard deviation 123
(SD) (Druckmann, 2007). In order to become a valid parameter combination in the 124
GC model, the error value was required to be less than two SDs away from the 125

experimental average of each feature.   126

A manual search for parameter sets fulfilling this requirement was very time-   127
consuming and could never be exhaustive. There are various automated parameter   128
search methods, such as gradient descent methods, genetic algorithms, simulated   129
annealing, and stochastic search methods, which make the search for parameters   130
more efficient (Vanier and Bower, 1999; Mitchell, 1998; Kirkpatrick *et al.*, 1983;   131
Press *et al.*, 2007). Since we were starting from a valid parameter combination,   132
we decided to use a gradient descent algorithm (Press *et al.*, 2007) in combination   133
with random parameter space exploration (see Methods). This method also led   134
to good parameter combinations within a few iteration steps when starting from   135
random parameter sets for which the model deviated from the experimental results.   136
By combining random parameter exploration with a gradient descent method,   137
parameter combinations could even be found when starting from initial parameter   138
sets for which the models produced no spikes at all (**Figure S2**).   139

Reduction of channel diversity   140

Electrophysiological signatures of neurons of the same class are often unique   141
allowing a loose classification of cell types by their electrophysiology. However,   142
the spiking mechanisms often include multiple ion channels with overlapping   143
functionality to achieve these specific spiking behaviours (Coetzee *et al.*, 2006;   144
Olypher and Calabrese, 2007; Marder, 2011; Bean, 2007; O'Leary *et al.*, 2014; Drion   145
et al., 2015; Goaillard and Marder, 2021). Thus, an important question is, how   146
many channels are functionally necessary for a given cell type. We addressed this   147
question in GCs whose membrane contains a large palette of voltage- and calcium-   148
dependent conductances (Beining *et al.*, 2017). The compact activity together with   149
the multitude of ion channels in the corresponding GC model (**Figure 1C**) suggests   150
that a reduction of channels without losing accurate model performance might be   151
possible. Therefore, we explored this possibility by incremental simplification of   152
the GC model. First, we reduced the number of voltage-dependent conductances   153

in the highly detailed multi-compartmental model of GCs by 6 channels (removing   154
Cav12, Cav13, Cav32, Kv11, Kv14, SK, **Figure 1C**, *Leftmost*). This left a total of 13 155
parameters when expression in the different regions of the neuron are taken into   156
account. Thereupon, we gradually reduced the number of remaining channels to   157
a minimum of 5 ion channels (9 total parameters, leaving only the leak channels   158
pas, as well as Kir21, Na8st, BK and Cav22) finding parameter combinations that   159
satisfied our cost function using the search algorithm (**Figure 1C**, *Center left*).   160

To visualise the contribution of individual currents to neuronal model activity, we   161
employed a recently developed method of plotting the time course of the relative   162
contribution of each ionic current (Alonso and Marder, 2019). Overall, as expected,   163
the electrophysiological activity of the different valid models in **Figure 1C** was   164
similar (for overview, see **Figure S3**). Despite the large variations in the number of   165
ion channels, the course of the total inward and outward current flow displayed   166
only slight changes between the three different baseline models (**Figure 1B, C**).   167
Since GCs have a relatively simple electrophysiological repertoire (nevertheless   168
responsible for sophisticated integration of excitatory and inhibitory information),   169
a small number of membrane time constants was sufficient to generate adequate   170
firing patterns. The presence of K^+ and Ca^{2+} channels with overlapping physio-   171
logical functionality ensured that many of the channels were not crucial for the   172
maintenance of functional activity. Only the composition of the inward and out-   173
ward currents differed. In the $5-$channel model, the calcium-sensitive potassium   174
channel (BK) took over the role that 8 different K^+ conductances had shared in the   175
non-reduced model (**Figure 1C**). BK thereby became the only remaining K^+ channel   176
overall. In interaction with the Ca^{2+} conductances (Cav22), the BK channel was   177
responsible for repolarising the membrane potential following an action potential   178
in the $5-$channel model.   179

Recent experimental and theoretical studies demonstrated that neurons can com-   180
pensate for pathological changes such as channel loss, genetic overexpression,   181
morphological changes or increased input activity by up- and downregulation of   182
the remaining ion channels (Guo *et al.*, 2005; Nerbonne *et al.*, 2008; Aizenman *et al.*,   183

2003; Turrigiano *et al.*, 1999; O'Leary *et al.*, 2010; Young *et al.*, 2009; Stegen *et al.*, 2012). ₁₈₄
This ability should be impaired in the reduced model where less redundancy exists. ₁₈₅
Indeed, we found that blocking the BK or N-type Cav22 channels in the full model ₁₈₆
was readily rescued by contributions from other channels (**Figures 1C**, *Right*). It is ₁₈₇
noticeable that the loss of the BK channel was compensated by a strong upregu- ₁₈₈
lation of another calcium-sensitive channel (SK), as well as of voltage-dependent ₁₈₉
potassium channels (Kv 7.2/3, Kv 1.1, Kv 2.1, **Figure S4**, *Left*). Neither loss of BK ₁₉₀
nor Cav22 could be compensated for in the reduced $5-$channel model since it had ₁₉₁
only one active gating mechanism per ion type. Even the $9-$channel model was not ₁₉₂
able to compensate for the pathological loss of Cav22 or BK. As expected, therefore, ₁₉₃
the full GC model's diversity contributed to the model's robustness with respect to ₁₉₄
the loss of specific ion channels through existing ion channel redundancies. ₁₉₅

Random parameter tuning as a viable approach to selecting GC model

Even though small changes in the ion channel expression level can already lead ₁₉₈
to drastic changes in neuronal activity, several experimental studies observed ₁₉₉
that intrinsic properties of nerve cells can vary considerably across neurons of ₂₀₀
the same type (Golowasch *et al.*, 2002; Golowasch and Marder, 1992; MacLean *et ₂₀₁
al.*, 2003; Swensen, 2005; Schulz *et al.*, 2006, 2007). Moreover, theoretical investi- ₂₀₂
gations demonstrated that indistinguishable network and single neuron activity ₂₀₃
can be obtained from a large variety of model parameter settings (Prinz *et al.*, ₂₀₄
2004; Golowasch *et al.*, 2002). This raises the question of whether the diversity of ₂₀₅
voltage- and calcium-dependent conductances has an effect on the variability of ₂₀₆
valid parameter sets in the GC model leading to realistic spiking activity. ₂₀₇

In order to check this, we first generated $20,000$ random model instances for each ₂₀₈
of the three baseline models by randomly sampling the individual conductance ₂₀₉
densities within a range between $0\times$ and $2\times$ the value in the baseline model. As ₂₁₀

the ohmic relations between current and voltage were consistent with experimental ₂₁₁ results in all cases (see **Figure S3B**), we did not change the densities of the leak ₂₁₂ channel or the inward-rectifying Kir21 channel, which primarily contribute to ₂₁₃ the passive properties of the neuron. The population of functional parameter ₂₁₄ combinations enabled us to calculate the Pearson's correlation coefficient r for all ₂₁₅ pairs of conductance density parameters. We found weak pairwise correlations ₂₁₆ indicating low dependencies between each pair of channels and thus increasing ₂₁₇ the robustness of the model (**Figure S5**). It is likely that higher-order correlations ₂₁₈ are more prevalent in the higher-dimensional models, allowing for more different ₂₁₉ solutions that compensate for fluctuations in the expression of a single channel. ₂₂₀ The strongest pairwise correlation was observed between the expression levels of ₂₂₁ the Na^+ channel in the soma and in the AIS ($r = -0.95$). The sodium channel is ₂₂₂ essential for spike initiation and its presence in different regions of the GC suggests ₂₂₃ that compensatory mechanisms could simply be instantiated by maintaining a ₂₂₄ balance between the same currents in different regions, which results in a significant ₂₂₅ anticorrelation. Interestingly, the reduced models showed stronger and different ₂₂₆ correlations between the channels than the full model. ₂₂₇

In our selection of random parameter combinations, we found suitable models cov- ₂₂₈ ering the entire sample range of the majority of parameters (**Figure 2**). In all cases, ₂₂₉ the most constrained parameter was the density of the $8-$state Na^+ channel. This ₂₃₀ channel models the behaviour of all Na^+ conductances using a single maximum ₂₃₁ conductance parameter (Schmidt-Hieber *et al.*, 2007), so it is unsurprising that the ₂₃₂ neuron's behaviour is more sensitive to changes in this maximum. In addition, the ₂₃₃ reduction of channel diversity in the $5-$channel model limited the variability of ₂₃₄ the calcium-dependent potassium channel BK (**Figure 2**, *Right*). Surprisingly, the ₂₃₅ overall percentage of randomly selected parameter combinations that were valid ₂₃₆ increased with the number of ion channels (**Figures 3A, B**, $\sim 0.7\%$ with 5 channels ₂₃₇ (for 9 total parameters), $\sim 3.3\%$ with 9 channels (13 parameters), and $\sim 5.7\%$ with ₂₃₈ 15 channels (27 parameters)). ₂₃₉

Figure 2. Valid parameter combinations in the fully complex model are well spread.
(*Top*) Activity traces of 4 randomly picked valid parameter combinations in each of the GC models
of different complexity. (*Bottom*) Coloured dots illustrate conductance densities of the four valid
parameter combinations shown in top traces. The grey violin plots delimit the entire range covered by
the valid parameter combinations. Conductances are weighted by the surface area of the corresponding
membrane regions.

The distribution of voltage- and calcium-activated channels in cell membranes   240
is under continuous regulation (Raj and van Oudenaarden, 2008; Gal *et al.*, 2010;   241
Marder *et al.*, 2014). On the one hand, the cell is subject to homeostatic regulation   242
maintaining its electrical activity despite changes in its environment and input. On   243
the other hand, the proteins are constantly exchanged during the lifetime of a cell.   244
In order to investigate the stability of the valid parameter combinations in the differ-   245
ent models in face of parameter perturbations due to e.g. protein exchange during   246
the lifetime of a cell, we performed random walks in the parameter space. Starting   247
from a valid parameter set that accurately reproduced the experimentally derived   248
behaviour, we iteratively changed each parameter by random steps between -5% 249
and $+5\%$ of the current parameter values (counting changes in all parameters as   250
one step). The random walk stopped as soon as the parameter combination became   251
invalid, i.e. the cost function for the resulting model increased beyond 2 standard   252

deviations away from experimental results. Interestingly, the average number 253
of possible random parameter changes before model failure increased with the 254
number of ion channels in the models (**Figure 3C**). 255

Toy model points to law of large numbers 256

As shown in the previous sections, we observed an increase in valid random param- 257
eter sets when biophysical models of neurons became more complex. One possible 258
explanation could be the fact that the more complex models included different ion 259
channels of a similar type. Since some of these ion channels show very similar 260
gating dynamics (see for example Cav22, Ca12 and Cav13, see **Figure 1**) their 261
functional contributions may be partially redundant. A theorem from probability 262
theory, namely the law of large numbers can play a role under such circumstances. 263
The law of large numbers states that increasing the number of samples (in our 264
case ion channels of a similar type) described by a random variable will move 265
the average over the samples closer to the expected mean value. For example, 266
throwing multiple fair dice with sides numbered between 1 and 6 and adding 267
the results will tend to give a result that is relatively closer to the expected value 268
(the number of dice multiplied by 3.5) as more dice are used. Since in our case 269
we sample conductances of similar ion channels, the average conductance would 270
therefore converge towards the starting parameter set that we know is functional. 271

Figure 3. Valid parameter combinations in the fully complex model are more stable as compared to reduced models.

A, Percentage of valid random parameter combinations in the total population in the $2-$fold range. **B**, Percentage of valid random parameter combinations in samples with different ranges around the valid reference parameter combination. Each sample contained $5,000$ parameter combinations. **C**, Random walk through the parameter space starting from valid combinations in models of different complexity. Relative percentage distribution of the maximum number of random steps the respective models could undergo without losing their valid GC spiking behaviour. Bin size is 4 steps. Asterisks indicate mean number of steps the corresponding models could undergo while maintaining realistic activity. Performed for $2,000$ repetitions per model. **A–C**, Colours were Green: full model, Orange: $9-$channel model, Purple: $5-$channel model. **D**, Reproduction of **A–C** with a toy model representing the model result as the average value of 1 (blue), 5 (red) and 8 (green) uniform random variables between 0 and 2. *Bottom panels*, Illustration of how the distribution of solutions becomes narrower when the number of variables is increased. This effect is explained by the law of large numbers while the Gaussian distribution results from the central limit theorem.

In order to illustrate this we designed a simple toy model using random variables for each parameter. Here, we represented each open parameter of the model by one random variable with a homogeneous probability of throwing any number between 0 and 2 corresponding to the parameter ranges used in the neuronal model between $0\times$ and $2\times$ the default value (**Figure 3D**, *Bottommost*). To keep things simple for explanatory purposes, we set the model outcome to be the statistical mean of the values of all separate random variables. The law of large numbers predicts a decreasing variance of the mean value with an increasing number of independent random variables as illustrated in the sketch at the bottom of **Figure 3D**. The central limit theorem in turn predicts a Gaussian distribution for this mean over a broad range of different probability distributions for each random variable separately. In analogy to our neuronal modelling, we then constrained valid parameter combinations by a cost function allowing a maximal distance of 0.015 from the mean value, i.e. 1, averaged over all random variables.

The analogy here is limited since, in contrast to the channels in the GC model, all variables in our toy model are functionally the same and independently regulated. Moreover, the GC compartmental model applies complex nonlinear and dynamic transformations of the starting parameter space, including distinct jumps in the cost function when the model no longer produces action potentials, to reach the cost (or function) space; in the toy model the parameter and function spaces are effectively indistinguishable. However, despite its simplicity, our toy model was able to qualitatively reproduce all results from our GC model in **Figure 3A–C** (**Figure 3D**). Adding correlations to the parameter space does not qualitatively change the results (**Figure S6A**). An important observation here is that the constraint on functionality implies negative correlations between the values of the individual random variables that make up valid points in the parameter space, despite these variables being generated independently or with positive correlations. In fact, under the toy model framework, the pairwise correlations within variables that produce valid models are almost completely independent of any correlations used to generate the overall population from which valid models are drawn (**Figure S6B**).

The output correlations are instead dependent on the number of variables, with 302
higher numbers of variables leading to weaker pairwise correlations. This result 303
agrees with the finding of stronger pairwise correlations between channel densities 304
in the 5-channel model compared to the full compartmental model (**Figure S5**). 305

The law of large numbers therefore provides a plausible explanation why a larger 306
number of random instances in the more complex neuron model would more 307
readily linger around their target functionality. 308

Additional model robustness through artificial ion channel iso- 309
forms 310

We have shown that the electrophysiological behaviour of GCs can be maintained 311
despite a reduction of ion channel diversity from 15 channels to 5 channels. How- 312
ever, our results also suggest that this loss of ion channels goes along with a 313
decrease in stability, a loss of compensatory opportunities, and a significant de- 314
crease in the valid model percentage within a randomised sample. From our toy 315
model based on probability theory we postulate that it might be the mere number 316
of ion channels that contribute to the increased robustness observed in the full 317
model rather than the particular ion channel composition present there. In order to 318
validate this hypothesis, we chose to start from the reduced model and increase 319
the number of ion channels in an artificial way to check whether we could recover 320
the robustness present in the realistic full model. 321

In order to establish a quantitative relation between channel diversity and model 322
stability in such a way, we scaled up the $5-$channel model's diversity by adding 323
more instances of the calcium (Cav22) and potassium channels (BK) remaining 324
in that model. These artificial isoforms of the existing ion channels distinguished 325
themselves from the original Cav22 and BK by randomised time constants (within a 326
two-fold range of the original parameters) to allow for different dynamics through 327
the new ion channel isoforms. 328

Figure 4. Artificial expansion of ion channel diversity recovers and enhances the proportion of valid parameter combinations in the reduced 5-channel model

A, Populations of expanded dentate GC models with $0-20$ added artificial ion channel isoforms. *Left panel,* The plot shows the percentage of functional parameter combinations in a population of randomly sampled channel densities. Black dots show the populations where all ion channels (including the $8-$state Markov chain modelled Na^+ channel) were sampled in a $0-2\times$ range. Blue dots show the populations where only potassium and calcium channels were sampled in a $0-2\times$ range. *Right panel,* Similar plot as in **Figure 3B** for the black models from the *left panel*. **B**, Similar overall analysis as in **A** but for a CA1 pyramidal cell model (Jarsky *et al.*, 2005).

To examine the proportion of valid parameter combinations with increasing num- ⁠329
ber of ion channels, we created a multitude of functional GC models with up to 20 ⁠330
additional ion channel isoforms (for 35 distinct channels in total). For each given ⁠331
number of ion channel isoforms, we randomly sampled all conductance values ⁠332
in a two fold range. Thereupon we selected the three parameter combinations ⁠333
with the best fitness value for each number of ion channel isoforms and improved ⁠334
their performance by applying a gradient descent algorithm. We then followed ⁠335
the same procedure as in **Figure 3**. Using this approach, the percentage of valid ⁠336
parameter combinations steadily increased with the number of additional ion chan- ⁠337
nel isoforms until reaching a plateau between 15 and 20 additional ion channel ⁠338
isoforms, for a total of 105 to 140 additional parameters (**Figure 4A**). To further ⁠339
generalise our findings in **Figure 4A** we have applied the same procedure to a ⁠340

different neuronal model type, one simulating a CA1 pyramidal neuron (Jarsky *et al.*, 2005; Cuntz *et al.*, 2021, **Figure 4B**). Viewed together, these results show the major contribution of ion channel diversity by demonstrating that scaling up the numbers of ion channels artificially in the reduced models leads to more frequent valid parameter combinations. This is in line with the law of large numbers.

Discussion

In this study, we explored the complex landscape of valid parameter combinations in a parameter space of a detailed multi-compartmental model of dentate GCs and its simplified versions with reduced numbers of ion channels (**Figure 1**). We used a population modelling approach (Gunay *et al.*, 2008; Marder, 2011; Britton *et al.*, 2013; Sekulic *et al.*, 2014) to find multiple ion channel parameter combinations for models that successfully reproduced the electrophysiological data (**Figures 2 and S1**). We show that the biologically realistic GC model (full model) with many redundant ion channel types was more robust to ion channel perturbations than valid models with reduced ion channel diversity. Importantly, noisy ion channel expression simulated by random parameter combinations produced $\sim 6\times$ more valid GC model instances in the full model as compared to the reduced models (**Figure 3**). The robustness in the reduced model was recovered when adding artificial isoforms of existing ion channels (**Figure 4**) indicating that it is indeed the number of channels that produces this effect. We argue that this increased robustness comes in part from a direct consequence of basic probability theory.

Robustness through ion channel degeneracy in complex GC models

Most neurons contain more than a dozen different ion channels. While early computational models implemented considerably fewer channels than known in biology, more and more models exist that contain a realistic number of mechanisms

(e.g. Beining *et al.*, 2017; Hay *et al.*, 2011). Although the different potassium chan- 366
nels in mammalian cortical neurons differ genetically, some are remarkably similar 367
in their functional contribution to the electrophysiological activity of neurons (Co- 368
etzee *et al.*, 2006; Drion *et al.*, 2015). This functional similarity is often referred to 369
as degeneracy (Goaillard and Marder, 2021) and is not a phenomenon restricted 370
to neurobiology (Edelman and Gally, 2001; Tononi *et al.*, 2002). Depending on 371
the computations a neuron should implement, its dynamics only need to cover 372
certain relevant time scales, e.g. in the form of different time constants of its gating 373
variables (Gjorgjieva *et al.*, 2016). Since five channels were sufficient to support re- 374
alistic voltage dynamics at relevant time scales, we were able to reduce the original 375
variety of ion channels without observing a significant loss in the performance of 376
the model. In our study, GCs with their compact electrophysiological repertoire 377
did not require a large variety of ion channels to reproduce their characteristic 378
activity patterns. To replicate the 9 experimentally derived spiking properties, the 379
models required only one active channel of each of the different subgroups of ion 380
channels (one Na^+-, one K^+- and one Ca^{2+}-channel, as well as the leak channels; 381
Figure 1C). 382

Experimental as well as theoretical studies from the last decades revealed that phar- 383
macological manipulations like the blockage or upregulation of intrinsic or synaptic 384
mechanisms, resulting in a pathological cellular activity on a short timescale, can 385
be compensated by up- and downregulation of the remaining conductances on 386
a long timescale (MacLean *et al.*, 2003; Swensen, 2005; O'Leary *et al.*, 2014; Drion 387
et al., 2015; Guo *et al.*, 2005; Nerbonne *et al.*, 2008; Stegen *et al.*, 2012; MacLean 388
et al., 2005). Interestingly, not all manipulations can be compensated by mecha- 389
nisms of homeostatic regulation (Zhang *et al.*, 2003; Yang *et al.*, 2022), indicating 390
differences in the capability of homeostatic compensation between ion channels as 391
well as types of neurons. As opposed to other studies using biophysically realistic 392
mechanisms of homeostatic intrinsic plasticity based on calcium signals (O'Leary 393
et al., 2013, 2014; Abbott and LeMasson, 1993; Golowasch *et al.*, 1999; Liu *et al.*, 394
1998; Franci *et al.*, 2020; see also Yang *et al.*, 2022), we decided to use a gradient 395

descent approach to investigate the large and complex parameter space of possible   396
intrinsic compensations. We chose this mathematical approach also because the   397
biophysical mechanisms of intrinsic plasticity are not yet fully understood in detail.   398
The implementation of biophysically incomplete mechanisms of intrinsic plasticity   399
could lead to unnecessary limitations on the regulatory mechanisms and erroneous   400
conclusions. A homeostatic mechanism based on a single feedback signal (O'Leary   401
et al., 2013, 2014) that has been suggested to play a role in model robustness was   402
not compatible with our model in our hands since it decreased ion channel degen-   403
eracy. This is in agreement with a recent study (Yang *et al.*, 2022) that provided   404
new insights into the complex relationship between ion channel diversity and   405
homeostatic co-regulation of ion channel densities. The study by Yang *et al.* (2022)   406
suggested the necessity of more than one master feedback regulator (i.e. more   407
regulators than just global calcium) for homeostatic feedback loops, which must   408
co-tune numerous degenerate and pleiotropic ion channels to achieve multiple   409
regulated functions or objectives (cf. Pallasdies *et al.*, 2021; Jedlicka *et al.*, 2022).   410
Viewed together, we believe that diversity and (multi-signal) feedback can act as   411
independent mechanisms to ensure viable and robust solutions to multi-objective   412
optimisation problems of neurons.   413

We demonstrated that the full GC model was capable of compensating the loss of   414
any potassium and calcium channels by up- and downregulation of the remaining   415
ion channels (**Figure 1C**). In contrast, the different reduced models relied on the   416
presence of certain indispensable ion channels, without which they could not   417
capture main electrophysiological characteristics of GCs. **Figure S4** shows that   418
there can be as much as a $20-$fold variability in the density of voltage-dependent   419
ion channels. Experimental studies have observed variations of a similar order of   420
magnitude as a result of compensatory mechanisms (MacLean *et al.*, 2003). The   421
ability of these models to compensate for losses of ion channels can be attributed   422
to the overlapping or degenerate physiological function of the present potassium   423
and calcium channels (Mishra and Narayanan, 2021).   424

The reduction of the diversity of gating mechanisms goes along with a loss of space   425

to manoeuvre in the process of achieving functional target activity (O'Leary *et al.*, 2013; Drion *et al.*, 2015). In case of a loss of the BK channel, several potassium channels (see **Figure S4**) were upregulated, and thus maintained the functional behaviour of the cell. In line with the concept of degeneracy (Druckmann, 2007; Aizenman *et al.*, 2003), the overlapping functionality of different channels enabled the neuron, depending on the given conditions, to achieve a target spiking behaviour in a number of different ways.

In addition, we tested the stability of the differently reduced models against random parameter perturbations, in order to simulate putative protein exchange during the lifetime of a cell. The ongoing protein replacement is one of the reasons for the continuous regulation of voltage- and calcium-dependent channels in cell membranes (Raj and van Oudenaarden, 2008; O'Leary *et al.*, 2014; Gal *et al.*, 2010). Although no homeostatic tuning mechanism with dynamic feedback was implemented, valid parameter combinations in the complete model were able to endure far more random parameter perturbations while maintaining realistic activity than the ones in the reduced models (**Figure 3C**). This is in agreement with experimental studies, which have shown that, although homeostatic tuning rules can compensate for many perturbations and knock-outs of ion channels, not all channel deletions and perturbations can be compensated for (Zhang *et al.*, 2003). A challenge for future experimental work will be to uncover the long-term effects of ion channel knock-outs in GCs in order to find out whether our theoretical results of the outstanding robustness of GCs against channel deletions can be observed in biology.

Random parameter selection as a viable fitting strategy for neurons

Like many biological processes, gene expression is a largely stochastic process resulting in considerable heterogeneity of mRNA and protein levels (Raj and van Oudenaarden, 2008; Gal *et al.*, 2010; Sigal *et al.*, 2006). This noise in gene expression is one reason for the cell-to-cell variability. However, noise in gene expression

could be harmful for achieving functional parameter sets of ion channel expression   454
during developmental maturation or during pathological perturbations. Neurons   455
are thought to target certain desired set points (or set ranges) in the output space   456
(i.e. function or behaviour space) corresponding to valid points or subspaces in the   457
high-dimensional parameter space of expression levels of ion channels (Jedlicka   458
et al., 2022). Our simulations show that the subspace around these target values   459
in parameter space tends to be more densely filled with functional model param-   460
eters than non-valid parameters (**Figure 3B**), particularly in higher dimensions.   461
Accordingly, despite fluctuations, high-dimensional models are more likely to   462
end up in functional subspaces. Even without the implementation of homeostatic   463
regulation processes, the chance of obtaining a functional ion channel expression   464
level is relatively high. This implies that the degeneracy between ion channel   465
types and isoforms supports robust excitability profiles in neurons despite ran-   466
dom fluctuations in the expression of ion channels. Our computational analysis   467
indicates that a complex high-dimensional parameter space supports the stabil-   468
ity of neuronal excitability against perturbations that would push neurons into   469
non-functional subspaces. The reason is that the topology of the high-dimensional   470
space increases the likelihood of neurons returning into functional subspaces by   471
random ion channel parameter adjustments. An interesting extension would be to   472
compare the efficiency of activity-dependent regulation (O'Leary *et al.*, 2014; Franci   473
et al., 2020; Yang *et al.*, 2022) implemented with single or multiple homeostatic error   474
signals (Yang *et al.*, 2022), with the multi-objective optimisation (Druckmann, 2007;   475
Van Geit *et al.*, 2008; Pallasdies *et al.*, 2021; Jedlicka *et al.*, 2022) that arises naturally   476
from stochastically exploring high-dimensional parameter spaces.   477

Due to the diversity of electrophysiological mechanisms, the cell is able to generate   478
valid electrophysiological activity by random selection of parameters with a high   479
chance of success despite stochastic fluctuations in the expression of channel-   480
coding genes. We showed that there was a clear relation between the number of   481
intrinsic mechanisms and the chance to obtain a valid set of parameters from a   482
random sample around a valid point in parameter space that produces functional   483

activity in output space (**Figures 3A, B, and 4**). Furthermore, we showed that   484
many other parameter combinations existed around such a functional point in   485
the parameter space that fulfilled our criteria for functional activity. While in a   486
random $0 - 2\times$ fold sample of the initial model, about $\sim 5.7\%$ of the parameter   487
combinations showed valid GC activity, this proportion decreased steadily to   488
$\sim 0.7\%$ with a reduction of the model (**Figure 3A**). In the closer surrounding of the   489
baseline models this difference was even more obvious. While in the unreduced   490
model in the close neighbourhood of $\pm 20\%$ of the initial parameter sets over 80% 491
of the models showed characteristic GC activity, in the heavily reduced model it   492
was only about 30% (**Figure 3B**).   493

Similar to Olypher and Calabrese (2007) and Achard and De Schutter (2006) we   494
showed that near each functional point in the parameter space many other pa-   495
rameter sets exist whose activity matches the activity of the original parameter   496
set (**Figure S7 – S9**). Instead of talking about parameter sets, one might rather   497
speak about subspaces that show functional behaviour. These subspaces can have   498
different densities of parameter sets showing characteristic electrophysiological   499
activity. This depends to a great extent on the diversity of the channels (**Figures 3A,**   500
B, and 4A, *Left panel*). Furthermore, different valid subspaces with the same   501
diversity differ in their density of functional solutions located in this subspace. In   502
order to be as robust as possible against perturbations and to simplify the process   503
of parameter fitting, it seems reasonable for a neuron to target as densely populated   504
a subspace as possible.   505

Ion channel correlations and random expression   506

When analysing the conductance values of the different types of ion channels in   507
the valid models, we observed that some pairs of ion channels shared significant   508
correlations (**Figure S5**, *Red squares*). This is in line with experimental studies   509
of cell-to-cell variations in ion channels showing that some ion channels are co-   510
expressed and might be co-regulated (Schulz *et al.*, 2006, 2007; Khorkova and   511

Golowasch, 2007; Tapia *et al.*, 2018; Iacobas *et al.*, 2019; Fujita *et al.*, 2020; Kodama *et* ⁠512
al., 2020). Future large-scale analysis of channel expression in real populations of ⁠513
GCs might validate the diversity of and correlations between expression levels in ⁠514
our population models. ⁠515

In our simulations, the ion channel correlations arose from constraints on the ⁠516
resultant functionality because our model-generating strategy sampled the ion con- ⁠517
ductance levels independently. Although our population modelling was inspired ⁠518
by random noise in gene expression, it does not imply that random noise is the only ⁠519
or predominant source of cell-to-cell variability in ion channel expression. Since the ⁠520
above mentioned experimental studies found ion channel co-expression, it is likely ⁠521
that a great amount of the cell-to-cell variability in ion channel expression is due ⁠522
to transcription regulatory mechanisms, and only to some extent to the unreliable ⁠523
and noisy nature of gene expression mechanisms. Moreover, the widespread ion ⁠524
channel co-variations, which suggest structured ion channel expression in high- ⁠525
dimensional space, might arise potentially from homeostatic feedback mechanisms ⁠526
(O'Leary *et al.*, 2013, 2014; Franci *et al.*, 2020; Yang *et al.*, 2022; see above). These ob- ⁠527
servations and models do not undermine our modelling strategy, but complement ⁠528
and extend our assumption that some of the variability in ion channel expression ⁠529
is due to intrinsic noise in the expression machinery. ⁠530

Probabilistic toy model and law of large numbers ⁠531

We have put forward the law of large numbers as a possible explanation for our ⁠532
observations in the GC model. As a consequence of the law of large numbers, a ⁠533
model containing more ion channels tends to exhibit a behaviour that is closer ⁠534
to its expected target behaviour (**Figure 3**). Accordingly, we were able to recover ⁠535
the amount of robustness observed in our full model when adding artificial ion ⁠536
channel isoforms (**Figure 4**). This is a strong indicator that indeed the number of ⁠537
ion channels and not their specific composition leads to the effect that we observed. ⁠538

However, this interpretation is not mutually exclusive to the complementary insight from biophysical modelling that 15-channel model is more robust than the 5-channel one due to the increasing timescale and voltage coverage with the increasing number of ion channels (due to the partial, but not complete, redundancy between similar ion channels). The abstract toy model does not account for these two (time and voltage-related) mechanistic aspects but offers an intuition for the impact of the number of ion channel instances and their stochastic variation. The increase in the number of identical random variables in the toy model is analogous to the increase in the number of random instances of different ion channels. The main biological insight from the toy model is that if neuron samples conductances of similar ion channels around a functional point in parameter space, with the increasing number of channels the average conductance will converge towards the valid parameter set that produces functional behaviour. In summary, both biophysical and toy models indicate that the large number of ion channel subtypes and isoforms expressed by a neuronal type supports the tuning and robustness of the electrophysiological phenotype.

Conclusions and outlook

Overall, our results suggest that the diversity of ion channels allows for increased robustness and higher flexibility of finding a solution in the complex parameter space of a neuron's excitability. It will be interesting to investigate whether our findings here translate to other biologically complex systems, in which case they will most likely affect our general understanding of how evolution deals with complex organisms.

Acknowledgments

This work was supported by BMBF (No. 01GQ1406 - Bernstein Award 2013 to HC, No. 031L0229 - HUMANEUROMOD to PJ) and by funds from the von Behring Röntgen Foundation to PJ. JT acknowledges support by the Johanna Quandt foundation. MS was supported by the International Max Planck Research School (IMPRS) for Neural Circuits in Frankfurt. The authors declare to have no competing financial interests.

Author contributions

M.S., A.D.B., A.G., J.T., P.J., and H.C. designed the study. M.S. performed the simulations and analysed the data. M.S., A.D.B., A.G. J.T., P.J., and H.C. wrote the paper.

Materials and methods

All simulations were performed in *Matlab 2017b* (Mathworks, Natick, MA, USA). Single neuron simulations were performed using *T2N* (Beining *et al.*, 2017, `www.treestoolbox.org/T2N`), a Matlab interface between the open source package *TREES toolbox* (Cuntz *et al.*, 2010, 2011, `www.treestoolbox.org`) and the *NEURON* simulation environment (Hines and Carnevale, 1997, `www.neuron.yale.edu`). Predefined functions from *TREES toolbox*, *T2N* as well as additional custom *Matlab* code were used to generate and analyse the models. All code will be made available on Zenodo upon publication.

The granule cell (GC) model

The GC model used in this study has been fully described in (Beining *et al.*, 2017). Briefly, the model was designed to reproduce passive and active GC properties as determined by voltage and current clamp experiments, dendritic patch recordings of bAPs, and intracellular calcium imaging. In order to reduce the number of parameters and to speed up simulations we simplified the morphology by deleting the artificially added axon. The loss of the axon was compensated by slight changes of the maximum conductances in the axon initial segment (AIS). Since the HCN channel in its original form had no influence on control GC activity, we did not take it into account. The compartment-specific distributions of ion channels are shown in Table S1. Detailed descriptions of the individual ion channels can be found in Beining *et al.* (2017). We used a realistic three-dimensional granule cell morphology from Schmidt-Hieber *et al.* (2007).

Stimulation protocols and cost function

Instead of using a single optimal error function, we decided to adopt a strategy that allows to take into account several potentially important properties of GC activity. To get a first impression of the "goodness of a model", we compared the experimental (Mongiat *et al.*, 2009) and the model spiking-properties following a $200ms$ current injection of 50 or $90pA$. The stimulation protocol was as follows: $50ms$ prerun without stimulation, followed by $200ms$ somatic current injection of 50 or $90pA$ followed by a $50ms$ long period without current injection.

We extracted the following 9 spiking properties (**Figures 1A**) from the raw traces of current injections with 50 and $90pA$:

1. Numbers of spikes fired within $200ms$ under current clamp.

2. Latency of first spike after stimulus onset in ms.

3. The voltage threshold was defined as the voltage at which the rate of change 607
   of membrane potential exceeded $15\frac{mV}{ms}$. 608

4. Average amplitude of spikes. 609

5. The fast after hyperpolarisation (fAHP) amplitude was calculated as the 610
   voltage difference between the spiking threshold and the minimum potential 611
   within $5ms$ after a spike. 612

6. Absolute value of fast after hyperpolarisation (fAHP) amplitude. 613

7. The action potential width was measured at half the height of the spike 614
   amplitude. 615

8. Interspike interval (ISI) in ms between the first and second spike during 616
   current clamp. 617

9. The adaptation index AI was calculated in the following manner: $AI = $ 618
   $1 - \frac{ISI_1}{ISI_{end}}$, where ISI_1 is the first and ISI_{end} the last ISI. 619

The spiking features for any given parameter combination in the model were then 620
compared with the same experimentally derived spiking features (Mongiat *et al.*, 621
2009) and expressed in units of standard deviation. This approach allowed us to 622
take into account the intrinsic variability of each feature separately. The overall 623
fitness F_i of spike feature i was defined as: 624

$$F_i = \frac{|SF_i - \overline{SF}_{i,exp}|}{SD_{i,exp}} \tag{1}$$

where $\overline{SF}_{i,exp}$ refers to the average value of the spike feature i and $SD_{i,exp}$ to the 625
standard deviation of the spike feature i across all recorded GCs. The value of the 626
spike feature of the corresponding model for a given parameter combination was 627
SF_i. For a parameter combination to be accepted as a valid combination, it was 628
required to fulfil the following condition: 629

$$P = max \left(\frac{|SF_i - \overline{SF}_{i,exp}|}{SD_{i,exp}} \right) < 2, \ for \ i = 1, 2, ..., 9 \tag{2}$$

The value of the Pareto efficiency P corresponded to the fitness F_i of the spiking   630
feature SF_i that deviated most from the experimental average.   631

The search algorithm   632

To search for parameter sets that match our criteria for valid GC activity we   633
combined random sampling with a gradient descent algorithm. In order to search   634
for local minima we used a conjugate gradient descent technique (Press *et al.*,   635
2007). Conjugate gradient descent techniques involve successive calculations of   636
local gradients followed by the exploration of the parameter space along a vector   637
derived from that gradient. Starting from a random or given point in the parameter   638
space, we calculated the gradients for each dimension with two sample points to   639
smooth the slopes. The algorithm evaluates the calculated gradients of the fitness   640
function in each dimension and moves in the direction of the steepest descent with   641
respect to the cost function. The sample points where calculated in steps of $\pm 5\%$ of   642
the corresponding parameter value. This procedure was then repeated until the   643
method converged to a local minimum of the corresponding Pareto efficiency P 644
(**Equation 2**). The successive line minimisation was done in conjugated directions,   645
so that the successive minimisations were as independent as possible. Theoretically,   646
this ensured that the parameter search found a local minimum of the target function   647
P. For some initial parameter combinations, large areas of the parameter space   648
were completely flat (i.e. the gradient was zero). This was especially the case   649
when the initial models showed no spiking activity (**Figure S2B**). In this case,   650
we increased the size of the iteration steps consecutively by $\pm 5\%$. If still (after   651
increasing the step size to $\pm 50\%$) no gradients other than zero were found or the   652
local minima did not fulfil the criteria of functional GC excitability, we randomised   653
the parameters in the next step in an iteratively increasing range (from $\pm 10\%$ of   654

the corresponding parameter values in steps of $\pm 10\%$ up to $\pm 50\%$). The gradient 655
descent algorithm was used to find the parameter settings of the reduced models. 656
Starting from the full model (**Figure 1C**, **Table S1, S2 and S3**), we gradually 657
reduced the number of ion channels, starting with the channels that influenced the 658
cost function the least. 659

Diversity expansion 660

In order to generate models with controllable amounts of ion channels we used the 661
reduced $5-$channel model as a basis. We then produced multiple instances each 662
of the remaining potassium (BK) and calcium (Cav22) channels. Each artificial 663
channel form obtained in such a way was associated with a randomised time 664
constant between $0\times$ and $2\times$ the value in the original GC model to obtain altered 665
dynamics. Furthermore, we randomised the conductances and applied the search 666
algorithm to reproduce characteristic GC activity to derive all base models with 667
different complexities in **Figure 4**. 668

Toy model 669

We created a toy model to test whether the law of large numbers is a plausible 670
explanation for the phenomena we observed in the GC model. In order to mimic 671
the distribution of functional overlapping ion channel expressions in a population 672
of GC models around a genetically targeted functional set point we used randomly 673
uniformly sampled variables between zero and two (**Figure 3D**). A valid toy model 674
is defined as having a smaller average deviation from the mean (targeted value) 675
than 0.015. By decreasing the sample range around the mean in steps of 0.1 down 676
to a sample range between 0.9 and 1.1 we change the intensity of fluctuations 677
around the target point (**Figure 3D**). 678

To expand the toy model to account for possible intrinsic correlations in the expres-

sion of ion channels (**Figure S6**), we used a Gaussian copula to impose a correlation structure on the random variables with uniform marginals and specified pairwise correlations. For a desired positive pairwise correlation ρ in a system of n variables we generated an $n \times n$ correlation matrix $\mathbf{R}$ with elements $\mathbf{R}_{i,j} = \rho$ if $i \neq j$ and $\mathbf{R}_{i,i} = 1$. If a random variable $\mathbf{u} = (u_1, u_2, ..., u_n)$ and each u_i is independently uniformly distributed in the range $[0, 1]$, then the correlated random variable $\mathbf{v}$ with uniform marginals on $[0, 1]$ is given by

$$\mathbf{v} = \boldsymbol{\Phi}_{\mathbf{R}}\big(\Phi^{-1}(u_1), \Phi^{-1}(u_2), ..., \Phi^{-1}(u_n)\big) \tag{3}$$

where $\boldsymbol{\Phi}_{\mathbf{R}}$ is the cumulative distribution function of a multivariate Gaussian distri- 679
bution in n-dimensions with mean 0 and covariance matrix $\mathbf{R}$ and Φ^{-1} is the inverse 680
cumulative distribution function of a standard univariate Gaussian. Multiplying $\mathbf{v}$ 681
by 2 maps it back to the same space as the uncorrelated toy model. 682

Hyperplanes 683

To learn more about the relationship of the set of valid models, we created linear 684
combinations of our best solutions. This method was adopted from Achard and 685
De Schutter (2006) and allowed us to better estimate whether the solutions lie 686
on a common low-dimensional manifold within the high-dimensional parameter 687
space of the GC model variants (**Figure S7 – S9**). As a first step, we created 688
linear combinations out of weighted sums of a pair of solutions. We weighted the 689
parameters of the respective model between 0.1 and 0.9 with a step size of 0.1. The 690
weighting of the second solution was chosen such that the sum of the weights was 691
equal to 1. As soon as the Pareto efficiency of all evaluated linear combinations 692
fulfilled the criteria for characteristic GC spiking, we assumed that the respective 693
models were connected. In the next step, we created linear combinations of three 694
different valid solutions to visualise the hyperplanes in two dimensions. We 695
used several triplets of valid parameter sets and weighted two of them with 696
values between -1.5 and 2.5 using a step size of 0.04. The corresponding grid of 697

combinations was visualised in a two-dimensional plot. The weighting of the third 698
selected parameter set was chosen in a way that the sum of all weights was equal 699
to 1. The hyperplanes consisted of several thousand points, whereby the parameter 700
sets with negative values were removed. As a result, each hyperplane had different 701
boundaries and thus a different size. Finally, for each of these points we ran 702
simulations and calculated their Pareto efficiency. The Pareto efficiency of the 703
models without spiking behaviour was set to 6, which explains the abrupt change 704
of colour on the right side of **Figure S7**. The colour selection of the plots allowed a 705
clear distinction between the valid (green) and the nonvalid (blue) models. 706

References 707

Abbott LF, LeMasson G (1993) Analysis of neuron models with dynamically 708
    regulated conductances. *Neural Computation* 5:823–842. 709

Achard P, De Schutter E (2006) Complex parameter landscape for a complex neuron 710
    model. *PLoS Computational Biology* 2:e94. 711

Aizenman CD, Akerman CJ, Jensen KR, Cline HT (2003) Visually driven reg- 712
    ulation of intrinsic neuronal excitability improves stimulus detection in vivo. 713
    Neuron 39:831–842. 714

Alonso LM, Marder E (2019) Visualization of currents in neural models with similar 715
    behavior and different conductance densities. *eLife* 8:e42722. 716

Bahl A, Stemmler MB, Herz AV, Roth A (2012) Automated optimization of a 717
    reduced layer 5 pyramidal cell model based on experimental data. *Journal of* 718
    Neuroscience Methods 210:22–34. 719

Bean BP (2007) The action potential in mammalian central neurons. *Nature Reviews* 720
    Neuroscience 8:451–465. 721

Beining M, Mongiat LA, Schwarzacher SW, Cuntz H, Jedlicka P (2017) T2N as a new tool for robust electrophysiological modeling demonstrated for mature and adult-born dentate granule cells. *eLife* 6:e26517.

Britton OJ, Bueno-Orovio A, Van Ammel K, Lu HR, Towart R, Gallacher DJ, Rodriguez B (2013) Experimentally calibrated population of models predicts and explains intersubject variability in cardiac cellular electrophysiology. *PNAS* 110:E2098–E2105.

Coetzee WA, Amarillo Y, Chiu J, Chow A, Lau D, Mccormack T, Morena H, Nadal MS, Ozaita A, Pountney D, Saganich M, Miera EVS, Rudy B (2006) Molecular diversity of K+ channels. *Annals of the New York Academy of Sciences* 868:233–255.

Connors BW, Gutnick MJ (1990) Intrinsic firing patterns of diverse neocortical neurons. *Trends in Neurosciences* 13:99–104.

Cuntz H, Bird AD, Mittag M, Beining M, Schneider M, Mediavilla L, Hoffmann FZ, Deller T, Jedlicka P (2021) A general principle of dendritic constancy: A neuron's size- and shape-invariant excitability. *Neuron* 109:3647–3662.e7.

Cuntz H, Forstner F, Borst A, Häusser M (2010) One rule to grow them all: A general theory of neuronal branching and its practical application. *PLoS Computational Biology* 6:e1000877.

Cuntz H, Forstner F, Borst A, Häusser M (2011) The TREES Toolbox—Probing the basis of axonal and dendritic Branching. *Neuroinformatics* 9:91–96.

Drion G, O'Leary T, Marder E (2015) Ion channel degeneracy enables robust and tunable neuronal firing rates. *PNAS* 112:E5361–E5370.

Druckmann S (2007) A novel multiple objective optimization framework for constraining conductance-based neuron models by experimental data. *Frontiers in Neuroscience* 1:7–18.

Edelman GM, Gally JA (2001) Degeneracy and complexity in biological systems. *PNAS* 98:13763–13768.

Franci A, O'Leary T, Golowasch J (2020) Positive dynamical networks in neuronal regulation: How tunable variability coexists with robustness. *IEEE Control Systems Letters* 4:946–951.

Fujita H, Kodama T, Du Lac S (2020) Modular output circuits of the fastigial nucleus for diverse motor and nonmotor functions of the cerebellar vermis. *Elife* 9:e58613.

Gal A, Eytan D, Wallach A, Sandler M, Schiller J, Marom S (2010) Dynamics of excitability over extended timescales in cultured cortical neurons. *Journal of Neuroscience* 30:16332–16342.

Gjorgjieva J, Drion G, Marder E (2016) Computational implications of biophysical diversity and multiple timescales in neurons and synapses for circuit performance. *Current Opinion in Neurobiology* 37:44–52.

Goaillard JM, Marder E (2021) Ion channel degeneracy, variability, and covariation in neuron and circuit resilience. *Annual Review of Neuroscience* Online ahead of print.

Golowasch J, Marder E (1992) Ionic currents of the lateral pyloric neuron of the stomatogastric ganglion of the crab. *Journal of Neurophysiology* 67:318–331.

Golowasch J, Casey M, Abbott LF, Marder E (1999) Network stability from activity-dependent regulation of neuronal conductances. *Neural Computation* 11:1079–1096.

Golowasch J, Goldman MS, Abbott LF, Marder E (2002) Failure of averaging in the construction of a conductance-based neuron model. *Journal of Neurophysiology* 87:1129–1131.

Gunay C, Edgerton JR, Jaeger D (2008) Channel density distributions explain spiking variability in the globus pallidus: A combined physiology and computer simulation database approach. *Journal of Neuroscience* 28:7476–7491.

Guo W, Jung WE, Marionneau C, Aimond F, Xu H, Yamada KA, Schwarz TL, Demolombe S, Nerbonne JM (2005) Targeted deletion of Kv4.2 eliminates Ito,f and

results in electrical and molecular remodeling, with no evidence of ventricular   776
hypertrophy or myocardial dysfunction. *Circulation Research* 97:1342–1350.   777

Hay E, Hill S, Schürmann F, Markram H, Segev I (2011) Models of neocortical layer   778
5b pyramidal cells capturing a wide range of dendritic and perisomatic active   779
properties. *PLoS Computational Biology* 7:e1002107.   780

Herrera-Valdez MA, McKiernan EC, Berger SD, Ryglewski S, Duch C, Crook S   781
(2013) Relating ion channel expression, bifurcation structure, and diverse firing   782
patterns in a model of an identified motor neuron. *Journal of Computational*   783
Neuroscience 34:211–229.   784

Hille B (2001) *Ion channels of excitable membranes* Oxford University Press.   785

Hines ML, Carnevale NT (1997) The NEURON simulation environment. *Neural*   786
Computation 9:1179–1209.   787

Iacobas DA, Iacobas S, Lee PR, Cohen JE, Fields RD (2019) Coordinated activity of   788
transcriptional networks responding to the pattern of action potential firing in   789
neurons. *Genes* 10:754.   790

Jarsky T, Roxin A, Kath WL, Spruston N (2005) Conditional dendritic spike prop-   791
agation following distal synaptic activation of hippocampal CA1 pyramidal   792
neurons. *Nature Neuroscience* 8:1667–1676.   793

Jedlicka P, Bird AD, Cuntz H (2022) Pareto optimality, economy–effectiveness   794
trade-offs and ion channel degeneracy: improving population modelling for   795
single neurons. *Open Biology* 12:14–23.   796

Keren N, Peled N, Korngreen A (2005) Constraining compartmental models using   797
multiple voltage recordings and genetic algorithms. *Journal of Neurophysiol-*   798
ogy 94:3730–3742.   799

Khorkova O, Golowasch J (2007) Neuromodulators, not activity, control coordi-   800
nated expression of ionic currents. *Journal of Neuroscience* 27:8709–8718.   801

Kirkpatrick S, Gelatt CD, Vecchi MP (1983) Optimization by simulated annealing. 802
    Science 220:671–680. 803

Kodama T, Gittis AH, Shin M, Kelleher K, Kolkman KE, McElvain L, Lam M, Du Lac 804
    S (2020) Graded coexpression of ion channel, neurofilament, and synaptic genes 805
    in fast-spiking vestibular nucleus neurons. *Journal of Neuroscience* 40:496–508. 806

Liu Z, Golowasch J, Marder E, Abbott LF (1998) A model neuron with activity- 807
    dependent conductances regulated by multiple calcium sensors. *Journal of Neuro-* 808
    science 18:2309–2320. 809

MacLean JN, Zhang Y, Goeritz ML, Casey R, Oliva R, Guckenheimer J, Harris- 810
    Warrick RM (2005) Activity-independent coregulation of IA and Ih in rhythmi- 811
    cally active neurons. *Journal of Neurophysiology* 94:3601–3617. 812

MacLean JN, Zhang Y, Johnson BR, Harris-Warrick RM (2003) Activity-independent 813
    homeostasis in rhythmically active neurons. *Neuron* 37:109–120. 814

Marder E (2011) Variability, compensation, and modulation in neurons and circuits. 815
    PNAS 108:15542–15548. 816

Marder E, Goaillard JM (2006) Variability, compensation and homeostasis in neuron 817
    and network function. *Nature Reviews Neuroscience* 7:563–574. 818

Marder E, O'Leary T, Shruti S (2014) Neuromodulation of circuits with variable pa- 819
    rameters: single neurons and small circuits reveal principles of state-dependent 820
    and robust neuromodulation. *Annual Review of Neuroscience* 37:329–346. 821

Mishra P, Narayanan R (2021) Ion-channel regulation of response decorrelation 822
    in a heterogeneous multi-scale model of the dentate gyrus. *Current Research in* 823
    Neurobiology 2:100007. 824

Mitchell M (1998) *An Introduction to Genetic Algorithms* MIT Press. 825

Mongiat LA, Espósito MS, Lombardi G, Schinder AF (2009) Reliable activation of 826
    immature neurons in the adult hippocampus. *PLoS One* 4:e5320. 827

Nerbonne JM, Gerber BR, Norris A, Burkhalter A (2008) Electrical remodelling maintains firing properties in cortical pyramidal neurons lacking KCND2-encoded A-type K+ currents. *Journal of Physiology* 586:1565–1579.

O'Leary T, van Rossum MC, Wyllie DJ (2010) Homeostasis of intrinsic excitability in hippocampal neurones: Dynamics and mechanism of the response to chronic depolarization. *Journal of Physiology* 588:157–170.

O'Leary T, Williams AH, Caplan JS, Marder E (2013) Correlations in ion channel expression emerge from homeostatic tuning rules. *PNAS* 110:E2645–E2654.

O'Leary T, Williams AH, Franci A, Marder E (2014) Cell types, network homeostasis, and pathological compensation from a biologically plausible ion channel expression model. *Neuron* 82:809–821.

Olypher AV, Calabrese RL (2007) Using constraints on neuronal activity to reveal compensatory changes in neuronal parameters. *Journal of Neurophysiology* 98:3749–3758.

Pallasdies F, Norton P, Schleimer JH, Schreiber S (2021) Neural optimization: Understanding trade-offs with pareto theory. *Current Opinion in Neurobiology* 71:84–91.

Press WH, Teukolsky SA, Vetterling WT, Flannery BP (2007) *Numerical Recipes* Cambridge University Press.

Prinz AA, Billimoria CP, Marder E (2003) Alternative to hand-tuning conductance-based models: construction and analysis of databases of model neurons. *Journal of Neurophysiology* pp. 3998 – 4015.

Prinz AA, Bucher D, Marder E (2004) Similar network activity from disparate circuit parameters. *Nature Neuroscience* 7:1345–1352.

Raj A, van Oudenaarden A (2008) Nature, nurture, or chance: stochastic gene expression and its consequences. *Cell* 135:216–226.

Rathour RK, Narayanan R (2019) Degeneracy in hippocampal physiology and   853
plasticity. *Hippocampus* 29:980–1022.   854

Rudy B (1988) Diversity and ubiquity of K channels. *Neuroscience* 25:729–749.   855

Schmidt-Hieber C, Jonas P, Bischofberger J (2007) Subthreshold dendritic signal   856
processing and coincidence detection in dentate gyrus granule cells. *Journal of*   857
Neuroscience 27:8430–8441.   858

Schulz DJ, Goaillard JM, Marder E (2007) Quantitative expression profiling of   859
identified neurons reveals cell-specific constraints on highly variable levels of   860
gene expression. *PNAS* 104:13187–13191.   861

Schulz DJ, Goaillard JM, Marder E (2006) Variable channel expression in identified   862
single and electrically coupled neurons in different animals. *Nature Neuro-*   863
science 9:356–362.   864

Sekulic V, Lawrence JJ, Skinner FK (2014) Using multi-compartment ensemble   865
modeling as an investigative tool of spatially distributed biophysical balances: 866
Application to hippocampal oriens-lacunosum/moleculare (O-LM) cells. *PLoS*   867
ONE 9:e106567.   868

Sigal A, Milo R, Cohen A, Geva-Zatorsky N, Klein Y, Liron Y, Rosenfeld N, Danon   869
T, Perzov N, Alon U (2006) Variability and memory of protein levels in human   870
cells. *Nature* 444:643–646.   871

Stegen M, Kirchheim F, Hanuschkin A, Staszewski O, Veh RW, Wolfart J (2012)   872
Adaptive intrinsic plasticity in human dentate gyrus granule cells during tempo-   873
ral lobe epilepsy. *Cerebral Cortex* 22:2087–2101.   874

Swensen AM (2005) Robustness of burst firing in dissociated Purkinje neurons   875
with acute or long-term reductions in sodium conductance. *Journal of Neuro-*   876
science 25:3509–3520.   877

Tapia M, Baudot P, Formisano-Tréziny C, Dufour MA, Temporal S, Lasserre M,   878
Marquèze-Pouey B, Gabert J, Kobayashi K, Goaillard JM (2018) Neurotransmitter   879

identity and electrophysiological phenotype are genetically coupled in midbrain dopaminergic neurons. *Scientific reports* 8:1–14.

Tononi G, Sporns O, Edelman GM (2002) Measures of degeneracy and redundancy in biological networks. *PNAS* 96:3257–3262.

Turrigiano G, Desai NS, Rutherford LC (1999) Plasticity in the intrinsic excitability of cortical pyramidal neurons. *Nature Neuroscience* 2:515–520.

Van Geit W, De Schutter E, Achard P (2008) Automated neuron model optimization techniques: a review. *Biological cybernetics* 99:241–251.

Vanier MC, Bower JM (1999) A comparative survey of automated parameter-search methods for compartmental neural models. *Journal of Computational Neuroscience* 7:149–171.

Yang J, Shakil H, Ratté S, Prescott SA (2022) Minimal requirements for a neuron to coregulate many properties and the implications for ion channel correlations and robustness. *Elife* 11:e72875.

Young CC, Stegen M, Bernard R, Müller M, Bischofberger J, Veh RW, Haas CA, Wolfart J (2009) Upregulation of inward rectifier K+ (Kir2) channels in dentate gyrus granule cells in temporal lobe epilepsy. *Journal of Physiology* 587:4213–4233.

Zhang Y, Oliva R, Gisselmann G, Hatt H, Guckenheimer J, Harris-Warrick RM (2003) Overexpression of a hyperpolarization-activated cation current (Ih) channel gene modifies the firing activity of identified motor neurons in a small neural network. *Journal of Neuroscience* 23:9059–9067.

Supporting information

Figure S1. Electrophysiological properties of mouse GCs.

Experimental data from (Mongiat *et al.*, 2009). **A**, Voltage traces of eight different GCs during $200ms$ current clamp injection of $90pA$. **B**, Frequency of action potentials elicited by $200ms$ lasting current injections (mean and standard deviation from raw traces, experimental standard deviation is shown as grey patches). **C**, Current-voltage (I–V) relationships (mean and standard deviation from raw traces, experimental standard deviation is shown as grey patches). **D**, Phase plots of the first action potential during $90pA$ current clamp. Modified from Figure 2 in Beining *et al.* (2017).

Figure S2. Gradient descent using multi-objective optimisation.
A, Temporal evolution of Pareto optimality (*top*, see Eq. 2) using the gradient descent method. Solutions are considered valid once their Pareto optimality drops below 2 (dashed line). Initial parameter combinations are random non-valid parameter combinations within a range between $0\times$ and $2\times$ the value in the reference parameter set. (*bottom*) Voltage traces of the model with initial parameter combinations (grey) and optimised parameters (green). **B**, Same as in **A**, but all initial parameter combinations were in a similar order of magnitude of Pareto optimality with corresponding models that did not even produce spikes.

Figure S3. Comparison of the different GC models in Figure 1C.

A–D, Similar panels as in **Figure S1** for the different models and respective parameter combinations as in **Figure 2A**.

Figure S4. Valid parameter combinations in models that compensate for the knock-out of the BK (Left) and Cav22 (Right) channel.

Valid parameter combinations in the fully complex model are well spread and more stable as compared to reduced models. Activity traces of 4 randomly picked valid parameter combinations in models successfully compensating the corresponding knock-out (Top). Coloured dots illustrate conductance densities of the four valid parameter combinations shown in top traces (Bottom). Violin plots show the probability distribution of valid parameter combinations. Conductances are weighted by the surface area of the corresponding membrane regions.

Figure S5. Correlations between pairs of channel conductances in the different populations.
Significant correlations are highlighted by red boxes (p-value < 0.01). Pairwise correlations in population of **A**, $15-$channel models, **B**, $9-$channel models, **C**, $5-$channel models.

Figure S6. Correlations in the toy model.
A, Effects of pairwise correlations on the proportion of valid models for different numbers of variables. The 1 variable model is not plotted as it is not affected by correlations. All models converge to the same point as their elements become perfectly correlated and the effective number of dimensions is reduced to 1. **B**, Observed output correlations in valid models as a function of the pairwise correlation used to generate the population from which valid models are drawn. For almost all input correlations the observed correlation depends only on the number of variables.

Figure S7. 2D illustrations of hyperplanes in the parameter space.
Hyperplane analysis inspired by Achard and De Schutter (2006) for the $15-$channel model. **A**, The hyperplane of **B** is shown in red as projection onto $g_{Na8st,AIS}$ vs. $g_{SK2,AIS}$ plane. 25 randomly chosen valid parameter combinations are represented by dots. The blue hyperplane is parallel to the red and is defined by the addition of 10% of the SD of all solutions (in every dimension). **B**, Hyperplane defined by the three individuals on the red line in **A**. The Fitness of all points is colour scaled. The three original individuals are highlighted as red dots. **C**, The red dots mark the places parallel to the 3 originally selected individuals.

Figure S8. 2D illustrations of hyperplanes in the parameter space.
Hyperplane analysis inspired by Achard and De Schutter (2006) for the $9-$channel model. **A**, The hyperplane of **B** is shown in red as projection onto $g_{Kv34,AIS}$ vs. $g_{Cav22,AIS}$ plane. 25 randomly chosen valid parameter combinations are represented by dots. The blue hyperplane is parallel to the red and is defined by the addition of 10% of the SD of all solutions (in every dimension). **B**, Hyperplane defined by the three individuals on the red line in **A**. The Fitness of all points is colour scaled. The three original individuals are highlighted as red dots. **C**, The red dots mark the places parallel to the 3 originally selected individuals.

Figure S9. 2D illustrations of hyperplanes in the parameter space.

Hyperplane analysis inspired by Achard and De Schutter (2006) for the $5-$channel model. **A**, The hyperplane of **B** is shown in red as projection onto $g_{na8st,AIS}$ vs. $g_{Cav22,AIS}$ plane. 25 randomly chosen valid parameter combinations are represented by dots. The blue hyperplane is parallel to the red and is defined by the addition of 10% of the SD of all solutions (in every dimension). **B**, Hyperplane defined by the three individuals on the red line in **A**. The Fitness of all points is colour scaled. The three original individuals are highlighted as red dots. **C**, The red dots mark the places parallel to the 3 originally selected individuals.

Name	AIS	Soma	Dendrite
pas	6.593×10^{-6}	1.385×10^{-5}	1.385×10^{-5}
Kir 2.1	6.741×10^{-5}	1.415×10^{-4}	1.415×10^{-4}
Na8st	0.614	0.1478	
Kv 1.1	2.76×10^{-4}		
Kv 1.4	1.77×10^{-2}		
Kv 2.1		0.0022	
Kv 3.4	0.6987		
Kv 4.2			0.0039
Kv 7.2/3	0.0031		
Cav 1.2	3.1×10^{-4}	7.1×10^{-5}	2×10^{-5}
Cav 1.3	5.48×10^{-6}	2.5×10^{-5}	3.7×10^{-6}
Cav 2.2	3.19×10^{-7}	7.4×10^{-5}	5.8×10^{-6}
Cav 3.2	1.22×10^{-5}	1.6×10^{-5}	3.8×10^{-5}
BK			
α	0.0018	9.3×10^{-4}	
β	0.51	0.0148	
SK2	1.1×10^{-5}	3.7×10^{-8}	8.5×10^{-7}

Table S1.   Summary of ion channel densities and models implemented in the $15-$channel model.

Ion channels and their expression profiles in the corresponding morphological compartments. Conductance densities are given in units of mS/cm^2.

Name	AIS	Soma	Dendrite
pas	6.593×10^{-6}	1.385×10^{-5}	1.385×10^{-5}
Kir 2.1	6.741×10^{-5}	1.415×10^{-4}	1.415×10^{-4}
Na8st	0.4925	0.0881	
Kv 2.1		0.0071	
Kv 3.4	0.0339		
Kv 7.2/3	0.0074		
Kv 4.2			0.0022
Cav 2.2	4.77×10^{-11}	4.5×10^{-4}	3.56×10^{-5}
BK			
α	1.25×10^{-7}	0.0043	
β	0.0148	0.0156	

Table S2. Summary of ion channel densities and models implemented in the $9-$**channel model.**

Ion channels and their expression profiles in the corresponding morphological compartments. Conductance densities are given in units of mS/cm^2.

Name	AIS	Soma	Dendrite
pas	6.593×10^{-6}	1.385×10^{-5}	1.385×10^{-5}
Kir 2.1	6.741×10^{-5}	1.415×10^{-4}	1.415×10^{-4}
Na8st	0.306	0.119	
Cav 2.2	5.82×10^{-15}	8.64×10^{-4}	1.22×10^{-4}
BK			
α	1.16×10^{-7}	0.0132	
β	1.321	0.0185	

Table S3. Summary of ion channel densities and models implemented in the $5-$channel model.

Ion channels and their expression profiles in the corresponding morphological compartments. Conductance densities are given in units of mS/cm^2.