Impaired bidirectional communication between interneurons and

oligodendrocyte precursor cells affects cognitive behavior

Lipao Fang[1], Na Zhao[1], Laura C. Caudal[1], Renping Zhao[2], Ching-Hsin Lin[3], Hsin-Fang

Chang[3], Nadine Hainz[4], Carola Meier[4], Bernhard Bettler[5], Wenhui Huang[1], Anja

Scheller[1], Frank Kirchhoff[1,*], Xianshu Bai[1,*]

1. Molecular Physiology, CIPMM, University of Saarland, 66421 Homburg, Germany

2. Biophysics, CIPMM, University of Saarland, 66421 Homburg, Germany

3. Cellular Physiology, CIPMM, University of Saarland, 66421 Homburg, Germany

4. Department of Anatomy and Cell Biology, University of Saarland, 66421 Homburg,

Germany

5. Department of Biomedicine, University of Basel, 4056 Basel, Switzerland

Correspondence to:

Prof. Dr. Frank Kirchhoff,

Molecular Physiology,

CIPMM, University of Saarland,

66421 Homburg, Germany

frank.kirchhoff@uks.eu

Dr. Xianshu Bai

Molecular Physiology,

CIPMM, University of Saarland,

66421 Homburg, Germany

xianshu.bai@uks.eu

Abstract

Cortical neural circuits are complex but very precise networks of balanced excitation and inhibition (E/I). Yet, the molecular and cellular mechanisms that form the E/I balance are just beginning to emerge. Here, using conditional GABA$_B$ receptor-deficient mice we identified a GABA/TNF-related cytokine (TNFSF12)-mediated bidirectional communication pathway between Parvalbumin-positive (PV$^+$) fast spiking interneurons and oligodendrocyte precursor cells (OPCs) that determines the density and function of interneurons in the developing medial prefrontal cortex (mPFC). Interruption of the GABAergic signaling to OPCs resulted in reduced myelination and hypoactivity of interneurons, strong changes of cortical network activities and impaired cognitive behavior. In conclusion, glial transmitter receptors are pivotal elements in finetuning distinct brain functions.

Introduction

The inhibition of cortical network activity is performed by interneurons that release the inhibitory transmitter γ-aminobutyric acid (GABA) which acts on ionotropic GABA$_A$ and metabotropic GABA$_B$ receptors (GABA$_B$R) [1, 2]. Alterations of interneuron cell density as well as concomitant changes of firing activity are often observed in several neuropsychiatric conditions [3-6]. In the mouse cortex, higher frequencies of spontaneous inhibitory postsynaptic currents (sIPSCs) were observed after increasing the interneuron density by transplantation of precursor cells [7]. Similarly, blocking of interneuron apoptosis caused a decrease of the excitation/inhibition (E/I) ratio [6, 8]. Interneuron density is a pivotal determinant of correct inhibitory circuits in the central nervous system (CNS). During development, a surplus of interneurons is generated that populates the cortical plate [9]. Subsequently, about 40 % of cortical inhibitory neurons are eliminated by programmed cell death during the first two postnatal weeks

[7]. One hypothesis links the apoptosis of supernumerary neurons with the onset of first neuron-neuron connections [10]. Only the neurons that receive sufficient neurotrophic signals from their target cells will survive [10, 11].

Oligodendrocyte precursor cells (OPCs) form synapses with interneurons as early as postnatal day (p) 4-5 [12]. Concomitantly, the incidence of interneuron apoptosis increases drastically after p5 and reaches its peak at p7 [7]. Right after, the interneuron-OPC connectivity accelerates till p10, which drives an immediate oligodendrocyte (OL) boom and subsequent interneuron myelination via OPC-$GABA_A$ and $GABA_B$ receptors [13-15]. Despite relatively short axons, myelination of cortical interneurons (mainly parvalbumin (PV)$^+$ fast-spiking interneurons, [16]) contributes significantly to the fine-tuning of local activity in the medial prefrontal cortex (mPFC) and cortex, ensuring proper performance of behavior [4, 17]. However, how the early communication between interneuron and OPC affect inhibitory network activity in the mPFC still remains unknown.

To assess the function of OPC $GABA_B$Rs for mPFC inhibition, we conditionally deleted the $GABA_B$ receptor subunit *gabbr1* selectively in OPCs at the end of the first postnatal week. By focusing on the medial prefrontal cortex (mPFC), we found that OPCs shape the inhibitory network by $GABA_B$ receptor and the cytokine tumor necrosis factor-like weak inducer of apoptosis (TWEAK) signaling pathways, thereby finetuning interneuron cell death and survival as well as its myelination onset. The functional and morphological changes of PV$^+$ interneurons observed in the adult mPFC of mutant mice generate a reduction of the inhibitory tone, which finally leads to impaired cognition and social behavior.

 Results

 GABA$_B$Rs of OPCs are required for oligodendrogenesis in the mPFC

To assess the function of GABA$_B$Rs for OPC-interneuron communication, we

generated Tg*(NG2-CreER^{T2}):GABA$_{B1}$R$^{fl/fl}$* mice (Fig. 1a) to conditionally knockout

(cKO) functional GABA$_B$R from OPCs and their progeny. We induced the cKO at

postnatal day 7 and 8 (p7/8) before the onset oligodendrocyte formation at p10-11 [13].

When analyzed at the age of 9 weeks (w), about 80 % of OPCs (platelet-derived

growth factor receptor α$^+$ (PDGFRα$^+$, Pα+))were found recombined (Pα$^+$tdT$^+$/Pα$^+$) in

the medial prefrontal cortex (mPFC), based on tdTomato (tdT) gene expression

(Fig. 1b, c).

To confirm and quantify the deletion of GABA$_B$R in OPCs, we performed Western blot

analysis with OPCs purified by magnetic-activated cell sorting (MACs-OPCs). In cKO

OPCs, GABA$_B$R expression was reduced by about 64 % compared with controls (ctl)

(Fig. 1d). Considering the substantial purity of MACS-sorted OPCs (85 %, data not

shown) and recombination efficiency (80 %), we concluded that a vast majority of

GABA$_B$Rs had been ablated from OPCs.

Next, we focused on the oligodendrocyte lineage and evaluated the contribution of

GABA$_B$Rs for OPC differentiation in the mPFC by immunostaining of PDGFRα and

adenomatous polyposis coli clone 1 (CC1) (Fig. 1e), established markers for OPCs

and oligodendrocytes, respectively. In the cKO mice, the density of OPCs

(PDGFRα$^+$Olig2$^+$) did not change (Fig. 1f), while that of mature oligodendrocytes

(CC1$^+$Olig2$^+$) was strongly reduced to 74 % (Fig. 1g), indicative of decreased OPC

differentiation to oligodendrocytes (Fig. 1h). Notably, in the corpus callosum the

densities of oligodendrocytes and OPCs, as well as their relative proportions did not

differ between ctl and cKO mice (Fig.1 f-h, Extended Data Fig. 1). These data suggest

that OPC-GABA$_B$Rs are required for OPC differentiation in the grey matter mPFC, but

not in the white matter corpus callosum.

We further analyzed structural aspects of myelination which is important for axonal

conductivity, i.e. the lengths of nodes (the gaps between paired contactin-associated

protein-positive (Caspr$^+$) segments) and paranodes (single Caspr$^+$ segments) by

immunostaining (Fig. 1i). In cKO mPFC, the paranodal length was increased (Fig. 1j)

while node length and density remained stable (Extended Data Fig. 2). In parallel, we

observed a reduction of myelin basic protein (MBP) expression at the mRNA

(Extended Data Fig. 2) and protein levels (Fig. 1k) in cKO mPFC, while the other

myelin protein, proteolipid protein (PLP) mRNA levels were not changed (Extended

Data Fig. 2). In the corpus callosum, however, myelination was unperturbed as

indicated by unchanged paranode length (Extended Data Fig. 2), node density

(Extended Data Fig. 2) and MBP expression (Fig. 1k). Therefore, our data show that

GABA$_B$Rs of OPCs promote differentiation and myelination in the mPFC, but do not

affect the corpus callosum.

Since OPCs continuously differentiate into oligodendrocytes [18-20], the cKO mice will

also generate oligodendrocytes lacking GABA$_B$Rs. To selectively target only

oligodendrocytes, we used Tg*(PLP-Cre^{ERT2}):GABA$_{B1}$R$^{fl/fl}$* mice (Extended Data Fig. 3) .

The gene deletion was induced by tamoxifen injections at p7/8 or at 4w. At the age of

9w, we did not observe differences in the OPC and oligodendrocyte population

between ctl and cKO mice (Extended Data Fig. 3), neither in the PFC nor in the corpus

callosum (Extended Data Fig. 3).

In summary, our data strongly suggest that GABA$_B$R is specifically required for early

OPC differentiation and subsequent myelination in the mPFC, but not in the corpus

callosum.

GABA$_B$Rs of OPCs promote the myelination of interneurons in the mPFC

Since callosal axons are primarily excitatory [21] and axons of inhibitory interneurons are

largely restricted to cortical areas of the same hemisphere (including mPFC, Fig.1l)

(Extended Data Fig. 4), we hypothesized that the observed myelin deficits were

restricted to interneurons. Required by their fast-spiking activity, parvalbumin (PV)$^+$

interneurons are the most abundantly myelinated interneurons [16, 22-24] of the cortex.

These neurons are most prominent at the layer 4 and 5 with majority of their axons

projecting to the layer 2/3 and 5 [25] (Fig. 1l, Extended Data Fig. 4). To evaluate their

myelination, we performed PV and MBP double immunostaining (Fig. 1l-n) and

analyzed the volume of myelin sheaths, by volume rendering of MBP immunolabel

covering PV$^+$ axons at the cortical layers 2/3 of mPFC. Indeed, the volume of myelin

sheaths wrapping PV neurons was reduced by 60 % in the cKO mPFC (Fig. 1n).

However, the myelination of excitatory neurons (in the corpus callosum) was not

affected as the g-ratio of myelinated axons and conduction velocity of compound action

potential did not differ between ctl and cKO mice (Extended Data Fig. 2). Thus, we

concluded that GABA$_B$R of OPCs are preferentially involved in the myelination of

inhibitory rather than excitatory neurons.

Postnatal deletion of GABA$_B$Rs in OPCs causes enhanced interneuron density in the

adult, but with reduced GABA release

While investigating the interneuronal myelination, we recognized that early ablation of

GABA$_B$R in OPCs (at p7/8, Fig. 2a) was associated with a 30 % increase of interneuron

density (10.4 vs 13.3 for ctl and cKO respectively) in the cortex as revealed by

immunostaining for PV (Fig. 2b). However, in the mPFC this increase did not result in

a concomitant increase of inhibitory input (Fig. 2c, d). In contrast, the frequency of

spontaneous inhibitory postsynaptic currents (sIPSCs) of pyramidal neurons in the

layer V of mPFC, where interneurons are abundantly located, was reduced by 40 %,

while the amplitude of sIPSCs remained unaffected in the mutant mPFC as well as the

firing rate and amplitude of excitatory postsynaptic currents (EPSCs).

Since neuronal activity can regulate myelination [22, 26], we asked whether the reduced

inhibitory firing rate contributed to the observed molecular and structural changes of

the myelin at interneuronal axons in the mutant mPFC. For that purpose, we further

analyzed the cKO mPFC of p10 mice (Fig. 2e) at the onset of myelination and when

interneuron-OPC communication peaks [13]. After induction of the cKO at p7/8, we

assessed the density of the vesicular GABA transporter (vGAT) close to OPC surfaces

as a quantitative readout of inhibitory presynaptic terminals by combined vGAT and

PDGFRα immunostaining [27] (Fig. 2f-i, Extended Data Fig. 5). We classified the vGAT

puncta into two subgroups based on the distance between vGAT staining and OPC

surface: '<20 nm' as vesicles targeting on postsynaptic OPCs (magenta), and '>20 nm'

as vesicles targeting on other cells (grey), assuming a width of about 20 nm for the

synaptic cleft [28]. And indeed, already at p10, i.e. only 3 days after the first tamoxifen

injection, the density and the volume of vGAT immunopuncta were reduced were

reduced to 75 % and 80 % in the cKO mPFC (Fig. 2f-i, Extended Data Fig. 5), including

a smaller volume of vGAT (Fig. 2h) and less puncta (Fig. 2i) at the OPC surface in

cKO mPFC, which can explain the attenuated firing rate of the interneurons in the adult.

These results indicate that interneurons exhibit lower activity and transmit less

GABAergic signals to OPCs in the cKO mPFC.

To test whether the reduced inhibitory tone affecting OPC differentiation specifically

correlates with the onset of myelination, we compared the OPC and oligodendrocyte

populations at p10 and p14 (Fig. 2j-m, Extended Data Fig. 6). At p10, the densities of

OPC and oligodendrocyte populations were still comparable in the cKO mPFC to

control mPFC, although the interneurons already expressed less vGAT (Fig. 2f-h).

However, at p14, the differentiation and cell numbers of oligodendrocytes dropped

strongly in the mutant mPFC (Fig. 2l, Extended Data Fig. 6), as did the MBP expression (Fig. 2m). Again, in the corpus callosum, OPC and oligodendrocyte population were not affected by the deletion of GABA$_B$Rs (Extended Data Fig. 1).

When we induced GABA$_B$R deletion in the young adult mice at the age of 4 weeks, both in the mPFC and corpus callosum, the density of oligodendrocyte as well as the MBP expression were not affected in the cKO mice (Extended Data Fig. 7).

In summary, our data strongly suggest that during early development of the mPFC deletion of GABA$_B$R in OPCs affects the inhibitory tone of interneurons, feeding back onto OPC differentiation and interneuron myelination.

OPCs induce interneuron apoptosis via GABA$_B$R-TWEAK signaling

As we had seen a surplus of interneurons in the mutant mPFC which could explain the abnormal neuronal activity [8, 29], we tested whether the interneuron numbers would be accurately controlled by programmed cell death early during development. After induction of GABA$_B$R in OPCs at p7/8, the extent of interneuron apoptosis was determined at p10 and p14 by immunostaining for PV and cleaved caspase-3 (CC-3, Extended Data Fig. 8), a well-established marker for apoptosis. As expected by the earlier experiments, we found 50 % reduction of PV$^+$/CC-3$^+$ interneurons at p10 suggestive of strongly reduced apoptosis of PV$^+$ interneurons in the mutant mPFC (Extended Data Fig. 8). Obviously, OPCs receive GABAergic input through GABA$_B$R and send back pro-apoptotic signals to interneurons.

To confirm the selective cell death of interneurons, we induced the cKO as early as p1 and p2 and compared the apoptosis of inhibitory neurons with that of excitatory ones at p5 and p7 (Fig. 3a). Interneuron apoptosis persists from p1 till p15 with a peak at p7 [7], while the time window for excitatory neuron apoptosis is rather narrow, between p2-5 [30]. Apoptosis of total interneurons (GAD67$^+$CC-3$^+$) as well as PV interneurons

(PV$^+$CC-3$^+$ left panel in Fig. 3b) was significantly reduced in cKO mPFC at p5 and p7

(Fig. 3c), while that of the excitatory neurons (CTIP$^+$CC-3$^+$, right panel in Fig. 3b;

TBR1$^+$CC-3$^+$) did not change at both time points (Fig. 3c, Extended Data Fig. 9).

Therefore, we concluded that OPCs send pro-apoptotic signals back to interneurons

at the early postnatal weeks.

To molecularly identify the critical apoptotic factor released by OPCs, we first tested

the expression levels of several peptides which had been associated with cell survival

or cell death in ctl and cKO mPFC (Extended Data Fig. 10). Gene deletion was induced

at p1/2, and mPFC was collected at p7, the peak of interneuron apoptosis (Extended

Data Fig. 10). We found only the expression of the cytokine tumor necrosis factor-like

weak inducer of apoptosis (TWEAK, also known as Apo3l or TNF superfamily member

12, TNFSF12) to be diminished about 30 % in cKO mPFC (Fig. 3d, Extended Data

Fig. 10). TWEAK induces an apoptotic cell death after binding to its cognate TWEAK

receptor (TWEAKR; TNF Receptor Superfamily Member 12A, TNFRSF12A; also

known as FN14) [31]. Subsequently, we tested MACS-sorted OPCs (collected from cKO

and ctl cortices at p7) (Fig. 3d) as well as Oli-neu cells (an OPC cell line) treated with

GABA$_B$R antagonist CGP 55845 for TWEAK. Also here, we observed TWEAK

downregulation (Extended Data Fig. 10). Three independent experiments, including

genetic deletion or pharmacological block of OPC GABA$_B$R decreased TWEAK

expression (Fig. 3d, Extended Data Fig. 10).

To confirm that OPC-derived TWEAK can indeed induce neuronal apoptosis, we

treated primary cortical neurons with conditioned medium obtained from OPC Oli-neu

cultures (Fig. 3e). A selective TWEAKR antagonist, L524-0366 was added to the

conditioned medium at 20 µM to block the binding of TWEAK on interneurons

(Fig. 3e, f). While the conditioned medium from the OPC cell line increased interneuron

apoptosis (Fig. 3g, h), co-treatment with the TWEAKR antagonist prevented the cell

death. These results further substantiate that OPCs induces interneuron apoptosis via

TWEAK secretion.

In summary, our results demonstrate that OPCs induce interneuron apoptosis by

releasing TWEAK upon GABA$_B$R activation. Blocking this pathway prevents the

apoptosis of interneurons. The concomitant dysregulation of interneuron density

results in a reduced inhibitory GABAergic tone and altered myelination in the mPFC.

Interrupted OPC-interneuron communication after GABA$_B$R loss in OPCs during

development generates cognitive impairment in adulthood

The mPFC is responsible for cognitive processes and their regulatory finetuning is

guaranteed by the E/I balance [32]. To test whether the selectively reduced interneuron

apoptosis and the associated structural myelin alterations can generate defects in

neural circuits and cognition, we further performed *in vivo* investigations using

electrophysiology and behavioral analysis in 9w old animals after gene ablation was

induced at P7/8 (Fig. 4a).

The patch-clamp recordings described above had revealed a reduced firing rate of

interneurons suggesting an imbalance of E/I in the mutant mPFC (Fig. 2c, d). Impaired

cortical   network   activity   of   the   mutant   mice   was   also   detected   in

electroencephalographical (EEG) recordings. The power of the gamma wave,

especially the lower gamma wave band (30-50 Hz) was slightly, but significantly

decreased by about 10% in cKO mice (Fig. 4b).

To evaluate the impact of circuit dysregulation in the mutant mPFC for the living animal,

we challenged mice for their cognitive performance employing tasks of social cognitive

behavior which had been associated with proper mPFC function, i.e. social novelty and

new object recognition [33]. In addition, nest building behavior is often introduced as a

gauge of general well-being of the mice and also being an early sign of cognition

decline [34]. We observed that the mutant mice built significantly poorer nests with less

than half score of the control animal nests, indicating putative cognition deficits of the

mutant mice (Fig. 4c). The three-chamber test was used to assess social cognition by

analyzing mice for a form of general sociability and their interest in social novelty

(Fig. 4d). First, the three-chamber test was used to monitor general social behavior,

i.e. preference of other mice over objects. Both, ctl and cKO mice exhibited more

interest in the mouse rather than the object during the first test phase, indicating no

disorder in sociability of the mutant mice (Fig. 4e, Extended Data Fig. 11). At the

second phase, a social novelty preference was assessed. Control animals investigated

the unfamiliar mouse more than the familiar one, while the cKO mice exhibited no novel

social preference as they spent similar time in both chambers (Fig. 4f, Extended Data

Fig. 11). Employing a new object recognition test (Fig. 4g), we again observed the

cognition defect of cKO mice. Control animals were able to distinguish the new object

replaced after a 40 min break from the habituation, while the cKO mice showed similar

interest to both of the old and new objects (right panel in Fig. 4g-i). Open field test

showed that the motor activity of the cKO mice (in terms of the speed and total distance)

was not affected (Fig. 4j-l).

Hence, we conclude that $GABA_BRs$ of OPCs are essential for cognition by modifying

inhibitory circuits in the mPFC.

Discussion

A balance of excitation and inhibition (E/I) in the medial prefrontal cortex (mPFC) is of key importance for mammalian cognition. Establishing correct cell densities and subsequent myelination have been identified as pivotal elements during CNS development [6, 17]. This has been well documented for fast-spiking parvalbumin-positive (PV+) interneurons. Here, we highlight OPCs as indispensable regulators of interneuron population and myelination in the mPFC by providing a molecular explanation and demonstrating how impaired bidirectional OPC-interneuron signaling affects social cognitive behavior. OPCs at the first two postnatal weeks release TWEAK to regulate interneuron apoptosis. Genetic ablation of GABA$_B$R in OPCs at the first postnatal week promoted interneuron survival. However, these neurons displayed hypoactivity, less contacts with OPCs during development and deteriorated myelin structures in the adulthood. Notably, these mutant mice exhibited severe cognitive defects in their social behavior.

During development, large populations of immature interneurons invade the cortex where their cell density is adjusted by programmed cell death [9, 10]. For assuring their survival, these neurons form connections with adjacent cells and receive retrograde signals from their targets [10, 11]. Interneuron apoptosis drastically increases at p7 [7], shortly after formation of interneuron-OPC synapses at p4-5 [12]. Based on this established knowledge, we hypothesized that OPCs could exert an apoptotic impact to their presynaptic partners. Indeed, our data provide strong evidence that, starting from postnatal day 5 on, OPCs elicit an apoptotic cascade of adjacent interneurons to adjust their density. In OPCs, the regulatory process employs a GABA$_B$R-TWEAK signaling pathway. TWEAK, TNF super family member 12 (TNFSF12), is TNF-like weak apoptotic factor. Membrane-bound and cleaved isoforms of TWEAK are both able to induce cell apoptosis by binding to the TWEAK receptor (TWEAKR, also known as FN14) on the neuronal side [31, 35, 36]. According to our data GABA$_B$R-TWEAK

signaling preferentially affects interneurons, suggesting this 'kill me' signal acting

highly localized. Indeed, TWEAK is specifically recruited to synapses where

TWEAKRs are expressed [37]. OPCs directly sense GABAergic signals at their

interneuron-OPC synapse. Subsequent signaling triggers TWEAK expression and

translocation to the OPC surface, where it could directly bind interneuronal TWEAKRs.

Alternatively, the TWEAK ectodomain could be proteolytically cleaved and act as a

soluble factor. However, this scenario could also happen at soma-somatic contact sites

between OPCs and interneurons [38, 39]. The regulatory impact appears to be highly

specific since the OPC-induced interneuron apoptosis was restricted to the first two

postnatal weeks, a well-established time window of interneuron apoptosis [7, 30]. This

also explains why the cKO induction at the age of 4w did not impact OPC differentiation

and myelination, although $GABA_BR$ expression peaks in OPCs at 4w (data not shown).

Please note, instead of a gradual decline, we found a slight increase of PV^+CC3^+ cells

at p14 compared to p10, probably due to the age-dependent increase of PV expression

described for fast spiking interneurons [40].

Neuronal overpopulation is often accompanied with suppressed activity [6, 8]. In the

OPC-$GABA_BR$ cKO mice, we observed excessive number of PV^+ interneurons (about

30 % increase) while attenuated vGAT puncta (reduced to 80 %), suggesting single

PV^+ interneuron exhibits suppressed activity (about 60% ≈ 80 %÷130 %). This was

also shown by the sIPSP recording which was decreased to 60 % in the cKO mPFC.

Differentiation of oligodendrocytes and subsequent myelination are strongly affected

by neuronal activity [22, 26, 41]. Indeed, we observed a suppression of the inhibitory tone

in the mPFC at p10 followed by a change in oligodendrocyte density detected at p14.

Concomitantly, the oligodendrocytes formed less myelin (only 40 % of control scenario)

around the interneuronal axons and altered paranodal structures in the adult mPFC

which are essential for precise action potential propagation. Such reduction is

attributed to the restrained PV^+ neuron activity (60 %) and oligodendrocyte population

(74 %) (60 %×74 % ≈ 40 %). Regardless, the morphological deterioration was only observed in the mPFC, but not in the corpus callosum composed axons of excitatory neurons only [21]. In line with these neuron-type specific observations, the qRT-PCR results indicated reduced levels of MBP, but not PLP mRNA expression in the mutant mPFC, both at p14 and 9w of age. Axons of interneurons and excitatory neurons are different in respect to the protein content of their myelin sheaths. MBP is preferentially expressed in the myelin of inhibitory axons, while PLP is more prominent in excitatory axons PLP [16].

An early loss of the GABA$_A$R γ2 subunit in OPCs reduced the firing rate of presynaptic PV$^+$ interneurons and caused a similar myelin defect of longer nodal structures in PV$^+$ neurons [4], the initially very different signal pathways of both, GABA$_A$R and GABA$_B$R, have to merge to affect myelin gene expression and OPC-axon contacts similarly. The GABA$_A$R signaling between interneurons and OPCs is relatively well established [4, 15, 42, 43]. E.g., the γ2 subunit of GABA$_A$R contributes to the maintenance of the OPC, but has no impact on oligodendrocyte formation [15]. However, the role of GABA$_B$Rs for OPC differentiation is complex. We observed that GABA$_B$R was essential for oligodendrogenesis at the early development but not at later stages. Similary, an *in vitro* study showed that activation of GABA$_B$R by the selective agonist baclofen could stimulate the differentiation of cortical OPCs prepared from p0-2 pups [14]. Since OPCs are heterogeneous with age [44], GABA$_B$R of OPCs could contribute to the developmental diversity.

Obviously, the GABAergic signaling of OPCs plays pivotal roles for the myelination of interneurons, which ensures precise finetuning of the local neural circuitry [45]. As seen in chandelier cells [6], excessive number of interneurons in the OPC-GABA$_B$R cKO mice exhibited suppressed activity, and reduced myelination in the mPFC. All these physiological and morphological abnormalities certainly contributed to cognition impairment. A proper E/I ratio in the prefrontal cortex, especially early E/I balance in

mPFC, is extremely important for social cognition [32, 46]. When the pyramidal neuron

activity was enhanced in the mPFC at early p7-11 caused a severe cognitive

dysfunction in the adulthood [46]. In addition, despite relatively short axons, myelination

of cortical interneurons (mainly PV$^+$ neurons [16]) contributes significantly to the

finetuning of local activity in the mPFC and cortex, insuring proper behavior

performance [4, 6, 17].

In conclusion, our study demonstrates that OPCs regulate inhibition in the mPFC via

GABA$_B$R/TWEAK signaling. During development, OPCs determine the density of

interneurons by adjusting their apoptosis. Subsequently, correct myelination of

interneurons, as an essential component of the network activity in the mPFC,

determines the cognition and social behavior.

References

1. Douglas, R.J. & Martin, K.A. Inhibition in cortical circuits. *Curr Biol* **19**, R398-402 (2009).
2. Bai, X., Kirchhoff, F. & Scheller, A. Oligodendroglial GABAergic Signaling: More Than Inhibition! *Neurosci Bull* (2021).
3. Marín, O. Interneuron dysfunction in psychiatric disorders. *Nat Rev Neurosci* **13**, 107-120 (2012).
4. Benamer, N., Vidal, M., Balia, M. & Angulo, M.C. Myelination of parvalbumin interneurons shapes the function of cortical sensory inhibitory circuits. *Nat Commun* **11**, 5151 (2020).
5. Ferguson, B.R. & Gao, W.J. PV Interneurons: Critical Regulators of E/I Balance for Prefrontal Cortex-Dependent Behavior and Psychiatric Disorders. *Front Neural Circuits* **12**, 37 (2018).
6. Wang, B.S.*, et al.* Retinal and Callosal Activity-Dependent Chandelier Cell Elimination Shapes Binocularity in Primary Visual Cortex. *Neuron* **109**, 502-515.e507 (2021).
7. Southwell, D.G.*, et al.* Intrinsically determined cell death of developing cortical interneurons. *Nature* **491**, 109-113 (2012).
8. Orduz, D.*, et al.* Developmental cell death regulates lineage-related interneuron-oligodendroglia functional clusters and oligodendrocyte homeostasis. *Nat Commun* **10**, 4249 (2019).
9. Lim, L., Mi, D., Llorca, A. & Marín, O. Development and Functional Diversification of Cortical Interneurons. *Neuron* **100**, 294-313 (2018).
10. Davies, A.M. Regulation of neuronal survival and death by extracellular signals during development. *EMBO J* **22**, 2537-2545 (2003).
11. Oppenheim, R.W. Cell death during development of the nervous system. *Annu Rev Neurosci* **14**, 453-501 (1991).
12. Vélez-Fort, M., Maldonado, P.P., Butt, A.M., Audinat, E. & Angulo, M.C. Postnatal switch from synaptic to extrasynaptic transmission between interneurons and NG2 cells. *J Neurosci* **30**, 6921-6929 (2010).
13. Orduz, D.*, et al.* Interneurons and oligodendrocyte progenitors form a structured synaptic network in the developing neocortex. *Elife* **4** (2015).
14. Serrano-Regal, M.P.*, et al.* Oligodendrocyte Differentiation and Myelination Is Potentiated via GABA. *Neuroscience* **439**, 163-180 (2020).
15. Balia, M., Benamer, N. & Angulo, M.C. A specific GABAergic synapse onto oligodendrocyte precursors does not regulate cortical oligodendrogenesis. *Glia* **65**, 1821-1832 (2017).
16. Micheva, K.D.*, et al.* A large fraction of neocortical myelin ensheathes axons of local inhibitory neurons. *Elife* **5** (2016).
17. Maas, D.A.*, et al.* Interneuron hypomyelination is associated with cognitive inflexibility in a rat model of schizophrenia. *Nat Commun* **11**, 2329 (2020).
18. Huang, W.*, et al.* Novel NG2-CreERT2 knock-in mice demonstrate heterogeneous differentiation potential of NG2 glia during development. *Glia* **62**, 896-913 (2014).
19. Kang, S.H., Fukaya, M., Yang, J.K., Rothstein, J.D. & Bergles, D.E. NG2+ CNS glial progenitors remain committed to the oligodendrocyte lineage in postnatal life and following neurodegeneration. *Neuron* **68**, 668-681 (2010).
20. Zhu, X.*, et al.* Age-dependent fate and lineage restriction of single NG2 cells. *Development* **138**, 745-753 (2011).
21. Fame, R.M., MacDonald, J.L. & Macklis, J.D. Development, specification, and diversity of callosal projection neurons. *Trends Neurosci* **34**, 41-50 (2011).
22. Stedehouder, J.*, et al.* Fast-spiking Parvalbumin Interneurons are Frequently Myelinated in the Cerebral Cortex of Mice and Humans. *Cereb Cortex* **27**, 5001-5013 (2017).

23.   Stedehouder, J.*, et al.* Local axonal morphology guides the topography of interneuron
myelination in mouse and human neocortex. *Elife* **8** (2019).
24.   Zonouzi, M.*, et al.* Individual Oligodendrocytes Show Bias for Inhibitory Axons in the
Neocortex. *Cell Rep* **27**, 2799-2808.e2793 (2019).
25.   Scala, F.*, et al.* Layer 4 of mouse neocortex differs in cell types and circuit organization
between sensory areas. *Nat Commun* **10**, 4174 (2019).
26.   Mitew, S.*, et al.* Pharmacogenetic stimulation of neuronal activity increases
myelination in an axon-specific manner. *Nat Commun* **9**, 306 (2018).
27.   Chaudhry, F.A.*, et al.* The vesicular GABA transporter, VGAT, localizes to synaptic
vesicles in sets of glycinergic as well as GABAergic neurons. *J Neurosci* **18**, 9733-9750 (1998).
28.   Dani, A., Huang, B., Bergan, J., Dulac, C. & Zhuang, X. Superresolution imaging of
chemical synapses in the brain. *Neuron* **68**, 843-856 (2010).
29.   Fang, W.Q.*, et al.* Overproduction of upper-layer neurons in the neocortex leads to
autism-like features in mice. *Cell Rep* **9**, 1635-1643 (2014).
30.   Wong, F.K.*, et al.* Pyramidal cell regulation of interneuron survival sculpts cortical
networks. *Nature* **557**, 668-673 (2018).
31.   Chicheportiche, Y.*, et al.* TWEAK, a new secreted ligand in the tumor necrosis factor
family that weakly induces apoptosis. *J Biol Chem* **272**, 32401-32410 (1997).
32.   Grossmann, T. The role of medial prefrontal cortex in early social cognition. *Front Hum*
Neurosci **7**, 340 (2013).
33.   Bicks, L.K., Koike, H., Akbarian, S. & Morishita, H. Prefrontal Cortex and Social
Cognition in Mouse and Man. *Front Psychol* **6**, 1805 (2015).
34.   Neely, C.L.C., Pedemonte, K.A., Boggs, K.N. & Flinn, J.M. Nest Building Behavior as an
Early Indicator of Behavioral Deficits in Mice. *J Vis Exp* (2019).
35.   Winkles, J.A. The TWEAK-Fn14 cytokine-receptor axis: discovery, biology and
therapeutic targeting. *Nat Rev Drug Discov* **7**, 411-425 (2008).
36.   Brown, S.A., Ghosh, A. & Winkles, J.A. Full-length, membrane-anchored TWEAK can
function as a juxtacrine signaling molecule and activate the NF-kappaB pathway. *J Biol Chem*
285, 17432-17441 (2010).
37.   Cheadle, L.*, et al.* Sensory Experience Engages Microglia to Shape Neural Connectivity
through a Non-Phagocytic Mechanism. *Neuron* **108**, 451-468.e459 (2020).
38.   Mangin, J.M., Kunze, A., Chittajallu, R. & Gallo, V. Satellite NG2 progenitor cells share
common glutamatergic inputs with associated interneurons in the mouse dentate gyrus. *J*
Neurosci **28**, 7610-7623 (2008).
39.   Boulanger, J.J. & Messier, C. Oligodendrocyte progenitor cells are paired with GABA
neurons in the mouse dorsal cortex: Unbiased stereological analysis. *Neuroscience* **362**, 127-
140 (2017).
40.   Larsen, R., Proue, A., Scott, E.P., Christiansen, M. & Nakagawa, Y. The Thalamus
Regulates Retinoic Acid Signaling and Development of Parvalbumin Interneurons in Postnatal
Mouse Prefrontal Cortex. *eNeuro* **6** (2019).
41.   Nagy, B., Hovhannisyan, A., Barzan, R., Chen, T.J. & Kukley, M. Different patterns of
neuronal activity trigger distinct responses of oligodendrocyte precursor cells in the corpus
callosum. *PLoS Biol* **15**, e2001993 (2017).
42.   Balia, M.*, et al.* Postnatal down-regulation of the GABAA receptor γ2 subunit in
neocortical NG2 cells accompanies synaptic-to-extrasynaptic switch in the GABAergic
transmission mode. *Cereb Cortex* **25**, 1114-1123 (2015).
43.   Passlick, S.*, et al.* Expression of the γ2-subunit distinguishes synaptic and
extrasynaptic GABA(A) receptors in NG2 cells of the hippocampus. *J Neurosci* **33**, 12030-12040
(2013).
44.   Spitzer, S.O.*, et al.* Oligodendrocyte Progenitor Cells Become Regionally Diverse and
Heterogeneous with Age. *Neuron* **101**, 459-471.e455 (2019).

45. Arancibia-Cárcamo, I.L*., et al.* Node of Ranvier length as a potential regulator of
myelinated axon conduction speed. *Elife* **6** (2017).
46. Bitzenhofer, S.H., Pöpplau, J.A., Chini, M., Marquardt, A. & Hanganu-Opatz, I.L. A
transient developmental increase in prefrontal activity alters network maturation and causes
cognitive dysfunction in adult mice. *Neuron* (2021).

Acknowledgement

We thank Daniel Schauenburg for excellent animal husbandry and Frank Rhode,

Alexander Grissmer and Samantha Bechet for experimental assistance. We are also

grateful to Prof. Jaqueline Trotter (University of Mainz, Mainz, Germany) for providing

the Oli-neu cell line and Prof. Leda Dimou (University of Ulm, Ulm, Germany) for

sharing MACs sorting protocols of OPCs. This work was supported by grants from the

Deutsche Forschungsgemeinschaft (SPP 1757, SFB 894 to FK; FOR2289 to AS, FK,

SPP1757 Young Investigator grant to XB), EraNet-Neuron BrIE to FK and

HOMFORexzellent2017 program to XB.

Author contributions

XB initiated the project. LF, XB, NZ, LC, NH, CL and AS performed experiments. LF,

XB, NZ, LC and RZ analyzed data. BB, WH. HC and CM provided materials. XB and

FK designed and supervised the study and wrote the manuscript with comments of the

other authors.

Competing interests

The authors declare no competing interests.

 Figure legends

Figure 1. Ablation of GABA_BRs in OPCs attenuates oligodendrocyte

differentiation and myelination of interneurons in the medial prefrontal cortex.

a, Genetic strategy and experimental schedule. *b,* Prefrontal cortex in coronal section

stained with PDGFRα. tdTomato$^+$ (tdT$^+$) cells indicate recombined cells. *b$_1$, b$_2$,*

magnified images of ctl and cKO medial prefrontal cortex (mPFC, highlighted in *b*). *c,*

Recombination efficiency of OPCs in ctl and cKO mPFC. (*ctl=4, cKO=5, unpaired t-*

test). *d,* Western blot analysis of GABA_BR subunit 1 in MACs sorted OPCs from cortex.

(*ctl=3, cKO=3, unpaired t-test*). *e,* Exemplary images of Olig2, PDGFRα and CC1

immunostaining in adult ctl and cKO mPFC. *f, g,* Quantification of OPC and

oligodendrocyte cell densities in mPFC and corpus callosum (cc). (mPFC: *ctl=7,*

cKO=8, cc: ctl=4, cKO=4, multiple t-tests). h, Quantification of the oligodendrocyte

proportion among the total lineage (mPFC: *ctl=4, cKO=4;* cc: *ctl=3, cKO=6, multiple t-*

tests). *i,* Immunostaining of Caspr and MBP in ctl and cKO mPFC. *j,* Quantitative

analysis of paranode length in mPFC (*ctl=4, cKO=4, multiple t-tests*). *k,* Western blot

analysis of MBP expression in mPFC and cc. (*ctl=4, cKO=3, multiple t-tests*). *l,*

Overview of coronal brain slices immunostained with parvalbumin (PV) and MBP

shows abundant PV$^+$ neurons are myelinated in mPFC. *m,* 3D reconstruction of

myelinated PV$^+$ axons in ctl and cKO mPFC using Imaris software. *n,* Relative volume

of MBP$^+$ myelin sheaths wrapping PV$^+$ axons (*normalized to ctl). (ctl=3, cKO=3,*

unpaired t-test). Scale bars in **b**=1 mm, **b$_2$, e**=10 μm, **i, m**=5 μm, **l**=200 μm.

Figure 2. In OPC-GABA_BR cKO mice impaired control of interneuronal density

and firing rate reduces oligodendrocyte differentiation and myelin gene

expression at the onset of myelination. *a,* Experimental schedule for *b-d*. *b,*

Immonostaining (left) and quantification (right) of PV$^+$ neurons in the mPFC at the age

of 9 weeks (*ctl=5, cK=9, unpaired t-test*). *c,* Exemplary traces of sIPSCs and sEPSCs

recorded at the layer V of mPFC. *d,* Quantification of frequency and amplitude of

sIPSC and sEPSC *(sIPSC: ctl=15 cells from 4 mice, cKO=15 cells from 5 mice; sEPSC:*

ctl=26 cells from 4 mice, cKO=21 cells from 5 mice, Mann Whitney test). **e,**

Experimental schedule for **f-m**. **f,** 3D reconstruction of OPCs and vGAT by PDGFRα

and vGAT immunostaining in ctl and cKO mPFC at p10 using Imaris software. vGAT

was classified into two subgroups: magenta vGATs have < 20nm distance from OPC

surface while grey vGATs have > 20nm. **g, h**, Density (**g**) and volume (**h**) of vGAT in

ctl and cKO mPFC. (*ctl=4, cKO=4; multiple t-tests*). **i,** Quantification of synaptic vGAT

(< 20nm) density per μm^2 of OPC surface. *(ctl=4, cKO=4, unpaired t-test).* **j,**

Immunostaining of OPCs and oligodendrocytes with PDGFRα, Olig2 and CC1 in p14

mPFC. **k, l,** Density of OPCs (**k**) and oligodendrocytes (**l**) in ctl and cKO mPFC at p10

and p14. (*p10 and p14: ctl=5, cKO=4; multiple t-tests*). **m,** Quantitative analysis of MBP

mRNA level in ctl and cKO mPFC at p14. (*ctl=4, cKO=4, unpaired t-test).* Scale bars

in **b**=50 µm, **f**=5 µm, **j**=10 µm.

Figure 3. During early CNS development the apoptosis of supernumerous

interneurons is mitigated upon impaired release of TWEAK from GABA$_B$R-

deficient OPC. a, Scheme of experimental plan. **b,** Representative images of

excitatory (CTIP) and inhibitory (PV$^+$) neurons co-stained with apoptotic marker CC-3

in p5 and p7 mPFC. **c,** Quantification of apoptotic neurons (CC-3$^+$) co-expressing

interneuron marker GAD67 and PV or excitatory neuronal marker CTIP and TBR1.

(GAD67: p5-ctl=3, p5-cKO=4; p7-ctl=3, p7-cKO=5) (PV: p5-ctl=5, p5-cKO=3; p7-ctl=3,

p7-cKO=5) (CTIP: p5-ctl=7, p5-cKO=4; p7-ctl=9, p7-cKO=3) (TBR1: p5-ctl=4, p5-

cKO=4; p7-ctl=4, p7-cKO=4) (multiple t-tests). **d,** Relative mRNA level of TWEAK in

the mPFC and MACs-OPCs from ctl and cKO mice *(tissue-ctl=4, tissue-cKO=6; MACs-*

ctl=4, MACs-cKO=3; multiple t-test). **e,** Experimental design for *in vitro* studies. **f,**

Immunolabeling of TWEAK receptor on 14 DIV PV$^+$ interneurons. **g,** Immunostaining

of apoptotic primary PV$^+$ neurons treated with conditional medium of Oli-neu cells co-

treated with or without TWEAKR antagonist L524-0366 (20 µM). **h,** Percentage of

apoptotic PV$^+$ neurons among total PV$^+$ neurons after the treatments *(n=3 independent*

experiments, one-way ANOVA, multiple comparison). Scale bars in **b**=5 µm, **f**=20 µm,

g=50 µm.

Figure 4.   In OPC-GABA$_B$R cKO mice perturbed electrical activity translates into

deficits of social behavior. *a,* Scheme of experimental plan. ***b,*** Representative

diagram of lower gamma band acquired by ECoG recordings. Violin plot shows

decreased gamma oscillation in the cKO mouse brain. (*ctl=24 from 4 mice, cKO=24*

from 4 mice, unpaired t-test). ***c,*** Impaired nest building ability of mutant mice. (*ctl=9,*

cKO=10, Mann Whitney test). ***d***, Scheme of three-chamber behavior test. ***e,***

Quantification of the sniffing time with mouse (***e***) or with unfamiliar mouse (***f***) among

total sniffing time (*ctl=9, cKO=9, unpaired t-tests*). ***g,*** Scheme of new object recognition

test. ***h,*** Assessment of object preference *(ctl=10, cKO=10)*. ***i,*** Percentage of sniffing

time with new object among total sniffing time. *(ctl=10, cKO=10, upaired t-test)*. ***j,***

Representative trajectory charts of the animals during the open field test. ***k, l,*** Both ctl

and cKO mice exhibited similar motor activities shown by the speed and distance

analysis. (*ctl=10, cKO=14, unpaired t-tests).*

 Supplementary figure legends

Supplementary Fig.1. OPC and oligodendrocyte density is not altered in the

corpus callosum of mutant mice. a, Scheme of experimental schedule (left panel)

and analyzed brain region in highlight, i.e. corpus callosum (cc, right panel). **b,** Corpus

callosal OPCs and oligodendrocytes of ctl and cKO mice were immunolabeled with

PDGFRα and CC1 respectively at the age of 2w and 9w. Olig2 was used as lineage

marker and tdTomato was indicator of recombined cells. **c,** OPC recombination

efficiency in ctl and cKO cc (*ctl=86.22±0.57 (n=3), cKO=76.17±3.10 (n=6)*, unpaired t-

test). **d, e,** Cell density of OPCs (PDGFRα$^+$) and oligodendrocytes (CC1$^+$) in cc of 2w-

and 9w-old mice were not changed. OPC: (*2w-ctl=77.02±4.21 (n=8), 2w-cKO=80±4.72*

(n=8), unpaired t-test) (*9w-ctl=30.01±3.68 (n=4), 9w-cKO=27.56±2.26 (n=4), unpaired*

t-test); oligodendrocytes (*2w-ctl=121.70±12.31 (n=8), 2w-cKO=97.62±10.54 (n=8),*

unpaired t-test) (*9w-ctl=296.90±32.83 (n=4), 9w-cKO=300.30±28.47 (n=4), unpaired*

t-test). **f,** Percentage of oligodendrocytes among total Olig2$^+$ cell population,

considered as OPC differentiation rate, was not changed at both ages (*2w-*

ctl=60.26±3.61(n=8), 2w-cKO=53.98±3.73 (n=8), unpaired t-test) (*9w-ctl=92.37±0.12*

(n=3), 9w-cKO=91.33±0.69 (n=6), unpaired t-test).

Supplementary Fig.2. Parameters of myelination remain largely unaltered in the

corpus callosum of mutant mice. a, Scheme of experimental schedule (left panel)

and analyzed brain region in highlight, i.e. mPFC and cc. **b,** Quantification of node

length of ctl and cKO mPFC. **c,** Paranode length is increased in the cKO mPFC. **d,**

Density of nodes remained same in the cKO mPFC. **e,** Relative mRNA level of MBP is

reduced in the cKO mPFC, while PLP mRNA is not affected (*ctl: n=4, cKO: n=3,*

unpaired t-test). **f,** Immunostaining of Caspr and MBP in cc of ctl and cKO mice. **g-i,**

Paranodal length and node density in ctl and cKO cc remained same. **j,** Electron

microscopic analysis of myelin ultrastructure in sagittal section of cc from ctl and cKO

mice. **k, l,** Quantification of g-ratios of axonal myelination shows unchanged

myelination in cKO cc. **m,** Compound action potential conduction velocity was not

altered indicated by patch clamp recordings. N1 indicates non-myelinated axonal

conduction velocity, while N2 is for myelinated axons.

Supplementary Fig.3. Ablation of GABA$_B$R in oligodendrocytes do not affect

formation and density of oligodendrocytes. a, Scheme of transgene construction.

b, Experimental schedule. **c,** Overview of CC1 immunostaining in a sagittal section of

9w old ctl mouse. Extensive tdTomato distribution shows a high recombination in

different brain regions of transgenic mouse. **d,** Magnified images from prefrontal cortex

(yellow box in **c**) and corpus callosum (white box in **c**). **e,** Quantification of

oligodendrocyte recombination efficiency in PFC were identical, while cKO mice

showed smaller recombination efficiency in cc. However, density of oligodendrocyte

cells did not differ between ctl and cKO mice in both PFC and cc. arrows: recombined

oligodendrocyte; triangles: non-recombined oligodendrocyte.

Supplementary Fig.4. Dense population of parvalbumin-expressing

interneurons in the adult mouse cortex. a, Overview of coronal brain slices

immunostained with parvalbumin and MBP shows abundant PV$^+$ neurons in the

somatosensory cortex. **b,** Magnified view (yellow box in **a**) shows the absence of PV$^+$

axons in the corpus callosum.

Supplementary Fig.5. At the onset of myelination, GABA$_B$R-deficient OPCs

receive less GABAergic input in the mPFC mutant mice. a, Experimental timeline

and analyzed brain region (mPFC) in highlight.**b,** Immunostaining of PDGFRα and

vGAT in the mPFC of p10 mice. **c,** Quantitative analysis of vGAT volume and the

distance to OPC surface (*ctl: n=4, cKO: n=4*). **d,** The volume of vGAT in cKO mPFC

was smaller than that in ctl group.

Supplementary Fig.6. The differentiation of GABA$_B$R-deficient OPCs is

suppressed in the mutant mPFC. a, Scheme of experimental schedule and analyzed

617 brain region (mPFC). **b,** Recombination efficiency of OPCs (% of $P\alpha^+tdt^+/P\alpha^+$) at p14

618 is identical in the ctl and cKO mPFC. **c,** Immunostaining and quantification of OPCs

619 incorporated with BrdU shows no change in OPC proliferation in the cKO mPFC at p14

620 (*ctl: n=4, cKO: n=5, unpaired t-test*). **d,** Immunostaining and quantification of

621 oligodendrocytes incorporated with BrdU shows decreased OPC differentiation in the

622 cKO mPFC at p14 (*ctl: n=4, cKO: n=5, unpaired t-test*). **e,** The portion of

623 oligodendrocytes among the oligodendrocyte lineage declined in cKO mPFC at p14

624 (*ctl: n=5, cKO: n=4, unpaired t-test*).

625 **Supplementary Fig.7. Controlled ablation of GABA$_B$R in OPCs of young adult**

626 **mice does not affect oligodendrogenesis. a,** Scheme of experimental schedule and

627 the analyzed brain regions (mPFC and cc). **b,** Immunostaining of OPCs (PDGFRα)

628 and oligodendrocyte (CC1) with Olig2 (lineage marker) at ctl and cKO mouse mPFC

629 and cc at the age of 9w. **c, d,** Cell densities of OPCs (PDGFRα$^+$) and oligodendrocytes

630 (CC1$^+$) of cKO animals were the same as those for ctl mice in both mPFC and cc.

631 mPFC. *(PDGFRα$^+$: (ctl=17.58±0.53 (n=4), cKO=18.69±1.58 (n=2), unpaired t-test));*

632 *(CC1$^+$: (ctl=9.05±1.38 (n=4), cKO=7.95±1.40 (n=2), unpaired t-test)); cc: (PDGFRα$^+$:*

633 *(ctl=35.38±2.25 (n=4), cKO=32.37±4.81 (n=2), unpaired t-test)); (CC1$^+$:*

634 *(ctl=311.81±43.39 (n=4), cKO=339.20±70.53 (n=2), unpaired t-test)).* **e,** The proportion

635 of oligodendrocyte among total lineage cells did not change, suggesting OPC

636 differentiation remained same in the mutant mouse mPFC and cc. mPFC:

637 *(ctl=30.49±5.13 (n=3), cKO=29.91±2.62 (n=2), unpaired t-test); cc: (ctl=89.52±1.30*

638 *(n=3), cKO=91.21±0.471 (n=2), unpaired t-test).* **f,** MBP expression is not altered in

639 the cKO mPFC at mRNA level (*ctl: n=4, cKO: n=3, unpaired t-test*).

640 **Supplementary Fig.8. The apoptosis of interneurons is mitigated in the mPFC of**

641 **mutant mice at p14. a,** Timeline of experiments and scheme of analyzed brain region

642 mPFC in highlight. **b,** Immunostaining of PV$^+$ neurons with apoptotic marker (cleaved

643 caspase-3) at p10 of ctl and cKO mPFC. **c,** Quantification of apoptotic PV$^+$ neurons at

p10 and p14 shows reduced PV$^+$ neuronal apoptosis in the cKO mPFC compared to

ctl (p10*: ctl=2.03±0.47 (n=6), cKO=0.92±0.34 (n=3), unpaired t-test;* p14:

ctl=3.70±1.21 (n=6), cKO=2.50±0.72 (n=9), unpaired t-test). **d,** Representative images

of apoptotic neurons at p14 of ctl and cKO mPFC. **e,** Quantification of apoptotic total

neurons (NeuN$^+$CC-3$^+$) at p14 *(ctl=2.76±0.42 (n=12), cKO=1.42±0.29 (n=9), unpaired*

t-test).

Supplementary Fig.9. Ablation of GABA$_B$R expression in OPCs does not affect

the apoptosis or survival of excitatory neurons. a, Scheme of experimental

schedule (left panel) and analyzed brain region in highlight. **b,** About 40 % of OPCs

are recombined (% of PDGFRα$^+$tdT$^+$/PDGFRα$^+$) at p7 both in the ctl and cKO mPFC.

c, d, Immunostainings and quantification of excitatory neurons (TBR1$^+$) with apoptotic

marker CC-3 in mPFC of ctl and cKO pups at p5 (**c**) and p7 (**d**) (p5: *ctl=1.64±0.39*

(n=4), cKO=1.74±0.24 (n=4), unpaired t-test; p10*: ctl=0.74±0.18 (n=4),*

cKO=0.67±0.16 (n=4), unpaired t-test).

Supplementary Fig.10. TWEAK expression in OPCs is regulated by GABA$_B$Rs. a,

Scheme of experimental design. **b,** Relative mRNA level of cytokines in the mPFC of

ctl and cKO mouse at p7. *(TNF-α: ctl=1±0.34 (n=4), cKO=1.23±0.36 (n=6), unpaired t-*

test); (TWEAK: ctl=1±0.10 (n=4), cKO=0.67±0.08 (n=6), unpaired t-test); (FasL:

ctl=1±0.57 (n=2), cKO=0.91±0.23 (n=4), unpaired t-test); (TNFSF7: ctl=1±1.00 (n=4),

cKO=0.93±0.67 (n=6), unpaired t-test); (Trp53: ctl=1±0.11 (n=4), cKO=1.14±0.08

(n=6), unpaired t-test); (NGF: ctl=1±0.24 (n=5), cKO=1.08±0.18 (n=6), unpaired t-test).

c, Western blot analysis of TWEAK expression in Oli-neu cells treated with

50 μM Baclofen at the presence or absence of 20 μM CGP 55845 (n=5 independent

experiments). The isoform in 36 KDa (blue box) was analyzed.

Supplementary Fig.11. OPC-GABA$_B$R cKO mice exhibit impaired social behavior.

a, b, Mutant mice stayed longer at the chamber of mouse than object, indicating that

sociability of the mutant mouse is not affected. **c, d,** Mutant mice spent similar time at

the chambers of familiar and unfamiliar mouse, suggesting impaired cognition of the

cKO mice (*ctl (n=9), cKO (n=9), paired t-test*).

Figure 1

Figure 2

Figure 3

a

Birth p1 2 5 7 Day ▼ Tam ▼ Analysis

b

Merge | CTIP | CC-3
ctl-p5

Merge | PV | CC-3
ctl-p7

cKO-p5

cKO-p7

c

Inhibitory neurons

GAD67⁺

p5: 0.027
p7: <0.001

PV⁺

p5: 0.011
p7: 0.014

Excitatory neurons

CTIP⁺

p5: 0.547
p7: 0.496

TBR1⁺

p5: 0.822
p7: 0.774

● ctl
● cKO

CC-3⁺ cells/1×10⁻²mm³

d

Relative Tweak mRNA level (normalized to ctls)

Tissue: 0.034
MACs-OPC: 0.068

e

Oli-neu

Conditioned medium (CM)

neuron
± L524-0366

f

TWEAKR | PV | merge

g

PV/CC-3

ctl+DMSO | CM+DMSO | CM+L524

h

% of PV⁺CC-3⁺/PV⁺

0.012
0.047

CM: − + +
L524: − − +

Figure 4

Supplementary figure 1

Supplementary figure 2

Supplementary figure 3

a

PLP promoter — CreERT2 x — GABA_B — ►loxP — Exon 7 — Exon 8 — ►loxP

x — Rosa26 — CAG — ►loxP — STOP — ►loxP — RATO

b

p7/8 — 4w — 9w — time

▼ 2 x tam
▼ 5 x tam
▼ analysis

c

CC1/ tdTomato

4w-9w   500 µm

d

CC1   tdT   Merge

PFC

cc   50 µm

e

PFC

% of CC1⁺tdT⁺/CC1⁺ cells

ctl
cKO

1w   4w

CC1⁺ cells/1x10⁻³ mm³

0.994   0.114

1w   4w

cc

% of CC1⁺tdT⁺/CC1⁺ cells

1w   4w

CC1⁺ cells/1x10⁻³ mm³

0.745   0.999

1w   4w

Supplementary figure 4

Supplementary figure 5

a

birth p7 p8 p10

▼ tam injection
▼ analysis

b

PDGFRα / vGAT

| ctl | cKO |

5 µm

c

vGAT volume (µm³)

Distance to OPC surface (µm)

d

% of volume distribution

● ctl
● cKO

Volume (µm³)

Supplementary figure 6

Supplementary figure 8

Supplementary figure 9

Supplementary figure 10

a

birth ▼p1 ▼p2 ▼p7

▼ tam injection
▼ analysis

b

Relative mRNA expression

0.675 TNF-α
0.034 TWEAK
0.865 FasL
0.936 TNFSF7
0.282 Trp-53
0.796 NGF

□ ctl
□ cKO

c

Bac ± CGP 55845

Oli-neu

| Bac | - | + | + | - | + | + |
| CGP | - | - | + | - | - | + |

KDa
48
35
25

Exp. 4   Exp. 5

KDa
48
35
25

TWEAK

α-tubulin   48

relative TWEAK level
(normalized to ctl)

0.254
0.043

● ctl
● Bac
● Bac + CGP

1
2 **Supplementary information for**
3

Impaired bidirectional communication between interneurons and oligodendrocyte

precursor cells affects cognitive behavior

Lipao Fang[1], Na Zhao[1], Laura C. Caudal[1], Renping Zhao[2], Ching-Hsin Lin[3], Hsin-Fang Chang[3],

Nadine Hainz[4], Carola Meier[4], Bernhard Bettler[5], Wenhui Huang[1], Anja Scheller[1], Frank

Kirchhoff[1,*], Xianshu Bai[1,*]

1. Molecular Physiology, CIPMM, University of Saarland, 66421 Homburg, Germany

2. Biophysics, CIPMM, University of Saarland, 66421 Homburg, Germany

3. Cellular Physiology, CIPMM, University of Saarland, 66421 Homburg, Germany

4. Department of Anatomy and Cell Biology, University of Saarland, 66421 Homburg, Germany

5. Department of Biomedicine, University of Basel, 4056 Basel, Switzerland

14
Correspondence to:
Prof. Dr. Frank Kirchhoff,
Molecular Physiology,
CIPMM, University of Saarland,
66421 Homburg, Germany
frank.kirchhoff@uks.eu
21
22 Dr. Xianshu Bai
23 Molecular Physiology,
24 CIPMM, University of Saarland,
25 66421 Homburg, Germany
26 xianshu.bai@uks.eu
27
28

Materials and Methods

Ethics statement

Animal husbandry and procedures were performed at the animal facility of CIPMM, University of

Saarland according to European and German guidelines for the welfare of experimental animals.

Animal experiments were approved by the Saarland state's "Landesamt für Gesundheit und

Verbraucherschutz" in Saarbrücken/Germany (animal license number: 65/2013, 12/2014,

34/2016, 36/2016 and 08/2021).

Animals

To conditionally knock out (cKO) GABA$_{B1}$ receptors in oligodendrocyte precursor cells (OPCs),

TgH(NG2-Cre^{ERT2})[1] mice were crossbred to GABA$_{B1}$$^{lox511/lox511}$ mice (flanking exon 7 and 8 of

gabbr1)[2]. Mice with genotypes of NG2$^{ct2/wt}$ x GABA$_{B1}$R$^{fl/fl}$ were used as cKO and the littermates

NG2$^{wt/wt}$ x GABA$_{B1}$R$^{fl/fl}$ or NG2$^{ct2/wt}$ x GABA$_{B1}$R$^{wt/wt}$ were controls after tamoxifen application. To

visualize the recombined cells, we crossbred the double transgenic mice with TgH(Rosa26-CAG-

^{fl}STOP^{fl}-tdTomato) (Rosa26-tdTomato) [3].

For oligodendrocyte specific deletion of GABA$_{B1}$R, we took advantage of TgN mice (PLP-Cre^{ERT2})

[4] crossbred with GABA$_{B1}$$^{lox511/lox511}$. Similarly, PLP$^{ct2/wt}$ x GABA$_{B1}$R$^{fl/fl}$ mice were used as cKO, while

PLP$^{wt/wt}$ x GABA$_{B1}$R$^{fl/fl}$ or PLP$^{ct2/wt}$ x GABA$_{B1}$R$^{wt/wt}$ were controls after tamoxifen application.

Mouse ages are indicated in the main text and figure legends. Behavioral tests were carried out

only with male mice at the age of 9 weeks, while both genders were used in the other experiments.

Tamoxifen administration

Tamoxifen was dissolved in Miglyol (3274, Caesar & Loretz GmbH) to a final concentration of

10 mg/ml. Tamoxifen was intraperitoneally injected to the mice depending on the body weight

(100 mg/kg body weight). The time points of injections are indicated in the figures. Only for the

pups treated at postnatal day 1 (p1) and 2, tamoxifen was injected to the lactating mother with

the same protocol [5]. For 4-week-old mice, tamoxifen was injected once per day for five

consecutive days [6].

Immunohistochemistry

Mice were perfused with PBS and 4 % paraformaldehyde (PFA). Mouse brains were post fixed

with 4 % PFA at 4°C overnight. Free floating brain slices (40 µm thickness) were cut in coronal or

sagittal sections using a Leica VT1000S vibratome. For immunocytochemistry, cells on coverslips

were fixed with ice cold 4 % PFA for 15 min. Slices or coverslips were incubated with blocking

solution containing 5 % horse serum and 0.5 % Triton X-100 at room temperature (RT) for 1 hour

(hr), followed by primary antibody incubation at 4 °C overnight and secondary antibody incubation

at RT for 2 hrs in blocking solution. Primary and secondary antibodies are listed in the

Supplementary table 1 and 2, respectively.

Magnetic cell separation (MACS) of OPCs

MACS sorting of OPCs was performed according to the manufacturer's instruction (Miltenyi

Biotec) with some modifications. Mice were perfused with cold Hank's balanced salt solution

without calcium and magnesium (HBSS, H6648, Gibco) and cortices were dissected in ice cold

HBSS. After the removal of debris (130-107-677, Miltenyi Biotec), cells were resuspended with

1 mL "re-expression medium" containing NeuroBrew-21 (1:50 in MACS neuro Medium) (130-093-

566 and 130-093-570, Milteny Biotec) and 200 mM L-glutamine (1:100, G7513, Sigma) at 37 °C

for 30 min. Cells were then incubated with Fc-receptor blocker for 10 min at 4 °C (provided with

CD140 microbeads kit), followed by a 15 min incubation with 10 µL microbeads mixture

composed of CD140 (130-101-502), NG2 (130-097-170) and O4 (130-096-670) in 1:1:1 at 4 °C.

For the purity assessment, cells were re-suspended with 1 mL "re-expression medium," seeded

on the coverslips which were coated with polyonithin (P2533, Sigma). After 2 hrs incubation at

37 °C with 5 % CO_2, cells were processed for immunocytochemistry against PDGFRα and DAPI.

For qRT-PCR or Western blot analysis, MACS sorted OPCs were lysed by RIPA buffer (89900,

Thermo Scientific).

Western blot analysis

Deeply anesthetized mice were perfused with cold PBS. Medial prefrontal cortices (mPFC) and corpus callosa (cc) were dissected in ice cold PBS. Tissues were homogenized with sucrose lysis buffer (320 mM sucrose, 10 mM Tris-HCl, 1 mM $NaHCO_3$ (pH=7.4); 1 mM $MgCl_2$), and MACS-sorted OPCs were lysed with RIPA buffer. Both buffers were supplemented with 1 X protease inhibitors (05892970001 Roche) and 1 X phosphatase inhibitor (04906837001, Roche). Protein (5 µg) was blotted onto nitrocellulose transfer membranes (QP0907015, qpore). Membrane was blocked with 5 % non-fat milk or 5 % BSA (A7906, Sigma) diluted in 0.1 % TBST. Primary antibodies were diluted with corresponding blocking buffer. Primary antibodies used for Western blot are listed in the **Supplementary table 3.**

Secondary antibodies were: HRP anti-mouse (1:2000,) and anti-rabbit (1:2000). Membranes were illuminated with WesternBright Chemilumineszenz Substrat Quantum kit (541015, Biozym) and documented with ChemiDoc-MP.

Quantitative real time PCR

Brain tissue or the MACS-sorted OPCs were homogenized as described above. NucleoSpin RNA Plus XS kit (740990.50, Macherey-Nagel) was used to extract mRNA and Omniscript kit (205113, QIAGEN) was used for reverse transcription. RT-PCR was performed using EvaGreen (27490, Axon) kit. Primer sequences for qRT-PCR are listed in **Supplementary table 4**.

Electrophysiology

Slice preparation: Mice were anesthetized by isoflurane before decapitation, and brain was quickly isolated and immersed in an ice-cold, oxygenated (5 % CO_2/95 % O_2, pH=7.4) solution containing (in mM) 87 NaCl, 3 KCl, 25 $NaHCO_3$, 1.25 NaH_2PO_4, 3 $MgCl_2$, 0.5 $CaCl_2$, 75 sucrose and 25 glucose. Coronal or semi-sagittal slices in 300 µm thickness were prepared with vibratome (Leica VT 1200S, Nussloch, Germany) and transferred to a nylon basket slice holder for incubation in artificial cerebral spinal fluid (ACSF) containing (in mM) 126 NaCl, 3 KCl,

25 $NaHCO_3$, 15 glucose, 1.2 NaH_2PO_4, 2 $CaCl_2$, and 2 $MgCl_2$ at 32 °C for 0.5 h. Subsequently,

slices were removed from the water bath and kept at RT with continuous oxygenation prior to use.

IPSCs and EPSCs recordings: Semi-sagittal slices were transferred to the recording chamber

that was continuously perfused with oxygenated ACSF containing 1 $MgCl_2$ and 2.5 $CaCl_2$ at a

flow rate of 2-5 mL/min. During recordings, 50 µM strychnine and 50 µM picrotoxin were added

to block inhibitory synaptic transmission. Pyramidal neurons were identified morphologically

(Axioskop 2 FS mot, Zeiss, Jena, Germany) with a 40x water immersion objective. Tissue was

detected with a QuantEM 512SC camera (Photometrics, Tucson, USA). Whole-cell membrane

currents were recorded by an EPC 10 USB amplifier (HEKA, Lambrecht, Germany), low pass

filtered at 3 kHz and data acquisition was controlled by Patchmaster software (HEKA). The

resistance of patch pipettes (7–9 ΩM) were fabricated from borosilicate capillaries (OD: 1.5 mm;

Sutter, USA) using a Micropipette Puller (Model P-97, Sutter Instrument Co., CA). Patch pipettes

were filled with the solution containing (in mM) 125 cesium gluconate, 20 tetraethylammonium

(TEA), 2 $MgCl_2$, 0.5 $CaCl_2$, 1 EGTA, 10 HEPES and 5 Na_2ATP (pH=7.2). Spontaneous excitatory

and inhibitory postsynaptic currents (sEPSCs, sIPSCs) of pyramidal neurons in mPFC were

recorded for 40 s in voltage-gated mode at a holding potential of -70 mV and +30 mV,

respectively. Currents above 10pA were analyzed with MATLAB. EPSCs of pyramidal neurons in

the layer V of mPFC were recorded by stimulating presynaptic axons in the layer II/III of mPFC

with a concentric bipolar microelectrode (MicroProbes, USA). Stimulus duration was 200 µs. To

estimate short-term plasticity, 40 trains of stimulation were applied to induce synaptic depression,

and the depressing trains were repeated with frequency of 10 Hz and an interval of 20 s to allow

for recovery from synaptic depression. The stimulation threshold was applied from 30 µA to

80 µA. IPSCs were conducted in the same region, however with a holding potential of 0 mV.

Compound action potential recordings: Compound action potentials were recorded in the

corpus callosum as previously described (Crawford et al., 2009). Micropipettes with 1-3 MΩ

resistance were filled with ACSF. Inward responses were evoked in current-clamp mode by

varying the intensity of stimulus pulses (0.2-4.0 mA) at 1 mm distance between recording and

stimulation electrodes, with a stimulus duration of 200 µs. The sample sweeps were acquired

every 5 s. Conduction velocity was estimated by changing the distance from 2.5 mm to 0.5 mm

between the stimulating and recording electrodes with a constant stimulus. To enhance the signal

to noise ratio, we averaged at least 15 successive sweeps. Data analysis was performed with

Igor pro 6.3.7.2 (WaveMetrics, Oregen, US).

All the experiments were conducted at RT (22 – 24 °C).

Data analysis: Data generated by Patch Master were loaded into MATLAB (Mathworks, MA,

USA) with a module adapted from sigTOOL (Lidierth, 2009). Evoked EPSC and IPSC traces from

the same cells were manually checked and pooled. The average of EPSC/IPSC traces from each

cell was used for analysis. Data analysis was performed using routines that were custom written

in Matlab.

EEG telemetry and analysis

Telemetric EEG transmitter implantation was adapted from Bedner and colleagues [7]. Mice were

implanted with telemetric EEG transmitters (DSI PhysioTel® ETA-F10, Harvard Biosciences, Inc.

Holliston, Massachusetts, USA) between 8 and 10 weeks of age. The animals were placed in a

stereotaxic frame (Robot stereotaxic, Neurostar, Tübingen, Germany) for implantation of depth

electrodes at 3.4 mm posterior to bregma and bilaterally 1.6 mm from the sagittal suture. After

post-surgical care and recovery, cages were placed on individual radio receiving plates (DSI

PhysioTel® RPC-1, Data Sciences International, St. Paul, USA) for synchronized EEG and video

recording (MediaRecorder Software, Noldus Information Technology, Wageningen, Netherlands).

EEG and video recording were performed for 24 hrs.

Data analysis: EEG traces were analysed with the Neuroscore software (Version 3.3.1., Data

Sciences International, St. Paul, USA). After applying a general Notch filter (50 Hz band stop),

relative power band values were determined with order 13 fast-Fourier transform in 10s epochs

and later averaged per hour of recording. Statistical analysis was performed with GraphPad

Prism.

Behavioral analysis

Several behavioral tests were performed with the same cohorts of mice (male, 9-week-old). The

order of the tests was designed from low to high invasiveness to reduce the interference from the

prior tests. The chamber or tested square was wiped with 75 % ethanol before tests to remove

odors.

Nest building test: The nest building test was carried out as previously described [8]. Mice were

moved into new single cages in the behavior testing room for habituation. After 24 h, a piece of

pressed cotton in size of 7x5 cm2 was placed in the cage. After 14 h, the shape and weight of the

cotton was recorded and unbiasedly scored according to the criteria [8].

Open field test: The mice were put in the open field maze, which measured 50 cm (length) x 50

cm (width) x 38 cm (height). Mice could move and explore freely for 10 min in the open field

square. In each test, the single mouse was put in the center of the square arena. The videos of

these 10 min were recorded (USB webcam) and analyzed (EthoVision XT 11.5, Noldus

Technology). Duration time in the center area (s), moved distance (cm) and speed (cm/s) were

determined [9]

New object recognition test: The mice could freely explore the two identical objects placed at a

distance of 8.5 cm from the side walls in two opposite corners of the apparatus for 10 min. After

40 min, one of the objects was replaced by a novel one, and the test mouse was allowed to

explore again. Preference index was defined as the percentage of the time exploring one identical

object within the total time exploring both objects. Recognition index was defined as the

percentage of the time exploring the novel object among the total time of exploring both objects.

Three chamber social behavior test: The three-chamber box was employed for social

behavioral studies [10]. Three chambers were equally sized and separated by two walls evenly

distributed in the box. A square door at the bottom center of each door allowed free running of

the mice within the three chambers. Two empty wire cages were placed at the side chambers,

leaving the center chamber empty. For habituation, mice were kept in the center chamber for

10 min, followed by a 10 min-habituation session with access to all three chambers. For the

sociability test, the test mouse was placed in the center chamber, while a mouse of similar age

was kept under the stainless wire cage in one of the side chambers. The other chamber contained

an empty wire cage. For 10 min, the test mouse could select freely all three chambers. For the

social novelty test, a novel mouse was placed under the other empty cage being an unfamiliar

mouse, and the prior mouse as familiar one. The test mouse was again allowed to freely explore

both animals for 10 min. The experiment was recorded with video camera and the time that the

test mouse spent in each chamber and the time of sniffing was analyzed.

Cell culture

Cell line Oli-neu: The murine oligodendroglial precursor cell line Oli-neu [11] was kindly provided

by Professor Jacqueline Trotter (University of Mainz). Undifferentiated Oli-neu cells were

incubated at 37 °C and 5 % CO_2 in poly-L-lysine (Merck) coated cell culture flasks (Greiner Bio

one) for expansion or cell culture dishes (Greiner Bio one) for experiments. Sato medium

consisting of DMEM high glucose medium (Fisher Scientific) with supplementation of 10 µg/ml

transferrin (Sigma), 10 µg/ml insulin (Santa Cruz), 100 µM putrescine (Sigma), 200 nM

progesterone (Sigma), 500 nM tri-iodo-thyrodine (Sigma), 220 nM sodium selenite (Sigma),

520 nM L-thyroxine (Sigma) and 1.5 % normal horse serum (Fisher Scientific) was used for

culturing and proliferation of cells. For experiments, $2x10^5$ cells were seeded in a 60 mm Petri

dish. 48 hours later, the medium was changed to fresh, with or without CGP 55845 (50 µM) for

24 hours. Cells were lysed with Qiazol (Qiagen) for qRT-PCR in a blind manner. Conditional

medium of control cells was collected for further neuronal treatment.

Primary culture of cortical neurons: Cortical neurons were isolated from p0 pups (C57BL/6

mouse strain). Briefly, the cortex was dissected from the whole brain in ice cold Earle's Balanced

Salt Solution (EBSS, Gibco). The cortex was then digested with 35 units papain (Worthington,

NJ) for 45 min at 37 °C, followed by gentle mechanical trituration. 1x105 cells were seeded on

25 mm glass coverslips in 24-well culture plates for further immunostaining. The glass coverslips

were pre-coated with a mixture of coating solution containing 17 mM acetic acid, poly-D-Lysine

(Sigma, P6407) and collagen I (Gibco, A1048301). Neurons were cultured in NBA culture medium

that contained 10 % FCS, 1 % penicillin-streptomycin, 1 % GlutaMAX and 2 % B-27 supplement

(Gibco) for 7 d at 37 °C with 5 % CO_2 incubator before the experiment. To examine the effect of

TWEAK on neurons, conditioned Oli-neu culture medium supernatant or Sato, was applied to

neuron NBA culture medium, with a ratio of 1:2 to NBA resulting in 1 ml medium per 24-well and

2 ml medium per 6-well. Cells were collected after 6 h with different culture condition with or

without 20 µM TWEAKR inhibitor L524-0366 (509374, Calbiochem) for further analysis.

Image acquisition and analysis

Brain slices were scanned with the fully automated slide scanner AxioScan.Z1 (Zeiss, Jena) and

LSM 710 confocal microscope (Zeiss, Jena). Cell counting was manually performed without bias

by using ZEN software (Zeiss, Jena). The lengths of nodes and paranodes were measured using

the 'straight' tool of Fiji software. The Imaris (version 9.6) with functions of volume analysis and

vesicle classification were used for MBP-PV (Fig. 1M, N) and vGAT study (Fig. F-I). Cell counting

and Imaris analysis were carried out in a blind manner.

Bromodeoxyuridine assay

Adult animals received 1 mg/ml bromodeoxyuridine (BrdU) (B5002, Sigma-Aldrich, St. Louis, MO)

dissolved in drinking water for seven consecutive days. For juvenile mice, single shot of 10 mg/ml

BrdU dissolved in 0.9 % NaCl were intraperitoneally injected to the p13 pups and analyzed at

p14.

Electron microscopy

Perfusion & dissection: Anesthetized (Ketamine/Xylazine in 0.9% NaCl solution (all: Bayer,

Leverkusen, Germany)) mice were intracardially perfused with buffered heparin (B. Braun,

Melsungen, Germany) solution (end concentration 50 IU/mL) followed by perfusion fixation with

4% (w/v) paraformaldehyde and 0.5% (w/v) glutaraldehyde in 0.1M cacodylate buffer. During

perfusion, the solutions were gradually cooled down from 37°C to ice cold temperatures. Whole

brains were dissected, immersed in fixative solution for 24h, and stored in 0.1M cacodylate buffer

at 4°C until further use.

Slicing: For vibratome slicing, whole brains were embedded in 10% gelatin and immersed in ice

cold 0.1M cacodylate buffer. 500 μm frontal sections were made using a vibratome (VT1000S,

Leica Microsystems, Wetzlar, Germany) at a frequency of 80Hz and 0.1mm/s speed. Slices

containing the region of interest were determined using a mouse brain atlas [12], and particular

brain areas (i.a. corpus callosum) were meticulously cut out using a fine scalpel.

Embedding & examination: The specimen of the different brain regions were embedded for

electron microscopy (1975) [13]. In short, samples were repeatedly rinsed in 0.1M cacodylate buffer,

osmicated in 2% osmium tetroxide in 0.1M cacodylate buffer for 1h, washed in distilled water, and

dehydrated in an ascending series of ethanol (70% to 100%) and acetone (100%, water-free). An

embedding in Epon resin (EMS, Hatfield, PA, USA) followed. After thorough polymerization, semi-

thin (300nm) and ultra-thin (65nm) sections were cut using an ultra-microtome (EM UC7, Leica

Microsystems, Wetzlar, Germany). Semi-thin sections were stained with Richardson blue staining

solution, and examined under a light microscope (DM2500, Leica Microsystems, Wetzlar,

Germany). Ultra-thin sections were contrasted with 5% lead citrate for 5min, examined with a

transmission electron microscope (Tecnai G2, FEI, Hillsboro, OR, USA), and randomly selected

areas were documented with a digital camera (MegaView III, Olympus, Shinjuku, Japan).

Statistics

Prior to statistical analysis, data were tested for outlier identification and Gaussian distribution

with GraphPad Prism 8. Statistical differences were indicated in the figures and legends. Each

data point in the graphs indicates biological individuals. Exceptions are indicated in the figure

legends.Data are shown as mean±SEM, except for violin plots (Fig. 2i and Fig. 4b). The median

and quartiles are indicated as thick and thin dashed lines, respectively.

References

1.   Huang, W*., et al.* Novel NG2-CreERT2 knock-in mice demonstrate heterogeneous differentiation potential of NG2 glia during development. *Glia* **62**, 896-913 (2014).

2.   Haller, C*., et al.* Floxed allele for conditional inactivation of the GABAB(1) gene. *Genesis* **40**, 125-130 (2004).

3.   Madisen, L*., et al.* A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. *Nat Neurosci* **13**, 133-140 (2010).

4.   Leone, D.P*., et al.* Tamoxifen-inducible glia-specific Cre mice for somatic mutagenesis in oligodendrocytes and Schwann cells. *Mol Cell Neurosci* **22**, 430-440 (2003).

5.   Huang, W*., et al.* During Development NG2 Glial Cells of the Spinal Cord are Restricted to the Oligodendrocyte Lineage, but Generate Astrocytes upon Acute Injury. *Neuroscience* **385**, 154-165 (2018).

6.   Jahn, H.M*., et al.* Refined protocols of tamoxifen injection for inducible DNA recombination in mouse astroglia. *Sci Rep* **8**, 5913 (2018).

7.   Bedner, P*., et al.* Astrocyte uncoupling as a cause of human temporal lobe epilepsy. *Brain* **138**, 1208-1222 (2015).

8.   Deacon, R.M. Assessing nest building in mice. *Nat Protoc* **1**, 1117-1119 (2006).

9.   Seibenhener, M.L. & Wooten, M.C. Use of the Open Field Maze to measure locomotor and anxiety-like behavior in mice. *J Vis Exp*, e52434 (2015).

10.   Kaidanovich-Beilin, O., Lipina, T., Vukobradovic, I., Roder, J. & Woodgett, J.R. Assessment of social interaction behaviors. *J Vis Exp* (2011).

11.   Jung, M*., et al.* Lines of murine oligodendroglial precursor cells immortalized by an activated neu tyrosine kinase show distinct degrees of interaction with axons in vitro and in vivo. *Eur J Neurosci* **7**, 1245-1265 (1995).

12.   Franklin, K. & Paxinos, G. *The mouse brain in stereotaxic coordinate* (Academic press, 2007).

13.   Glauert, A.M. *Fixation, Dehydration and Embedding of Biological Specimens. Practical methods in electron microscopy.* (North-Holland Publishing Company, 1975).

292 **Supplementary table 1: Primary antibodies used for immunostaining**

Antibodies	Host	Dilutions	Cat.No.	Company	PRID
PDGFRα	goat	1:500	AF1062	R&D Systems	AB_2236897
adenomatous polyposis coli clone 1(CC1)	mouse	1:200	OP80	Calbiochem	AB_2057371
Olig2	rabbit	1:500	AB9610	Millipore	AB_570666
MBP	mouse	1:500	SMI99	Biolegend	AB_10120130
Parvalbumin	rabbit	1:1000	PV 25	Swant	AB_10000344
Parvalbumin	mouse	1:500	P3088	Sigma	AB_477329
Caspr	rabbit	1:500	ab34151	abcam	AB_869934
DsRed	rabbit	1:1000	632496	Clontec	AB_10013483
BrdU	rat	1:1000	ab6326	abcam	AB_305426
Cleaved Caspase-3	rabbit	1:200	9661	Cell Signaling Technology	AB_2341188
Cleaved Caspase-3	mouse	1:100	STJ9744	St. John's Laboratory	
GAD67	mouse	1:500	MAB5406	Millipore	AB_2278725
NeuN	mouse	1:500	MAB377	Millipore	AB_2298772
CTIP2	rat	1:200	650601	Biolegend	AB_10896795
TBR1	rabbit	1:500,	49661	Cell Signaling Technology	AB_2799364
vGAT	mouse	1:500	13002	Synaptic Systems	AB_887871
FN14	mouse	1:50	314108	Biolegend	AB_2810477

294 **Supplementary table 2: Secondary antibodies used for immunostaining**

Antibodies	Fluorophore	Cat. No.	Company
Donkey anti-mouse	Alexa Fluor® 488		Thermofischer
	Alexa Fluor® 546	A10036	
	Alexa Fluor® 647	A31571	
	DyLight® 755	SA5-10171	Invitrogen
Donkey anti-rabbit	Alexa Fluor® 488	A21206	Thermofischer
	Alexa Fluor® 546	A10040	
	Alexa Fluor® 647	A31573	
	Alexa Fluor® 790	A11374	
Donkey anti-goat	Alexa Fluor® 488	A11055	Thermofischer
	Alexa Fluor® 546	A11056	
	Alexa Fluor® 647	A21447	
	Alexa Fluor® 750	ab175744	abcam
Donkey anti-rat	DyLight-755	SA5-10031	Thermofischer
DAPI (25ng/ml**)**		A10010010	biochmica

296 Secondary antibodies were diluted 1:1000 with the blocking buffer.

298   **Supplementary Table 3. Primary antibodies used for Western blot**

Antibodies	Host	Dilutions	Cat.No.	Company	PRID
GABA$_{B1}$R	mouse	1:500	ab55051	abcam	AB_941703
GAPDH	mouse	1:2000	G8795	Sigma	AB_1078991
MBP	mouse	1:1000	SMI99	Biolegend	AB_10120130
Tubulin-α	rabbit	1:2000	T6074	Sigma	AB_477582
TWEAK	rabbit	1:500	NBP1-76695	Novus Biologicals	AB_11036308

301　**Supplementary Table 4. Primers used for qRT-PCR**

Gene	Forward	Reverse
GABA_{B1}R	5'-CGAAGCATTTCCAACATGAC-3'	5'-CAAGGCCCAGATAGCATCATA-3'
β-actin	5'-CTTCCTCCCTGGAGAAGAGC-3'	5'-ATGCCACAGGATTCCATACC-3'
TNF-α	5'-GACGTGGAACTGGCAGAAGAG-3'	5'-TTGGTGGTTTGTGAGTGTGAG-3'
TWEAK	5'-CCGCCAGATTGGGGAATTTAC-3'	5'-AGTCCAAAGTAGGTTAGGAAGGG-3'
FasL	5'-TCCGTGAGTTCACCAACCAAA-3'	5'-GGGGGTTCCCTGTTAAATGGG-3'
TNFSF7	5'-TGTAGCGGACTACTCAGTAAGC-3'	5'-TGGGGTCCTTCCGAGGAAC-3'
Trp53	5'-CTCTCCCCCGCAAAAGAAAAA-3'	5'-CGGAACATCTCGAAGCGTTTA-3'
NGF	5'-TGATCGGCGTACAGGCAGA-3'	5'-GCTGAAGTTTAGTCCAGTGGG-3'