Impaired bidirectional communication between interneurons and

oligodendrocyte precursor cells affects cognitive behavior

Lipao Fang[1], Na Zhao[1], Laura C. Caudal[1], Renping Zhao[2], Ching-Hsin Lin[3], Hsin-

Fang Chang[3], Nadine Hainz[4], Carola Meier[4], Bernhard Bettler[5], Wenhui Huang[1], Anja

Scheller[1], Frank Kirchhoff[1,*], Xianshu Bai[1,*]

1. Molecular Physiology, CIPMM, University of Saarland, 66421 Homburg, Germany

2. Biophysics, CIPMM, University of Saarland, 66421 Homburg, Germany

3. Cellular Physiology, CIPMM, University of Saarland, 66421 Homburg, Germany

4. Department of Anatomy and Cell Biology, University of Saarland, 66421 Homburg,

Germany

5. Department of Biomedicine, University of Basel, 4056 Basel, Switzerland

Correspondence to:

Prof. Dr. Frank Kirchhoff,

Molecular Physiology,

CIPMM, University of Saarland,

66421 Homburg, Germany

frank.kirchhoff@uks.eu

Dr. Xianshu Bai

Molecular Physiology,

CIPMM, University of Saarland,

66421 Homburg, Germany

xianshu.bai@uks.eu

Abstract

Cortical neural circuits are complex but very precise networks of balanced excitation and inhibition (E/I). Yet, the molecular and cellular mechanisms that form the E/I balance are just beginning to emerge. Here, using conditional $GABA_B$ receptor-deficient mice we identified a GABA/TNF-related cytokine (TNFSF12)-mediated bidirectional communication pathway between Parvalbumin-positive (PV^+) fast spiking interneurons and oligodendrocyte precursor cells (OPCs) that determines the density and function of interneurons in the developing medial prefrontal cortex (mPFC). Interruption of the GABAergic signaling to OPCs resulted in reduced myelination and hypoactivity of interneurons, strong changes of cortical network activities and impaired cognitive behavior. In conclusion, glial transmitter receptors are pivotal elements in finetuning distinct brain functions.

Introduction

The inhibition of cortical network activity is performed by interneurons that release the inhibitory transmitter γ-aminobutyric acid (GABA) which acts on ionotropic $GABA_A$ and metabotropic $GABA_B$ receptors [1, 2]. Alterations of interneuron cell density as well as concomitant changes of firing activity are often observed in several neuropsychiatric conditions [3-6]. In the mouse cortex, higher frequencies of spontaneous inhibitory postsynaptic currents (sIPSCs) were observed after increasing the interneuron density by transplantation of precursor cells [7]. Similarly, blocking of interneuron apoptosis caused a decrease of the excitation/inhibition (E/I) ratio [6, 8]. Interneuron density is a pivotal determinant of correct inhibitory circuits in the central nervous system (CNS). During development, a surplus of interneurons is generated that populates the cortical plate [9]. Subsequently, about 40 % of cortical inhibitory neurons are eliminated by programmed cell death during the first two

postnatal weeks [7]. One hypothesis links the apoptosis of supernumerary neurons

with the onset of first neuron-neuron connections [10]. Only the neurons that receive

sufficient neurotrophic signals from their target cells will survive [10, 11].

Oligodendrocyte precursor cells (OPCs) form synapses with interneurons as early as

postnatal day (p) 4-5 [12]. Concomitantly, the incidence of interneuron apoptosis

increases drastically after p5 and reaches its peak at p7 [7]. Right after, the

interneuron-OPC connectivity accelerates till p10, which drives an immediate

oligodendrocyte (OL) boom and subsequent interneuron myelination via OPC-$GABA_A$

and $GABA_B$ receptors [13-15]. Despite relatively short axons, myelination of cortical

interneurons (mainly parvalbumin (PV)$^+$ fast-spiking interneurons, [16]) contributes

significantly to the fine-tuning of local activity in the medial prefrontal cortex (mPFC)

and cortex, ensuring proper performance of behavior [4, 17]. However, how the early

communication between interneuron and OPC affect inhibitory network activity in the

mPFC still remains unknown.

To assess the function of OPC $GABA_B$Rs for mPFC inhibition, we conditionally

deleted the $GABA_B$ receptor subunit *gabbr1* selectively in OPCs at the end of the first

postnatal week. By focusing on the medial prefrontal cortex (mPFC), we found that

OPCs shape the inhibitory network by $GABA_B$ receptor and the cytokine tumor

necrosis factor-like weak inducer of apoptosis (TWEAK) signaling pathways, thereby

finetuning interneuron cell death and survival as well as its myelination onset. The

functional and morphological changes of PV$^+$ interneurons observed in the adult

mPFC of mutant mice generate a reduction of the inhibitory tone, which finally leads

to impaired cognition and social behavior.

Results

$GABA_B$Rs of OPCs are required for oligodendrogenesis in the mPFC

To assess the function of $GABA_B$Rs for OPC-interneuron communication, we

generated Tg*(NG2-CreER^{T2}):GABA$_{B1}$R$^{fl/fl}$* mice (Fig. 1a) to conditionally knockout

(cKO) functional $GABA_B$R from OPCs and their progeny. We induced the cKO at

postnatal day 7 and 8 (p7/8) before the onset oligodendrocyte formation at p10-11 [13].

When analyzed at the age of 9 weeks (w), about 80 % of OPCs (platelet-derived

growth factor receptor α$^+$ (PDGFRα$^+$, Pα+))were found recombined (Pα$^+$tdT$^+$/Pα$^+$) in

the medial prefrontal cortex (mPFC), based on tdTomato (tdT) gene expression

(Fig. 1b, c).

To confirm and quantify the deletion of $GABA_B$R in OPCs, we performed Western

blot analysis with OPCs purified by magnetic-activated cell sorting (MACs-OPCs). In

cKO OPCs, $GABA_B$R expression was reduced by about 64 % compared with controls

(ctl) (Fig. 1d). Considering the substantial purity of MACS-sorted OPCs (85 %, data

not shown) and recombination efficiency (80 %), we concluded that a vast majority of

$GABA_B$Rs had been ablated from OPCs.

Next, we focused on the oligodendrocyte lineage and evaluated the contribution of

$GABA_B$Rs for OPC differentiation in the mPFC by immunostaining of PDGFRα and

adenomatous polyposis coli clone 1 (CC1) (Fig. 1e), established markers for OPCs

and oligodendrocytes, respectively. In the cKO mice, the density of OPCs

(PDGFRα$^+$Olig2$^+$) did not change (Fig. 1f), while that of mature oligodendrocytes

(CC1$^+$Olig2$^+$) was strongly reduced by 26 %(Fig. 1g), indicative of decreased OPC

differentiation to oligodendrocytes (Fig. 1h). Notably, in the corpus callosum the

densities of oligodendrocytes and OPCs, as well as their relative proportions did not

differ between ctl and cKO mice (Fig.1 f-h, Extended Data Fig. 1). These data

suggest that OPC-GABA$_B$Rs are required for OPC differentiation in the grey matter

mPFC, but not in the white matter corpus callosum.

We further analyzed structural aspects of myelination which is important for axonal

conductivity, i.e. the lengths of nodes (the gaps between paired contactin-associated

protein-positive (Caspr$^+$) segments) and paranodes (single Caspr$^+$ segments) by

immunostaining (Fig. 1i). In cKO mPFC, the paranodal length was increased (Fig. 1j)

while node length and density remained stable (Extended Data Fig. 2). In parallel, we

observed a reduction of myelin basic protein (MBP) expression at the mRNA

(Extended Data Fig. 2) and protein levels (Fig. 1k) in cKO mPFC, while the other

myelin protein, proteolipid protein (PLP) mRNA levels were not changed (Extended

Data Fig. 2). In the corpus callosum, however, myelination was unperturbed as

indicated by unchanged paranode length (Extended Data Fig. 2), node density

(Extended Data Fig. 2) and MBP expression (Fig. 1k). Therefore, our data show that

GABA$_B$Rs of OPCs promote differentiation and myelination in the mPFC, but do not

affect the corpus callosum.

Since OPCs continuously differentiate into oligodendrocytes [18-20], the cKO mice will

also generate oligodendrocytes lacking GABA$_B$Rs. To selectively target only

oligodendrocytes, we used Tg*(PLP-Cre^{ERT2}):GABA$_{B1}$R$^{fl/fl}$* mice (Extended Data

Fig. 3) . The gene deletion was induced by tamoxifen injections at p7/8 or at 4w. At

the age of 9w, we did not observe differences in the OPC and oligodendrocyte

population between ctl and cKO mice (Extended Data Fig. 3), neither in the PFC nor

in the corpus callosum (Extended Data Fig. 3).

In summary, our data strongly suggest that GABA$_B$R is specifically required for early

OPC differentiation and subsequent myelination in the mPFC, but not in the corpus

callosum.

GABA_BRs of OPCs promote the myelination of interneurons in the mPFC

Since callosal axons are primarily excitatory [21] and axons of inhibitory interneurons

are largely restricted to cortical areas of the same hemisphere (including mPFC,

Fig.1l) (Extended Data Fig. 4), we hypothesized that the observed myelin deficits

were restricted to interneurons. Required by their fast-spiking activity, parvalbumin

(PV)[+] interneurons are the most abundantly myelinated interneurons [16, 22-24] of the

cortex. These neurons are most prominent at the layer 4 and 5 with majority of their

axons projecting to the layer 2/3 and 5 [25] (Fig. 1l, Extended Data Fig. 4). To evaluate

their myelination, we performed PV and MBP double immunostaining (Fig. 1l-n) and

analyzed the volume of myelin sheaths, by volume rendering of MBP immunolabel

covering PV[+] axons at the cortical layers 2/3 of mPFC. Indeed, the volume of myelin

sheaths wrapping PV neurons was reduced by 60 % in the cKO mPFC (Fig. 1n).

However, the myelination of excitatory neurons (in the corpus callosum) was not

affected as the g-ratio of myelinated axons and conduction velocity of compound

action potential did not differ between ctl and cKO mice (Extended Data Fig. 2). Thus,

we concluded that GABA_BR of OPCs are preferentially involved in the myelination of

inhibitory rather than excitatory neurons.

Postnatal deletion of GABA_BRs in OPCs causes enhanced interneuron density in the

adult, but with reduced GABA release

While investigating the interneuronal myelination, we recognized that early ablation

of GABA_BR in OPCs (at p7/8, Fig. 2a) was associated with a 20 % increase of

interneuron density in the cortex as revealed by immunostaining for PV (Fig. 2b).

However, in the mPFC this increase did not result in a concomitant increase of

inhibitory input (Fig. 2c, d). In contrast, the frequency of spontaneous inhibitory

postsynaptic currents (sIPSCs) of pyramidal neurons in the layer V of mPFC, where

interneurons are abundantly located, was reduced by 50 %, while the amplitude of

sIPSCs remained unaffected in the mutant mPFC as well as the firing rate and

amplitude of excitatory postsynaptic currents (EPSCs).

Since neuronal activity can regulate myelination [22, 26], we asked whether the reduced

inhibitory firing rate contributed to the observed molecular and structural changes of

the myelin at interneuronal axons in the mutant mPFC. For that purpose, we further

analyzed the cKO mPFC of p10 mice (Fig. 2e) at the onset of myelination and when

interneuron-OPC communication peaks [13]. After induction of the cKO at p7/8, we

assessed the density of the vesicular GABA transporter (vGAT) close to OPC

surfaces as a quantitative readout of inhibitory presynaptic terminals by combined

vGAT and PDGFRα immunostaining [27] (Fig. 2f-i, Extended Data Fig. 5). We

classified the vGAT puncta into two subgroups based on the distance between vGAT

staining and OPC surface: '<20 nm' as vesicles targeting on postsynaptic OPCs

(magenta), and '>20 nm' as vesicles targeting on other cells (grey), assuming a width

of about 20 nm for the synaptic cleft [28]. And indeed, already at p10, i.e. only 3 days

after the first tamoxifen injection, the density and the volume of vGAT immunopuncta

were reduced were reduced to 75 % and 80 % in the cKO mPFC (Fig. 2f-i, Extended

Data Fig. 5), including a smaller volume of vGAT (Fig. 2h) and less puncta (Fig. 2i) at

the OPC surface in cKO mPFC, which can explain the attenuated firing rate of the

interneurons in the adult. These results indicate that interneurons exhibit lower

activity and transmit less GABAergic signals to OPCs in the cKO mPFC.

To test whether the reduced inhibitory tone affecting OPC differentiation specifically

correlates with the onset of myelination, we compared the OPC and oligodendrocyte

populations at p10 and p14 (Fig. 2j-m, Extended Data Fig. 6). At p10, the densities of

OPC and oligodendrocyte populations were still comparable in the cKO mPFC to

control mPFC, although the interneurons already expressed less vGAT (Fig. 2f-h).

However, at p14, the differentiation and cell numbers of oligodendrocytes dropped

strongly in the mutant mPFC (Fig. 2l, Extended Data Fig. 6), as did the MBP

expression (Fig. 2m). Again, in the corpus callosum, OPC and oligodendrocyte

population were not affected by the deletion of $GABA_BRs$ (Extended Data Fig. 1).

When we induced $GABA_BR$ deletion in the young adult mice at the age of 4 weeks,

both in the mPFC and corpus callosum, the density of oligodendrocyte as well as the

MBP expression were not affected in the cKO mice (Extended Data Fig. 7).

In summary, our data strongly suggest that during early development of the mPFC

deletion of $GABA_BR$ in OPCs affects the inhibitory tone of interneurons, feeding back

onto OPC differentiation and interneuron myelination.

OPCs induce interneuron apoptosis via $GABA_BR$-TWEAK signaling

As we had seen a surplus of interneurons in the mutant mPFC which could explain

the abnormal neuronal activity [8, 29], we tested whether the interneuron numbers

would be accurately controlled by programmed cell death early during development.

After induction of $GABA_BR$ in OPCs at p7/8, the extent of interneuron apoptosis was

determined at p10 and p14 by immunostaining for PV and cleaved caspase-3 (CC-3,

Extended Data Fig. 8), a well-established marker for apoptosis. As expected by the

earlier experiments, we found 50 % reduction of $PV^+/CC-3^+$ interneurons at p10

suggestive of strongly reduced apoptosis of PV^+ interneurons in the mutant mPFC

(Extended Data Fig. 8). Obviously, OPCs receive GABAergic input through $GABA_BR$

and send back pro-apoptotic signals to interneurons.

To confirm the selective cell death of interneurons, we induced the cKO as early as

p1 and p2 and compared the apoptosis of inhibitory neurons with that of excitatory

ones at p5 and p7 (Fig. 3a). Interneuron apoptosis persists from p1 till p15 with a

peak at p7 [7], while the time window for excitatory neuron apoptosis is rather narrow,

between p2-5 [30]. Apoptosis of total interneurons ($GAD67^+CC-3^+$) as well as PV

interneurons (PV$^+$CC-3$^+$ left panel in Fig. 3b) was significantly reduced in cKO mPFC

at p5 and p7 (Fig. 3c), while that of the excitatory neurons (CTIP$^+$CC-3$^+$, right panel

in Fig. 3b; TBR1$^+$CC-3$^+$) did not change at both time points (Fig. 3c, Extended Data

Fig. 9). Therefore, we concluded that OPCs send pro-apoptotic signals back to

interneurons at the early postnatal weeks.

To molecularly identify the critical apoptotic factor released by OPCs, we first tested

the expression levels of several peptides which had been associated with cell

survival or cell death in ctl and cKO mPFC (Extended Data Fig. 10). Gene deletion

was induced at p1/2, and mPFC was collected at p7, the peak of interneuron

apoptosis (Extended Data Fig. 10). We found only the expression of the cytokine

tumor necrosis factor-like weak inducer of apoptosis (TWEAK, also known as Apo3l

or TNF superfamily member 12, TNFSF12) to be diminished about 30 % in cKO

mPFC (Fig. 3d, Extended Data Fig. 10). TWEAK induces an apoptotic cell death

after binding to its cognate TWEAK receptor (TWEAKR; TNF Receptor Superfamily

Member 12A, TNFRSF12A; also known as FN14) [31]. Subsequently, we tested

MACS-sorted OPCs (collected from cKO and ctl cortices at p7) (Fig. 3d) as well as

Oli-neu cells (an OPC cell line) treated with GABA$_B$R antagonist CGP 55845 for

TWEAK. Also here, we observed TWEAK downregulation (Extended Data Fig. 10).

Three independent experiments, including genetic deletion or pharmacological block

of OPC GABA$_B$R decreased TWEAK expression (Fig. 3d, Extended Data Fig. 10).

To confirm that OPC-derived TWEAK can indeed induce neuronal apoptosis, we

treated primary cortical neurons with conditioned medium obtained from OPC Oli-neu

cultures (Fig. 3e). A selective TWEAKR antagonist, L524-0366 was added to the

conditioned medium at 20 µM to block the binding of TWEAK on interneurons

(Fig. 3e, f). While the conditioned medium from the OPC cell line increased

interneuron apoptosis (Fig. 3g, h), co-treatment with the TWEAKR antagonist

prevented the cell death. These results further substantiate that OPCs induces

interneuron apoptosis via TWEAK secretion.

In summary, our results demonstrate that OPCs induce interneuron apoptosis by

releasing TWEAK upon $GABA_BR$ activation. Blocking this pathway prevents the

apoptosis of interneurons. The concomitant dysregulation of interneuron density

results in a reduced inhibitory GABAergic tone and altered myelination in the mPFC.

Interrupted OPC-interneuron communication after $GABA_BR$ loss in OPCs during

development generates cognitive impairment in adulthood

The mPFC is responsible for cognitive processes and their regulatory finetuning is

guaranteed by the E/I balance [32]. To test whether the selectively reduced interneuron

apoptosis and the associated structural myelin alterations can generate defects in

neural circuits and cognition, we further performed *in vivo* investigations using

electrophysiology and behavioral analysis in 9w old animals after gene ablation was

induced at P7/8 (Fig. 4a).

The patch-clamp recordings described above had revealed a reduced firing rate of

interneurons suggesting an imbalance of E/I in the mutant mPFC (Fig. 2c, d).

Impaired cortical network activity of the mutant mice was also detected in

electroencephalographical (EEG) recordings. The power of the gamma wave,

especially the lower gamma wave band (30-50 Hz) was slightly, but significantly

decreased by about 10% in cKO mice (Fig. 4b).

To evaluate the impact of circuit dysregulation in the mutant mPFC for the living

animal, we challenged mice for their cognitive performance employing tasks of social

cognitive behavior which had been associated with proper mPFC function, i.e. social

novelty and new object recognition [33]. In addition, nest building behavior is often

introduced as a gauge of general well-being of the mice and also being an early sign

of cognition decline [34]. We observed that the mutant mice built significantly poorer

nests with less than half score of the control animal nests, indicating putative

cognition deficits of the mutant mice (Fig. 4c). The three-chamber test was used to

assess social cognition by analyzing mice for a form of general sociability and their

interest in social novelty (Fig. 4d). First, the three-chamber test was used to monitor

general social behavior, i.e. preference of other mice over objects. Both, ctl and cKO

mice exhibited more interest in the mouse rather than the object during the first test

phase, indicating no disorder in sociability of the mutant mice (Fig. 4e, Extended

Data Fig. 11). At the second phase, a social novelty preference was assessed.

Control animals investigated the unfamiliar mouse more than the familiar one, while

the cKO mice exhibited no novel social preference as they spent similar time in both

chambers (Fig. 4f, Extended Data Fig. 11). Employing a new object recognition test

(Fig. 4g), we again observed the cognition defect of cKO mice. Control animals were

able to distinguish the new object replaced after a 40 min break from the habituation,

while the cKO mice showed similar interest to both of the old and new objects (right

panel in Fig. 4g-i). Open field test showed that the motor activity of the cKO mice (in

terms of the speed and total distance) was not affected (Fig. 4j-l).

Hence, we conclude that $GABA_B Rs$ of OPCs are essential for cognition by modifying

inhibitory circuits in the mPFC.

Discussion

A balance of excitation and inhibition (E/I) in the medial prefrontal cortex (mPFC) is

of key importance for mammalian cognition. Establishing correct cell densities and

subsequent myelination have been identified as pivotal elements during CNS

development [6, 17]. This has been well documented for fast-spiking parvalbumin-

positive (PV[+]) interneurons. Here, we highlight OPCs as indispensable regulators of

interneuron population and myelination in the mPFC by providing a molecular

explanation and demonstrating how impaired bidirectional OPC-interneuron signaling

affects social cognitive behavior. OPCs at the first two postnatal weeks release

TWEAK to regulate interneuron apoptosis. Genetic ablation of $GABA_BR$ in OPCs at

the first postnatal week promoted interneuron survival. However, these neurons

displayed hypoactivity, less contacts with OPCs during development and deteriorated

myelin structures in the adulthood. Notably, these mutant mice exhibited severe

cognitive defects in their social behavior.

During development, large populations of immature interneurons invade the cortex

where their cell density is adjusted by programmed cell death [9, 10]. For assuring their

survival, these neurons form connections with adjacent cells and receive retrograde

signals from their targets [10, 11]. Interneuron apoptosis drastically increases at p7 [7],

shortly after formation of interneuron-OPC synapses at p4-5 [12]. Based on this

established knowledge, we hypothesized that OPCs could exert an apoptotic impact

to their presynaptic partners. Indeed, our data provide strong evidence that, starting

from postnatal day 5 on, OPCs elicit an apoptotic cascade of adjacent interneurons

to adjust their density. In OPCs, the regulatory process employs a $GABA_BR$-TWEAK

signaling pathway. TWEAK, TNF super family member 12 (TNFSF12), is TNF-like

weak apoptotic factor. Membrane-bound and cleaved isoforms of TWEAK are both

able to induce cell apoptosis by binding to the TWEAK receptor (TWEAKR, also

known as FN14) on the neuronal side [31, 35, 36]. According to our data $GABA_BR$-

TWEAK signaling preferentially affects interneurons, suggesting this 'kill me' signal

acting highly localized. Indeed, TWEAK is specifically recruited to synapses where

TWEAKRs are expressed [37]. OPCs directly sense GABAergic signals at their

interneuron-OPC synapse. Subsequent signaling triggers TWEAK expression and

translocation to the OPC surface, where it could directly bind interneuronal

TWEAKRs. Alternatively, the TWEAK ectodomain could be proteolytically cleaved

and act as a soluble factor. However, this scenario could also happen at soma-

somatic contact sites between OPCs and interneurons [38, 39]. The regulatory impact

appears to be highly specific since the OPC-induced interneuron apoptosis was

restricted to the first two postnatal weeks, a well-established time window of

interneuron apoptosis [7, 30]. This also explains why the cKO induction at the age of 4w

did not impact OPC differentiation and myelination, although $GABA_BR$ expression

peaks in OPCs at 4w (data not shown). Please note, instead of a gradual decline, we

found a slight increase of PV^+CC3^+ cells at p14 compared to p10, probably due to the

age-dependent increase of PV expression described for fast spiking interneurons [40].

During the same time period of interneuronal apoptosis a substantial portion of OPCs

arriving in the cortex are eliminated as well [41]. This first wave of OPCs share the

same cellular origin with a 60 % majority of interneurons, i.e. $Nkx2.1^+$ precursors from

the medial ganglionic eminence (MGE) and the embryonic preoptic area (ePOA) [9, 41].

However, whether a causal link exists between common origin and

$GABA_BR$/TWEAK-mediated interneuron-OPC signaling requires further analysis.

Differentiation of oligodendrocytes and subsequent myelination are strongly affected

by neuronal activity [22, 26, 42]. Indeed, when we induced $GABA_BR$ loss at p7, we

observed a suppression of the inhibitory tone in the mPFC early in development (p10;

attenuated expression of vGAT in interneurons and less vGAT puncta on the surface

of OPCs), before a change in oligodendrocyte density could be detected at p14.

Concomitantly, the oligodendrocytes formed less myelin around the interneuronal

axons and altered paranodal structures in the adult mPFC which are essential for

precise action potential propagation. The morphological deterioration was only

observed in the mPFC, but not in the corpus callosum composed axons of excitatory

neurons only [21]. In line with these neuron-type specific observations, the qRT-PCR

results indicated reduced levels of MBP, but not PLP mRNA expression in the mutant

mPFC, both at p14 and 9w of age. Axons of interneurons and excitatory neurons are

different in respect to the protein content of their myelin sheaths. MBP is

preferentially expressed in the myelin of inhibitory axons, while PLP is more

prominent in excitatory axons PLP [16].

An early loss of the $GABA_AR$ γ2 subunit in OPCs reduced the firing rate of

presynaptic PV[+] interneurons and caused a similar myelin defect of longer nodal

structures in PV[+] neurons [4], the initially very different signal pathways of both,

$GABA_AR$ and $GABA_BR$, have to merge to affect myelin gene expression and OPC-

axon contacts similarly. The $GABA_AR$ signaling between interneurons and OPCs is

relatively well established [4, 15, 43, 44]. E.g., the γ2 subunit of $GABA_AR$ contributes to the

maintenance of the OPC, but has no impact on oligodendrocyte formation [15].

However, the role of $GABA_BRs$ for OPC differentiation is complex. We observed that

$GABA_BR$ was essential for oligodendrogenesis at the early development but not at

later stages. Similary, an *in vitro* study showed that activation of $GABA_BR$ by the

selective agonist baclofen could stimulate the differentiation of cortical OPCs

prepared from p0-2 pups [14]. Since OPCs are heterogeneous with age [45], $GABA_BR$ of

OPCs could contribute to the developmental diversity.

Obviously, the GABAergic signaling of OPCs plays pivotal roles for the myelination of

interneurons, which ensures precise finetuning of the local neural circuitry [46]. As seen

in chandelier cells [6], excessive number of interneurons in the OPC-$GABA_BR$ cKO

mice exhibited suppressed activity, and reduced myelination in the mPFC. All these

physiological and morphological abnormalities certainly contributed to cognition

impairment. A proper E/I ratio in the prefrontal cortex, especially early E/I balance in

mPFC, is extremely important for social cognition [32, 47]. When the pyramidal neuron

activity was enhanced in the mPFC at early p7-11 caused a severe cognitive

dysfunction in the adulthood [47]. In addition, despite relatively short axons, myelination

of cortical interneurons (mainly PV^+ neurons [16]) contributes significantly to the

finetuning of local activity in the mPFC and cortex, insuring proper behavior

performance [4, 6, 17].

In conclusion, our study demonstrates that OPCs regulate inhibition in the mPFC via

$GABA_BR$/TWEAK signaling. During development, OPCs determine the density of

interneurons by adjusting their apoptosis. Subsequently, correct myelination of

interneurons, as an essential component of the network activity in the mPFC,

determines the cognition and social behavior.

References

1.   Douglas, R.J. & Martin, K.A. Inhibition in cortical circuits. *Curr Biol* **19**, R398-402

(2009).

2.   Bai, X., Kirchhoff, F. & Scheller, A. Oligodendroglial GABAergic Signaling: More Than

Inhibition! *Neurosci Bull* (2021).

3.   Marín, O. Interneuron dysfunction in psychiatric disorders. *Nat Rev Neurosci* **13**,

107-120 (2012).

4.   Benamer, N., Vidal, M., Balia, M. & Angulo, M.C. Myelination of parvalbumin

interneurons shapes the function of cortical sensory inhibitory circuits. *Nat Commun* **11**,

5151 (2020).

5.   Ferguson, B.R. & Gao, W.J. PV Interneurons: Critical Regulators of E/I Balance for

Prefrontal Cortex-Dependent Behavior and Psychiatric Disorders. *Front Neural Circuits* **12**, 37

(2018).

6.   Wang, B.S., *et al.* Retinal and Callosal Activity-Dependent Chandelier Cell Elimination

Shapes Binocularity in Primary Visual Cortex. *Neuron* **109**, 502-515.e507 (2021).

7.   Southwell, D.G., *et al.* Intrinsically determined cell death of developing cortical

interneurons. *Nature* **491**, 109-113 (2012).

8.   Orduz, D., *et al.* Developmental cell death regulates lineage-related interneuron-

oligodendroglia functional clusters and oligodendrocyte homeostasis. *Nat Commun* **10**, 4249

(2019).

9.   Lim, L., Mi, D., Llorca, A. & Marín, O. Development and Functional Diversification of

Cortical Interneurons. *Neuron* **100**, 294-313 (2018).

10.   Davies, A.M. Regulation of neuronal survival and death by extracellular signals

during development. *EMBO J* **22**, 2537-2545 (2003).

11.   Oppenheim, R.W. Cell death during development of the nervous system. *Annu Rev*

Neurosci **14**, 453-501 (1991).

12.   Vélez-Fort, M., Maldonado, P.P., Butt, A.M., Audinat, E. & Angulo, M.C. Postnatal
switch from synaptic to extrasynaptic transmission between interneurons and NG2 cells. *J*
Neurosci **30**, 6921-6929 (2010).

13.   Orduz, D.*, et al.* Interneurons and oligodendrocyte progenitors form a structured
synaptic network in the developing neocortex. *Elife* **4** (2015).

14.   Serrano-Regal, M.P.*, et al.* Oligodendrocyte Differentiation and Myelination Is
Potentiated via GABA. *Neuroscience* **439**, 163-180 (2020).

15.   Balia, M., Benamer, N. & Angulo, M.C. A specific GABAergic synapse onto
oligodendrocyte precursors does not regulate cortical oligodendrogenesis. *Glia* **65**, 1821-
1832 (2017).

16.   Micheva, K.D.*, et al.* A large fraction of neocortical myelin ensheathes axons of local
inhibitory neurons. *Elife* **5** (2016).

17.   Maas, D.A.*, et al.* Interneuron hypomyelination is associated with cognitive
inflexibility in a rat model of schizophrenia. *Nat Commun* **11**, 2329 (2020).

18.   Huang, W.*, et al.* Novel NG2-CreERT2 knock-in mice demonstrate heterogeneous
differentiation potential of NG2 glia during development. *Glia* **62**, 896-913 (2014).

19.   Kang, S.H., Fukaya, M., Yang, J.K., Rothstein, J.D. & Bergles, D.E. NG2+ CNS glial
progenitors remain committed to the oligodendrocyte lineage in postnatal life and following
neurodegeneration. *Neuron* **68**, 668-681 (2010).

20.   Zhu, X.*, et al.* Age-dependent fate and lineage restriction of single NG2 cells.
Development **138**, 745-753 (2011).

21.   Fame, R.M., MacDonald, J.L. & Macklis, J.D. Development, specification, and
diversity of callosal projection neurons. *Trends Neurosci* **34**, 41-50 (2011).

22.   Stedehouder, J.*, et al.* Fast-spiking Parvalbumin Interneurons are Frequently
Myelinated in the Cerebral Cortex of Mice and Humans. *Cereb Cortex* **27**, 5001-5013 (2017).

23.   Stedehouder, J., *et al.* Local axonal morphology guides the topography of

interneuron myelination in mouse and human neocortex. *Elife* **8** (2019).

24.   Zonouzi, M., *et al.* Individual Oligodendrocytes Show Bias for Inhibitory Axons in the

Neocortex. *Cell Rep* **27**, 2799-2808.e2793 (2019).

25.   Scala, F., *et al.* Layer 4 of mouse neocortex differs in cell types and circuit

organization between sensory areas. *Nat Commun* **10**, 4174 (2019).

26.   Mitew, S., *et al.* Pharmacogenetic stimulation of neuronal activity increases

myelination in an axon-specific manner. *Nat Commun* **9**, 306 (2018).

27.   Chaudhry, F.A., *et al.* The vesicular GABA transporter, VGAT, localizes to synaptic

vesicles in sets of glycinergic as well as GABAergic neurons. *J Neurosci* **18**, 9733-9750 (1998).

28.   Dani, A., Huang, B., Bergan, J., Dulac, C. & Zhuang, X. Superresolution imaging of

chemical synapses in the brain. *Neuron* **68**, 843-856 (2010).

29.   Fang, W.Q., *et al.* Overproduction of upper-layer neurons in the neocortex leads to

autism-like features in mice. *Cell Rep* **9**, 1635-1643 (2014).

30.   Wong, F.K., *et al.* Pyramidal cell regulation of interneuron survival sculpts cortical

networks. *Nature* **557**, 668-673 (2018).

31.   Chicheportiche, Y., *et al.* TWEAK, a new secreted ligand in the tumor necrosis factor

family that weakly induces apoptosis. *J Biol Chem* **272**, 32401-32410 (1997).

32.   Grossmann, T. The role of medial prefrontal cortex in early social cognition. *Front

Hum Neurosci* **7**, 340 (2013).

33.   Bicks, L.K., Koike, H., Akbarian, S. & Morishita, H. Prefrontal Cortex and Social

Cognition in Mouse and Man. *Front Psychol* **6**, 1805 (2015).

34.   Neely, C.L.C., Pedemonte, K.A., Boggs, K.N. & Flinn, J.M. Nest Building Behavior as an

Early Indicator of Behavioral Deficits in Mice. *J Vis Exp* (2019).

35.   Winkles, J.A. The TWEAK-Fn14 cytokine-receptor axis: discovery, biology and

therapeutic targeting. *Nat Rev Drug Discov* **7**, 411-425 (2008).

36.   Brown, S.A., Ghosh, A. & Winkles, J.A. Full-length, membrane-anchored TWEAK can

function as a juxtacrine signaling molecule and activate the NF-kappaB pathway. *J Biol Chem*

285, 17432-17441 (2010).

37.   Cheadle, L.*, et al.* Sensory Experience Engages Microglia to Shape Neural

Connectivity through a Non-Phagocytic Mechanism. *Neuron* **108**, 451-468.e459 (2020).

38.   Mangin, J.M., Kunze, A., Chittajallu, R. & Gallo, V. Satellite NG2 progenitor cells share

common glutamatergic inputs with associated interneurons in the mouse dentate gyrus. *J*

Neurosci **28**, 7610-7623 (2008).

39.   Boulanger, J.J. & Messier, C. Oligodendrocyte progenitor cells are paired with GABA

neurons in the mouse dorsal cortex: Unbiased stereological analysis. *Neuroscience* **362**, 127-

140 (2017).

40.   Larsen, R., Proue, A., Scott, E.P., Christiansen, M. & Nakagawa, Y. The Thalamus

Regulates Retinoic Acid Signaling and Development of Parvalbumin Interneurons in

Postnatal Mouse Prefrontal Cortex. *eNeuro* **6** (2019).

41.   Kessaris, N.*, et al.* Competing waves of oligodendrocytes in the forebrain and

postnatal elimination of an embryonic lineage. *Nat Neurosci* **9**, 173-179 (2006).

42.   Nagy, B., Hovhannisyan, A., Barzan, R., Chen, T.J. & Kukley, M. Different patterns of

neuronal activity trigger distinct responses of oligodendrocyte precursor cells in the corpus

callosum. *PLoS Biol* **15**, e2001993 (2017).

43.   Balia, M.*, et al.* Postnatal down-regulation of the GABAA receptor γ2 subunit in

neocortical NG2 cells accompanies synaptic-to-extrasynaptic switch in the GABAergic

transmission mode. *Cereb Cortex* **25**, 1114-1123 (2015).

44.   Passlick, S.*, et al.* Expression of the γ2-subunit distinguishes synaptic and

extrasynaptic GABA(A) receptors in NG2 cells of the hippocampus. *J Neurosci* **33**, 12030-

12040 (2013).

45.   Spitzer, S.O.*, et al.* Oligodendrocyte Progenitor Cells Become Regionally Diverse and

Heterogeneous with Age. *Neuron* **101**, 459-471.e455 (2019).

46.   Arancibia-Cárcamo, I.L.*, et al.* Node of Ranvier length as a potential regulator of

myelinated axon conduction speed. *Elife* **6** (2017).

47.   Bitzenhofer, S.H., Pöpplau, J.A., Chini, M., Marquardt, A. & Hanganu-Opatz, I.L. A

transient developmental increase in prefrontal activity alters network maturation and

causes cognitive dysfunction in adult mice. *Neuron* (2021).

Acknowledgement

We thank Daniel Schauenburg for excellent animal husbandry and Frank Rhode,

Alexander Grissmer and Samantha Bechet for experimental assistance. We are also

grateful to Prof. Jaqueline Trotter (University of Mainz, Mainz, Germany) for providing

the Oli-neu cell line and Prof. Leda Dimou (University of Ulm, Ulm, Germany) for

sharing MACs sorting protocols of OPCs. This work was supported by grants from

the Deutsche Forschungsgemeinschaft (SPP 1757, SFB 894 to FK; FOR2289 to AS,

FK, SPP1757 Young Investigator grant to XB), EraNet-Neuron BrlE to FK and

HOMFORexzellent2017 program to XB.

Author contributions

XB initiated the project. LF, XB, NZ, LC, NH, CL and AS performed experiments. LF, XB, NZ, LC and RZ analyzed data. BB, WH. HC and CM provided materials. XB and FK designed and supervised the study and wrote the manuscript with comments of the other authors.

Competing interests

The authors declare no competing interests.

 Figure legend

Figure 1. Ablation of GABA$_B$Rs in OPCs attenuates oligodendrocyte

differentiation and myelination of interneurons in the medial prefrontal cortex.

a, Genetic strategy and experimental schedule. *b,* Exemplary image of prefrontal

cortex in coronal section stained with PDGFRα. tdTomato$^+$ (tdT$^+$) cells indicate

recombined cells. *b$_1$, b$_2$,* magnified images of ctl and cKO medial prefrontal cortex

(mPFC, highlighted in *b*) stained with PDGFRα and tdTomato expression. *c,*

Recombination efficiency of OPCs in ctl and cKO mPFC. (*ctl=4, cKO=5, unpaired t-*

test). *d,* Western blot analysis of GABA$_B$R subunit 1 in MACs sorted OPCs from

cortex shows reduction of GABA$_{B1}$R expression in cKO OPCs. (*ctl=3, cKO=3,*

unpaired t-test). *e,* Exemplary images of Olig2 (oligodendrocyte lineage), PDGFRα

(OPCs) and CC1 (oligodendrocyte) immunostaining in adult ctl and cKO mPFC. *f, g,*

Quantification of OPC and oligodendrocyte cell densities in mPFC and corpus

callosum (cc). (mPFC: *ctl=7, cKO=8, cc: ctl=4, cKO=4, multiple t-tests). h,*

Quantification of the oligodendrocyte proportion among the total lineage (mPFC:

ctl=4, cKO=4; cc: ctl=3, cKO=6, multiple t-tests). i, Immunostaining of Caspr and

MBP in ctl and cKO mPFC. *j,* Quantitative analysis of paranode length in mPFC

(*ctl=4, cKO=4, multiple t-tests*). *k,* Western blot analysis of MBP expression in mPFC

and cc. (*ctl=4, cKO=3, multiple t-tests*). *l,* Overview of coronal brain slices

immunostained with parvalbumin (PV) and MBP shows abundant PV$^+$ neurons are

myelinated in mPFC. *m,* 3D reconstruction of myelinated PV$^+$ axons in ctl and cKO

mPFC using Imaris software. *n,* Relative volume of MBP$^+$ myelin sheats wrapping

PV$^+$ axons (*normalized to ctl). (ctl=3, cKO=3, unpaired t-test*).

Figure 2. In OPC-GABA$_B$R cKO mice impaired control of interneuronal density

and firing rate reduces oligodendrocyte differentiation and myelin gene

expression at the onset of myelination. *a,* Experimental schedule for *b-d*. *b*,

Immonostaining (left) and quantification (right) of PV$^+$ neurons in the mPFC at the

age of 9 weeks (*ctl=5, cK=9, unpaired t-test*). *c,* Exemplary traces of sIPSCs and

sEPSCs recorded at the layer V of mPFC. *d,* Quantification of frequency and

amplitude of sIPSC and sEPSC *(sIPSC: ctl=15 cells from 4 mice, cKO=15 cells from*

5 mice; sEPSC: ctl=26 cells from 4 mice, cKO=21 cells from 5 mice, Mann Whitney

test). *e,* Experimental schedule for *f-m*. *f,* 3D reconstruction of OPCs and vGAT by

PDGFRα and vGAT immunostaining in ctl and cKO mPFC at p10 using Imaris

software. vGAT was classified into two subgroups: magenta indicates vGATs which

have < 20nm distance from OPC surface; grey vGATs have > 20nm from OPC

surface. *g, h*, Density (*g*) and volume (*h*) of vGAT in ctl and cKO mPFC. (*ctl=4,*

cKO=4; multiple t-tests). *i,* Quantification of synaptic vGAT (< 20nm) density per μm^2

of OPC surface. *(ctl=4, cKO=4, unpaired t-test).* *j,* Immunostaining of OPCs and

oligodendrocytes with PDGFRα, Olig2 and CC1 in p14 mPFC. *k, l*, Density of OPCs

(*k*) and oligodendrocytes (*l*) in ctl and cKO mPFC at p10 and p14. (*p10 and p14:*

ctl=5, cKO=4; multiple t-tests). *m,* Quantitative analysis of MBP mRNA level in ctl

and cKO mPFC at p14. (*ctl=4, cKO=4, unpaired t-test).*

Figure 3. During early CNS development the apoptosis of supernumerous

interneurons is mitigated upon impaired release of TWEAK from GABA$_B$R-

deficient OPC. *a,* Scheme of experimental plan. *b,* Representative images of

excitatory (CTIP) and inhibitory (PV$^+$) neurons co-stained with apoptotic marker CC-3

in p5 and p7 mPFC. *c,* Quantification of apoptotic neurons (CC-3$^+$) co-expressing

interneuron marker GAD67 and PV or excitatory neuronal marker CTIP and TBR1.

(GAD67: p5-ctl=3, p5-cKO=4; p7-ctl=3, p7-cKO=5) (PV: p5-ctl=5, p5-cKO=3; p7-ctl=3,

p7-cKO=5) (CTIP: p5-ctl=7, p5-cKO=4; p7-ctl=9, p7-cKO=3) (TBR1: p5-ctl=4, p5-

cKO=4; p7-ctl=4, p7-cKO=4) (multiple t-tests). *d,* Relative mRNA level of TWEAK in

the mPFC and MACs-OPCs from ctl and cKO mice *(tissue-ctl=4, tissue-cKO=6;*

MACs-ctl=4, MACs-cKO=3; multiple t-test). *e,* Experimental design for *in vitro* studies.

f, Immunolabeling of TWEAK receptor on 14 DIV PV$^+$ interneurons. *g,*

Immunostaining of apoptotic primary PV$^+$ neurons treated with conditional medium of

Oli-neu cells co-treated with or without TWEAKR antagonist L524-0366 (20 µM). *h,*

Percentage of apoptotic PV$^+$ neurons among total PV$^+$ neurons after the treatments

(n=3 independent experiments, one-way ANOVA, multiple comparison).

Figure 4.   In OPC-GABA$_B$R cKO mice perturbed electrical activity translates

into deficits of social behavior. *a,* Scheme of experimental plan. *b,* Representative

diagram of lower gamma band acquired by ECoG recordings. Violin plot shows

decreased gamma oscillation in the cKO mouse brain. (*ctl=24 from 4 mice, cKO=24*

from 4 mice, unpaired t-test). *c,* Impaired nest building ability of mutant mice. (*ctl=9,*

cKO=10, Mann Whitney test). *d*, Scheme of three-chamber behavior test. *e,*

Quantification of the sniffing time with mouse (*e*) or with unfamiliar mouse (*f*) among

total sniffing time (*ctl=9, cKO=9, unpaired t-tests*). *g,* Scheme of new object

recognition test. *h,* Assessment of object preference *(ctl=10, cKO=10). i,* Percentage

of sniffing time with new object among total sniffing time. *(ctl=10, cKO=10, upaired t-*

test). *j,* Representative trajectory charts of the animals during the open field test. *k, l,*

Both ctl and cKO mice exhibited similar motor activities shown by the speed and

distance analysis. (*ctl=10, cKO=14, unpaired t-tests).*

 Supplementary Figure legends

Supplementary Fig.1. OPC and oligodendrocyte density is not altered in the

corpus callosum of mutant mice. a, Scheme of experimental schedule (left panel)

and analyzed brain region in highlight, i.e. corpus callosum (cc, right panel). **b,**

Corpus callosal OPCs and oligodendrocytes of ctl and cKO mice were

immunolabeled with PDGFRα and CC1 respectively at the age of 2w and 9w. Olig2

was used as lineage marker and tdTomato was indicator of recombined cells. **c,**

OPC recombination efficiency in ctl and cKO cc (ctl=86.22±0.57 (n=3),

cKO=76.17±3.10 (n=6), unpaired t-test). **d, e,** Cell density of OPCs (PDGFRα$^+$) and

oligodendrocytes (CC1$^+$) in cc of 2w- and 9w-old mice were not changed. OPC: (2w-

ctl=77.02±4.21 (n=8), 2w-cKO=80±4.72 (n=8), unpaired t-test) (9w-ctl=30.01±3.68

(n=4), 9w-cKO=27.56±2.26 (n=4), unpaired t-test); oligodendrocytes (2w-

ctl=121.70±12.31 (n=8), 2w-cKO=97.62±10.54 (n=8), unpaired t-test) (9w-

ctl=296.90±32.83 (n=4), 9w-cKO=300.30±28.47 (n=4), unpaired t-test). **f,** Percentage

of oligodendrocytes among total Olig2$^+$ cell population, considered as OPC

differentiation rate, was not changed at both ages (2w-ctl=60.26±3.61(n=8), 2w-

cKO=53.98±3.73 (n=8), unpaired t-test) (9w-ctl=92.37±0.12 (n=3), 9w-

cKO=91.33±0.69 (n=6), unpaired t-test).

Supplementary Fig.2. Parameters of myelination remain largely unaltered in the

corpus callosum of mutant mice. a, Scheme of experimental schedule (left panel)

and analyzed brain region in highlight, i.e. mPFC and cc. **b,** Quantification of node

length of ctl and cKO mPFC. **c,** Paranode length is increased in the cKO mPFC. **d,**

Density of nodes remained same in the cKO mPFC. **e,** Relative mRNA level of MBP

is reduced in the cKO mPFC, while PLP mRNA is not affected (ctl: n=4, cKO: n=3,

unpaired t-test). **f,** Immunostaining of Caspr and MBP in cc of ctl and cKO mice. **g-i,**

Paranodal length and node density in ctl and cKO cc remained same. **j,** Electron

microscopic analysis of myelin ultrastructure in sagittal section of cc from ctl and cKO

mice. **k, l,** Quantification of g-ratios of axonal myelination shows unchanged

myelination in cKO cc. **m,** Compound action potential conduction velocity was not

altered indicated by patch clamp recordings. N1 indicates non-myelinated axonal

conduction velocity, while N2 is for myelinated axons.

Supplementary Fig.3. Ablation of GABA$_B$R in oligodendrocytes do not affect

formation and density of oligodendrocytes. a, Scheme of transgene construction.

b, Experimental schedule. **c,** Overview of CC1 immunostaining in a sagittal section of

9w old ctl mouse. Extensive tdTomato distribution shows a high recombination in

different brain regions of transgenic mouse. **d,** Magnified images from prefrontal

cortex (yellow box in **c**) and corpus callosum (white box in **c**). **e,** Quantification of

oligodendrocyte recombination efficiency in PFC were identical, while cKO mice

showed smaller recombination efficiency in cc. However, density of oligodendrocyte

cells did not differ between ctl and cKO mice in both PFC and cc. arrows:

recombined oligodendrocyte; triangles: non-recombined oligodendrocyte.

Supplementary Fig.4. Dense population of parvalbumin-expressing

interneurons in the adult mouse cortex. a, Overview of coronal brain slices

immunostained with parvalbumin and MBP shows abundant PV$^+$ neurons in the

somatosensory cortex. **b,** Magnified view (yellow box in **a**) shows the absence of PV$^+$

axons in the corpus callosum.

Supplementary Fig.5. At the onset of myelination, GABA$_B$R-deficient OPCs

receive less GABAergic input in the mPFC mutant mice. a, Experimental timeline

and analyzed brain region (mPFC) in highlight.**b,** Immunostaining of PDGFRα and

vGAT in the mPFC of p10 mice. **c,** Quantitative analysis of vGAT volume and the

distance to OPC surface (ctl: n=4, cKO: n=4). **d,** The volume of vGAT in cKO mPFC

was smaller than that in ctl group.

Supplementary Fig.6. The differentiation of GABA$_B$R-deficient OPCs is suppressed in the mutant mPFC. a, Scheme of experimental schedule and analyzed brain region (mPFC). **b,** Recombination efficiency of OPCs (% of PDGFRα$^+$tdt$^+$/PDGFRα$^+$) at p14 is identical in the ctl and cKO mPFC. **c,** Immunostaining and quantification of OPCs incorporated with BrdU shows no change in OPC proliferation in the cKO mPFC at p14 (*ctl: n=4, cKO: n=5, unpaired t-test*). **d,** Immunostaining and quantification of oligodendrocytes incorporated with BrdU shows decreased OPC differentiation in the cKO mPFC at p14 (*ctl: n=4, cKO: n=5, unpaired t-test*). **e,** The portion of oligodendrocytes among the oligodendrocyte lineage declined in cKO mPFC at p14 (*ctl: n=5, cKO: n=4, unpaired t-test*).

Supplementary Fig.7. Controlled ablation of GABA$_B$R in OPCs of young adult mice does not affect oligodendrogenesis. a, Scheme of experimental schedule and the analyzed brain regions (mPFC and cc). **b,** Immunostaining of OPCs (PDGFRα) and oligodendrocyte (CC1) with Olig2 (lineage marker) at ctl and cKO mouse mPFC and cc at the age of 9w. **c, d,** Cell densities of OPCs (PDGFRα$^+$) and oligodendrocytes (CC1$^+$) of cKO animals were the same as those for ctl mice in both mPFC and cc. mPFC: *(PDGFRα$^+$: (ctl=17.58±0.53 (n=4), cKO=18.69±1.58 (n=2), unpaired t-test)) (CC1$^+$: (ctl=9.05±1.38 (n=4), cKO=7.95±1.40 (n=2), unpaired t-test));* cc: *(PDGFRα$^+$: (ctl=35.38±2.25 (n=4), cKO=32.37±4.81 (n=2), unpaired t-test)) (CC1$^+$: (ctl=311.81±43.39 (n=4), cKO=339.20±70.53 (n=2), unpaired t-test)).* **e,** the Proportion of oligodendrocyte among total lineage cells did not change, suggesting OPC differentiation remained same in the mutant mouse mPFC and cc. mPFC: *(ctl=30.49±5.13 (n=3), cKO=29.91±2.62 (n=2), unpaired t-test); cc: (ctl=89.52±1.30 (n=3), cKO=91.21±0.471 (n=2), unpaired t-test).***f,** MBP expression is not altered in the cKO mPFC at mRNA level. (ctl: n=4, cKO: n=3, unpaired t-test).

Supplementary Fig.8. The apoptosis of interneurons is mitigated in the mPFC of mutant mice at p14. a, Timeline of experiments and scheme of analyzed brain

region mPFC in highlight. **b,** Immunostaining of PV$^+$ neurons with apoptotic marker

(cleaved caspase-3) at p10 of ctl and cKO mPFC. **c,** Quantification of apoptotic PV$^+$

neurons at p10 and p14 shows reduced PV$^+$ neuronal apoptosis in the cKO mPFC

compared to ctl (p10*: ctl=2.03±0.47 (n=6), cKO=0.92±0.34 (n=3), unpaired t-test;* p14:

ctl=3.70±1.21 (n=6), cKO=2.50±0.72 (n=9), unpaired t-test). **d,** Representative

images of apoptotic neurons at p14 of ctl and cKO mPFC. **e,** Quantification of

apoptotic total neurons (NeuN$^+$CC-3$^+$) at p14 *(ctl=2.76±0.42 (n=12), cKO=1.42±0.29*

(n=9), unpaired t-test).

Supplementary Fig.9. Ablation of GABA$_B$R expression in OPCs does not affect

the apoptosis or survival of excitatory neurons. a, Scheme of experimental

schedule (left panel) and analyzed brain region in highlight. **b,** About 40 % of OPCs

are recombined (% of PDGFRα$^+$tdT$^+$/PDGFRα$^+$) at p7 both in the ctl and cKO mPFC.

c, d, Immunostainings and quantification of excitatory neurons (TBR1$^+$) with

apoptotic marker CC-3 in mPFC of ctl and cKO pups at p5 (**c**) and p7 (**d**) (p5:

ctl=1.64±0.39 (n=4), cKO=1.74±0.24 (n=4), unpaired t-test; p10*: ctl=0.74±0.18 (n=4),*

cKO=0.67±0.16 (n=4), unpaired t-test).

Supplementary Fig.10. TWEAK expression in OPCs is regulated by GABA$_B$Rs.

a, Scheme of experimental design. **b,** Relative mRNA level of cytokines in the mPFC

of ctl and cKO mouse at p7 *(TNF-α: ctl=1±0.34 (n=4), cKO=1.23±0.36 (n=6),*

unpaired t-test) (TWEAK: ctl=1±0.10 (n=4), cKO=0.67±0.08 (n=6), unpaired t-test)

(FasL: ctl=1±0.57 (n=2), cKO=0.91±0.23 (n=4), unpaired t-test) (TNFSF7: ctl=1±1.00

(n=4), cKO=0.93±0.67 (n=6), unpaired t-test) (Trp53: ctl=1±0.11 (n=4),

cKO=1.14±0.08 (n=6), unpaired t-test) (NGF: ctl=1±0.24 (n=5), cKO=1.08±0.18 (n=6),

unpaired t-test). **c,** Western blot analysis of TWEAK expression in Oli-neu cells

treated with 50 µM Baclofen at the presence or absence of 20 µM CGP 55845 (n=5

independent experiments). The isoform in 36 KDa (blue box) was analyzed.

Supplementary Fig.11. OPC-GABA$_B$R cKO mice exhibit impaired social

behavior. a, b, Mutant mice stayed longer at the chamber of mouse than object,

indicating that sociability of the mutant mouse is not affected. **c, d,** Mutant mice spent

similar time at the chambers of familiar and unfamiliar mouse, suggesting impaired

cognition of the cKO mice (ctl (n=9), cKO (n=9), paired t-test).

Figure 1

Figure 2

Figure 3

a

birth — p1 2 — 5 — 7 — day — ▼ tam / ▼ analysis

b

| merge (ctl-p5) | CTIP / CC-3 | merge (ctl-p7) | PV / CC-3 |
| cKO-p5 | | cKO-p7 | 5 μm |

c

Inhibitory neurons — Excitatory neurons

GAD67+: p5 0.027, p7 <0.001
PV+: p5 0.011, p7 0.014
CTIP+: p5 0.547, p7 0.496
TBR1+: p5 0.822, p7 0.774

CC-3+ cells/1x10^-2mm^3

● ctl   ● cKO

d

relative Tweak mRNA level (normalized to ctls)

Tissue 0.034, MACs-OPC 0.068

e

Oli-neu → conditioned medium (CM) → neuron ± L524-0366

f

TWEAKR | PV | merge

g

PV/CC-3

ctl+DMSO | CM+DMSO | CM+L524 | 10 μm

h

% of PV+CC-3+/PV+

CM: − + +
L524: − − +

0.012   0.047

Figure 4

Supplementary figure 1

Supplementary figure 2

Supplementary figure 3

a

PLP promoter → CreERT2 x GABA_B — *loxP* ▶ Exon 7 — Exon 8 — *loxP* ▶

x Rosa26 — CAG — *loxP* ▶ STOP — *loxP* ▶ RATO

b

p7/8 — 4w — 9w — time

▼ 2 x tam
▼ 5 x tam
▼ analysis

c

CC1/ tdTomato
4w-9w
500 µm

d

CC1 | tdT | Merge

PFC

cc

50 µm

e

PFC

% of CC1⁺tdT⁺/CC1⁺ cells vs 1w, 4w
ctl (blue), cKO (red)

CC1⁺ cells/1x10⁻³ mm³ — 1w: 0.994, 4w: 0.114

CC

% of CC1⁺tdT⁺/CC1⁺ cells vs 1w, 4w

CC1⁺ cells/1x10⁻³ mm³ — 1w: 0.745, 4w: 0.999

Supplementary figure 4

Supplementary figure 5

a

birth   p7   p8   p10

▼ tam injection
▼ analysis

b

PDGFRα /vGAT

ctl   cKO

5 µm

c

vGAT volume (µm³)

Distance to OPC surface (µm)

d

% of volume distribution

Volume (µm³)

ctl
cKO

Supplementary figure 6

Supplementary figure 8

Supplementary figure 9

a birth p1 p2 p5 p7 ▼ tam injection ▼ analysis

b PDGFRα / tdTomato — PFC — ctl, cKO — Pα, tdT — 50 μm — Recombination efficiency — ● ctl ● cKO

c TBR1/CC-3 — p1/2-p5-TBR1 — ctl, cKO — TBR1, CC-3 — 5 μm — CC-3⁺ TBR1⁺ cells/1x10⁻² mm³ — 0.822

d TBR1/CC-3 — p1/2-p7-TBR1 — ctl, cKO — TBR1, CC-3 — 5 μm — CC-3⁺ TBR1⁺ cells/1x10⁻² mm³ — 0.801

Supplementary figure 10

a

birth p1 p2 p7

▼ tam injection
▼ analysis

b

Relative mRNA expression

TNF-α TWEAK FasL TNFSF7 Trp-53 NGF

0.675 0.034 0.865 0.936 0.282 0.796

☐ ctl
☐ cKO

c

Bac ± CGP 55845

Oli-neu

Bac - + + - + +
CGP - - + - - +

Exp. 4 Exp. 5

KDa   KDa
48   48
35   35
25   25

TWEAK

α-tubulin   48

relative TWEAK level
(normalized to ctl)

0.254
0.043

● ctl
● Bac
● Bac + CGP

Supplementary figure 11