Running Head: Allopolyploid networks with phased sequences

Benefits and Limits of Phasing Alleles for Network Inference of Allopolyploid Complexes

George P. Tiley[1,*], Andrew A. Crowl[2], Paul S. Manos[2], Emily B. Sessa[3], Claudia Solís-Lemus[4],

Anne D. Yoder[2], J. Gordon Burleigh[3]

[1]Royal Botanic Gardens Kew, Richmond, TW9 3AE, UK

[2]Department of Biology, Duke University, Durham NC, 27708, USA

[3]Department of Biology, University of Florida, Gainesville FL, 32611, USA

[4]Wisconsin Institute for Discovery and Department of Plant Pathology, University of Wisconsin –

Madison, Madison WI, 53706, USA

[*]Author for correspondence: george.tiley@gmail.com

Abstract

Accurately reconstructing the reticulate histories of polyploids remains a central challenge for understanding plant evolution. Although phylogenetic networks can provide insights into relationships among polyploid lineages, inferring networks may be hindered by the complexities of homology determination in polyploid taxa. We use simulations to show that phasing alleles from allopolyploid individuals can improve phylogenetic network inference under the multispecies coalescent by obtaining the true network with fewer loci compared to haplotype consensus sequences or sequences with heterozygous bases represented as ambiguity codes. Phased allelic data can also improve divergence time estimates for networks, which is helpful for evaluating allopolyploid speciation hypotheses and proposing mechanisms of speciation. To achieve these outcomes in empirical data, we present a novel pipeline that leverages a recently developed phasing algorithm to reliably phase alleles from polyploids. This pipeline is especially appropriate for target enrichment data, where depth of coverage is typically high enough to phase entire loci. We provide an empirical example in the North American *Dryopteris* fern complex that demonstrates insights from phased data as well as the challenges of network inference. We establish that our pipeline (PATÉ: Phased Alleles from Target Enrichment data) is capable of recovering a high proportion of phased loci from both diploids and polyploids. These data may improve network estimates compared to using haplotype consensus assemblies by accurately inferring the direction of gene flow, but statistical non-identifiability of phylogenetic networks poses a barrier to inferring the evolutionary history of reticulate complexes.

INTRODUCTION

Polyploidy, or whole-genome duplication, occurs throughout the eukaryotic tree of life.

However, its evolutionary significance is especially evident in plants, with recent estimates

suggesting up to 35% of vascular plant species are of recent polyploid origin (Wood et al. 2009;

Barker et al. 2016). Despite advances in genomic data generation and numerous studies

investigating the role of whole-genome duplication in plant speciation and local adaptation

(reviewed in Soltis et al. 2014), polyploidy remains a central challenge for the field of

phylogenetics. One persistent problem when analyzing sequence data from polyploid taxa, and

especially allopolyploids, is identifying the alleles and divergent homeolog copies from parental

lineages. Most bioinformatic tools for processing next generation sequence data were

developed with diploids, or specifically humans, in mind. These approaches often collapse

variable homeolog sequences into a single consensus sequence for *de novo* assemblies or

assume the organism is diploid when performing genotyping and phasing for reference-based

assembly. For polyploids, this can create chimeric sequences that may interfere with

phylogenetic reconstruction and obscure evidence of polyploidy and its origins. Using allelic

data that more accurately capture the complex genomic histories of polyploids should enable

analyses to incorporate divergent signals from polyploid loci into phylogenomic inference,

distinguish allopolyploidy from autopolyploidy, and identify parental taxa. However, few studies

have examined the potential benefits of using phased versus unphased data to reconstruct

polyploid histories (Kamneva et al. 2017; Oberprieler et al. 2017; Eriksson et al. 2018), and

there are few formal methods and little guidance for phasing alleles from polyploid taxa. Here

we explore the value of using phased data to reconstruct polyploid networks, leveraging fast

polyploid phasing algorithms (Xie et al. 2016) to develop a bioinformatic pipeline that can phase

alleles from polyploids using target enrichment sequence data.

Previous studies have suggested phasing alleles is crucial for accurate evolutionary

reconstruction of reticulate complexes, at least when sampling relatively few loci (e.g., 4 to 10;

Rothfels et al. 2017; Eriksson et al. 2018). However, it remains challenging to genotype (e.g.

Blischak et al. 2018b) and phase next generation sequencing data from polyploids using short

reads, in which individual variants may lack physical linkage information (e.g. Schrinner et al.

2020). Methods exist to genotype consensus loci from target enrichment data, but these have

been either limited to diploids (Kates et al. 2018; Andermann et al. 2019) or used manual

curation of variants with polyploids where both parental populations are available (Eriksson et

al. 2018). Otherwise, obtaining phased sequence data for polyploids has largely depended on

long-read sequencing to recover complete haplotype sequences (e.g., Rothfels et al. 2017) or

cloning PCR products (e.g., Sessa et al. 2012; Oberprieler et al. 2017).

Target enrichment (or HybSeq), in which specific regions of the genome are isolated and

sequenced (Albert et al. 2007; Gnirke et al. 2009; Faircloth et al. 2012; Lemmon et al. 2012), is

an increasingly common method for collecting large-scale phylogenomic datasets, and these

data can provide insights into the evolutionary history of reticulate complexes (e.g., Crowl et al.

2020; Karimi et al. 2020) and sources of gene tree discordance (e.g., Crowl et al. 2017;

Morales-Briones et al. 2018; Stull et al. 2020). Probe kits for target enrichment have been

developed in many land plant lineages (Wolf et al. 2018; Johnson et al. 2019; Liu et al. 2019;

Breinholt et al. 2021), and there are bioinformatic pipelines available for custom probe design

(e.g., Jantzen et al. 2020). The most common approach to assemble phylogenetic datasets from

target enrichment data has been to use haplotype consensus assemblies from various pipelines

(e.g. Faircloth 2016; Johnson et al. 2016; Andermann et al. 2018; Breinholt et al. 2018). The *de*

novo assembly algorithms within these pipelines often consider only the most frequent

nucleotide sequence and discard the alternatives (Bankevich et al. 2012; Iqbal et al. 2012; Luo

et al. 2012). This results in loci in which variable positions are collapsed to a single base call

(haplotype consensus loci), losing information related to heterozygosity. While this may be

appropriate, or at least benign, for phylogenetic analyses of diploid taxa (Kates et al. 2018), it

may pose substantial problems when attempting to investigate the evolutionary history of

polyploid taxa or reticulate lineages. Some of the before-mentioned pipelines can recover

heterozygosity information and phase sequences by mapping reads back to their haplotype

consensus sequences (Andermann et al. 2018), but this ability is, at the time of writing, limited

to diploids.

Recovering phased haplotype sequences from polyploid individuals, especially

allopolyploids with distinct subgenomes, should be valuable when estimating phylogenetic

networks that likely are more accurate representations of reticulate evolution than a bifurcating

tree. Simulations and empirical analyses have suggested that phylogenetic networks can

recover the reticulate histories of polyploid lineages with few loci, at least when gene tree

discordance due to incomplete lineage sorting (ILS; Hudson et al. 1983; Pamilo and Nei 1988)

is low (Oberprieler et al. 2017) using parsimony methods (Huber et al. 2006; Lott et al. 2009), or

with moderate ILS when it is explicitly modeled (Jones et al. 2013). Contemporary phylogenetic

network models and software packages jointly consider gene tree variation due to allele

sampling error as described by the multispecies coalescent (MSC; Rannala and Yang 2003)

and gene flow modeled as episodic introgression events (Solis-Lemus and Ané 2016; Wen et al.

2016; Zhang, Ogilvie et al. 2018, Flouri et al. 2020). Depending on the complexity and goals of

the research question, these methods can search for networks with a constrained number of

reticulation events using quartet-based maximum pseudolikelihood (Solis-Lemus and Ané 2016;

Wen et al. 2018) or a full-likelihood Bayesian model where the number of reticulations is a

parameter (Wen et al. 2018; Zhang et al. 2018). Also, it is possible to estimate model

parameters (e.g., divergence times, population sizes, and the fraction of introgressed genes) on

a fixed species network using a full-likelihood Bayesian model that allows efficient computation

with large numbers of loci (Flouri et al. 2020). We refer to these network models from here on as

the multispecies coalescent with introgression (MSci), consistent with Flouri et al. (2020),

although other names have been used, such as the network multispecies coalescent (NMSC;

Zhu and Degnan 2017) and multispecies network coalescent (MSNC; Wen et al. 2016). We

emphasize that using network approaches to investigate polyploid complexes is not novel (e.g.,

Huber et al. 2006; Lott et al. 2009; Jones et al. 2013; Crowl et al. 2017), but the difficulty of

collecting appropriate genomic data from polyploids for such analyses has limited their use.

To address the issues outlined above, we have developed a pipeline, PATÉ (Phased

Alleles from Target Enrichment data), that can phase genotyping data for individuals of a known

ploidy without the need for sampling their parental lineages. PATÉ was designed with scalability

and population-level sampling in mind for target enrichment projects where deep coverage from

paired-end Illumina data allows calling of high-quality variants. In this study, we first use

simulations to explore the ability of network approaches to reconstruct the history of

allopolyploidy in the presence of ILS, and whether phasing the data affects the accuracy of the

reconstruction. We show that using phased allelic data can improve network estimation and

divergence time estimation compared to using haplotype consensus sequences, but also

highlight scenarios where phasing may not be necessary or beneficial. For an empirical

example of how PATÉ output can be used in studies of reticulate evolution, we compared

phased (either all alleles or only one) and unphased (either haplotype consensus or IUPAC

ambiguity codes) data to infer the evolutionary history of the North American *Dryopteris* fern

complex, a model system for reticulate polyploid evolution (Sessa et al. 2012a; Sessa et al.

2012b), using new targeted enrichment data. The system includes four diploid species, as well

as one extinct diploid, that have formed five allopolyploids in which there is high confidence in

the parent-progeny relationships (Fig. 1), although numerous sterile allopolyploid species have

also been reported within the complex (Montgomery and Paulton 1981). The allopolyploid

species have relatively ancient origins, with the best estimates placing hybridization events

between six and 13 Ma (Sessa et al. 2012b). PATÉ is capable of recovering phased haploid

sequences almost always the length of the original haplotype consensus assembly from

polyploid individuals, although the switch errors among these haplotype sequences is unknown.

Networks inferred from phased data more accurately represent some relationships within the

North American *Dryopteris* complex than unphased strategies, but statistical identifiability

problems confound all data types.

Figure 1 — Hypothesized Relationships among North American *Dryopteris*. Synthesis of results from Sessa et al. 2012a and Sessa et al. 2012b. A) Links between shapes show the putative parents and their allopolyploid derivatives. Black circles are diploids, squares are tetraploids, and the hexagon is the one hexaploid species in the group. *Dryopteris semicristata* is presumed extinct. B) Placement of allopolyploids in the context of the backbone relationships among diploids. Tetraploids are indicated with solid orange lines and the hexaploid with dotted lines. The grey line denoting a sister relationship between *D. semicristata* and *D. expansa* reflects one possible placement for the extinct taxon based on previous analyses (Sessa et al., 2012b).

MATERIALS & METHODS

Testing the Effects of Phasing on Network Inference through Simulation

Simulating phased and unphased sequence data for an allopolyploid — We simulated gene trees and their nucleotide sequence data using the MSC model (Rannala and Yang 2003) with BPP v4.4.0 (Flouri et al. 2020) under a network with five tips, such that one taxon was a hybrid of the ancestors of two other tip taxa (Fig. 2). All simulations used the HKY85 (Hasegawa et al 1985) model of sequence evolution with a transition-transversion rate ratio of 3 and

gamma-distributed among-site rate heterogeneity with a shape parameter of 0.6. Equilibrium

frequencies were 0.3, 0.2, 0.3, and 0.2 for nucleotides A, C, G, and T. These parameters for the

sequence evolution model were based on the observed means of parameter distributions

across gene trees for our empirical *Dryopteris* data (see Materials and Methods: *Analyses of a*

Species Complex with Allopolyploidy). The allopolyploid species *E* was treated as two lineages

(*E* sister to *B* and *F* sister to *C*) whose alleles were pooled to form the hybrid species *E* at time

τ_h. This makes species *E* a tetraploid hybrid with the parents the ancestors of *B* and *C*. The

mutation-scaled effective population size (θ) was constant among lineages and set at 0.01.

Assuming a per-generation mutation rate (μ) of 1×10^{-8} and one year per generation yields an

effective population size (N_e) of 250,000 and hybridization age of 10 Ma (Fig. 2). We also

simulated data such that θ was 0.001 (hybridization at 1 Ma) or 0.1 (hybridization at 100 Ma).

Investigating such extreme values can generate general expectations for downstream

estimation under a wide range of parameters, but it also has some biological salience as there

is increasing evidence for reticulate evolution early in the diversification of plant orders and

families (Sun et al. 2015; Stull et al. 2020; Liu et al. 2022).

Figure 2 — Species network used for simulation. The divergence times in expected substitutions per site are given for each node, and *h* is the hybrid node where two alleles enter from both *t* and *v*. E is an allotetraploid while other species are diploid. Absolute divergence times were changed by using values of θ at 0.001, 0.01, and 0.1. ILS was increased by multiplying θ by a factor of 2, 3, or 4 while keeping all τ_u fixed to their initial values.

Because the distance between speciation nodes is 0.01 substitutions per site and $\theta =$

0.01 (Fig. 2), there are two coalescent units between nodes, $T = \frac{\tau}{\left(\frac{\theta}{2}\right)}$ (Yang 2006), which yields a

low probability (0.09) of gene tree discordance due to ILS (Hudson 1983). We incorporated

more ILS into the simulation by keeping the node heights fixed but increasing θ to 0.02, 0.03,

and 0.04. This increased the probability of a non-speciodentric split among all nodes to 0.25,

0.34, and 0.4, respectively. With three different starting values of θ and τ, this gave a total of 12

simulation conditions. While our simulations do not explore the limits of gene tree discordance,

they allow us to learn about some general features of increasing ILS on network inference with

different data types. We simulated 1000 gene trees and their sequences, 500 base pairs (bp) in

length, under the MSC 100 times for each of the 12 conditions. We also explored the effects of

sampling fewer genes on downstream analyses. For each replicate of 1000 gene trees and their

sequences, we randomly sampled 400, 40, and four genes without replacement.

All simulations sampled haploid data. Two haploid sequences were sampled for each

diploid species, and four haploid sequences were sampled for the allopolyploid species *E,*

where two sequences came from each parental lineage. We then investigated unphased data in

three ways. First, to generate unphased genotype sequences, the simulated haploid sequences

for each species were collapsed into a single sequence in which heterozygous sites were

represented by IUPAC ambiguity codes (genotype). The allopolyploid species was not restricted

to only biallelic sites. Second, we generated haploid consensus sequences where for each

variable site only one base was randomly chosen and retained (consensus). This could

represent a case in which read coverage across a locus is highly uneven such that a haploid

sequence is actually a chimera of two or more alleles. Finally, we simply picked one phased

haploid sequence, which is possible when one parental haplotype has a majority of reads for a

locus (pick one). This scenario where only one parent's sequence would be recovered in the

offspring could be anticipated in real data due to subgenome dominance (e.g., Buggs et al.

2014; Bird et al. 2018). In practice, we expect most *de novo* assemblers to generate output in

between the haploid consensus data and pick one data.

Inferring species networks with phased and unphased simulated data — We estimated

species networks with the SNaQ function (Solís-Lemus and Ané 2016) from PhyloNetworks

v0.12.0 (Solís-Lemus et al. 2017) using Julia v.1.4.1 (Bezanson et al. 2015) from either the true

gene trees used to simulate the data or gene trees estimated from the phased, pick one,

genotype, and consensus sequence data. For estimated trees, we used IQTREE v1.6.10

(Nguyen et al. 2015) with the same substitution model used to simulate the data. Each SNaQ

analysis used the species tree (A,((B,E),(C,D))) as the starting tree and allowed zero, one, or

two reticulation events. Each analysis included ten independent optimizations of the

pseudolikelihood score. We considered increasing numbers of reticulations if there was a

pseudolikelihood score improvement of two or greater compared to the previous model with one

fewer allowed hybrid edges. Note that pseudolikelihoods violate standard likelihood-based

model selection criteria such as likelihood ratio tests or using AIC weights, and slope heuristics

(Baudry et al. 2012) are often used instead. The choice of a cutoff here was necessary to

streamline processing across thousands of simulations. We compared the estimated networks

with one reticulation to the true network with the *hardwiredClusterDistance* function (Huson et

al. 2010) in PhyloNetworks. This allowed us to score the number of replicates that 1) recovered

the correct number of reticulations and 2) matched the true network when the number of

reticulations was set to one. We estimated networks for samples of four, 40, 400, and 1000

gene trees for each of the 100 replicates from the 12 simulation conditions.

Effect of phasing on divergence time estimation — We also used our simulated phased

and unphased data to estimate divergence times under the MSci model (Flouri et al. 2020)

using BPP v4.4.0. Here, we estimate divergence times (τ), population sizes (θ), and the

introgression probability, or proportion of immigrant loci in the receiving population, (φ) on the

correct fixed species network. φ was originally represented by γ (Meng and Kubatko 2009) and

referred to as an inheritance probability (Yu et al. 2011), but we use φ and the term

introgression probability to be consistent with BPP (Flouri et al. 2020). Our MSci analyses used

diffuse priors on τ_r and θ, with a mean on their simulated values and $\varphi \sim \beta(1,1)$. Phased, pick

one, and consensus sequences were treated as haploids while genotype sequences were

treated as unphased diploids and used for analytical integration over phases (Gronau et al.

2011) implemented in BPP. Although this is not correct for the tetraploid, it is arguably more

appropriate than treating all of the genotype sequences as haploids with ambiguous states.

Each Markov chain Monte Carlo (MCMC) analysis collected 10,000 posterior samples, saving

every 100 generations, while discarding the first 100,000 generations (i.e., 10% of the total run)

as burnin. Scripts for simulation and subsequent analyses of simulated data are available in

Dryad (https://doi.org/10.5061/dryad.5qfttdz53).

A Phasing Pipeline for Polyploids

Target enrichment data — The success of empirical studies in which phasing was

informative about hybridization or introgression events (e.g., Kamneva et al. 2017; Oberprieler

et al. 2017; Eriksson et al. 2018) motivated us to use phased data to infer reticulate evolutionary

histories of polyploids. We were aware of few instances of phasing genomic or phylogenomic

data in polyploids, except in cases where chromosome-level whole-genome assemblies have

characterized subgenomes in allopolyploid crops (Yang et al. 2017; Colle et al. 2019) or tree-

based methods that are dependent on the sampling of parental lineages (Freyman et al. 2020;

Nauheimer et al. 2020). We designed PATÉ for target enrichment data because of the

availability of such data for many taxa, but it is applicable to other types of data with paired-end

Illumina reads.

The end product of many *de novo* target enrichment assembly pipelines (such as

HybPiper; Johnson et al. 2016) is a single consensus sequence for each locus for each

individual. Allelic variation may be represented by ambiguous nucleotide codes within the single

consensus sequence or lost when the pipeline outputs the haplotype consensus sequence

where the majority vote from a collection of reads is used. We use these existing *de novo*

assembly pipelines as a starting point to provide the reference sequence for each locus for each

individual and leverage a recent phasing algorithm with high-quality variant calls to recover

phased haplotype sequences for taxa with known ploidy levels. Ploidy levels are well

established and relationships have previously been characterized for individuals in our

Dryopteris analyses (Sessa 2012b).

Phasing alleles within loci — PATÉ (Supplementary Fig. S1) starts with assembled

target enrichment loci, such as the supercontig files output from HybPiper (Johnson et al. 2016),

that contain a single haplotype consensus sequence from each individual per locus. Reads for

each individual are then realigned to their consensus locus using BWA v0.7.17 (Li and Durbin

2009). PCR duplicates are flagged with MarkDuplicates in Picard v2.9.2

(http://broadinstitute.github.io/picard), and variant calls are computed with HaplotypeCaller in

GATK v.4.1.4 (McKenna et al. 2010). We applied the following hard filters with VariantFiltration

in GATK: (1) the variant quality divided by the depth of the alternate allele (QD) should be two or

greater; (2) a Fisher Exact test statistic based on a contingency table of read direction and

whether that read has the reference or alternate allele (FS) should be 60 or less; (3) the root

mean square of mapping quality scores (MQ) should be 40 or greater; (4) an approximation of

the Rank Sum Test for variant position in a read should be negative eight or greater to avoid

variants consistently towards the end of reads; (5) the fraction of reads with the alternate allele

(AF) should be between 5% and 95%, and the total read depth of the site (DP) should be at

least ten. The filter expressions are as follows: (1) $QD < 2.0$, (2) $FS > 60.0$, (3) $MQ < 40.0$, (4)

ReadPosRankSum < -8.0, (5) $AF < 0.05$, $AF > 0.95$, (6) $DP < 10$. These loosely follow

community recommendations on filters for germline variant discovery (DePristo et al. 2011).

Notably, we do not perform quality score recalibration or filter on the mapping quality rank sum,

as we anticipate allopolyploids could have a lower mapping quality associated with an alternate

allele due to sequence divergence or structural variation among homeologous chromosomes.

We also consider a very narrow window for filtering on allele frequency. Because increasing

ploidy levels will generate smaller anticipated ratios of alternate to reference alleles, coupled

with sequencing error and read stochasticity, we only aim to remove the most extreme cases.

For example, if almost all reads support the alternate allele at a site, it is difficult to diagnose if

the error lies in the consensus assembly or the read alignment. In these cases, only the

reference site is retained, and the variant does not pass the allele frequency filter. However, the

allele frequency filter could be removed if investigators are focused on organisms with higher

ploidy levels. Additional options for pipeline configuration are available on GitHub

(https://github.com/gtiley/Phasing).

Biallelic SNPs that pass filters are then phased with H-PoPG v.0.2.0 (Xie et al. 2016). H-

PopG solves a heuristic phasing problem efficiently using dynamic programming. Although not

guaranteed to be an optimal solution, H-PoPG has been shown to have high accuracy while

also being fast (Xie et al. 2016; He et al. 2018; Moeinzadeh et al. 2020). Phasing variants in

polyploids is difficult because for n variants and k ploidy, there are $2^{n-1}(k-1)^n$ possible ways

to link the sites together. H-PoPG evaluates possible solutions efficiently by grouping reads into

k groups in such a way that differences within the groups are minimized. Focusing on target

enrichment data also constrains the complexity of the phasing problem compared to whole-

genome alignments because haplotype blocks are constrained to about 1000 bp. We then used

the phased variants to create individual allele sequences, where invariable sites are filled in

based on the reference sequence. In cases where there is no linkage information to phase

across the entire locus, we retain the phasing only for the longest block. The variants for the

shorter haplotype blocks can be collapsed into IUPAC ambiguity codes or treated as missing

data based on the investigator's preferences. PATÉ outputs analysis-ready fasta files with

multiple alleles per species. Variants are only phased within loci; we do not attempt to assign

loci to parental subgenomes. While this may complicate analyses of concatenated multi-locus

datasets, it is ideal for the MSC that assumes free recombination between loci and can leverage

multiple alleles per species for estimating θ. Assigning alleles to parental subgenomes should

be possible with PATÉ outputs where both parental lineages are sampled and there is sufficient

divergence for gene tree estimation.

Analyses of a Species Complex with Allopolyploidy

The North American wood fern complex (Dryopteris) — We tested PATÉ using new

target enrichment data from nine North American *Dryopteris* species, including both

allotetraploid and allohexaploid taxa, with well-studied reticulate relationships (Sessa et al.

2012a, b), as well as two outgroup taxa from the sister genus *Polystichum*. All putative parental

lineages are represented in our dataset, with the exception of a hypothesized extinct lineage (*D.*

semicristata; Sessa et al. 2012b). We sampled two or three individuals for each *Dryopteris*

taxon (Table S1). The target enrichment data were generated from the GoFlag 408 flagellate

land plant probe set (Breinholt et al. 2021) at RAPID Genomics (Gainesville, FL). The target

regions for this probe set are 408 exons found in 229 single or low-copy nuclear genes. We

generated haplotype consensus assemblies for each with HybPiper (Johnson et al. 2016). The

resulting supercontig sequences, which includes the exon region targeted by probes for

enrichment and flanking intron regions, became our reference sequences for genotyping and

phasing with PATÉ. We aligned both phased and haplotype consensus (i.e., the reference

supercontig) sequences with MUSCLE v.3.8.31 with default settings (Edgar 2004). For

comparison with simulation results, we generated a "pick one" dataset where only one phased

haplotype per locus was randomly selected from each individual. A "genotype" dataset was

generated by collapsing allelic variation into IUPAC codes for each individual as a locus.

Three species tests — We first explored the value of phasing data when estimating

reticulate relationships among three species, where we sampled two extant parental lineages

and their putative hybrid. We examined the three such extant relationships in the North

American *Dryopteris* complex:

    1) Two diploid parental lineages (*D. ludoviciana* and *D. goldiana*) and their putative

332
       allotetraploid descendent (*D. celsa*).

2) Two diploid parental lineages (*D. expansa* and *D. intermedia*) and their putative

allotetraploid descendent (*D. campyloptera*).

3) One tetraploid parent (*D. cristata*), one diploid parent (*D. goldiana*), and their putative

allohexaploid descendent (*D. clintoniana*).

These three scenarios represent different ages of parental divergence, with a common ancestor

near 10 Ma for *D. ludoviciana* and *D. goldiana*, 15 Ma for *D. expansa* and *D. intermedia*, and 30

Ma for *D. cristata* and *D. goldiana* (Sessa et al. 2012b). We used both a full-likelihood Bayesian

approach and a topology-based pseudolikelihood approach to estimate the correct species

relationships from the different data types. First, using BPP v.4.4.0 (Flouri et al. 2020), we

estimated log-marginal likelihoods (ln *mL*) with stepping-stone sampling (Xie et al. 2011) for the

three possible rooted 3-taxon trees and network models that imply differences in the timing and

direction of gene flow among species (Supplementary Fig. S2). Each ln *mL* estimate used 24

steps, and each step had a posterior sample of 10,000, saving every 30 generations after a

30,000 generation burnin (10% of the total run). The ln *mL* values were then used to calculate

the six model probabilities following equation 1 using the bppr v.0.6.3 R package

(https://github.com/dosreislab/bppr). A percentile-based 95% confidence interval was generated

for each model probability by bootstrapping posterior distributions 1,000 times to investigate

uncertainty in model selection.

352
$$P(model) = \frac{exp(ln\ mL_{model}\ -\ ln\ mL_{max})}{\Sigma_i[exp(ln\ mL_i\ -\ ln\ mL_{max})]} \quad Equation\ 1$$

We performed four independent runs for each model for each data type prior to marginal

likelihood estimation to evaluate mixing and diagnose potential convergence issues. The

combined posteriors from the models that represent *a priori* hypotheses of relationships

between allopolyploids and their parental lineages were used estimate all τ and θ for

comparisons across data types. Notably, BPP can analytically integrate over possible phases of

sequence data when given IUPAC codes representing heterozygous sites (Gronea et al. 2011),

which is as accurate as perfectly phased data for diploids (Huang et al. 2022). In comparison,

standard gene tree estimation under the Felsenstein likelihood (Felsenstein 1981) treat IUPAC

codes as uncertainty and integrate over the possible tip states at sites independently.

For our topology-based method, we used the SNaQ function (Solís-Lemus and Ané 2016) from

PhyloNetworks v0.12.0 (Solís-Lemus et al. 2017) to test the presence and placement of gene

flow between the three species, while also including the two *Polystichum* outgroups. Gene trees

were estimated for each data type across the three cases with IQ-TREE v1.6.10 (Nguyen et al.

2015), using ModelFinder (Kalyaanamoorthy et al. 2017) for model selection on the

unpartitioned alignments. The starting tree was obtained with ASTRAL III v5.6.3 (Zhang et al.

2018) from the collection of ML gene trees. We tested the presence of zero, one, or two

reticulations based on a two pseudolikelihood point cutoff for accepting an additional

reticulation. Each analysis used ten independent optimizations. For SNaQ and ASTRAL

analyses, all haplotypes were mapped to their respective species, as opposed to individuals.

Nine species tests — We investigated the differences in networks from SNaQ across

data types for the nine-species complex. This complex involves multiple reticulation events on

an edge, and thus the true network is not identifiable with topology-based methods (Solis-

Lemus and Ané 2016). Although full-likelihood methods should not suffer from the some of the

identifiability issues (Yu and Nakhleh 2015), they can impose a heavy computational burden

(Wen et al. 2018; Zhang et al. 2018). Despite the inherent identifiability issues, it is worthwhile to

examine the performance of scalable topology-based methods in such contexts to see if they

will identify some of the correct reticulate relationships and if they also recover erroneous ones.

Thus, we performed analyses of the nine-species complex and two outgroups across the four

data types with SNaQ as described above, but we allowed up to six reticulation events. We

used ASTRAL III v5.6.3 (Zhang et al. 2018) to generate the starting species tree for network

estimation using gene trees inferred from IQ-TREE v1.6.10 (Nguyen et al. 2015) with the best

model selected by ModelFinder (Kalyaanamoorthy et al. 2017). 100 standard bootstrap

replicates were performed for each gene tree, which were used to evaluate gene tree

incongruence and informativeness with respect to the ASTRAL species tree from phased data

for the consensus, genotype, and pick one gene trees with phyparts (Smith et al. 2015). To

assess node support for SNaQ, we performed 100 non-parametric bootstrap replicates, in which

both loci and bootstrap trees per locus were sampled with replacement. Each bootstrap

replicate was distributed independently to hasten the computation using a Perl script that

accommodates multiple alleles available on GitHub (https://github.com/gtiley/bootSNaQ-MA).

RESULTS

Simulation Shows Benefits of Phased Data

Network inference — In our simulation results, all data types performed well estimating the

correct number of reticulations (Supplementary Fig. S3). The analyses typically converged to

the true number of reticulations with 40 or more gene trees, except when using gene trees

estimated from genotype data with a 25% ($\theta = 2 \times \tau_h$) chance of a discordant gene tree.

However, using phased data provided more accurate estimates of the placement of the

reticulation edge in comparison to the genotype, consensus, and pick one data (Fig. 3). When

the true gene trees were used, which have information about the allopolyploid's hybridization

event (i.e., the allele sequences are sister to their respective parents in every tree), the correct

network can be inferred with 40 loci regardless of the age of hybridization or degree of ILS. The

gene trees estimated from phased data perform equally well, aside from requiring a few more

gene trees for recent hybridization events with high ILS ($\theta = 4 \times \tau_h$). Analyses based on

genotype, consensus, and pick one data almost always recover the true network when sampling

400 or more loci, but can miss the true network over half the time with 40 or fewer loci.

However, the genotype data failed to converge to the true network with increasing loci for a

hybridization age of 10 Ma with low to moderate amounts of ILS. Otherwise, varying degrees of

ILS had a negligible effect on network estimation among data types, and estimated gene trees,

aside from those from phased data, were less efficient when the hybridization age was more

recent.

Figure 3 — Proportion of simulation replicates that correctly identify the allopolyploid lineage using the SNaQ function of PhyloNetworks. The x-axes are the number of loci sampled for each simulation. The y-axes are the proportions of correct networks. Results are based on networks estimated with a single reticulation, even if that network was considered less optimal than networks with zero or two reticulations. We saved the true gene trees from the simulations, while we estimated gene trees with the phased, genotype, consensus, and pick one data types.

Divergence times — Using phased data, whether using all alleles or only one, typically resulted in accurate estimation of the hybridization age (Fig. 4). When divergence was low, there was a small amount of estimation error, such that the phased data underestimated and the pick one data overestimated the hybridization age, but the pattern suggests estimates could converge to the true values with more loci. Interestingly, convergence towards the true value appeared more efficient with higher ILS (Fig. 4). The phased and pick one data could otherwise accurately estimate the hybridization time with 40 or fewer loci across all other divergence and ILS scenarios. For analyses with genotype and consensus data, as the number of loci increased and uncertainty in the posterior was reduced, the posterior mean did not converge to the true

estimate, and the simulated value was not within the highest posterior density (HPD) interval.

There was always a small degree of estimation error with the genotype data despite using

analytical phasing, and the consensus data overestimated the hybridization age between 1.5x

and 2x when using 400 loci across all simulation scenarios. These biases from genotype or

consensus data were also propagated for the parent nodes of the hybridization event across all

levels of ILS for $\tau_h = 0.001$ (Supplementary Figs. S4-S7), $\tau_h = 0.01$ (Supplementary Figs. S8-

S11), and $\tau_h = 0.1$ (Supplementary Figs. S12-S15). Nodes from speciation events could always

be estimated accurately with high precision by the phased and pick one data types provided

enough loci were sampled. The phased data was more efficient, always converging to the true

value by 40 loci, while the pick one data required 400 to recover τ_u when the sequence

divergence was low at $\tau_h = 0.001$. There was a small degree of estimation error among

speciation nodes for the genotype and consensus data, and the patterns suggested they might

not converge to the true values with more loci. However, speciation times were overestimated

by genotype and consensus data for $\tau_h = 0.001$ (Supplementary Figs. S4-S7) and $\tau_h = 0.01$

(Supplementary Figs. S8-S11), but underestimated by the genotype data and overestimated by

the consensus data for $\tau_h = 0.1$ (Supplementary Figs. S12-S15). Across all simulations, there

was little effect of increasing ILS aside from increasing HPD intervals, but the uncertainty was

negligible when 400 loci were sampled.

Analyses of target enrichment data from Dryopteris

Recovery of Phased Loci — On average, 62% of loci sequenced for an individual were

phased with eight variants passing filters per locus per individual (Supplementary Table S2).

The ploidy level appears to be strongly associated with the number of phased loci. Among

diploids, only 32% of loci were phased; the other 68% of diploid loci were either homozygous or

had too few linked variants for phasing. For tetraploids and hexaploids, 87% and 94% of loci,

respectively, were phased such that two or more phased haplotype sequences could be

recovered. Among loci where phasing was possible, variants were almost always resolved as a

single haplotype block, as opposed to being split into two or more blocks because not enough

reads were available to physically link variants. For polyploids, the number of unique phased

haplotype sequences most frequently matched the ploidy level, except in the case of a single *D.*

campyloptera individual (B087-D08), which also had relatively few recovered loci. Phasing data

only extended sequence alignment length by about two base pairs relative to the haplotype

consensus sequences (Supplementary Table S3). However, phased data more than tripled the

number of parsimony informative sites per alignment, similar to the increase in number of tips

(Supplementary Table S3).

Placing a single reticulation event — For our three-species full-likelihood analyses with

the MSci model, all data types decisively supported the putative reticulation hypothesis in the

cases of the allotetraploids *D. celsa* or *D. campyloptera* (Table 1). However, only the genotyped

data, coupled with the analytical phasing technique of BPP, chose the putatively correct model

for the allohexaploid *D. clintoniana* (Table 1). The phased and pick one data types selected the

network with the diploid *D. goldiana* as the hybrid descendent of the allohexaploid *D. clintoniana*

and allotetraploid *D. cristata*, while the best model for consensus data was a binary tree with *D.*

clintoniana sister to *D. cristata* (Table 1). However, model probability confidence intervals

suggest a great deal of uncertainty underlies model selection results for the allohexaploid case,

with the upper end of the interval for the putatively correct model at 0.603, 0.582, and 0.996 for

the phased, consensus, and pick one data, respectively (Table 1). Model parameters typically

converged across all analyses with the exception of the phased data for the *D. clintoniana* triplet

(Supplementary Fig. S16), which was due to one MCMC run despite convergence of the lnL

values.

The simulation experiments established some expectations that the phased and pick

one data would provide unbiased estimates of divergence times for nodes involved in the

hybridization events, while the genotyped and consensus data would provide under- and

Table 1 — Model Probabilities for Possible Topological Hypotheses Using the MSci model.

Hybrid	Model	Phased Model Probability	Genotype Model Probability	Consensus Model Probability	Pick One Model Probability
D. celsa	1	8.02×10^{-142} (3.62×10^{-146}, 3.31×10^{-137})	1.95×10^{-173} (1.03×10^{-177}, 2.78×10^{-169})	5.79×10^{-168} (3.15×10^{-172}, 8.7×10^{-164})	4.32×10^{-105} (1.99×10^{-109}, 2.27×10^{-101})
	2	3.18×10^{-126} (4.89×10^{-131}, 1.34×10^{-121})	5.44×10^{-136} (2.17×10^{-140}, 7.99×10^{-132})	1.66×10^{-113} (9.18×10^{-118}, 2.83×10^{-109})	2.78×10^{-84} (1.24×10^{-88}, 2.14×10^{-80})
	3	3.33×10^{-124} (5.19×10^{-129}, 1.66×10^{-119})	5.77×10^{-149} (1.63×10^{-153}, 1.31×10^{-144})	5.66×10^{-162} (6.09×10^{-166}, 9.31×10^{-158})	1.44×10^{-109} (6.87×10^{-114}, 2.06×10^{-105})
	4	1.36×10^{-146} (5.71×10^{-151}, 5.14×10^{-142})	6.98×10^{-149} (2.24×10^{-153}, 1.62×10^{-144})	2.89×10^{-152} (1.19×10^{-156}, 4.33×10^{-148})	1.44×10^{-109} (8.54×10^{-114}, 8.43×10^{-106})
	5	**1 (1, 1)**	**1 (1, 1)**	**1 (1, 1)**	**1 (1, 1)**
	6	3.69×10^{-113} (1.31×10^{-117}, 1.45×10^{-108})	1.59×10^{-143} (7.07×10^{-148}, 3.22×10^{-139})	1.97×10^{-118} (1.83×10^{-122}, 4.43×10^{-114})	1.46×10^{-92} (1.12×10^{-96}, 8.06×10^{-89})
D. campyloptera	1	1.42×10^{-251} (1.6×10^{-256}, 5.32×10^{-247})	3.77×10^{-251} (2.11×10^{-255}, 6.16×10^{-247})	3.15×10^{-154} (3.78×10^{-158}, 5.25×10^{-150})	1.52×10^{-130} (5.8×10^{-135}, 3.11×10^{-126})
	2	1.03×10^{-210} (1.12×10^{-215}, 8.01×10^{-206})	2.54×10^{-226} (1.33×10^{-230}, 2.41×10^{-222})	1.69×10^{-110} (9.98×10^{-115}, 1.72×10^{-106})	1.11×10^{-90} (5.02×10^{-95}, 2.23×10^{-86})
	3	2.16×10^{-242} (2.64×10^{-247}, 6.06×10^{-238})	1.53×10^{-184} (7.18×10^{-189}, 5.03×10^{-180})	1.24×10^{-129} (6.49×10^{-134}, 2.39×10^{-125})	5.72×10^{-123} (5.18×10^{-127}, 1.1×10^{-118})
	4	1.22×10^{-133} (7.33×10^{-139}, 7.63×10^{-129})	1.87×10^{-193} (6.47×10^{-198}, 2.12×10^{-189})	2.8×10^{-134} (1.98×10^{-138}, 7.92×10^{-130})	4.19×10^{-114} (2.78×10^{-118}, 6.67×10^{-110})
	5	**1 (1, 1)**	**1 (1, 1)**	**1 (1, 1)**	**1 (1, 1)**
	6	4.96×10^{-173} (6.57×10^{-178}, 3.7×10^{-168})	1.88×10^{-189} (9.15×10^{-194}, 1.93×10^{-185})	5.11×10^{-103} (2.49×10^{-107}, 4.81×10^{-99})	5.44×10^{-86} (3.65×10^{-90}, 1.32×10^{-81})
D. clintoniana	1	3.44×10^{-110} (5.27×10^{-115}, 1.25×10^{-105})	6.01×10^{-65} (2.21×10^{-69}, 1.99×10^{-60})	2.62×10^{-12} (8.24×10^{-17}, 1.82×10^{-8})	1.52×10^{-13} (3.47×10^{-18}, 4.54×10^{-10})
	2	1.44×10^{-04} (1.46×10^{-9}, 0.826)	1.91×10^{-47} (4.69×10^{-52}, 3×10^{-43})	4.49×10^{-9} (1.83×10^{-13}, 4.18×10^{-5})	5.84×10^{-6} (9.84×10^{-11}, 7.03×10^{-2})
	3	4.6×10^{-134} (6.97×10^{-139}, 5.65×10^{-130})	2.31×10^{-55} (5.51×10^{-60}, 5.35×10^{-51})	**1 (4.84×10^{-2}, 1)**	2.23×10^{-12} (5.82×10^{-17}, 1.38×10^{-8})
	4	1.23×10^{-115} (3.09×10^{-120}, 2.69×10^{-111})	2.18×10^{-27} (3.04×10^{-32}, 3.68×10^{-23})	*1.88×10^{-4} (1.05×10^{-8}, 0.728)*	2.02×10^{-27} (3.5×10^{-32}, 1.34×10^{-23})
	5	*4.23×10^{-5} (7.74×10^{-10}, 0.603)*	**1 (1, 1)**	*1.73×10^{-4} (6.85×10^{-09}, 0.582)*	*5.7×10^{-3} (1.31×10^{-7}, 0.996)*
	6	**1 (3.81×10^{-2}, 1)**	2.48×10^{-31} (6.11×10^{-36}, 5.76×10^{-27})	1.04×10^{-15} (6.71×10^{-20}, 6.87×10^{-12})	**0.994 (3.6×10^{-3}, 1)**

Note: Topologies of the six models that enumerate the possible binary and reticulate evolutionary histories are displayed in Figure S2. The best model for each data type and triplet is shown in bold. A percentile-based 95% confidence interval is given for each probability by bootstrapping the posteriors.

overestimates, respectively. However, phased data generally produced younger mean τ

estimates for the *D. celsa* triplet with reduced credible intervals relative to other data types, but

much older mean τ estimates with more uncertainty for the *D. campyloptera* and *D. clintoniana*

triplets (Fig. 5; Supplementary Table S4). Similarly, there was inconsistency in θ estimates

observed among data types across triplets, with notably larger credible intervals for phased data

in the *D. celsa* and *D. clintoniana* triplets. Estimates of φ were sensitive to data type across

triplets, such that the mean φ suggested the major contribution to the focal allopolyploid from

phased data was *D. goldiana* into *D. celsa*, *D. intermedia* into *D. campyloptera*, and *D. goldiana*

into *D. clintoniana*, with other data types sometimes disagreeing (Fig. 5). Credible intervals

around φ always spanned $\varphi = 0.5$, but estimates for genotype, consensus, and phased data

suggested some biased gene retention for the *D. celsa* and *D. clintoniana* triplets

(Supplementary Table S4).

503   When performing a network search based on gene tree distributions with SNaQ, all data

types recovered the putatively correct relationship among *D. celsa* and its diploid parental

lineages (Fig. 6a). For *D. campyloptera*, all data types correctly detected one hybrid edge

(Supplementary Table S5), but they got the direction of gene flow wrong (Fig. 6b). All data types

also detected one hybrid edge for the allohexaploid *D. clintoniana* (Supplementary Table S5),

but there was disagreement over the underlying network. Both the phased and consensus data

invoked a ghost lineage sister to the three sampled species*,* such that gene flow was from the

ghost into the allotetraploid *D. cristata*. While not unreasonable given the putatively extinct *D.*

semiscristata and missing *D. ludoviciana* from the sample, the relationship between *D.*

clintoniana and its parental lineages was undetected (Fig. 6c). The genotype and pick one data

recovered the correct network, but reversed the direction of gene flow from allohexaploid *D.*

clintoniana into allotetraploid *D. cristata* (Fig 6c).

Figure 5 — *Dryopteris* **Divergence Times, Population Sizes, and Introgression Probabilities under the MSci Model.** Points are posterior means and error bars show the 95% HPD intervals. Divergence times are measured as the expected number of substitutions per site.

Figure 6 — Network search results for three-taxon *Dryopteris* cases with SNaQ. The major topology edge is indicated in blue and the minor edge (direction of introgression) is shown in gold. The introgression probability (φ) is shown for each data type. In cases where a) *D. celsa* or b) *D. campyloptera* were the allotetraploids under investigation, all data types recovered the same species network. When c) *D. clintoniana* was the allohexaploid lineage from the allotetraploid *D. cristata* and diploid *D. goldiana* progenitors, different networks were recovered by the phased or consensus versus genotype or pick one data types.

525    *Inferring relationships among a complex with multiple reticulation events* — The nine

526 *Dryopteris* species SNaQ network analysis with phased data identified two out of five

527 hypothesized reticulation events correctly along with one incorrect reticulation edge (Fig. 7a;

528 Supplementary Table S6). The analysis correctly identified the diploids *D. ludoviciana* and *D.*

529 *goldiana* as parents to the allotetraploid *D. celsa*, as well as the diploids *D. expansa* and *D.*

530 *intermedia* as parents to the allotetraploid *D. campyloptera*. A hybrid edge from the common

531 ancestor of *D. celsa*, *D. ludoviciana*, and *D. goldiana* to the common ancestor of *Polystichum*

532 was also recovered with an introgression probability of 0.335, although with low bootstrap

533 support. The parents of *D. celsa* were also detected by the other data types, but all had some

534 degree of error. The genotype data found a reticulation edge between the allotetraploid *D.*

535 *carthusiana* and diploid parent *D. intermedia*, but reversed the direction of gene flow (Fig. 7b). A

536 reticulation edge was also detected between *D. cristata* and the common ancestor of *D.*

537 *clintoniana*, *D. goldiana*, *D. celsa*, and *D. ludoviciana*, which is not completely unreasonable

538 given the allotetraploid *D. cristata* is believed to be a parent of *D. clintoniana*. The consensus

539 data only detected two reticulation edges (Supplementary Table S6), with the second edge

540 correctly showing gene flow from the diploid parent *D. intermedia* into the allotetraploid *D.*

541 *carthusiana*, provided the other parent *D. semicristata* is presumed extinct obscuring the

542 network interpretation (Fig. 7c). The pick one data, similar to the genotype data, inferred a

543 reticulation edge between *D. carthusiana* and *D. intermedia* with the wrong direction of gene

544 flow (Fig. 7d). The pick one data also detected a reticulation edge from *D. cristata* into the

545 common ancestor of *D. carthusiana D. intermedia*, *D. campyloptera*, and *D. expansa*, which

546 lacks a clear interpretation other than invoking missing lineages underlying complexity of the

547 backbone topology (Fig. 1). In the one case where all data types detected the correct

548 relationship regarding *D. celsa*, the introgression probability was approximately 0.5 with phased

549 data while between 0.37 and 0.39 with others (Fig. 7; Supplementary Fig. S17).

Figure 7 — Networks for all North American *Dryopteris* inferred from SNaQ. Colors indicate major and minor (reticulation) edges, and which minor edges were hypothesized or an unexpected result. Introgression probabilities for each minor edge are provided. Numbers on branches subtending nodes are bootstrap support values. Results are shown separately for a) phased data where all haplotype sequences are included in gene trees, b) genotyped data with a single sequence for individual with IUPAC ambiguity codes, c) haplotype consensus sequences, and d) a single haplotype sequence representing each individual among gene trees.

Differences in the network estimates among data types may be explained by their

underlying spectra of quartet concordance factors (Supplementary Fig. S18). While spectra

were overall similar among data types, some deviations from expectations under ILS alone were

stronger in the phased data versus others (Supplementary Fig. S18). Analyses of gene tree

incongruence among the consensus (Supplementary Fig. S19), genotype (Supplementary Fig.

S20), and pick one (Supplementary Fig. S21) data showed similar levels of conflict among the

different data types with respect to a species tree estimated from the phased data (such

analyses with multiple alleles per species is not possible). The frequency of conflicting splits

with high bootstrap support was similar across datatypes for all nodes (Supplementary Fig.

S22), and low-support conflicts represented on average 56% of splits within *Dryopteris*.

DISCUSSION

New phylogenetic network methods offer the promise of elucidating the often complex

reticulate histories of polyploid lineages, even in the presence of ILS. Our results demonstrate

that under some conditions phasing polyploid target enrichment data can improve the accuracy

of such network inferences as well as divergence time estimates for networks, and we describe

a novel pipeline (PATÉ) to address the difficult problem of phasing polyploid data. Although

PATÉ could handle different types of genomic data, such as transcriptomes and whole

genomes, target enrichment data are ideal for computational phasing because they can yield

high (i.e. greater than 100x) and even coverage within a locus. This was demonstrated in our

Dryopteris analyses when almost all loci were recovered as single haplotype blocks

(Supplementary Table S2). Because MSC methods treat loci independently and assume free

recombination between loci, it is not necessary to assign individual loci to parental subgenomes

for network analyses. However, the allele sequences output by PATÉ can also be used as input

for other tools such as *Homologizer* (Freyman et al. 2023) or *HybPhaser* (Nauheimer et al.

2020), which attempt to phase across loci and assign parental subgenomes. Our methods also

enable population genomic studies of polyploids (Tiley et al. 2024) where accurate estimation of

site frequency spectra can be used for demographic analyses otherwise complicated by

polyploidy (e.g., Excoffier et al. 2013; Liu and Fu 2020) or SNP-based network inference in the

absence of variation suitable for gene tree estimation (Blischak et al. 2018a; Olave and Meyer

2020).

Promises and Limits of Phasing

The prospect of using alleles from phased genomic data presents an exciting step

towards revealing the evolutionary history of polyploids, which remains a critical impediment

within the plant evolution community (McKain et al. 2018). New strategies for explicitly

addressing this challenge are emerging (Freyman et al. 2023; Nauheimer et al. 2020; Mendez-

Reneau et al. 2023), and PATÉ can be a useful tool by phasing variants for many individuals

while leveraging genotype quality information. Our simulations demonstrated that phasing can

improve estimates of reticulate evolutionary relationships using network methods. In

simulations, phased data with all alleles represented in gene trees could more efficiently recover

the placement and directionality of hybrid edges than the genotyped, haplotype consensus, or

single haplotype sequences (Fig. 3). The advantages of using phased data for network

estimation diminished in simulations as the number of loci increased (Fig. 3), and this was

evident from our empirical analyses of *Dryopteris* triplets, which produced similar results across

data types with 330 or 328 loci (Table 1; Fig. 6), except for the correct inference of allopolyploid

and parent relationships with genotyped data in the case of *D. clintoniana* (Table 1).

Both our simulations (Fig. 4; Supplementary Figs. S4-S15) and previous investigations

(Anderman et al. 2019) suggested phased data can be important for unbiased divergence time

estimation, especially estimating the origin of a hybrid lineage (Fig. 4). Our empirical analyses

also demonstrated how the timing of introgression τ_h and other parameters can be greatly

affected by the data type used (Fig. 5). In our *Dryopteris* analyses, τ_h for *D. celsa* was at least in

the same order-of-magnitude among data types, which had partially overlapping credible

intervals, while phased τ_h estimate *for D. campyloptera* was nearly four times older than the

other data types (Fig. 5; Supplementary Table S4). This was surprising given that simulations

suggested haplotype consensus data should generate older estimates than phased data (Fig.

4). The *D. clintoniana* τ_h estimate from phased data was also much older than other data types,

albeit with a wide 95% HPD interval. There were noticeable mixing problems in our *D.*

clintoniana full-likelihood analyses, which makes it difficult to evaluate whether the uncertainty

was due to extreme differences in θ among phased alleles within *D. clintoniana* and/or the

computational difficulty of the Bayesian methods with increasing sequence data. In all cases,

the differences between τ_h among data types is not consistent with results from the simulation

experiments, which suggested the consensus data should overestimate age compared to

phased or pick one data, with genotyped data having slightly younger estimates than phased

data. This highlights the difficulties of simulating data that capture the complexities of empirical

data and makes deciding which estimate is more reliable difficult. For example, the uncertainty

of τ_h with phased data for *D. celsa* and *D. campyloptera* triplets was reduced compared to the

consensus and pick one estimates, but greater uncertainty was observed for τ and θ at other

nodes (Fig. 5). A potential compromise from simulations and empirical analyses is the use of

analytical phasing of polyploid data, as done with our genotyped data for BPP analyses.

Analytical phasing (Gronau et al. 2011; Flouri et al. 2020) has been shown to perform well in

diploids (Huang et al. 2022), and any potential biases from ignoring ploidy seem outweighed by

getting the *D. clintoniana* model correct (Table 1) as well as seemingly reasonable parameter

estimates under the MSci model with tight credible intervals (Fig. 5).

Our analyses also suggest that phasing may be problematic when the parental lineages

are deeply diverged. Although the phased data were able to accurately estimate the age of

older speciation nodes in simulations (Supplementary Figs. S4-S15), as phylogenetic

information is lost from multiple hits, the influence of the prior in full likelihood analyses should

become stronger. Deeper divergences will likely generate more switch errors among phased

haplotype sequences generated by PATÉ (Xie et al. 2016), as there will be more opportunities

for paralogy, structural variants, and sequence divergence to confound the read mapping,

genotyping, and phasing processes. When enough reads are available to call high-confidence

variants, we suggest that phasing with PATÉ can improve network estimation based on the

correct identification of the allotetraploids *D, celsa* and *D. campyloptera* and their parental

lineages in the analyses of the North American *Dryopteris* complex (Fig. 7a); however, phasing

alone cannot overcome some inherent challenges of network estimation.

Challenges of Network Estimation

Our analyses highlight the difficulty of estimating the evolutionary histories of reticulate

complexes, regardless of data type. The full-likelihood implementation of the MSci model

appears to be useful when the number of tips and plausible models for investigation are low, but

these methods may not be practical for generating hypotheses and exploring unknown

relationships for large numbers of taxa (e.g., Zhang et al. 2018). Quartet-based methods are

much faster and accurate when model assumptions are met (Solis-Lemus and Ané 2016) and

show similar accuracy to full-likelihood methods for estimating introgression probabilities (Flouri

et al. 2020). However, there are scenarios where the true networks are non-identifiable (Yu and

Nakhleh 2015; Solís-Lemus and Ané 2016). For example, when multiple introgression events

affect the same lineage, such that there are overlapping cycles in a network (Solís-Lemus and

Ané 2016), the expected quartet distribution under the MSci model becomes a poor fit for the

empirical data (Cai and Ané 2020), and the network estimated may be incorrect. These effects

were evident in our empirical analyses where the relationship between *D. campyloptera*, *D.*

expansa, and *D. intermedia* was missing in the nine-species analysis (Fig. 7), which we expect

is due to a reticulation edge present between *D. intermedia* and *D. carthusiana*. Phasing

sequence data adds information that can improve estimates (Supplementary Table S3), but

unsampled or extinct lineages, such as the hypothesized *D. semicristata,* can create significant

barriers to recovering the true evolutionary history of reticulate complexes, regardless of how

many loci or individuals are available. Another identifiability problem is getting the direction of

gene flow correct (Yu and Nakhleh 2015; Thawornwattana et al. 2023), and these scenarios

were evident in our quartet-based analyses of a triplet that reversed gene flow from the

allotetraploid *D. campyloptera* into the diploid *D. expansa* (Fig. 6d) or in the analyses of the full

complex that reversed gene flow from the allotetraploid *D. carthusiana* into the diploid *D.*

intermedia (Fig. 7b and 7d).

Insights into Dryopteris Evolution

The North American *Dryopteris* complex has been well-characterized through the study

of multi-locus nuclear and chloroplast phylogenies, morphology, cytological observations,

chromatography, and isozyme analyses (reviewed in Sessa et al. 2012b). This makes it a useful

system for testing our phasing pipeline, and our analyses add nuance to our understanding of

some of the relationships among *Dryopteris* species. For example, our results with phased data

indicated *D. campyloptera* has received more loci from *D. intermedia* than *D. expansa* (Fig. 7a),

and *D. intermedia* is likely the maternal progenitor of *D. campyloptera* (Sessa et al. 2012b).

Following the allopolyploidy event, the *D. intermedia* subgenome may have been dominant,

providing a selective advantage for *D. campyloptera* in its distribution at the time (Bird et al.

2018). The phased nine-taxon analyses also supported the hypothesis of the unsampled diploid

lineage *D. semicristata*, based on the placement of *D. carthusiana* as sister to the rest of

Dryopteris, and the general sensitivity of *D. carthusiana* and *D. cristata* in the phylogenetic

backbone across data types (Fig. 7). Additional insights may be gained by a more complete

sampling of the genus, as the North American *Dryopteris* complex is not a clade (Sessa et al.

2012b), which should alleviate some of the identifiability problems. Analyses of the North

American *Dryopteris* complex alone highlight the need for caution when interpreting networks,

which are increasingly prominent plant evolutionary research.

Does Phasing Matter?

Combining phased data with recent network methods can help confront a major

challenge of plant phylogenetics: resolving the complex histories of polyploids. While our

analyses showed haplotype consensus sequences can be adequate for resolving single

reticulation events where both parental lineages are sampled, phased sequences may provide

more information when only a few loci are available for cases where sequence divergence is not

too high, while more generally improving divergence time estimates (see also Huang et al.

2022). Analytical phasing approaches (Gronau et al. 2011; Flouri et al. 2020) provide a way

forward for full-likelihood methods while avoiding the potential for switch errors from read-based

phasing as done in PATÉ, but these will become difficult with many tips. This highlights the

need for continued development of models that explicitly consider polyploidization mechanisms

(Jones et al. 2013) while addressing scalability and limiting assumptions for contemporary

genomic data (Yan et al. 2022). Some reticulate complexes will be difficult to disentangle with

any data when confronted with statistical identifiability problems, such as networks with multiple

reticulations on a single edge (Solís-Lemus et al. 2016) when using quartet-based methods, and

our *Dryopteris* analyses demonstrate that caution is needed when interpreting biological

processes from phylogenetic networks in the absence of corroborating evidence. As haplotype-

phased genomes from long-read data become increasingly available, the discussion of phase

resolution will become trivial, but progress on models that leverage such data and improve

interpretation of processes, especially in polyploids that maintain distinct subgenomes, from

networks is still needed.

DATA AVAILABILITY

PATÉ is available through GitHub (https://github.com/gtiley/Phasing) and can be run on any

UNIX environment after installing basic genotyping software and H-PoPG. Simulated and

empirical data supporting findings and files for replicating some analyses are available from the

Dryad Digital Repository: https://doi.org/10.5061/dryad.5qfttdz53. Raw Fastq reads for

Dryopteris individuals are available through the NCBI SRA database and are associated with

BioProject PRJNA725004. Individual SRA Identifiers are available in Supplementary Table S1.

SUPPLEMENTARY MATERIAL

Data and supplementary material is available from the Dryad Digital Repository:

https://doi.org/10.5061/dryad.5qfttdz53

ACKNOWLEDGEMENTS

The authors thank A.M. Duffy, K. Imwattana, M. Nieto-Lugilde, B.T. Piatkowski, and A.J.

Shaw for helpful discussions and providing feedback on the manuscript. We also thank M.G

Johnson, J. Mendez Reneau, L. Nauheimer, and C.J. Rothfels for discussions and sharing their

strategies for phasing sequence data. We are grateful to the people who made data collection

possible; we used *Dryopteris* samples collected by C.J. Rothfels, M.A. Sundue, and W. Testo,

and DNA extractions were performed by S.B. Carey and E. Lockwood. The manuscript was

improved through helpful comments from the associate editor S.D. Smith and three anonymous

reviewers.

FUNDING

Funding was provided by National Science Foundation awards DEB-2038213 to AAC

and PSM, and from DEB-1541506 to JGB and EBS. This work was also partially supported by

the Department of Energy award DE-SC0021016 to CSL. ADY and GPT gratefully acknowledge

support from Duke University. This project has received funding from the European Union's

Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie grant

agreement No. 101026923, awarded to GPT.

LITERATURE CITED

742 **LITERATURE CITED**

743 Albert T.J., Molla M.N., Muzny D.M., Nazareth L., Wheeler D., Song X., Richmond T.A., Middle
744 C.M., Rodesch M.J., Packard C.J., Weinstock G.M., Gibbs R.A. 2007. Direct selection of human
745 genomic loci by microarray hybridization, *Nature Methods*. 4:903-905.
746
747 Andermann T., Cano A., Zizka A., Bacon C., Antonelli A. 2018. SECAPR-a bioinformatics
748 pipeline for the rapid and user-friendly processing of targeted enriched Illumina sequences, from
749 raw reads to alignments. *PeerJ* 6:e5175.
750
751 Andermann T., Fernandes A.M., Olsson U., Topel M., Pfeil B., Oxelman B., Aleixo A., Faircloth
752 B.C., Antonelli A. 2019. Allele Phasing Greatly Improves the Phylogenetic Utility of
753 Ultraconserved Elements. *Syst. Biol.* 68:32-46.
754
755 Baaijens J.A., Schonhuth A. 2019. Overlap graph-based generation of haplotigs for diploids and
756 polyploids. *Bioinformatics* 35:4281-4289.
757
758 Bankevich A., Nurk S., Antipov D., Gurevich A.A., Dvorkin M., Kulikov A.S., Lesin V.M.,
759 Nikolenko S.I., Pham S., Prjibelski A.D., Pyshkin A.V., Sirotkin A.V., Vyahhi N., Tesler G.,
760 Alekseyev M.A., Pevzner P.A. 2012. SPAdes: a new genome assembly algorithm and its
761 applications to single-cell sequencing. *J. Comput. Biol.* 19:455-477.
762
763 Barker M.S., Arrigo N., Baniaga A.E., Li Z., Levin D.A. 2016. On the relative abundance of
764 autopolyploids and allopolyploids. *New Phytol.* 210:391-398.
765
766 Baudry J.-P., Maugis C., Michel B. 2012. Slope heuristics: overview and implementation.
767 *Statistics and Computing*. 22:455-470.
768
769 Berger E., Yorukoglu D., Peng J., Berger B. 2014. HapTree: a novel Bayesian framework for
770 single individual polyplotyping using NGS data. *PLoS Comput. Biol.* 10:e1003502.
771
772 Bezanson J., Edelman A., Karpinski S., Shah V.B. 2015. Julia: A Fresh Approach to Numerical
773 Computing. *arXiv*1411.1607v4.
774
775 Bird K.A., VanBuren R., Puzey J.R., Edger P.P. 2018. The causes and consequences of
776 subgenome dominance in hybrids and recent polyploids. *New Phytol.* 220:87-93.
777
778 Blischak P.D., Chifman J., Wolfe A.D., Kubatko, L.S. 2018a. HyDe: a Python package for
779 genome-scale hybridization detection. *Syst. Biol.*, 67:821-829.
780
781 Blischak PD, Kubatko LS, Wolfe AD. 2018b. SNP genotyping and parameter estimation in
782 polyploids using low-coverage sequencing data. *Bioinformatics* 34:407-415.
783
784 Breinholt J.W., Carey S.B., Tiley G.P., Davis E.C., Endara L., McDaniel S.F., Neves L.G., Sessa
785 E.B., von Konrat M., Chantanaorrapint S., Fawcett S., Ickert-Bond S.M., Labiak P.H., Larraín J.,
786 Lehnert M., Lewis L.R., Nagalingum N.S., Patel N., Rensing S.A., Testo W., Vasco A., Villarreal
787 J.C., Williams E.W., Burleigh J.G. 2021. A target enrichment probe set for resolving the
788 flagellate plant tree of life. *Appl. Plant Sci.* 9:e11406.
789

790 Breinholt J.W., Earl C., Lemmon A.R., Lemmon E.M., Xiao L., Kawahara A.Y. 2018. Resolving
791 relationships among the megadiverse butterflies and moths with a novel pipeline for anchored
792 phylogenomics. *Syst. Biol.* 67:78-93.
793
794 Buggs J.A., Wendel J.F., Doyle J.J., Soltis D.E., Soltis P.S., Coate J.E. 2014. The legacy of
795 diploid progenitors in allopolyploid gene expression patterns. *Philos. Trans. R. Soc. Lond. B*
796 *Biol. Sci.* 368:20130354.
797
798 Cai R., Ané C. 2020. Assessing the fit of the multi-species network coalescent to multi-locus
799 data. *Bioinformatics* btaa863. doi: https://doi.org/10.1093/bioinformatics/btaa863
800
801 Colle M., Leisner C.P., Wai C.M., Ou S., Bird K.A., Wang J., Wisecaver J.H., Yocca A.E., Alger
802 E.I., Tang H., Xiong Z., Callow P., Ben-Zvi G., Brodt A., Baruch K., Swale T., Shiue L., Song G.-
803 Q., Childs K.L., Schilmiller A., Vorsa N., Buell C.R., VanBuren R., Jiang N., Edger P.P. 2019.
804 Haplotype-phased genome and evolution of phytonutrient pathways of tetraploid blueberry.
805 *GigaScience* 8:giz012.
806
807 Crowl A.A., Myers C., Cellinese N. 2017. Embracing discordance: Phylogenomic analyses
808 provide evidence for allopolyploidy leading to cryptic diversity in a Mediterranean Campanula
809 (Campanulaceae) clade. *Evolution* 71:913-922.
810
811 Crowl A.A., Manos P.S., McVay J.D., Lemmon A.R., Lemmon E.M., Hipp A.L. 2020. Uncovering
812 the genomic signature of ancient introgression between white oak lineages (*Quercus*). *New*
813 *Phytologist* 226: 1158–1170.
814
815 DePristo M., Banks E., Poplin R., Garimella K., Maguire J., Hartl C., Philippakis A., del Angel G.,
816 Rivas M.A., Hanna M., McKenna A., Fennell T., Kernytsky A., Sivachenko A., Cibulskis K.,
817 Gabriel S., Altschuler S., Daly M. 2011. A framework for variation discovery and genotyping
818 using next-generation DNA sequencing data. *Nat. Genet.* 43:491-498.
819
820 Edgar R.C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
821 throughput. *Nucleic Acids Res.* 32:1792-1797.
822
823 Eriksson J.S., de Sousa F., Bertrand Y.J.K., Antonelli A., Oxelman B., Pfeil B.E. 2018. Allele
824 phasing is critical to revealing a shared allopolyploid origin of Medicago arborea and M.
825 strasseri (Fabaceae). *BMC Evol. Biol.* 18:9.
826
827 Excoffier L., Dupanloup I., Huerta-Sanchez E., Sousa V.C., Foll M. 2013. Robust demographic
828 inference from genomic and SNP data. *PLoS Genet.* 9:e1003905.
829
830 Faircloth B.C. 2016. PHYLUCE is a software package for the analysis of conserved genomic
831 loci. *Bioinformatics* 32:786-788.
832
833 Faircloth B.C., McCormack J.E., Crawford N.G., Harvey M.G., Brumfield R.T., Glenn T.C. 2012.
834 Ultraconserved elements anchor thousands of genetic markers spanning multiple evolutionary
835 timescales. *Syst. Biol.* 61:717-726.
836
837 Felsenstein J. 1981. Evolutionary trees from DNA sequences: A maximum likelihood approach.
838 *Journal of Molecular Evolution*. 17:368-376.
839

840   Flouri T., Jiao X., Rannala B., Yang Z. 2020. A Bayesian Implementation of the Multispecies
841   Coalescent Model with Introgression for Phylogenomic Analysis. *Mol. Biol. Evol.* 37:1211-1223.
842

843   Flouri T., Jiao X., Rannala B., Yang Z.. 2018. Species Tree Inference with BPP Using Genomic
844   Sequences and the Multispecies Coalescent. *Mol. Biol. Evol.* 35:2585-2593.
845

846   Freyman W.A,. Johnson M.G., Rothfels C.J. 2023. Homologizer: Phylogenetic phasing of gene
847   copies into polyploid subgenomes. *Methods in Ecology and Evolution*. 14:1230-1244.
848

849   Gnirke A., Melnikov A., Maguire J., Rogov P., LeProust E.M., Brockman W., Fennell T.,
850   Giannoukos G., Fisher S., Russ C., Gabriel S., Jaffe D.B., Lander E.S., Nusbaum C. 2009.
851   Solution hybrid selection with ultra-long oligonucleotides for massively parallel targeted
852   sequencing. *Nature Biotechnology*. 27:182-189.
853

854   Gronau I., Hubisz M.J., Gulko B., Danko C.G., Siepel A. 2011. Bayesian inference of ancient
855   human demography from individual genome sequences. *Nat. Genet.* 43:1031-1034.
856

857   He D., Saha S., Finkers R., Parida L. 2018. Efficient algorithms for polyploid haplotype phasing.
858   *BMC Genomics* 19:110.
859

860   Huang J., Bennett J., Flouri T., Leaché A.D., Yang Z. 2022. Phase resolution of heterozygous
861   sites in diploid genomes is important to phylogenomic analysis under the multispecies
862   coalescent model. *Syst Biol.* 71:334-352.
863

864   Huang J., Flouri T., Yang Z. 2020. A simulation study to examine the information content in
865   phylogenomic datasets under the multispecies coalescent model. *Mol. Biol. Evol.* 37:3211-3224.
866

867   Huber K.T., Oxelman B., Lott M., Moulton V. 2006. Reconstructing the evolutionary history of
868   polyploids from multi-labelled trees. *Mol. Biol. Evol.* 23:1784-1791.
869

870   Hudson R.R. 1983. Testing the Constant-Rate Neutral Allele Model with Protein Sequence
871   Data. *Evolution* 37:203-217.
872

873   Huson D.H., Rupp R., Scornavacca C. 2010. Phylogenetic networks: concepts, algorithms and
874   applications. Cambridge University Press.
875

876   Iqbal Z., Caccamo M., Turner I., Flicek P., McVean G. 2012. De novo assembly and genotyping
877   of variants using colored de Bruijn graphs. *Nat. Genet.* 44:226-232.
878

879   Jantzen J.R., Amarasinghe P., Folk R.A., Reginato M., Michelangeli F.A., Soltis D.E., Cellinese
880   N., Soltis P.S. 2020. A two-tier bioinformatic pipeline to develop probes for target capture of
881   nuclear loci with applications in Melastomataceae. *Appl. Plant Sci.* 8:e11345.
882

883   Ji J., Jackson D.J., Leaché A.D., Yang Z. 2022. Power of Bayesian and heuristic tests to detect
884   cross-species introgression with reference to gene flow in the *Tamias quadrivittatus* group of
885   North American chipunks. *Syst Biol.* syac077.
886

887   Johnson M.G., Gardner E.M., Liu Y., Medina R., Goffinet B., Shaw A.J., Zerega N.J., Wickett
888   N.J. 2016. HybPiper: Extracting coding sequence and introns for phylogenetics from high-
889   throughput sequencing reads using target enrichment. *Appl. Plant Sci.* 4:1600016.
890

891 Johnson M.G., Pokorny L., Dodsworth S., Botigue L.R., Cowan R.S., Devault A., Eiserhardt
892 W.L., Epitawalage N., Forest F., Kim J.T., Leebens-Mack J.H., Leitch I.J., Maurin O., Soltis
893 D.E., Soltis P.E., Wong G.K.-S., Baker W.J., Wickett N.J. 2019. A Universal Probe Set for
894 Targeted Sequencing of 353 Nuclear Genes from Any Flowering Plant Designed Using k-
895 Medoids Clustering. *Syst. Biol.* 68:594-606.
896
897 Jones G., Sagitov S., Oxelman B. 2013. Statistical inference of allopolyploid species networks in
898 the presence of incomplete lineage sorting. *Syst. Biol.* 62:467-478.
899
900 Jukes T.H., Cantor C.R. 1969. Evolution of protein molecules. In: Munro H.N., editor.
901 Mammalian Protein Metabolism. New York, NY: Acedemic Press. p. 21-132.
902
903 Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., von Haeseler A., Jermiin L.S. 2017.
904 ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* 14:587-
905 589.
906
907 Kamneva O.K., Syring J., Liston A., Rosenberg N.A. 2017. Evaluating allopolyploid origins in
908 strawberries (*Fragaria*) using haplotypes generated from target capture sequencing. *BMC Evol*
909 *Biol*. 17:180.
910
911 Karimi N., Grover C.E., Gallagher J.P., Wendel J.F., Ané C., Baum D.A. 2020. Reticulate
912 Evolution Helps Explain Apparent Homoplasy in Floral Biology and Pollination in Baobabs
913 (Adansonia; Bombacoideae; Malvaceae). *Syst. Biol.* 69:462-478.
914
915 Kass R.E., Raftery A.E. 1995. Bayes factors. *Journal of the American Statistical Association*.
916 90:773-795.
917
918 Kates H.R., Johnson M.G., Gardner E.M., Zerega N.J.C., Wickett N.J. 2018. Allele phasing has
919 minimal impact on phylogenetic reconstruction from targeted nuclear gene sequences in a case
920 study of Artocarpus. *Am. J. Bot.* 105:404-416.
921
922 Lemmon A.R., Emme S.A., Lemmon E.M. 2012. Anchored hybrid enrichment for massively
923 high-throughput phylogenomics. *Syst. Biol.* 61:727-744.
924
925 Li H., Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.
926 *Bioinformatics* 25:1754-1760.
927
928 Liu X., Fu Y.X. 2020. Stairway Plot 2: demographic history inference with folded SNP frequency
929 spectra. *Genome Biol.* 21:280.
930
931 Liu Y., Johnson M.G., Cox C.J., Medina R., Devos N., Vanderpoorten A., Hedenas L., Bell N.E.,
932 Shevock J.R., Aguero B., Quandt D., Wickett N.J., Shaw A.J., Goffinet B. 2019. Resolution of
933 the ordinal phylogeny of mosses using targeted exons from organellar and nuclear genomes.
934 *Nat. Commun.* 10:1485.
935
936 Liu X.-Q., Xia X.-M., Chen L., Wang X-Q. 2022. Phylogeny and evolution of Cupressaceae:
937 Updates on intergeneric realtionships and new insights on ancient intergeneric hybridization.
938 *Mol. Phylogenet. Evol.* 177:107606.
939
940 Lott M., Spillner A., Huber K.T., Moulton V. 2009. PADRE: a package for analyzing and
941 displaying reticulate evolution. *Bioinformatics* 25:1199-1200.

942
943 Luo R., Liu B., Xie Y., Li Z., Huang W., Yuan J., He G., Chen Y., Pan Q., Liu Y., Tang J., Wu G.,
944 Zhang H., Shi Y., Liu Y., Yu C., Wang B., Lu Y., Han C., Cheung D.W., Yiu S.-M., Peng S.,
945 Xiaoqian Z., Liu G., Liao X., Li Y., Yang H., Wang J., Lam T-W., Wang J. 2012. SOAPdenovo2:
946 an empirically improved memory-efficient short-read de novo assembler. *GigaScience* 1:18.
947
948 McKain M.R., Johnson M.G., Uribe-Convers S., Eaton D., Yang Y. 2018. Practical
949 considerations for plant phylogenomics. *Appl. Plant Sci.* 6:e1038.
950
951 McKenna A., Hanna M., Banks E., Sivachenko A., Cibulskis K., Kernytsky A., Garimella K.,
952 Altshuler D., Gabriel S., Daly M., DePristo M.A. 2010. The Genome Analysis Toolkit: a
953 MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.*
954 20:1297-1303.
955
956 Mendez-Reneau J., Burleigh J.G., Sigel E.M. 2023. Target capture methods offer ingsight into
957 the evolution of rapidly diverged taxa and resolve allopolyploid homeologs in the fern genus
958 *Polypodium* s.s. *Systematic Botany* 48:96-109.
959
960 Meng C., Kubatko L.S. 2009. Detecting hybrid speciation in the presence of incomplete lineage
961 sorting using gene tree incongruence: a model. *Theor. Popul. Biol.* 75:35-45.
962
963 Moeinzadeh M.H., Yang J., Muzychenko E., Gallone G., Heller D., Reinert K., Haas S., Vingron
964 M. 2020. Ranbow: A fast and accurate method for polyploid haplotype reconstruction. *PLoS*
965 *Comput. Biol.* 16:e1007843.
966
967 Montgomery J.D., Paulton E.M. 1981. *Dryopteris* in North America. *Fiddlehead Forum* 8:25-31.
968
969 Morales-Briones D.F., Liston A., Tank D.C. 2018. Phylogenomic analyses reveal a deep history
970 of hybridization and polyploidy in the Neotropical genus Lachemilla (Rosaceae). *New Phytol.*
971 218:1668-1684.
972
973 Nauheimer L., Weigner N., Joyce E., Crayn D., Clarke C., Nargar K. 2020. HybPhaser: a
974 workflow for the detection and phasing of hybrids in target capture datasets. bioRxiv. doi:
975 https://doi.org/10.1101/2020.10.27.354589.
976
977 Nguyen L.-T., Schmidt H.A., von Haeseler A., Minh B.Q. 2015. IQ-TREE: a fast and effective
978 stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* 32:268-
979 274.
980
981 Oberprieler C., Wagner F., Tomasello S., Konowalik K. 2017. A permutation approach for
982 inferring species networks from gene trees in polyploid complexes by minimising deep
983 coalescences. *Methods in Ecology and Evolution* 8:835-849.
984
985 Olave M., Meyer A. 2020. Implementing Large Genomic Single Nucleotide Polymorphism Data
986 Sets in Phylogenetic Network Reconstructions: A Case Study of Particularly Rapid Radiations of
987 Cichlid Fish. *Syst. Biol.* 69:848-862.
988
989 Pamilo P., Nei M. 1988. Relationships between gene trees and species trees. *Mol. Biol. Evol.*
990 5:568-583.
991

992 Rothfels C.J., Pryer K.M., Li F.-W. 2017. Next-generation polyploid phylogenetics: rapid
993 resolution of hybrid polyploid complexes using PacBio single-molecule sequencing. *New Phytol.*
994 213:413-429.
995
996 Saada O.A., Tsouris A., Freidrich A., Schachrer J. 2020. nPhase: An accurate and contiguous
997 phasing method for polyploids. *bioRxiv* doi:https://doi.org/10.1101/2020.07.24.219105.
998
999 Schrinner S.D., Mari R.S., Ebler J., Rautianen M., Seillier L., Reimer J.J., Usadel B., Marschall
1000 T., Klau G.W. 2020. Haplotype threading: accurate polyploid phasing from long reads. *Genome*
1001 *Biology* 21:252.
1002
1003 Sessa E.B., Zimmer E.A., Givnish T.J. 2012a. Reticulate evolution on a global scale: a nuclear
1004 phylogeny for New World Dryopteris (Dryopteridaceae). *Mol. Phylogenet. Evol.* 64:563-581.
1005
1006 Sessa E.B., Zimmer E.A., Givnish T.J. 2012b. Unraveling reticulate evolution in North American
1007 Dryopteris (Dryopteridaceae). *BMC. Evol. Biol.* 12:104.
1008
1009 Solis-Lemus C., Ané C. 2016. Inferring Phylogenetic Networks with Maximum Pseudolikelihood
1010 under Incomplete Lineage Sorting. *PLoS Genet.* 12:e1005896.
1011
1012 Solis-Lemus C., Bastide P., Ané C. 2017. PhyloNetworks: A Package for Phylogenetic
1013 Networks. *Mol. Biol. Evol.* 34:3292-3298.
1014
1015 Soltis, D.E., Visger C.J., Soltis P.S. 2014. The polyploidy revolution then...and now: Stebbins
1016 revisited. *American Journal of Botany* 101: 1057–1078.
1017
1018 Stull G.W., Soltis P.S., Soltis D.E., Gitzendanner M.A., Smith S.A. 2020. Nuclear phylogenomic
1019 analyses of asterids conflict with plastome trees and support novel relationships among major
1020 lineages. *Am. J. Bot.* 107:790-805.
1021
1022 Sun M., Soltis D.E., Soltis P.S., Zhu X., Burleigh J.G., Chen Z. 2015. Deep phylogenetic
1023 incongruence in the angiosperm clade *Rosidae*. *Mol. Phylogenet. Evol.* 83:156-166.
1024
1025 Thawornwattana Y., Huang J., Flouri T., Mallet J., Yang Z. 2023. Inferring the direction of
1026 introgression using genomic sequence data. *Mol Biol Evol*. 40:msad178.
1027 https://doi.org/10.1093/molbev/msad178
1028
1029 Tiley G.P., Crowl A.A., Almary T.O.M., Luke W.R.Q., Solofondranohatra C.L., Besnard G.,
1030 Lehmann C.E.R., Yoder A.D., Vorontsova M.S. 2024. Genetic variation in *Loudetia simplex*
1031 supports the presence of ancient grasslands in Madagascar. *Plants People Planet* 6:315-329.
1032
1033 Wen D., Yu Y., Nakhleh L. 2016. Bayesian Inference of Reticulate Phylogenies under the
1034 Multispecies Network Coalescent. *PLoS Genet.* 12:e1006006.
1035
1036 Wen D., Yu Y., Zhu J., Nakhleh L. 2018. Inferring Phylogenetic Networks Using PhyloNet. *Syst.*
1037 *Biol.* 67:735-740.
1038
1039 Wolf P.G., Robison T.A., Johnson M.G., Sundue M.A., Testo W.L., Rothfels C.J. 2018. Target
1040 sequence capture of nuclear-encoded genes for phylogenetic analysis in ferns. *Appl. Plant Sci.*
1041 6:e01148.
1042

1043   Wood T.E., Takebayashi N., Barker M.S., Mayrose I., Greenspoon P.B., Rieseberg L.H. 2009.
1044   The frequency of polyplid speciation in vascular plants. *Proc. Natl. Acad. Sci. U. S. A.*
1045   106:13875-13879.
1046
1047   Xie W., Lewis P.O., Fan Y., Kuo L., Chen M.H. 2011. Improving marginal likelihood estimation
1048   for Bayesian phylogenetic model selection. *Syst. Biol.* 60:150-160.
1049
1050   Xie M., Wu Q., Wang J., Jiang T. 2016. H-PoP and H-PoPG: heuristic partitioning algorithms for
1051   single individual haplotyping of polyploids. *Bioinformatics* 32:3735-3744.
1052
1053   Yan Z., Cao Z., Liu Y., Ogilvie H.A., Nakhleh L. 2022. Phylogenetic networks in the presence of
1054   polyploid complexes. *Syst Biol.* 71:706-720.
1055
1056   Yang J., Moeinzadeh M-H., Kuhl H., Helmuth J., Xiao P., Haas S., Liu G., Zheng J., Sun Z., Fan
1057   W., Deng G., Wang H., Hu F., Zhao S., Fernie A.R., Boerno S., Timmermann B., Zhang P.,
1058   Vingron M. 2017. Haplotype-resolved sweet potato genome traces back its hexaploidization
1059   history. *Nat Plants* 3:696-703.
1060
1061   Yang Z. 2006. Computational molecular evolution: Oxford University Press.
1062
1063   Yu Y., Nakhleh L. 2015. A maximum pseudo-likelihood approach for phylogenetic networks.
1064   *BMC Genomics*. 16:S10.
1065
1066   Yu Y., Than C., Degnan J.H., Nakhleh L. 2011. Coalescent histories on phylogenetic networks
1067   and detection of hybridization despite incomplete lineage sorting. *Syst. Biol.* 60:138-149.
1068
1069   Zhang C., Ogilvie H.A., Drummond A.J., Stadler T. 2018. Bayesian Inference of Species
1070   Networks from Multilocus Sequence Data. *Mol. Biol. Evol.* 35:504-517.
1071
1072   Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree
1073   reconstruction from partially resolved gene trees. *BMC Bioinformatics* 19:153.
1074
1075   Zhu S., Degnan J.H. 2017. Displayed trees do not determine distinguishability under the
1076   network multispecies coalescent. *Syst. Biol.* 66:283-298.
1077

 FIGURE CAPTIONS

Figure 1 — Hypothesized Relationships among North American *Dryopteris*. Synthesis of

results from Sessa et al. 2012a and Sessa et al. 2012b. A) Links between shapes show the

putative parents and their allopolyploid derivatives. Black circles are diploids, squares are

tetraploids, and the hexagon is the one hexaploid species in the group. *Dryopteris semicristata*

is presumed extinct. B) Placement of allopolyploids in the context of the backbone relationships

among diploids. Tetraploids are indicated with solid orange lines and the hexaploid with dotted

lines. The grey line denoting a sister relationship between *D. semicristata* and *D. expansa*

reflects one possible placement for the extinct taxon based on previous analyses (Sessa et al.,

2012b).

1088

1089 **Figure 2 — Species network used for simulation.** The divergence times in expected

1090 substitutions per site are given for each node, and *h* is the hybrid node where two alleles enter

1091 from both *t* and *v*. E is an allotetraploid while other species are diploid. Absolute divergence

1092 times were changed by using values of θ at 0.001, 0.01, and 0.1. ILS was increased by

1093 multiplying θ by a factor of 2, 3, or 4 while keeping all τ_u fixed to their initial values.

1094

Figure 3 — Proportion of simulation replicates that correctly identify the allopolyploid

lineage using the SNaQ function of PhyloNetworks. The x-axes are the number of loci

sampled for each simulation. The y-axes are the proportions of correct networks. Results are

based on networks estimated with a single reticulation, even if that network was considered less

optimal than networks with zero or two reticulations. We saved the true gene trees from the

simulations, while we estimated gene trees with the phased, genotype, consensus, and pick one

data types.

1102

Figure 4 — Estimating the Timing of Hybridization with the MSci Model. Divergence times

are for node *h* in Figure 2. Divergence times are measured as the expected number of

substitutions per site. The dashed line represents the true simulated values. Points are posterior

means, and error bars are 95% HPD intervals, averaged across 100 replicates. Results are not

shown for the genotype data when $\tau_h = 0.1$ and $\theta = 4\tau_h$ because posterior means went towards

infinity and numerically unstable values that cannot be displayed.

1109

1110   **Figure 5 — *Dryopteris* Divergence Times, Population Sizes, and Introgression**

1111   **Probabilities under the MSci Model.** Points are posterior means and error bars show the 95%

1112   HPD intervals. Divergence times are measured as the expected number of substitutions per

1113   site.

1114

Figure 6 — Network search results for three-taxon *Dryopteris* cases with SNaQ. Colors

indicate the major and minor edge. The arrow from the minor edge shows the direction of

introgression. The introgression probability (φ) is shown for each data type. In cases where a)

D. celsa or b) *D. campyloptera* were the allotetraploids under investigation, all data types

recovered the same species network. When c) *D. clintoniana* was the allohexaploid lineage from

the allotetraploid *D. cristata* and diploid *D. goldiana* progenitors, different networks were

recovered by the phased or consensus versus genotype or pick one data types.

1122

Figure 7 — Networks for all North American *Dryopteris* inferred from SNaQ. Colors

indicate major and minor (reticulation) edges, and which minor edges were hypothesized or an

unexpected result. Introgression probabilities for each minor edge are provided. Numbers on

branches subtending nodes are bootstrap support values. Results are shown separately for a)

phased data where all haplotype sequences are included in gene trees, b) genotyped data with

a single sequence for individual with IUPAC ambiguity codes, c) haplotype consensus

sequences, and d) a single haplotype sequence representing each individual among gene trees.

1130