Running Head: Phased Target Enrichment for Polyploids

Phasing Alleles Improves Network Inference with Allopolyploids

George P. Tiley[1,†,*], Andrew A. Crowl[1,†], Paul S. Manos[1], Emily B. Sessa[2], Claudia Solís-

Lemus[3], Anne D. Yoder[1], J. Gordon Burleigh[2]

[1]Department of Biology, Duke University, Durham NC, 27708, USA

[2]Department of Biology, University of Florida, Gainesville FL, 32611, USA

[3]Wisconsin Institute for Discovery and Department of Plant Pathology, University of Wisconsin –

Madison, Madison WI, 53706, USA

[†]These authors contributed equally

[*]Author for correspondence: george.tiley@duke.edu

Abstract

Accurately reconstructing the reticulate histories of polyploids remains a central

challenge for understanding plant evolution. Although phylogenetic networks can provide

insights into relationships among polyploid lineages, inferring networks may be hampered by the

complexities of homology determination in polyploid taxa. We use simulations to show that

phasing alleles from allopolyploid individuals can improve inference of phylogenetic networks

under the multispecies coalescent. Phased allelic data can also improve divergence time

estimates for networks, which is helpful for evaluating allopolyploid speciation hypotheses and

proposing mechanisms of speciation. To achieve these outcomes, we present a novel pipeline

that leverages a recently developed phasing algorithm to reliably phase alleles from polyploids.

This pipeline is especially appropriate for target enrichment data, where depth of coverage is

typically high enough to phase entire loci. We provide an empirical example in the North

American *Dryopteris* fern complex that demonstrates how phasing can help reveal the mode of

polyploidization and improve network inference. We establish that our pipeline (PATÉ: Phased

Alleles from Target Enrichment data) is capable of recovering a high proportion of phased loci

from both diploids and polyploids, and that these data improve network estimates compared to

using haplotype consensus assemblies. This approach is shown to be especially effective in

reticulate complexes where there are multiple hybridization events. The pipeline is available at:

https://github.com/gtiley/Phasing.

Key words: Introgression; Hybridization; Reticulate Evolution; Multispecies Coalescent;

Divergence Time Estimation; Polyploidy; Target Enrichment; *Dryopteris*

INTRODUCTION

The phenomenon of polyploidy, or whole-genome duplication, occurs throughout the tree of life. Nowhere, perhaps, is its evolutionary significance more evident than in plants, with recent estimates suggesting up to 35% of vascular plant species are of recent polyploid origin (Wood et al. 2009; Barker et al. 2016). Despite advances in genomic data generation and a long-term interest in understanding the role of whole-genome duplication in driving plant speciation and local adaptation (reviewed in Soltis et al. 2014), polyploids remain a central challenge for the field of phylogenetics. One persistent problem when analyzing sequence data from polyploid taxa, and especially allopolyploids, is identifying the alleles and divergent homeolog copies from parental lineages. Most bioinformatic tools for processing next generation sequence data were developed with diploids, or specifically humans, in mind. These approaches often collapse variable homeolog sequences into a single consensus sequence for *de novo* assemblies or assume the organism is diploid when performing genotyping and phasing for reference-based assembly. For polyploids, this creates chimeric sequences that may interfere with phylogenetic reconstruction and obscure signals of polyploidy and polyploid mode-of-origin. Using allelic data that more accurately capture the complex genomic histories of polyploids should enable the incorporation of divergent signals from polyploid loci into phylogenomic inference, distinguish allopolyploidy from autopolyploidy, and identify parental taxa. However, few studies have examined the potential benefits of using phased versus unphased data to reconstruct polyploid histories, and there are few formal methods and little guidance for phasing alleles from polyploid taxa. Here we explore the value of using phased data to reconstruct polyploid networks leverage recent algorithmic advances in polyploid phasing (Xie et al. 2016) to develop a bioinformatic pipeline that can phase alleles from polyploids using target enrichment sequence data.

Previous studies of reticulate complexes have suggested phasing alleles is crucial for accurate evolutionary reconstruction, at least when sampling relatively few loci (e.g., 4 to 10;

Rothfels et al. 2017; Eriksson et al. 2018). The applications of phased sequencing data for

phylogenomic studies of polyploid complexes are thus enticing; however, it remains challenging

to genotype and phase next generation sequencing data from polyploids. Methods exist to

genotype consensus loci from target enrichment data, but these have been either limited to

diploids (Kates et al. 2018; Andermann et al. 2019) or manual curation of variants with

polyploids where both parental populations are available (Eriksson et al. 2018). Otherwise,

obtaining phased sequence data for polyploids has largely depended on costly long-read

sequencing to recover complete haplotype sequences (e.g., Rothfels et al. 2017), or cloning

PCR products (e.g., Sessa et al. 2012; Oberprieler et al. 2017).

Target enrichment (or HybSeq), where specific regions of the genome are isolated and

sequenced (Faircloth et al. 2012; Lemmon et al. 2012), is an increasingly common method for

collecting large-scale phylogenomic datasets, and these data can provide insights into the

evolutionary history of reticulate complexes (e.g., Crowl et al. 2017; Karimi et al. 2020) and

sources of gene tree discordance (e.g., Morales-Briones et al. 2018; Stull et al. 2020). Probe

kits for target enrichment have been developed in many land plant lineages (Wolf et al. 2018;

Johnson et al. 2019; Liu et al. 2019; Breinholt et al. 2021), and there are bioinformatic pipelines

available for custom probe design (e.g., Jantzen et al. 2020). The most common approach to

assemble phylogenetic datasets from target enrichment data has been to use *de novo* assembly

pipelines (Faircloth 2016; Johnson et al. 2016; Andermann et al. 2018; Breinholt et al. 2018).

The assembly algorithms within these pipelines typically treat variable base calls as sequencing

errors and consider only the most frequent nucleotide sequence while discarding the

alternatives (Bankevich et al. 2012; Iqbal et al. 2012; Luo et al. 2012). This results in loci in

which variable positions are collapsed to a single base call (haplotype consensus loci), losing

information related to heterozygosity. While this may be appropriate, or at least benign, for

phylogenetic analyses of diploid taxa (Kates et al. 2018), it may pose substantial problems when

attempting to investigate the evolutionary history of polyploid taxa or reticulate lineages.

While phylogenetic studies in plants often infer strictly bifurcating trees, the complexities

of allopolyploid evolution can be represented more accurately using networks. Simulations and

empirical analyses have suggested that phylogenetic networks can recover the reticulate

histories of polyploid lineages with few loci, at least when gene tree discordance due to

incomplete lineage sorting (ILS; Hudson et al. 1983; Pamilo and Nei 1988) is low (Oberprieler et

al. 2017) using parsimony methods (Huber et al. 2006; Lott et al. 2009), or even with moderate

ILS when explicitly modeled (Jones et al. 2013). Contemporary phylogenetic network models

and software packages jointly consider gene tree variation due to allele sampling error as

described by the multispecies coalescent (MSC; Rannala and Yang 2003) and gene flow

modeled as episodic introgression events (Solis-Lemus and Ané 2016; Wen et al. 2016; Zhang,

Ogilvie et al. 2018, Flouri et al. 2020). Depending on the complexity and goals of the research

question, these methods can search for networks with a constrained number of reticulation

events using quartet-based maximum pseudolikelihood (Solis-Lemus and Ané 2016; Wen et al.

2018) or a full-likelihood Bayesian model where the number of reticulations is a parameter (Wen

et al. 2018; Zhang et al. 2018). Also, it is possible to estimate model parameters (divergence

times, population sizes, and the fraction of introgressed genes) on a fixed species network using

a full-likelihood Bayesian model that allows efficient computation with large numbers of loci

(Flouri et al. 2020). We refer to these network models from here on as the multispecies

coalescent with introgression (MSci), consistent with Flouri et al. (2020); although, other names

have been used, such as the network multispecies coalescent (NMSC; e.g Zhu and Degnan

2017) and multispecies network coalescent (MSNC; e.g., Wen et al. 2016). We emphasize that

network approaches to investigate polyploid complexes are not novel (e.g., Huber et al. 2006;

Lott et al. 2009; Jones et al. 2013), but the difficulty of collecting appropriate genomic data from

polyploids for such analyses has limited their use.

To address the issues outlined above, we have developed a pipeline, PATÉ (Phased

Alleles from Target Enrichment data), that can phase genotyping data for individuals of a known

ploidy without the need for sampling their parental lineages. PATÉ was designed with scalability

and population-level sampling in mind for target enrichment projects where deep coverage from

paired-end Illumina data allow calling of high-quality variants. In this study, we first use

simulations to explore the ability of network approaches to reconstruct the history of

allopolyploidy in the presence of ILS, and whether phasing the data affects the accuracy of the

reconstruction. We show that using phased allelic data can improve network estimation and

divergence time estimation compared to using haplotype consensus sequences, but also

highlight scenarios where phasing may not be necessary or beneficial. Next, we describe the

individual steps used by PATÉ to phase target enrichment data. For an empirical example of the

benefits of PATÉ, we compare phased and unphased (haplotype consensus) data to infer the

evolutionary history of the North American *Dryopteris* complex, a model system for reticulate

polyploid evolution (Sessa et al. 2012a; Sessa et al. 2012b), using new targeted enrichment

data. The system includes four diploid species, as well as one extinct diploid, that have formed

five allopolyploids in which there is high confidence in the parent-progeny relationships (Fig. 1),

although numerous sterile allopolyploid species have also been reported within the complex

(Montgomery and Paulton 1981). The allopolyploid species have relatively ancient origins, with

the best estimates placing hybridization events between six and 13 Ma (Sessa et al. 2012b).

PATÉ is largely successful in recovering phased haploid sequences from polyploid individuals,

and networks inferred from phased data more accurately represent the North American

Dryopteris complex than those inferred from unphased data.

MATERIALS & METHODS

Testing the Effects of Phasing on Network Inference through Simulation

Simulating phased and unphased sequence data for an allopolyploid — We simulated

gene trees and their nucleotide sequence data using the MSC model with BPP v1.4.1 (Flouri et

al. 2018) under a five-species network (Fig. 2). All simulations used the Jukes-Cantor (Jukes

and Cantor 1969) model of sequence evolution with no rate heterogeneity. The allopolyploid

species E was treated as two lineages (E sister to B and F sister to C) whose alleles were

pooled to form the hybrid species E at time τ_h. This makes species E a tetraploid hybrid with the

parents B and C. θ was constant among lineages and set at 0.01. Assuming a per-generation

mutation rate (μ) of 1×10^{-8} and one year per generation yields an effective population size

(N_e) of 250,000 and root age of 10 Ma for the simulation network (Fig. 2). We also simulated

data under a shallow divergence scenario, in which all node ages were divided by 10, and a

deep divergence scenario, in which all node ages were multiplied by 10. This changes the root

age (τ_r) to 0.01 (1 Ma) and 1.0 (100 Ma), respectively.

Because the distance between speciation nodes is 0.025 and $\theta = 0.01$ for the simulation

network (Fig. 2), there are five coalescent units between nodes, $T = \frac{\tau}{\left(\frac{\theta}{2}\right)}$ (Yang 2006), which

implies a near-zero probability of gene tree discordance due to ILS (Hudson 1983). We

incorporated ILS into the simulation by reducing the node heights τ_u and τ_s to 0.0375 and 0.05

(for a low level of ILS) and 0.03 and 0.035 (for a moderate level of ILS). This increases the

probability of ILS at node u and s to 0.05 and 0.25, respectively. Thus, we used a total of nine

simulation conditions that combined three levels of evolutionary distance and three levels of ILS.

While our simulations do not explore extreme levels of gene tree discordance, they allow us to

learn about some general features of increasing ILS on network inference with different data

types. We simulated 1000 gene trees and their sequences, 500bp in length, under the MSC for

each of the nine conditions. We also explored the effects of sampling fewer genes on

downstream analyses. For each replicate of 1000 gene trees and their sequences, we randomly

sampled 400, 40, and four genes without replacement.

All simulations sampled haploid data. Two haploid sequences were sampled for each

diploid species and four haploid sequences were sampled for the allopolyploid species E, where

two sequences came from each parental lineage. We then investigated unphased data in three

ways. First, to generate unphased genotype sequences, the simulated haploid sequences for

each species were collapsed into a single sequence in which heterozygous sites were

represented by IUPAC ambiguity codes (genotype). The allopolyploid species was not restricted

to only biallelic sites. Second, we generated haploid consensus sequences, where for each

variable site, only one base was randomly retained (consensus). This could represent a case in

which read coverage across a locus is highly uneven such that a haploid sequence is actually a

chimera of two or more alleles. Finally, we simply picked one phased haploid sequence, which

is possible when one parental haplotype has a majority of reads for a locus (pick one). This

scenario where only one parent's sequence would be recovered in the offspring could be

anticipated in real data due to subgenome dominance (e.g., Buggs et al. 2014; Bird et al. 2018).

In practice, we expect most *de novo* assemblers to generate output in between the haploid

consensus data and pick one data.

Inferring species networks with phased and unphased simulated data — We estimated

species networks with PhyloNetworks v0.12.0 (Solis-Lemus et al. 2017) using Julia v.1.4.1

(Bezanson et al. 2015) from either the true gene trees used to simulate the data, or gene trees

estimated from the phased or unphased sequence data. For estimated trees, we used IQTREE

v1.6.10 (Nguyen et al. 2015) with the same model used for simulation. Each PhyloNetworks

analysis used the species tree (A,((B,E),(C,D))) as the starting tree and allowed zero, one, or

two reticulation events. Each analysis included ten independent optimizations of the

pseudolikelihood score. We considered larger numbers of reticulations an improvement if they

were two or more pseudolikelihood units lower than the best model. We compared the

estimated networks with one reticulation to the true network with the *hardwiredClusterDistance*

function (Huson et al. 2010) in PhyloNetworks. This allowed us to score the number of

replicates that 1) recovered the correct number of reticulations and 2) matched the true network

when the number of reticulations was set to one. We estimated networks for samples of four,

40, 400, and 1000 gene trees for each of the 30 replicates for each of the nine simulation

divergence and ILS conditions.

Effect of phasing on divergence time estimation — We also used our simulated phased

and unphased data to estimate divergence times under the MSci model (Flouri et al. 2020)

using BPP v4.2.9. Here, we estimate divergence times (τs), population sizes (θs), and the

proportion of introgressed loci (φ) on the correct fixed species network. MSCi analyses used

diffuse priors on τ_r and θ with a mean on their simulated values with $\varphi \sim \beta(1,1)$. Phased,

consensus, and pick one sequences were treated as haploids while genotype sequences were

treated as unphased diploids and used the analytical phasing (Gronau et al. 2011) implemented

in BPP. Although this is not correct for the tetraploid, it is arguably more appropriate than

treating all of the genotype sequences as haploid. Each Markov chain Monte Carlo (MCMC)

analysis collected 10,000 posterior samples, saving every 100 generations, while discarding the

first 100,000 generations (i.e., 10% of the total run) as burnin. All scripts for simulation and

subsequent analyses of simulated data are available in Dryad (X).

A Phasing Pipeline for Polyploids

Target enrichment data — We were motivated by the general premise of using phased

data to infer reticulate evolutionary histories of polyploids based on the success of empirical

studies where phasing was informative about hybridization or introgression events (e.g.,

Oberprieler et al. 2017; Eriksson et al. 2018). We were aware of few instances of phasing

genomic or phylogenomic data in polyploids, except in cases where chromosome-level whole-

genome assemblies have characterized subgenomes in allopolyploid crops (Yang et al. 2017;

Colle et al. 2019) or emerging results that are dependent on the sampling of parental lineages

(Freyman et al. 2020; Nauheimer et al. 2020). We designed PATÉ for target enrichment data

because of the availability of such data for many of taxa, but it is applicable to other types of

data with paired-end Illumina reads.

The end product of a *de novo* target enrichment assembly pipeline (such as HybPiper;

Johnson et al. 2016) generally is a single consensus sequence for each locus for each

individual. Allelic variation may be represented by ambiguous nucleotide codes within the single

consensus sequence or lost when the pipeline outputs the haplotype consensus sequence

where the majority vote from a collection of reads is used. We use these existing *de novo*

assembly pipelines as a starting point to provide the reference sequence for each locus for each

individual and leverage a recent phasing algorithm with high-quality variant calls to recover

phased haplotype sequences for taxa with known ploidy levels. Ploidy levels were well-

characterized for individuals in our *Dryopteris* analyses, but for unknown systems there are

existing methods to estimate ploidy directly from target enrichment data (Weiss et al. 2018;

Viruel et al. 2019) in the absence of other sources, such as flow cytometry (Farhat et al. 2019).

Phasing alleles within loci — PATÉ (Fig. 3) starts with assembled target enrichment loci,

such as the supercontig files output from HybPiper (Johnson et al. 2016) that contain a single

haplotype consensus sequence from each individual per locus. Reads for each individual are

then realigned to their consensus locus using BWA v0.7.17 (Li and Durbin 2009). PCR

duplicates are flagged with MarkDuplicates in Picard v2.9.2

(http://broadinstitute.github.io/picard), and variant calls are computed with HaplotypeCaller in

GATK v.4.1.4 (McKenna et al. 2010). We applied the following hard filters with VariantFiltration

in GATK: (1) QD < 2.0, (2) FS > 60.0, (3) MQ < 40.0, (4) ReadPosRankSum < -8.0, (5) AF <

0.05 || AF > 0.95. These loosely follow community recommendations on filters for germline

variant discovery (DePristo et al. 2011). Notably, we do not perform quality score recalibration

or filter on the mapping quality rank sum, as we anticipate allopolyploids could have a lower

mapping quality associated with an alternate allele due to sequence divergence or structural

variation among homeologous chromosomes. We also consider a very narrow window for

filtering on allele frequency. Because increasing ploidy levels will generate smaller anticipated

ratios of alternate to reference alleles, coupled with sequencing error and read stochasticity, we

only aim to remove the most extreme cases. For example, if almost all reads support the

alternate allele at a site, it is difficult to diagnose if the error lies in the consensus assembly or

the read alignment. In these cases, only the reference site is retained, and the variant does not

pass the allele frequency filter. However, the allele frequency filter could be removed if

investigators are focused on organisms with extremely high ploidy levels.

Biallelic SNPs that pass filters are then phased with H-PoPG v.0.2.0 (Xie et al. 2016). H-

PopG solves a heuristic phasing problem efficiently using dynamic programming. Although not

guaranteed to be an optimal solution, H-PoPG has been shown to have high accuracy while

also being fast (Xie et al. 2016; He et al. 2018; Moeinzadeh et al. 2020). Phasing variants in

polyploids is difficult because for n variants and k ploidy, there are $2^{n-1}(k-1)^n$ possible ways

to link the sites together. H-PoPG evaluates possible solutions efficiently by grouping reads into

k groups in such a way that differences within the groups are minimized. Focusing on target

enrichment data also constrains the complexity of the phasing problem compared to whole-

genome alignments (i.e., haplotype blocks are constrained to about 1000 bp). We then used the

phased variants to create individual allele sequences, where invariable sites are filled in based

on the reference sequence. In cases where there is no linkage information to phase across the

entire locus, we retain the phasing only for the longest block. The variants for the shorter

haplotype blocks can be collapsed into IUPAC ambiguity codes or treated as missing data

based on the investigator's preferences. PATÉ outputs analysis-ready fasta files with multiple

alleles per species. Variants are only phased within loci; we do not attempt to assign loci to

parental subgenomes. While this may complicate analyses of concatenated multi-locus

datasets, it is ideal for the MSC that assumes free recombination between loci and can leverage

multiple alleles per species for estimating θs. Those interested in concatenated analyses can

use other recent approaches that assign gene copies to parental subgenomes (Freyman et al.

2020; Nauheimer et al. 2020).

Analyses of a Species Complex with Allopolyploidy

The North American wood fern complex (Dryopteris) — We tested PATÉ using new

target enrichment data from nine North American *Dryopteris* species, including both

allotetraploid and allohexaploid taxa, with well-studied reticulate relationships (Sessa et al.

2012a, b) as well as two outgroup taxa from the sister genus *Polystichum*. All putative parental

lineages are represented in our dataset, with the exception of a hypothesized extinct lineage (*D.*

semicristata; Sessa et al. 2012b). We sampled two or three individuals for each *Dryopteris*

taxon (Table S1). The target enrichment data were generated from the GoFlag 408 flagellate

land plant probe set (Breinholt et al. 2021) at RAPID Genomics (Gainesville, FL). The target

regions for this probe set are 408 exons found in 229 single or low-copy nuclear genes. We

generated haplotype consensus assemblies for each with HybPiper (Johnson et al. 2016). The

resulting supercontig sequences became our reference sequences for genotyping and phasing

with PATÉ. We aligned both phased and unphased (i.e., the reference supercontig sequences)

with MUSCLE with default settings (Edgar 2004).

Three species tests — We first explored the value of phasing data when estimating

reticulate relationships among three species, including two diploid parental lineages (*D.*

expansa and *D. intermedia*) and their putative allotetraploid descendent (*D. campyloptera*). The

two diploid parents last shared a common ancestor during the Late Eocene and Early Miocene,

approximately 23 Ma (Sessa et al. 2012b). We used both a full-likelihood Bayesian approach

and a topology-based pseudolikelihood approach to estimate the correct species relationships

from phased and unphased data. First, using BPP v.4.1.4 (Flouri et al. 2020), we estimated log-

marginal likelihoods (ln *mL*) with stepping-stone sampling (Xie et al. 2011) for the three possible

rooted three taxon trees and twelve possible network models that imply differences for the

timing and direction of allopolyploidy and the presence of unsampled ancestral lineages

(Supplementary Fig. S1). Each ln *mL* estimate used 24 steps, and each step had a posterior

sample of 10,000, saving every 100 generations after a 100,000 generation burnin (10% of the

total run). The ln *mL* values were then used to calculate the fifteen model probabilities following

equation 1 (e.g., Beerli et al. 2019).

$$P(model) = \frac{exp(ln\ mL_{model}\ -\ ln\ mL_{max})}{\Sigma_i[exp(ln\ mL_i\ -\ ln\ mL_{max})]} \quad Equation\ 1$$

We repeated analyses for 30 random subsets of 40 and then four loci to explore the

effects of the number of loci on inferring allopolyploidy. Next, we used PhyloNetworks to test the

presence and placement of gene flow between the three species. For that analysis, we also

included the sequences from the two *Polystichum* outgroups. IQ-TREE v1.6.10 (Nguyen et al.

2015) and model selection by ModelFinder (Kalyaanamoorthy et al. 2017) was used to estimate

gene trees for the phased and unphased data. The starting tree was obtained with ASTRAL III

v5.6.3 (Zhang, Rabiee et al. 2018). We tested the presence of zero, one, or two reticulations

with slope heuristics (Solis-Lemus and Ané 2016). Each analysis used ten independent

optimizations. In addition to the dataset of all loci, we analyzed the same 30 random subsets of

40 and four loci from the marginal likelihood analyses.

Nine species tests — We also investigated the differences in results from phased versus

consensus sequences when estimating a network for the nine-species complex, which involves

multiple reticulation events on an edge and thus should be difficult for network estimation (Solis-

Lemus and Ané 2016). Because the increased number of species and complexity of reticulation

in *Dryopteris,* evaluation with marginal likelihoods was not computationally feasible. Instead, we

performed analyses of the nine-species complex and two outgroups with phased and unphased

haplotype consensus data with PhyloNetworks as described above, but we allowed up to six

reticulation events. We used ASTRAL III v5.6.3 (Zhang, Rabiee et al. 2018) to generate the

starting species tree for network estimation using gene trees inferred from IQ-TREE v1.6.10

(Nguyen et al. 2015) with the best model selected by ModelFinder (Kalyaanamoorthy et al.

2017).

RESULTS

Simulation Shows Benefits of Phased Data

Network inference — In our simulation results, both phased (i.e., the haplotypic allele

sequences) and unphased data (i.e., genotype, consensus, and pick one) performed well when

the goal is only to detect the correct number of reticulations (Supplementary Fig. S2). The only

case where analyses did not converge to the true number of reticulations was in the presence of

moderate ILS and high nucleotide divergence; however, this appears largely due to gene tree

estimation error, as analyses using the true simulated gene trees greatly outperformed those

using gene trees estimated from the simulated sequence data. However, using phased data

provides more accurate estimates of the placement of the reticulation edge in comparison to

using genotype data, and to a lesser extent, consensus sequences (Fig. 4). When the true gene

trees were used, which have information about the allopolyploid's hybridization event (i.e., the

allele sequences are sister to their respective parents in every tree), the correct network can be

inferred with 40 or fewer loci when nucleotide divergence is moderate. The gene trees

estimated from phased data perform equally well, although they require a few more gene trees

when nucleotide divergence is low and ILS moderate. Analyses using the genotype data almost

never recover the true network for these medium and low divergence scenarios, regardless of

the amount of ILS. Analyses based on pick one data perform similarly well to the phased and

true data when sampling 400 loci, but they are less accurate for four or 40 loci at low and

medium divergences. Analyses using the consensus data perform poorly for low numbers of loci

under a low divergence and no ILS scenario, but they are capable of recovering the true

reticulation when sampling 400 or more loci for the other low and medium divergence cases. In

the high divergence simulations, all data types could infer the true network if enough genes

were sampled, but analyses with the phased data required fewer genes than others.

Divergence times — Using phased data also improves divergence time estimates when

nucleotide divergence is low, but not when divergence is moderate or high (Fig. 5). When

divergence was low, the timing of reticulate events was accurately estimated when using

phased haplotypic data, but was overestimated when using genotype and especially consensus

data. For analyses with genotype and consensus data, as the number of loci increased and

uncertainty in the posterior was reduced, the posterior mean did not converge to the true

estimate and the simulated value was not within the highest posterior density (HPD) interval.

Additionally, all other nodes in the species network were overestimated with genotype or

consensus data, while the phased data were capable of recovering the simulated divergence

times (Supplementary Figs. S3-S5). Aside from some cases with the consensus sequences, all

four data types performed similarly with four loci; however, this is likely due to the posterior

being dominated by the prior in the absence of enough data. There was little improvement in

divergence time estimates when going from 40 to 400 loci, aside from further reduction in the

HPD intervals.

For medium sequence divergence, there was little difference between the phased and

unphased data. Phased sequences slightly underestimated the timing of hybridization while

unphased data slightly overestimated the timing of these events. However, phased data

accurately estimated the age of older speciation nodes that were again systematically

overestimated with unphased data (Supplementary Figs. S6-S8). At a high level of nucleotide

divergence, genotype data were capable of accurately estimating all divergence time

parameters while the phased data underestimated the timing of hybrid events (Fig. 5;

Supplementary Figs. S9-S11). Notably, the pick one data performed very well for all divergence

time estimation scenarios. Divergence times were not strongly affected by increasing levels of

ILS, likely because estimates were performed with the MSci model and we did not explore very

high ILS scenarios, but age estimates improved slightly for the genotypes, consensus, and pick

one data with increasing ILS.

Analyses of target enrichment data from Dryopteris

Recovery of Phased Loci — On average, 62% of loci sequenced for an individual were

phased with eight variants passing filters (Table S2). The ploidy level appears to be strongly

associated with the number of phased loci. Among diploids, only 30% of loci were phased; the

other 70% of diploid loci were either homozygous or had too few linked variants for phasing. For

tetraploids and hexaploids, 87% and 94% of loci, respectively, were phased such that two or

more phased haplotype sequences could be recovered. Among loci where phasing was

possible, variants were almost always resolved as a single haplotype block, as opposed to

being split into two or more blocks because not enough reads were available to physically link

variants. For polyploids, the number of phased haplotype sequences most frequently matched

the ploidy level except in the case of a single *D. campyloptera* individual (B087-D08), which also

had few recovered loci. Phasing data only extended sequence alignment length by about four

base pairs, but it more than doubled the number of parsimony informative sites (Table S3).

Placing a single reticulation event — For our three-species full-likelihood analyses with

the MSCi, both phased and unphased data recovered the anticipated reticulation hypothesis,

identifying *D. campyloptera* as an allotetraploid with the two diploid parental lineages *D.*

expansa and *D. intermedia*, when using all loci (Table 1). Model probabilities show decisive

support for a scenario where there are two unsampled ancestral populations that were the

progenitors of *D. campyloptera*, as opposed to *D. campyloptera* being a hybrid species with

extant *D. expansa* and *D. intermedia* as parents. All model parameters converged for both

phased (Supplementary Figs. S12 and S13) and unphased data (Supplementary Figs. S14 and

S15). Divergence time estimates were older in the analysis of phased data, although the relative

order of divergence events was the same using phased and unphased data (Fig. 6). There was

more uncertainty in the θ estimates, especially for the alloplyploid species *D. campyloptera* and

the two ancestral populations of the parental lineages that formed the polyploid ancestor

(Supplementary Figure S16). Repeating the analyses with fewer loci did not always produce the

same result, but there was either decisive support for a model or multiple plausible models that

all had the correct species relationships and direction of introgression for both phased and

unphased data when using 40 loci (Fig. 7). The only difference between models was the

presence or absence of ancestral θs for unsampled lineages. Analyses with four loci produced

less reliable results for both phased and unphased data. The four-locus analyses of phased and

unphased data found some non-negligible model probability for trees without hybridization or

networks where hybridization was incorrect in nine and twelve out of 30 replicates, respectively

(Fig. 7).

When performing a network search based on gene tree distributions, both the phased

and unphased data were able to recover the hypothesized allopolyploidy event (Fig. 8;

Supplementary Table S5). Both analyses inferred the major branch to be *D. intermedia* with the

minor branch from *D. expansa*. Phased data estimated a slightly smaller inheritance probability

compared to the unphased data. These findings from PhyloNetworks are consistent with

parameter estimation under the MSci model, where phased data estimated a slightly smaller

mean φ_h compared to unphased data (Table S4). When sampling 100 replicates of 40 loci, 98%

of replicates for phased data and 100% for unphased data were capable of detecting a single

hybrid event in the data. When sampling four loci, this drops to 66% and 85%, respectively (Fig.

8). Phased data more frequently recovered the correct network (58%) compared to unphased

data (38%) with 40 loci; however, the converse is true for four loci, with 34% correct for phased

and 41% correct for unphased (Fig. 8). Unphased data also got the network wrong more

frequently than phased data, such that phased data had the direction of introgression incorrect

in 36% and 12% of replicates while unphased data were incorrect in 60% and 38% replicates for

40 and four loci, respectively (Fig. 8).

Inferring relationships among a complex with multiple reticulation events — The network

recovered for phased data identified three out of five hypothesized reticulation events among

the nine *Dryopteris* species (Fig. 9; Supplementary Table S5). The allotetraploid *D. celsa* was

correctly identified with diploid *D. ludoviciana* and *D. goldiana* as parents. Analysis of the

phased data detected a low level of gene flow from the common ancestor of this clade into

tetraploid *D. cristata,* which has *D. ludoviciana* as one hypothesized parent while the other

parent lineage (*D. semicristata*) is assumed to be extinct (Sessa et al. 2012b). *Dryopteris*

carthusiana is another tetraploid that is assumed to share the extinct common ancestor with *D.*

cristata, but has experienced introgression from *D. intermedia*, with a high inheritance

probability of 0.43 (Fig. 9). However, the phased data missed the putative allotetraploid case of

D. campyloptera, despite the strong evidence for this reticulation event in our earlier three-taxon

analyses. Our network with phased data also failed to identify the putative reticulate

evolutionary history of *D. clintoniana*, an allohexaploid where allotetraploid *D. cristata* and

diploid *D. goldiana* are assumed to be the parents (Sessa et al. 2012b). The spectra of quartet

concordance factors were overall similar between the phased and unphased data

(Supplementary Fig. S17), but the phased data were arguably more accurate.

Although the phased data were not successful in recovering all hypothesized reticulate

relationships, they performed better than the unphased data. Analyses of unphased data were

capable of finding the allotetraploid history of *D. celsa* with an inheritance probability similar to

the phased data (Fig. 9). Hybridization between *D. intermedia* and *D. carthusiana* was also

detected; however, the directionality was reversed, with gene flow going from the allotetraploid

into the diploid. A third reticulation edge was found in the unphased analysis, from the common

ancestor of *Dryopteris* into the common ancestor of *D. clintoniana* and its sister clade. This

hybrid edge is difficult to reconcile because of the hypothesized extinct common ancestor that

contributed to both *D. cristata* and *D. carthusiana*. *Dryopteris clintoniana* is correctly placed in

the major species tree topology, as a grade between *D. cristata* and *D. goldiana*. This

reticulation edge from the *Dryopteris* common ancestor may reflect the extinct lineage and a

high degree of gene tree variation.

DISCUSSION

New phylogenetic network methods offer the promise of elucidating the often complex

reticulate histories of polyploid lineages, even in the presence of ILS. Our results demonstrate

that phasing polyploid target enrichment data can improve the accuracy of such network

inferences as well as divergence time estimates for the networks, and we describe a novel

pipeline (PATÉ) to address the difficult problem of phasing polyploid data. Although PATÉ could

handle different types of genomic data, such as transcriptomes and whole genomes, target

enrichment data are ideal for investigation because they often yield high and even coverage

across loci. Because MSC methods assume treat loci independently and assume free

recombination between loci, it is not necessary to assign individual loci to parental subgenomes.

However, the allele sequences output by PATÉ can also be used as input for the recently

developed *Homologizer* (Freyman et al. 2020) or *HybPhaser* (Nauheimer et al. 2020), which

attempt to phase across loci and recover parental subgenomes. Our methods also enable

population genomic studies of polyploids where accurate estimation of site frequency spectra

can be used for demographic analyses otherwise complicated by polyploidy (e.g., Excoffier et

al. 2013; Liu and Fu 2020) or SNP-based network inference in the absence of variation suitable

for gene tree estimation (Blischak et al 2018; Olave and Meyer 2020).

Promises and Pitfalls of Phasing

The prospect of using alleles from phased genomic data presents an exciting step

towards revealing the evolutionary history of polyploids, which remains a critical impediment

within the plant evolution community (McKain et al. 2018). Strategies for explicitly addressing

this challenge are only now emerging (Freyman et al. 2020; Nauheimer et al. 2020), and PATÉ

can be a useful tool by phasing variants for many individuals while leveraging genotyping

information. Our simulations demonstrate that phasing can improve estimates of reticulate

evolutionary relationships using network methods. Phased data can more accurately recover

the placement and directionality of hybrid edges than various types of unphased data in

simulations (Fig. 4) and empirical analyses with limited numbers of loci (Fig. 7). The advantages

of using phased versus unphased data for network estimation decrease when a large number of

loci are sampled (Table 1; Fig. 8).

Perhaps an underappreciated aspect of phased data is their ability to improve

divergence times estimates (Fig. 5; Supplementary Figs. S3-S11; Anderman et al. 2019). Our

empirical analyses also demonstrated how the timing of introgression τ_h can be greatly affected

by whether phased or consensus data are used. In our *Dryopteris* analyses, the estimate from

phased data was nearly four times older than the estimate from unphased data (Supplementary

Table S4). The direction of this difference was unanticipated, because our simulations

suggested the consensus data should overestimate age compared to phased data. This

highlights the difficulties of simulating data that capture real complexities and makes deciding

which estimate is more reliable somewhat difficult; however, the uncertainty of τ_h for haplotype

consensus data reflected in its posteriors (Supplementary Fig. S14) gives us more confidence in

the phased estimates (Supplementary Fig. S12). Because we used the MSci model for

divergence time estimation, we did not observe any effect of ILS on age estimates in our

simulations, but if we were using concatenation methods to date the divergence times for the

allopolyploids two subgenomes, nodes affected by ILS should be overestimated (Tiley et al.

2020).

In most cases phasing appears to be beneficial, but it may be problematic when the

parental lineages are deeply diverged. Although the phased data were able to accurately

estimate the age of older speciation nodes (Supplementary Figs. S3-S11), as phylogenetic

information is lost from multiple hits, the influence of the prior becomes more substantial. These

deep divergence simulation scenarios may border on being biologically unrealistic, as identifying

an allopolyploid and its two parental lineages becomes more difficult over time due to extinction

and population genetic processes, but it provides some expectations for the performance of

phased and unphased data in the presence of high nucleotide divergence. Our simulations

showed that when alleles are phased but only one is sampled, as in our pick one simulations,

network and divergence time estimation is similar to having all phased alleles present. We also

showed where consensus data can perform poorly through simulation (Figs. 3 and 4) as well as

empirical analyses where the direction of introgression was more frequently reversed in a

simple example of two parental lineages and an allopolyploid (Fig. 8). When enough reads are

available to call high-confidence variants, we suggest that phasing with PATÉ can improve

network and divergence time estimation for species complexes with low to moderate sequence

divergence. However, when investigating very ancient hybrid events, unphased genotype data

may be preferential, and using analytical approaches that integrate over phases (Gronau et al.

2011; Flouri et al. 2020) may outperform analyses with phased sequences because allele-

specific information no longer captures shared ancestry with parental lineages and haplotype

blocks become smaller due to recombination.

Challenges of Network Estimation

Our analyses highlight the difficulty of estimating the evolutionary histories of reticulate

complexes using phylogenetic methods, regardless of data type. The full-likelihood

implementation of the MSci model appears to be useful for limited cases, but these methods are

computationally demanding. Thus, they may not be practical for generating hypotheses and

exploring unknown relationships for large numbers of taxa (e.g., Zhang et al. 2018). Quartet-

based methods are fast and accurate when there are limited numbers of reticulations (Solis-

Lemus and Ané 2016) and show similar accuracy to full-likelihood methods for estimating

introgression probabilities (Flouri et al. 2020). However, there are scenarios where true network

topologies become non-identifiable (Solis-Lemus and Ané 2016). For example, when multiple

introgression events affect the same lineage, the expected quartet distribution under the MSci

model becomes a poor fit for the empirical data (Cai and Ané 2020) and the network estimated

may be incorrect. These effects were evident in our empirical analyses where the relationship

between *D. campyloptera*, *D. expansa*, and *D. intermedia* was missing in the nine-species

analysis (Fig. 9), which we expect is due to a reticulation edge present between *D. intermedia*

and *D. carthusiana*. Phasing sequence data adds information that can improve estimates

(Supplementary Table S3), but unsampled or extinct lineages, such as the hypothesized *D.*

semicristata, can create significant barriers to recovering the true evolutionary history of

reticulate complexes, regardless of how many loci or individuals are available.

Insights into Dryopteris Evolution

The North American *Dryopteris* complex has been well-characterized through the study

of multi-locus nuclear and chloroplast phylogenies, morphology, cytological observations,

chromatography, and isozyme analyses (reviewed in Sessa et al. 2012b). This makes it a useful

system for testing our phasing pipeline, and our analyses add nuance to our understanding of

some of the relationships among *Dryopteris* species. For example, our results indicate *D.*

campyloptera has received slightly more loci from *D. intermedia* than *D. expansa* (Fig. 8), and

D. intermedia is thought to be the likely maternal progenitor (Sessa et al. 2012b). Although our

analysis of marginal likelihoods for all target enrichment loci suggested the presence of

unsampled ancestral populations (Table 1), the age of introgression and divergence of the *D.*

campyloptera ancestral population from the *D. intermedia* and *D. expansa* parental lineages is

consistent with hybrid speciation rather than a lineage that was isolated from *D. intermedia* and

received more recent gene flow from *D. expansa* (Fig. 6). Following the allopolyploidy event, the

D. intermedia genome was likely dominant, providing some selective advantage for *D.*

campyloptera in its distribution at the time (Bird et al. 2018). Similar insights can be gained from

the nine-taxon analyses, which suggests *D. ludoviciana* is the dominant genome in *D. celsa*.

Dryopteris ludoviciana is the maternal parent of *D. celsa*, again suggesting some bias in

retaining homeologous alleles from the maternal lineage, which provides the chloroplast

genome in ferns (Sessa et al. 2012b).

The nine-taxon analyses also support the hypothesis of the unsampled diploid lineage *D.*

semicristata, based on the placement of *D. carthusiana* as sister to the rest of *Dryopteris* in the

phased nine-taxon analyses, but with *D. carthusiana* having received over 40% of its genes

from *D. intermedia* more recently. Our analyses suggest that *D. cristata* did not have *D.*

ludoviciana as a progenitor, but rather an unsampled common ancestor of *D. goldiana* and *D.*

ludoviciana. In the case of *D. cristata*, both parental diploid lineages, including *D. semicristata,*

may have gone extinct.

Conclusions

Combining phased data with recent network methods holds much promise for

confronting a major challenge of plant phylogenetics: resolving the complex histories of

polyploids. The PATÉ pipeline can enhance systematic, speciation genomic, and population

genomic analyses of groups containing polyploids. While haplotype consensus sequences may

be adequate for resolving single reticulation events where both parents are sampled, using

phased sequences can improve inferences of more complicated allopolyploid events,

demonstrating how allelic variation can be leveraged for MSC methods that account for

reticulation. Still, some reticulate complexes can be difficult to disentangle with any data when

there are multiple hybrid events involving the same branch. PATÉ is available through GitHub

(https://github.com/gtiley/Phasing) and can be run on any UNIX environment after installing

basic genotyping software and H-PoPG.

DATA AVAILABILITY

PATÉ is freely available through GitHub at https://github.com/gtiley/Phasing. Simulated

and empirical data supporting findings and files for replicating analyses are available from the

Dryad Digital Repository: (X). Raw Fastq reads for *Dryopteris* individuals are available through

the NCBI SRA database and are associated with BioProject PRJNA725004. Individual SRA

Identifiers are available in Supplementary Table S1.

SUPPLEMENTARY MATERIAL

Data available from the Dryad Digital Repository: (X).

ACKNOWLEDGEMENTS

The authors thank A.M. Duffy, K. Imwattana, M. Nieto-Lugilde, B.T. Piatkowski, and A.J.

Shaw for helpful discussions and providing feedback on the manuscript. We also thank M.G

Johnson, J. Mendez Reneau, and L. Nauheimer for discussions and sharing their strategies for

phasing sequence data. We are grateful to the people who made data collection possible; we

used *Dryopteris* samples collected by C.J. Rothfels, M.A. Sundue, and W. Testo, and DNA

extractions were performed by S.B. Carey and E. Lockwood.

FUNDING

Funding was provided by National Science Foundation awards DEB-2038213 to AAC

and PSM, and from DEB-1541506 to JGB and EBS. This work was also partially supported by

the Department of Energy award DE-SC0021016 to CSL. ADY and GPT gratefully acknowledge

support from Duke University.

608 **LITERATURE CITED**

609 Andermann T., Cano A., Zizka A., Bacon C., Antonelli A. 2018. SECAPR-a bioinformatics
610 pipeline for the rapid and user-friendly processing of targeted enriched Illumina sequences, from
611 raw reads to alignments. *PeerJ* 6:e5175.
612
613 Andermann T., Fernandes A.M., Olsson U., Topel M., Pfeil B., Oxelman B., Aleixo A., Faircloth
614 B.C., Antonelli A. 2019. Allele Phasing Greatly Improves the Phylogenetic Utility of
615 Ultraconserved Elements. *Syst. Biol.* 68:32-46.
616
617 Baaijens J.A., Schonhuth A. 2019. Overlap graph-based generation of haplotigs for diploids and
618 polyploids. *Bioinformatics* 35:4281-4289.
619
620 Bankevich A., Nurk S., Antipov D., Gurevich A.A., Dvorkin M., Kulikov A.S., Lesin V.M.,
621 Nikolenko S.I., Pham S., Prjibelski A.D., Pyshkin A.V., Sirotkin A.V., Vyahhi N., Tesler G.,
622 Alekseyev M.A., Pevzner P.A. 2012. SPAdes: a new genome assembly algorithm and its
623 applications to single-cell sequencing. *J. Comput. Biol.* 19:455-477.
624
625 Barker M.S., Arrigo N., Baniaga A.E., Li Z., Levin D.A. 2016. On the relative abundance of
626 autopolyploids and allopolyploids. *New Phytol.* 210:391-398.
627
628 Beerli P., Mashayekhi S., Sadeghi M., Khodaei M., Shaw K. 2019. Population Genetic Inference
629 With MIGRATE. *Curr. Protoc. Bioinformatics* 68:e87.
630
631 Berger E., Yorukoglu D., Peng J., Berger B. 2014. HapTree: a novel Bayesian framework for
632 single individual polyplotyping using NGS data. *PLoS Comput. Biol.* 10:e1003502.
633
634 Bezanson J., Edelman A., Karpinski S., Shah V.B. 2015. Julia: A Fresh Approach to Numerical
635 Computing. *arXiv*1411.1607v4.
636
637 Bird K.A., VanBuren R., Puzey J.R., Edger P.P. 2018. The causes and consequences of
638 subgenome dominance in hybrids and recent polyploids. *New Phytol.* 220:87-93.
639
640 Blischak P.D., Chifman J., Wolfe A.D., Kubatko, L.S. 2018. HyDe: a Python package for
641 genome-scale hybridization detection. *Syst. Biol.*, 67:821-829.
642
643 Breinholt J.W., Carey S.B., Tiley G.P., Davis E.C., Endara L., McDaniel S.F., Neves L.G., Sessa
644 E.B., von Konrat M., Chantanaorrapint S., Fawcett S., Ickert-Bond S.M., Labiak P.H., Larraín J.,
645 Lehnert M., Lewis L.R., Nagalingum N.S., Patel N., Rensing S.A., Testo W., Vasco A., Villarreal
646 J.C., Williams E.W., Burleigh J.G. 2021. A target enrichment probe set for resolving the
647 flagellate plant tree of life. *Appl. Plant Sci.* 9:e11406.
648
649 Breinholt J.W., Earl C., Lemmon A.R., Lemmon E.M., Xiao L., Kawahara A.Y. 2018. Resolving
650 relationships among the megadiverse butterflies and moths with a novel pipeline for anchored
651 phylogenomics. *Syst. Biol.* 67:78-93.
652
653 Buggs J.A., Wendel J.F., Doyle J.J., Soltis D.E., Soltis P.S., Coate J.E. 2014. The legacy of
654 diploid progenitors in allopolyploid gene expression patterns. *Philos. Trans. R. Soc. Lond. B*
655 *Biol. Sci.* 368:20130354.
656

657   Cai R., Ané C. 2020. Assessing the fit of the multi-species network coalescent to multi-locus
658   data. *Bioinformatics* btaa863. doi: https://doi.org/10.1093/bioinformatics/btaa863
659

660   Colle M., Leisner C.P., Wai C.M., Ou S., Bird K.A., Wang J., Wisecaver J.H., Yocca A.E., Alger
661   E.I., Tang H., Xiong Z., Callow P., Ben-Zvi G., Brodt A., Baruch K., Swale T., Shiue L., Song G.-
662   Q., Childs K.L., Schilmiller A., Vorsa N., Buell C.R., VanBuren R., Jiang N., Edger P.P. 2019.
663   Haplotype-phased genome and evolution of phytonutrient pathways of tetraploid blueberry.
664   *GigaScience* 8:giz012.
665

666   Crowl A.A., Myers C., Cellinese N. 2017. Embracing discordance: Phylogenomic analyses
667   provide evidence for allopolyploidy leading to cryptic diversity in a Mediterranean Campanula
668   (Campanulaceae) clade. *Evolution* 71:913-922.
669

670   DePristo M., Banks E., Poplin R., Garimella K., Maguire J., Hartl C., Philippakis A., del Angel G.,
671   Rivas M.A., Hanna M., McKenna A., Fennell T., Kernytsky A., Sivachenko A., Cibulskis K.,
672   Gabriel S., Altschuler S., Daly M. 2011. A framework for variation discovery and genotyping
673   using next-generation DNA sequencing data. *Nat. Genet.* 43:491-498.
674

675   Edgar R.C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
676   throughput. *Nucleic Acids Res.* 32:1792-1797.
677

678   Eriksson J.S., de Sousa F., Bertrand Y.J.K., Antonelli A., Oxelman B., Pfeil B.E. 2018. Allele
679   phasing is critical to revealing a shared allopolyploid origin of Medicago arborea and M.
680   strasseri (Fabaceae). *BMC Evol. Biol.* 18:9.
681

682   Excoffier L., Dupanloup I., Huerta-Sanchez E., Sousa V.C., Foll M. 2013. Robust demographic
683   inference from genomic and SNP data. *PLoS Genet.* 9:e1003905.
684

685   Faircloth B.C. 2016. PHYLUCE is a software package for the analysis of conserved genomic
686   loci. *Bioinformatics* 32:786-788.
687

688   Faircloth B.C., McCormack J.E., Crawford N.G., Harvey M.G., Brumfield R.T., Glenn T.C. 2012.
689   Ultraconserved elements anchor thousands of genetic markers spanning multiple evolutionary
690   timescales. *Syst. Biol.* 61:717-726.
691

692   Farhat P., Hidalgo O., Robert T., Siljak-Yakovlev S., Leitch I.J., Adams R.P., Bou Dagher-
693   Kharrat M. 2019. Polyploidy in the Conifer Genus *Juniperus*: An Unexpectedly High Rate. *Front.*
694   *Plant Sci.* 10:676.
695

696   Flouri T., Jiao X., Rannala B., Yang Z. 2020. A Bayesian Implementation of the Multispecies
697   Coalescent Model with Introgression for Phylogenomic Analysis. *Mol. Biol. Evol.* 37:1211-1223.
698

699   Flouri T., Jiao X., Rannala B., Yang Z.. 2018. Species Tree Inference with BPP Using Genomic
700   Sequences and the Multispecies Coalescent. *Mol. Biol. Evol.* 35:2585-2593.
701

702   Freyman W.A,. Johnson M.G., Rothfels C.J. 2020. Homologizer: Phylogenetic phasing of gene
703   copies into polyploid subgenomes. *bioRxiv* doi: https://doi.org/10.1101/2020.10.22.351486.
704

705   Gronau I., Hubisz M.J., Gulko B., Danko C.G., Siepel A. 2011. Bayesian inference of ancient
706   human demography from individual genome sequences. *Nat. Genet.* 43:1031-1034.
707

708 He D., Saha S., Finkers R., Parida L. 2018. Efficient algorithms for polyploid haplotype phasing.
709 *BMC Genomics* 19:110.
710
711 Huang J., Flouri T., Yang Z. 2020. A simulation study to examine the information content in
712 phylogenomic datasets under the multispecies coalescent model. *Mol. Biol. Evol.* 37:3211-3224.
713
714 Huber K.T., Oxelman B., Lott M., Moulton V. 2006. Reconstructing the evolutionary history of
715 polyploids from multi-labelled trees. *Mol. Biol. Evol.* 23:1784-1791.
716
717 Hudson R.R. 1983. Testing the Constant-Rate Neutral Allele Model with Protein Sequence
718 Data. *Evolution* 37:203-217.
719
720 Huson D.H., Rupp R., Scornavacca C. 2010. Phylogenetic networks: concepts, algorithms and
721 applications. Cambridge University Press.
722
723 Iqbal Z., Caccamo M., Turner I., Flicek P., McVean G. 2012. De novo assembly and genotyping
724 of variants using colored de Bruijn graphs. *Nat. Genet.* 44:226-232.
725
726 Jantzen J.R., Amarasinghe P., Folk R.A., Reginato M., Michelangeli F.A., Soltis D.E., Cellinese
727 N., Soltis P.S. 2020. A two-tier bioinformatic pipeline to develop probes for target capture of
728 nuclear loci with applications in Melastomataceae. *Appl. Plant Sci.* 8:e11345.
729
730 Johnson M.G., Gardner E.M., Liu Y., Medina R., Goffinet B., Shaw A.J., Zerega N.J., Wickett
731 N.J. 2016. HybPiper: Extracting coding sequence and introns for phylogenetics from high-
732 throughput sequencing reads using target enrichment. *Appl. Plant Sci.* 4:1600016.
733
734 Johnson M.G., Pokorny L., Dodsworth S., Botigue L.R., Cowan R.S., Devault A., Eiserhardt
735 W.L., Epitawalage N., Forest F., Kim J.T., Leebens-Mack J.H., Leitch I.J., Maurin O., Soltis
736 D.E., Soltis P.E., Wong G.K.-S., Baker W.J., Wickett N.J. 2019. A Universal Probe Set for
737 Targeted Sequencing of 353 Nuclear Genes from Any Flowering Plant Designed Using k-
738 Medoids Clustering. *Syst. Biol.* 68:594-606.
739
740 Jones G., Sagitov S., Oxelman B. 2013. Statistical inference of allopolyploid species networks in
741 the presence of incomplete lineage sorting. *Syst. Biol.* 62:467-478.
742
743 Jukes T.H., Cantor C.R. 1969. Evolution of protein molecules. In: Munro H.N., editor.
744 Mammalian Protein Metabolism. New York, NY: Acedemic Press. p. 21-132.
745
746 Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., von Haeseler A., Jermiin L.S. 2017.
747 ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* 14:587-
748 589.
749
750 Karimi N., Grover C.E., Gallagher J.P., Wendel J.F., Ané C., Baum D.A. 2020. Reticulate
751 Evolution Helps Explain Apparent Homoplasy in Floral Biology and Pollination in Baobabs
752 (Adansonia; Bombacoideae; Malvaceae). *Syst. Biol.* 69:462-478.
753
754 Kates H.R., Johnson M.G., Gardner E.M., Zerega N.J.C., Wickett N.J. 2018. Allele phasing has
755 minimal impact on phylogenetic reconstruction from targeted nuclear gene sequences in a case
756 study of Artocarpus. *Am. J. Bot.* 105:404-416.
757

758 Lemmon A.R., Emme S.A., Lemmon E.M. 2012. Anchored hybrid enrichment for massively
759 high-throughput phylogenomics. *Syst. Biol.* 61:727-744.
760
761 Li H., Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.
762 *Bioinformatics* 25:1754-1760.
763
764 Liu X., Fu Y.X. 2020. Stairway Plot 2: demographic history inference with folded SNP frequency
765 spectra. *Genome Biol.* 21:280.
766
767 Liu Y., Johnson M.G., Cox C.J., Medina R., Devos N., Vanderpoorten A., Hedenas L., Bell N.E.,
768 Shevock J.R., Aguero B., Quandt D., Wickett N.J., Shaw A.J., Goffinet B. 2019. Resolution of
769 the ordinal phylogeny of mosses using targeted exons from organellar and nuclear genomes.
770 *Nat. Commun.* 10:1485.
771
772 Lott M., Spillner A., Huber K.T., Moulton V. 2009. PADRE: a package for analyzing and
773 displaying reticulate evolution. *Bioinformatics* 25:1199-1200.
774
775 Luo R., Liu B., Xie Y., Li Z., Huang W., Yuan J., He G., Chen Y., Pan Q., Liu Y., Tang J., Wu G.,
776 Zhang H., Shi Y., Liu Y., Yu C., Wang B., Lu Y., Han C., Cheung D.W., Yiu S.-M., Peng S.,
777 Xiaoqian Z., Liu G., Liao X., Li Y., Yang H., Wang J., Lam T-W., Wang J. 2012. SOAPdenovo2:
778 an empirically improved memory-efficient short-read de novo assembler. *GigaScience* 1:18.
779
780 McKain M.R., Johnson M.G., Uribe-Convers S., Eaton D., Yang Y. 2018. Practical
781 considerations for plant phylogenomics. *Appl. Plant Sci.* 6:e1038.
782
783 McKenna A., Hanna M., Banks E., Sivachenko A., Cibulskis K., Kernytsky A., Garimella K.,
784 Altshuler D., Gabriel S., Daly M., DePristo M.A. 2010. The Genome Analysis Toolkit: a
785 MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.*
786 20:1297-1303.
787
788 Moeinzadeh M.H., Yang J., Muzychenko E., Gallone G., Heller D., Reinert K., Haas S., Vingron
789 M. 2020. Ranbow: A fast and accurate method for polyploid haplotype reconstruction. *PLoS*
790 *Comput. Biol.* 16:e1007843.
791
792 Montgomery J.D., Paulton E.M. 1981. *Dryopteris* in North America. *Fiddlehead Forum* 8:25-31.
793
794 Morales-Briones D.F., Liston A., Tank D.C. 2018. Phylogenomic analyses reveal a deep history
795 of hybridization and polyploidy in the Neotropical genus Lachemilla (Rosaceae). *New Phytol.*
796 218:1668-1684.
797
798 Nauheimer L., Weigner N., Joyce E., Crayn D., Clarke C., Nargar K. 2020. HybPhaser: a
799 workflow for the detection and phasing of hybrids in target capture datasets. bioRxiv. doi:
800 https://doi.org/10.1101/2020.10.27.354589.
801
802 Nguyen L.-T., Schmidt H.A., von Haeseler A., Minh B.Q. 2015. IQ-TREE: a fast and effective
803 stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* 32:268-
804 274.
805
806 Oberprieler C., Wagner F., Tomasello S., Konowalik K. 2017. A permutation approach for
807 inferring species networks from gene trees in polyploid complexes by minimising deep
808 coalescences. *Methods in Ecology and Evolution* 8:835-849.

809
810 Olave M., Meyer A. 2020. Implementing Large Genomic Single Nucleotide Polymorphism Data
811 Sets in Phylogenetic Network Reconstructions: A Case Study of Particularly Rapid Radiations of
812 Cichlid Fish. *Syst. Biol.* 69:848-862.
813
814 Pamilo P., Nei M. 1988. Relationships between gene trees and species trees. *Mol. Biol. Evol.*
815 5:568-583.
816
817 Rothfels C.J., Pryer K.M., Li F.-W. 2017. Next-generation polyploid phylogenetics: rapid
818 resolution of hybrid polyploid complexes using PacBio single-molecule sequencing. *New Phytol.*
819 213:413-429.
820
821 Saada O.A., Tsouris A., Freidrich A., Schachrer J. 2020. nPhase: An accurate and contiguous
822 phasing method for polyploids. *bioRxiv* doi:https://doi.org/10.1101/2020.07.24.219105.
823
824 Sessa E.B., Zimmer E.A., Givnish T.J. 2012a. Reticulate evolution on a global scale: a nuclear
825 phylogeny for New World Dryopteris (Dryopteridaceae). *Mol. Phylogenet. Evol.* 64:563-581.
826
827 Sessa E.B., Zimmer E.A., Givnish T.J. 2012b. Unraveling reticulate evolution in North American
828 Dryopteris (Dryopteridaceae). *BMC. Evol. Biol.* 12:104.
829
830 Solis-Lemus C., Ané C. 2016. Inferring Phylogenetic Networks with Maximum Pseudolikelihood
831 under Incomplete Lineage Sorting. *PLoS Genet.* 12:e1005896.
832
833 Solis-Lemus C., Bastide P., Ané C. 2017. PhyloNetworks: A Package for Phylogenetic
834 Networks. *Mol. Biol. Evol.* 34:3292-3298.
835
836 Soltis, D.E., Visger C.J., Soltis P.S. 2014. The polyploidy revolution then...and now: Stebbins
837 revisited. *American Journal of Botany* 101: 1057–1078.
838
839 Stull G.W., Soltis P.S., Soltis D.E., Gitzendanner M.A., Smith S.A. 2020. Nuclear phylogenomic
840 analyses of asterids conflict with plastome trees and support novel relationships among major
841 lineages. *Am. J. Bot.* 107:790-805.
842
843 Tiley G.P., Poelstra J.W., dos Reis M., Yang Z., Yoder A.D. 2020. Molecular Clocks without
844 Rocks: New Solutions for Old Problems. *Trends Genet.* 36:845-856.
845
846 Viruel J., Conejero M., Hidalgo O., Pokorny L., Powell R.F., Forest F., Kantar M.B., Soto Gomez
847 M., Graham S.W., Gravendeel B., Wilkin P., Leitch I.J. 2019. A Target Capture-Based Method
848 to Estimate Ploidy from Herbarium Specimens. *Front. Plant Sci.* 10:937.
849
850 Weiss C.L., Pais M., Cano L.M., Kamoun S., Burbano H.A. 2018. nQuire: a statistical framework
851 for ploidy estimation using next generation sequencing. *BMC Bioinformatics* 19:122.
852
853 Wen D., Yu Y., Nakhleh L. 2016. Bayesian Inference of Reticulate Phylogenies under the
854 Multispecies Network Coalescent. *PLoS Genet.* 12:e1006006.
855
856 Wen D., Yu Y., Zhu J., Nakhleh L. 2018. Inferring Phylogenetic Networks Using PhyloNet. *Syst.*
857 *Biol.* 67:735-740.
858

859   Wolf P.G., Robison T.A., Johnson M.G., Sundue M.A., Testo W.L., Rothfels C.J. 2018. Target
860   sequence capture of nuclear-encoded genes for phylogenetic analysis in ferns. *Appl. Plant Sci.*
861   6:e01148.
862
863   Wood T.E., Takebayashi N., Barker M.S., Mayrose I., Greenspoon P.B., Rieseberg L.H. 2009.
864   The frequency of polyplid speciation in vascular plants. *Proc. Natl. Acad. Sci. U. S. A.*
865   106:13875-13879.
866
867   Xie M., Wu Q., Wang J., Jiang T. 2016. H-PoP and H-PoPG: heuristic partitioning algorithms for
868   single individual haplotyping of polyploids. *Bioinformatics* 32:3735-3744.
869
870   Xie W., Lewis P.O., Fan Y., Kuo L., Chen M.H. 2011. Improving marginal likelihood estimation
871   for Bayesian phylogenetic model selection. *Syst. Biol.* 60:150-160.
872
873   Yang J., Moeinzadeh M-H., Kuhl H., Helmuth J., Xiao P., Haas S., Liu G., Zheng J., Sun Z., Fan
874   W., Deng G., Wang H., Hu F., Zhao S., Fernie A.R., Boerno S., Timmermann B., Zhang P.,
875   Vingron M. 2017. Haplotype-resolved sweet potato genome traces back its hexaploidization
876   history. *Nat Plants* 3:696-703.
877
878   Yang Z. 2006. Computational molecular evolution: Oxford University Press.
879
880   Zhang C., Ogilvie H.A., Drummond A.J., Stadler T. 2018. Bayesian Inference of Species
881   Networks from Multilocus Sequence Data. *Mol. Biol. Evol.* 35:504-517.
882
883   Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree
884   reconstruction from partially resolved gene trees. *BMC Bioinformatics* 19:153.
885
886   Zhu S., Degnan J.H. 2017. Displayed trees do not determine distinguishability under the
887   network multispecies coalescent. *Syst. Biol.* 66:283-298.
888

Topology	Phased Marginal lnL	Unphased Marginal lnL	Phased Model Probability	Unphased Model Probability
1	-613405.3154	-552720.6776	1.006E-278	3.2574E-214
2	-613223.0755	-552534.6540	1.4073E-199	2.0039E-133
3	-613366.4128	-552644.7603	7.9026E-262	3.0431E-181
4	-612866.6368	-552245.2218	8.86633E-45	1.00164E-07
5	-612828.7348	-552243.7091	2.56082E-28	4.54611E-07
6	-612822.4509	-552238.6462	1.37227E-25	7.18535E-05
7†	-612765.2027	-552229.1054	1	0.999927592
8	-613211.8583	-552484.8320	1.047E-194	8.6956E-112
9	-613168.7562	-552479.3732	5.4825E-176	2.0419E-109
10	-613102.1509	-552454.2002	4.6266E-147	1.74796E-98
11	-613079.9316	-552426.8867	2.0654E-137	1.27244E-86
12	-613379.6350	-552538.6937	1.4303E-267	3.5276E-135
13	-613390.9585	-552533.6798	1.7287E-272	5.3085E-133
14	-613272.7561	-552561.3165	3.7359E-221	5.2787E-145
15	-613247.7503	-552566.6741	2.7054E-210	2.4873E-147

891 **Table 1 — Marginal Likelihoods for Possible Topological Hypotheses.** Topologies of the 15

892 models are displayed in Figure S1.

893 †*a priori* allopolyploid hypothesis

Figure 1 — Hypothesized Relationships among North American *Dryopteris*. Synthesis of results from Sessa et al. 2012a and Sessa et al. 2012b. A) Links between shapes show the putative parents and their allopolyploid derivatives. Black circles are diploids, squares are tetraploids, and the hexagon is the one hexaploid species in the group. *Dryopteris semicristata* is presumed extinct. B) Placement of allopolyploids in the context of the backbone relationships among diploids. Tetraploids are indicated with solid orange lines and the hexaploid with dotted lines. The grey line denoting a sister relationship between *D. semicristata* and *D. expansa* reflects one possible placement for the extinct taxon based on previous analyses (Sessa et al., 2012b).

Figure 2 — Species network used for simulation. The divergence times in expected

substitutions per site are given for each node, and *h* is the hybrid node where two alleles enter

from both *t* and *v*. E is an allotetraploid while other species are diploid. Nucleotide divergence

was reduced by dividing all τ by 10 or increased by multiplying all τ by 10. ILS was increased by

halving the distance between τ_h and τ_u and between τ_u and τ_s either once (for the medium ILS

condition) or twice (for the high ILS condition).

Figure 3 — PATÉ Phasing Pipeline. Overview of data input, output, and steps taken to phase

alleles. Input data are assumed to be paired-end Illumina reads and reference sequences for

each individual are required (consensus loci from HybPiper can be used). The ploidy of each

individual must be specified. Only biallelic sites are used for phasing.

Figure 4 — Proportion of simulations that correctly identify the allopolyploid lineage. The x-axes are the number of loci sampled for each simulation. Th y-axes are the proportions of correct networks. Results are based on networks estimated with a single reticulation, even if that network was considered less optimal than networks with zero or two reticulations. We saved the true gene trees from the simulations, while we estimated gene trees with the phased and unphased (Genotype, Consensus, and Pick One) data.

Figure 5 — Estimating the Timing of Introgression. Divergence times are for node *h* in

Figure 2. The low divergence simulation corresponds to the y-axis units of $\tau_h \times 10^{-2}$ while the

high divergence case is represented by τ_h. Divergence times are measured as the expected

number of substitutions per site. The dashed line represents the true simulated values. Points

are posterior means, and error bars are 95% HPD intervals, averaged across 30 replicates.

Figure 6— *Dryopteris* Divergence Times under the MSci Model. Divergence times are

measured as the expected number of substitutions per site. The x-axis shows estimates from

phased data and the y-axis shows estimates from unphased (haplotype consensus) data for the

inset network (equivalent to model 7 from Supplementary Fig. S1). The dashed one-to-one line

shows where older age estimates are consistently obtained from phased data. Error bars on

points show the 95% HPD intervals for phased and unphased data.

Figure 7— Probabilities for three-taxon *Dryopteris* MSci Models with fewer Loci. Marginal

likelihoods were estimated for the 15 MSci models (Supplementary Fig. S1). Model weights

were used to obtain probabilities. We consider model probability greater than 0.95 as decisive

evidence in favor of a model, and a probability less than 0.05 is evidence against a model. We

considered probabilities between 0.05 and 0.95 to be ambiguous. Models four through seven (in

bold) all have the correct reticulate relationships between the parental diploid lineages and the

allopolyploid. Models one through three do not have introgression, and models eight through 15

have incorrect introgression events. Each row represents one of 30 sampling replicates.

Figure 8 — Network search results for three-taxon *Dryopteris* example. Both phased and unphased haplotype consensus data recovered the same network topology with introgression occurring in the expected direction. The major topology is indicated by the blue edge and the minor edge (direction of introgression) is shown in tan. The inheritance probability γ was slightly higher in the haplotype consensus data (γ_{unphased} = 0.455). Bar plots show the proportion of 100 replicates when sampling four or 40 loci that correctly detect one reticulation based on the pseudolikelihood scores (top), correctly estimate the network with introgression going from one of the diploids into *D. campyloptera* (middle), and estimate a network where the direction of introgression is from *D. campyloptera* into one of the diploids (bottom).

Figure 9 — Networks for Nine-Taxon *Dryopteris* example. Both data types recovered

optimal networks with three reticulation events. The major topology edge is blue and the minor

(reticulation) edges shown in tan, with the direction of introgression flowing into the major edge.

The position of *D. carthusiana* changes in the major topology between phased and unphased

data but the relationships are otherwise the same. All three reticulation events in the phased

data are plausible, but in the unphased data, the direction of introgression from *D. carthusiana*

into *D. intermedia* is incorrect and the reticulate edge from the common ancestor of *Dryopteris* is

difficult to reconcile. Inheritance probabilities for each introgression event are shown next to

reticulation edges.