The central nucleus of the amygdala and gustatory cortex assess affective valence during CTA learning and expression in male and female rats

Alyssa Bernanke[1], Elizabeth Burnette[1], Justine Murphy[1], Nathaniel Hernandez[1], Sara Zimmerman[1], Q. David Walker[1], Rylee Wander[1], Samantha Sette[1], Zackery Reavis[1], Christopher Armstrong[1], Mary Louise Risher[2], Cynthia Kuhn[1]

1 Department of Pharmacology and Cancer Biology

Duke University Medical Center

Durham NC

2 Department of Biomedical Sciences

Joan C. Edwards School of Medicine

Marshall University

Huntington West Virginia

Abstract

This study evaluated   behavior (Boost® intake, LiCl-induced behaviors, ultrasonic vocalizations (USVs), task performance) and c-fos activation during conditioned taste aversion (CTA) to understand how males and female rats assess the relative danger or safety of a stimulus in learning and performing a task. Females drank more Boost® than males but engaged in comparable aversive behaviors after LiCl treatment. Males produced 55 kHz USVs (indicating positive valence), when anticipating Boost® and inhibited these calls after pairing with LiCl. Females produced 55 kHz USVs based on their estrous cycle but were more likely to make 22 kHz USVs than males (indicating negative valence) after pairing with LiCl.   C-fos responses were similar in males and females after Boost® or LiCl. Females engaged the gustatory cortex more than males during the reinforcing task and males engaged the amygdala more than females in both the reinforcing and devalued tasks. RNAscope identified a population of D1-receptor expressing cells in the CeA that responded to Boost® and D2 receptor-expressing cells that responded to LiCl. The present study suggests that males and females differentially process the affective valence of a stimulus to produce the same goal-directed behavior.

Introduction

The ability to properly evaluate the relative safety of a stimulus is critical to survival across species. Dysfunction of this mechanism contributes to a range of mental health disorders, including anxiety disorders such as generalized anxiety, obsessive compulsive disorders, and post-traumatic stress disorders [1-5]. Anxiety disorders are one of the most common neuropsychiatric disorders, with a lifetime prevalence of 28.8% [6]. Perturbations in the ability to assess the relative danger or safety of a situation is a key component that drives anxiety disorders.

Interrogation of the neural circuits that represent normal and maladaptive safety evaluation traditionally rely on fear conditioning paradigms, in which a neutral cue is paired with an aversive stimulus to produce a conditioned response [7-9]. These fear-learning circuits offer important insight into the formation and retrieval of fear memory [10-13]. However, these paradigms are limited in their ability to address how an animal balances the safety and/or danger of a stimulus, a process that more closely recapitulates real-world decision-making.

Here we use conditioned taste aversion (CTA) to model the complexity of this decision-making process. CTA is a classical conditioning paradigm which pairs a palatable substance with an aversive visceral experience, such as the emetic agent lithium chloride (LiCl), producing an aversion to the substance at subsequent exposure. Unlike many other classical conditioning paradigms, in which an aversive stimulus is paired with a neutral cue, the use of a palatable substance in CTA requires the animal to process the affective valence of the reward versus the aversion. This provides an opportunity to understand neural processes by which this decision-making occurs.

We used several modifications of typical approaches to better capture the ability of this task to assess affective valence. We used a nutritive substance (chocolate-flavored Boost®) in non-food deprived animals, which more closely represents CTA outside the confines of a laboratory [14]. While current studies of CTA focus primarily on performance of the task, this study aimed to better capture both the emotional valence and the neural mechanisms that respond to each step of the CTA process. We accomplished this goal by evaluating both behavior and c-fos responses to the appetitive stimulus, (chocolate Boost®), the aversive stimulus (LiCl), the performance of the CTA task in the control condition in which animals expect only the appetitive stimulus, and the performance of the task when Boost® was paired with LiCl. By including the

acute conditions, as well as the appetitive task, we were better able to capture previously underappreciated neural activation in brain regions relevant to the CTA paradigm.

We also included females in the present study. Animal models of neuropsychiatric disorders rarely use female subjects. Indeed, only 2% of CTA studies utilize female rodents [15]. Females are disproportionately affected by anxiety-related disorders, including post-traumatic stress disorder, obsessive-compulsive disorder, generalized anxiety, and phobias [5, 16-20]. Despite these known sex differences, female rodents are critically understudied in animal models of neuropsychiatric disorders. We aimed to determine if any sex differences in the CTA response reflected sex differences in responses to the aversive stimulus, palatable stimulus, or recall and performance of the task.

In this study, we characterized behavioral and neural responses to CTA in males and females. Behavioral observations of the aversive stimulus included pica, ptosis, and lying on belly (LOB) as an indication of nausea in LiCl-treated rats. We also quantitated ultrasonic vocalizations (USVs) as a measure of emotional valence on task day. Rats engage in two distinct call patterns: short 55-kHz (which range from 30-70 kHz and are associated with positive emotional valence), and longer 22-kHz calls (associated with negative emotional valence). They will emit 55 kHz calls during behaviors such as play, mating, social approach, and in anticipation of a reward [21-24]. We therefore hypothesized that animals would engage in these calls on task day when anticipating Boost® that was not devalued with LiCl injection. Conversely, 22-kHz calls are exhibited in contexts of social withdrawal (e.g. after mating [25]) and to warn the colony of potential predators [26-28]. Female rats are more likely to exhibit 22-kHz calls when in the presence of cat urine [29]. We hypothesized that rats would engage in these calls on task day when anticipating Boost® that was previously paired with LiCl.

We used the immediate early gene c-fos as a marker of neuronal activity at each stage of CTA: in response to the acute rewarding and aversive stimuli, and during CTA expression. We analyzed 10 brain regions known to be associated with the response to reward and/or LiCl. We selected more rostral regions that are associated with the decision-making process in CTA, rather than the more caudal regions which serve to transmit signals that are interpreted by higher-order areas of the brain. Our c-fos data prompted us to more closely examine specific cell types in the amgydala using RNAScope, a novel mRNA *in situ* hybridization technique.

The following studies showed that animals taken through CTA assign valence to their experimental condition, characterized behaviors that correlate with CTA expression, and identified distinct neuronal cell types that are responsive to rewarding and aversive stimuli.

Materials and Methods

Animals

Male and female Sprague Dawley rats [post-natal day (PN) 60, Charles River Laboratories, Raleigh, NC] were received 1 week before behavior tests. Rats were housed by sex in ventilated plastic cages with ad libitum PM5001 Rodent Chow and water on a 7 am-7pm light/dark cycle. Females were not selected based on estrous cycle stage but cycle was determined by lavage on the final day of behavioral testing. A single lavage was performed as repeated lavages can establish place preference for the location of lavage and suppress behavioral responses to other reinforcers [30].

Conditioned Taste Aversion Protocol

Animals were conditioned to ip injections with NaCl for 3-4 consecutive days prior to experimental procedures. The conditioned taste aversion protocol (Fig. 1a) is conducted over 3 days. On day 0, animals are water deprived overnight and then conditioned to test cage. Water bottles are weighed and placed on cages with spouts pointing way. They are allowed to habituate

to their new environment for 30 minutes, after which they are given access to water for 60 minutes in the test cage, to condition them to drinking from new bottles. Water bottles are weighed again, and consumption is measured.

On day 1, non-water deprived animals were placed in the same test cage and allowed to habituate for 30 minutes. They were then offered Boost® and allowed to drink for 60 minutes. Boost® consumption was measured. If animals drank a minimum of 1 mL of Boost®, they were then injected with LiCl (38 mg/kg, 0.15M concentration) or the equivalent volume of NaCl. They were observed in the test cage for 60 minutes for LiCl behaviors.

On day 1, animals were returned to the test cage and allowed to habituate for 30 minutes. They were then offered Boost® and allowed to drink for 60 minutes. Boost® consumption was measured. Animals were then given a dose of LiCl (38 mg/kg) or the equivalent of NaCl. This dose was selected in order to achieve robust c-fos response for interrogation of the neural network. They were observed for 60 minutes for LiCl-related behaviors. They were then returned to their home cages and allowed to drink and eat ad libitum.

On day 2, non-water deprived animals were returned to the same test cage and allowed to habituate for 30 minutes. They were then given a 2-bottle test, offering either Boost® or water for 60 minutes. Boost® and water consumption were measured and ratio of day 2/day 1 (D2/D1) consumption was calculated. After 60 minutes of re-exposure to the Boost®, animals were perfused with formalin and brains collected for c-fos analysis. Brains were post-fixed in formalin overnight at 4C, then cryoprotected in 30% sucrose solution (30% sucrose, 70% PBS) for 3-4 days. Brains were flash-frozen in brain molds in ethanol/dry ice bath, in 2:1 30% sucrose/tissue freezing media solution. They were stored at -80°C until ready to cut for immunohistochemistry or RNAScope.

Animals whose brains were analyzed for the acute stimulus were taken through day 0 as described. On day 1, they were allowed to habituate for 30 minutes in the test cage before receiving a dose of LiCl or the equivalent of NaCl or were allowed to drink Boost® for one hour, after which they were perfused, and brains collected for c-fos analysis (see Table 1).

Table 1

Acute Stimulus	Conditioned Stimulus
Boost®	Boost® paired with NaCl
LiCl (38 mg/kg)	Boost® paired with LiCl
NaCl (control)	Context only control

Behavior Scoring

Three behaviors were time-sampled as a behavioral estimation of nausea [pica, ptosis, lying on belly (LOB)]. As rats do not vomit, they will often attempt to dilute an ingested toxin through pica, the consumption of non-food substances. LiCl behaviors were assessed using a modified version of published measures: Pica and LOB are well known LiCl behaviors [31-33], and ptosis was included following the observation that it occurred more frequently in LiCl-treated animals. Animals are observed for two consecutive 15-second blocks (30 seconds total) every 5 minutes for 1 hour. Animals were given a score of 0-3 as described in table 2.

Table 2.

Behavior	Score = 0	Score = 1	Score = 2	Score = 3
Pica	No pica	Some pica while still moving around	Pica in a crouched position, but still moved or looked around	Remained crouched, engaged in pica for most of 15-second interval
LOB	No LOB	Some LOB, still in crouched position	Some LOB, more flattened posture	Entirely flattened posture
Ptosis	No ptosis	Ptosis for some of interval (<5 seconds)	Ptosis for most of interval (>5 seconds, <10 seconds)	Ptosis for entire interval

Scoring of Ultrasonic Vocalizations

In a within-groups design, rats were split by sex, treatment [NaCl + H_2O (Control), NaCl + Boost®, and LiCl + Boost®] on Day 2 of CTA experiment (Test day). USVs were recorded with an Echo Meter Touch 2, from *Wildlife Acoustics*, an accessory that attaches to a lightning port on the Apple iPhone. The Echo Meter Touch Bat Detector iOS app (v 2.7.20) was used to record the vocalizations. Recordings were uploaded from the smartphone to a secure online server. Manual scoring of the USVs during the 10-minute recording period was done using the visualization software from *Wildlife Acoustics*, Kaleidoscope Pro Analysis. Scoring was performed by a blinded observer in batches of 1-minute intervals. The scorer determined how often the rat engaged in >10 55 kHz vocalizations, or >1 22 kHz vocalizations in each interval. Scores were calculated as a percentage of the 10 1-minute blocks in which the vocalizations occurred.

Immunohistochemistry

Brains were cut on a Leica CM3050S cryostat at 30 uM and stored in 1:1 TBS/glycerol solution at -20℃ until ready to stain. To stain, slices were first washed in 0.2% Triton-X in TBS solution 3 times. Slices were then blocked for 1 hour in solution of 0.3% Triton-X and 5% normal goat serum (NGS) in PBS. Slices were stained overnight in 0.3% Triton-X, 5% NGS, and 1:20,000 anti-c-fos antibody (Abcam - ab190289) with gentle shaking at 4℃. Slices were then washed 3 times in solution of 0.3% Triton-X and 5% NGS in PBS (10 minutes first wash, 30 minutes second wash, 40 minutes third wash). Slices were then stained with secondary antibody (AlexaFluor 488, Invitrogen – A-11034) in 0.3% Triton-X, 5% NGS, and 1:200 secondary antibody for 2 hours. Slices were then washed in 1X PBS (10 minutes first wash, 30 minutes second wash, and 40 minutes third wash), with DAPI (R&D Systems 5748) added at a dilution of 1:10,000 for last 10 minutes of second wash. Slices were mounted on VWR Superfrost® Plus microscope slides with a drop of Vecta-Shield anti-fade medium (Vector Laboratories 101098-042).

Vaginal lavage and estrus cycle analysis

A vaginal lavage smear was performed on all female test subjects prior to transcardial perfusion. Vaginal cytology review was performed to determine the phase of estrous cycle. The method of estrous cycle analysis as previously described [30]. Animals were grouped into high estradiol states (proestrus/estrus) and low estradiol states (metestrus/diestrus).

RNAScope

Brains were cut at 14 uM on a Leica CM3050S cryostat and placed on a Superfrost® Plus Micro Slide. 3 consecutive slices were placed on each slide. Slides were collected ~90 uM apart. Slides were desiccated at -20℃ for 20 minutes, then placed into -80℃ with desiccants for long term storage. Slides were stained within 1 month of collection for optimal signal.

When ready to stain, slides were placed in 1X PBS for 5 minutes to remove OTC then placed in the HybEZ™ oven for 30 minutes at 60°C. They were then transferred to cold PFA for 15 minutes at 4°C, after which they were taken through the RNAScope procedure as outlined by ACDBio, using Protease IV. Probes were for c-fos (403591), Drd1 (317031-C2), and Drd2 (315641-C3). Secondary antibodies were obtained from Akoya (OPAL 690, OPAL 520, and OPAL 570) and diluted to a concentration of 1:750.

Imaging and analysis

Slides were imaged with an Axio Imager upright microscope at 20x (16 z-stacks 2 uM apart) for IHC or 20x (8 z-stacks 1 uM apart) for RNAScope. Images were z-projected for max intensity using FIJI (ImageJ) software and c-fos+ neurons were manually counted. 2-6 slices per region were imaged and counted per animal. Counts were averaged per animal.

Drugs

LiCl was purchased from Sigma Aldrich (203637). LiCl was dissolved in distilled sterile water to a concentration of 0.15M. Sterile isotonic saline (NaCl, 0.9%; 0.15M) was used as a control injection.

Statistics

All results were analyzed by ANOVA with posthoc Fisher's exact test corrected for multiple comparisons using the statistical package NCSS. Behavioral results were analyzed by 2-way ANOVA (sex x treatment). C-fos responses were analyzed by 3-way repeated measures ANOVA (sex and condition as between measures and brain area as repeated measure). All conditions yielded area as a main effect, and second level ANOVAs were run for each brain area independently. N for all behavior and c-fos experiments was estimated by power analysis and

estimated variance based on previous experience with CTA behavior and c-fos analysis of multiple brain regions (refs).

Results

Behavioral responses during conditioned taste aversion

The CTA experimental design is described in Figure 1a. To first characterize CTA expression, a dose-response curve was established in both sexes. Animals received either NaCl (0.15M) or LiCl (19 mg/kg, 38 mg/kg, or 80 mg/kg) (Figure 1b). LiCl-induced behavior was measured as described in Table 1. Statistics show a main effect of sex [$F_{(1,111)}$=6.05, p=0.015] and a main effect of treatment [$F_{(3,111)}$=29.31, p< 0.001] but no interaction of treatment x sex. We found that males inhibited Boost® intake more than females after pairing with the low dose of 19 mg/kg LiCl.

Female rats drank more Boost® than males at initial exposure when adjusting for body weight, with a main effect of sex [$F_{(1,146)}$=21.91, p< 0.001, (Fig. 1c)]. Both male and female rats generally increased the amount of a novel palatable substance they will consume at subsequent exposure. This ratio of consumption was similar in males and females despite the propensity for females to drink more on first exposure.

To determine whether estrous cycle might contribute to CTA behavior, cycle was determined by vaginal lavage on task day. Cycle stage did not have a significant effect on drinking behavior to the novel sweet tastant (Fig. 1d), nor did it influence CTA expression on task day (Fig. 1e).

Figure 1. (A) Protocol for CTA paradigm. (B) Dose response curve to LiCl (n=7-23). (C) Boost® consumption on first exposure by sex (n=48-100). (D) Estrous cycle and day 1 Boost® consumption (n=8-9). (E) Estrous cycle and CTA expression on task day (n=10-14). All data expressed as mean ± SEM.

LiCl-induced behaviors

Lying on belly (LOB) was the only nausea behavior measured in this study that correlated with CTA expression, Pearson correlation = -0.3701 and p < 0.0434 (Fig. 2b, c). Pica

was measured after a range of doses of LiCl: 0 mg/kg (NaCl), 19, 38, and 80 mg/kg. Both male and female rats engaged in pica equally across doses of LiCl. 2-way ANOVA showed an effect of treatment [$F_{(3,158)}=20.79$, $p< 0.001$] and no effect of sex or interaction of treatment x sex (Fig. 2a). Pica events at 80 mg/kg were higher than 38 and 19 mg/kg. Male and female rats also exhibited ptosis comparably after LiCl injection (Fig. 2d). Neither of these behaviors correlated with CTA behavior as measured by the ratio of Boost® consumption on day 2 as compared to day 1.

Figure 2. (A) Pica events time-sampled over 1 hour (n=26-44) after 38 mg/kg LiCl or NaCl. Data shown as mean ± SEM. (B) Pica events after increasing doses of LiCl (0, 19, 38, and 80 mg/kg) in male and female rats (n=6-45) Data shown as mean ± SEM. *=different from NaCl (0 mg/kg). #=different from 19 mg/kg and 38 mg/kg. (C) LOB was measured for 1 hour after 38 mg/kg LiCl or NaCl (n=18-33). Data shown as mean ± SEM. (D) LOB correlation with D2/D1 ratio (E) Ptosis events measured over 1 hour after 38 mg/kg LiCl or NaCl (n=18-33).

Ultrasonic vocalizations during conditioned taste aversion task

Rats were taken through the CTA protocol as described. As vocalizations typically occur

in anticipation of a reward, we recorded USVs during the first 10 minutes of the habituation

process on Day 2, before offering the 2-bottle test.

We found that male rats reliably vocalized at 55 kHz (Fig. 3b) in anticipation of Boost®

if it had been paired with NaCl [$F_{(4.52, 1)}$ $p < 0.038$ effect of treatment, Fig. 3a]. Female rat

vocalizations were substantially influenced by estrous cycle, with high-estradiol states associated

with an increased likelihood of engaging in 55 kHz calls, and low-estradiol states decreasing the

frequency of 55 kHz calls, regardless of experimental condition [$F_{(1,27)} = 6.35$, $p < 0.018$, Fig.

3f]. By task, females showed an effect of estrous cycle with no effect of treatment [$F_{(1,24)} = 7.38$,

$p < 0.012$, Fig. 3e]. One-way ANOVA for each low- and high-estradiol states individually

showed females in high-estradiol states increased 55 kHz vocalizations in the Boost® task

compared to LiCl task [$F_{(1,12)} = 5.20$, $p < 0.042$, Fig. 3e]. There was no effect of treatment in the

low-estradiol group.

There was a significant correlation of 22 kHz calls (Fig. 3d) in the LiCl-paired

environment by 2-way ANOVA, with a main effect treatment [$F_{(1,59)} = 9.48$, $p < 0.0032$, Fig. 3c].

Splitting out by sex due to pre-planned contrast showed that females significantly increased 22

kHz calls in the LiCl-paired environment [$F_{(1,34)} = 9.17$, $p < 0.005$], while males were not

significant.

Figure 3. USVs were recorded for first 10 minutes during CTA task. (A) 55 kHz calls in male and female rats during CTA expression (n=11-19). (B) 22 kHz warning calls during CTA expression in male and female rats (n=12-19). (C) Representative 55 kHz call (D) Representative 22 kHz call (E) 55 kHz vocalizations in females by cycle and condition (n=6-8) (F) 55 kHz calls in females by cycle only

The c-fos response to acute and conditioned stimuli

We evaluated c-fos expression to better understand the neural circuit involved in the development and expression of CTA. c-fos analysis was performed across several brain regions associated with the response to rewarding and aversive stimuli (Table 2).

Table 3

	Acute NaCl		Acute Boost®		Acute LiCl		Control Task		Boost® task		LiCl Task	
	F	M	F	M	F	M	F	M	F	M	F	M
vmPFC	244±37	229±72	244±20	239±33	266±41	272±51	246±37	241±32	268±52	212±31	249±44	221±43
aIC	183±32	187±40	198±20	215±22	212±50	206±27	232±32	193±14	223±31	176±26	227±21	174±24
gIC_5/6	100±38	87±44	197±29	178±77	206±44	206±70	81±45	86±26	162±53	145±23	144±55	99±72
gIC_5/6_L4	134±66	85±28	262±23	253±82	255±23	267±89	148±82	98±64	225±69	184±27	121±45	79±43
NAc_C	91±30	98±35	133±7	98±20	100±62	109±29	110±1	76±18	123±26	99±34	101±21	64±29
NAc_S	117±39	123±36	125±19	107±33	171±78	141±68	120±26	90±17	153±38	113±41	124±52	118±52
SON	7±5	14±8	11±9	10±12	112±34	112±3	10±10	2±1	21±6	27±27	5±2	45±70
BLA	34±8	32±8	56±16	53±8	67±25	58±15	52±19	24±5	61±10	42±3	52±6	41±15
CeA	44±22	42±26	133±40	108±33	191±73	199±58	30±7	42±60	142±19	115±60	62±48	48±28
PVN	113±51	103±26	122±51	143±29	230±102	226±96	95±36	107±28	91±20	74±45	141±39	134±70

c-fos immunoreactivity during CTA acquisition and expression

We performed 3-way repeated measures ANOVA (sex x treatment, area as a repeated measure) on all experimental conditions, which revealed a main effect of sex [$F_{(1,62)}=11.46$, p=<0.001], a main effect of treatment [$F_{(5,45)}=5.29$, p=0.025], a main effect of area [$F_{(9,62)}=207.52$, p <0.001], and an interaction of treatment x area [$F_{(45,62)}=7.75$, p< 0.001].

We then performed 3-way repeated measures ANOVA for the acute and conditioned stimuli separately. For the acute condition, 3-way repeated measures ANOVA (treatment x sex, area as a repeated measure) revealed a main effect of treatment [$F_{(2,31)}=21.27$, p < 0.001] and area [$F_{(9,31)}=84.95$, p < 0.001], and an interaction of treatment x area [$F_{(9,31)}= 8.46$, p <

0.005], with no effect of sex for the acute stimuli. Acute Boost® and acute LiCl groups both differ from NaCl control. We conducted lower level 2-way ANOVA (sex x Rx) for areas that are implicated functionally in CTA and/or showed statistically relevant findings.

As discussed in the introduction, the central nucleus of the amygdala (CeA) is known to responds to aversive visceral stimuli such as LiCl and, more recently, rewarding stimuli. We found the CeA increased both to Boost® and LiCl [$F_{(2,30)}$= 31.39, p< 0.001 effect of treatment] similarly in male and female rats (Fig. 4a). The basolateral amygdala (BLA), known for its role in assessing the valence of a stimulus, showed an effect of treatment [$F_{(2,17)}$= 9.06, p< 0.0021], with no effect of sex (Fig. 4b). Post-hoc analysis showed both Boost® and LiCl groups were increased compared to NaCl control.

The granular insula (gIC_5/6) receives both taste and visceral inputs [34-41], and is divided into several layers. We considered layer IV (gIC_5/6_4) and layers V/VI (gIC_5/6_5/6). c-Fos in gIC_5/6_5/6 increased with both Boost® and LiCl (Fig. 4e), showing a significant effect of treatment [gIC_5/6; $F_{(2,29)}$=14.52, p< 0.001].   c-Fos in gIC_4 likewise increased in both Boost® and LiCl groups compared to NaCl control (Fig. 4f). The gIC_4 showed an effect of treatment [$F_{(2,17)}$=15.62, p> 0.001].

The supraoptic nucleus (SON) responds to acute LiCl [42, 43]. It is the primary site of oxytocin production, which is known to have emetic and anorectic properties [44-47]. The c-fos in SON response increased markedly in response to LiCl [$F_{(2,16)}$=54.36, p< 0.001 effect of treatment], with no response to Boost® (Fig. 4j). c-Fos in the paraventricular nucleus (PVN), which contains receptors for oxytocin and is the primary location for release of stress-related peptides such as corticotropin-releasing hormone and vasopressin [48-51], likewise increased specifically in response to the aversive stimulus [$F_{(2,27)}$=8.44, p< 0.001, Fig. 4k].

Interoceptive signals and contextual stimuli converge in the ventromedial frontal cortex (vmPFC) and agranular insula (aIC) [38, 52-58]. However, no significant effects were found in the vmPFC or aIC in any condition, and so these will not be discussed further.

The nucleus accumbens responds to rewarding stimuli, typically in behavioral paradigms that utilize deprivation. The accumbens shell is more responsive to rewarding stimuli, while the accumbens core responds to conditioning of rewarding stimuli [59-65]. No significant effects were found in either the accumbens core (NAc_C) or shell (NAc_S) during the acute stimuli.

c-fos immunoreactivity during CTA expression

3-way repeated measures ANOVA (sex x treatment x area) of the CTA task conditions revealed a main effect of sex [$F(7.42,1)$, $p< 0.011$] and area [$F(138.09,9)$, $p< 0.001$] and an interaction of treatment x area [$F(6.73,18)$, $p< 0.001$]. Females were more responsive than males overall. We then ran second order ANOVAs of treatment x sex in individual areas, and treatment x area in each sex.

C-fos in the CeA increased in response to the Boost® task. It showed an effect of treatment [$F(2,29)=24.95$, $p< 0.001$] and barely missed an effect of sex [$F(1,29)=3.96$, $p< 0.056$]. The Boost® task differed from the control task, both globally and in each sex individually. c-Fos in the BLA showed an effect of sex [$F(1,18)=17.34$, $p < 0.001$]. This was driven largely by high baseline c-fos immunoreactivity in the control task for females. Lower-level ANOVA by sex showed a significant effect of condition in males only, with both Boost® task and LiCl task differing from control.

The gIC_5/6 showed an effect of treatment [$F(2,31)=5.29$, $p< 0.011$], with the Boost® task differing from the control task. This difference is driven by the female rats, which show increases in c-fos response in both the Boost® task and LiCl task compared to control (Fig. 4e).

c-fos in the gIC_4 showed an effect of treatment [F(2,17)=4.39, p< 0.014], and no effect of sex (Fig. 4f). The Boost® task showed increased c-fos expression compared to the LiCl task globally. The females showed a statistically significant increase in c-fos expression in the Boost® task compared to LiCl task conditions.

Females had higher c-fos expression in the NAc_C than males. It showed an effect of sex [F(1,31)=12.31, p< 0.001] but no effect of treatment (data not shown). Females had higher c-fos expression than males, both globally and in the Boost® task specifically. c-fos in the NAc_S showed an effect of sex [n=4, F(1,18)=14.52, p< 0.001], with no effect of treatment (data not shown).

Neither the PVN nor the SON responded during the task conditions.

A

Central Nucleus of the Amygdala

B

Basolateral Amygdala

C

D <u>Central Nucleus of the Amygdala</u>

E

Granular Insula

c-fos+ cells

NaCl Boost(R) LiCl NaCl Boost(R) LiCl Control Task Boost(R) Task LiCl Task Control Task Boost(R) Task LiCl Task

F

Gustatory Cortex

c-fos+ cells

NaCl Boost(R) LiCl NaCl Boost(R) LiCl Control Task Boost(R) Task LiCl Task Control Task Boost(R) Task LiCl Task

G

H Granular Insula

Control Boost® LiCl

Acute

CTA Task

Gustatory Cortex

Control Boost® LiCl

Acute

J

Supraoptic Nucleus

K

Paraventricular Nucleus

L

Figure 4. (A) c-fos response in the CeA across conditions (B) c-fos response in the BLA across conditions (C) Atlas image of CeA and BLA. (D) Representative images. (E) c-fos response in the gIC_5/6 across conditions. (F) c-fos response in the GIC_4 across conditions (G) Atlas image of gIC_5/6 and GIC_4. (H) Representative images. (J) c-fos expression in the PVN across conditions (K) c-fos expression in the SON across conditions. (L) Atlas image of SON and PVN. (M) Representative images.

Responses of dopamine D1- and D2-receptor expressing cells in the amygdala to rewarding and aversive stimuli

The role of the amygdala in the response to acute aversive stimuli such as LiCl is well-established [42, 43, 61, 66-69]. A thorough study by Tonegawa, et. al., examined the expression patterns and responses to both fear and reward of multiple neuronal cell types in the CeA and BLA [70], laying an important groundwork for further interrogation into the function of specific neuronal cell types. The BLA is responsible for assessing the salience of a signal, whether it be rewarding or aversive [70-76]. BLA outputs to the CeA promote defensive and appetitive behaviors [70].

Dopamine receptors are present throughout the amygdala, but there is little known about their reactivity to reinforcing or aversive stimuli. We used RNAscope to visualize D1 and D2 dopamine receptors (Drd1 and Drd2) throughout the amygdala. We found that Drd1+ and Drd2+ cells occupy largely distinct anatomical locations within the amygdala. The CeA is further divided into three subnuclei: the central medial (CeM), central lateral (CeL), and the capsular (CeC). Consistent with previous studies, we found that the CeC and CeL were composed primarily of Drd+ cells, while the CeM contained primarily Drd1+ cells (Fig 5a). The BLA, in contrast to the CeA, represents a nearly homogenous population of D1-receptor expressing cells, with few D2-receptor expressing cells (Fig. 5a).

Using the novel mRNA *in situ* hybridization method RNAScope, the co-expression of c-fos and D1 or D2 receptors was characterized in each of the acute conditions, as well as during CTA expression, in the BLA and CeA. A representative image of the amygdala is shown in Fig. 5a.

A

B

BLA

Blue = DAPI
Green = c-fos
Red = drd1
Magenta = drd2

C <u>CeA</u>

c-fos+drd1   c-fos+drd2   Merge

Acute LiCl

Blue = DAPI
Green = c-fos
Red = drd1
Magenta = drd2

CeM
CeL

Acute Boost®

CeM
CeL

Acute NaCl

CeM
CeL

Figure 5. (A) Distribution of Drd1+ and Drd2+ cells across the amygdala. BLA = basolateral amygdala. CeA = central nucleus of the amygdala. ASTA = amygdalostriatal transition area. (B) Representative images of RNAScope in acute LiCl and acute Boost® conditions in the CeA. (C) Representative images of RNAScope in acute LiCl and Boost® conditions in the BLA. (D) RNAScope analysis of co-expression of c-fos and Drd1+/Drd2+ cells in the basolateral amygdala across conditions. (E) RNAScope analysis of co-expression of c-fos and Drd1+/Drd2+ cells in the central nucleus of the amygdala across conditions.

Responses of D1-receptor expressing cells in the BLA to acute and conditioned rewarding and aversive stimuli

Representative images for the BLA are shown in Fig. 5b and results in Fig. 5d. The BLA expressed c-fos in response to both rewarding and aversive stimuli, and during CTA expression in male rats only. The Drd2 cells, which represented a small fraction of total cells, did not respond in any stimulus-specific pattern.

We performed 2-way ANOVA (sex x treatment) for total c-fos mRNA. In the acute conditions, total c-fos mRNA expression in the BLA showed a main effect of treatment [F(2,17)=32.00, p< 0.001] and an effect of treatment by sex [F(2,17)=5.73, p< 0.013] for the acute stimuli. Posthoc analysis showed Boost® and LiCl conditions differed from NaCl control, globally and for both sexes individually. We next considered the co-expression of c-fos+/Drd1+ cells. 2-way ANOVA (sex x treatment) revealed an effect of treatment [F(2,22)=4.03, p< 0.005] and an interaction of treatment x sex [F(2,22)=4.03, p< 0.037]. Posthoc analyses revealed a difference between the acute Boost® and LiCl conditions compared to NaCl control, both globally and by each sex individually.

For the task condition, 2-way ANOVA revealed a main effect of treatment [F(2,23)=5.66, p< 0.001] and an interaction of sex x treatment [F(2,23)=9.46, p< 0.0016]. There was no main effect of sex. Posthoc analysis revealed both the Boost® and LiCl tasks differed from the control task. By sex, males in the Boost® and LiCl tasks differed from their respective controls. Females did not differ from controls, which had high levels of baseline c-fos.

The same pattern emerged when considering c-fos+/Drd1+ co-expressing cells. There was an effect of treatment [F(2,23)=8.59, p< 0.0024] and an interaction of sex x treatment [F(2,23)=9.64, p 0.0014]. There was no main effect of sex. As with total c-fos, the Boost® and LiCl task conditions showed greater c-fos mRNA expression than the control. As with total c-fos only males differed in the Boost® and LiCl task conditions from their respective controls.

The CeA is responsive to rewarding and aversive stimuli in a cell-specific manner

Representative images for the CeA are shown in Fig. 5c and results in Fig. 5e. Unlike the BLA, the CeA responded to rewarding and aversive stimuli in a cell-specific manner. In the acute condition, ANOVA revealed an effect of treatment [F(2,17)=30.32 p < 0.001] for total c-

fos+ cells. There was no main effect of sex or treatment x sex interaction. Posthoc analysis revealed the NaCl differed from both the Boost® and LiCl conditions.

When considering c-fos+/Drd1+ co-expressing cells, there was an effect of treatment [F(2,17)=100.22, p=< 0.001]. There was no main effect of sex or interaction of sex x treatment. The acute Boost® condition differed from the acute NaCl condition, with robust expression of double positive cells (c- fos and Drd1). There was no difference between the acute LiCl and acute NaCl condition.

The opposite association of c-fos and condition was found for Drd2 expressing cells. By ANOVA there was an effect of treatment [F(2,17)=38.63, p < 0.001]. There was no effect of sex or interaction of treatment x sex. The acute LiCl condition showed significant expression of c-fos in Drd2+ cells, but not the acute Boost® condition.

In the task condition, there was an effect of treatment [F(2,18)=40.24, p=<0.001], with no effect of sex or treatment x sex interaction. There was significant mRNA expression of c-fos in the Boost® task, but not the LiCl task condition. When considering c-fos+/Drd1+ cells, there was an effect of treatment [F(2,18)=53.15, p < 0.001]. As with the acute condition, there was significant overlap between c-fos+ and Drd1+ cells.

Altogether, these data showed that Drd1+ cells are differentially responsive to the rewarding stimulus in the CeA, while Drd2+ cells are differentially responsive to the aversive stimulus in the CeA. Following treatment with acute LiCl, c-fos response showed both anatomical and neuronal cell type specificity, with activation occurring primarily in the CeL and CeC, and co-expressing with Drd2-expressing cells. Conversely, Boost® produced a c-fos response in the CeM and CeL and co-expressed with Drd1-expressing cells. Increases in c-fos and c-fos/drd1 co-expression were observed in both acute and conditioned Boost®.

Discussion

Conditioned taste aversion is most often characterized as a task which reflects avoidance/aversion. By assessing behavioral and neural responses to both the reinforcing and aversive stimulus, the present findings reveal how animals balance the competing signals of safety or danger to learn and perform this task. Interrogation of the neural network revealed that stimuli activate circuits in a stimulus- and cell type-specific manner, suggesting that opposing stimuli activate parallel, but distinct, neural pathways.

Behavioral responses to rewarding and aversive stimuli

CTA is an essential evolutionary behavior to prevent further consumption of foods that may be toxic and should be avoided. Critical to the CTA paradigm is the devaluation of the hedonic stimulus when it is paired with an aversive visceral stimulus. For example, delivering only an aversive stimulus like footshock can reduce CS consumption without changing palatability, while aversive *visceral* experience reduces both consumption and palatability, as measured by orofacial responses to the CS [77, 78]. Here, we use USVs to show that the behavior is more complex than simply showing aversion. Animals are both attaching a hedonic value to the rewarding stimulus and signaling the danger of the devalued stimulus.

This is the first study to comprehensively evaluate nausea behaviors in both male and female rats in CTA. We found that male and female rats showed similar nausea behaviors after LiCl administration. Of the three acute nausea behaviors we observed, only lying on belly correlated with CTA expression, suggesting this behavior may be an expression of more severe nausea than pica. Further studies in CTA and other nausea-relevant paradigms may prove useful in characterizing how this behavior correlates to nausea.

USVs proved to be a useful tool to assess both positive (55 kHz) and negative (22 kHz) response to stimuli during CTA expression. Animals expressed 55 kHz calls in anticipation of Boost® and inhibited 55 kHz calls, while increasing 22 kHz calls when the Boost® was devalued with LiCl. The expression of USVs offers insight into the valuation animals are assigning to the Boost® and to the LiCl. Increases in 55 kHz in the context previously paired with Boost® suggests animals are assigning a positive valuation to the hedonic stimulus. The converse inhibition of 55 kHz USVs reinforces the notion that USVs can be correlated with an expected outcome. Interpretation of female 55 kHz calls was complicated by a marked effect of estrous cycle, as vocalization patterns reflected high-estradiol (more vocal) and low-estradiol (less vocal) states. Conversely, females were more likely to engage in 22 kHz calls than males. It is unclear whether this result suggests that females are more likely to attach a negative valuation to LiCl-paired Boost® or that they are more likely, in general, to elicit warning calls to alert the colony of nearby dangers. The latter hypothesis is supported by studies showing females are more likely than males to increase 22 kHz calls in the presence of a predator [79], especially considering their comparable sensitivity to CTA. USVs provide additional insight not only into the animal's valuation of the stimulus, but to how they communicate this valuation to others in the colony.

The c-fos response to rewarding and aversive stimuli

CTA is a unique form of learning in that the stimuli do not need the same temporal proximity as other operant learning and behavioral conditioning paradigms [80, 81]. The nature of digestion is such that visceral malaise may occur minutes to hours after the ingestion of a toxic substance. Hedonic and aversive inputs follow similar pathways: caudal regions, such as the nucleus of the solitary tract and parabrachial nucleus, increase c-fos in response to both LiCl

and sucrose [82, 83]. These circuits then converge in brain regions like the amygdala to assess valence. In other words, we hypothesize that proximity may be found within the circuit itself, reducing the necessity for events to overlap in time for learning to occur.

This schema was borne out in our data. The neural circuit underlying the response to the acute stimuli overlapped in key brain regions, including the BLA, the CeA, the gIC_5/6, and the gIC_4. In contrast to brain areas that respond to both acute stimuli, the PVN and SON uniquely responded to the aversive stimuli. Sampling of more caudal areas may prove useful in delineating the cell type-specific circuit in future studies. Regions such as the NTS [84-91] and PBN [67, 68, 83, 85, 91, 92] which were omitted in the present study to focus on rostral brain areas are known to respond to both aversive (emetic) and rewarding stimuli, and future use of cell-type specific probes will allow more complete dissection of this circuit .

Our data show that the gustatory cortex (GC, which processes taste) overlaps significantly with the visceral insular cortex (VIC, which processes visceral stimuli) show significant overlap in c-fos expression [40, 93]. The GC reacts to novel taste and lesions are known to cause a decrease in neophobia [36, 94-96]. Additionally, lesions to the GC can attenuate CTA on first trial, a deficit that disappears with repeated CS-US pairings [37]. These studies raise the question whether the loss of neophobia acts as a latent inhibition, reducing CTA by making the CS appear familiar [97]. An alternative explanation from our studies is the possibility that lesions of the GC likewise destroy cells that are part of the VIC, and thereby reduce the nausea effects of LiCl. Behavioral studies of nausea behavior would help to clarify this question. Additionally, our findings of cell-specific c-fos responses in the CeA suggested that future studies which interrogate cell-specific responses in areas like the highly heterogenous

insular cortex to rewarding and aversive stimuli in a cell-specific manner will provide new insight into integration of affective valence with motivated behavior.

CTA is often characterized as analogous to fear learning. However, CTA involves the convergence of two stimuli of opposing valence, resulting in complex decision-making rather than a fear-induced defensive response. The RNAScope data showing that the response to rewarding and aversive stimuli show cell- and anatomical specificity in the CeA suggests that other regions may show similar stimulus-specificity when specific cell populations are evaluated. How these two neural networks converge to produce the memory of the experience, and the subsequent decision-making process that guides the animal's behavior, is an important area of future study.

We were surprised to see there was no activation of the nucleus accumbens shell after acute Boost®, or the core during expression of the task. One explanation is that most studies showing such activation increased the reward of the hedonic substance by water and/or food deprivation during CTA acquisition, while we did our studies in animals that were not deprived. Studies show deprived animals express higher levels of c-fos in these key brain areas compared to satiated animals [62, 64].

We found no sex differences in the neural activation in response to the acute stimuli in any of 10 brain regions we assessed. This was a surprising result, as females drink more Boost® at first exposure and are relatively less sensitive to developing CTA. It is possible that our limited analysis did not sample a critical area that accounts for these differences. A more comprehensive study of the neural circuit that includes other brain regions associated with CTA, such as the bed nucleus of the stria terminalis, ventral tegmental area, or parabrachial nucleus may be of value.

The c-fos response during CTA expression

We used a novel experimental paradigm for studying the neural circuits engaged during the acute rewarding and aversive stimuli, as well as during the reinforcing task (expectation of Boost®) and CTA task (expectation of Boost® after pairing with LiCl). Neural activation during the Boost® task was similar to that of the acute Boost®, while activation during the LiCl task differed considerably from its acute stimulus. The neural circuit of animals that received Boost® that was not devalued with LiCl is reinforced with repeated offerings of the hedonic stimulus. We therefore saw a similar neural activation during expression of the task When this same stimuluswass devalued with LiCl, drinking behavior was reduced, and the circuit was nhibited.

During CTA expression, females broadly expressed higher levels of c-fos than males. By area, they were significantly more responsive in the gIC_5/6 and GIC_4 during the Boost® task. These data are consistent with their behavior, as they tend to drink more Boost® compared to males. Cell-type analysis would benefit our understanding of how females differentially process sweet taste, a critical area of study in the field of eating disorders [98-106]. This result is particularly interesting in light of their relative lack of USV expression in response to the rewarding stimulus. While males are more unequivocal about the "goodness" of Boost®, as evidenced by their consistent expression of 55 kHz USVs, females in low-estradiol states do not express 55 kHz USVs, despite drinking comparably.

Dopamine receptors in the amygdala respond to rewarding and aversive stimuli

Our studies showed that Drd1+ neurons were responsive to the rewarding stimulus, while Drd2+ cells were responsive to the aversive stimulus. This is the first time these cell types have been studied in the context of CTA. Tonegawa et. al. [70] showed that silencing Drd1+ neurons in the CeA inhibited feeding behavior in mice. However, other studies demonstrate that direct

blockade of Drd1 in the CeA did not affect feeding behavior, suggesting these neurons are being activated by something other than dopamine. Drd1+ neurons highly co-express with 3 other neuropeptides: tachykinin 2, neurotensin, and somatostatin. Although these neuronal cell types do not affect behavior when manipulated on their own, Tonegawa shows that these cell types may work collectively to coordinate feeding behavior.

Only about 30% of c-fos+ neurons in the CeA co-expressed with Drd1+ cells in response to the appetitive stimuli, suggesting additional cell types may also be important for the processing of the reward. A similar percentage of Drd2+ receptors were responsive to LiCl. A more thorough examination of the CeA in response to a food reward and aversive stimulus would be useful in delineating the cell types that process this stimulus.

Contrary to our findings in the CeA, Drd1+ neurons in the BLA responded to both rewarding and aversive stimuli, with limited input from Drd2+ neurons. These findings suggest that Drd1+ neurons may receive innervation from more caudal areas that respond specifically to these stimuli [107]. Additionally, the BLA is known to respond to rewarding or aversive stimuli in a location-specific manner, with the rostral BLA responding to aversive stimuli and the caudal BLA responding to rewarding stimuli. The dual responses observed in the present study may reflect the location at which we sampled the BLA, which was in the central zone where these two stimuli converge [108].

In summary these findings support our hypothesis that CTA is best characterized as a task in which animals must balance rewarding and aversive stimuli, and therefore is not simply a model of fear learning [109-113]. As such, it has significant relevance as a task which reveals how affective valence is integrated into decisions about behavior. The USV analysis showing inhibition of 55 kHz calls and increased 22 kHz calls in the LiCl-paired context suggests that

additional neural centers may be involved in assigning valence during the CTA task that are not captured by LiCl behaviors or our c-fos analysis. Our c-fos analysis of rostral brain regions revealed a significant role of the CeA, gustatory cortex, and granular insula in processing of the reinforcing and aversive stimuli, and increased global c-fos expression in females overall during the task. Further characterization of both the acute and conditioned circuits in males and females will help clarify how these stimuli are differentially processed between the sexes. Our data found a novel role for specific cell types in the CeA and BLA that have not previously been studied. These results suggest that similar cell-specific analysis in cortical areas will continue to reveal details of the neural circuit(s) engaged while animals learn this type of task.

References

1. Apergis-Schoute, A.M., et al., *Neural basis of impaired safety signaling in Obsessive Compulsive Disorder.* Proc Natl Acad Sci U S A, 2017. **114**(12): p. 3216-3221.
2. Padgaonkar, N.T., et al., *Neural correlates of emotional reactivity and regulation in youth with and without anxiety.* Depress Anxiety, 2021.
3. Stegmann, Y., et al., *Individual differences in human fear generalization-pattern identification and implications for anxiety disorders.* Transl Psychiatry, 2019. **9**(1): p. 307.
4. Via, E., et al., *Ventromedial prefrontal cortex activity and pathological worry in generalised anxiety disorder.* Br J Psychiatry, 2018. **213**(1): p. 437-443.
5. Young, L.J. and D.W. Pfaff, *Sex differences in neurological and psychiatric disorders.* Front Neuroendocrinol, 2014. **35**(3): p. 253-4.
6. Kessler, R.C., et al., *Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication.* Arch Gen Psychiatry, 2005. **62**(6): p. 593-602.
7. Delprato, D.J. and R.W. Thompson, *Effect of electroconvulsive shock on passive avoidance learning with high and low intensity footshock.* Psychol Rep, 1965. **17**(1): p. 209-10.
8. Dennis, M., *Sex-dependent and sex-independent neural control of reactivity to electric footshock in the rat.* Exp Neurol, 1972. **37**(2): p. 256-68.
9. Davis, M. and D.I. Astrachan, *Conditioned fear and startle magnitude: effects of different footshock or backshock intensities used in training.* J Exp Psychol Anim Behav Process, 1978. **4**(2): p. 95-103.
10. Parent, M.B., E. Avila, and J.L. McGaugh, *Footshock facilitates the expression of aversively motivated memory in rats given post-training amygdala basolateral complex lesions.* Brain Res, 1995. **676**(2): p. 235-44.
11. Berlau, D.J. and J.L. McGaugh, *Basolateral amygdala lesions do not prevent memory of context-footshock training.* Learn Mem, 2003. **10**(6): p. 495-502.
12. Malin, E.L. and J.L. McGaugh, *Differential involvement of the hippocampus, anterior cingulate cortex, and basolateral amygdala in memory for context and footshock.* Proc Natl Acad Sci U S A, 2006. **103**(6): p. 1959-63.
13. Uysal, N., et al., *Acute footshock-stress increases spatial learning-memory and correlates to increased hippocampal BDNF and VEGF and cell numbers in adolescent male and female rats.* Neurosci Lett, 2012. **514**(2): p. 141-6.
14. Anderson, R.I., et al., *Stress, kappa manipulations, and aversive effects of ethanol in adolescent and adult male rats.* Neuroscience, 2013. **249**: p. 214-22.
15. Lebron-Milad, K. and M.R. Milad, *Sex differences, gonadal hormones and the fear extinction network: implications for anxiety disorders.* Biol Mood Anxiety Disord, 2012. **2**: p. 3.
16. Wickens, M.M., D.A. Bangasser, and L.A. Briand, *Sex Differences in Psychiatric Disease: A Focus on the Glutamate System.* Front Mol Neurosci, 2018. **11**: p. 197.
17. Ogrodniczuk, J.S., et al., *Body talk: Sex differences in the influence of alexithymia on physical complaints among psychiatric outpatients.* Psychiatry Res, 2018. **261**: p. 168-172.
18. Green, T., S. Flash, and A.L. Reiss, *Sex differences in psychiatric disorders: what we can learn from sex chromosome aneuploidies.* Neuropsychopharmacology, 2019. **44**(1): p. 9-21.
19. Bangasser, D.A. and R.J. Valentino, *Sex differences in stress-related psychiatric disorders: neurobiological perspectives.* Front Neuroendocrinol, 2014. **35**(3): p. 303-19.
20. Kokras, N. and C. Dalla, *Sex differences in animal models of psychiatric disorders.* Br J Pharmacol, 2014. **171**(20): p. 4595-619.
21. Costa, G., et al., *Influence of dopamine transmission in the medial prefrontal cortex and dorsal striatum on the emission of 50-kHz ultrasonic vocalizations in rats treated with amphetamine:*

Effects on drug-stimulated and conditioned calls. Prog Neuropsychopharmacol Biol Psychiatry, 2020. **97**: p. 109797.

22. Demaestri, C., H.C. Brenhouse, and J.A. Honeycutt, *22 kHz and 55 kHz ultrasonic vocalizations differentially influence neural and behavioral outcomes: Implications for modeling anxiety via auditory stimuli in the rat.* Behav Brain Res, 2019. **360**: p. 134-145.

23. Mulvihill, K.G. and S.M. Brudzynski, *Effect of microinjections of dopamine into the nucleus accumbens shell on emission of 50kHz USV: Comparison with effects of d-amphetamine.* Pharmacol Biochem Behav, 2019. **176**: p. 23-32.

24. Willey, A.R. and L.P. Spear, *Development of anticipatory 50 kHz USV production to a social stimuli in adolescent and adult male Sprague-Dawley rats.* Behav Brain Res, 2012. **226**(2): p. 613-8.

25. Cagiano, R., et al., *Mediation of rat postejaculatory 22 kHz ultrasonic vocalization by dopamine D2 receptors.* Pharmacol Biochem Behav, 1989. **34**(1): p. 53-8.

26. Reno, J.M., et al., *Negative Affect-Associated USV Acoustic Characteristics Predict Future Excessive Alcohol Drinking and Alcohol Avoidance in Male P and NP Rats.* Alcohol Clin Exp Res, 2017. **41**(4): p. 786-797.

27. Taylor, J.O., C.M. Urbano, and B.G. Cooper, *Differential patterns of constant frequency 50 and 22 khz usv production are related to intensity of negative affective state.* Behav Neurosci, 2017. **131**(1): p. 115-126.

28. Thakore, N., et al., *Alcohol enhances unprovoked 22-28 kHz USVs and suppresses USV mean frequency in High Alcohol Drinking (HAD-1) male rats.* Behav Brain Res, 2016. **302**: p. 228-36.

29. Litvin, Y., D.C. Blanchard, and R.J. Blanchard, *Rat 22kHz ultrasonic vocalizations as alarm cries.* Behav Brain Res, 2007. **182**(2): p. 166-72.

30. Walker, Q.D., et al., *Vaginal lavage attenuates cocaine-stimulated activity and establishes place preference in rats.* Pharmacol Biochem Behav, 2002. **73**(4): p. 743-52.

31. Mitchell, D., et al., *Poison induced pica in rats.* Physiol Behav, 1976. **17**(4): p. 691-7.

32. Watson, P.J. and C. Leitner, *Patterns of increased and decreased ingestive behavior after injections of lithium chloride and 2-deoxy-D-glucose.* Physiol Behav, 1988. **43**(6): p. 697-704.

33. Meachum, C.L. and I.L. Bernstein, *Behavioral conditioned responses to contextual and odor stimuli paired with LiCl administration.* Physiol Behav, 1992. **52**(5): p. 895-9.

34. Kosar, E., H.J. Grill, and R. Norgren, *Gustatory cortex in the rat. II. Thalamocortical projections.* Brain Res, 1986. **379**(2): p. 342-52.

35. Kosar, E., H.J. Grill, and R. Norgren, *Gustatory cortex in the rat. I. Physiological properties and cytoarchitecture.* Brain Res, 1986. **379**(2): p. 329-41.

36. Gallo, M., G. Roldan, and J. Bures, *Differential involvement of gustatory insular cortex and amygdala in the acquisition and retrieval of conditioned taste aversion in rats.* Behav Brain Res, 1992. **52**(1): p. 91-7.

37. Lin, J.Y., J. Arthurs, and S. Reilly, *Gustatory insular cortex, aversive taste memory and taste neophobia.* Neurobiol Learn Mem, 2015. **119**: p. 77-84.

38. King, M.S., *Distribution of Fos-immunoreactive neurons in the gustatory cortex elicited by intra-oral infusion of taste solutions in conscious rats.* Brain Res, 2018. **1683**: p. 67-77.

39. Sticht, M.A., et al., *Endocannabinoid regulation of nausea is mediated by 2-arachidonoylglycerol (2-AG) in the rat visceral insular cortex.* Neuropharmacology, 2016. **102**: p. 92-102.

40. Cechetto, D.F. and C.B. Saper, *Evidence for a viscerotopic sensory representation in the cortex and thalamus in the rat.* J Comp Neurol, 1987. **262**(1): p. 27-45.

41. Allen, G.V., et al., *Organization of visceral and limbic connections in the insular cortex of the rat.* J Comp Neurol, 1991. **311**(1): p. 1-16.

42. Spencer, C.M. and T.A. Houpt, *Dynamics of c-fos and ICER mRNA expression in rat forebrain following lithium chloride injection.* Brain Res Mol Brain Res, 2001. **93**(2): p. 113-26.

43. Spencer, C.M., et al., *Area postrema lesions attenuate LiCl-induced c-Fos expression correlated with conditioned taste aversion learning.* Physiol Behav, 2012. **105**(2): p. 151-60.

44. Herisson, F.M., et al., *Oxytocin Acting in the Nucleus Accumbens Core Decreases Food Intake.* J Neuroendocrinol, 2016. **28**(4).

45. Klockars, O.A., et al., *Oxytocin administration in the basolateral and central nuclei of amygdala moderately suppresses food intake.* Neuroreport, 2018. **29**(6): p. 504-510.

46. Head, M.A., et al., *Effect of Oxytocin on Hunger Discrimination.* Front Endocrinol (Lausanne), 2019. **10**: p. 297.

47. Wald, H.S., et al., *NTS and VTA oxytocin reduces food motivation and food seeking.* Am J Physiol Regul Integr Comp Physiol, 2020. **319**(6): p. R673-R683.

48. Piekut, D.T., S. Pretel, and C.D. Applegate, *Activation of oxytocin-containing neurons of the paraventricular nucleus (PVN) following generalized seizures.* Synapse, 1996. **23**(4): p. 312-20.

49. Olszewski, P.K., et al., *Opioids affect acquisition of LiCl-induced conditioned taste aversion: involvement of OT and VP systems.* Am J Physiol Regul Integr Comp Physiol, 2000. **279**(4): p. R1504-11.

50. Pirnik, Z. and A. Kiss, *Detection of oxytocin mRNA in hypertonic saline Fos-activated PVN neurons: comparison of chromogens in dual immunocytochemical and in situ hybridization procedure.* Endocr Regul, 2002. **36**(1): p. 23-30.

51. Pirnik, Z., B. Mravec, and A. Kiss, *Fos protein expression in mouse hypothalamic paraventricular (PVN) and supraoptic (SON) nuclei upon osmotic stimulus: colocalization with vasopressin, oxytocin, and tyrosine hydroxylase.* Neurochem Int, 2004. **45**(5): p. 597-607.

52. Nasser, H.M. and G.P. McNally, *Neural correlates of appetitive-aversive interactions in Pavlovian fear conditioning.* Learn Mem, 2013. **20**(4): p. 220-8.

53. Uematsu, A., et al., *Correlation Between Activation of the Prelimbic Cortex, Basolateral Amygdala, and Agranular Insular Cortex During Taste Memory Formation.* Cereb Cortex, 2015. **25**(9): p. 2719-28.

54. Hernadi, I., et al., *Alterations of conditioned taste aversion after microiontophoretically applied neurotoxins in the medial prefrontal cortex of the rat.* Brain Res Bull, 2000. **53**(6): p. 751-8.

55. Hefner, K., et al., *Impaired fear extinction learning and cortico-amygdala circuit abnormalities in a common genetic mouse strain.* J Neurosci, 2008. **28**(32): p. 8074-85.

56. Gonzalez, M.C., et al., *Medial prefrontal cortex dopamine controls the persistent storage of aversive memories.* Front Behav Neurosci, 2014. **8**: p. 408.

57. Ramirez-Lugo, L., et al., *Choice Behavior Guided by Learned, But Not Innate, Taste Aversion Recruits the Orbitofrontal Cortex.* J Neurosci, 2016. **36**(41): p. 10574-10583.

58. Hurley, S.W. and R.M. Carelli, *Activation of Infralimbic to Nucleus Accumbens Shell Pathway Suppresses Conditioned Aversion in Male But Not Female Rats.* J Neurosci, 2020. **40**(36): p. 6888-6895.

59. Yim, C.Y. and G.J. Mogenson, *Response of nucleus accumbens neurons to amygdala stimulation and its modification by dopamine.* Brain Res, 1982. **239**(2): p. 401-15.

60. Prado-Alcala, R. and R.A. Wise, *Brain stimulation reward and dopamine terminal fields. I. Caudate-putamen, nucleus accumbens and amygdala.* Brain Res, 1984. **297**(2): p. 265-73.

61. Yasoshima, Y., T.R. Scott, and T. Yamamoto, *Memory-dependent c-Fos expression in the nucleus accumbens and extended amygdala following the expression of a conditioned taste aversive in the rat.* Neuroscience, 2006. **141**(1): p. 35-45.

62.   Moscarello, J.M., O. Ben-Shahar, and A. Ettenberg, *Dynamic interaction between medial prefrontal cortex and nucleus accumbens as a function of both motivational state and reinforcer magnitude: a c-Fos immunocytochemistry study.* Brain Res, 2007. **1169**: p. 69-76.

63.   Yamamoto, T., *Central mechanisms of roles of taste in reward and eating.* Acta Physiol Hung, 2008. **95**(2): p. 165-86.

64.   Moscarello, J.M., O. Ben-Shahar, and A. Ettenberg, *Effects of food deprivation on goal-directed behavior, spontaneous locomotion, and c-Fos immunoreactivity in the amygdala.* Behav Brain Res, 2009. **197**(1): p. 9-15.

65.   Alejandro Borja, G.P., et al., *Accumbens and amygdala in taste recognition memory: The role of d1 dopamine receptors.* Neurobiol Learn Mem, 2020. **174**: p. 107277.

66.   Lamprecht, R. and Y. Dudai, *Differential modulation of brain immediate early genes by intraperitoneal LiCl.* Neuroreport, 1995. **7**(1): p. 289-93.

67.   Yamamoto, T., et al., *Neural substrates for conditioned taste aversion in the rat.* Behav Brain Res, 1994. **65**(2): p. 123-37.

68.   Yamamoto, T., *Neural mechanisms of taste aversion learning.* Neurosci Res, 1993. **16**(3): p. 181-5.

69.   Olson, B.R., et al., *c-Fos Expression in Rat Brain and Brainstem Nuclei in Response to Treatments That Alter Food Intake and Gastric Motility.* Mol Cell Neurosci, 1993. **4**(1): p. 93-106.

70.   Kim, J., et al., *Basolateral to Central Amygdala Neural Circuits for Appetitive Behaviors.* Neuron, 2017. **93**(6): p. 1464-1479 e5.

71.   Namburi, P., et al., *A circuit mechanism for differentiating positive and negative associations.* Nature, 2015. **520**(7549): p. 675-8.

72.   Gore, F., et al., *Neural Representations of Unconditioned Stimuli in Basolateral Amygdala Mediate Innate and Learned Responses.* Cell, 2015. **162**(1): p. 134-45.

73.   Kim, J., et al., *Antagonistic negative and positive neurons of the basolateral amygdala.* Nat Neurosci, 2016. **19**(12): p. 1636-1646.

74.   Zhang, X. and B. Li, *Population coding of valence in the basolateral amygdala.* Nat Commun, 2018. **9**(1): p. 5195.

75.   O'Neill, P.K., F. Gore, and C.D. Salzman, *Basolateral amygdala circuitry in positive and negative valence.* Curr Opin Neurobiol, 2018. **49**: p. 175-183.

76.   Zhang, X., J. Kim, and S. Tonegawa, *Amygdala Reward Neurons Form and Store Fear Extinction Memory.* Neuron, 2020. **105**(6): p. 1077-1093 e7.

77.   Inui, T., et al., *The Basolateral Nucleus of the Amygdala Executes the Parallel Processes of Avoidance and Palatability in the Retrieval of Conditioned Taste Aversion in Male Rats.* eNeuro, 2019. **6**(4).

78.   Pelchat, M.L., et al., *Quality of acquired responses to tastes by Rattus norvegicus depends on type of associated discomfort.* J Comp Psychol, 1983. **97**(2): p. 140-53.

79.   Blanchard, R.J., et al., *Twenty-two kHz alarm cries to presentation of a predator, by laboratory rats living in visible burrow systems.* Physiol Behav, 1991. **50**(5): p. 967-72.

80.   Hinderliter, C.F. and J.R. Misanin, *Context familiarity and delayed conditioned taste aversion in young-adult and old-age rats.* Percept Mot Skills, 1993. **77**(3 Pt 2): p. 1403-6.

81.   Foy, M.R. and J.G. Foy, *Reversal of long-delay conditioned taste aversion learning in rats by sex hormone manipulation.* Integr Physiol Behav Sci, 2003. **38**(3): p. 203-13.

82.   Chen, K., et al., *c-Fos expression in rat brainstem following intake of sucrose or saccharin.* Front Med, 2011. **5**(3): p. 294-301.

83.   Yamamoto, T., et al., *C-fos expression in the rat brain after intraperitoneal injection of lithium chloride.* Neuroreport, 1992. **3**(12): p. 1049-52.

84.   Swank, M.W., G.E. Schafe, and I.L. Bernstein, *c-Fos induction in response to taste stimuli previously paired with amphetamine or LiCl during taste aversion learning.* Brain Res, 1995. **673**(2): p. 251-61.

85.   Thiele, T.E., M.F. Roitman, and I.L. Bernstein, *c-Fos induction in rat brainstem in response to ethanol- and lithium chloride-induced conditioned taste aversions.* Alcohol Clin Exp Res, 1996. **20**(6): p. 1023-8.

86.   Houpt, T.A., et al., *c-Fos induction in the rat nucleus of the solitary tract by intraoral quinine infusion depends on prior contingent pairing of quinine and lithium chloride.* Physiol Behav, 1996. **60**(6): p. 1535-41.

87.   Travers, S.P. and R. Norgren, *Coding the sweet taste in the nucleus of the solitary tract: differential roles for anterior tongue and nasoincisor duct gustatory receptors in the rat.* J Neurophysiol, 1991. **65**(6): p. 1372-80.

88.   Spray, K.J., C.B. Halsell, and I.L. Bernstein, *c-Fos induction in response to saccharin after taste aversion learning depends on conditioning method.* Brain Res, 2000. **852**(1): p. 225-7.

89.   Kinzeler, N.R. and S.P. Travers, *Licking and gaping elicited by microstimulation of the nucleus of the solitary tract.* Am J Physiol Regul Integr Comp Physiol, 2008. **295**(2): p. R436-48.

90.   Roussin, A.T., et al., *Taste coding in the nucleus of the solitary tract of the awake, freely licking rat.* J Neurosci, 2012. **32**(31): p. 10494-506.

91.   Weiss, M.S., J.D. Victor, and P.M. Di Lorenzo, *Taste coding in the parabrachial nucleus of the pons in awake, freely licking rats and comparison with the nucleus of the solitary tract.* J Neurophysiol, 2014. **111**(8): p. 1655-70.

92.   Fu, O., et al., *SatB2-Expressing Neurons in the Parabrachial Nucleus Encode Sweet Taste.* Cell Rep, 2019. **27**(6): p. 1650-1656 e4.

93.   Gal-Ben-Ari, S. and K. Rosenblum, *Molecular Mechanisms Underlying Memory Consolidation of Taste Information in the Cortex.* Frontiers in Behavioral Neuroscience, 2012. **5**(87).

94.   Rosenblum, K., N. Meiri, and Y. Dudai, *Taste memory: the role of protein synthesis in gustatory cortex.* Behav Neural Biol, 1993. **59**(1): p. 49-56.

95.   Geddes, R.I., et al., *Gustatory insular cortex lesions disrupt drug-induced, but not lithium chloride-induced, suppression of conditioned stimulus intake.* Behav Neurosci, 2008. **122**(5): p. 1038-50.

96.   Soto, A., et al., *c-Fos activity in the insular cortex, nucleus accumbens and basolateral amygdala following the intraperitoneal injection of saccharin and lithium chloride.* Neurosci Lett, 2017. **647**: p. 32-37.

97.   Reilly, S. and M.A. Bornovalova, *Conditioned taste aversion and amygdala lesions in the rat: a critical review.* Neurosci Biobehav Rev, 2005. **29**(7): p. 1067-88.

98.   Frank, G.K., et al., *Altered structural and effective connectivity in anorexia and bulimia nervosa in circuits that regulate energy and reward homeostasis.* Transl Psychiatry, 2016. **6**(11): p. e932.

99.   Frank, G.K.W., et al., *Association of Brain Reward Learning Response With Harm Avoidance, Weight Gain, and Hypothalamic Effective Connectivity in Adolescent Anorexia Nervosa.* JAMA Psychiatry, 2018. **75**(10): p. 1071-1080.

100.   Drewnowski, A., et al., *Taste and eating disorders.* Am J Clin Nutr, 1987. **46**(3): p. 442-50.

101.   Sunday, S.R. and K.A. Halmi, *Taste perceptions and hedonics in eating disorders.* Physiol Behav, 1990. **48**(5): p. 587-94.

102.   Chao, A.M., et al., *A Systematic Review of Taste Differences Among People With Eating Disorders.* Biol Res Nurs, 2020. **22**(1): p. 82-91.

103.   Drewnowski, A., *Taste responsiveness in eating disorders.* Ann N Y Acad Sci, 1989. **575**: p. 399-408; discussion 408-9.

104. Frank, S., S. Kullmann, and R. Veit, *Food related processes in the insular cortex.* Front Hum Neurosci, 2013. **7**: p. 499.
105. Friederich, H.C., et al., *Neurocircuit function in eating disorders.* Int J Eat Disord, 2013. **46**(5): p. 425-32.
106. Alonso-Alonso, M., *Brain, reward, and eating disorders: a matter of taste?* Am J Psychiatry, 2013. **170**(10): p. 1082-5.
107. Hernandez, V.S., et al., *Hypothalamic Vasopressinergic Projections Innervate Central Amygdala GABAergic Neurons: Implications for Anxiety and Stress Coping.* Front Neural Circuits, 2016. **10**: p. 92.
108. McLaughlin, R.J. and S.B. Floresco, *The role of different subregions of the basolateral amygdala in cue-induced reinstatement and extinction of food-seeking behavior.* Neuroscience, 2007. **146**(4): p. 1484-94.
109. Liu, C., J. Showalter, and P.S. Grigson, *Ethanol-induced conditioned taste avoidance: reward or aversion?* Alcohol Clin Exp Res, 2009. **33**(3): p. 522-30.
110. Roitman, M.F., et al., *Hedonic and nucleus accumbens neural responses to a natural reward are regulated by aversive conditioning.* Learn Mem, 2010. **17**(11): p. 539-46.
111. Yamamoto, T. and K. Ueji, *Brain mechanisms of flavor learning.* Front Syst Neurosci, 2011. **5**: p. 76.
112. Verendeev, A. and A.L. Riley, *Relationship between the rewarding and aversive effects of morphine and amphetamine in individual subjects.* Learn Behav, 2011. **39**(4): p. 399-408.
113. Verendeev, A. and A.L. Riley, *Conditioned taste aversion and drugs of abuse: history and interpretation.* Neurosci Biobehav Rev, 2012. **36**(10): p. 2193-205.