Comparative meta-proteomic analysis for the identification of novel plasmodesmata proteins and regulatory cues.

1   Philip Kirk[1], Sam Amsbury[2], Liam German[1], Rocio Gaudioso-Pedraza[1], Yoselin

2   Benitez-Alfonso[1*]

4   [1] Centre for Plant Science, School of Biology, University of Leeds, Leeds LS2 9JT, UK.

5   [2] Department of Animal and Plant Science, University of Sheffield, Sheffield, S10 2TN, UK.

7   *** Correspondence:** y.benitez-alfonso@leeds.ac.uk; phone: +44 113 343 2811

Summary

A major route for cell-to-cell signaling is via cell wall-embedded pores termed plasmodesmata

(PD) forming the symplasm. PD regulate many aspects of plant development and responses to

the environment however, our understanding of what factors affect their structure and

permeability is limited. In this paper, a meta-analysis is presented as a tool for the identification

of conditions affecting PD transport and *in silico* generation of PD proteomes for species of

interest. The custom-built pipeline searches the whole genome for protein structural features

and conserved domains identified on experimental proteomes and use it to predict PD

candidates in 22 compatible plant species. Using the *in silico* proteome and microarray

analysis, interactions between PD genes and conditions affecting PD function are identified.

High salinity and osmotic stress affect a significant number of PD candidate genes and we

provide evidence that these conditions regulate symplasmic transport of GFP. Using the

pipeline, the *in silico* PD proteome for *Medicago truncatula* was generated, as an example of

a plant in which experimental data is not available. The identification of a candidate receptor

like protein was experimentally validated in *M. truncatula* transgenic roots expressing

fluorescently tagged protein fusion. Together the results highlight the power of our newly

designed tool in the identification of new factors and proteins influencing PD in diverse plant

species.

Keywords: *Arabidopsis thaliana*, cell-to-cell communication, *Medicago truncatula,* phloem

unloading, plasmodesmata proteome**,** symplasmic transport, transcriptomic analysis.

Introduction

Without careful coordination of the processes occurring in distant cells, plants would be

incapable of responding effectively to changes in the environment, to the availability of

resources for growth or to defend themselves against predators. Unsurprisingly, plants have

evolved a myriad of long- and short- distance signalling pathways underpinning their ability to

adapt and thrive in diverse conditions on Earth. A major route for signalling is the symplast,

the continuous cytoplasmic connections established by cell walls-embedded pores termed

plasmodesmata (PD). PD are dynamic structures tightly controlled to regulate intercellular

signalling during development and in response to the environment (Li et al., 2021). Despite

their importance, many questions remain unanswered especially regarding the molecular

composition and the conditions that affect their function.

Over the last decade, proteomic analysis of PD-enriched membrane fractions has greatly
improved our understanding of what proteins may localise to and associate with PD
(Fernandez-Calvino et al., 2011; Park et al., 2017; Leijon et al., 2018; Brault et al., 2019).
Localisation of labelled protein fusions using confocal, FRET-FLIM (Förster resonance energy
transfer by fluorescence lifetime imaging) or transmission electron microscopy has
experimentally confirmed PD association for around 60 proteins in the model plant *Arabidopsis*
thaliana (Table S1). The PD proteomes include callose metabolic enzymes, such as callose
synthase 3 (CALS3) and the PD-located β-(1,3)–glucanases PdBGs and AtBG_ppap (Sagi *et*
al., 2005; Vatén *et al.*, 2011; Benitez-Alfonso *et al.*, 2013). It also comprises signalling proteins
and kinases such as the PD-Located Protein family (PDLPs) and the Lysin Motif domain-
containing glycosylphosphatidylinositol-anchored protein 2 (LYM2) (Thomas *et al.*, 2008;
Faulkner *et al.*, 2013). Proposed functions for many of these factors was recently reviewed in
the context of cell-cell connectivity (Amsbury *et al.*, 2018; Dorokhov *et al.*, 2019).
Detection in PD proteomes is not proof of PD localisation as preparations of PD domains
within cell walls are often contaminated with non-PD structures such as the plasma membrane
(PM), cell walls and the endoplasmic reticulum (ER) (Brault *et al.*, 2019). The detection of
proteins belonging to the ER is often attributed, not always verified, to the desmotubule (DT),
a tubular structure that runs through PD connecting the ER of neighbouring cells. To eliminate
contaminants, proteomic data from PM fractions is used but this is far from an ideal
solution (Leijon et al., 2018; Brault et al., 2019). Contamination issues aside, the isolation of
PD enriched fractions has been optimised for cell cultures which do not reflect the complexity
and dynamic composition of PD associated with plant development and responses to the
environment. Phylogenetic analysis can be used to identify family members expressed in
distinct tissues but this is time consuming and predictions for diverse plant species is not always
succesful.
In this paper, we present a meta-analysis to overcome some of the limitations imposed by
proteomic screens. Comparing four published proteomes, we observed considerable overlap in
protein subfamily composition. Based on this, we developed an analysis pipeline (implemented
in R and named Plasmodesmata In silico Proteome 1 or PIP1; available as a resource) to rapidly
generate candidate PD proteome lists for compatible species (currently 22 species). Overlap
between predictions and experimental data demonstrates the validity of this approach. PD
proteome lists were obtained *in silico* for the well-studied *A. thaliana* and for *Medicago*
truncatula, a species for which a PD proteome has not been experimentally determined.
Bioinformatic analysis of public transcriptomes allowed us to generate co-expression tables

suggesting novel interactions between PD genes and conditions potentially affecting PD

function. Using this approach and knowledge, we demonstrate the effect of salinity and high

osmotic potential in the symplasmic unloading of GFP and confirm PD localization for a novel

receptor like protein in *M. truncatula*. Together, the results highlight the power of our newly

designed tool in the identification of new factors affecting PD in diverse plant species.

Material and Methods

<u>Plant material and growth conditions</u>

Seeds of *A. thaliana* plants expressing p*SUC2*::GFP were surface sterilised with ethanol.

Control ATS media was prepared as described by Wilson et al. (1990) with 0.8% (w/v) agar

(Type E, Sigma- Aldrich). When required, plates were prepared using ATS-based media

supplemented with 3% (w/v) polyethylene glycol (MW 8000) or with 75 mM NaCl as

indicated. Seeds were stratified at 4°C for 4 days before being transferred to long day light

conditions (22°C, 16 h day, 150 mE/m2/s) for growth.

For *p35S::*Medtr1g073320-YFP transgenic roots, Gateway cloning was used to generate the

vector following manufacture's instruction (Invitrogen, USA). In brief, primers were designed

to amplify Medtr1g073320 with linkers compatible for cloning into pDNR221 by BP reaction

(Medtr1g0733201-Attb1: GGG GAC AAG TTT GTA CAA AAA AGC AGG CTC CAT GTT

TTG ATT CTC TCT CCA; Medtr1g0733201-Attb2: GGG GAC CAC TTT GTA CAA GAA

AGC TGG GTA CCA CAA ATC TCT TTC AGC AAA AA). Positive pDNR clones were

confirmed by sequencing and used in LR reaction with the destination vector pB7YWG2

(Karimi *et al.*, 2002). The *p35S::*Medtr1g073320-YFP vector was amplified in *E. coli* and

expressed in *Agrobacterium rhizogenes* for transformation.

Wild-type A17 *M. truncatula* seeds were lightly scarified with sandpaper and sterilised for 3

minutes in a 10% sodium hypochlorite solution, washed with water and left undisturbed in

water for 4 hours. The seeds were transferred to agar-water plates and left in the dark for 7 days

at 4 °C. Plates were transferred to RT overnight and *A. rhizogenes* carrying the *p35S*-

Medtr1g073320-YFP was used for root transformation as described by (Boisson-Dernier *et al.*,

2001). After two to four weeks post-transformation, transgenic roots expressing YFP fusions

were identified using fluorescent microscopy and selected for confocal imaging.

<u>PD proteome meta-analysis</u>

PD proteomic data for *A. thaliana*, *Nicotiana benthamiana* and *Populus trichocarpa* were

retrieved from their original publications (Fernandez-Calvino *et al.*, 2011; Park *et al.*, 2017;

Leijon *et al.*, 2018; Brault *et al.*, 2019) and subfamilies annotated based on PANTHER16 (Mi

et al., 2021). The proteomes were incorporated into a pipeline using a custom-built R script

(https://github.com/PhilPlantMan/PIP1). Instructions on how to use the pipeline are included

in the GitHub repository along with instructions on how to customise pipeline parameters

depending on user requirements. Necessary databases including the PD proteomes used here

are packaged with the script.

The script dependencies include the R library 'biomartr' (Drost and Paszkowski, 2017).

Protein features were predicted using the R library 'ragp' which integrates multiple tools for

glycophosphatidylinositol anchor (GPI) and secretory signal peptide (SP) predictions

(Emanuelsson *et al.*, 2007; Käll *et al.*, 2007; Pierleoni *et al.*, 2008; Almagro Armenteros *et al.*,

2019; Gíslason *et al.*, 2019; Dragićević *et al.*, 2020). Genes were classified as having a GPI

and/or SP if at least one tool returned true for that feature. Transmembrane domains (TM)

domains prediction was acquired from the ENSEMBL database annotation via the R package

'biomartr' using TMHMM (Krogh *et al.*, 2001). Predictions for N-myristoylation, S-

farnesylation, S-geranylgeranylation, S-palmitoylation and S-nitrosylation were made using

tools available by the Cuckoo workgroup (Xue *et al.*, 2010; Xie *et al.*, 2016; Ning *et al.*, 2020).

For protein feature enrichment analysis, the tools described above were used to predict features

for the whole *A. thaliana* genome (Araport11). Fisher's exact test was used to determine

statistical significance (cutoff provided in figure legends). To be fully pipeline-compatible for

proteome input or to generate candidate gene output, the species must be listed in both Ensembl

Plant databases (used by biomartr to retrieve sequence information) and in PANTHER16 (used

to retrieve subfamily annotation) (Drost & Paszkowski, 2017; Howe *et al.*, 2020; Mi *et al.*,

2021). Currently, there are 22 compatible plant species (Table S2). For non compatible species,

such as *N. benthamiana*, Arabidopsis orthologues can use. This enable integration of the PD

proteome of Park et al. (2017) in the pipeline.

Lists of genes were compared by drawing a Venn or Euler diagram using the R library 'eulerr'.

The significance of the overlap between candidate lists and proteomes was determined using

bootstrap analysis as follows. Sets of genes the same length as a candidate list and a proteome

were randomly sampled from Araport11. The overlap in genes between the samples was

recorded and repeated for *n* cycles ($n = 10,000$). Probability (p) was calculated as the proportion

of cycles that attained an overlap at least as large as was observed between the candidate list

and the proteome. The size of the overlap by chance was given as the median overlap in random

samples over *n* cycles.

 <u>Expression analysis</u>

The gene correlation dataset 'Ath-u.c1-0' was downloaded from the ATTEDII database

(Obayashi *et al.*, 2018). Optimal gene order and the corresponding dendrogram were computed

using hierarchical clustering. The dendrogram was cut at an optimised height (h = 16) that gave

a sufficient number/size of clusters (k = 151). These processes were performed in base R.

Enrichment of genes within clusters from candidate lists and verified genes were determined

using pairwise comparisons with Fisher's exact test (p<0.05, holm-adjusted) via the R library

'rcompanion '.

Publicly available microarrays for a subset of conditions were independently analysed.

Microarray datasets (Affymetrix GeneChip Arabidopsis ATH1-121501) were downloaded

from EBI ArrayExpress. For each experiment, expression data were normalised using the

robust multi-array average (RMA) method and log_2 transformed with the R package 'oligo'

(Carvalho & Irizarry, 2010). Principal component analysis was used to identify and exclude

outlier arrays and experiments with insufficient biological replicates. Genes with low levels

of expression were filtered out. A design matrix was constructed for each experiment and a

linear model applied using the R package 'limma' (Ritchie *et al.*, 2015). Differential expression

of genes and a multiple comparison correction were determined using empirical bayes statistics

via the package 'limma' and the results filtered by gene IDs. Heatmaps were constructed using

the R package 'ComplexHeatmap' (Gu *et al.*, 2016).

<u>Confocal microscopy imaging for the analysis of GFP diffusion and protein localization</u>

Seedlings of *A. thaliana* expressing *pSUC2::*GFP were mounted on glass slides in 10 μg/ml

propidium iodide (Sigma- Aldrich). Root tips were imaged using an LSM 800 upright confocal

microscope (Zeiss, Germany). Profiles of fluorescence were determined using line and

profiling tools in ImageJ. Lateral profiles were taken across the transition zone and rootward

profiles started from the basal/apical meristem transition zone ending 150 μm towards the root

tip. Fluorescence of lateral and rootward profiles were scaled between 0 and 1. Fluorescence

across each lateral profile was binned (bins = 100) to compensate for small differences in root

width. Fluorescence profiles of at least 6 plants per treatment were aggregated by calculating

the mean (±SD) for position along the profile and plotted.

Transgenic roots expressing p*35S*::Medtr1g073320-YFP were stained with aniline blue

fluorochrome (Biosupplies, Australia) in 0.1 M K_3PO_4 (pH 12) for callose detection. Roots

were imaged by confocal microscopy (LSM-880, Zeiss) using 488 nm excitation laser for YFP

and emission at 505–530 nm. Aniline blue was imaged with 405 nm excitation and emission
at 463 nm.
172
Results

<u>Workflow for the prediction of PD proteomes via meta-analysis of conserved subfamilies</u>
<u>and structural domains.</u>

Proteomic data obtained from *A. thaliana* (Fernandez-Calvino *et al.*, 2011; Brault *et al.*, 2019),
N. benthamiana (Park *et al.*, 2017) and *P. trichocarpa* (Leijon *et al.*, 2018) PD-enriched
fractions were fed into a custom-built R-based pipeline released as a resource with this article
(https://github.com/PhilPlantMan/PIP1). The workflow is described in Figure 1 and in the
Material and Methods section. First, subfamilies of genes identified in at least one of the PD
proteomes mentioned above were annotated using PANTHER16. Subfamilies classifications
are used to extract all genes making the *in silico* proteome for a particular plant of interest. For
some applications, using families classifications may be more appropriate and PIP1 prompts
users whether to generate candidates based on family or subfamily identifiers. Data presented

Figure 1. Pipeline to determine candidate plasmodesmata (PD) proteome. 4 proteome databases have been determined experimentally by Fernandez-Calvino et al., 2011; Park et al., 2017; Brault et al., 2019 and Leijon et al., 2018. A list was compiled of subfamilies of proteins identified in the different experimental proteome databases using PANTHER16. Genes belonging to these subfamilies were extracted for the target plant species. These were classified by whether its subfamily was present in at least two PD experimental proteomes or just one. Further classification was based on whether genes were predicted to have a signal peptide (SP), transmembrane domain (TM) or glycosylphosphatidylinositol anchor (GPI) as these features are enriched in verified PD genes (Figure S1). As a result, 4 list of genes were obtained (A-D) representing the predicted PD proteome for the target plant.

in this paper are based on candidate generation using subfamily identifiers as a more restrictive

approach leading to shorter candidate lists (see example for Arabidopsis below). The pipeline

categorises the output by whether the subfamily is present in one or multiple PD proteomes

and based on predictions of distinctive features identified in PD verified genes (Figure 1).

Features of membrane-bound proteins were predicted for the list of PD verified genes in Table

S1 using multiple online platforms as described in Material and Methods. When compared to

the Arabidopsis whole proteome, PD-localised proteins are overrepresented in predicted GPI

anchors, s-geranylgeranylation, s-palmitoylation, SP and TM domains (Figure S1). The

presence of a SP in combination with either a GPI and/or TM domain returned the highest

proportion of verified genes thus these features were chosen for gene categorisation.

As a result, the pipeline outputs 4 lists of genes: list A are genes of subfamilies present in more

than one PD proteome containing either a SP, GPI or TM domains; list B are also present in

multiple proteomes but lacking the membrane localising features, list C and D are genes in

subfamilies found in a single proteome either with (list C) and without (list D) the predicted

SP, GPI or TM features (Figure 1). Using this approach, 1882 gene subfamilies were identified

in at least one PD proteome (Table S3). As expected, there is substantial overlap between

subfamilies identified in the different proteomes, with 311 subfamilies identified in more than

one proteome (Table S3), forming list A and B. As PD verified genes usually display membrane

targeting features, list A should be considered the priority list but, since only four proteomes

are available, list C cannot be discarded. In support of this prediction, list A contains

subfamilies with known PD activities such as enzymes involved in callose metabolism (BG

and CALS) and receptors involved in signalling (such as PDLPs).

Applying PIP1 to proteome predictions in *Arabidopsis thaliana* and poplar.

For *Arabidopsis*, the *in silico* proteome comprises list A with 206 genes (158 subfamilies), list

B: 208 genes (152 subfamilies), list C: 751 genes (597 subfamilies) and D: 1117 genes (802

subfamilies) (Table S4). Most known PD genes are isolated from studies in *A. thaliana*. To

validate our approach, we determined the number of PD verified genes identified by the

pipeline. Generated candidate lists contain 60.7% of the genes known to localise at PD

identified either using fluorescent translational fusions and/or isolated in independent, non-

proteomic studies (compare Tables S4 and S1). These include the beta-1,3 glucanase

AtBG_ppap and C2 calcium/lipid-binding plant phosphoribosyltransferase family protein

MTCP6 and MTCP9 identified in the Brault *et al.* (2019) proteome and proteins identified in

independent studies such as the leucine-rich repeat receptor-like serine/threonine-protein

kinase BAM2 (Rosas-Diaz *et al.*, 2018). We failed to output known PD proteins such as CRINKLY4 (ACR4) or the PD-located protein 5 (PDLP5) likely because these subfamilies are low expressed or absent in cell cultures. Extending the output to family members instead of just subfamily members increased the coverage of known PD genes to 83.6% but consequently increased the total candidate gene size to over 9000 genes (Table 1). Genes in list A and C included in families containing at least one previously verified PD-localised protein were identified as the most likely targets for future experiments (Table S5). In Arabidopsis, this list comprises 55 new proteins.

Table 1. Verified PD genes and candidates identified in the Arabidopsis *in silico* proteome. The number of genes identified when using either subfamilies (subfam) or families (fam) classifications in PIP1 is provided.

In silico proteome list	Verified PD genes (subfam/fam)[a]	N° of candidates (subfam/fam)[b]
A	18/41	206/2345
B	1/4	208/1432
C	18/5	751/1961
D	0/1	1117/3935

[a] For a full list of verified PD genes and the publications that report them consult table S1.

[b] The lists of candidates identified using PIP1 and subfamilies classifications is provided is table S4.

Despite lacking features for PD localisation, list B contains a considerable number of genes identified in more than one PD proteome. The number of known PD proteins identified in list B and D suggest they might contain contaminants but it could also contain PD proteins with no predicted membrane targeting domains such as Remorin 2 REM1.2 (Huang *et al.*, 2019). We explored the hypothesis that list B might contain mobile proteins unintentionally captured within PD fractions. To test this hypothesis, the candidate PD gene lists were compared to the mobile proteome identified in *Cuscuta australis* (dodder) parasitising *A. thaliana* (Liu *et al.*, 2020). Candidate mobile proteins were identified in all candidate lists but list B had the largest overrepresentation (20.2%) with an overlap over 10x greater than would be expected by chance (Figure S2).

To evaluate the strength of the pipeline further, we generated candidate PD lists for *P. trichorcarpa*. 1032 out of 1148 genes/proteins identified in the PD-enriched fraction of poplar cell cultures (Leijon *et al.*, 2018) were predicted in the candidate lists generated with our pipeline. The lists contained 182 (list A), 129 (list B), 397 (list C) and 289 (list D) of the genes identified in the PD-enriched fraction (Leijon *et al.*, 2018). For list A and C the *in silico*

proteome is 50% larger than the experimental proteome, reflecting inclusion of proteins not
expressed in cell cultures. 116 genes identified in the experimental proteome are excluded
because they are not annotated within PANTHER subfamilies. When the poplar proteome is
excluded as input in the pipeline, the overlap between the candidates lists and the poplar PD
proteome remains high; 20x larger than would be expected by chance significant (Figure S3).
In summary, our results validate the use of PIP1 as a new resource to generate candidate PD
genes, and potentially mobile proteins, without the cost of time and resources associated with
growing and maintaining cell cultures carrying out their biochemical isolations, fractionations
and protein sequencing. The *in silico* proteome list is larger than experimental proteomes
because there are not constrictions associated with protein expression in cell cultures.
253
Drought and salinity regulate the expression of PD candidates and symplasmic transport.
The generation of PD proteomes not only enables target gene selection but, in combination
with transcriptomics, can reveal conditions that regulate PD genes. To establish whether this
is the case, co-expression data extracted from ATTED-II database was ordered by hierarchical
clustering to identify co-expressed genes in the *in silico* proteomes. For *A. thaliana*, two
clusters were found to be significantly over-represented (relative to the wider ATTED-II gene
set) in PD-verified genes: cluster 87 and cluster 100 (Figure 2 A, B). 11 genes from list A and
18 from list C were found in cluster 87, whereas cluster 100 contained 16 genes of list C (see
Table S5 for examples). The expression of PD-candidates in clusters 87 and 100 was also
analysed using publicly available microarrays as described in Materials and Methods. Figure
2C and Figure S4 show that the expression of the candidate genes overlaps with PD-marker
proteins such as AtBG_ppap, PDCB (PD callose binding proteins) and PDLP family members.

Figure 2: Expression analysis of PD candidates and PD verified genes. *Arabidopsis thaliana* co-expression data from ATTEDII were subset into clusters based on hierarchical clustering (k = 151). Representation analysis of the total ATTEDII database, candidate lists A-D and verified PD subfamilies was performed on each cluster; (A) cluster 87 and (B) cluster 100 show significant overrepresentation in verified subfamily members. Number underneath each bar = number of genes within that cluster. Bars with differing letters above are significantly different (Fisher's exact test, Holm corrected, p ≤ 0.05). (C) Differential expression (log$_2$FC, ranges from blue: downregulated to red: upregulated in relation to control) of genes in clusters 87 and 100 (only list A, C and verified genes are shown) determined using public microarrays from experiments pertaining to osmotic stress. Other treatments are presented in Supplemental figure 4. Columns labels bottom = ArrayExpress accession codes followed by a reference number. See Supplemental table 6 for further information. (left) Rows are grouped by cluster and then (dendrogram) ordered by hierarchical clustering. (right) Gene ID and properties. Asterisks in cells denote predictions on membrane targeting features (e.g. is predicted to have a SP). SP = Signal peptide, GPI = glycosylphosphatidylinositol anchor, TM = transmembrane domain. Cell colour above each column = plant material sampled.

When comparing the expression of PD-verified and PD-candidate genes in a variety of biotic and abiotic stress conditions (Figure S4), osmotic stress (drought, desiccation, elevated NaCl and exogenous treatment with PEG) was found to be a strong regulator (Figure 2C). To determine if these changes in expression reflect conditions affecting symplasmic transport, cell-to-cell connectivity was tested in root tips of transgenic *A. thaliana* seedlings expressing cytosolic GFP, driven by the phloem companion cell specific promoter *Sucrose Symporter 2 SUC2* (p*SUC2*::GFP) (Imlau *et al.*, 1999).

p*SUC2*::GFP seeds were germinated and grown in control conditions, 3% PEG and 75mM NaCl (Hunter *et al.*, 2019). Media water potential (at 25 °C) was estimated by adding the

Figure 3: GFP diffusion in the root meristem of *Arabidopsis thaliana* exposed to PEG and sodium chloride. (A) Seeds expressing p*SUC2*::GFP were grown on ATS control media, ATS supplemented with 3% (w/v) polyethylene glycol (PEG) or with 75 mM NaCl. Roots were counterstained with propidium iodide (PI) and images sequentially collected 4 days post-germination (4 dpg) at 561 nm (PI shown in grey) and 488 nm (green: GFP). Estimated water potential (MPa, see main text) is labelled on top of the representative pictures. (B-C) Transects highlighted by magenta boxes in A were used to determine GFP fluorescence profiles using mean grey values (AU: arbitrary units). (B) shows transects across the transition zone (TZ) while (C) shows transects rootward starting from the transition/proliferation zone (TZ/PZ) towards the root tip. (D-F) Similar experiments were carried out with seedlings germinated on ATS media and transferred for 1 day (1dpt) to 3% PEG or 75 mM NaCl. Scale bar = 100 μm. Charts show traces representing min-max normalised GFP fluorescence along the transect for N>6 plants (mean ± SE).

estimated solute potentials of individual medium components based on empirical and modelled
data (Robinson & Stokes, 1949; Ghashghaie *et al.*, 1991; Gopal & Iwama, 2007). The water
potential was reduced in both 3% PEG and 75mM NaCl (Figure 3A). Roots were imaged using
confocal microscopy and both lateral and rootward diffusion of GFP from phloem companion
cells were measured using mean grey values in Image J. The short and long term effects of
changes in water potential were evaluated by comparing changes in GFP distribution in roots
grown for 4 days directly in PEG and NaCl media (Figure 3A) and roots grown for 4 days in
control media and transferred for 24h to PEG and NaCl media (Figure 3D). There were no
noticeable changes in lateral distribution of GFP across primary roots grown permanently in
PEG and NaCl relative to control conditions (Figure 3 A, B) but the rootward profiles, show a
decrease in fluorescence in both, 3% PEG and 75mM NaCl (Figure 3C). The effect is stronger
in NaCl rootward fluorescence profiles, particularly between 50 μm - 150 μm towards the tip
where relative fluorescence drops over 13% in 75 mM NaCl and less than 2% in control
conditions (Figure 3C).
No significant differences in GFP distribution were observed 24 hours after transfer from
control to 3% PEG conditions (Figure 3 D-F) but fluorescence was diminished in the rootward
profile of roots transferred to 75 mM NaCl (Figure 3F). Over the length of 150 μm from the
tip, relative fluorescence decreased 11% in plants transferred to 75 mM NaCl. Lateral profiles
in roots transferred to 75 mM NaCl show a higher fluorescence (likely auto-fluorescence) in
the epidermis (Figure 3E).
Taken together, these data suggest that regulation of PD candidates in osmotic stress is
translated into changes in symplasmic connectivity demonstrating the potential for using
combined proteomic and transcriptomic meta-analysis as a method to determine conditions
regulating PD function and therefore symplasmic communication.
300
A meta-analysis approach predicts plasmodesmata proteins in *Medicago truncatula*.
One of the advantages to the pipeline outlined here is that it can be applied to species for which
an experimentally determined PD proteome is not publicly available. To demonstrate this, the
pipeline was used to generate candidate PD gene lists for *M. truncatula* (Table S6). List A and
C comprised 1018 genes belonging to subfamilies represented in at least one experimental
proteome and displaying membrane targeting features (SP, GPI or TM). These lists include *M.*
truncatula orthologues of the CALS, PDCB and PDLP genes (Table S7).

Figure 4: Expression analysis of *Medicago truncatula* PD candidates in rhizobia and nitrate transcriptomes. Gene expression analysis of PD candidates (lists A and C only) and verified PD genes BG2 and SUNN (see table S6). Differential gene expression (log₂FC) was determined using public microarray data of experiments relating to nitrate and rhizobia inoculation in root tissue. Columns labels bottom= ArrayExpress accession codes followed by a reference number. See Supplemental table 8 for further information. (left dendrogram) Rows are ordered by hierarchical clustering. Asterisks in cells denote that gene has a membrane targeting feature or are verified. SP = Signal peptide, GPI = glycophosphatidylinositol anchor, TM = transmembrane domain. Red arrow indicates the position of MTR_1g073320, which is used in further analysis.

Transcriptomic profiling was used to determine clusters of genes co-regulated within the predicted proteome lists A and C. Published microarray data from early rhizobia inoculation, various nitrate concentrations and nodulating roots (Table S8) were re-analysed. These conditions were selected as they have been previously shown to regulate PD genes and symplasmic communication (Haney & Long, 2010; Crook *et al.*, 2016; Gaudioso-Pedraza *et al.*, 2018). The β-(1,3)-glucanase, MtBG2 (Gaudioso-Pedraza et al., 2018), and the Super Numeric Nodule receptor (SUNN, Crook *et al.*, 2016) are, to our knowledge, the only verified PD-localised proteins in *M. truncatula*. A cluster containing MtBG2 and SUNN was identified comprising 70 genes from list A and C (Figure 4). Genes in this cluster share a similar

expression pattern, particularly in roots post-inoculated with rhizobia (E-MEXP-1097).

Interestingly, four genes encoding receptor-like proteins show a strikingly similar expression

profile to MtBG2. These are MTR_5g083910, MTR_4g014070, MTR_1g073320 and

MTR_2g011180. MTR_1g073320 (also known as Medtr1g073320) was selected for further

studies because it contains sequence domains characteristic of PDLP family members (Table

S7).

Figure 5: The *Medicago truncatula* protein Medtr1g073320 co-localizes with callose at PD microdomains. *Agrobacterium rhizogenes* transformed roots expressing Medtr1g073320 fused with YFP were counterstained with aniline blue to reveal callose deposition. Roots were imaged with a confocal microscope and (A) excitation laser 561 nm (green: Medtr1g073320-YFP) and (B) 405 nm (shown in magenta: aniline blue). Merged images are shown in C and co-localisation events highlighted with arrows. Bottom panels (A'-C') are magnification of the area highlighted by the white box in the top panels. Scale bar = 20 µm.

To established Medtr1g073320 localisation, C-terminal YFP fusions were created and

introduced in *M. truncatula* roots using *A. rhizogenes*- mediated transformation. One week old

transgenic roots were identified and counterstained with aniline blue which fluoresce in the

presence of callose- a cell wall glucan component associated with PD. Confocal microscope

images show Medtr1g073320-YFP in a punctate pattern on the cell periphery, co-localising

with callose deposits indicating PD targeting (Figure 5).

Altogether, the results demonstrate how the pipeline, implemented in our custom build R script,

can be used to predict the PD proteome in plant species where this information has not been

experimentally derived, opening doors to the identification of novel PD proteins participating

in the regulation of cell-to-cell communication.

Discussion

Many aspects of PD function and regulation are not well understood despite the increasing

evidence of the important role these structures play in plant signalling, organ development and

response to physiological and environmental cues (Petit *et al.*, 2020; Vu *et al.*, 2020; Li *et al.*,

2021). Proteomic data is lacking for most plant species due to difficulties in isolating clean PD

fractions and because association to PD domains can be quite dynamic, varying with

environmental conditions and tissue types (Grison *et al.*, 2019; Hunter *et al.*, 2019; Cheval *et*

al., 2020). These factors limit our current knowledge of what proteins associate with PD and

how their properties and function are controlled.

We created a pipeline that exploits the overlap in subfamily composition of existing PD

proteomes to generate candidate PD genes (Figure 1). This allows the generation of *in silico*

proteomes for 22 plant species which are represented in both PANTHER16 and in Ensembl

Plant databases. The gene lists were classified according to their presence in one or more PD

proteomes and according to predicted membrane targeting features overrepresented in verified

PD proteins. Based on our categorisation, list A (subfamilies present in multiple proteomes and

displaying predicted membrane targeting features) is more likely to contain PD localised

proteins but genes in list C (subfamilies present in a single proteome) can not be excluded as

only four PD proteomes are sequenced so far. Supporting our predictions, list A and C

contained 36 out of the 60 proteins reported to target PD in *A. thaliana,* 46 if using families for

in silico predictions (Table 1). The pipeline demonstrated to be effective in predicting the

poplar proteome showing significant overlap with the experimentally determined PD enriched

fraction (Fig. S3).

Combining proteomic with transcriptomic analysis we identified drought, osmotic and salinity

stresses as conditions affecting the expression of clusters of genes identified in the Arabidopsis

in silico proteomes. The diffusion in root meristems of GFP, expressed in companion cells

under the SUC2 promoter, was restricted after continuous (4 days) and short (24h) exposure to

3% PEG and 75mM NaCl (Figure 3), suggesting that the expression of PD candidates is linked

to the regulation of symplastic transport. We determined for the first time the *in silico* proteome

of *M. truncatula* using PIP1 and exploited published microarrays to study gene expression

during rhizobia infection and nodulation as conditions reported to affect symplasmic transport

(Gaudioso-Pedraza *et al.*, 2018). This strategy led us to identify the receptor protein

Medtr1g073320, which we confirmed localises at PD when expressed as fluorescent fusion in

transgenic roots (Figure 5). Together our results validate the use of the PIP1 pipeline in

combination with transcriptomic analysis for the identification of new PD proteins and
conditions influencing symplasmic communication.

Our approach relies on subfamily annotation within the PANTHER database. For the 40
reference plant genomes included in PANTHER16, family/subfamily annotation coverage is
between 60-95% depending on species (Mi *et al.*, 2021). For *A. thaliana* and poplar, the
coverage is at 89% and 81% respectively but for *M. truncatula*, coverage is only 67%. Genes
without a PANTHER classification will not be included in the output of the pipeline. We expect
our approach may have less predictive power in monocots or non-angiosperms, as experimental
proteomes are all from dicot species. It is unknown to what extend the composition of PD
differs between monocots and dicots, especially as they differ in cell walls composition
(Calderan-Rodrigues *et al.*, 2019). The experimental determination of more PD proteomes
including representative monocot species and the identification of PD proteins by independent
approaches will improve *in silico* predictions as these can be easily added as inputs to PIP1.
Adding, as input data, subfamilies of PD proteins identified by different means can get around
the limitations associated with the cell culture origin of experimental proteomes. Researchers
already use sequence - domain analysis, phylogeny and transcriptomics to study families of PD
proteins and to identify members expressed outside cell cultures (Thomas *et al.*, 2008; Simpson
et al., 2009; Gaudioso-Pedraza *et al.*, 2018). Our comparative meta-analysis provides a
platform to systematically apply this approach enabling *in silico* predictions of whole PD
proteome based in multiple instead of single experiments.

Applying the pipeline to *A. thaliana* or poplar generates a list of candidates larger than
experimentally determined because expression in cell cultures is not a pre-requisite for gene
identification. For example, the Arabidopsis *in silico* proteome contains BAM2, the Germin-
like protein PDGLP1 and CALNEXIN2 (CNX2), which are not present in any of the
experimental proteomes but localised at PD in independent studies (Fernandez-Calvino *et al.*,
2011; Liu *et al.*, 2017; Rosas-Diaz *et al.*, 2018; Brault *et al.*, 2019). The candidate lists also
includes subfamilies/proteins lacking predicted membrane targeting features (SP, TM and GPI)
(list B and D). Overlap between list B (subfamilies present in multiple proteomes) and the
mobile proteome, recently reported in dodder parasitising *A. thaliana* (Liu *et al.*, 2020),
suggest that these might be subfamilies of proteins identified while in transit via PD. List B
might also include proteins with unusual membrane targeting mechanisms or with features
poorly predicted by the available platforms. This is the case for REMORIN2, for example,
which localises to PD independently of the secretory pathway and lacks a predicted SP, GPI or
TM (Reymond *et al.*, 1996; Gronnier *et al.*, 2017; Huang *et al.*, 2019). Based on structural

predictions and proteomic count, the lists serve as a tool to prioritise the screening, instead of

rejecting, candidate PD proteins.

Besides the obvious reasons of obtaining *in silico* proteomes as a tool for the identification of

new PD genes, we propose that transcriptomic analysis of the candidate lists can predict

conditions affecting PD transport. Here, we showed that osmotic and salinity stresses modify

the expression of clusters of PD candidates in Arabidopsis and that symplasmic transport of

GFP is affected in these conditions. This aligns well with recent publications reporting changes

in symplasmic transport and in PD protein localisation in response to salt and mannitol (Grison

et al., 2019; Hunter *et al.*, 2019). It is not clear if PD function is affected by changes in the

localisation of PD proteins or by changes in turgor pressure or, more likely, to a combination

of these (Hernández-Hernández *et al.*, 2019). The identification and phenotypic

characterisation of mutants in PD genes and of relevant symplasmic mobile molecules are

necessary to mechanistically understand the biological role of PD in the plant response to these

abiotic stresses.

Our past research indicates that PD are regulated in response to rhizobia infection, a mechanism

mediated by the degradation of the cell wall polysaccharide callose by the MtBG2 protein

(Gaudioso-Pedraza *et al.*, 2018). PDLP regulation has been tightly linked to callose deposition

in Arabidopsis (Lee *et al.*, 2011; Wang *et al.*, 2013; Caillaud *et al.*, 2014). Here, we identified

Medtr1g073320 as a PD-located member of a subfamily of PDLP proteins (Thomas *et al.*,

2008) in *M. truncatula*. Expression tables suggest that Medtr1g073320 is co-regulated with

SUNN and MtBG2 during rhizobia infection and nodulation. This finding opens doors for

research on the mechanisms involving callose and PD regulation in the establishment of

nitrogen fixation in legumes, an important process for sustainable agriculture.

To summarise, an R-based tool has been made available to integrate the data obtained from PD

proteomes and enable the identification of new PD genes in a variety of plant species. Together

with transcriptome analysis, the pipeline becomes a useful tool to identify conditions affecting

PD composition and function. Our approach works with all plant species and genes with

subfamilies annotated in both PANTHER16 and Ensembl Plant databases and will only expand

as more new and updated UniProt Reference Proteomes are added (Drost & Paszkowski, 2017;

Howe *et al.*, 2020; Mi *et al.*, 2021). The pipeline can help in prioritising the targets for

validation from experimental proteomes and can predict PD proteins for species where PD

proteomic information is not available. The pipeline is publicly accessible and can be easily

modified by the user to add new sequenced proteomes and/or experimentally verified genes

thus will serve as a long-lasting resource for the plant community.

Acknowledgements

The authors would like to thank Hendrik Swiegers for his help in testing the pipeline and Dr. Eva Deinum and Dr. Michael Wilson for critical reading and help shaping the manuscript. This work was supported by UKRI Future Leader Fellowship (MR/T04263X/1), Leverhulme Trust Grant RPG-2016-136, a University of Leeds Gosden Studentship and a BBSRC DTP (BB/M011151/1).

Author Contributions

P.K. designed and generated the pipeline, cross-referenced with published proteomics and databases and performed experiments and analysed data for *A. thaliana*. S.A. designed the R script for transcriptomic analysis; L.G. helped acquiring images and analysing data for *M. truncatula*. R.G.P designed and tested the construct for Medtr1g073320, generated transgenic roots and imaged them. Y.B.A. was in charge of conceptualisation, supervision, data curation, and writing- reviewing and editing the manuscript with the help of P.K. All authors discussed the results, edited and commented on the manuscript.

Data Availability Statement

PIP1, the R-based pipeline described here, will be available for download along with associated database from GitHub (https://github.com/PhilPlantMan/PIP1) upon publication. Medtr1g073320 construct will be provided upon request.

References

Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S, Von Heijne G, Nielsen H. 2019. SignalP 5.0 improves signal peptide predictions using deep neural networks. *Nature Biotechnology* **37**(4): 420-423.

Amsbury S, Kirk P, Benitez-Alfonso Y. 2018. Emerging models on the regulation of intercellular transport by plasmodesmata-associated callose. *Journal of Experimental Botany* **69**(1): 105-115.

Benitez-Alfonso Y, Faulkner C, Pendle A, Miyashima S, Helariutta Y, Maule A. 2013. Symplasmic intercellular connectivity regulates lateral root patterning. *Developmental cell* **26**(2): 136-147.

Boisson-Dernier A, Chabaud M, Garcia F, Bécard G, Rosenberg C, Barker DG. 2001. Agrobacterium rhizogenes-transformed roots of Medicago truncatula for the study of nitrogen-fixing and endomycorrhizal symbiotic associations. *Molecular Plant-Microbe Interactions* **14**(6): 695-700.

Brault ML, Petit JD, Immel F, Nicolas WJ, Glavier M, Brocard L, Gaston A, Fouché M, Hawkins TJ, Crowet JM, et al. 2019. Multiple C2 domains and transmembrane region proteins (MCTP s) tether membranes at plasmodesmata. *EMBO reports* **20**(8).

Caillaud M-C, Wirthmueller L, Sklenar J, Findlay K, Piquerez SJM, Jones AME, Robatzek S, Jones JDG, Faulkner C. 2014. The Plasmodesmal Protein PDLP1 Localises to Haustoria-Associated Membranes during Downy Mildew Infection and Regulates Callose Deposition. *PLoS Pathogens* **10**(11): e1004496.

Calderan-Rodrigues MJ, Guimarães Fonseca J, De Moraes FE, Vaz Setem L, Carmanhanis Begossi A, Labate CA. 2019. Plant Cell Wall Proteomics: A Focus on Monocot Species, Brachypodium distachyon, Saccharum spp. and Oryza sativa. *International Journal of Molecular Sciences* **20**(8): 1975.

Carvalho BS, Irizarry RA. 2010. A framework for oligonucleotide microarray preprocessing. *Bioinformatics* **26**(19): 2363-2367.

Cheval C, Samwald S, Johnston MG, De Keijzer J, Breakspear A, Liu X, Bellandi A, Kadota Y, Zipfel C, Faulkner C. 2020. Chitin perception in plasmodesmata characterizes submembrane immune-signaling specificity in plants. *Proceedings of the National Academy of Sciences*: 201907799.

Crook AD, Schnabel EL, Frugoli JA. 2016. The systemic nodule number regulation kinase SUNN inMedicago truncatulainteracts with MtCLV2 and MtCRN. *The Plant Journal* **88**(1): 108-119.

Dorokhov YL, Ershova NM, Sheshukova EV, Komarova TV. 2019. Plasmodesmata Conductivity Regulation: A Mechanistic Model. *Plants* **8**(12): 595.

Dragićević MB, Paunović DM, Milica, Todorović SI, Simonović AD. 2020. ragp: Pipeline for mining of plant hydroxyproline-rich glycoproteins with implementation in R. *Glycobiology* **30**(1): 19-35.

Drost H-G, Paszkowski J. 2017. Biomartr: genomic data retrieval with R. *Bioinformatics*: btw821.

Emanuelsson O, Brunak S, Von Heijne G, Nielsen H. 2007. Locating proteins in the cell using TargetP, SignalP and related tools. *Nature protocols* **2**(4): 953.

Faulkner C, Petutschnig E, Benitez-Alfonso Y, Beck M, Robatzek S, Lipka V, Maule AJ. 2013. LYM2-dependent chitin perception limits molecular flux via plasmodesmata. *Proceedings of the National Academy of Sciences* **110**(22): 9166-9170.

Fernandez-Calvino L, Faulkner C, Walshaw J, Saalbach G, Bayer E, Benitez-Alfonso Y, Maule A. 2011. Arabidopsis plasmodesmal proteome. *PLOS one* **6**(4): e18880.

Gaudioso-Pedraza R, Beck M, Frances L, Kirk P, Ripodas C, Niebel A, Oldroyd GED, Benitez-Alfonso Y, de Carvalho-Niebel F. 2018. Callose-Regulated Symplasmic Communication Coordinates Symbiotic Root Nodule Development. *Curr Biol* **28**(22): 3562-3577 e3566.

Ghashghaie J, Brenckmann F, Saugier B. 1991. Effects of agar concentration on water status and growth of rose plants cultured in vitro. *Physiologia Plantarum* **82**(1): 73-78.

Gíslason MH, Nielsen H, Armenteros JJA, Johansen AR. 2019. Prediction of GPI-Anchored proteins with pointer neural networks. *bioRxiv*: 838680.

Gopal J, Iwama K. 2007. In vitro screening of potato against water-stress mediated through sorbitol and polyethylene glycol. *Plant Cell Rep* **26**(5): 693-700.

Grison MS, Kirk P, Brault ML, Wu XN, Schulze WX, Benitez-Alfonso Y, Immel F, Bayer EM. 2019. Plasma Membrane-Associated Receptor-like Kinases Relocalize to Plasmodesmata in Response to Osmotic Stress. *Plant Physiology* **181**(1): 142-160.

Gronnier J, Crowet J-M, Habenstein B, Nasir MN, Bayle V, Hosy E, Platre MP, Gouguet P, Raffaele S, Martinez D, et al. 2017. Structural basis for plant plasma membrane protein dynamics and organization into functional nanodomains. *eLife* **6**.

Gu Z, Eils R, Schlesner M. 2016. Complex heatmaps reveal patterns and correlations in multidimensional genomic data. *Bioinformatics* **32**(18): 2847-2849.

Haney CH, Long SR. 2010. Plant flotillins are required for infection by nitrogen-fixing bacteria. *Proceedings of the National Academy of Sciences* **107**(1): 478-483.

Hernández-Hernández V, Benítez M, Boudaoud A. 2019. Interplay between turgor pressure and plasmodesmata during plant development. *Journal of Experimental Botany*.

Howe KL, Contreras-Moreira B, Nishadi, Maslen G, Akanni W, Allen J, Alvarez-Jarreta J, Barba M, Bolser DM, Cambell L, et al. 2020. Ensembl Genomes 2020—enabling non-vertebrate genomic research. *Nucleic Acids Research* **48**(D1): D689-D695.

Huang D, Sun Y, Ma Z, Ke M, Cui Y, Chen Z, Chen C, Ji C, Tran TM, Yang L, et al. 2019. Salicylic acid-mediated plasmodesmal closure via Remorin-dependent lipid organization. *Proc Natl Acad Sci U S A* **116**(42): 21274-21284.

Hunter K, Kimura S, Rokka A, Tran HC, Toyota M, Kukkonen JP, Wrzaczek M. 2019. CRK2 Enhances Salt Tolerance by Regulating Callose Deposition in Connection with PLDα1. *Plant Physiology* **180**(4): 2004-2021.

Imlau A, Truernit E, Sauer N. 1999. Cell-to-Cell and Long-Distance Trafficking of the Green Fluorescent Protein in the Phloem and Symplasmic Unloading of the Protein into Sink Tissues. *The Plant Cell* **11**(3): 309-322.

Käll L, Krogh A, Sonnhammer EL. 2007. Advantages of combined transmembrane topology and signal peptide prediction—the Phobius web server. *Nucleic Acids Research* **35**(suppl_2): W429-W432.

Karimi M, Inzé D, Depicker A. 2002. GATEWAY™ vectors for Agrobacterium-mediated plant transformation. *Trends in plant science* **7**(5): 193-195.

Krogh A, Larsson B, Von Heijne G, Sonnhammer ELL. 2001. Predicting transmembrane protein topology with a hidden markov model: application to complete genomes11Edited by F. Cohen. *Journal of Molecular Biology* **305**(3): 567-580.

Lee JY, Wang X, Cui W, Sager R, Modla S, Czymmek K, Zybaliov B, van Wijk K, Zhang C, Lu H, et al. 2011. A plasmodesmata-localized protein mediates crosstalk between cell-to-cell communication and innate immunity in Arabidopsis. *Plant Cell* **23**(9): 3353-3373.

Leijon F, Melzer M, Zhou Q, Srivastava V, Bulone V. 2018. Proteomic analysis of plasmodesmata from populus cell suspension cultures in relation with callose biosynthesis. *Frontiers in Plant Science* **9**: 1681.

Li ZP, Paterlini A, Glavier M, Bayer EM. 2021. Intercellular trafficking via plasmodesmata: molecular layers of complexity. *Cellular and Molecular Life Sciences* **78**(3): 799-816.

Liu DY, Smith PM, Barton DA, Day DA, Overall RL. 2017. Characterisation of Arabidopsis calnexin 1 and calnexin 2 in the endoplasmic reticulum and at plasmodesmata. *Protoplasma* **254**(1): 125-136.

Liu N, Shen G, Xu Y, Liu H, Zhang J, Li S, Li J, Zhang C, Qi J, Wang L, et al. 2020. Extensive Inter-plant Protein Transfer between Cuscuta Parasites and Their Host Plants. *Molecular Plant* **13**(4): 573-585.

Mi H, Ebert D, Muruganujan A, Mills C, Albou L-P, Mushayamaha T, Thomas PD. 2021. PANTHER version 16: a revised family classification, tree-based classification tool, enhancer regions and extensive API. *Nucleic Acids Research* **49**(D1): D394-D403.

Ning W, Jiang P, Guo Y, Wang C, Tan X, Zhang W, Peng D, Xue Y. 2020. GPS-Palm: a deep learning-based graphic presentation system for the prediction of S-palmitoylation sites in proteins. *Briefings in Bioinformatics*.

Obayashi T, Aoki Y, Tadaka S, Kagaya Y, Kinoshita K. 2018. ATTED-II in 2018: A Plant Coexpression Database Based on Investigation of the Statistical Property of the Mutual Rank Index. *Plant and Cell Physiology* **59**(1): e3-e3.

Park S-H, Li F, Renaud J, Shen W, Li Y, Guo L, Cui H, Sumarah M, Wang A. 2017. NbEXPA1, an α-expansin, is plasmodesmata-specific and a novel host factor for potyviral infection. *The Plant Journal* **92**(5): 846-861.

Petit JD, Li ZP, Nicolas WJ, Grison MS, Bayer EM. 2020. Dare to change, the dynamics behind plasmodesmata-mediated cell-to-cell communication. *Current Opinion in Plant Biology* **53**: 80-89.

Pierleoni A, Martelli PL, Casadio R. 2008. PredGPI: a GPI-anchor predictor. *BMC bioinformatics* **9**(1): 392.

Reymond P, Kunz B, Paul-Pletzer K, Grimm R, Eckerskorn C, Farmer EE. 1996. Cloning of a cDNA encoding a plasma membrane-associated, uronide binding phosphoprotein with physical properties similar to viral movement proteins. *The Plant Cell* **8**(12): 2265-2276.

Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. 2015. limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Research* **43**(7): e47-e47.

Robinson RA, Stokes RH. 1949. Tables of osmotic and activity coefficients of electrolytes in aqueous solution at 25 C. *Transactions of the Faraday Society* **45**: 612-624.

Rosas-Diaz T, Zhang D, Fan P, Wang L, Ding X, Jiang Y, Jimenez-Gongora T, Medina-Puche L, Zhao X, Feng Z. 2018. A virus-targeted plant receptor-like kinase promotes cell-to-cell spread of RNAi. *Proceedings of the National Academy of Sciences* **115**(6): 1388-1393.

Sagi G, Katz A, Guenoune-Gelbart D, Epel BL. 2005. Class 1 reversibly glycosylated polypeptides are plasmodesmal-associated proteins delivered to plasmodesmata via the Golgi apparatus. *The Plant Cell* **17**(6): 1788-1800.

Simpson C, Thomas C, Findlay K, Bayer E, Maule AJ. 2009. An Arabidopsis GPI-anchor plasmodesmal neck protein with callose binding activity and potential to regulate cell-to-cell trafficking. *The Plant Cell* **21**(2): 581-594.

Thomas CL, Bayer EM, Ritzenthaler C, Fernandez-Calvino L, Maule AJ. 2008. Specific targeting of a plasmodesmal protein affecting cell-to-cell communication. *PLoS biol* **6**(1): e7.

Vatén A, Dettmer J, Wu S, Stierhof Y-D, Miyashima S, Yadav SR, Roberts CJ, Campilho A, Bulone V, Lichtenberger R. 2011. Callose biosynthesis regulates symplastic trafficking during root development. *Developmental cell* **21**(6): 1144-1155.

Vu MH, Iswanto ABB, Lee J, Kim J-Y. 2020. The Role of Plasmodesmata-Associated Receptor in Plant Development and Environmental Response. *Plants* **9**(2): 216.

Wang X, Sager R, Cui W, Zhang C, Lu H, Lee JY. 2013. Salicylic Acid Regulates Plasmodesmata Closure during Innate Immune Responses in Arabidopsis. *The Plant Cell* **25**(6): 2315-2329.

Wilson AK, Pickett FB, Turner JC, Estelle M. 1990. A dominant mutation inArabidopsis confers resistance to auxin, ethylene and abscisic acid. *Molecular and General Genetics MGG* **222**(2-3): 377-383.

Xie Y, Zheng Y, Li H, Luo X, He Z, Cao S, Shi Y, Zhao Q, Xue Y, Zuo Z, et al. 2016. GPS-Lipid: a robust tool for the prediction of multiple lipid modification sites. *Scientific Reports* **6**(1): 28249.

Xue Y, Liu Z, Gao X, Jin C, Wen L, Yao X, Ren J. 2010. GPS-SNO: Computational Prediction of Protein S-Nitrosylation Sites with a Modified GPS Algorithm. *PLOS one* **5**(6): e11290.