The gut parasite *Nosema ceranae* impairs olfactory learning in bumblebees

Tamara Gómez-Moracho[1,*,a], Tristan Durand[1,*], Mathieu Lihoreau[1]

[1] Research Center on Animal Cognition (CRCA), Center for Integrative Biology (CBI); CNRS, University Paul Sabatier, Toulouse, France

* These authors contributed equally to this work

[a] Author for correspondence: Tamara Gómez-Moracho (tamara.gomez@univ-tlse3.fr)

Abstract

Pollinators are exposed to numerous parasites and pathogens when foraging on flowers. These biological stressors may affect critical cognitive abilities required for foraging. Here, we tested whether exposure to *Nosema ceranae*, one of the most widespread parasite of honey bees also found in wild pollinators, impacts cognition in bumblebees. We investigated different forms of olfactory learning and memory using conditioning of the proboscis extension reflex. Seven days after feeding parasite spores, bumblebees showed lower performance in absolute and differential learning, and reversal learning than controls. Long-term memory was also slightly reduced. The consistent effect of *N. ceranae* exposure across different types of olfactory learning indicates that its action was not specific to particular brain areas or neural processes. We discuss the potential mechanisms by which *N. ceranae* impairs bumblebee cognition and the broader consequences for populations of pollinators.

Keywords : Bumble bees ; *Bombus terrestris*; learning and memory ; PER

1. Introduction

Pollinators, such as bees, rely on a rich cognitive repertoire to collect pollen and nectar on flowers. These include associative learning and memories of floral traits like odours, shapes, colours, and textures, to identify profitable resources (Giurfa, 2015; Menzel, 2012), and spatial cues to navigate (Collett et al., 2013). Any disruption of these cognitive abilities by environmental stressors can considerably reduce the foraging performances of bees, ultimately compromising brood development and survival (Klein et al., 2017).

In particular, foraging bees are exposed to a number of parasites that can affect their physiology and behaviour (Gómez-Moracho et al., 2017). The microsporidia *Nosema ceranae* is one of the most prevalent parasites of bees worldwide with a large range of hosts including honey bees (Higes et al., 2006), bumblebees (Plischuk et al., 2009), solitary bees (Ravoet et al., 2014), but also other flower visitors like wasps (Porrini et al., 2017). Insects get infected by ingesting parasite spores from contaminated water or pollen (Higes et al., 2008), or during physical contacts with contaminated individuals (Smith, 2012). The spores invade the gut epithelial cells of the bee where they develop (Holt et al., 2013). *Nosema* degenerates the gut epithelium of the host (Higes et al., 2007), alters its metabolism (Mayack and Naug, 2009) and disrupts the immune response (Antúnez et al., 2009). In honey bees, this causes a disease (nosemosis) believed to contribute to colony collapse disorder (Cox-Foster et al., 2007).

Infected honey bees also show impaired navigation (Wolf et al., 2014) and increased flight activity (Dussaubat et al., 2013) suggesting their cognitive abilities are affected. Few studies have explored this possibility and their results so far are mixed (Bell et al., 2020; Charbonneau et al., 2016; Gage et al., 2018; Piiroinen et al., 2016; Piiroinen and Goulson, 2016). Most of them were conducted on honey bees using Pavlovian olfactory conditionings of the proboscis extension reflex (PER) in which harnessed bees are trained to associate an odour, or a combination of odours, with a sucrose reward (Takeda, 1961). Honey bees showed reduced absolute learning (when an odour is paired to a reward) between days 7 and 23 after parasite exposure ((Bell et al., 2020; Gage et al., 2018; Piiroinen and Goulson, 2016); but see (Charbonneau et al., 2016)). Short-term memory was impaired after 7 and 15 days (Gage et al., 2018), but not after 9 (Piiroinen and Goulson, 2016) or 23 days (Bell *et al.*, 2020). Long-

term memory was affected after 7 days but not after 15 days (Gage et al., 2018). Only two studies explored these effects on bumblebees. One suggests a slight impairment of absolute learning (Piiroinen and Goulson, 2016), and both report an absence of effect on memory (Piiroinen et al., 2016, Piiroinen and Goulson 2016). Note however that in these two studies less than 3% of the bumblebees exposed to the parasite showed signs of infection after the behavioural tests (i.e. PCR positive).

Given the expanding geographical distribution of *N. ceranae* worldwide (Klee et al., 2007) and its prevalence in domesticated and wild bees (Plischuk et al., 2009; Porrini et al., 2017; Ravoet et al., 2014), clarifying its influence on host cognition has become an urgent matter. In particular, other critical forms of learning, such as the ability to associate one of two odours with a reward (differential learning) and reverse this association (reversal learning) have so far been unexplored. These types of learning are essential to discriminate flowers. They involve different brain centers and may thus be more or less impacted by the parasite if its effect is specific (Boitard et al., 2015; Giurfa and Sandoz, 2012).

Here we build on a recently established method yielding high the rates of experimental infection by *N. ceranae* (Gómez-Moracho et al., 2021) to study the impact of the parasite on different cognitive tasks in bumblebees. We used PER conditioning to compare the olfactory learning and memory performances of control bumblebees, bumblebees exposed to the parasite, and bumblebees contaminated by the parasites (PCR positive) at seven days post exposure.

2. Material and methods

2.1. Bumblebees

We used bumblebees (*B. terrestris*) from 14 commercial colonies acquired from Biobest (Westerlo, Belgium). Before the experiments, we verified the absence of *N. ceranae* (Martín-Hernández *et al.*, 2007), and other common parasites (*N. bombi* (Klee *et al.*, 2006); *Crithidia bombi* (Schmid-Hempel and Tognazzo, 2010)) in a PCR using 15 bumblebees from each colony. We maintained bumblebees in their original colonies with *ad libitum* access to the

syrup provided by the manufacturer and germ-free pollen (honey bee collected pollen exposed
to UV light for 12 hours), in a room at 25±1°C under a 12 h light:12 h dark photocycle.

2.2.   *N. ceranae* spores

We obtained fresh spores from naturally infected honey bee colonies (*Apis mellifera*)
maintained at our experimental apiary (University Toulouse III, France). To prepare spore
solutions, we dissected the gut of 15 honey bees and crushed them in 15 mL of distilled H_2O.
We confirmed by PCR the presence of *N. ceranae* and the absence of *N. apis* (another
common parasite of honey bees) in each homogenate (Martín-Hernández et al., 2007), and
purified them following standard protocols (Fries et al., 2013). We centrifuged homogenates
in aliquots of 1 mL at 5,000 rpm for 5 minutes and re-suspended the pellet in 500 µL of dH_2O
by vortexing. This was repeated three times to obtain a spore solution of 85% purity (Fries et
al., 2013). We counted *N. ceranae* spores in an improved Neubauer haemocytometer
(Cantwell, 1970) in a light microscope (x400) and adjusted the spore inoculum to 15,000
spores/µL in 20% (w/w) of sucrose solution. Spore solutions were used within the same week
they were purified.

2.3.   Parasite exposure and experimental conditions

We exposed bumblebees to *N. ceranae* as described in Gómez-Moracho *et al.*, (2021).
Briefly, we confined individual bumblebees in a Petri dish during 5 h without food. We then
exposed some bumblebees to a 20 µL drop of inoculum containing 300,000 *N. ceranae*
spores. Control bumblebees received 20 µL of sucrose solution (20% w/w). We only used
bumblebees that consumed the entire drop of sucrose within the next 2 h. We then allocated
bumblebees into microcolonies of 20-25 individuals, containing a gravity feeder with *ad*
libitum access to food (Kraus et al., 2019). Since diet can affect host-parasites relationships
(Frost et al., 2008) we provided bumblebees an artificial diet with a protein to carbohydrates
ratio of 1:207 previously shown to elicit highest *N. ceranae* prevalence in bumblebees
(Gómez-Moracho et al., 2021). The diet was made with a fixed total amount of nutrients of
170 g/L (protein + carbohydrates) and 0.5% of vitamin mixture for insects (Sigma, Germany).

Carbohydrates were supplied as sucrose (Euromedex, France). Proteins consisted in a mixture

of casein and whey (4:1) (Nutrimuscle, Belgium) (Gómez-Moracho et al., 2021). We kept

bumblebee microcolonies in a room at 25±1 °C with a 12 h light: 12 h dark photoperiod until

the behavioural tests. Every day, we renewed the diet and removed dead bumblebees.

2.4.   Behavioural experiments

We tested the cognitive performances of bumblebees using PER at day 7 after parasite

exposure. The day before the behavioural tests we kept diet to low levels (~200

µL/bumblebee) to keep bumblebees motivated for PER experiments. Three hours before the

behavioural tests we collected bumblebees, chilled them in ice for 5 min and restrained them

in a modified 2 mL Eppendorf tube (hereafter, capsule) that we cut in length to fit each

bumblebee (adapted from Toda *et al.*, (2009); Figure 1A). Bumblebees were tested in the

horizontal position and could slightly move inside the tube. We found these conditions better

suited to perform PER experiments with bumblebees than the classical vertical harnessing

used for honey bees (Giurfa and Sandoz, 2012), in which bumblebees appeared paralysed

(unpublished data). Once in the capsule, we kept the bumblebees in the dark, in an incubator

at 28ºC, with no access to food. All bumblebees that finished the conditioning protocols were

kept at -20°C for later analyses of their infection status through PCR.

2.5.   Sucrose responsiveness

We tested the sucrose responsiveness of bumblebees to control for potential influences of *N.*

ceranae on their reward perception or feeding motivation. We presented seven sucrose

solutions to each bumblebee, from concentrations of 0% (pure water) to 60% (w/w), with

increments of 10% (Graystock et al., 2013). For each concentration, we touched the antennae

of the bumblebee with a toothpick soaked in the corresponding sucrose solution to elicit PER.

We presented solutions in an increasing concentration gradient with an inter-trial interval of 5

minutes between concentrations. We scored a positive response if bumblebees extended their

proboscis after a solution presentation. We discarded bumblebees responding to water (i.e. 0%

sucrose solution) to avoid the effect of thirst on sucrose responsiveness (Baracchi et al.,

2018). The sum of the number of PER responses divided by the number of concentration
tested formed the individual gustatory score (Scheiner et al., 2013).

2.6.   Conditioning experiments

All experiments shared the same general protocol (Figure S1). An encapsulated bumblebee
(Figure 1A) was placed 1.5 cm ahead of an automated conditioning setup (described in
(Aguiar et al., 2018)) delivering a continuous stream of odourless air at 1.2 ml/s to which
specific odours were selectively added (Raiser et al., 2017). We used two odorants as
conditioned stimulus (CS): nonanal and phenylacetaldehyde (Palottini et al., 2018), in a 1:100
dilution in mineral oil. Before conditioning, we tested the responsiveness of bees to sucrose
by touching both antennae with a toothpick soaked in 50% (w/w) sucrose solution without
allowing them to lick. Bumblebees extending their proboscis were considered motivated and
kept for the experiments. Conditioning trials (Figure S1c) consisted in 15 seconds of
odourless airflow, followed by 6 seconds of CS, and 3 seconds of unconditioned stimulus
(US) (i.e. 50% sucrose solution applied with a toothpick on the bumblebee's antennae), with 2
seconds of overlap between CS+US, and 20 seconds of odourless airflow (Aguiar et al.,
2018). The inter-trial interval was 10 minutes. An air extractor was placed behind the
bumblebee to prevent odorant accumulation during CS delivery. Bumblebees extending their
proboscis within 3 seconds of US presentation (i.e. 2s CS+US and 1s US) were allowed to
lick the toothpick soaked in sucrose (50%, w/w). In unrewarded trials (see reversal learning
and memory tests) no US was applied. We scored a conditioned response if the bumblebee
extended its proboscis to the odour delivery before sucrose presentation (Figure S1c).
Bumblebees that responded to the odour in the first conditioning trial were discarded from the
analyses. We used conditioned responses to calculate three individual scores for each
bumblebee, describing its performance during conditioning (i.e. acquisition score), at the end
of conditioning (i.e. learning score) or during memory retrieval (i.e. memory score)
(Monchanin et al., 2020; see details below). Exposed and control bumblebees were always
conditioned in parallel.

2.6.1. Absolute learning, short-term memory and long-term memory

We tested the effects of *N. ceranae* on the ability of bumblebees to associate an odour with a reward. This form of learning only requires peripheral brain centers, i.e. the antennal lobes (Giurfa and Sandoz, 2012). We trained bumblebees in a spaced 3-trial absolute conditioning learning (Figure S1a) that was shown to generate robust long-term memory in bees (Menzel et al., 2001). We used the same rewarded odour (CS+) during training of a given bumblebee, but both nonanal and phenylacetaldehyde were used as a CS+ for different bumblebees. Only bumblebees that responded to the US in the three trials (i.e. efficient bumblebees) were kept for the analyses (Table S2). For each of these bees we calculated an acquisition score (sum of PER responses (i.e. 0-2) divided by the number of trials (i.e. 3)), and a learning score (1 if they responded to CS+ in the last trial or 0).

We tested memory retention in efficient bumblebees that made at least one conditioned response in either one of the last two trials. We performed tests either 1 h (i.e. short-term memory; STM) or 24 h (i.e. long-term memory; LTM) after the last acquisition trial. For tests performed after 24 h, bumblebees were fed until satiation with 50% (w/w) sucrose solution right after conditioning. In bees, LTM is dependent on protein synthesis whereas STM is not (Menzel, 2001). Studying these two types of memories was thus a mean to explore whether exposure to *N. ceranae* interfered with protein synthesis. We presented bumblebees the two odorants without any reward: the odour used as a CS+ to test for memory formation, and the second odour as a novel odorant (NOd), to control for potential generalization (Villar et al., 2020). For example, when nonanal was used as CS+, phenylacetaldehyde was used as a NOd, and *vice versa*. Memory formation was considered when bumblebees extended their proboscis in response to the CS+ but not to the NOd, and thus had a memory score of 1 (0 otherwise). Just after the memory test, we tested the motivation of bumblebees by touching their antennae with a toothpick soaked with 50% (w/w) sucrose solution. Bumblebees that did not respond to the US were discarded for the analyses (i.e. 14.06% in LTM; Table S2).

2.6.2. Reversal learning

We tested the effects of *N. ceranae* on the ability of bumblebees to learn to discriminate two
odours and reverse the task. This form of learning involves a first phase of differential
conditioning that requires the antennal lobes but is not dependant on central brain centers, i.e.
mushroom bodies, and a second phase of reversal learning that requires both the antennal
lobes and the mushroom bodies (Boitard et al., 2015). In the differential learning phase (phase
1), we trained bumblebees to discriminate between two odours. This phase consisted in 10-
trial blocks (Figure S1b), five with each odour that was either paired with the US (A+) or
unpaired (B-), presented in a pseudo-random order. The rewarded and unrewarded odours
were randomized on different training days. Bumblebees that did not respond to US in two or
more trials were discarded for the analyses and for reversal phase. In the reversal learning
phase (phase 2) we trained bumblebees to invert the first learnt contingency. The reversal
phase started 1h after the end of the differential phase. Here we trained bumblebees in 12
trial-blocks, six with each odour presented in a pseudo-random order. The previously
rewarded odour was not associated with a reward anymore (A-) while the previously
unrewarded odour became rewarded (B+). To start with the same level of learning, we
analysed reversal learning in bumblebees that responded to A- either in the first and/or second
presentation of this odour during the reversal phase (Table S3). We analysed the performance
of bumblebees in each phase separately by attributing them an acquisition score (sum of all
trials where the bee responded to CS+ but not to CS-, divided by the number of trials) and a
learning score (1 if the bumblebee responded to CS+ but not to CS- in the last trial) for each
phase.

2.7.   Infection status

We assessed the infection status of bumblebees that finished the tests in a PCR using the
primers 218MITOC (Martín-Hernández et al., 2007). For each bumblebee, the entire gut was
extracted, homogenised in sterile dH2O and vortexed with 2 mm glass beads (Labbox
Labware, Spain). Genomic DNA was extracted using Proteinase K (20 mg/mL; Euromedex,
France) and 1 mM of Tris-EDTA Buffer (pH = 8). A sample with *N. ceranae* spores was
included in each round of extraction as positive control. PCRs were performed with the Taq
Polymerase Direct Loading Buffer (5 U/µL; MP Biomedicals, CA) following manufacturer's

instructions. We used a final volume of 25 µL with 0.4 µM of each pair of primers (Martín-
Hernández et al., 2007), 200 µM of dNTPs (Jena Biosciences, Germany), 0.48 µg/µL of BSA
(Sigma, Germany) and 2.5 µL of DNA sample. PCR reactions were carried out in a S1000™
Thermal Cycler (Biorad, CA). Thermal conditions were 94 °C for 2 min, 35 cycles of 94 °C
for 30 s, 61.8ºC for 45 s and 72 °C for 2 min, and a final step of 72 °C for 7 min. The length
of PCR products (i.e. 218 pb) was checked in a 1.2% agarose gel electrophoresis stained with
SYBR Safe DNA Stain (Edvotek, Washington DC). Positive and negative controls of PCR
were run in parallel. Based on the PCR results we classified bumblebees in three different
infection statuses, i.e. control, exposed negative or exposed positive. Bumblebees that were
not exposed to the parasite but nevertheless showed a positive result in PCR were excluded
from the analyses (i.e. 6.26%; 23 out of 367 control bumblebees; Tables S1, S2 and S3).

2.8.   Statistical analyses

All analyses were conducted in RStudio (version 1.4.1106). We evaluated the effect of
parasite exposure and infection on gustatory, acquisition, learning and memory scores. The
gustatory scores and the acquisition scores (i.e. absolute and reversal conditioning) were
standardized and analysed in a Generalized Linear Mixed Model (GLMM) (package
[glmmTMB]; (Brooks et al., 2017)). The response to the first and last sucrose concentrations,
as well as the learning and memory scores were analysed in a binomial GLMM (package
[lme4]; (Bates et al., 2015)). In all models, bee infection status was used as a fixed factor,
while bee identity and odorant were used as random factors. We performed Tukey post-hoc
pairwise comparisons (package [multcomp]; (Hothorn et al., 2008)) to assess the relationship
between the three bee statuses. Raw data are available online (doi:10.5281/zenodo.4376362).

3. Results

3.1.   Parasite exposure did not influence sucrose responsiveness

We tested responsiveness to different sucrose concentrations in 73 bumblebees (43 controls,
20 exposed negative, 10 exposed positive; Table S1). Overall, 71.23% of the bumblebees

responded to sucrose (Figures 1B and C). This percentage was similar across the three
infection statuses (X^2 = 2.236, df = 2, p = 0.326). Bumblebees of the three infection statuses
responded similarly to the lowest sucrose concentration (Figure 1 B; GLMM, X^2 = 0.920, df =
2, p = 0.631), reached similar gustatory scores (Figure 1C; GLMM, X^2 = 1.686, df = 2, p =
0.430) and similar final level of response to the highest sucrose concentration (Figure 1B;
GLMM, X^2 = 0.217, df = 2, p = 0.897). Therefore, exposure to *N. ceranae* affected neither the
sucrose sensitivity nor the feeding motivation of bumblebees.

Figure 1. Odour delivery set up and sucrose responsiveness. A) Photo of a bumblebee
during odour conditioning. The bumblebee is placed inside the capsule in front of the air
delivery set up. After the odour is delivered, sucrose is presented with a toothpick to the
bumblebees which extends its proboscis to drink the reward. **B)** Proportion of control (blue),
exposed negative (yellow) and exposed positive (red) bumblebees responding to an increase
gradient of sucrose concentrations. Pie chart represents the percentage of exposed negative
and exposed positive bees (n total=30). **C)** Violin plots show gustatory score of bumblebees
as the sum of all responses for each bumblebee. Black diamonds represent the mean score for
each infection status. White dots represent the score of each individual. n is the sample size.
Letters above violin plots represent significant differences between status (GLMM; p < 0.05).
n is the sample size.

3.1.   Parasite exposure reduced absolute learning but not memory

We analysed absolute conditioning in 420 bumblebees (141 controls, 228 exposed negative,
51 exposed positive; Table S2). The proportion of bumblebees showing a conditioned
response increased over trials in the three infection statuses (Figure 2A). Exposed bumblebees
(either negative or positive) had significantly lower acquisition scores (Figure 2B; GLMM,
status: $X^2 = 31.558$, df = 2, p < 0.001) and learning scores (Figure 2C; GLMM, status, $X^2 = $
16.037, df = 2, p < 0.001) than controls. Both acquisition and learning scores were similar for
negative and positive exposed bumblebees (Tukey p > 0.05; Table S4).

Figure 2**. Absolute learning and memory. A)** Learning curves show the proportion of
control (blue), exposed negative (yellow) and exposed positive (red) bumblebees extending
their proboscis to the odour during conditioning. Pie chart shows the percentage of negative
and positive exposed bumblebees that finished the conditioning (N total = 279). **B-C)** Violin
plots for acquisition score (B; sum of correct responses divided by the number of trials for
each bumblebee) and learning score (C; correct response at the last trial). Black diamonds
represent the mean score for each infection status. White dots represent the score of each
individual. **D)** Short-term memory. Bar plots show the proportion of bumblebees responding
to odour 1 h after training. **E)** Long-term memory. Bar plots show the proportion of

bumblebees responding to odour 24 h after training. Letters above violin plots and bar plots

represent significant differences between status (GLMM; p < 0.05). n is the sample size.

We analysed short-term and long-term memory formation of efficient bumblebees that

responded to odour at least once in any of the last two last trials (41.52% in short-term

memory and 50% in long-term memory; Table S2). A high proportion of the bumblebees

(82.02%±0.011, mean±SE) remembered the rewarded odour 1h after training. This proportion

was similar in the three infection statuses (GLMM, status: $X^2 = 0.013$, df = 2, p = 0.993),

indicating that *N. ceranae* did not affect short-term memory (Figure 2D). A much lower

proportion of the bumblebees (34.35%±0.06, mean ±SE) remembered the rewarded odour

after 24 h. Exposed bumblebees (either negative or positive) showed a lower proportion of

correct responses than controls, although this was not significantly different (Figure 2E;

GLMM, status: $X^2 = 2.265$, df = 2, p = 0.322), presumably due to the low amount of exposed

positive individuals (9 bumblebees).

3.2.   Parasite exposure reduced differential and reversal learning

We analysed differential learning in 125 bumblebees (64 controls, 41 exposed negative, 20

exposed positive; Table S3). Bumblebees in the three infection statuses increased their

response to A+ over trials, but not to B- (Figure 3A), indicating that they were equally able to

discriminate the odours. However, parasite exposure affected acquisition scores (GLMM,

status, $X^2 = 22.039$, df = 2, p < 0.001; Figure 3B). Exposed bumblebees (positive or negative)

showed similar acquisition scores (Tukey test: p > 0.05; Table 1) and these scores were

significantly lower than the scores of controls (Tukey test: p < 0.05; Table 1). Parasite

exposure also affected the final level of learning (GLMM, status, $X^2 = 11.168$, df = 2, p =

0.003; Figure 2C). Exposed positive bumblebees had the lowest learning score. Their score

was significantly lower than that of controls (Tukey test: p = 0.002; Table 1) but similar to

that exposed negative bumblebees (Tukey test: p = 0.133; Table 1). Thus overall, exposure to

N. ceranae reduced differential learning performances.

Figure 3. Reversal learning. **A-C)** Differential learning phase. **A)** Percentage of PER responses to rewarded (A+, circle) and unrewarded (B-, square) odours by control (blue), exposed negative (yellow) and exposed positive (red) bumblebees. Pie chart shows the percentage of exposed negative and positive bumblebees that finished conditioning (n total = 67). Violin plots of **B)** acquisition scores (i.e. sum of the correct responses divided by the number of trials for each bee) and **C)** learning scores (i.e. performance of bumblebees at last trial) of bumblebees with different infection statuses. Black diamonds represent the mean score of each status. White dots are the scores for each individual. **D-F)** Reversal learning phase. **D)** Curves show the increase in the percentage of PER response to B+ over A- over trials. **E)** Acquisition score. **F)** learning scores. Letters above violin plots show significant differences between infection statuses in the acquisition and learning scores (GLMM, p < 0.05). n is the sample size.

We analysed reversal learning in 108 bumblebees that finished the differential phase, as they responded to A- in any of the two first presentations of this odour during reversal phase (56

controls, 36 exposed negative, 16 exposed positive; Table S3). All bumblebees reduced their

response to A- in favour of B+ over trials (Figure 3D). However, bumblebees with different

infection statuses reached significantly different acquisition scores (Figure 3E; GLMM,

status, X^2 = 6.042, df = 2, p = 0.048). Exposed positive bumblebees showed slightly lower

acquisition scores than controls (Tukey test: p = 0.049; Table 1) and exposed negative

bumblebees (Tukey test: p = 0.056, Table 1), suggesting they had a lower ability to reverse

the task. By the end of training, bumblebees of all infection statuses reversed the task, as

shown by the lack of difference between their learning scores (Binomial GLMM, status: X^2 =

3.073, df = 2, p = 0.215). Thus overall, exposure to *N. ceranae* also impaired the reversal

phase of reversal learning. We found no effect of *N. ceranae* exposure on either phase of

reversal learning when bumblebees were tested at 2 days post exposure, suggesting that stress

due to parasite exposure or parasite infection requires longer time to be established (Text S1).

4. Discussion

Bees are exposed to a number of parasites that can affect cognitive abilities supporting crucial

behaviour (Koch et al., 2017; Schmid-Hempel, 2013). Previous studies exploring the effect of

N. ceranae on absolute olfactory learning and memory in bees reported contrasting results,

presumably because of differences in conditioning protocols and infection rates across studies

(Bell et al., 2020; Charbonneau et al., 2016; Gage et al., 2018; Piiroinen et al., 2016; Piiroinen

and Goulson, 2016). Here, we ran a suite of standard olfactory cognitive assays showing that

feeding bumblebees spores of this parasite consistently impairs different types of olfactory

learning but not memory seven days after exposure. This suggests that the influence of *N.*

ceranae exposure on bumblebee cognition is not specific to a particular brain area.

Exposure to *N. ceranae* in food clearly impaired the ability of bumblebees to associate an

odour with a reward (absolute learning), discriminate two odours (differential learning), and

learn an opposite association (reversal learning). These are fundamental cognitive operations

a bee must display to efficiently forage on flowers (Giurfa and Sandoz, 2012). This result

agrees with a previous study reporting a reduced absolute learning in *N. ceranae* exposed

bumblebees (Piiroinen and Goulson, 2016). Here we provide evidence that these reduced

cognitive performances were not due to an effect of the parasite on odour perception or
motivation to feed, and give insight on where the parasite might act in the nervous system. In
bees, sucrose perception through the antennae, olfactory absolute learning, differential
learning and reversal learning all require processing of olfactory information through the
antennal lobes (Giurfa and Sandoz, 2012) while reversal learning also requires functional
mushroom bodies (Boitard et al., 2015). The fact that all types of learning were impaired and
that sucrose sensitivity was not, indicates that *N. ceranae* did not specifically target the
antennal lobes or the mushroom bodies. Rather, it likely impacted the learning processes in
general.

By contrast, we found no evidence that *N. ceranae* influenced memory. During training
animals learn and form short-term memories that are later consolidated and transformed into
stable long-term memories (Menzel and Muller, 1996) after protein synthesis (Menzel, 2001).
In our experiments, *N. ceranae* neither impaired short-term nor long-term memory. Nearly all
exposed bumblebees (either positive or negative in PCR) remembered the association of the
odour with the reward 1 h after training. However, they showed slightly lower performance
than controls at 24 h. This tendency was not significant and may result from low sample size
as only 29.44% of the efficient exposed bumblebees were eligible to analyse their memory
(i.e. efficient bumblebees, with at least one conditioned response during learning, that
responded to sucrose after memory test).

It has recently been questioned whether bumblebees are natural hosts of *N. ceranae* based on
the lack of evidence of parasitic forms inside host cells (Gisder et al., 2020). Several studies
have nevertheless reported *N. ceranae* in wild bumblebees at low (e.g. 4.76%; (Sinpoo,
2018)) and high prevalence (e.g. 72%; (Arbulo et al., 2015)). Whether or not bumblebees are
suitable hosts for *N. ceranae* replication is confirmed in future studies, our results imply that
bumblebees are strongly impacted by an acute exposure to the parasite. Such exposure may be
extremely frequent in nature due to the high prevalence of *N. ceranae* in honey bees (Runckel
et al., 2011)) that contaminate flowers with spores through physical contact or in their faeces
(Graystock et al., 2015). Our protocol of parasite exposure significantly increased the
infection rate of bumblebees to 28% in comparison to previous studies (Piiroinen et al., 2016;
Piiroinen and Goulson, 2016), which allowed the evaluation of cognitive traits in bumblebees

in the three infection statuses. Bumblebees that tested positive to *N. ceranae* showed a
tendency for lower cognitive performances than negative exposed bumblebees. They reached
the lowest learning during the absolute conditioning and barely discriminated odours,
suggesting that infection may interfere with some aspects of cognition, however further
experiments are needed to tackle this specific question.

Through which mechanism may the parasite impair learning? Ingestion of *N. ceranae* spores
exerts a stress that can reduce cognition. *N. ceranae* is known to alter the immune system of
bees, for example by modulating the expression of antimicrobial peptides (Antúnez et al.,
2009; Botías et al., 2020; Sinpoo, 2018). Stimulation of the immune system with non-
pathogenic elicitors, as lipopolysaccharides (LPS), was shown to reduce learning abilities in
honey bees, that were less able to associate an odour with a reward (Mallon et al., 2003), and
bumblebees, that showed lower performances in odour (Mobley and Gegear, 2018) and colour
differential learning tasks (Mobley and Gegear, 2018). It is thus possible that the observed
effects of *N. ceranae* exposure on bumblebee cognition were caused by an activation of the
immune response. In honey bees, infection with *N. ceranae* was shown to downregulate the
expression of genes in the brain (Doublet et al., 2016), some of which are linked to olfaction
(Badaoui et al., 2017; Doublet et al., 2016), potentially leading to changes in behaviour and
cognition. Interestingly, none of these effects were observed at two days post exposure (Text
S1). Whether *N. ceranae* triggers the immune response at this time is unknown. In honey
bees, the earliest effects described so far were observed after three days (Chaimanee et al.,
2012).

Since olfactory learning is essential for foraging, this sublethal effect of *N. ceranae* exposure
on bumblebee cognition can compromise colony foraging and development. Commercial and
wild bumblebee colonies exhibit physiological and behavioural differences as a result of
different selective pressures (Velthuis and Doorn, 2006), and may, therefore, show different
susceptibility to parasites. Parasite loads in the field can range from a few to thousands spores
(Meana et al., 2010). Here we used a substantially higher spore loads of *N. ceranae* to infect
commercially reared bumblebees than the actual infection rates found in wild bumblebees
(e.g. 6800 spores per individual (Graystock et al., 2013)). Further studies are thus needed to
analyse the effects of different concentrations of *N.* ceranae spores, and their possible

interactions with other stressors in the field in order to assess their real impact on wild
pollinators.

5. Acknowledgements

We thank Philipp Heeb, Cristian Pasquaretta and Maria Eugenia Villar for constructive
discussions, Charline Schartier for helping running some experiments, and Martin Giurfa for
lending the automated PER conditioning setup.

6. Funding
TGM was funded by a postdoctoral grant of the Fyssen foundation. While writing, TGM, TD
and ML were funded by grants of the Agence Nationale de la Recherche (POLLINET ANR-
16-CE02-0002-01, 3DNaviBee ANR-19-CE37-0024, BEE-MOVE ANR-20-ERC8-0004-01),
the European Regional Development Fund (ECONECT), and the Agence de la Transition
Ecologique (LOTAPIS) to ML.

7. References

Aguiar, J.M.R.B.V., Roselino, A.C., Sazima, M., Giurfa, M., 2018. Can honey bees
discriminate between floral-fragrance isomers? J. Exp. Biol. 221, jeb180844.
https://doi.org/10.1242/jeb.180844

Antúnez, K., Martín-Hernández, R., Prieto, L., Meana, A., Zunino, P., Higes, M., 2009.
Immune suppression in the honey bee (*Apis mellifera*) following infection by *Nosema*
ceranae (Microsporidia). Environ. Microbiol. 11, 2284–2290. https://doi.org/10.1111/j.1462-
2920.2009.01953.x

Arbulo, N., Antúnez, K., Salvarrey, S., Santos, E., Branchiccela, B., Martín-Hernández, R.,
Higes, M., Invernizzi, C., 2015. High prevalence and infection levels of *Nosema ceranae* in
bumblebees *Bombus atratus* and *Bombus bellicosus* from Uruguay. J. Invertebr. Pathol. 130,

165–168. https://doi.org/10.1016/j.jip.2015.07.018

Badaoui, B., Fougeroux, A., Petit, F., Anselmo, A., Gorni, C., Cucurachi, M., Cersini, A.,
Granato, A., Cardeti, G., Formato, G., Mutinelli, F., Giuffra, E., Williams, J.L., Botti, S.,
2017. RNA-sequence analysis of gene expression from honeybees (*Apis mellifera*) infected
with *Nosema ceranae*. PLOS ONE 12, e0173438.
https://doi.org/10.1371/journal.pone.0173438

Baracchi, D., Rigosi, E., de Brito Sanchez, G., Giurfa, M., 2018. Lateralization of sucrose
responsiveness and non-associative learning in honeybees. Front. Psychol. 9, 425.
https://doi.org/10.3389/fpsyg.2018.00425

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects models
using **lme4**. J. Stat. Softw. 67. https://doi.org/10.18637/jss.v067.i01

Bell, H.C., Montgomery, C.N., Benavides, J.E., Nieh, J.C., 2020. Effects of *Nosema ceranae*
(Dissociodihaplophasida: Nosematidae) and flupyradifurone on olfactory learning in honey
bees, *Apis mellifera* (Hymenoptera: Apidae). J. Insect Sci. 20.
https://doi.org/10.1093/jisesa/ieaa130

Boitard, C., Devaud, J.-M., Isabel, G., Giurfa, M., 2015. GABAergic feedback signaling into
the calyces of the mushroom bodies enables olfactory reversal learning in honey bees. Front.
Behav. Neurosci. 9, 1–13. https://doi.org/10.3389/fnbeh.2015.00198

Botías, C., Jones, J.C., Pamminger, T., Bartomeus, I., Hughes, W.O.H., Goulson, D., 2020.
Multiple stressors interact to impair the performance of bumblebee (*Bombus terrestris*)
colonies. J. Anim. Ecol. 1365-2656.13375. https://doi.org/10.1111/1365-2656.13375

Brooks, M., E., Kristensen, K., Benthem, K., J. ,van, Magnusson, A., Berg, C., W., Nielsen,
A., Skaug, H., J., Mächler, M., Bolker, B., M., 2017. glmmtmb balances speed and flexibility
among packages for zero-inflated Generalized Linear Mixed Modeling. R J. 9, 378.
https://doi.org/10.32614/RJ-2017-066

Cantwell, G., 1970. Standard methods for counting *Nosema* spores. Am. Bee J. 110, 222–223.

Chaimanee, V., Chantawannakul, P., Chen, Y., Evans, J.D., Pettis, J.S., 2012. Differential

expression of immune genes of adult honey bee (*Apis mellifera*) after inoculated by *Nosema*

ceranae. J. Insect Physiol. 58, 1090–1095. https://doi.org/10.1016/j.jinsphys.2012.04.016

Charbonneau, L.R., Hillier, N.K., Rogers, R.E.L., Williams, G.R., Shutler, D., 2016. Effects

of *Nosema apis*, *N. ceranae*, and coinfections on honey bee (*Apis mellifera*) learning and

memory. Sci. Rep. 6, 22626. https://doi.org/10.1038/srep22626

Collett, M., Chittka, L., Collett, T.S., 2013. Spatial memory in insect navigation. Curr. Biol.

23, R789–R800. https://doi.org/10.1016/j.cub.2013.07.020

Cox-Foster, D.L., Conlan, S., Holmes, E.C., Palacios, G., Evans, J.D., Moran, N.A., Quan, P.-

L., Briese, T., Hornig, M., Geiser, D.M., Martinson, V., vanEngelsdorp, D., Kalkstein, A.L.,

Drysdale, A., Hui, J., Zhai, J., Cui, L., Hutchison, S.K., Simons, J.F., Egholm, M., Pettis, J.S.,

Lipkin, W.I., 2007. A metagenomic survey of microbes in honey bee Colony Collapse

Disorder. Science 318, 283–287. https://doi.org/10.1126/science.1146498

Doublet, V., Paxton, R.J., McDonnell, C.M., Dubois, E., Nidelet, S., Moritz, R.F.A., Alaux,

C., Le Conte, Y., 2016. Brain transcriptomes of honey bees (*Apis mellifera*) experimentally

infected by two pathogens: Black queen cell virus and *Nosema ceranae*. Genomics Data 10,

79–82. https://doi.org/10.1016/j.gdata.2016.09.010

Dussaubat, C., Maisonnasse, A., Crauser, D., Beslay, D., Costagliola, G., Soubeyrand, S.,

Kretzchmar, A., Le Conte, Y., 2013. Flight behavior and pheromone changes associated to

Nosema ceranae infection of honey bee workers (*Apis mellifera*) in field conditions. J.

Invertebr. Pathol. 113, 42–51. https://doi.org/10.1016/j.jip.2013.01.002

Fries, I., Chauzat, M.-P., Chen, Y.-P., Doublet, V., Genersch, E., Gisder, S., Higes, M.,

McMahon, D.P., Martín-Hernández, R., Natsopoulou, M., Paxton, R.J., Tanner, G., Webster,

T.C., Williams, G.R., 2013. Standard methods for *Nosema* research. J. Apic. Res. 52, 1–28.

https://doi.org/10.3896/IBRA.1.52.1.14

Frost, P.C., Ebert, D., Smith, V.H., 2008. Bacterial infection changes the elemental

composition of *Daphnia magna*. J. Anim. Ecol. 77, 1265–1272.

https://doi.org/10.1111/j.1365-2656.2008.01438.x

Gage, S.L., Kramer, C., Calle, S., Carroll, M., Heien, M., DeGrandi-Hoffman, G., 2018.
Nosema ceranae parasitism impacts olfactory learning and memory and neurochemistry in
honey bees (*Apis mellifera*). J. Exp. Biol. 221, jeb161489. https://doi.org/10.1242/jeb.161489

Gisder, S., Horchler, L., Pieper, F., Schüler, V., Šima, P., Genersch, E., 2020. Rapid
gastrointestinal passage may protect *Bombus terrestris* from becoming a true host for *Nosema*
ceranae. Appl. Environ. Microbiol. 86. https://doi.org/10.1128/AEM.00629-20

Giurfa, M., 2015. Learning and cognition in insects: learning and insect cognition. Wiley
Interdiscip. Rev. Cogn. Sci. 6, 383–395. https://doi.org/10.1002/wcs.1348

Giurfa, M., Sandoz, J.-C., 2012. Invertebrate learning and memory: fifty years of olfactory
conditioning of the proboscis extension response in honeybees. Learn. Mem. 19, 54–66.
https://doi.org/10.1101/lm.024711.111

Gómez-Moracho, T., Durand, T., Pasquaretta, C., Heeb, P., Lihoreau, M., 2021. Artificial
diets modulate infection rates by *Nosema ceranae* in bumblebees. Microorganisms 9, 158.
https://doi.org/10.3390/microorganisms9010158

Gómez-Moracho, T., Heeb, P., Lihoreau, M., 2017. Effects of parasites and pathogens on bee
cognition: Bee parasites, pathogens and cognition. Ecol. Entomol. 42, 51–64.
https://doi.org/10.1111/een.12434

Graystock, P., Goulson, D., Hughes, W., 2015. Parasites in bloom: flowers aid dispersal and
transmission of pollinator parasites within and between bee species. Proc. R. Soc. B Biol. Sci.
282. https://doi.org/10.1098/rspb.2015.1371

Graystock, P., Yates, K., Darvill, B., Goulson, D., Hughes, W.O.H., 2013. Emerging dangers:
Deadly effects of an emergent parasite in a new pollinator host. J. Invertebr. Pathol. 114, 114–
119. https://doi.org/10.1016/j.jip.2013.06.005

Higes, M., García-Palencia, P., Martín-Hernández, R., Meana, A., 2007. Experimental
infection of *Apis mellifera* honeybees with *Nosema ceranae* (Microsporidia). J. Invertebr.
Pathol. 94, 211–217. https://doi.org/10.1016/j.jip.2006.11.001

Higes, M., Martín, R., Meana, A., 2006. *Nosema ceranae*, a new microsporidian parasite in
honeybees in Europe. J. Invertebr. Pathol. 92, 93–95. https://doi.org/10.1016/j.jip.2006.02.005

Higes, M., Martín-Hernández, R., Botías, C., Bailón, E.G., González-Porto, A.V., Barrios, L.,
del Nozal, M.J., Bernal, J.L., Jiménez, J.J., Palencia, P.G., Meana, A., 2008. How natural
infection by *Nosema ceranae* causes honeybee colony collapse. Environ. Microbiol. 10,
2659–2669. https://doi.org/10.1111/j.1462-2920.2008.01687.x

Holt, H.L., Aronstein, K.A., Grozinger, C.M., 2013. Chronic parasitization by *Nosema*
microsporidia causes global expression changes in core nutritional, metabolic and behavioral
pathways in honey bee workers (*Apis mellifera*). BMC Genomics 14, 799.
https://doi.org/10.1186/1471-2164-14-799

Hothorn, T., Bretz, F., Westfall, P., 2008. Simultaneous inference in general parametric
models. Biom. J. Biom. Z. 50, 346–363. https://doi.org/10.1002/bimj.200810425

Klee, J., Besana, A.M., Genersch, E., Gisder, S., Nanetti, A., Tam, D.Q., Chinh, T.X., Puerta,
F., Ruz, J.M., Kryger, P., Message, D., Hatjina, F., Korpela, S., Fries, I., Paxton, R.J., 2007.
Widespread dispersal of the microsporidian *Nosema ceranae*, an emergent pathogen of the
western honey bee, *Apis mellifera*. J. Invertebr. Pathol. 96, 1–10.
https://doi.org/10.1016/j.jip.2007.02.014

Klee, J., Tek Tay, W., Paxton, R.J., 2006. Specific and sensitive detection of *Nosema bombi*
(Microsporidia: Nosematidae) in bumble bees (*Bombus spp.*; Hymenoptera: Apidae) by PCR
of partial rRNA gene sequences. J. Invertebr. Pathol. 91, 98–104.
https://doi.org/10.1016/j.jip.2005.10.012

Klein, S., Cabirol, A., Devaud, J.-M., Barron, A.B., Lihoreau, M., 2017. Why bees are so
vulnerable to environmental stressors. Trends Ecol. Evol. 32, 268–278.
https://doi.org/10.1016/j.tree.2016.12.009

Koch, H., Brown, M.J., Stevenson, P.C., 2017. The role of disease in bee foraging ecology.
Curr. Opin. Insect Sci. 21, 60–67. https://doi.org/10.1016/j.cois.2017.05.008

Kraus, S., Gómez-Moracho, T., Pasquaretta, C., Latil, G., Dussutour, A., Lihoreau, M., 2019.

Bumblebees adjust protein and lipid collection rules to the presence of brood. Curr. Zool. 65,

437–446. https://doi.org/10.1093/cz/zoz026

Mallon, E.B., Brockmann, A., Schmid-Hempel, P., 2003. Immune response inhibits

associative learning in insects. Proc. R. Soc. Lond. B Biol. Sci. 270, 2471–2473.

https://doi.org/10.1098/rspb.2003.2456

Martín-Hernández, R., Meana, A., Prieto, L., Salvador, A.M., Garrido-Bailon, E., Higes, M.,

2007. Outcome of Colonization of *Apis mellifera* by *Nosema ceranae*. Appl. Environ.

Microbiol. 73, 6331–6338. https://doi.org/10.1128/AEM.00270-07

Mayack, C., Naug, D., 2009. Energetic stress in the honeybee *Apis mellifera* from *Nosema*

ceranae infection. J. Invertebr. Pathol. 100, 185–188.

https://doi.org/10.1016/j.jip.2008.12.001

Meana, A., Martín-Hernández, R., Higes, M., 2010. The reliability of spore counts to

diagnose *Nosema ceranae* infections in honey bees. J. Apic. Res. 49, 212–214.

https://doi.org/10.3896/IBRA.1.49.2.12

Menzel, R., 2012. The honeybee as a model for understanding the basis of cognition. Nat.

Rev. Neurosci. 13, 758–768. https://doi.org/10.1038/nrn3357

Menzel, R., 2001. Searching for the memory trace in a mini-brain, the honeybee. Learn. Mem.

8, 53–62. https://doi.org/10.1101/lm.38801

Menzel, R., Muller, U., 1996. Learning and memory in honeybees: from behavior to neural

substrates. Annu. Rev. Neurosci. 19, 379–404.

https://doi.org/10.1146/annurev.ne.19.030196.002115

Menzel, Randolf, Manz, G., Menzel, Rebecca, Greggers, U., 2001. Massed and spaced

learning in honeybees: the role of cs, us, the intertrial interval, and the test interval. Learn.

Mem. 8, 198–208. https://doi.org/10.1101/lm.40001

Mobley, M.W., Gegear, R.J., 2018. Immune-cognitive system connectivity reduces

bumblebee foraging success in complex multisensory floral environments. Sci. Rep. 8, 5953.

https://doi.org/10.1038/s41598-018-24372-5

Monchanin, C., Drujont, E., Devaud, J.-M., Lihoreau, M., Barron, A.B., 2020. Heavy metal
pollutants have additive negative effects on honey bee cognition. bioRxiv 2020.12.11.421305.
https://doi.org/10.1101/2020.12.11.421305

Palottini, F., Estravis Barcala, M.C., Farina, W.M., 2018. Odor learning and its experience-
dependent modulation in the South American native bumblebee *Bombus atratus*
(hymenoptera: apidae). Front. Psychol. 9, 603. https://doi.org/10.3389/fpsyg.2018.00603

Piiroinen, S., Botías, C., Nicholls, E., Goulson, D., 2016. No effect of low-level chronic
neonicotinoid exposure on bumblebee learning and fecundity. PeerJ 4, e1808.
https://doi.org/10.7717/peerj.1808

Piiroinen, S., Goulson, D., 2016. Chronic neonicotinoid pesticide exposure and parasite stress
differentially affects learning in honeybees and bumblebees. Proc. R. Soc. B Biol. Sci. 283,
20160246. https://doi.org/10.1098/rspb.2016.0246

Plischuk, S., Martín-Hernández, R., Prieto, L., Lucía, M., Botías, C., Meana, A.,
Abrahamovich, A.H., Lange, C., Higes, M., 2009. South American native bumblebees
(Hymenoptera: Apidae) infected by *Nosema ceranae* (Microsporidia), an emerging pathogen
of honeybees (*Apis mellifera*). Environ. Microbiol. Rep. 1, 131–135.
https://doi.org/10.1111/j.1758-2229.2009.00018.x

Porrini, M.P., Porrini, L.P., Garrido, P.M., de Melo e Silva Neto, C., Porrini, D.P., Muller, F.,
Nuñez, L.A., Alvarez, L., Iriarte, P.F., Eguaras, M.J., 2017. *Nosema ceranae* in South
American native stingless bees and social wasp. Microb. Ecol. 74, 761–764.
https://doi.org/10.1007/s00248-017-0975-1

Raiser, G., Galizia, C.G., Szyszka, P., 2017. A high-bandwidth dual-channel olfactory
stimulator for studying temporal sensitivity of olfactory processing. Chem. Senses 42, 141–
151. https://doi.org/10.1093/chemse/bjw114

Ravoet, J., De Smet, L., Meeus, I., Smagghe, G., Wenseleers, T., de Graaf, D.C., 2014.
Widespread occurrence of honey bee pathogens in solitary bees. J. Invertebr. Pathol. 122, 55–

58. https://doi.org/10.1016/j.jip.2014.08.007

Runckel, C., Flenniken, M.L., Engel, J.C., Ruby, J.G., Ganem, D., Andino, R., DeRisi, J.L.,
2011. Temporal analysis of the honey bee microbiome reveals four novel viruses and seasonal
prevalence of known viruses, *Nosema*, and *Crithidia*. PLoS ONE 6, e20656.
https://doi.org/10.1371/journal.pone.0020656

Scheiner, R., Abramson, C.I., Brodschneider, R., Crailsheim, K., Farina, W.M., Fuchs, S.,
Grünewald, B., Hahshold, S., Karrer, M., Koeniger, G., Koeniger, N., Menzel, R., Mujagic,
S., Radspieler, G., Schmickl, T., Schneider, C., Siegel, A.J., Szopek, M., Thenius, R., 2013.
Standard methods for behavioural studies of *Apis mellifera*. J. Apic. Res. 52, 1–58.
https://doi.org/10.3896/IBRA.1.52.4.04

Schmid-Hempel, P., 2013. Evolutionary parasitology: the integrated study of infections,
immunology, ecology, and genetics. Oxford University Press.

Schmid-Hempel, R., Tognazzo, M., 2010. Molecular divergence defines two distinct lineages
of *Crithidia bombi* (Trypanosomatidae), parasites of bumblebees. J. Eukaryot. Microbiol. 57,
337–345. https://doi.org/10.1111/j.1550-7408.2010.00480.x

Sinpoo, C., 2018. Impact of *Nosema ceranae* and *Nosema apis* on individual worker bees of
the two host species (*Apis cerana* and *Apis mellifera*) and regulation of host immune
response. J. Insect Physiol. 8.

Smith, M.L., 2012. The honey bee parasite *Nosema ceranae*: transmissible via food
exchange? PLOS ONE 7, e43319. https://doi.org/10.1371/journal.pone.0043319

Takeda, K., 1961. Classical conditioned response in the honey bee. J. Insect Physiol. 6, 168–
179. https://doi.org/10.1016/0022-1910(61)90060-9

Toda, N.R.T., Song, J., Nieh, J.C., 2009. Bumblebees exhibit the memory spacing effect.
Naturwissenschaften 96, 1185–1191. https://doi.org/10.1007/s00114-009-0582-1

Velthuis, H.H.W., Doorn, A. van, 2006. A century of advances in bumblebee domestication
and the economic and environmental aspects of its commercialization for pollination.

Apidologie 37, 421–451. https://doi.org/10.1051/apido:2006019

Villar, M.E., Marchal, P., Viola, H., Giurfa, M., 2020. Redefining single-trial memories in the

honeybee. Cell Rep. 30, 2603-2613.e3. https://doi.org/10.1016/j.celrep.2020.01.086

Wolf, S., McMahon, D.P., Lim, K.S., Pull, C.D., Clark, S.J., Paxton, R.J., Osborne, J.L.,

2014. So near and yet so far: harmonic radar reveals reduced homing ability of *Nosema*

infected honeybees. PLoS ONE 9, e103989. https://doi.org/10.1371/journal.pone.0103989

650