1 **Systematic temporal analysis of transcriptomes using *TrendCatcher* identifies**
2 **early and persistent neutrophil activation as an early hallmark of severe**
3 **COVID-19**
4

5 Xinge Wang[1,2,3], Mark Sanborn[1,2,3], Yang Dai[1] and Jalees Rehman[1,2,3*]
6
7 [1]Department of Biomedical Engineering, University of Illinois Colleges of Engineering and
8 Medicine, Chicago, IL, USA
9
10 [2]Department of Pharmacology and Regenerative Medicine, University of Illinois College of
11 Medicine, Chicago, IL, USA
12
13 [3]Division of Cardiology, Department of Medicine, University of Illinois College of Medicine,
14 Chicago, IL, USA
15
16 *Please address correspondence to Jalees Rehman (jalees@uic.edu)
17
The University of Illinois at Chicago, College of Medicine (M/C 868)
835 S. Wolcott Ave. Rm. E403, Chicago, IL 60612, USA
Office phone: +1 312-996-5552; Fax: +1 312-996-1225

Keywords:
RNA-sequencing
Single cell RNA-sequencing
Temporal analysis
Gene expression dynamics
Differential expression
Genomics
Transcriptomics
Pathway analysis
COVID-19
Immunity
Inflammation

Abstract

Studying the dynamic shifts in gene expression during disease progression may provide important insights into the biological mechanisms that distinguish adaptive and maladaptive responses. Existing computational tools for the analysis of transcriptomic data are not designed to optimally identify distinct temporal patterns when analyzing dynamic differentially expressed genes (DDEGs). Moreover, there is also a lack of methods to assess and visualize the temporal progression of biological pathways or gene sets mapped from time course transcriptomic datasets. In this study, we developed the open-source R package *TrendCatcher* (https://github.com/jaleesr/TrendCatcher), which applies the smoothing spline ANOVA model and break point searching strategy to estimate dynamic signals. *TrendCatcher* identifies and visualizes distinct dynamic transcriptional gene signatures and biological processes from sequential datasets. We used *TrendCatcher* to perform a systematic temporal analysis of COVID-19 peripheral blood transcriptomic datasets, including both whole blood bulk RNA-seq and PBMC scRNA-seq time course data. *TrendCatcher* uncovered the early and persistent activation of neutrophils as well as impaired type I interferon (IFN-I) signaling in circulating cells as early hallmark of patients who progressed to develop severe COVID-19, whereas no such patterns were identified in individuals receiving SARS-CoV-2 vaccinations or patients with mild COVID-19. These results underscore the importance of systematic temporal analysis of gene expression during disease progression to identify biomarkers and possible therapeutic targets.

Introduction

Time-course transcriptomic profiling has been widely used to study and model dynamic biological processes in cells (Bar-Joseph, Gitter, & Simon, 2012). By profiling mRNA levels during consecutive time points, researchers can infer dynamic responses to various external cues that cannot be observed by looking at only initial and terminal states. Recent improvements in high-throughput RNA sequencing (RNA-seq) technologies, including single-cell RNA-sequencing (scRNA-seq) provide viable approaches to study dynamic gene expression changes (Hwang, Lee, & Bang, 2018). Especially scRNA-seq allows for the in-depth analysis of temporal changes in distinct cell populations, thus providing insights into the heterogeneity and dynamics of responses to environmental cues or pathogenic stimuli. However, the analysis and visualization of longitudinal bulk RNA-seq data or scRNA-seq can be computationally challenging.

Currently, there are two predominant strategies for the analysis of sequential transcriptomic datasets. One strategy treats the sampling time points as categorical variables and is based on generalized linear models (GLMs). GLM-based packages include DESeq2 (Love, Huber, & Anders, 2014), edgeR (McCarthy, Chen, & Smyth, 2012) and limma (Ritchie et al., 2015). A complementary strategy is to treat time as a continuous variable and fit the time expression data into a spline-like model. These methods include DESeq2Spline (DESeq2 adopted with spline model for temporal RNA-seq datasets) fitting, ImpulseDE2 (Fischer, Theis, & Yosef, 2018) and Next maSigPro (Nueda, Tarazona, & Conesa, 2014). The former strategies focus on the magnitude of change instead of the time order of gene expression, and may also suffer from a relative loss of statistical testing power, especially if many time points are assessed (Fischer et al., 2018). The latter strategy increases the power of detecting dynamic genes, but is based on strong model assumptions which are not optimally suited for multiphasic responses, such as gene trajectory patterns that reflect initial acute stimuli followed by counter-regulatory compensatory responses. Furthermore, there is a lack of tools that leverage existing knowledge of functional pathway databases to infer and visualize pathway trajectories instead of individual gene trajectories only.

We found this is challenging in a complex disease, such as coronavirus disease 19 (COVID-19) caused by the SARS-CoV-2 virus infection (Zhou et al., 2020). COVID-19 is characterized by distinct disease progression patterns that suggest diverse host immune responses (Wu & McGoogan, 2020). Patients with severe disease exhibit profound inflammatory responses and immunopathology (Pedersen & Ho, 2020). COVID-19 immunophenotyping studies involve a

large number of time points from corresponding RNA-seq and scRNA seq datasets

(Bergamaschi et al., 2021; Bernardes et al., 2020). Existing approaches for temporal

transcriptomic analysis either do not take the temporal sequence into account when identifying

dynamic differentially expressed genes (DDEGs) using pairwise comparison between time

points, nor systematically analyze pathways that are dysregulated at defined time points.

In this study, we developed *TrendCatcher*, an open-source R package tailored for longitudinal

bulk RNA-seq and scRNA-seq analysis. *TrendCatcher* uses a framework that combines the

smoothing spline ANOVA model and break point searching strategy, which identifies inflection

points when gene expression trends reverse. We show that *TrendCatcher* outperformed

commonly used methods for longitudinal RNA-seq analysis when using simulated time course

data for benchmarking. We also analyzed bulk RNA-seq and scRNA-seq gene expression

profiles of peripheral blood cells in COVID-19 patients at various disease time points.

TrendCatcher allowed us to identify and visualize dynamic gene expression signature profiles

in peripheral blood that were associated with poor disease outcomes during the early phases

of disease, and could thus serve as novel mechanistic targets as well as early biomarkers for

patient prognostication.

Results

TrendCatcher accurately identifies DDEGs in simulated datasets

First, we tested the prediction performance of the *TrendCatcher* platform (**Figure 1A**) using a set of simulated time course RNA-seq datasets, because simulated data provides defined standards to assess the accuracy of novel analytical platforms. We considered a comprehensive collection of gene temporal trajectory patterns to simulate a set of realistic data with biological characteristics (Bar-Joseph et al., 2012). We embedded 10,000 simulated trajectories with varied temporal patterns, including 90% non-dynamic trajectories, 2.5% monotonous transition trajectories (continuously increasing or continuously decreasing gene expression levels throughout the time course), 2.5% impulse shaped single break point trajectories (only one temporal inflection point, i.e. up-peak-down or down-trough-up), 2.5% two break point trajectories and 2.5% three break point trajectories (multimodal dynamic response, e.g., a combination of 2 or more basic types of trajectories). Compared to DESeq2, DESeq2Spline and ImpulseDE2, *TrendCatcher* had the highest area under the ROC curve (AUC) in a mixed simulated dataset for time-course data with 7 time points (**Figure 1B**). We also tested each model's performance on a varying number of time points, including 3, 5, 7, 9 and 11 time points. As shown in **Figure 1C**, *TrendCatcher* had the highest prediction AUC across all time points, with the AUC values range from 0.88 to 0.90.

We next evaluated the prediction performance for each type of temporal trajectories. As shown in (**Figure 1 – Supplement 1**), although other three methods achieved slightly higher accuracy than *TrendCatcher* in monotonic trajectories, their AUCs dropped markedly when more complicated trajectories were embedded. DESeq2 only achieved AUC values of 0.49 to 0.62 for both bi-phasic trajectory and multimodal trajectory. The DESeq2Spline approach using a spline curve fitting model also dropped to an AUC of approximately 0.7 once multiphasic trajectories were introduced. These results suggests that existing approaches for longitudinal or time course analyses are well-suited for monotonic trajectories (continuously up or continuously down) but that *TrendCatcher* maybe more broadly applicable, because it identifies monotonic, biphasic (up-down, down-up) and multi-phasic shifts in gene expression, which are especially important in complex pathological setting when initial biological responses are followed by counter-regulatory adaptive or maladaptive responses.

TrendCatcher identifies rapid but transient upregulation of interferon signaling in peripheral blood following SARS-CoV-2 infection in a non-human primate model

To define the key dynamic gene signatures associated with SARS-CoV-2 infection in peripheral blood, we first analyzed the global transcriptomics profiles from a non-human primate dataset (Aid et al., 2020), in which samples were collected on Days 0 (uninfected controls), 1, 2, 4, 7, 10 and 14 following the live SARS-CoV-2 inoculation. This experiment in non-human primates had the advantage of clearly defining the timing of inoculation and following the time course of dynamic genes. *TrendCatcher* identified 962 DDEGs out of 12,754 total genes, accounting for 7.6% of total expression, suggesting that over 90% of the expressed genes in the peripheral blood remain close to the baseline expression levels even in the setting of the SARS-CoV-2 infection. We observed two major types of dynamic trajectories: (a) 167 genes followed a biphasic "0D-2D up, 2D-14D down" pattern, with their expression level peaking at Day 2 and gradually returning close to baseline levels at Day 14 (**Figure 2A**). These dynamic genes were primarily associated with host defense biological pathways, such as defense response to viruses, regulation of viral life cycle and type I interferon signaling pathways (**Figure 2B**). (b) 263 genes follow a "0D-14D down" pattern (**Figure 2A**). This set of genes followed a monotonous trajectory, with their expression gradually decreasing until Day 14. Interestingly, we found these genes were primarily associated with mitochondrial ATP synthesis and oxidative phosphorylation, suggesting that disruption of mitochondrial respiration and ATP generation may be an important feature of the circulating cells transcriptomic shift in a SARS-CoV-2 infection (**Figure 2C**). *TrendCatcher* assigned trajectory pattern types to all dynamic genes, and provided hierarchical pie charts to visualize the composition of trajectory patterns (**Figure 2-Supplement1 A**).

To systematically assess and visualize the dynamic programming of the top biological pathways associated with SARS-CoV-2 infection, *TrendCatcher* generated a *TimeHeatmap*. The *TimeHeatmap* function of *TrendCatcher* visualizes shifts in pathways by displaying the mean-fold change of individual DDEGs in a given pathway at defined time points while also depicting the number of dynamic genes and the percentage of dynamic genes within that pathway (**Figure 2D**). This quantifies the pathway level shifts over time, and allows for the magnitude of pathway change to be visualized. Together with the number and fraction of dynamic genes in a given pathway, to gauge the relative importance of the pathway during the time course.

During initial infection (Day 0 to Day2), pathways related to innate immune response and
interferon pathways were highly upregulated. Examples are upregulation of pathways such as
defense response to virus and regulation of innate immune response increased with an
average log2 fold-change of 2.57 and 1.76 within the first day. Mucosal immune response,
antimicrobial humoral response and killing of cells of other organisms were activated during
the later stage of infection (Day 4 to Day 7), with an average log2 fold-change around 2. On
the other hand, mitochondrial ATP synthesis coupled electron transport and protein targeting
to ER, on the other hand, were gradually down-regulated until Day 14. Dynamic gene
signatures from type I interferon signaling pathway and mitochondrial ATP synthesis coupled
electron transport were shown using traditional heatmaps in **Figure 2-Supplement 1 B,C**.
The temporal analysis of this non-human primate dataset highlights the rapidity of interferon
signaling activation btu also underscores that this initial burst of immune activation is transient
and followed by a gradual downregulation of the antiviral interferon responses during the first
week following infection.

Increased generation of immunoglobulin synthesis in plasma B cells as early as Day 1 of symptom onset in patients diagnosed with SARS-CoV-2 infection

Next, we analyzed the longitudinal gene expression profiles of peripheral blood mononuclear cells (PBMCs) obtained from patients diagnosed with a SARS-CoV-2 infection who were admitted to the hospital, but predominantly had uncomplicated disease progression with 4 out 5 patients showing only mild symptoms (Zhu et al., 2020) (**Figure 3A**). The study performed single cell RNA-seq analysis on PBMCs, thus allowing for a cell-type specific analysis of gene expression shifts in distinct B cell subtypes, T cell subtypes and NK cells. We adopted the cell-type labels from the original study and generated pseudo-bulk RNA-seq datasets for each cell type in order to quantify changes in DDEGs for a specific cell type. We also adopted the time annotation using stages, by binning the disease processes of COVID-19 patients from symptom onset to discharge into stages 0, 1, 2, 3 and 4. We defined Stage 0 as the time point of samples obtained from healthy controls and stages 1, 2, 3 and 4 represented Day 1 to Day 16 from symptom onset. We only applied *TrendCatcher* to cell types containing more than 1,000 cells at any given stage to ensure the robustness of the results.

TrendCatcher identified 400 DDEGs in memory B cells, 213 DDEGs in naive B cells, 1,413 DDEGs in plasma B cells, 398 DDEGs in CD4$^+$ T cells, 645 DDEGs in CD8$^+$ T cells, 423 DDEGs in MAIT, 1,161 DDEGs in naive T cells and 667 DDEGs in NK cells. The *TimeHeatmap* of plasma B cells visualized the most dynamic biological pathways. Importantly, this temporal analysis of single cell RNA-seq data showed how rapidly plasma B cells ramp up the upregulation Fc-gamma receptor signaling and immunoglobulin synthesis as early as stage 1 (which corresponds to Day 1 of symptom onset) (**Figure 3B**). However, not all immunoglobulin genes are upregulated during the same temporal phase. As shown in (**Figure 3-Supplement 1**), genes involved in immunoglobulin synthesis show distinct temporal patterns. Due to the comparatively lower number of plasma B cells, the prominence of such increased immunoglobulin changes may be diluted in peirpheral blood bulk RNA-seq analysis or PBMC bulk RNA-seq analysis. However, temporal analysis of all peripheral blood mononuclear cell types, across the whole time course demonstrated increases in type I interferon signaling and defense responses to viruses as the most prominent changes over time (**Figure 3C**), thus mirroring the responses to SARS-CoV-2 we observed in the peripheral blood of non-human primates (**Figure 2**). To define cell-type specific temporal dynamics, we generated *TimeHeatmaps* for individual PBMC cell types and found significant upregulation of Type I interferon signaling in T cells, NK cells and memory B cells during Stage 1 but

222   subsequent downregulation by Stage 2 of the disease in patients with mild symptoms (**Figure**

223   **3-Supplement 2A-C, 3A-D**).

TrendCatcher identifies early and persistent neutrophil activation as a hallmark of severe COVID-19

We then assessed whether a systematic temporal analysis of gene expression trends could be used to distinguish disease severity and prognosis of COVID-19 patients. We thus applied _TrendCatcher_ to a time course human whole blood bulk RNA-seq dataset (Bergamaschi et al., 2021), which contained longitudinal whole blood transcriptomes from COVID-19 patients with mild, moderate and severe clinical outcomes. _TrendCatcher_ identified 77 DDEGs from the mild group, 226 DDEGs from the moderate group, and 1,205 DDEGs from the severe group. Only 42 DDEGs were shared among these three groups (**Figure 4A**), and these were primarily associated with B cell mediated immunity, Fc-gamma receptor signaling pathways, humoral immune response and lymphocyte mediated immunity (**Figure 4B**). However, _TrendCatcher_ identified 978 DDEGs uniquely shown in the severe COVID-19 patient group. Importantly, these genes were strongly enriched for neutrophil-related biological pathways, including neutrophil activation and neutrophil mediated immunity. These severe-disease associated genes also included genes found in pathways such as myeloid cell differentiation, reactive oxygen species metabolic process, and positive regulation of cytokine production (**Figure 4B**).

To systematically characterize which biological processes in whole blood RNA-seq samples were most dynamic in severe COVID-19 patients and how they progress over time, we applied the _TrendCatcher_'s _TimeHeatmap_ function. As seen in the non-human primate PBMC and the human COVID-19 single cell RNA-seq datasets, we again observed the upregulation of genes involved in the response to virus, humoral immune response and type I interferon signaling pathway within Week 1 of disease onset (**Figure 4C**). Importantly, some dynamic biological responses were only enriched in the group of patients which subsequently progressed to severe COVID-19, including neutrophil activation (117 DDEGs, 23.4% of the corresponding GO term), blood coagulation (36 DDEGs, 10.5% of the corresponding GO term), and regulation of response to cytokine stimulus and respiratory burst (**Figure 4C, Figure 4-Supplement 1A-B**). For instance, neutrophil activation gene expression increased by a mean of 1.3 log2-units (approximately 2.5-fold increase in mean gene expression) within Week 1, increased continuously until Week 4, and only very gradually decreased in surviving patients by Week 7. However, the summation of the averaged log2 fold-change from the _TimeHeatmap_ was larger than zero, which indicates that the neutrophil activation may not have returned fully to baseline levels by 7 weeks. To confirm this, we applied LOESS smooth curve fitting to all neutrophil activation DDEGs identified from the three severity groups. LOESS fitting confirmed

that severe COVID-19 patients showed markedly higher neutrophil activation at the early

stage of infection (Week 1- Week 2), and also remained highly activated even after 7 weeks.

Such a persistent neutrophil activation was not observed in either mild or moderate COVID-

19 patient groups (**Figure 4D**).

We also observed severe COVID-19 patients showed evidence of greater humoral immune

response gene upregulation (**Figure 4E**), as well as markedly higher upregulation blood

coagulation genes (**Figure 4F**) and respiratory burst genes (**Figure 4G**), which remained

upregulated even after Week 6. These were not found to be upregulated in patients with mild

or moderate COVID-19 disease. These DDEGs and dynamic biological pathways may thus

be suited to serve as early biomarkers that distinguish severe COVID-19 from mild and

moderate COVID. All these dynamic gene signatures from neutrophil activation, blood

coagulation and respiratory burst pathways were listed using traditional heatmap (**Figure 4-**

Supplement 2A-C).

Early impaired IFN-I signaling in PBMC provides a hallmark of severe COVID-19

Next, to define the cell-type specific dynamic gene signatures and biological processes, we used *TrendCatcher* to analyze a human PBMC single cell RNA-seq dataset time course dataset in which patients were categorized as having either moderate or severe COVID-19 (Liu et al., 2021). Importantly, as this dataset only contained mRNA from peripheral blood mononuclear cells, it lacked mRNA from neutrophils. *TrendCatcher* generated "pseudo-bulk" mRNA profiles for each cell type in order to perform the analysis of gene expression dynamics in a cell-type specific manner. *TrendCatcher* identified more dynamic shifts in almost all cell types from severe COVID-19 patients than moderate groups (**Supplement Table 1, 2**). To identify dynamic responsive processes unique to the severe group, we compared GO enrichment for each cell type between severe and moderate. As shown in **Figure 5A**, all innate immune cells and some adaptive immune cells (including B cells, CD8$^+$ T cells) from moderate COVID-19 were highly enriched in type I interferon signaling, negative regulation of viral process and defense to virus, whereas this was not observed significantly enriched in severe COVID-19. Severe COVID-19 patients were highly dynamic in MAPK cascade, NF-κB signaling, T cell receptor signaling and positive regulation of cytokine production for both NK cells and CD4$^+$ T cells. For monocytes and DC (dendritic cells), no uniquely enriched dynamic biological processes were observed in severe COVID-19 patients versus moderate COVID-19 patients.

Furthermore, we also found the extent of the interferon signaling response to be the key distinguishing feature between moderate and severe COVID-19. To quantitatively compare the trajectory differentiation of the type I interferon signaling pathway, we performed the LOESS smooth fitting to the DDEGs identified from this pathway. As shown in **Figure 5B**, there is a profound separation between these two groups, with early strong activation of IFN-I signaling in NK cells, monocytes, B cells and CD8$^+$ T cells in moderate COVID-19 patients but not in severe COVID-19 patients. Patients with moderate COVID-19 showed activation of type I interferon signaling pathways whereas patients with severe COVID-19 had a blunted activation of type I interferon signaling. This is also shown in the cell-type specific *TimeHeatmap*. As shown in **Figure C,D**, although NK cells from both moderate and severe COVID-19 patients demonstrated activation of type I interferon response within the first week, moderate COVID-19 exhibited a stronger activation than severe group, with average 1.62 log2 fold-change compared to 0.87 log2 fold-change. In CD8$^+$ T cells, only moderate COVID-19 groups were observed to have a strong type I interferon response within the first week. On the other hand, CD8$^+$ T cells in patients who would go on to develop severe COVID-19, showed upregulation of cell proliferation and cell differentiation genes instead. Together, these data

311   suggest that a robust early Type I interferon response in PBMCs is associated with reduced

312   severity of COVID-19.

TrendCatcher identifies metabolic gene expression shifts in NK cells as a hallmark response to COVID-19 vaccination

We next applied _TrendCatcher_ to a longitudinal human PBMC single cell RNA-seq vaccination dataset (Arunachalam et al., 2021), to provide insights into how the immune system physiologically responds to mRNA vaccines over time. The vaccination study collected single-cell PBMCs from 56 healthy volunteers vaccinated with the Pfizer-BioNtech at Day 0, 1, 2, 7, 21, 22, 28, and 42 after the vaccination. We processed each patient's scRNA-seq individually. We clustered cells using the Seurat algorithm (Hao et al., 2021) and annotated the cell types using SingleR (Aran et al., 2019). Then we remove cell types containing less than 1,000 cells for each time point across all samples. As **Figure 6A** shown, it is the UMAP of one patient's PBMCs single cell data at Day 0.

TrendCatcher identified 650 DDEGs in NK cells, 450 in B cells, 23 in CD8$^+$ T cells, 62 in monocytes and only 6 in CD4$^+$ T cells. This indicates NK cells exhibit a strong dynamic gene expression shift after vaccination. After comparing the GO enrichment analysis across cell types, we found NK cells gene expression shifts were enriched in metabolic processes in response to the Pfizer-BioNtech SARS-CoV-2 mRNA vaccine, such as regulation of cellular amino acid metabolism and ATP metabolism (**Figure 6B**). As shown in **Figure 6C**, a _TimeHeatmap_ of NK cells shows 54 DDEGs, which account for nearly half of the ATP metabolic process pathway genes and exhibited a mean of 0.44 log2 fold upregulation after the first dose of vaccination. These findings indicate the reprogramming of metabolism in NK cells may be an indicator of an intact vaccine response. Another hallmark of the vaccine response was the activation of interferon pathways, including type I interferon signaling pathway and response to interferon-gamma (**Figure 6-Supplement 1A**). Importantly, the _TimeHeatmap_ demonstrates that these gene expression shifts were very transient, usually decreasing by Day 2 or Day 7, and were again upregulated when the subjects received the booster vaccine at Day 21. Interferon pathways showed an average increase of 0.71 and 1.42 log2 fold-change. In adaptive immune cells, we also observed strong type I interferon signaling pathways activated in both B cells and CD8$^+$ T cells (**Figure 6-Supplement 1B,C**). Additionally, B cells demonstrated upregulation of antigen processing and protein synthesis (**Figure 6-Supplement 1B**), which were upregulated only for the first day, and then began decreasing. It is noteworthy that early upregulation of interferon pathways in the PBMCs of vaccine recipients mirrors that seen in patients with moderate COVID-19 infection, consistent with the notion that early and transient interferon upregulation is a hallmark of a healthy immune response to the SARS-CoV-2 infection.

Discussion

Temporal analysis of gene expression is gaining importance in the analysis of complex dynamic processes such as disease progression. Besides gene dynamic pattern characterization, time-course gene expression data are also used to infer regulatory and signaling relationships among genes (Krieger, Lupo, Levy, & Barkai, 2020; Yuan & Bar-Joseph, 2019). Integrating with other different types of measurements, such as pathology and infection over time helps disentangle the complex dynamic processes and possible underlying mediators (Graw, 2021). Thus, accurate identification of dynamic differentially expressed genes (DDEGs) in a time course RNA-seq or single cell RNA-seq study can help identify time-dependent disease mechanisms, adaptive and maladaptive molecular signatures as well as potential biomarkers that may be associated with disease severity.

In recent years, tools have been implemented to characterize time course RNA-seq data; however, these tools were focused on either bulk RNA-seq datasets (Fischer et al., 2018; Love et al., 2014; Nueda et al., 2014) and few infer trajectories from scRNA-seq (Qiu et al., 2017; Van den Berge et al., 2020). Compared to methods that analyze time course data without considering the sequential nature of time points, modeling time as a continuous function avoids a relative loss of statistical testing power, especially when many multiple time points were studied (Fischer et al., 2018). There is also a need for methods which combine DDEGs identification with the visualization of dynamic pathway shifts.

In this study, we developed an open-source R-software package to perform temporal analysis in longitudinal bulk RNA-seq and scRNA-seq datasets. *TrendCatcher* can identify DDEGs and infer the trajectories for each DDEG. By combining a time-window sliding strategy on inferred gene trajectories and leveraging annotated biological pathway databases, *TrendCatcher* can infer and visualize the most dynamic biological pathways in response to the external stimuli. Furthermore, by utilizing the *TimeHeatmap* function, *TrendCatcher* can help researchers identify the magnitude and dynamic nature of pathways shifts. Using simulated datasets for benchmarking, *TrendCatcher* achieved higher accuracy (AUC = 0.9) compared to other commonly used methods for the analysis of temporal gene expression datasets, when analyzing three or more time points. The advantage of using *TrendCatcher* was especially apparent in the setting of biphasic or multiphasic temporal trajectories, in which gene expression levels can change their trend of upregulation or downregulation during the time course.

Human immune responses to SARS-CoV-2 infection are highly dynamic and time dependent
upregulation as well as downregulation of the immune responses are key to ensure optimal
host defense (Chen et al., 2020; Gong et al., 2020; Mehta et al., 2020; Qin et al., 2020). A
recent study of B cell dynamics highlighted how clonal lineages expand along infection
processes (Montague et al., 2021) and a parallel study of T cell memory clones identified the
dynamics of T-cell memory clones (Minervina et al., 2021). Understanding the dynamics of
the COVID-19 immune response could form the basis of developing therapies that are
appropriate for a given time window (Goyal, Cardozo-Ojeda, & Schiffer, 2020). Personalized
medicine or precision medicine are gaining traction by developing therapies tailored to patients
based on their genotype and phenotype, but personalization likely also requires tailoring
therapies based on the temporal phase of a disease. These studies highlight the need for
time-course or longitudinal analyses of the host responses to the SARS-CoV-2 infection.
397
In this study, we applied *TrendCatcher* to systematically analyze sequential blood samples
from either non-human primates infected with SARS-CoV-2, human patients with COVID-19
of varying severity or human subjects who received a SARS-CoV-2 mRNA vaccine.
TrendCatcher identified dynamic gene expression and biological pathway signatures for (a)
SARS-CoV-2 infection progression over time, (b) severe COVID-19 vs. moderate or mild
COVID-19 and (c) COVID-19 vaccine recipients versus control. *TrendCatcher* established
response to virus, humoral immune response and type I interferon signaling pathway
activation across peripheral blood cell types in mild, moderate and severe COVID-19.
However, we found temporal patterns of gene expression shifts that were unique in the severe
COVID-19 patients. Severe COVID-19 was associated with marked activation of neutrophils
as well as blunted type I interferon signaling as early as week 1. Importantly, severe COVID-
19 was associated with persistent activation of neutrophils and genes regulating coagulation
for as long as 6 weeks, underscoring that the temporal analysis by *TrendCatcher* identified
hallmarks of severe COVID-19 in peripheral blood samples.
412
Although roles of monocytes and macrophages in severe COVID-19 have been well
demonstrated (Merad & Martin, 2020), less is known about neutrophils, which are also an
important element of innate immunity, in COVID-19 (Laforge et al., 2020; Tomar, Anders,
Desai, & Mulay, 2020). As one of the first responders to microbial threats, neutrophils infiltrate
tissues and produce inflammatory molecules to facilitate pathogen clearance. Neutrophils are
particularly active in highly vascularized organs, such as lungs and kidneys, which are prime
targets in COVID-19 (Libby & Luscher, 2020). However, dysregulated neutrophil response,
such as prolonged activation may cause damage to vessels and underlying parenchymal
(Bardoel, Kenny, Sollberger, & Zychlinsky, 2014; Metzemaekers et al., 2021;

Papayannopoulos, 2018). Recently, some studies also focused on the kinetics of circulating

neutrophils during COVID-19 progression, and the key role neutrophils play in COVID-19

pathology (Kaiser et al., 2021; Metzemaekers et al., 2021; Radermecker et al., 2020). In our

study, we observed strongly neutrophil activation which was unique to the severe COVID-19

group. Importantly, we showed that neutrophils remained highly activated in surviving patients

with severe COVID-19 even after seven weeks. The increased activation of coagulation genes

in the whole blood mRNA studies in severe COVID-19 patients also points to another feature

that is associated with severe COVID-19, thrombotic and embolic complications such as

strokes (Ackermann et al., 2020; Aid et al., 2020; Rapkiewicz et al., 2020). We believe the

analysis of the RNA-seq and scRNA-seq data support neutrophil activation and upregulation

of coagulation genes as defining features of severe COVID-19, highlighting their role as early

biomarkers to provide prognostic information and thus optimally treat patients who have a

higher likelihood of progressing to severe disease. Importantly, our results also raise questions

about the role of neutrophils and of coagulation in what is referred to as "long COVID" or "post-

acute sequelae of COVID" (Carfi, Bernabei, Landi, & Gemelli Against, 2020; Huang et al.,

2021; Nalbandian et al., 2021), because we observed that the activation persists for as long

as seven weeks after the initial infection.

Our temporal analysis showed that severe COVID-19 single-cell PBMCs were characterized

by impaired type I interferon signaling at the onset of infection (Week 1), compared to the

moderate COVID-19 group. This impaired type I interferon signaling was identified in innate

cells, such as NK cells and monocytes, and adaptive cells, such as B cell and CD8[+] T cells.

Type I interferons, which are essential for antiviral immunity (McNab, Mayer-Barber, Sher,

Wack, & O'Garra, 2015; Muller et al., 1994), have been shown to be upregulated in COVID-

19 (Blanco-Melo et al., 2020; Hadjadj et al., 2020). Other studies also suggested that impaired

type I interferon signaling may promote severe COVID-19 and that interferon therapy could

be used as therapy in severe COVID-19 (Darazam et al., 2021). However, since these studies

were not designed for longitudinal analyses, the timing of when to intervene on IFN signaling

was not clear. Traditional analyses of gene expression data are not suited for the identification

of temporal windows and thus do not address whether impaired IFN production is present at

the onset of infection, whether it was delayed, or whether IFN production is exhausted after

an initial activation (Hadjadj et al., 2020). *TrendCatcher* addressed this question by showing

the dynamic timing of type I IFN from both the moderate group and severe group.

In our COVID-19 vaccination single cell PBMC temporal data analysis, we also identified

prominent metabolic shifts in NK cells. Cellular metabolism is recognized as an important

factor that can determine the fate and function of lymphocytes (Loftus & Finlay, 2016; O'Neill,

Kishton, & Rathmell, 2016). Certain metabolic pathways have been shown to have direct
immunoregulatory roles (Loftus & Finlay, 2016). Activated NK cells engage in a robust
metabolic response, which is required for normal effector functions (O'Brien & Finlay, 2019),
and our data suggests that assessing NK metabolic shifts may be an intriguing alternative to
assessing vaccine responsiveness, which does not rely on antibody levels which can be highly
variable in laboratory assays.

In conclusion, we have developed the novel *TrendCatcher* R package platform designed for
time course RNA-seq data analysis which identifies and visualizes distinct dynamic
transcriptional programs. When applied to real whole blood bulk RNA-seq time course
datasets from COVID-19 patients, we observed that patients with severe COVID-19 showed
gene expression profiles consistent with profound neutrophil activation early during the
progression of the disease (starting from the first week of symptom onset). This indicates that
neutrophil activation may be a key early biomarker and possible mediator of severe COVID-
19. When applied to scRNA-seq time course vaccination data of PBMCs (peripheral blood
mononuclear cells), *TrendCatcher* identified specific shifts in NK cell metabolic gene
expression programs. This raises important questions about the role of NK cells in vaccine
efficacy. The distinct kinetics of gene expression shifts identified by *TrendCatcher* in COVID-
19 patients may allow for the identification of therapeutic targets during distinct temporal
phases.

Acknowledgments
The studies were supported by NIH grants R01-HL149300 (to JR), P01-HL151327 (to JR),
and T32-HL139439 (to MS).

Competing interests
J.R. is a reviewing editor of eLife. There are no other competing interests to report.

Author contributions
X.W. and J.R. designed the study, X.W. performed the analyses and wrote the first draft of the
manuscript, J.R. co-wrote the initial draft of the manuscript. M.S. and Y.D. provided critical
revisions of the manuscript.

Materials and Methods

TrendCatcher framework

The main components of the *TrendCatcher* framework are shown in **Figure 1**. *TrendCatcher* requires two main inputs: the raw count table C of a temporal study with a dimension of $m \times n$, where m denotes the number of genes and n denotes the number of samples, and a user-defined baseline time variable T, such as "0 hour". Since samples may have different sequencing depths and batch effect, *TrendCatcher* integrates with limma (Ritchie et al., 2015) and provides preprocessing steps, such as batch correction and normalization. For single-cell RNA sequencing datasets, *TrendCatcher* extracts cells for each cell type annotated in the meta data slot of Seurat object and convert it into cell-type specific "pseudobulk" time course RNA library. Based on a user-specified threshold, relatively low abundant genes are removed from the count table, and reads are normalized and batch effects are removed. *TrendCatcher*'s core algorithm is composed of five main steps: (a) baseline fluctuation confidence interval estimation, (b) model dynamic longitudinal count, (c) time point dynamic p-value calculation, (d) gene-wise dynamic p-value calculation, and (e) break point screening and gene-wise dynamic pattern assignment. Mathematical details will be expanded in the following sections. For the output of *TrendCatcher*, there are mainly two components: a master table and a set of functions for versatile visualization purposes. The master table contains all the dynamic details of each single gene, including its dynamic p-value, its break point location time, and its dynamic trajectory pattern. In addition to the master table, *TrendCatcher* produces five main types of visualizations: (a) a figure showing the observed counts and fitted splines of each gene, (b) genes trajectories from each trajectory pattern group (c) a hierarchical pie chart that represents trajectory pattern composition, (d) a *TimeHeatmap* to infer trajectory dynamics of top dynamic biological pathways and (e) a two-sided bar plot to show the top most positively and negatively changed (averaged accumulative log2FC) biological pathways.

Baseline fluctuation confidence interval estimation

We assumed that the observed number of RNA-seq reads count from the baseline time (e.g., t=0 hour) $X_{i,t_{baseline}}$ was generated from a negative binomial distribution (Equation 1):

$$X_{i,t_{baseline}} \sim NB(\mu_{i,t_{baseline}}, \varphi_i) \tag{1}$$

Where gene $i = 1, \dots, n$ and $\mu_{i,t_{baseline}}$ is the mean count of gene i at baseline time, and φ_i is the dispersion factor. First, the dispersion factor φ_i was pre-estimated as a constant hyperparameter for each gene with DESeq2 (Love et al., 2014), as shown in Equation 2. Here, $\sigma_i(t)$ is the variance at time t.

$$\sigma_i(t)^2 = \mu_i(t) + \varphi_i * \mu_i(t)^2 \tag{2}$$

525   Then, $\mu_{i,t_{baseline}}$ was estimated using maximum likelihood from Equation 3.

$$\hat{\mu}_{i,t_{baseline}} = argmax\ L_{NB}(u,\varphi) = argmax\ \frac{\Gamma(\varphi+x)}{x!\,\Gamma(\varphi)} \left(\frac{\mu}{\mu+\varphi}\right)^x \left(\frac{\varphi}{\varphi+\mu}\right)^\varphi \tag{3}$$

526   Based on the negative binomial distribution and the estimated mean count for baseline time,

527   we constructed the 90% confidence interval $[X_{i,t_{baseline},0.05}, X_{i,t_{baseline},0.95}]$ as the baseline count

528   fluctuation interval (Equation 4(a) and Equation 4(b)).

$$P\left(x_{i,t_{baseline}} \le X_{i,t_{baseline},0.05}\ |\ \mu_{i,t_{baseline}}, \varphi_i\right) = 0.05 \tag{4a}$$

$$P\left(x_{i,t_{baseline}} \ge X_{i,t_{baseline},0.95}\ |\ \mu_{i,t_{baseline}}, \varphi_i\right) = 0.95 \tag{4b}$$

Model dynamic longitudinal count estimation

To model the time-dependent gene expression value, we applied a smoothing spline ANOVA

model (C, 2014; Metwally et al., 2018) with a negative binomial family constraint to fit the reads

from samples across non-baseline multiple time points. The random variable $X_{i,t(t \ne t_{baseline})}$ is

assumed to follow the NB distribution in Equation 5, with a positive integer α represents

number of failures before the αth success in a sequential of Bernoulli trials, and $p(t) \in (0,1)$

represents the success probability.

$$X_{i,t(t \ne t_{baseline})} \sim NB(\alpha, p(t)) \tag{5}$$

537   The log-likelihood given a time-course observed count $x = \{x_{i,t(t \ne t_{baseline})}\}_{i=1,\dots,n;\ t=1,\dots,T}$ is

538   calculated as Equation 6.

$$L = log\ L(\alpha, p(t)\ |\ X = x\ )$$

$$= \sum_{t=1}^{T}\ \sum_{i=1}^{n}\ [x_{i,t(t \ne t_{baseline})}\ log\ (1 - p(t) + \alpha log p(t) + log\Gamma\left(\alpha + x_{i,t(t \ne t_{baseline})}\right) - loglog\Gamma(\alpha)] + C \tag{6}$$

539   Taking the logit link and model time effect, we define the logit link η

$$\eta = log\ \frac{p(t)}{1 - p(t)} \tag{7}$$

540   To allow for flexibility in the estimation of η , and find the best trade-off between goodness of

541   fit and the smoothness of the spline curve, soft constraints of the form $J(\eta)$ is added to the

542   minus log-likelihood, with the smoothing parameter $\lambda > 0$.

$$-L + \lambda\ J(\eta) \tag{8}$$

543   The solution to the optimization of Equation 8 leads to a smoothing fitting to the reads from

544   samples across different non-baseline time points. The estimated mean of $\mu_{i,t(t \neq t_{baseline})}$ can

545   be estimated using Equation 9.

$$\mu_{i,t(t \neq t_{baseline})} = \hat{\alpha}e^{\hat{\eta}(t)} = \frac{\hat{\alpha}\hat{p}(t)}{1 - \hat{p}(t)} \tag{9}$$

Gene's dynamic p-value calculation

548   To calculate gene's non-baseline dynamic signal significance, each gene's non-baseline

549   estimated mean count $\mu_{i,t(t \neq t_{baseline})}$ was tested against the baseline fluctuation interval.

550   Based on Equation 10(a) and Equation 10(b), for each gene at each single non-baseline time

551   point, a dynamic time p-value was calculated.

$$If \; \mu_{i,t(t \neq t_{baseline})} \geq \mu_{i,t_{baseline}}, \; then \; p_{i,t} = \int P(x \geq \mu_{i,t(t \neq t_{baseline})} \,|\, \mu_{i,t_{baseline}}, \varphi_i) \tag{10a}$$

$$If \; \mu_{i,t(t \neq t_{baseline})} \leq \mu_{i,t_{baseline}}, \; then \; p_{i,t} = \int P(x \leq \mu_{i,t(t \neq t_{baseline})} \,|\, \mu_{i,t_{baseline}}, \varphi_i) \tag{10b}$$

552   Then, we applied Fisher's combined probability test method to calculate a gene-wise dynamic

553   p-value (Equation 11):

$$\chi^2_{i,2t} \sim -2 \sum_{t=0}^{T} ln\,(p_{i,t}) \tag{11}$$

Trajectory pattern assignment

555   First, we connect all the significant dynamic signal time points, with a p-value threshold less

556   than 0.05. Then we applied a break point searching strategy to capture the gene expression

557   change trend. The definition of break point is defined using Equation 12(a) and Equation 12(b).

$$If \; \mu_{i,t} > \mu_{i,t_{next}} \; AND \; \mu_{i,t} > \mu_{i,t_{previous}}, break \; point \; type \; I. \tag{12a}$$

$$If \; \mu_{i,t} < \mu_{i,t_{next}} \; AND \; \mu_{i,t} < \mu_{i,t_{previous}}, break \; point \; type \; II. \tag{12a}$$

558   There are two types of break points: type I means gene up-regulated followed by a down

559   regulation, and type II means gene expression level down regulated and then followed by an

560   upregulation. By screening along the break point, the master-pattern and sub-pattern were

561   assigned to each gene.

 TimeHeatmap **enrichment analysis and Two-sided bar plot**

To build the *TimeHeatmap* for visualizing the biological pathway enrichment change over time,

we designed a window-sliding strategy to capture all the up-regulated or down-regulated

genes within each time interval. If we denote time vector as t_j and $j \in 1, ..., T$, each time

interval is denoted as $[t_{j-1}, t_j]$. We found N_{up} up-regulated genes and N_{down} down-regulated

genes within the time window $[t_{j-1}, t_j]$, then Fisher's exact test was performed to obtain the

Gene Ontology (GO) term enrichment with the corresponding time interval for up-regulated

genes and down-regulated genes separately. Users can select the top most enriched

biological pathways (ranked by enrichment p-value) for each time interval (by default is 10).

Then for each selected GO term within the corresponding time window, we calculated the

averaged log 2 fold-change (Avg_log2FC) of all the DDEGs from this GO term. A series of

Avg_log2FC values over time characterize the trajectory dynamics of the corresponding

biological pathway, it is defined as biological pathway trajectory inference in this study. The

summation of the series of Avg_log2FC estimates the averaged accumulative log 2 fold-

change (GO_mean_logFC) for the corresponding GO term. *TrendCatcher* ranks biological

pathways based on their dynamic magnitude inferred from the GO_mean_logFC value. Users

are free to choose the top most positively and negatively changed (averaged accumulative

log2FC) biological pathways. Besides GO enrichment analysis (Yu, Wang, Han, & He, 2012),

TrendCatcher also packages Enrichr (Kuleshov et al., 2016) biological pathway databases.

Visualization is constructed using the ComplexHeatmap (Gu, Eils, & Schlesner, 2016)

package.

Simulated dataset

To mimic the real biological RNA-seq dataset, we only allowed ~10% of the genes to be

dynamic responsive genes. In this study, we embedded 5 different types of trajectories into

the temporal RNA-seq simulated datasets, including non-dynamic trajectory (~90%),

monotonic trajectory (~2.5%), bi-phase trajectory (~2.5%), and multimodal trajectory (~5%)

including 2 break-point and 3 break-point trajectory. Each type of trajectory was constructed

by adding negative binomial distribution noise to the embedded trajectory count. For example,

for monotonic trajectory, we defined the first and last time-points' RNA expression level

change to be 0.5-2 log2 fold-change, then we added negative binomial distribution noise to

the embedded monotonic trajectory. To check how the AUC changes as time-course study

extends longer, we sampled 3, 5, 7, 9 and 11 different number of time points with evenly time

intervals, and randomly sampled 3 replicates for each time point. To validate how the

prediction AUC varies across different types of trajectories, we embedded ~10% of the genes

to have non-dynamic trajectories, and ~90% of the genes have a specific dynamic trajectory
(monotonic, bi-phase or multimodal).

Pseudobulk RNA library construction
To construct "pseudo-bulk" RNA library from single cell RNA-seq datasets, all cells for each
cell type in a given sample were computationally "pooled" by summing all counts for a given
gene. Since pseudobulk libraries composed of few cells are not likely modeled properly, we
removed cell types containing less than 1k cells in this study. Lowly expressed genes were
removed for each cell type as well, using the filterByExpr function from *edgeR* R package
(McCarthy et al., 2012). Gene counts were transformed using the log of the counts per million
(cpm) and library size was normalized using calcNormFactors function with the method =
"RLE".

Gene set enrichment of DDEGs
Gene sets enrichment analysis in this study was performed using clusterProfiler (Yu et al.,
2012) R package, and enrichment comparison from multiple groups were visualized using the
compareCluster function.

Dataset	Organism	Number of time points	Study	Figure	Dataset Source	Reference
Bulk Peripheral Blood (Aid et al.)	Macaca mulatta	7	Covid vs. Normal	Figure 2	GSE156701	(Aid et al., 2020)
scRNA PBMC (Zhu et al.)	Homo Sapien	5	Covid vs. Normal	Figure 3	https://db.cngb.org/searc h/project/CNP0001102/	(Zhu et al., 2020)
Bulk Peripheral Blood (Bergamaschi et al.)	Homo Sapien	8	Severe vs. Moderate	Figure 4	https://www.covid19cellatl as.org/patient/citiid/	(Bergamaschi et al., 2021)
scRNA PBMC (Liu et al.)	Homo Sapien	4	Severe vs. Moderate	Figure 5	GSE161918	(Liu et al., 2021)
scRNA PBMC (Arunachalam et al.)	Homo Sapien	8	Vaccination	Figure 6	GSE171964 and GSE169159	(Arunachalam et al., 2021)

617   Table 1. Datasets presented in this study

Innate immune cells					
NK		Monocyte		DC	
Moderate	Severe	Moderate	Severe	Moderate	Severe
708	1,179	1,479	1,845	22	17

620 Supplement Table 1. Number of DDEGs for each innate immune cell type identified by

621 *TrendCatcher*. Using FDR less than 0.05.

Adaptive immune cells							
B		CD4[+] T		CD8[+] T		MAIT	
Moderate	Severe	Moderate	Severe	Moderate	Severe	Moderate	Severe
969	1,344	190	1,060	634	817	71	13

624 Supplement Table 2. Number of DDEGs for each adaptive immune cell type identified by

625 *TrendCatcher*. Using FDR less than 0.05.

References

Ackermann, M., Verleden, S. E., Kuehnel, M., Haverich, A., Welte, T., Laenger, F., . . . Jonigk, D. (2020). Pulmonary Vascular Endothelialitis, Thrombosis, and Angiogenesis in Covid-19. *New England Journal of Medicine, 383*(2), 120-128. doi:10.1056/NEJMoa2015432

Aid, M., Busman-Sahay, K., Vidal, S. J., Maliga, Z., Bondoc, S., Starke, C., . . . Barouch, D. H. (2020). Vascular Disease and Thrombosis in SARS-CoV-2-Infected Rhesus Macaques. *Cell, 183*(5), 1354-+. doi:10.1016/j.cell.2020.10.005

Aran, D., Looney, A. P., Liu, L. Q., Wu, E., Fong, V., Hsu, A., . . . Bhattacharya, M. (2019). Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic macrophage. *Nature Immunology, 20*(2), 163-+. doi:10.1038/s41590-018-0276-y

Arunachalam, P. S., Scott, M. K. D., Hagan, T., Li, C. F., Feng, Y. P., Wimmers, F., . . . Pulendran, B. (2021). Systems vaccinology of the BNT162b2 mRNA vaccine in humans. *Nature, 596*(7872), 410-+. doi:10.1038/s41586-021-03791-x

Bar-Joseph, Z., Gitter, A., & Simon, I. (2012). Studying and modelling dynamic biological processes using time-series gene expression data. *Nat Rev Genet, 13*(8), 552-564. doi:10.1038/nrg3244

Bardoel, B. W., Kenny, E. F., Sollberger, G., & Zychlinsky, A. (2014). The Balancing Act of Neutrophils. *Cell Host & Microbe, 15*(5), 526-536. doi:10.1016/j.chom.2014.04.011

Bergamaschi, L., Mescia, F., Turner, L., Hanson, A. L., Kotagiri, P., Dunmore, B. J., . . . Smith, K. G. C. (2021). Longitudinal analysis reveals that delayed bystander CD8(+) T cell activation and early immune pathology distinguish severe COVID-19 from mild disease. *Immunity, 54*(6), 1257-+. doi:10.1016/j.immuni.2021.05.010

Bernardes, J. P., Mishra, N., Tran, F., Bahmer, T., Best, L., Blase, J. I., . . . Deutsch, C.-O. I. (2020). Longitudinal Multi-omics Analyses Identify Responses of Megakaryocytes, Erythroid Cells, and Plasmablasts as Hallmarks of Severe COVID-19. *Immunity, 53*(6), 1296-+. doi:10.1016/j.immuni.2020.11.017

Blanco-Melo, D., Nilsson-Payant, B. E., Liu, W. C., Uhl, S., Hoagland, D., Moller, R., . . . tenOever, B. R. (2020). Imbalanced Host Response to SARS-CoV-2 Drives Development of COVID-19. *Cell, 181*(5), 1036-+. doi:10.1016/j.cell.2020.04.026

C, G. (2014). Smoothing spline ANOVA models: R package gss. *J Stat Softw., 58*(5), 25.

Carfi, A., Bernabei, R., Landi, F., & Gemelli Against, C.-P. A. (2020). Persistent Symptoms in Patients After Acute COVID-19. *Jama-Journal of the American Medical Association, 324*(6), 603-605. doi:10.1001/jama.2020.12603

Chen, G., Wu, D., Guo, W., Cao, Y., Huang, D., Wang, H. W., . . . Ning, Q. (2020). Clinical and immunological features of severe and moderate coronavirus disease 2019. *Journal of Clinical Investigation, 130*(5), 2620-2629. doi:10.1172/jci137244

Darazam, I. A., Shokouhi, S., Pourhoseingholi, M. A., Irvani, S. S. N., Mokhtari, M., Shabani, M., . . . Khoshkar, A. (2021). Role of interferon therapy in severe COVID-19: the COVIFERON randomized controlled trial. *Scientific Reports, 11*(1). doi:10.1038/s41598-021-86859-y

Fischer, D. S., Theis, F. J., & Yosef, N. (2018). Impulse model-based differential expression analysis of time course sequencing data. *Nucleic Acids Res, 46*(20), e119. doi:10.1093/nar/gky675

Gong, J., Dong, H., Xia, Q. S., Huang, Z. Y., Wang, D. K., Zhao, Y., . . . Lu, F. E. (2020). Correlation analysis between disease severity and inflammation-related parameters in patients with COVID-19: a retrospective study. *Bmc Infectious Diseases, 20*(1). doi:10.1186/s12879-020-05681-5

Goyal, A., Cardozo-Ojeda, E. F., & Schiffer, J. T. (2020). Potency and timing of antiviral therapy as determinants of duration of SARS-CoV-2 shedding and intensity of inflammatory response. *Science Advances, 6*(47). doi:10.1126/sciadv.abc7112

Graw, F. (2021). DISEASE Deciphering the triad of infection, immunity and pathology. *Elife, 10*. doi:10.7554/eLife.72379

Gu, Z. G., Eils, R., & Schlesner, M. (2016). Complex heatmaps reveal patterns and
correlations in multidimensional genomic data. *Bioinformatics, 32*(18), 2847-2849.
doi:10.1093/bioinformatics/btw313
Hadjadj, J., Yatim, N., Barnabei, L., Corneau, A., Boussier, J., Smith, N., . . . Terrier, B.
(2020). Impaired type I interferon activity and inflammatory responses in severe
COVID-19 patients. *Science, 369*(6504), 718-+. doi:10.1126/science.abc6027
Hao, Y. H., Hao, S., Andersen-Nissen, E., Mauck, W. M., Zheng, S. W., Butler, A., . . . Satija,
R. (2021). Integrated analysis of multimodal single-cell data. *Cell, 184*(13), 3573-+.
doi:10.1016/j.cell.2021.04.048
Huang, C. L., Huang, L. X., Wang, Y. M., Li, X., Ren, L. L., Gu, X. Y., . . . Cao, B. (2021). 6-
month consequences of COVID-19 in patients discharged from hospital: a cohort
study. *Lancet, 397*(10270), 220-232. doi:10.1016/s0140-6736(20)32656-8
Hwang, B., Lee, J. H., & Bang, D. (2018). Single-cell RNA sequencing technologies and
bioinformatics pipelines. *Experimental and Molecular Medicine, 50*.
doi:10.1038/s12276-018-0071-8
Kaiser, R., Leunig, A., Pekayvaz, K., Popp, O., Joppich, M., Polewka, V., . . . Stark, K.
(2021). Self-sustaining IL-8 loops drive a prothrombotic neutrophil phenotype in
severe COVID-19. *Jci Insight, 6*(18). doi:10.1172/jci.insight.150862
Krieger, G., Lupo, O., Levy, A. A., & Barkai, N. (2020). Independent evolution of transcript
abundance and gene regulatory dynamics. *Genome Research, 30*(7), 1000-1011.
doi:10.1101/gr.261537.120
Kuleshov, M. V., Jones, M. R., Rouillard, A. D., Fernandez, N. F., Duan, Q., Wang, Z., . . .
Ma'ayan, A. (2016). Enrichr: a comprehensive gene set enrichment analysis web
server 2016 update. *Nucleic Acids Res, 44*(W1), W90-97. doi:10.1093/nar/gkw377
Laforge, M., Elbim, C., Frere, C., Hemadi, M., Massaad, C., Nuss, P., . . . Becker, C. (2020).
Tissue damage from neutrophil-induced oxidative stress in COVID-19 (Jul,
10.1038/s41577-020-0407-1, 2020). *Nature Reviews Immunology, 20*(9), 579-579.
doi:10.1038/s41577-020-00425-7
Libby, P., & Luscher, T. (2020). COVID-19 is, in the end, an endothelial disease. *European
Heart Journal, 41*(32), 3038-3044. doi:10.1093/eurheartj/ehaa623
Liu, C., Martins, A. J., Lau, W. W., Rachmaninoff, N., Chen, J. G., Imberti, L., . . . Clinicians,
C. (2021). Time-resolved systems immunology reveals a late juncture linked to fatal
COVID-19. *Cell, 184*(7), 1836-+. doi:10.1016/j.cell.2021.02.018
Loftus, R. M., & Finlay, D. K. (2016). Immunometabolism: Cellular Metabolism Turns
Immune Regulator. *Journal of Biological Chemistry, 291*(1), 1-10.
doi:10.1074/jbc.R115.693903
Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol, 15*(12), 550.
doi:10.1186/s13059-014-0550-8
McCarthy, D. J., Chen, Y., & Smyth, G. K. (2012). Differential expression analysis of
multifactor RNA-Seq experiments with respect to biological variation. *Nucleic Acids
Res, 40*(10), 4288-4297. doi:10.1093/nar/gks042
McNab, F., Mayer-Barber, K., Sher, A., Wack, A., & O'Garra, A. (2015). Type I interferons in
infectious disease. *Nature Reviews Immunology, 15*(2), 87-103. doi:10.1038/nri3787
Mehta, P., McAuley, D. F., Brown, M., Sanchez, E., Tattersall, R. S., Manson, J. J., &
Speciality, H. L. H. A. (2020). COVID-19: consider cytokine storm syndromes and
immunosuppression. *Lancet, 395*(10229), 1033-1034. doi:10.1016/s0140-
6736(20)30628-0
Merad, M., & Martin, J. C. (2020). Pathological inflammation in patients with COVID-19: a
key role for monocytes and macrophages. *Nature Reviews Immunology, 20*(6), 355-
362. doi:10.1038/s41577-020-0331-4
Metwally, A. A., Yang, J., Ascoli, C., Dai, Y., Finn, P. W., & Perkins, D. L. (2018).
MetaLonDA: a flexible R package for identifying time intervals of differentially
abundant features in metagenomic longitudinal studies. *Microbiome, 6*(1), 32.
doi:10.1186/s40168-018-0402-y

Metzemaekers, M., Cambier, S., Blanter, M., Vandooren, J., de Carvalho, A. C., Malengier-
Devlies, B., . . . Proost, P. (2021). Kinetics of peripheral blood neutrophils in severe
coronavirus disease 2019. *Clinical & Translational Immunology, 10*(4).
doi:10.1002/cti2.1271
Minervina, A. A., Komech, E. A., Titov, A., Koraichi, M. B., Rosati, E., Mamedov, I. Z., . . .
Pogorelyy, M. V. (2021). Longitudinal high-throughput TCR repertoire profiling
reveals the dynamics of T-cell memory formation after mild COVID-19 infection. *Elife,*
10. doi:10.7554/eLife.63502
Montague, Z., Lv, H. B., Otwinowski, J., DeWitt, W. S., Isacchini, G., Yip, G. K., . . . Mok, C.
K. P. (2021). Dynamics of B cell repertoires and emergence of cross-reactive
responses in patients with different severities of COVID-19. *Cell Reports, 35*(8).
doi:10.1016/j.celrep.2021.109173
Muller, U., Steinhoff, U., Reis, L. F. L., Hemmi, S., Pavlovic, J., Zinkernagel, R. M., & Aguet,
M. (1994). FUNCTIONAL-ROLE OF TYPE-I AND TYPE-II INTERFERONS IN
ANTIVIRAL DEFENSE. *Science, 264*(5167), 1918-1921.
doi:10.1126/science.8009221
Nalbandian, A., Sehgal, K., Gupta, A., Madhavan, M. V., McGroder, C., Stevens, J. S., . . .
Wan, E. E. Y. (2021). Post-acute COVID-19 syndrome. *Nature Medicine, 27*(4), 601-
615. doi:10.1038/s41591-021-01283-z
Nueda, M. J., Tarazona, S., & Conesa, A. (2014). Next maSigPro: updating maSigPro
bioconductor package for RNA-seq time series. *Bioinformatics, 30*(18), 2598-2602.
doi:10.1093/bioinformatics/btu333
O'Brien, K. L., & Finlay, D. K. (2019). Immunometabolism and natural killer cell responses.
Nature Reviews Immunology, 19(5), 282-290. doi:10.1038/s41577-019-0139-2
O'Neill, L. A. J., Kishton, R. J., & Rathmell, J. (2016). A guide to immunometabolism for
immunologists. *Nature Reviews Immunology, 16*(9), 553-565.
doi:10.1038/nri.2016.70
Papayannopoulos, V. (2018). Neutrophil extracellular traps in immunity and disease. *Nature*
Reviews Immunology, 18(2), 134-147. doi:10.1038/nri.2017.105
Pedersen, S. F., & Ho, Y. C. (2020). SARS-CoV-2: a storm is raging. *Journal of Clinical*
Investigation, 130(5), 2202-2205. doi:10.1172/jci137647
Qin, C., Zhou, L. Q., Hu, Z. W., Zhang, S. Q., Yang, S., Tao, Y., . . . Tian, D. S. (2020).
Dysregulation of Immune Response in Patients With Coronavirus 2019 (COVID-19)
in Wuhan, China. *Clinical Infectious Diseases, 71*(15), 762-768.
doi:10.1093/cid/ciaa248
Qiu, X. J., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H. A., & Trapnell, C. (2017).
Reversed graph embedding resolves complex single-cell trajectories. *Nature*
Methods, 14(10), 979-+. doi:10.1038/nmeth.4402
Radermecker, C., Detrembleur, N., Guiot, J., Cavalier, E., Henket, M., d'Emal, C., . . .
Marichal, T. (2020). Neutrophil extracellular traps infiltrate the lung airway, interstitial,
and vascular compartments in severe COVID-19. *Journal of Experimental Medicine,*
217(12). doi:10.1084/jem.20201012
Rapkiewicz, A. V., Mai, X. C., Carsons, S. E., Pittaluga, S., Kleiner, D. E., Berger, J. S., . . .
Reynolds, H. R. (2020). Megakaryocytes and platelet-fibrin thrombi characterize
multi-organ thrombosis at autopsy in COVID-19: A case series. *Eclinicalmedicine,*
24. doi:10.1016/j.eclinm.2020.100434
Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., & Smyth, G. K. (2015).
limma powers differential expression analyses for RNA-sequencing and microarray
studies. *Nucleic Acids Res, 43*(7), e47. doi:10.1093/nar/gkv007
Tomar, B., Anders, H. J., Desai, J., & Mulay, S. R. (2020). Neutrophils and Neutrophil
Extracellular Traps Drive Necroinflammation in COVID-19. *Cells, 9*(6).
doi:10.3390/cells9061383
Van den Berge, K., de Bezieux, H. R., Street, K., Saelens, W., Cannoodt, R., Saeys, Y., . . .
Clement, L. (2020). Trajectory-based differential expression analysis for single-cell
sequencing data. *Nature Communications, 11*(1). doi:10.1038/s41467-020-14766-3

Wu, Z. Y., & McGoogan, J. M. (2020). Characteristics of and Important Lessons From the
Coronavirus Disease 2019 (COVID-19) Outbreak in China Summary of a Report of
72 314 Cases From the Chinese Center for Disease Control and Prevention. *Jama-*
Journal of the American Medical Association, 323(13), 1239-1242.
doi:10.1001/jama.2020.2648
Yu, G., Wang, L. G., Han, Y., & He, Q. Y. (2012). clusterProfiler: an R package for
comparing biological themes among gene clusters. *OMICS, 16*(5), 284-287.
doi:10.1089/omi.2011.0118
Yuan, Y., & Bar-Joseph, Z. (2019). Deep learning for inferring gene relationships from
single-cell expression data. *Proceedings of the National Academy of Sciences of the*
United States of America, 116(52), 27151-27158. doi:10.1073/pnas.1911536116
Zhou, F., Yu, T., Du, R. H., Fan, G. H., Liu, Y., Liu, Z. B., . . . Cao, B. (2020). Clinical course
and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a
retrospective cohort study. *Lancet, 395*(10229), 1054-1062. doi:10.1016/s0140-
6736(20)30566-3
Zhu, L. N., Yang, P. H., Zhao, Y. Z., Zhuang, Z. K., Wang, Z. F., Son, R., . . . Liu, W. J.
(2020). Single-Cell Sequencing of Peripheral Mononuclear Cells Reveals Distinct
Immune Response Landscapes of COVID-19 and Influenza Patients. *Immunity,*
53(3), 685-+. doi:10.1016/j.immuni.2020.07.009

Figure 1. **Overview and benchmarking of of *TrendCatcher*.** (A) *TrendCatcher* framework. *TrendCatcher* preprocess input data, which includes creating cell-type specific "pseudobulk" datasets for temporal analysis when single-cell data is used. *TrendCatcher*'s core algorithm is composed of five main steps. *TrendCatcher*'s output includes four main types of visualizations and DDEG identification (numbered 1-5). (B) *TrendCatcher* prediction ROC for a 7 time point simulated dataset compared to DESeq2, DESeq2Spline and ImpulseDE2, with mixed trajectories. (C) *TrendCatcher* prediction performance (AUC) across different number of time points, from 3 to 11 time points. *TrendCatcher*'s AUC values across time points from 3 to 11 are, 0.90, 0.92, 0.90, 0.89 and 0.88.

Figure 1 - Supplement 1. (A) Prediction performance of *TrendCatcher* across varying numbers of time points for monotonic trajectories. (B) Prediction performance of *TrendCatcher* across varying numbers of time points for biphasic trajectories. (C) Prediction performance of *TrendCatcher* across varying numbers of time points for multimodal trajectories. DESeq2 is shown in green, DESeq2Spline is shown in yellow, ImpulseDE2 is shown in purple, *TrendCatcher i*s shown in red.

Figure 2. Dynamic gene expression in peripheral blood following SARS-CoV-2 inoculation in a non-human primate model. (A) Analysis of the two predominant trajectory patterns in the non-human primate peripheral blood RNA-seq data from Day 0 to Day 14. The top-left figure represents 167 DDEGs following an up-down expression pattern, which peaked at Day 2 then slowly decreased until Day 14. The top-right figure represents their expression using a traditional z-score normalized heatmap. The bottom-left figure represents 263 DDEGs following a monotonic down-regulated trajectory pattern and their gene expression values were represented in the corresponding heatmap on the right. (B) Top 3 Gene ontology (GO) enrichment analysis pathways using 167 DDEGs from trajectory pattern "0D-2D Up, 2D-14D Down". (C) Top 3 Gene ontology (GO) enrichment analysis pathways using 263 DDEGs from trajectory pattern "0D-14D Down". (D) *TimeHeatmap* of the top 15 dynamic pathways and their dynamic time windows visualizes the temporal patterns. Each column represents a time window. The "%GO" column represents the percentage of DDEGs found in the corresponding pathway. The "nDDEG" column represents number of DDEGs found in the corresponding pathway. Color represents the averaged log fold-change of the DDEGs within each time window for the corresponding pathway.

Figure 2-Supplement 1. (A) Hierarchical pie chart shows the composition of trajectory patterns of DDEGs identified in a non-human primate bulk peripheral blood mRNA (Aid et al.). (B) Dynamic gene signatures for SARS-CoV-2 infection from type I interferon signaling pathway. (C) Dynamic gene signatures response for SARS-CoV-2 infection from mitochondrial ATP synthesis coupled electron transport pathway. Color represents normalized z-score for gene expression.

Figure 3. Cell-type specific dynamic gene expression in peripheral blood mononuclear cells following SARS-CoV-2 infection in patients. (A) UMAP visualization of single cell RNA-seq PBMC dataset (Zhu et al.) with annotated cell types from the original study. (B) *TimeHeatmap* of top dynamic biological pathway from plasma B cells. Each column represents a time window (stage). Stage 0 represents uninfected baseline. Each row represents the most dynamic biological pathways. Color represents the averaged log fold-change of the DDEGs within each time window from the corresponding pathway. (C) Top GO enrichment comparison analysis using DDEGs from each cell type. Dot size represents gene ratio. Dot color represents GO enrichment p-value. The number in the x-axis labels represent the number of DDEGs for each cell type.

Figure 3-Supplement 1. Heatmap showing B cell's 38 DDEGs identified from immunoglobulin production process. Color represents normalized z-score of gene expression.

Figure 3-Supplement 2. (A) *TimeHeatmap* of Memory B cells. (B) *TimeHeatmap* of Naive B cells. (C) *TimeHeatmap* of NK cells.

Figure 3-Supplement 3. (A) *TimeHeatmap* of Naive T cells. (B) *TimeHeatmap* of CD4⁺ T cells. (C) *TimeHeatmap* of CD8⁺ T cells. (D) *TimeHeatmap* of MAIT cells.

Figure 4. Temporal analysis of whole blood RNA-seq data in patients grouped according to disease severity. (A) Venn diagram of DDEGs identified from three COVID-19 severity groups. (B) Top GO enrichment from shared DDEGs across three groups compared with top GO enrichment from severe group. Dot size represents number of DDEGs enriched from the corresponding biological pathway. (C) *TimeHeatmap* of the top dynamic pathways from the severe group. Each column represents a time window, with W depicting week. (D-E) LOESS curve fitting of DDEGs identified in the severe COVID-19 group of the (D) neutrophil activation pathway, (E) humoral immune response pathway, (F) blood coagulation pathway and (G) respiratory burst pathway. Red curves represent the severe group, blue curves represent the moderate group, and green curves represent the mild group.

A

Mild COVID-19

	0W-1W	1W-2W	2W-3W	3W-4W	4W-5W	5W-6W	6W-7W	%GO	nDDEG
Fc-gamma receptor signaling pathway	1.71	0.49	-0.03	-0.74	-0.6	-0.31	-0.25	18.6	26
B cell mediated immunity	1.68	0.6	-0.06	-0.73	-0.59	-0.3	-0.25	13.3	30
lymphocyte mediated immunity	1.68	0.6	-0.06	-0.73	-0.59	-0.3	-0.25	8.3	30
phagocytosis	1.68	0.6	-0.21	-0.65	-0.59	-0.3	-0.25	7.9	30
humoral immune response	1.63	0.56	0.25	-0.85	-0.58	-0.29	-0.24	8.4	32
protein targeting to ER			0.73	-1.82	1		0.53	4.2	5

Avg_log2FC
2
1
0
-1
-2

B

Moderate COVID-19

	0W-1W	1W-2W	2W-3W	3W-4W	4W-5W	5W-6W	6W-7W	%GO	nDDEG
type I interferon signaling pathway	2.34	-0.63	-0.57	-0.52	-0.47	-0.42	-0.4	26.3	25
humoral immune response	1.53	1.14	-0.87	-0.99	-0.51	-0.27	-0.28	15.5	59
lymphocyte mediated immunity	1.48	1.29	-0.86	-1.06	-0.54	-0.24	-0.22	13.9	50
B cell mediated immunity	1.47	1.32	-0.87	-1.09	-0.54	-0.23	-0.22	20.8	47
Fc-gamma receptor signaling pathway	1.43	1.42	-0.89	-1.12	-0.54	-0.21	-0.2	28.6	40
phagocytosis	1.39	1.23	-0.87	-1.04	-0.52	-0.23	-0.2	13.4	51

Avg_log2FC
4
2
0
-2
-4

Figure 4-Supplement 1. (A) *TimeHeatmap* of top dynamic GO terms found in severe COVID-19 shown in mild group. (B) *TimeHeatmap* of top dynamic GO terms found in severe COVID-19 shown in moderate group.

Figure 4-Supplement 2. (A) Heatmap of severe COVID-19 DDEGs identified from the neutrophil activation pathway. The heatmap is ordered by week following infection with healthy controls (HC) on the left. Color represents Z-score normalized expression values. Each column represents one sample.

Figure 4-Supplement 2. (B) Severe COVID-19 DDEGs identified from the blood coagulation pathway. (C) Severe COVID-19 DDEGs identified from respiratory burst pathway. The heatmap is ordered by week following infection with healthy controls (HC) on the left. Color represents Z-score normalized expression values.

Figure 5. Temporal analysis of single cell RNA-seq data of PBMCs from patients with either moderate or severe COVID-19. (A) Dot plot showing GO enrichment comparison between severe COVID-19 and moderate COVID-19 for each cell type. The size of the dot represents gene count. (B) LOESS curve fitting on DDEGs identified from the type I interferon pathway using TrendCatcher from moderate COVID-19 and severe COVID-19. Blue color indicates moderate group and red color indicates severe group. (C) *TimeHeatmap* of NK cells from moderate COVID-19. (D) *TimeHeatmap* of NK cells from severe COVID-19. (E) *TimeHeatmap* of CD8+T cells from moderate COVID-19. (F) *TimeHeatmap* of CD8+T cells from severe COVID-19. Each column represents a time window with W depicting week.

C

	0D-1D	1D-2D	2D-7D	7D-21D	21D-22D	22D-28D	28D-42D	%GO	nDDEG
type I interferon signaling pathway	0.6	0.2	0.05	0.27	0.11	-0.06	-0.54	16.8	16
regulation of cell cycle phase transition	0.59	0	0.02	-0.16	0	-0.03	-0.19	10.5	50
response to interferon-gamma	0.58	0.02	0.08	0.25	0.22	0.14	-0.47	10.2	20
interferon-gamma-mediated signaling pathway	0.55	0.03	0.1	0.31	0.26	0.13	-0.63	13.2	12
Fc receptor signaling pathway	0.49	0	0.02	-0.01	0.01	0.02	-0.38	18	43
regulation of innate immune response	0.47	-0.01	-0.03	0.02	0.05	-0.01	-0.34	14	44
antigen processing and presentation of peptide antigen via MHC class I	0.46	-0.01	-0.02	-0.04	0.01	-0.03	-0.26	34.3	34
regulation of cellular amino acid metabolic process	0.46	-0.01	-0.02	-0.05	0	-0.03	-0.25	43.5	27
ATP metabolic process	0.44	-0.01	0.02	-0.08	-0.01	-0.04	-0.18	17.3	54
oxidative phosphorylation	0.42	0	0.01	-0.09	-0.02	-0.04	-0.17	27.7	41

Figure 6. Temporal analysis of PBMC single cell RNA-seq data from human subjects receiving the SARS-CoV-2 mRNA vaccine. (A) UMAP of the single-cell transcriptional profile of patient 2047 at Day 0. Cell types were auto-annotated by SingleR. (B) Dot plot of comparison of the top GO terms enriched from cell-type specific DDEGs. *TrendCatcher* identified 650, 62, 450, 6 and 23 DDEGs from NK cells, monocytes, B cells, CD4+ T cells and CD8+ T cells. Size of the dot represents gene ratio from the enrichment analysis corresponding to each cell type. (C) T*imeHeatmap* of NK cells. Each column represents a time window measured by days. The second dose was administered at Day 21.

Figure 6-Supplement 1. (A) *TimeHeatmap* of monocytes. (B) *TimeHeatmap* of B cells. (C) *TimeHeatmap* of CD8+T cells. Day 1 is the first dose of vaccination, Day 21 is the second dose of vaccination.