<u>Manuscript type</u>: Research Advances
<u>Title:</u>
Identification of organ-specific transcriptomic shifts in the vasculature during
systemic inflammation using TrendCatcher
<u>Authors:</u>

Xinge Wang[1], Mark Sanborn[1, 2, 3], Yang Dai[1] and Jalees Rehman[1, 2, 3*]
Xinge Wang: xwang234@uic.edu, Mark Sanborn (msanbo2@uic.edu), Yang Dai:
(yangdai@uic.edu), Jalees Rehman: jalees@uic.edu
[1] Department of Bioengineering, The University of Illinois College of Engineering and
College of Medicine, Chicago, IL, USA
11
12 [2] Department of Pharmacology and Regenerative Medicine, University of Illinois College
13 of Medicine, Chicago, IL, USA
14
15 [3] Division of Cardiology, Department of Medicine, University of Illinois College of
16 Medicine, Chicago, IL, USA
17
18 *Please address correspondence to: Jalees Rehman (jalees@uic.edu)
19
20 The University of Illinois at Chicago, College of Medicine (M/C 868)
21 835 S. Wolcott Ave. Rm. E403, Chicago, IL 60612, USA
22 Office phone: +1 312-996-5552; Fax: +1 312-996-1225

23
<u>Keywords:</u>
Endothelial cells
Endothelial heterogeneity
RNA-seq
Differential expression
Dynamic programming
Vascular biology
Inflammation
Transcription
Translation
Systems biology
Pathway analysis
Brain endothelium
Heart endothelium
Lung endothelium

Abstract

A recent analysis of endothelial cell (EC) gene expression suggests that ECs of distinct organs vary in their gene expression profiles and respond distinctly to the systemic inflammatory stimulus of endotoxemia (Jambusaria et al., 2020). There is a need to systematically define tissue-specific gene expression dynamics in response to inflammation but such an analysis is in part limited by the availability of appropriate algorithms to analyze differential expression across a time course. Here, we present *TrendCatcher* (https://jaleesr.github.io/TrendCatcher_1.0.0/), an R package designed for time course RNA-seq data analysis which identifies distinct dynamic transcriptional programs. When applied to ECs, we observed that approximately 85-95% of EC genes in all three organs followed a biphasic response following endotoxemia. The rapid upregulation of innate immune response and bacterial response genes occurred within 6 hours in all three vascular beds but the subsequent upregulation of reparative EC mitosis and cell cycle genes occurred most rapidly in lung ECs (24-48h) and was most delayed in cardiac ECs (72h-168h). The distinct kinetics of EC inflammatory injury and regeneration identify vascular-bed specific temporal windows for targeted therapeutics.

<u>Introduction</u>

Endothelial cells (ECs) form the inner cellular lining of all blood vessels and establish a barrier

between the circulating blood and the underlying parenchyma (Aird, 2012; Bautch, 2011). ECs

perform a wide range of critical functions in many physiological processes, such as maintaining

intravascular homeostasis, trafficking nutrients and waste between blood and the parenchymal

tissues, as well as initiating and regulating immune responses (Liao, 2013). Despite sharing

numerous essential roles in physiology, ECs exhibit distinct molecular and functional properties

across different tissue types and vascular locations (Nolan et al., 2013; Potente & Mäkinen, 2017).

EC heterogeneity has been validated and characterized in both within-organ and organ-specific

scenarios. For example, studies focused on characterizing EC heterogeneity within single organs

(Sabbagh et al., 2018; Su et al., 2018; Vanlandewijck et al., 2018) have revealed that local cues

from the tissue microenvironment, such as metabolites, cell-cell interactions, and cell-matrix

interactions, are critical for EC heterogeneity (Lukowski et al., 2019; Potente & Mäkinen, 2017; N.

Yucel et al., 2020). The heterogeneity of the microvascular endothelium across organs is apparent

in the distinct gene expression profiles of ECs from each major organ (Jambusaria et al., 2020;

Kalucka et al., 2020; Nolan et al., 2013; Paik et al., 2020).

To investigate the transcriptomic basis of EC heterogeneity, unbiased transcriptomic

profiling is essential for providing precise insights into the underpinning molecular mechanisms.

Bulk RNA sequencing (RNA-seq) and single-cell RNA sequencing (scRNA-seq) provide viable

approaches to study global gene expression. Studies using scRNA-seq have characterized

transcriptomic signatures of distinct EC subpopulations and the imprinted transcriptome (Kalucka

et al., 2020; Paik et al., 2020). However, the isolation of defined cell populations from their

physiological tissue context is required to prepare single-cell suspensions (Eberwine, Sul, Bartfai,

& Kim, 2014). Disaggregation of ECs for single-cell analysis or for flow cytometry cell sorting

results in the loss of homeostatic outside-in signaling cues from neighboring cells and the

extracellular matrix, and could thus alter gene expression profiles (Haimon et al., 2018; van den

Brink et al., 2017). The complementary approach of RiboTag-based translatome profiling avoids

cell removal from their native niche and enables the direct isolation of tissue-specific ribosome-

attached mRNAs without cell dissociation (Cleuren et al., 2019; Sanz et al., 2009). More

importantly, this genetic approach allows the direct probing of EC heterogeneity *in situ*, so it offers

the recovery and sequence analysis to evaluate the program directly from intact organs (Cleuren

et al., 2019). A recent study of the translatome in the cardiac endothelium demonstrated that the

tissue-specific endothelial gene expression profile is a reflection of signature open chromatin

regions within the cardiac endothelium, which are lost once the cardiac endothelium is removed

from its native environment and cultured (N. Yucel et al., 2020). Furthermore, RiboTag

complements conventional global bulk RNA-seq and scRNA-seq EC transcriptomic analysis by

focusing on mRNAs undergoing translation whereas global RNA-Seq analysis and single-cell

RNA-Seq analysis provide profiles reflecting total mRNA in a cell (Jambusaria et al., 2020).

Recent studies on organ-specific EC transcriptomic profiles have focused on differences

between ECs at baseline but there is a growing interest in how organ-specific EC heterogeneity

changes in response to a stressor such as inflammation. Which biological responses show similar

changes in magnitude and kinetics across endothelial cells from distinct organs and which stress

responses demonstrate organ-specific patterns? This has been addressed in part in recent

studies (Ajami et al., 2017; Chong, Yu, Brighton, Bear, & Bautch, 2017; Gunawardana et al., 2021;

Tombor et al., 2021), but there is a need for a systematic and unbiased analysis of the temporal

kinetics of endothelial gene expression across multiple organs. The endothelial RiboTag

transcriptome data collected at 6 different time points following induction of endotoxemia from the

heart, brain and lung (Jambusaria et al., 2020) is ideally suited to perform a systematic temporal

analysis. Such temporal dynamic characterization would capture the timing of cellular stress

response programs and provide insights into the heterogeneous organ-specific kinetics of

endothelial injury and repair.

Analysis of time-course RNA-seq data represents a computational challenge. Currently,

there are mainly two strategies to address these challenges. One strategy treats sampling time

points as categorical variables and is based on generalized linear models (GLMs). For example,

the software packages DESeq2(Love, Huber, & Anders, 2014), edgeR (McCarthy, Chen, &

Smyth, 2012), and limma (Ritchie et al., 2015) use this strategy. These methods focus on the

magnitude of change instead of the time order of gene expression, and may also suffer from a

relative loss of statistical testing power, especially if many time points are assessed (Fischer,

Theis, & Yosef, 2018). A complementary strategy is to treat time as a continuous variable and fit

the time expression data into a spline-like model. Such methods include DESeq2Spline (DESeq2

adopted with spline model for temporal RNA-seq datasets) fitting, ImpulseDE2 (Fischer et al.,

2018) and Next maSigPro (Nueda, Tarazona, & Conesa, 2014). We found the latter strategy

increased the power of detecting dynamic genes, but it may suffer from a strong model

assumption that oversimplifies the complicated dynamic biological scenarios, such as multiphasic

responses in which gene expression patterns can oscillate between rapid increases and

decreases to reflect initial responses as well as counter-regulatory compensatory responses.

In our present study, we developed TrendCatcher, a versatile R package tailored for time-

course RNA-seq analysis. TrendCatcher uses a framework that combines the smooth spline

ANOVA model and break point searching strategy which identifies inflection points when gene

expression trends reverse. We show that TrendCatcher outperformed the leading methods for

time-course RNA-Seq analysis by using simulated time course data. TrendCatcher has the unique

ability to show users changes in biological pathway enrichment over time, which highlights the

distinct transcriptional programs that are associated with dynamic biological processes. We

applied TrendCatcher to the gene expression profiles of ECs from the brain, heart, and lung EC

and identified the kinetics of injury, repair, and regeneration programs.

Results

TrendCatcher accurately identifies dynamic differentially expressed genes (DDEGs) in

simulated datasets

First, we tested the prediction performance of the TrendCatcher platform (**Figure 1**) using a set

of simulated time course RNA-seq datasets, because simulated data provides defined standards

to assess the accuracy of novel analytical platforms. We considered a comprehensive collection

of gene temporal trajectory patterns to simulate a set of realistic data with biological

characteristics (Bar-Joseph, Gitter, & Simon, 2012). We embedded 1,000 constant trajectories

(non-dynamic genes), 500 linear trajectories (monotonous state transition, i.e. continuously

increasing or continuously decreasing gene expression levels throughout the time course), 500

impulse shaped single break point trajectories (only one temporal inflection point, i.e. up-peak-

down or down-trough-up), 500 two break point trajectories and 500 three break point trajectories

(more complicated dynamic response, e.g., a combination of 2 or more basic types of trajectories).

As shown in **Figure 2A-C**, compared to DESeq2, DESeq2Spline and ImpulseDE2, TrendCatcher

achieved higher accuracy in a mixed simulated dataset. We also tested each model's

performance on a varying number of time points. As shown in **Figure 2D**, TrendCatcher had the

highest prediction accuracy across all time points tested, with accuracies of more than 75%.

We also generated 1,000 gene trajectories for each type of dynamic profile to compare

each model's performance across different trajectory types. As shown in **Figure 2—figure**

supplement 1A-B, for basic trajectory patterns, such as linear trajectory and impulse shaped

trajectory, TrendCatcher achieved accuracies of 80.8% for linear trajectory and 75.3% for impulse

shaped trajectories. As shown in **Figure 2—figure supplement 1C-D**, 2 to 3 break points

represent the combination of basic temporal trends, such as an impulse response leading to

another sustained response. TrendCatcher had higher prediction accuracies of 94.2% for two

break points trajectories and 91.4% for three break points trajectories and thus outperformed the

other three models. Interestingly, DESeq2 and DESeq2Spline works well only when the trajectory

pattern is simple, but we observed their accuracies dropped for more general trajectory types. For

example, DESeq2Spline prediction accuracy dropped to 62.46% in the three break point dynamic

pattern. These results suggest that TrendCatcher can maintain its accuracy even in the setting of

various complex temporal patterns (with multiple break or inflection points), which may occur in

the setting of early biological responses followed by counter-regulatory adaptive responses.

TrendCatcher identifies in situ organ-specific DDEGS and their temporal patterns

composition during inflammatory injury

We applied TrendCatcher to the previously reported RiboTag dataset of the endothelial

translatome in the murine heart, brain and lungs during baseline and endotoxemia (Jambusaria

et al., 2020) to identify dynamic inflammatory response genes. The RiboTag dataset

(GSE136848) is an RNA-seq dataset of *in situ* endothelial cell response to systemic injury induced

by sublethal intraperitoneal bacterial endotoxin lipopolysaccharide (LPS) at six time points and

across three distinct vascular beds during homeostasis and inflammation. We analyzed the global

transcriptomics profiles at 0 hour (baseline), 6 hours, 24 hours, 48 hours, 72 hours and 168 hours

after LPS-injury to the brain, heart and lung. TrendCatcher identified 2,047 significantly dynamic

genes in brain ECs (**Supplementary file 1**), 2,044 dynamic genes in heart ECs (**Supplementary**

file 2), and 2,632 dynamic genes in lung ECs (**Supplementary file 3**). They accounted for 11.3%,

13.4% and 16.2% of the total expression profiles of brain, heart and lung, respectively, suggesting

that over 80% of the expressed genes in the endothelium only fluctuate around baseline

expression level in response to the profound inflammatory injury.

TrendCatcher also assigned trajectory pattern types to all dynamic genes identified from

post LPS-injury, and provided hierarchical pie charts to visualize the composition of trajectory

patterns across the three organs. Each dynamic gene's trajectory pattern was characterized using

a hierarchical scheme, a master-pattern type and a sub-pattern type. A master-pattern type

describes the basic type of gene expression trend, for example, "up-down" represents a biphasic

response, upregulation followed by downregulation. The sub-pattern type shows the time-

dependent dynamic behavior of a gene transcriptional response. For example,

"0h_up_24h_down_168h" belongs to an "up-down" master-pattern type and represents peak

gene upregulation at 24 hours, followed by downregulation until 168 hours.

We observed the same 8 major master-patterns across the three organs (**Figure 3A**) but

with sub-pattern composition shifts (**Figure 3—figure supplement 1-3**), especially for "up-down"

biphasic response with distinct upregulation time. As the inner hierarchical pie charts of **Figure**

3B-D show, we found over 85-95% of all the dynamic genes from brain, heart and lung ECs had

a biphasic response to LPS-inflammation. The majority of the biphasic response genes were

upregulated first, suggesting that there is a clear asymmetry in favor of endothelial gene

upregulation instead of gene repression during inflammatory injury. We found that heart and lung

ECs tended to have more early-upregulated genes at the first 6 hours (**Figure 3C-D)**, as 51.6%

of DDEGs in heart EC and 40% of DDEGs in lung ECs reached the expression peak at 6 hours.

In contrast, in brain ECs only 20.3% of DDEGs reached peak expression at 6 hours. Brain EC

DDEGs tended to gradually increase their expression level, such that 20% and 20.6% reached

peak expression 24 hours and 48 hours after induction of inflammatory LPS-injury.

204
205
Organ-universal and organ-specific inflammatory shift using genes from the same

temporal pattern

We further investigated whether dynamic genes across brain, heart and lung ECs share similar

biological functions by exploring organ-universal inflammation pathways. We assessed the

overlap among the three EC types for each sub-pattern gene trajectory and performed a gene

ontology (GO) enrichment analysis. We found 143 "universal" (i.e. across ECs in all three organs)

dynamic genes (**Figure 4A, Supplementary file 4**) in the early 6-hour activation sub-pattern

("0h_up_6h_down_168h") which were enriched for self-defense-related pathways. These

included response to lipopolysaccharide (19 genes), defense response to virus (27 genes) and

response to interferon-beta (14 genes), response to interferon-gamma (17 genes) (**Figure 4B**).

These shared early activation pathways suggest an organ-universal mechanism to LPS-induced

inflammatory injury, in which ECs across multiple major organs activate self-defense pathways

shortly after injury, with a strong enrichment of interferon response genes.

We also assessed organ-specific pathways that were upregulated early on at 6 hours after

LPS injury. Brain-specific inflammatory responses were enriched for increased neutrophil

migration and regulation of angiogenesis (**Figure 4C**). However, pathways related to Wnt

signaling were downregulated (**Figure 4-figure supplement 1B**), which is important because

Wnt signaling is a key regulator of the integrity of the blood brain barrier (BBB) (Laksitorini,

Yathindranath, Xiong, Hombach-Klonisch, & Miller, 2019). Early-peaking heart-specific EC

responses were highly enriched in ribosome assembly and ribosome biogenesis processes

suggesting global upregulation of protein translation (**Figure 4D**), but showed an early

downregulation in cytokinesis pathways (**Figure 4-figure supplement 1C**). In lung ECs, we

observed the unique upregulation of leukocyte, neutrophil, and granulocyte migration and

chemotaxis, and pathways related to calcium ion homeostasis (**Figure 4E**), whereas endothelial

cell migration gene expression was downregulated (**Figure 4-figure supplement 1D**).

At 24h, brain ECs upregulated ribosome biogenesis processes, mirroring the upregulation

of these processes that we had observed in heart ECs at the earlier 6h time point (**Figure 4-figure**

supplement 2B). At this time point, heart ECs concomitantly upregulated ER stress response

pathways as well as proteasomal degradation pathways (**Figure 4-figure supplement 2C**). In

lung ECs, the bulk of upregulated genes occurred at the 6h time point but we also identified

selected genes peaking later at 24h, however they were also related to pro-inflammatory

processes such as leukocyte migration (**Figure 4-figure supplement 2D**), suggesting that

compared to the heart and the brain, the lung endothelium showed the most prominent sustained

upregulation of pro-inflammatory genes during 6h-24h.

Time-Heatmap discloses organ-specific dynamic programming of endothelial cells

Next, we developed a Time-Heatmap function in TrendCatcher to analyze and visualize the time-dependent dynamic programming. Instead of grouping genes solely based on the trajectory sub-pattern types, which uncovers very few enriched biological pathways, TrendCatcher groups genes based on a derivative manner. It treats neighboring time points as a time window and compares each gene's expression level within the time window to its previous break point expression level. In this way, genes sharing the same expression dynamic within a time window were grouped, independent of their whole dynamic trajectory pattern.

We first applied this approach to the lung EC dataset which has exhibited the most prominent and sustained upregulation of inflammatory genes. As shown in **Figure 5**, TrendCatcher captured distinct phases of lung EC inflammation: 1) inflammatory activation and injury responses, 2) EC proliferation and regeneration as well as 3) EC migration and angiogenesis. The enrichment of host-defense pathways, such as innate immune response and response to bacterium, were strongly activated within the 0h-6h time window, and then subsequently downregulated. This is followed by an increase in cell cycle related pathways over the 24h-48h time window, such as cell division and regulation of mitotic cell cycle phase transition. The upregulation of EC migration and angiogenesis during the late phase (72h-168h) suggests that after inflammatory injury and proliferation, repair of the lung endothelial barrier likely involves EC migration.

We also applied the Time-Heatmap analysis to brain ECs and heart ECs. We observed a similar juxtaposition of early host-defense responses followed by EC proliferation and regeneration (**Figure 5—figure supplement 1 and 2**). However, the regenerative responses in heart ECs showed a significantly delayed activation which was primarily seen at 168h, i.e., 1 week after the injury.

For mechanistic analyses, it is also critical to identify individual genes which drive the biological processes which is why we developed a function allowing for a visualization of fold-

change (logFC) expression of enriched gene sets from selected target GO terms. As shown in

Figure 6A, during the 0-6h time window the GO term defense response was enriched in brain,

heart and lung ECs, but in each vascular bed different sets of dynamic genes were involved. For

brain ECs (**Figure 6A**), the highly dynamic DDEGs were tumor necrosis factor (Tnf), aconitate

decarboxylase 1 (Acod1), interleukin 1 receptor (Il1r2), sialic acid binding Ig-like lectin E (Siglece),

chemokine ligand 1 (Cxcl1), interleukin 6 (Il6) and matrix metallopeptidase 12 (Mmp12); For heart

ECs (**Figure 6B**), the highly DDEGs are interleukin 18 receptor accessory protein (Il18rap), sialic

acid binding Ig-like lectin E (Siglece), interleukin 1 receptor (Il1r2), interleukin 6 (Il6), chemokine

ligand 5 (Cxcl5) and N-formyl peptide receptor 2 (Fpr2). Whereas, for lung ECs (**Figure 6C**), top

dynamic corresponding genes were triggering receptor expressed on myeloid cells 3 (Trem3),

arginase 1 (Arg1), interleukin 1 receptor type 2 (Il1r2), interleukin 10 (Il10), alpha 2-HS

glycoprotein (Ahsg). These organ-specific dynamic EC response genes may provide novel

vascular-bed novel therapeutic targets.

Organ-specific baseline differentially expressed genes (DEGs) temporal behavior

following endotoxemia

We next focused on the top organ-specific highly expressed differentially expressed genes

(DEGs) identified during homeostatic conditions (Jambusaria et al., 2020) to see whether these

organ-specific EC signature genes remain stable over time in the setting of inflammation or

whether they are also dynamic. For each organ, we focused on the study of the top 10 RiboTag

signature genes ranked by their log fold change (logFC) value compared to the other two organs

(**Figure 7A, Supplementary file 5**). We found the following selected genes which constitute the

organ-specific EC signatures during homeostasis are modulated during inflammatory injury. For

brain ECs, 7 out of 10 highly expressed brain EC signature genes were disrupted by the

inflammatory process (**Figure 7A, Supplementary file 5**), most of which showed rapid

downregulation. Only apolipoprotein D followed an "up-down" pattern, which was upregulated in

the first 72 hours after inflammation and returned to baseline after 72 hours. For heart ECs (**Figure**

7B, Supplementary file 5), butyrophilin like 9 (Btnl9) and aquaporin-7(Aqp7) followed a

monotonic increase pattern and increased almost 2-fold at 168 hours after inflammatory

activation. For lung ECs (**Figure 7C, Supplementary file 5**), only resistin like alpha (Retnla) was

upregulated in the first 6 hours and then downregulated to baseline. In contrast to brain ECs, the

top 10 signature genes for heart and lung ECs remained stable during the injury phase suggesting

that brain ECs exhibit the greatest vulnerability to losing their homeostatic brain EC signature in

the face of systemic LPS injury.

<u>Discussion</u>

ECs serve as a highly dynamic interface under normal physiological conditions and their organ-

specific heterogeneity is essential for organ development, homeostasis, and tissue regeneration

(Bautch, 2011). The control of tissue vasculature by ECs is vital for disease states ranging from

inflammation to cancer (Augustin & Koh, 2017). ECs regulate inflammation by preventing

coagulation, controlling blood flow, and trafficking proteins from blood to tissues (Pober & Sessa,

2007). RiboTag RNA-seq profiling enables *in situ* assessment of organ-specific endothelial

translatomes (Cleuren et al., 2019), and by performing this analysis across time points, we were

able to track how each organ's endothelial translatomes changed over time in response to a

stressor. Uncovering how organ-specific ECs respond to inflammation in a time-dependent

manner may uncover new therapeutic targets and the corresponding temporal windows.

Using our computational platform, TrendCatcher, we identified time-dependent endothelial

inflammatory response genes in different organs post endotoxemia, which could be broadly

grouped into shared universal response genes across all three endothelial beds as well as

response genes that were unique to specific endothelial beds. Approximately 85-95% of all organ-

specific dynamic genes responded with biphasic trajectories, and most of the genes responding

to LPS-injury reached their peak within 48 hours. We also observed significant universal

upregulated inflammatory responses across all endothelial beds, which were host-defense-

related pathways. The early host-defense-related dynamic inflammatory markers suggest an

organ-universal self-defense mechanism, in which ECs from various vascular beds activate pro-

inflammatory cytokines and likely facilitate the migration of immune cells to the infected sites.

We observed brain-specific EC downregulation of Wnt pathway genes during early

inflammation. Emerging evidence shows crosstalk between oligodendrocytes and ECs, which is

mediated by signaling through the Wingless and Int-1(WNT)/beta-catenin pathway (Manukjan,

Ahmed, Fulton, Blankesteijn, & Foulquier, 2020). Dysregulation of Wnt signaling results in a

breakdown of blood brain barrier function (BBB) (Laksitorini et al., 2019; Martowicz et al., 2019),

which is why our observation that Wnt pathway genes were predominantly downregulated in brain

ECs is especially intriguing. It may indicate that Wnt downregulation mediates LPS-induced

compromise of BBB function and constitutes a point of vulnerability for the brain endothelium.

Since BBB integrity can be compromised by endotoxemia and is thought to exacerbate

neurodegenerative diseases such as Alzheimer's disease (Barton et al., 2019; L. M. Liu, Wan,

Xia, Kalionis, & Li, 2014) the identification of Wnt pathway gene downregulation in our unbiased

temporal analysis offers important new therapeutic targets for restoring BBB integrity in

endotoxemia.

One of the key features of bacteremia and endotoxemia is the compromise of cardiac

function (Kakihana, Ito, Nakahara, Yamaguchi, & Yasuda, 2016; G. Yucel et al., 2017). On the

one hand, the myocardium responds with hyperdynamic function especially in the setting of

systemic vasodilation to maintain systemic perfusion (Kalogeris, Baines, Krenz, & Korthuis, 2012),

but prolonged endotoxemia or bacteremia can also lead to stress cardiomyopathy (Jafarzadeh,

Thomas, Warren, Gill, & Fraser, 2016; Roivainen et al., 2000). As there is growing evidence that

cardiac ECs are key regulators of cardiac function and that loss of cardiac contractility is in part

due to a compromise of the cardiac endothelium (Potente & Mäkinen, 2017; N. Yucel et al., 2020),

the gene expression changes we observed in cardiac ECs may reflect these distinct phases of

endotoxemia or bacteremia. The early rise in ribosome biogenesis genes is followed by ER stress

pathway upregulation, which suggests that ER stress in the cardiac endothelium may contribute

to myocardial dysfunction during prolonged endotoxemia, and later, at 24 hours, heart EC

activated genes were enriched in the endoplasmic reticulum (ER).

Lung ECs are of special interest during systemic inflammation and endotoxemia because

lung vascular injury is a core pathogenic mechanism in the development of acute respiratory

distress syndrome (ARDS). Leukocytes have been shown to migrate to the lung microvasculature

and interstitium during acute lung injury (ALI) (Claser et al., 2019; Senaldi, Vesin, Chang, Grau,

& Piguet, 1994 Grau, & Piguet, 1994) and the upregulation of calcium signaling increases lung

endothelial permeability (Tiruppathi, Minshall, Paria, Vogel, & Malik, 2002), which further

exacerbates lung injury by leading to unfettered influx of immune cells and fluid into the lung

alveolar space. Importantly, TrendCatcher identified upregulation of lung EC cell cycle pathways

at 48h-72h, which coincides with what was recently reported for the kinetics of lung EC

regeneration after endotoxemic injury (M. L. Liu et al., 2019).

In summary, we have developed a novel computational and visualization tool,

TrendCatcher, to facilitate the interpretation of time-course RNA-seq data. TrendCatcher can

identify dynamic differentially expressed genes (DDEGs), assign dynamic trajectory patterns to

DDEGs, reveal dynamic pattern composition across different experiments, and visualize changes

in biological pathway enrichment across multiple time-points. Compared to existing computational

tools, it achieved higher prediction power in dynamic gene identification from simulated datasets.

Since TrendCatcher assesses distinct time points sequentially and thus maintains a temporal

hierarchy, it may be better suited for the identification of complex temporal trajectories than

current methods. By allowing users to set fold-change thresholds as well as define a priori gene

sets of interest, TrendCatcher is versatile and can be used to mine a variety of temporal RNA-seq

datasets. By applying it to EC transcriptomics, we identified injury and regeneration kinetics

across various vascular-beds which could pave the way for assessing novel therapeutic targets

during defined temporal windows.

Materials and Methods

TrendCatcher framework

The main components of the TrendCatcher framework are shown in **Figure 1**. TrendCatcher

requires two main inputs: the raw count table C of a temporal study with a dimension of $m \times n$,

where m denotes the number of genes and n denotes the number of samples, and a user-defined

baseline time vector T, such as "0 hour". Since samples may have different sequencing depths

and batch effect, TrendCatcher integrates with limma (Ritchie et al., 2015) and provides

preprocessing steps. Based on a user-specified threshold, relatively low abundant genes are

removed from the count table, and reads are normalized and batch effects are removed.

TrendCatcher's core algorithm is composed of five main steps: (a) baseline fluctuation confidence

interval estimation, (b) model dynamic longitudinal count, (c) time point dynamic p-value

calculation, (d) gene-wise dynamic p-value calculation, and (e) break point screening and gene-

wise dynamic pattern assignment. Mathematical details will be expanded in the following sections.

For the output of TrendCatcher, there are mainly two components: a master table and a set of

functions for versatile visualization purposes. The master table contains all the dynamic details of

each single gene, including its dynamic p-value, its break point location time, and its dynamic

trajectory pattern. In addition to the master table, TrendCatcher produces three main types of

visualizations: (a) a figure showing the observed counts and fitted splines of each gene, (b) a

hierarchical pie chart that represents trajectory pattern composition, (c) and a Time-Heatmap of

dynamic biological pathway enrichment change over time.

Baseline fluctuation confidence interval estimation

We assumed that the observed number of RNA-seq reads count from the baseline time (e.g., t=0

hour) $X_{i,t_{baseline}}$ was generated from a negative binomial distribution (Equation 1):

$$X_{i,t_{baseline}} \sim NB(\mu_{i,t_{baseline}}, \varphi_i) \tag{1}$$

398 Where gene $i = 1, \ldots, n$ and $\mu_{i,t_{baseline}}$ is the mean count of gene i at baseline time, and φ_i is the

399 dispersion factor. First, the dispersion factor φ_i was pre-estimated as a constant hyperparameter

400 for each gene with DESeq2(Love et al., 2014), as shown in Equation 2. Here, σ_i is the variance.

$$\sigma_i(t)^2 = \mu_i(t) + \varphi_i * \mu_i(t)^2 \tag{2}$$

401 Then, $\mu_{i,t_{baseline}}$ was estimated using maximum likelihood from Equation 3.

$$\hat{\mu}_{i,t_{baseline}} = argmax\, L_{NB}(u, \varphi) = argmax\, \frac{\Gamma(\varphi + x)}{x!\, \Gamma(\varphi)} \left(\frac{\mu}{\mu + \varphi}\right)^x \left(\frac{\varphi}{\varphi + \mu}\right)^\varphi \tag{3}$$

402 Based on negative binomial distribution and the estimated mean count for baseline time, we

403 constructed the 95% confidence interval $[X_{i,t_{baseline},0.05}, X_{i,t_{baseline},0.95}]$ as the baseline count

404 fluctuation interval (Equation 4(a) and Equation 4(b)).

$$P\left(x_{i,t_{baseline}} \le X_{i,t_{baseline},0.05} \mid \mu_{i,t_{baseline}}, \varphi_i\right) = 0.05 \tag{4a}$$

$$P\left(x_{i,t_{baseline}} \ge X_{i,t_{baseline},0.95} \mid \mu_{i,t_{baseline}}, \varphi_i\right) = 0.95 \tag{4b}$$

406 **Model dynamic longitudinal count estimation**

407 To model the time dependent gene expression value, we applied a smoothing spline ANOVA

408 model (Gu, 2013; Metwally et al., 2018) with a negative binomial family constrain to fits the reads

409 from samples across non-baseline multiple time points. The random variable $X_{i,t(t \ne t_{baseline})}$ is

410 assumed to follow NB distribution in Equation 5, with a positive integer α represents number of

411 failures before the αth success in a sequential of Bernoulli trials, and $p(t) \in (0,1)$ represents the

412 success probability.

$$X_{i,t(t \ne t_{baseline})} \sim NB(\alpha, p(t)) \tag{5}$$

413 The log likelihood given a time-course observed count $x = \{x_{i,t(t \ne t_{baseline})}\}_{i=1,\ldots,n;\, t=1,\ldots,T}$ is

414 calculated as Equation 6.

$$L = log\ L(\alpha, p(t)\ |\ X = x\ )$$

$$= \sum_{t=1}^{T} \sum_{i=1}^{n} [x_{i,t(t \neq t_{baseline})}\ log\ (1 - p(t) + \alpha logp(t) + log\Gamma(\alpha + x_{i,t(t \neq t_{baseline})}) - \quad (6)$$

$$loglog\Gamma(\alpha)] + C$$

415 Taking the logit link and model time effect, we define the logit link η

$$\eta = log\ \frac{p(t)}{1 - p(t)} \qquad (7)$$

416 To allow for flexibility in the estimation of η , and find the best trade-off between goodness of fit

417 and the smoothness of the spline curve, soft constraints of the form $J(\eta)$ is added to the minus

418 log-likelihood, with the smoothing parameter $\lambda > 0$.

$$-L + \lambda\ J(\eta) \qquad (8)$$

419 The solution to the optimization of Equation 8 leads to a smoothing fitting to the reads from

420 samples across different non-baseline time points. The estimated mean of $\mu_{i,t(t \neq t_{baseline})}$ can be

421 estimated using Equation 9.

$$\mu_{i,t(t \neq t_{baseline})} = \hat{\alpha}e^{\hat{\eta}(t)} = \frac{\hat{\alpha}\hat{p}(t)}{1 - \hat{p}(t)} \qquad (9)$$

423 **Gene's dynamic p-value calculation**

424 To calculate gene's non-baseline dynamic signal significance, each gene's non-baseline

425 estimated mean count $\mu_{i,t(t \neq t_{baseline})}$ was tested against the baseline fluctuation interval. Based

426 on Equation 10(a) and Equation 10(b), for each gene at each single non-baseline time point, a

427 dynamic time p-value was calculated.

$$If\ \mu_{i,t(t \neq t_{baseline})} \geq \mu_{i,t_{baseline}},\ then\ p_{i,t} = \int P(x \geq \mu_{i,t(t \neq t_{baseline})}\ |\ \mu_{i,t_{baseline}}, \varphi_i) \qquad (10a)$$

$$If\ \mu_{i,t(t \neq t_{baseline})} \leq \mu_{i,t_{baseline}},\ then\ p_{i,t} = \int P(x \leq \mu_{i,t(t \neq t_{baseline})}\ |\ \mu_{i,t_{baseline}}, \varphi_i) \qquad (10b)$$

428 Then, we applied Fisher's combined probability test method to calculate a gene-wise

429 dynamic p-value (Equation 11):

$$\chi_{i,2t}^2 \sim -2 \sum_{t=0}^{T} ln\,(p_{i,t})$$
(11)

430 **Trajectory pattern assignment**

431 First, we connect all the significant dynamic signal time points, with a p-value threshold

432 less than 0.05. Then we applied a break point searching strategy to capture the gene

433 expression change trend. The definition of break point is defined using Equation 12(a)

434 and Equation 12(b).

$$If\ \mu_{i,t} > \mu_{i,t_{next}}\ \&\ \mu_{i,t} > \mu_{i,t_{previous}}\ ,break\ point\ type\ I.$$
(12a)

$$If\ \mu_{i,t} < \mu_{i,t_{next}}\ \&\ \mu_{i,t} < \mu_{i,t_{previous}}\ ,break\ point\ type\ II.$$
(12a)

There are two types of break points, type I means gene up-regulated followed by a down

regulation and type II means gene expression level down regulated and then followed by

an upregulation. By screening along the break point, the master-pattern and sub-pattern

were assigned to each gene.

Time-Heatmap enrichment analysis

To build the Time-Heatmap for visualizing the biological pathway enrichment change over

time, we designed a window-sliding strategy to capture all the up-regulated or down-

regulated genes for each time interval. If we denote time vector as t_j and $j \in 1,...,T$, each

time interval is denoted as $[t_{j-1}, t_j]$. We found N_{up} up-regulated genes and N_{down} down-

regulated genes within the time window $[t_{j-1}, t_j]$, then a Fisher's exact test was performed

to obtain the Gene Ontology (GO) term enrichment with the corresponding time interval.

Besides GO enrichment analysis (Yu, Wang, Han, & He, 2012), TrendCatcher also

packed Enrichr (Kuleshov et al., 2016) biological pathway databases.

Simulated dataset

We embedded 5 different types of trajectories into the temporal RNA-seq simulated

datasets, including constant trajectory, linear trajectory, impulse trajectory, 2 break-point

trajectory and 3 break-point trajectory. Each type of trajectory was constructed by adding

negative binomial distribution noise to the embedded trajectory count. We mixed 1,000

constant trajectories with the other 4 types of dynamic trajectories (500 from each type).

Then, we used a dynamic filter to trajectories to further distinguish constant trajectories

versus strong dynamic trajectories. We used log2FC range over the time course less than

0.5 as a threshold to relabel constant trajectories and strong dynamic trajectories. We

sampled 6 to 10 different number of time points with evenly time intervals, and randomly

sampled 3 replicates for each time point.

RiboTag dataset preprocessing and batch correction

We downloaded RiboTag RNA-seq dataset from NCBI-GEO website with the access

number GSE136848. We applied limma R package for library normalization and batch

correction to the Brain EC, Heart EC and Lung EC respectively. For gene symbol

annotation, we used ENSEMBL features from biomaRt (Durinck, Spellman, Birney, &

Huber, 2009) package.

<u>Acknowledgments</u>

The studies were supported by NIH grants R01-HL152515 (to JR), R01-HL154538 (to JR), P01-

HL060678(to JR) and R01-HL126516 (to JR).

References

Aird, W. C. (2012). Endothelial cell heterogeneity. *Cold Spring Harb Perspect Med, 2*(1), a006429. doi:10.1101/cshperspect.a006429

Ajami, N. E., Gupta, S., Maurya, M. R., Nguyen, P., Li, J. Y. S., Shyy, J. Y. J., . . . Subramaniam, S. (2017). Systems biology analysis of longitudinal functional response of endothelial cells to shear stress. *Proceedings of the National Academy of Sciences of the United States of America, 114*(41), 10990-10995. doi:10.1073/pnas.1707517114

Augustin, H. G., & Koh, G. Y. (2017). Organotypic vasculature: From descriptive heterogeneity to functional pathophysiology. *Science, 357*(6353). doi:10.1126/science.aal2379

Bar-Joseph, Z., Gitter, A., & Simon, I. (2012). Studying and modelling dynamic biological processes using time-series gene expression data. *Nat Rev Genet, 13*(8), 552-564. doi:10.1038/nrg3244

Barton, S. M., Janve, V. A., McClure, R., Anderson, A., Matsubara, J. A., Gore, J. C., & Pham, W. (2019). Lipopolysaccharide Induced Opening of the Blood Brain Barrier on Aging 5XFAD Mouse Model. *Journal of Alzheimers Disease, 67*(2), 503-513. doi:10.3233/jad-180755

Bautch, V. L. (2011). Stem cells and the vasculature. *Nature Medicine, 17*(11), 1437-1443. doi:10.1038/nm.2539

Chong, D. C., Yu, Z. X., Brighton, H. E., Bear, J. E., & Bautch, V. L. (2017). Tortuous Microvessels Contribute to Wound Healing via Sprouting Angiogenesis. *Arteriosclerosis Thrombosis and Vascular Biology, 37*(10), 1903-+. doi:10.1161/atvbaha.117.309993

Claser, C., Nguee, S. Y. T., Balachander, A., Wu Howland, S., Becht, E., Gunasegaran, B., . . . Renia, L. (2019). Lung endothelial cell antigen cross-presentation to CD8. *Nat Commun, 10*(1), 4241. doi:10.1038/s41467-019-12017-8

Cleuren, A. C. A., van der Ent, M. A., Jiang, H., Hunker, K. L., Yee, A., Siemieniak, D. R., . . . Ginsburg, D. (2019). The in vivo endothelial cell translatome is highly heterogeneous across vascular beds. *Proceedings of the National Academy of Sciences of the United States of America, 116*(47), 23618-23624. doi:10.1073/pnas.1912409116

Durinck, S., Spellman, P. T., Birney, E., & Huber, W. (2009). Mapping identifiers for the integration of genomic datasets with the R/Bioconductor package biomaRt. *Nat Protoc, 4*(8), 1184-1191. doi:10.1038/nprot.2009.97

Eberwine, J., Sul, J. Y., Bartfai, T., & Kim, J. (2014). The promise of single-cell sequencing. *Nat Methods, 11*(1), 25-27. doi:10.1038/nmeth.2769

Fischer, D. S., Theis, F. J., & Yosef, N. (2018). Impulse model-based differential expression analysis of time course sequencing data. *Nucleic Acids Res, 46*(20), e119. doi:10.1093/nar/gky675

Gu, C. (2013). *Smoothing Spline ANOVA Models*: Springer New York.

Gunawardana, H., Romero, T., Yao, N., Heidt, S., Mulder, A., Elashoff, D. A., & Valenzuela, N. M. (2021). Tissue-specific endothelial cell heterogeneity contributes to unequal inflammatory responses. *Sci Rep, 11*(1), 1949. doi:10.1038/s41598-020-80102-w

Haimon, Z., Volaski, A., Orthgiess, J., Boura-Halfon, S., Varol, D., Shemer, A., . . . Jung, S. (2018). Re-evaluating microglia expression profiles using RiboTag and cell isolation strategies. *Nat Immunol, 19*(6), 636-644. doi:10.1038/s41590-018-0110-6

Jafarzadeh, S. R., Thomas, B. S., Warren, D. K., Gill, J., & Fraser, V. J. (2016). Longitudinal Study of the Effects of Bacteremia and Sepsis on 5-year Risk of Cardiovascular Events. *Clinical Infectious Diseases, 63*(4), 495-500. doi:10.1093/cid/ciw320

Jambusaria, A., Hong, Z., Zhang, L., Srivastava, S., Jana, A., Toth, P. T., . . . Rehman, J. (2020). Endothelial heterogeneity across distinct vascular beds during homeostasis and inflammation. *Elife, 9*. doi:10.7554/eLife.51413

Kakihana, Y., Ito, T., Nakahara, M., Yamaguchi, K., & Yasuda, T. (2016). Sepsis-induced myocardial dysfunction: pathophysiology and management. *Journal of Intensive Care, 4*, 10. doi:10.1186/s40560-016-0148-1

Kalogeris, T., Baines, C. P., Krenz, M., & Korthuis, R. J. (2012). CELL BIOLOGY OF ISCHEMIA/REPERFUSION INJURY. In K. W. Jeon (Ed.), *International Review of Cell and Molecular Biology, Vol 298* (Vol. 298, pp. 229-317). San Diego: Elsevier Academic Press Inc.

Kalucka, J., de Rooij, L. P. M. H., Goveia, J., Rohlenova, K., Dumas, S. J., Meta, E., . . . Carmeliet, P. (2020). Single-Cell Transcriptome Atlas of Murine Endothelial Cells. *Cell, 180*(4), 764-779.e720. doi:10.1016/j.cell.2020.01.015

Kuleshov, M. V., Jones, M. R., Rouillard, A. D., Fernandez, N. F., Duan, Q., Wang, Z., . . . Ma'ayan, A. (2016). Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Res, 44*(W1), W90-97. doi:10.1093/nar/gkw377

Laksitorini, M. D., Yathindranath, V., Xiong, W., Hombach-Klonisch, S., & Miller, D. W. (2019). Modulation of Wnt/beta-catenin signaling promotes blood-brain barrier phenotype in cultured brain endothelial cells. *Scientific Reports, 9*, 13. doi:10.1038/s41598-019-56075-w

Liao, J. K. (2013). Linking endothelial dysfunction with endothelial cell activation. *J Clin Invest, 123*(2), 540-541. doi:10.1172/JCI66843

Liu, L. M., Wan, W. B., Xia, S. J., Kalionis, B., & Li, Y. M. (2014). Dysfunctional Wnt/beta-catenin signaling contributes to blood-brain barrier breakdown in Alzheimer's disease. *Neurochemistry International, 75*, 19-25. doi:10.1016/j.neuint.2014.05.004

Liu, M. L., Zhang, L. H., Marsboom, G., Jambusaria, A., Xiong, S. Q., Toth, P. T., . . . Malik, A. B. (2019). Sox17 is required for endothelial regeneration following inflammation-induced vascular injury. *Nature Communications, 10*, 14. doi:10.1038/s41467-019-10134-y

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol, 15*(12), 550. doi:10.1186/s13059-014-0550-8

Lukowski, S. W., Patel, J., Andersen, S. B., Sim, S. L., Wong, H. Y., Tay, J., . . . Khosrotehrani, K. (2019). Single-Cell Transcriptional Profiling of Aortic Endothelium Identifies a Hierarchy from Endovascular Progenitors to Differentiated Cells. *Cell Rep, 27*(9), 2748-2758.e2743. doi:10.1016/j.celrep.2019.04.102

Manukjan, N., Ahmed, Z., Fulton, D., Blankesteijn, W. M., & Foulquier, S. (2020). A Systematic Review of WNT Signaling in Endothelial Cell Oligodendrocyte Interactions: Potential Relevance to Cerebral Small Vessel Disease. *Cells, 9*(6). doi:10.3390/cells9061545

Martowicz, A., Trusohamn, M., Jensen, N., Wisniewska-Kruk, J., Corada, M., Ning, F. C., . . . Nyqvist, D. (2019). Endothelial beta-Catenin Signaling Supports Postnatal Brain and Retinal Angiogenesis by Promoting Sprouting, Tip Cell Formation, and VEGFR (Vascular Endothelial Growth Factor Receptor) 2 Expression. *Arteriosclerosis Thrombosis and Vascular Biology, 39*(11), 2273-2288. doi:10.1161/atvbaha.119.312749

McCarthy, D. J., Chen, Y., & Smyth, G. K. (2012). Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res, 40*(10), 4288-4297. doi:10.1093/nar/gks042

Metwally, A. A., Yang, J., Ascoli, C., Dai, Y., Finn, P. W., & Perkins, D. L. (2018). MetaLonDA: a flexible R package for identifying time intervals of differentially abundant features in metagenomic longitudinal studies. *Microbiome, 6*(1), 32. doi:10.1186/s40168-018-0402-y

Nolan, D. J., Ginsberg, M., Israely, E., Palikuqi, B., Poulos, M. G., James, D., . . . Rafii, S. (2013). Molecular signatures of tissue-specific microvascular endothelial cell heterogeneity in organ maintenance and regeneration. *Dev Cell, 26*(2), 204-219. doi:10.1016/j.devcel.2013.06.017

Nueda, M. J., Tarazona, S., & Conesa, A. (2014). Next maSigPro: updating maSigPro bioconductor package for RNA-seq time series. *Bioinformatics, 30*(18), 2598-2602. doi:10.1093/bioinformatics/btu333

Paik, D. T., Tian, L., Williams, I. M., Rhee, S., Zhang, H., Liu, C., . . . Wu, J. C. (2020). Single-Cell RNA Sequencing Unveils Unique Transcriptomic Signatures of Organ-Specific Endothelial Cells. *Circulation, 142*(19), 1848-1862. doi:10.1161/CIRCULATIONAHA.119.041433

Pober, J. S., & Sessa, W. C. (2007). Evolving functions of endothelial cells in inflammation. *Nat Rev Immunol, 7*(10), 803-815. doi:10.1038/nri2171

Potente, M., & Mäkinen, T. (2017). Vascular heterogeneity and specialization in development and disease. *Nat Rev Mol Cell Biol, 18*(8), 477-494. doi:10.1038/nrm.2017.36

Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., & Smyth, G. K. (2015). limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res, 43*(7), e47. doi:10.1093/nar/gkv007

Roivainen, M., Viik-Kajander, M., Palosuo, T., Toivanen, P., Leinonen, M., Saikku, P., . . . Manttari, M. (2000). Infections, inflammation, and the risk of coronary heart disease. *Circulation, 101*(3), 252-257. doi:10.1161/01.cir.101.3.252

Sabbagh, M. F., Heng, J. S., Luo, C., Castanon, R. G., Nery, J. R., Rattner, A., . . . Nathans, J. (2018). Transcriptional and epigenomic landscapes of CNS and non-CNS vascular endothelial cells. *Elife, 7*. doi:10.7554/eLife.36187

Sanz, E., Yang, L., Su, T., Morris, D. R., McKnight, G. S., & Amieux, P. S. (2009). Cell-type-specific isolation of ribosome-associated mRNA from complex tissues. *Proc Natl Acad Sci U S A, 106*(33), 13939-13944. doi:10.1073/pnas.0907143106

Senaldi, G., Vesin, C., Chang, R., Grau, G. E., & Piguet, P. F. (1994). Role of polymorphonuclear neutrophil leukocytes and their integrin CD11a (LFA-1) in the pathogenesis of severe murine malaria. *Infect Immun, 62*(4), 1144-1149. doi:10.1128/IAI.62.4.1144-1149.1994

Su, T., Stanley, G., Sinha, R., D'Amato, G., Das, S., Rhee, S., . . . Red-Horse, K. (2018). Single-cell analysis of early progenitor cells that build coronary arteries. *Nature, 559*(7714), 356-362. doi:10.1038/s41586-018-0288-7

Tiruppathi, C., Minshall, R. D., Paria, B. C., Vogel, S. M., & Malik, A. B. (2002). Role of Ca2+ signaling in the regulation of endothelial permeability. *Vascul Pharmacol, 39*(4-5), 173-185. doi:10.1016/s1537-1891(03)00007-7

Tombor, L. S., John, D., Glaser, S. F., Luxán, G., Forte, E., Furtado, M., . . . Dimmeler, S. (2021). Single cell sequencing reveals endothelial plasticity with transient mesenchymal activation after myocardial infarction. *Nature Communications, 12*. doi:https://doi.org/10.1038/s41467-021-20905-1

van den Brink, S. C., Sage, F., Vértesy, Á., Spanjaard, B., Peterson-Maduro, J., Baron, C. S., . . . van Oudenaarden, A. (2017). Single-cell sequencing reveals dissociation-induced gene expression in tissue subpopulations. *Nat Methods, 14*(10), 935-936. doi:10.1038/nmeth.4437

Vanlandewijck, M., He, L., Mäe, M. A., Andrae, J., Ando, K., Del Gaudio, F., . . . Betsholtz, C. (2018). A molecular atlas of cell types and zonation in the brain vasculature. *Nature, 554*(7693), 475-480. doi:10.1038/nature25739

Yu, G., Wang, L. G., Han, Y., & He, Q. Y. (2012). clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS, 16*(5), 284-287. doi:10.1089/omi.2011.0118

Yucel, G., Zhao, Z. H., El-Battrawy, I., Lan, H., Lang, S., Li, X., . . . Akin, I. (2017). Lipopolysaccharides induced inflammatory responses and electrophysiological dysfunctions in human-induced pluripotent stem cell derived cardiomyocytes. *Scientific Reports, 7*, 13. doi:10.1038/s41598-017-03147-4

622   Yucel, N., Axsom, J., Yang, Y., Li, L., Rhoades, J. H., & Arany, Z. (2020). Cardiac endothelial
623   cells maintain open chromatin and expression of cardiomyocyte myofibrillar genes. *Elife,*
624   *9.* doi:10.7554/eLife.55730
625
626

627 **<u>Figures</u>**

629
630
631   Figure 1. TrendCatcher Framework
632

634
635
636 **Figure 2.** TrendCatcher prediction performance. (**A**) Prediction accuracy compared to DESeq2,

637 DESeq2Spline and ImpulseDE2 model with mixed types of trajectory patterns. (**B**) Prediction

638 recall from the same simulated dataset with (**A**). (**C**) Prediction precision from the same simulated

639 dataset with (**A**). **(D)** Prediction accuracy compared to DESeq2, DESeq2Spline and ImpulseDE2

640 model with various number of time points, from 6 to 10, using mixed types of trajectory patterns.

	Brain ECs	Heart ECs	Lung ECs
Up_down	1,456	1,712	1,601
Down_up	506	210	660
up	35	94	183
Down_up_down	30	2	16
Up_down_up_down	9	15	92
Up_down_up	8	8	38
Down	2	2	13
Down_up_down_up	1	1	29
Total	2,047	2,044	2,632

Figure 3. Dynamic gene composition comparison among brain, heart and lung ECs. **(A)** Number of different types of dynamic gene trajectories belonged to each master-pattern. **(B)** Hierarchical pie chart of brain EC, inner pie is master-pattern composition, outer pie is sub-pattern composition. **(C)** Hierarchical pie chart of heart EC. **(D)** Hierarchical pie chart of lung EC.

649
Figure 4. Organ-universal inflammatory response, and organ-specific inflammatory response for

trajectory sub-pattern, "0h_up_6h_down_168h". **(A)** Venn diagram of dynamic genes from sub-

pattern "0h_up_6h_down_18h" across brain, heart and lung ECs. **(B)** GO enrichment for the

overlapped 143 dynamic genes. X-axis is the number of genes, y-axis is GO term, p-value is the

enrichment p-value for each GO term. **(C)** GO enrichment analysis for 106 genes only found in

brain EC. **(D)** GO enrichment analysis for 629 genes only found in heart EC. **(E)** GO enrichment

analysis for 702 genes only found in lung EC.

Figure 5. Time-heatmap of lung ECs. Each column represents a time window, each row represents a significantly enriched (p-value <0.05) GO term. Red shade represents GO term are upregulated compared to the previous break point. Blue shade represents GO term are down regulated compared to the previous break point. Percentage number within each tile represents the percentage of genes belongs to the corresponding GO term are dynamic and captured by TrendCatcher. Shade of the color represents the enrichment p-value, the darker the shade is, the more significantly enriched from GO enrichment analysis.

Figure 6. GO-Heatmap. For time window 0h-6h, from selected pathways, including "regulation of defense response", "leukocyte migration", "myeloid leukocyte migration", "leukocyte chemotaxis", "granulocyte chemotaxis", "cellular response to chemokine", "chemokine-mediated signaling pathway", "angiogenesis", "sprouting angiogenesis", "response to bacterium", "leukocyte mediated immunity". Dynamic genes from each organ. **(A)** Brain EC. **(B)** Heart EC. **(C)** Lung EC. The color shade represent log fold change magnitude. Red means up-regulated, blue means down-regulated.

Figure 7. Top 10 organ-specific RiboTag EC signature genes dynamic trajectories. (**A**) Brain ECs top 10 signature genes trajectories, with gene symbol, dynamic p-value from TrendCatcher and its trajectory pattern. (**B**) Heart ECs top 10 signature genes trajectories. (**C**) Lung ECs top 10 signature genes trajectories. *dyn.p.adj* is the adjusted dynamic p-value from TrendCatcher. Black dots are observed count data for each time point for each gene. Red line is smoothed ANOVA spline fitting. Grey lines are baseline fluctuation intervals constructed using constant NB model.

684 **Figure Supplement**
685

686
687
Figure 2-figure supplement 1. TrendCatcher prediction performance across different types of
trajectories. (**A**) Prediction accuracy (left), recall (top-right) and precision (bottom-right),
compared to DESeq2, DESeq2Spline and ImpulseDE2 model with linear trajectories.
(**B**) Prediction accuracy (left), recall (top-right) and precision (bottom-right), compared
to DESeq2, DESeq2Spline and ImpulseDE2 model with impulse trajectories. (**C**) Prediction
accuracy (left), recall (top-right) and precision (bottom-right), compared
to DESeq2, DESeq2Spline and ImpulseDE2 model with two break points trajectories.
(**D**) Prediction accuracy (left), recall (top-right) and precision (bottom-right), compared
to DESeq2, DESeq2Spline and ImpulseDE2 model with three break points trajectories.
697

698
699 **Figure 3–figure supplement 1.** Top sub-pattern dynamic gene trajectories in lung ECs. Ranked
700 by the number of sub-pattern trajectory type. "0h up 48h down" means gene upregulated in the
701 first 48 hours and then downregulated from 48 hours to 168 hours. X-axis is time (hour), y-axis is
702 the log2 count of RNA-seq reads.
703

704
705 **Figure 3–figure supplement 2.** Top sub-pattern dynamic gene trajectories in brain ECs. Ranked

706 by the number of sub-pattern trajectory type. X-axis is time (hour), y-axis is the log2 count of RNA-

707 seq reads

709
710 **Figure 3–figure supplement 3.** Top sub-pattern dynamic gene trajectories in heart ECs. Ranked

711 by the number of sub-pattern trajectory type. X-axis is time (hour), y-axis is the log2 count of RNA-

712 seq reads.

Figure 4–figure supplement 1. Organ-universal inflammatory response, and organ-specific

inflammatory response for trajectory sub-pattern, "0h_down_6h_up_168h". (**A**) Venn diagram of

dynamic genes from sub-pattern "0h_down_6h_up_168h" across brain, heart and lung ECs. (**B**)

GO enrichment analysis for 90 genes only found in brain EC. (**C**) GO enrichment analysis for 101

genes only found in heart EC. (**D**) GO enrichment analysis for 543 genes only found in lung EC.

Figure 4–figure supplement 2. Organ-universal inflammatory response, and organ-specific inflammatory response for trajectory sub-pattern, "0h_up_24h_down_168h". (**A**) Venn diagram of dynamic genes from sub-pattern "0h_up_24h_down_168h" across brain, heart and lung ECs. (**B**) GO enrichment analysis for 307 genes only found in brain EC. (**C**) GO enrichment analysis for 332 genes only found in heart EC. (**D**) GO enrichment analysis for 110 genes only found in lung EC.

Figure 5–figure supplement 1. Time-heatmap of brain ECs. Each column represents a time
window, each row represents a significantly enriched (p-value <0.05) GO term. Red shade
represents GO term are upregulated compared to the previous break point. Blue shade
represents GO term are down regulated compared to the previous break point. Percentage
number within each tile represents the percentage of genes belongs to the corresponding GO
term are dynamic and captured by TrendCatcher. Shade of the color represents the enrichment
p-value, the darker the shade is, the more significantly enriched from GO enrichment analysis.

Figure 5–figure supplement 2. Time-heatmap of heart ECs. Each column represents a time window, each row represents a significantly enriched (p-value <0.05) GO term. Red shade represents GO term are upregulated compared to the previous break point. Blue shade represents GO term are down regulated compared to the previous break point. Percentage number within each tile represents the percentage of genes belongs to the corresponding GO term are dynamic and captured by TrendCatcher. Shade of the color represents the enrichment p-value, the darker the shade is, the more significantly enriched from GO enrichment analysis.