Comprehensive PAM prediction for CRISPR-Cas systems reveals evidence for spacer sharing, preferred strand targeting and conserved links with CRISPR repeats

Jochem NA Vink[1,2], Jan HL Baijens[1,2] and Stan JJ Brouns[1,2]

[1] Department of Bionanoscience, Delft University of Technology, Delft, Netherlands

[2] Kavli Institute of Nanoscience, Delft, Netherlands

Abstract

The adaptive CRISPR-Cas immune system stores sequences from past invaders as spacers in CRISPR arrays and thereby provides direct evidence that links invaders to hosts. Mapping CRISPR spacers has revealed many aspects of CRISPR biology, including target requirements such as the protospacer adjacent motif (PAM). However, studies have so far been limited by a low number of mapped spacers in the database. By using vast metagenomic sequence databases, we mapped one third (~70,000) of more than 200,000 unique CRISPR spacers from a variety of microbes, and derived a catalog of more than one hundred unique PAM sequences associated with specific CRISPR subtypes. These PAMs were further used to correctly assign the orientation of CRISPR arrays, revealing conserved patterns between the last nucleotides of the CRISPR repeat and PAM. From the curated CRISPR arrays dataset we could also deduce CRISPR subtype specific preferences for targeting either template or coding strand of open reading frames. While some DNA-targeting systems (e.g. Type I-E and Type II systems) prefer the template strand and avoid mRNA, other DNA- and RNA-targeting systems (i.e. Type I-A, I-B and Type III systems) prefer the coding strand and mRNA. In addition, we found large scale evidence that both CRISPR adaptation machinery and CRISPR arrays are shared between different CRISPR-Cas systems.

23 This could lead to simultaneous DNA- and RNA targeting of invaders, which may be effective at

24 combating mobile genetic invaders.

Introduction

CRISPR, an adaptive immune system provides heritable defence in the form of spacers, which are short

nucleic acid sequences (28-36 bp) obtained from previous encounters with mobile genetic elements

(MGE). These are stored in the bacterial or archaeal chromosome in CRISPR arrays (Jackson et al.,

2017). CRISPR arrays contain spacers flanked on both sides by repeat sequences (~30 bp) and are

transcribed as a single RNA, and subsequently processed into multiple crRNAs. crRNAs can be loaded

into effector complexes formed by Cas proteins, that subsequently scan the cell for nucleic acid targets.

Base pairing between the spacer and target nucleic acids (protospacer) allows the specific binding of

effector complexes to targets, which are then destroyed (Brouns et al., 2008; Marraffini, 2015). CRISPR-

Cas systems are widespread in bacteria and archaea, with 42% of bacterial and 85% of archaeal genomes

containing a CRISPR system (Makarova et al., 2020).

Both acquisition of new spacers (CRISPR adaptation) and target inactivation (CRISPR interference) are

carried out by specialized sets of Cas proteins. *Cas* genes likely have originated from Casposons

(Krupovic et al., 2014), a family of self-replicating transposons, and have since evolved many new genes

and gene variants (Makarova et al., 2020). Based on the evolutionary classification of their *cas* genes,

there are two classes of CRISPR-Cas systems. Class I systems contain crRNA-effector complexes made

up of multiple subunits, while effector complexes of Class II systems are encoded by a single *cas* gene

(Makarova et al., 2020). The two classes are further divided into six types, where each type is further

divided into subtypes. The different types and subtypes do not occur homogeneously in nature, with

Class II systems being nearly exclusive to bacteria (Makarova et al., 2020). More than 95% of CRISPR

systems found in complete genomes are one of the first three types: Type I, II or III (Pourcel et al.,

2020).

CRISPR systems can be studied on a mechanistic or on a functional level. Mechanistic features describe

how CRISPR systems are able to fulfil their role. The mechanisms through which CRISPR systems

operate are diverse. For example, some CRISPR systems defend the cell by targeting DNA (e.g. Type

I, II, IV and V), whereas other CRISPR types target invader RNA (e.g. Type III and VI) (Makarova et

al., 2020). Another important mechanistic feature is the presence of a protospacer adjacent motif (PAM),

which DNA-targeting systems require to differentiate self from non-self (Gleditzsch et al., 2019; Hale

et al., 2009; Mojica et al., 2009). Furthermore, the PAM is an important feature in the target search

process of DNA-targeting systems within the cell (Vink et al., 2020; Xue et al., 2017). This motif

sequence flanking the crRNA-pairing site, between one and five nucleotides long, not only differs

between subtypes, but can also differ between *cas* gene orthologs within the same subtype, for example

Cas9 variants (Gasiunas et al., 2020).

Functional features describe what purposes CRISPR systems fulfil within the cell. There is evidence for

some CRISPR functioning beyond adaptive immunity (Westra et al., 2014), however even within the

context of an adaptive immune system, CRISPR systems can serve different roles (e.g. as a first line of

defence, or as an activator of other immune system pathways). This can be a reason why 23% of

genomes with CRISPR systems contain more than one subtype (Bernheim et al., 2020), in spite of their

costs (Nobrega et al., 2020; Vale et al., 2015). There are preferred combinations of certain subtypes,

suggesting that there is an added benefit of having a specific combination of different subtypes present

in the cell. The added benefit might consist of cooperativity between systems by formation of different

lines of defence, avoidance of type-specific CRISPR inhibition by MGE or coupling of abortive

infections mechanisms (Bernheim et al., 2020; Hoikkala et al., 2021; Pawluk et al., 2017; Silas et al.,

2017). On the other hand, some CRISPR systems are specialized to protect from certain invaders, which

may require multiple co-occurring systems to be present in a single genome to protect from different

types of invaders. Type IV systems that co-occur together with Type I systems primarily target plasmids

(Pinilla-Redondo et al., 2020) and Type III systems have been shown to be able to target a class of

phages that other Type I and V systems cannot (Malone et al., 2020; S. D. Mendoza et al., 2020),

indicating that specialization in targets is a potential reason for co-occurrence of different subtypes.

Through cooperation and specialization, co-occurring subtypes can function complementary.

The functional and mechanistic features described above have been demonstrated experimentally for

several microbial model systems, and these are often of specific interest to applications such as genome-

editing. High-throughput assays to identify the PAM of CRISPR systems have been developed, but

remain laborious (Gasiunas et al., 2020; Walton et al., 2021). The full diversity of PAM and other

mechanistic and functional features of CRISPR-Cas systems in nature remain understudied. To improve

our knowledge on mechanistic and functional features of single and co-occurring CRISPR systems

beyond the model organisms, we relied on vast metagenomic sequence databases to computationally

find targets for spacers from diverse bacteria and archaea. We mapped a third of the unique spacers to a

target in publicly available metagenome sequence databases. We used the flanking regions of found

spacer targets to build an initial PAM catalog of more than a hundred unique PAMs, and for more than

half of the spacers in CRISPRCasDB (Pourcel et al., 2020). This was then employed to assign the correct

orientation of transcription of CRISPR arrays, giving access to target strand information of invaders,

further improving PAM predictions, and uncovering conserved links between repeat ends and PAM.

Through the quantification of the spacers targeting template or coding strands we found that the

preference for one of these strands is subtype specific, and indicates that some DNA-targeting systems

(Type I-E, Type II-A and Type II-C) avoid RNA while other DNA- and RNA-targeting systems

preferentially target RNA (Type I-A, Type I-B and Type III systems). We found spacers in co-occurring

CRISPR systems to be compatible with both PAM and strand requirements, indicating that they may be

shared between systems and will lead to both DNA and RNA targeting. Lastly, we identified three

categories of multi-effector compatible spacers, which meet the PAM and strand requirements of co-

occurring DNA and RNA-targeting systems.

Results

Blast analysis finds matches for 32% of spacers from CRISPRCasDb

The first step in our analysis was to select a set of CRISPR spacers and find potential matches to these

sequences in DNA sequence databases. To this end, we selected the previously described

CRISPRCasDB, which contained all spacers from 4266 complete bacterial and archaeal genomes

(Pourcel et al., 2020). The spacers from CRISPRCasDB were then mapped to sequences from the NCBI

nucleotide database as well as metagenomic databases with high number of prokaryotic or their virus

sequences. Matches between spacers and sequences from the databases were found using BLASTn

(Altschul et al., 1990). The matches were then filtered using an optimized approach which increased the

number of matches while keeping the false positives to a minimum (Methods, Supplementary figure

1A). As indication of the false positive rate, we determined that for the matches found in NCBI

nucleotide database 1% were eukaryotic or eukaryotic virus sequences, with 10% of matches in

prokaryotic viral sequences and the majority (88%) corresponding to prokaryotic genome sequences

(Supplementary figure 1A). This specificity towards prokaryotic sequences in a database that contains

predominantly (83%) eukaryotic sequences shows that even though false positive hits cannot be

excluded, the false positive rate is low.

From the 221,850 total unique spacers analysed, this optimized filtering approach resulted in 72,099

spacers (32% of total) with at least one match (Figure 1A), of which 31,327 spacers (15% of total) had

a match in the NCBI nucleotide database (Figure 1B). The fraction of spacers with matches differed

greatly between different genera, with *Streptococcus*, *Pseudomonas* and *Staphylococcus* among the

genera with the highest fraction of matches (77%, 69% and 64% respectively) and *Calothrix*, *Nostoc*

and *Thermosipho* among the lowest (4%, 4% and 3% respectively) (Figure 1C). Genera with high spacer

matches typically occurred in well-sampled environments (human-associated), whereas the genera with

lower matches occurred in what appear to be poorly sampled environments (soil, oceanic). A previous

study (Shmakov et al., 2017) which looked for spacer matches in the NCBI nucleotide database found

matches for 7% of spacers, using a more stringent 95% sequence identity and 95% coverage cut off as

filtering thresholds. This difference in the fraction of spacers with matches in the NCBI nucleotide

database indicates the added benefit and importance of our more sensitive filtering process.

Additionally, the number of sequences in the database has increased in recent years from ~230 billion

to ~700 billion bases. The most important factor for the increase in the number of spacers with matches

however was the use of metagenomic databases, as the majority of unique spacer matches derived from

these databases (Figure 1B, Supplementary Figure 1B).

To find the subtypes of the spacers, we aligned the CRISPR repeat sequences to repeat sequences with

known subtypes, based on the method described by Bernheim et al., 2020. With the exception of subtype

II-B for which we extracted 453 spacers, all analysed subtypes from Type I, II and III systems contained

more than a thousand spacers (Figure 1D). An exceptionally high fraction of spacers with matches was

found for subtypes II-A (63%) and II-C (53%), while subtype I-A, subtype I-D, and Type III subtypes

had notably lower fractions of spacer matches than average (15%, 11% and 20% respectively). The

differences in fractions of matches found between subtypes may be due to their phylogenetic

distributions, where well-sampled genera have different subtypes than poorly sampled genera (see

above). However, even within well-sampled genera the fraction of spacers with matches differs between

subtypes, with Type III subtypes having fewer hits on average (22%) than other subtypes (38%). Overall,

the large number of spacers with matches revealed sets of sequences that were targeted by each CRISPR-

Cas subtype, which were then used to study mechanistic and functional aspects of CRISPR defence.

Alignment of protospacer flanks reveals 114 unique subtype-specific PAMs covering

55% of spacers

One of the important mechanistic features of CRISPR defence for DNA targeting systems (type I, II, IV

and VI) is PAM recognition (Deveau et al., 2008; Mojica et al., 2009; Shah et al., 2013). The first PAM

was discovered in the alignment of bacteriophage sequences that were targeted by *Streptococcus* spacers

(Bolotin et al., 2005). Later studies revealed more PAMs or the effect of mutant versions of the PAM

(Anders et al., 2014; Fischer et al., 2012; Leenay et al., 2016; Musharova et al., 2019). We expand on

these known PAMs that are limited to well-studied organisms by predicting new PAMs based on the

alignment of the flanks of spacer matches (protospacers). The potential of this method for large scale

PAM predictions was shown in a previous bioinformatics study (Mendoza & Trinh, 2018), with a key

limiting factor being the number of spacers with matching targets. It was also previously shown that

PAMs, acquisition machinery and repeat clusters co-evolve (Shah et al., 2013). We therefore increased

the number of spacers with matches within one group by clustering spacers based on repeat similarity

(>90% nucleotide identity and same repeat length). The sensitivity of PAM detection depends on the

information content of the nucleotide positions of the PAM (signal) compared to the information content

of the other flanking positions (noise). We found that clustering based on repeat similarity increased

signal to noise ratio for PAM detection compared to clustering based on species-subtype (e.g.

Escherichia coli I-E) or genus-subtype (e.g. *Pseudomonas* I-F). We furthermore found that spacers

originating from organisms with very high or low GC-contents, displayed increased noise. We thus

further increased the signal to noise ratio by adjusting the expected frequency of flanking nucleotides

based on the average GC-content of the spacers within the cluster (Supplementary Figure 2). The flanks

of unique hits within each cluster can subsequently be aligned, and with enough spacer hits, the

information content reliably reveals the PAM sequence and position relative to the protospacer (Figure

2).

This clustering approach together with our large number of hits led to a PAM prediction for 123,144

spacers (55% of all spacers; Supplementary File 1 and 2). For Type I and Type IV the PAM is known

to occur in the 5' (upstream) flank of the protospacer, while Type II systems have their PAM in the 3'

(downstream) flank of the protospacer (Jackson et al., 2017) (Figure 2A). This well characterized feature

of the PAM therefore allows the unique possibility to correctly orient CRISPR arrays given the rules

described above. To measure the accuracy of CRISPR array orientation predictions, we compared

predictions to experimentally determined orientations from a recent study using transcriptome

sequencing (TOP) to determine the direction of transcription of arrays (Houenoussi et al., 2020). The

7968 experimentally inferred spacer orientations were the same as our predictions in 85% of cases, while

only 33% of TOP predicted spacer orientations were the same as the CRISPRCasDb prediction

(Supplementary data). We furthermore found that many Type I and Type III repeats for which we

predicted the orientation based on the PAM, contained the 3'-end motif ATTGAAAC of their repeat

(Supplementary Figure 3) described previously (Lange et al., 2013). This conserved motif is transcribed

and forms the 5' handle of the crRNA and is held by crRNA-effector complexes. Altogether, these

findings indicate that the position of the PAM is a reliable indicator for the orientation of the CRISPR

array, and can be used to annotate CRISPR array information, giving access to features such as spacer

acquisition chronology and strandedness.

Type I PAMs are shorter and closer to the protospacer than Type II PAMs

Sequence logos of alignments of Type I (Figure 2A) recover previously known PAMs including the subtype I-E AWG PAM found in *Escherichia* and subtype I-F CC PAM found in *Pseudomonas* (Leenay & Beisel, 2017), but also many previously undescribed PAMs. They are generally short (2-3 nt) and are well defined (high information content/bit score). For Type II PAMs, we found both short, well defined PAM motifs (such as *Streptococcus* II-A) as well as longer PAMs with less conserved PAM motifs (Figure 2B). Poorly conserved PAM motifs could be caused by a variation of PAMs used within the same repeat cluster or by the promiscuity of PAM recognition in Type II systems (Crawley et al., 2018). Additionally, many Type II PAMs consisted of multiple consecutive nucleotides of the same kind in a row, such as NAAAA (*Capnocytophaga* II-C). A low nucleotide conservation and repetitive nucleotide identity of a sequence motif can be caused by ambiguity in PAM distance to the protospacer, as this ambiguity will spread the nucleotide conservation over a larger range of positions from the protospacer. PAMs were found with the closest conserved nucleotide ranging 2 to 5 nucleotides away from the protospacer. The first nucleotide position on the 3' end of the protospacer was always found to be an N for Type II PAMs. For a minority of subtype II-A and subtype II-C repeat clusters, a distinct lack of PAM was found (Supplementary figure 4B). As some Cas9 proteins from subtypes II-A and II-C can target RNA independently from a PAM sequence (Strutt et al., 2018), this RNA targeting could contribute to natural PAM-less variants that may inspire engineered PAM-less variants (Walton et al., 2020). Alternatively, PAM usage might be highly variable in these specific repeat clusters and could therefore obscure distinct PAM motifs. Overall, we found 114 unique PAMs (PAM-subtype combinations; Supplementary Data File 1), of which 43 PAMs in Type I systems (Supplementary Table 1), with each subtype containing at least two different PAMs and subtype I-B containing 12 different PAMs.

43% of Type III repeat clusters contain a PAM

Like the PAM-less Type II variants, some Type III repeat clusters were devoid of a PAM. This is expected, as RNA-targeting systems do not require a PAM to find a target (Figure 2C, Supplementary Figure 4), and rely on the Protospacer Flanking Sequence (PFS) to avoid self targeting (Deng et al.,

2013; Elmore et al., 2016). Interestingly, other repeat clusters contained PAMs that appeared to be the

same as Type I PAMs, which raised the question, why these clusters contained a PAM. We compared

the PAM detection frequency for clusters with at least 25 unique spacer hits (Figure 2D). For Type I

subtypes and subtype II-A, the majority of repeat clusters have a defined PAM, whereas for Type II-C

and Type III systems the number of PAM-containing repeat clusters was lower, with Type III-A having

the lowest (16%) and III-B the highest (56%) fraction of PAM-containing repeat clusters in Type III

systems. As it was previously shown that Type III systems often lack their own acquisition machinery

(Makarova et al., 2015), we hypothesized that the PAM found in Type III repeat clusters originates from

the spacer acquisition machinery that Type I systems share with Type III systems. We observed that the

PAM frequency in Type III clusters that lack their own acquisition machinery is high (95%; Figure 2E),

whereas the PAM frequency is low in Type III clusters that contain their own *cas1-cas2* genes (8%).

This supports the hypothesis that the PAM in Type III arrays originates from Type I spacer acquisition

modules functioning in *trans*.

Conserved patterns between PAM and repeats

PAMs usually differ from the ends of CRISPR repeats, which allows for self-nonself discrimination

(Leenay et al., 2016; Mojica et al., 2009; Westra et al., 2013). Type III and other RNA-targeting CRISPR

systems do not require a PAM, but many do require mismatching between the repeat end and the

protospacer flanking sequence (PFS) (Johnson et al., 2019; Marraffini & Sontheimer, 2010). Given these

previous observations, we wanted to investigate if there are conserved links between repeat ends and

PAM of individual systems (Figure 3A), and whether Type III PAMs that originate from Type I spacer

acquisition modules are also compatible with Type III PFS requirements.

We collected all unique repeat-PAM sequence combinations in our dataset and compared the repeat

nucleotide with the corresponding PAM nucleotide in each position. For Type I systems (Figure 3B) we

found that the -3 and -2 nucleotide of the repeat can be a strong predictor of the corresponding PAM

nucleotide, where a -3C in the repeat would lead to a -3A in the PAM, -3G to -3T, -3T to -3A. At the

middle position a -2C would lead to a -2A in the PAM. (Figure 3B). The most common -2 and -3 repeat

nucleotide is an A, in which case the PAM nucleotide mostly is either a T or a C. For the -1 position,

the nucleotide identity of the PAM sequence cannot be predicted directly from the repeat sequence.

For Type II systems, most nucleotide positions can accommodate two or three PAM nucleotides

(Supplementary Figure 5A). In +2 and +3 positions, the most common repeat nucleotide (T),

accommodates either an A or G PAM nucleotide, which is analogous to the most common nucleotide in

Type I systems (-3 and -2 adenine), which tends to co-occur with a C or T PAM nucleotide. For Type

III systems, the variation of repeat nucleotides is smaller, but generally similar combinations are found

as in Type I systems (Supplementary Figure 5B). Overall, the most conserved repeat-PAM co-

occurrence patterns are found in the -2 and -3 positions of the Type I and Type III arrays.

These co-occurrence patterns suggest that in most cases the PAM that is used and selected for differs

from the repeat. However previous studies have shown that in some cases, part of the repeat sequence

is PAM -derived (Swarts et al., 2012). We then asked in what CRISPR subtypes the PAM matches the

corresponding repeat nucleotide for each of the spacer flanking positions. When we counted the

occurrence of a matching PAM, we found that this only occurred frequently in the -1 position of Type

I-C (35%) and Type I-E (48%; Figure 3C). We found that these matches are associated with repeats that

have TTC PAMs in Type I-C and AAG PAMs in Type I-E, which could indicate that the C of Type I-

C repeat sequences is PAM-derived, as was similarly demonstrated for the G of AAG PAMs in Type I-

E (Swarts et al., 2012).

In other positions and CRISPR types, >98% of the repeat-PAM combinations did not match each other,

which shows that the general patterns between repeats and PAMs, and perhaps mechanism of self- vs

nonself discrimination is conserved in all subtypes. In Type III systems all cases demonstrate

mismatches between PAM and repeat, which is a requirement of functional Type III spacers (Johnson

et al., 2019; Marraffini & Sontheimer, 2010). This finding demonstrates that the PAMs of Type III array

spacers acquired with Type I acquisition modules are compatible with PFS requirements of Type III

systems.

Strand bias for the template or coding strand is subtype specific

Our method has revealed a large number of newly identified PAMs and has shown that Type III systems which lack their own acquisition machinery and co-occur with Type I systems, almost always contain a PAM. The presence of a PAM in these systems could enable Type I systems to use the spacers stored in Type III arrays as they are compatible with the PAM requirements of Type I effector complexes. Furthermore Type III effector complexes could benefit from a PAM-selecting acquisition module, as it excludes spacers with repeat-PAM matches (Figure 3C).

Besides the PFS, another requirement for type III spacers is that the spacer comes from the correct strand, as these complexes can only bind to the RNA transcripts. We wondered whether some species indeed use Type I and III dual functionality CRISPR arrays, as PAM-dependent DNA targeting and PAM-independent mRNA targeting are not mutually exclusive. We therefore asked whether spacers of DNA-targeting systems are also compatible with Type III surveillance complexes, if they happened to be picked from the correct strand.

To determine the potential ability of crRNA to target RNA, we measured the strand bias by counting the spacers that targeted the coding or template strand of predicted open reading frames (ORFs) (Figure 4A). As spacers targeting the template strand are unable to base pair the transcribed RNA, the fraction of spacers targeting the coding strand serves as an estimate of the RNA targeting ability of the crRNA. For example, in *Moraxella* IIIB arrays, a significant bias for the coding strand was found (88%, $p<e^{-11}$) (Figure 4B). This bias allows Type III effectors carrying crRNA from those spacers to bind to their target RNA. However, also I-C spacers in *Moraxella*, for whose effectors this is not strictly required, show significant bias for the coding strand ($p<e^{-3}$), indicating a selection for RNA-targeting spacers.

For *Escherichia* subtype I-E, 977 spacer matches inside ORFs were found, of which 611 (63%) targeted the template strand (Figure 4C), showing a significant bias for targeting the template strand ($p<e^{-14}$) potentially avoiding RNA. No significant strand bias was found for *Escherichia* subtype I-F (43% template strand, p=0.11), suggesting that strand bias is CRISPR subtype specific.

Analysis of our complete dataset revealed general trends in the strand preferences for each subtype

(Figure 4D, E). The strongest strand bias was found in Type III systems with an average of 65% of the

spacers matching the coding strand (coding strand:template strand ~ 2:1). This result demonstrates that

there is selection in Type III systems for spacers to target the transcribed RNA. This selection can

originate at the adaptation stage by dedicated adaptation machinery selecting from RNA/coding strands

such as RT-Cas1 (Silas et al., 2016) or at the interference stage, where only functional RNA-targeting

spacers are retained in the population (Artamonova et al., 2020). The strand biases we found are

consistent with our curated CRISPR array orientation predictions, because an incorrect CRISPR array

orientation prediction would obscure strand specific targeting. Type I-A and Type I-B also displayed

significant strand bias for the coding strand although at lower levels (60% and 55%; $p<e^{-9}$ and $p<e^{-14}$

respectively).

Contrary to the Type III, Type I-A and I-B systems, we found a significant strand bias towards the

template strand in in subtype I-E, Type IV and Type II systems, with the strongest bias found in subtype

II-A (59%) and subtype I-E (57%). Given the high number of spacers in these groups the chance of

observing this bias by chance is small ($p <e^{-23}$ and $p <e^{-69}$ respectively), again suggesting avoidance of

RNA.

Co-occurrence of Type I and Type III systems lead to PAM and strand targeting

compatibility

As we noticed that Type III spacers were compatible with Type I PAMs in multiple cases, we next asked

whether Type I spacers are compatible with RNA targeting in microbes with co-occurring Type I and

III systems. We measured the strand bias of Type I spacers in genomes containing either combination

of Type I, Type II and Type III surveillance complexes (Figure 4F). No significant strand bias was found

for Type I spacers in the presence of Type I and/or Type II surveillance complexes. However, in the

presence of Type I and Type III surveillance complexes, Type I spacers had a slight but significant

coding strand bias (55%, $p<e^{-14}$). This might be caused by increased selection pressure to keep RNA

targeting spacers in the presence of RNA targeting surveillance complexes. This would suggest that

spacers are selected to be compatible for both Type I and Type III effector complexes in such situations.

For Type II spacers, the presence of Type III did not significantly change the strand bias (Figure 4G).

Given the natural tendency of Type II spacers to bias towards the template strand (Figure 4E), these

findings suggest that Type II spacers are less compatible with co-occurring Type III effector complexes

than Type I spacers.

Three distinct categories of co-occurring multi-effector compatible arrays exist

The findings above indicate that subtype specific preferences exist for either the template or coding

strand of the DNA. These preferences might enable or preclude compatibility between the spacers of

co-occurring subtypes. We categorised all multi-effector compatible arrays that can be used by effector

complexes from different subtypes. This means for co-occurring DNA-targeting systems these arrays

need to have a PAM that can be used in both systems, whereas for co-occurrence of a DNA- target

CRISPR-Cas system with an RNA targeting system, the arrays present in the genome need to both have

the correct PAM and have a bias for the coding strand.

Overall, we can distinguish three main categories of co-occurring CRISPR-Cas systems in which spacers

are compatible for multiple effectors (Figure 5A, Supplementary File 3).

The first category, exemplified by *Paenibacillus larvae* SAG 10367, consists of two co-occurring DNA-

targeting systems which have their own adaptation machinery and their own repeat sequences. This is

the smallest category and has been found in seven genomes (Figure 5A, B; Type I-A-Type I-B: 5, Type

I-B-Type I-C: 2). We furthermore found 45 genomes of *Listeria*, which contain a Type I-B system and

a Type II-A system from which the spacers have PAMs that might be compatible with both I-B and II-

A effector complexes if the arrays are transcribed bi-directionally (CCN and NGG).

The second category, exemplified by *Clostridium botulinum* MAP 5, consists of a co-occurring DNA-

targeting and RNA-targeting, with distinct repeat sequences but a commonly shared acquisition

machinery (Figure 5A, C). We have only found evidence for multi-effector compatibility in co-occurring

Type I and Type III systems. The Type I array in this category has a strand bias which indicates that the

Type III effector complexes can use these spacers whereas the Type III arrays have the same PAM

sequence as the Type I arrays allowing the Type I effector complexes to use these spacers. This category

has been found in 17 genomes, mostly containing a Type I-B and a Type III system but also two genomes

were found with a Type I-E and a Type III system.

The third category, exemplified by *Sulfolobus acidocaldarius* SUSAZ, consists of a co-occurring DNA-

targeting and RNA-targeting system, with shared repeat sequences and shared acquisition machinery

(Figure 5A, D). In this category the single repeat cluster present has a PAM and coding strand bias. This

is the most common category of multi-effector compatible arrays which we detected in 85 genomes. It

consists of co-occurring Type III systems with either a Type I-B (82%) but also Type I-A (13%) and

Type I-C (5%).

Taken together, our data indicate that multi-effector compatible arrays are most prevalent between Type

I and Type III systems. Within the Type I systems, the most common subtype to use multi-effector

compatible arrays is Type I-B, but also Type I-A, Type I-C and Type I-E use these arrays. The Type III

systems that use compatible arrays lack their own adaptation machinery, however repeat clusters in these

co-occurring systems display a strand bias that suggests selection for RNA-targeting spacers. The

information content is similarly strong for PAMs in Type III arrays as in Type I arrays, which

demonstrates that the PAM is equally strong selected for Type I as shared Type III arrays.

Discussion

In this study we have matched CRISPR spacers of complete genomes of bacteria and archaea with their

targets in (meta)genome databases and subsequently analysed the genomic flanks of the protospacers.

We computationally found targets for 32% of CRISPR spacers from thousands of bacterial and archaeal

genomes. This is a major increase in spacer targets compared to previous studies and is due to our

sensitive filtering process and use of metagenomic databases (Shmakov et al., 2017). We found that

Type III spacers had the highest fraction of unknown targets of any CRISPR type. This was not solely

caused by the phylogenetic or environmental occurrence of Type III systems, because the fraction of

Type III spacers with unknown targets within a genus was typically higher than that of other types. This

means that the targets of Type III systems are either under sampled, or that Type III spacers contain

more mismatches to their targets, making them harder to find computationally. Recently, a single new

study doubled the number of known RNA viruses including phages (Wolf et al., 2020), while another

study greatly increased the number of known single-stranded RNA phages (Callanan et al., 2020),

indicating that RNA phages have been poorly sampled. We predict the fraction of spacers with matches

to increase with increasing numbers of available metagenomic data, especially including more RNA

viruses and more data from poorly sampled environments.

By analysing the flanks of the spacer hits in great depth, we have generated a vast catalog of PAM

sequences for each CRISPR repeat cluster. The repeat sequence is a good predictor of parts of the PAM

sequence, and outperformed clustering based on genus-subtype classifications. This finding is

corroborated by the position-wise comparison of PAM and repeat nucleotides, which shows certain

repeat nucleotides predict PAM nucleotides. This may be helpful to either predict the PAM from scratch,

or to further experimentally determine the PAM while reducing the degeneracy at certain positions,

limiting the predicted PAM sequence space. The mismatch between repeat and PAM nucleotides

generally holds, except for the Type I-E and Type I-C, where for some repeat clusters the repeat

nucleotide matches the PAM at the -1 position. The most common PAMs of these systems (TTC for I-

C; AAG for I-E) are also complementary to each other. These findings indicate Type I-C systems could

have a similar mechanism of spacer acquisition with a PAM-derived last repeat nucleotide as in Type I-

E (Swarts et al., 2012), even though these systems do not share related Cas1 proteins (Makarova et al.,

2011) or repeat structures (Lange et al., 2013).

The PAM catalog can be used to predict the PAM for arrays in newly sequenced genomes and

metagenomic contigs if they contain repeats that are closely related to the repeats in our database, which

gives access to unexplored mechanistic and biotechnological potential. For repeats that are not in our

database, the nucleotide identities of the repeat in the spacer flanking positions can be used to predict,

with lesser certainty, which PAM it could have and select certain CRISPR systems of interest for further

study.

Furthermore, the position of the PAM in the target is a reliable indicator for the orientation of

transcription of CRISPR arrays. Correct prediction of transcription of CRISPR arrays gives access to

measuring chronology of invader encounters and strand specific targeting of CRISPR-Cas systems,

which is especially relevant for RNA targeting CRISPR. The spacers of Type III systems, which target

RNA, have a bias towards targeting coding strands, making them capable of base pairing and thereby

targeting RNA. Unexpectedly we also found several subtypes with a preference for the template strand

(I-E and Type II). The reason for this type of strand bias is not yet clear, but we pose that this could be

caused by a selection for spacers that do not target RNA (RNA avoidance), as DNA-targeting with these

spacers might be impacted by inactivating complementary RNA (Jore et al., 2011). In addition, there

might be a difference in binding or dislodging of crRNA effector complexes from the template strand

vs coding strand by RNA polymerase (Clarke et al., 2018; Vink et al., 2020).

We have categorized multi-effector compatible CRISPR arrays whether they share the same repeats

and/or acquisition machinery and whether only DNA, or both DNA and RNA are targeted. DNA-

targeting systems that use multi-effector compatible arrays generally have their own acquisition

machinery and the low frequency of this co-occurrence in nature might indicate that this is not actively

selected for. It needs to be experimentally verified whether the spacers in these compatible arrays are

actually shared between complexes. However, some crRNA sharing between DNA systems has already

been observed experimentally, so it's therefore likely to be found for more systems (Majumdar et al.,

2015).

Multi-effector compatible arrays are much more common in co-occurring DNA- and RNA-targeting

systems and the strand bias that occurs in Type I arrays indicates that Type III effector complexes are

using these spacers and thereby creating selection pressure on the RNA binding potential of the

transcribed crRNA. It also seems that the most commonly co-occurring Type I systems (I-A, I-B and I-

C) that use compatible arrays, also have the largest coding strand bias. Whether this strand bias is

induced by the presence of Type III or whether these subtypes by their nature have a strand preference

and therefore became more commonly compatible with Type III systems is not yet clear. Interestingly,

many of the subtype combinations that share PAMs also co-occur more often than expected by chance,

suggesting they have positive epistatic interactions (Bernheim et al., 2020). Furthermore, repeat

sequences of type I-A and I-B are in same repeat families with Type III repeats, providing further

indications of their compatibility (Lange et al., 2013).

The experimentally determined spacer sharing in *Marinomonas mediterranea* (Silas et al., 2017)

described previously does not fall within the categories in this study as the Type III system has its own

adaptation machinery. In this case, the systems are not mutually compatible because the Type I systems

cannot use the Type III spacers due to a lack of PAM, which we have not further investigated in this

study. Also the other previously experimentally described spacer sharing systems in *Pyrococcus*

(Majumdar et al., 2015) and *Flavobacterium* (Hoikkala et al., 2021) were not found due to a lack of

sufficient hits, which demonstrates that these bio-informatic analysis likely underestimate the number

of systems that can cooperate.

The discovery of multi-effector spacer compatibility in a large number of genomes in this study together

with previous experimental evidence of spacer sharing of RNA and DNA-targeting systems (Deng et

al., 2013; Majumdar et al., 2015; Silas et al., 2017) shows that there is selection pressure to share spacers

cooperatively within arrays. The evolutionary benefits of such cooperativity could be profound. Firstly,

as two subtypes generally have different mismatch tolerance (Anderson et al., 2015; Fineran et al., 2014;

Manica et al., 2013), targeting the same sequence with two subtypes can reduce the probability of escape

mutation. Secondly a combination of an RNA and DNA targeting systems can provide multiple layers

of defence, where RNA-targeting might give more time for DNA-targeting systems to destroy the

invader before the cell is taken over (Vink et al., 2020). Thirdly the length of arrays in a genome has

recently been shown to be limited by auto-immunity (H. Chen et al., 2021). By sharing spacers, each

subtype is supplied with a maximum diversity of spacers while self-targeting costs are minimized. Lastly

the different mechanisms these systems use allows for complementary and distinct benefits. The priming

mechanism (Datsenko et al., 2012; Nicholson et al., 2019), unique to DNA targeting systems can

accelerate spacer acquisition for both systems, whereas cOA signaling pathways (Kazlauskiene et al.,

2017; Niewoehner et al., 2017), unique to Type III, could activate defence systems that benefit both

systems.

Altogether this study highlights the wealth of information that can be retrieved by analysing the targets

of CRISPR spacers on a large scale. It furthermore demonstrates under what conditions CRISPR-Cas

systems can cooperate and provides a large catalog of PAM predictions and targeted MGEs awaiting

further study.

Materials and Methods

CRISPR spacers and sequence data

221 089 spacers along with information on *cas* gene presence, genome and repeat sequence were

obtained from CRISPRCasDb (Pourcel et al., 2020) in February 2020 and the taxonomy of the genomes

was obtained from NCBI Taxonomy database (Federhen, 2012). We created our own sequence database

by combining all sequences from the NCBI nucleotide database (Benson et al., 2018; Pruitt et al., 2005),

environmental nucleotide database (Sayers et al., 2009), PHASTER (Arndt et al., 2016), Mgnify

(Mitchell et al., 2020) , IMG/M (I. M. A. Chen et al., 2017), IMG/Vr (Paez-Espino et al., 2019),

HuVirDb (Soto-Perez et al., 2019), HMP database (Peterson et al., 2009), and data from Pasolli et al.,

2019. All databases were accessed in February 2020.

Subtypes were predicted based on the repeat sequences using the subtype predictions and method

described by Bernheim et al., 2020, where the subtype of a spacer was inferred by the similarity of its

repeat sequence to repeat sequences with known subtype (74% identity threshold to infer subtype).

Blast hits and filtering

Hits between spacers and sequences from the aforementioned databases were obtained using the

command line blastn program (Altschul et al., 1990) version 2.10.0, which was run with parameters

word_size 10, gapopen 10, penalty 1 and an e-value cutoff of 1, to find as many potential targets as

possible. These blast hits were then filtered to remove hits of spacers inside CRISPR arrays and false

positive hits found by chance. Hits inside CRISPR arrays were detected by aligning the repeat sequence

of the spacer to the flanking regions of the spacer hit (23 nucleotides on both sides). This alignment was

done using the globalxs function from the Biopython pairwise2 package (Cock et al., 2009) with -3 gap

open and -3 gap extend parameters. If more than 13 nucleotides were identical in the alignment of at

least one flank, the hit was suspected to fall inside a CRISPR array and was filtered out.

To minimize the number of hits found by chance, we filtered hits based on the fraction of spacer

nucleotides that hit the target sequence, as this metric considers both the sequence identity and the

coverage of the spacer by the blast hit. In a first step, only hits with this fraction higher than 90% were

kept. To find targets for even more spacers while keeping the number of false positives low, we included

a second step where hits with a fraction higher than 80% were kept if another spacer from the same

genus hit the same contig or genome in the first step. This second step did not introduce hits on any new

contigs or genomes and was based on the assumption that multiple spacers from the same genus hitting

the same contig or genome is unlikely to be caused by chance. Finally, we removed spacers that were

shorter than 27 nucleotides (54 spacers) and removed 7 spacers that were hitting aspecifically, such as

inside ribosomal RNAs or tRNAs. This left 72,099 unique spacers with target hits for downstream

analysis.

Protospacer flank alignment for orientation and PAM predictions

The PAM is known to occur on the 5' end of the protospacer for Type I, Type IV and V CRISPR-Cas

systems, and on the 3' end for Type II systems (Collias & Beisel, 2021; Jackson et al., 2017). We used

this property to predict the orientation of transcription of CRISPR arrays and sequence of crRNA. The

PAM sides were compared to the nucleotide conservation in the flanking regions of the spacer hits and

the spacer orientations were predicted such that the flank with the greater conservation matched the

known PAM side.

To measure the nucleotide conservation in the flanking regions, data from multiple spacers was

combined based on the subtype and repeat sequences of the spacers. Highly similar repeat sequences

from the same subtype were clustered using CD-HIT (Fu et al., 2012) with a 90% identity threshold.

We hypothesized that similar repeat sequences would be used in a similar orientation and would utilize

the same PAM sequences, as coevolution of PAM, repeat and Cas1 and Cas2 sequences has been shown

previously (Alkhnbashi et al., 2014; Lange et al., 2013). For each repeat cluster the flanking regions of

the spacer hits were aligned. To equally weigh each spacer within the repeat cluster, irrespective of the

number of blast hits, consensus flanks were obtained per spacer. These consensus flanks contained the

most frequent nucleotide per position of the flanking regions. From the alignment of consensus flanks

the nucleotide conservation, or information content, in each flank was calculated in bitscore (Schneider

& Stephens, 1990) using the Sequence logo python package. We corrected for GC-content of the

targeted sequences by calculating the expected occurrences of each nucleotide based on the GC-content

of the spacer sequences. To minimize the number of orientation predictions based on little or noisy data,

we only predicted the orientation for repeat clusters when the alignment of consensus flanks consisted

of at least 10 unique protospacers. Furthermore, the information content of at least two positions was

higher than 0.3 bitscore and higher than 5 times the median bitscore calculated from 23-nt flanks on

both sides. These parameters were chosen as strictly as possible, while still yielding orientation

predictions for the highest number of spacers.

Using the orientation predictions described above, we predicted the PAMs for each repeat cluster by

checking which nucleotide positions were conserved. To minimize PAM predictions based on noise, we

only predicted the PAM for repeat clusters where the alignment of consensus flanks consisted of at least

10 unique protospacers. A nucleotide position was predicted to be part of the PAM when higher than

0.5 bitscore and higher than 10 times the median bitscore. These parameters were chosen as strictly as

possible, while maximizing the number of repeat clusters with PAM predictions and minimizing the

number of unique PAMs predicted.

We subsequently categorized and counted multi-effector compatible spacers in the following ways.

Firstly by an occurrence of multiple repeat clusters with different subtype classification that both

contained the same PAM, for two DNA targeting clusters (category I) or a DNA and a RNA targeting

cluster (category II). Secondly if multiple *cas* gene clusters from different subtypes were in the vicinity

of a single repeat cluster and their genomes did not further contain other arrays linked to these *cas* gene

clusters they were counted as a third category multi-effector compatible array.

Coding versus template strand targeting analysis

For each spacer target inside an open reading frame (ORF), we determined if the spacer targets the

coding (DNA and RNA) or template strand (DNA-only) during transcription. The ORFs and its

orientation were predicted using Prodigal (Hyatt et al., 2010) for one target sequence per spacer. The

target sequence of each spacer was selected as the longest hit sequence in the NCBI nucleotide database,

excluding 'other sequences', or, if no such sequence was hit, the longest hit sequence in metagenomics

database. Using our spacer orientation predictions for Type I, II and IV spacers, and the orientation

predictions from CRISPRCasDb for the other spacers, we checked if the spacer target (blast hit

orientation) was on the coding or template strand of the predicted ORF. To test for significant bias

towards either the temperate or the coding strand, a two-sided tailed binomial test was performed with

an expected probability of 0.5.

Data and Material availability

533   The datasets on which the analysis is based have been submitted as Supplementary Files. Scripts to

534   reproduce figures are available on request.

Acknowledgements

537   The authors thank Christine Pourcel and Pierre-Albert Charbit for supplying the CRISPRCasDB in a

538   spacer-based format and all members of the Brouns groups for input during group discussions. S.B. is

539   supported by a Vici grant of the Netherlands Organisation for Scientific Research (VI.C.182.027;

540   NWO).

Author contributions

543   S.B. and J.V. conceived and supervised the project; J.V. gathered databases; J.V. and J.B. wrote

544   analysis scripts; J.V., J.B. and S.B. wrote the manuscript.

Declaration of Interests

The authors declare no competing financial interests.

Literature cited

Alkhnbashi, O. S., Costa, F., Shah, S. A., Garrett, R. A., Saunders, S. J., & Backofen, R. (2014). CRISPRstrand: Predicting repeat orientations to determine the crRNA-encoding strand at CRISPR loci. *Bioinformatics*. https://doi.org/10.1093/bioinformatics/btu459

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*. https://doi.org/10.1016/S0022-2836(05)80360-2

Anders, C., Niewoehner, O., Duerst, A., & Jinek, M. (2014). Structural basis of PAM-dependent target DNA recognition by the Cas9 endonuclease. *Nature*, *513*(7519), 569–573. https://doi.org/10.1038/nature13579

Anderson, E. M., Haupt, A., Schiel, J. A., Chou, E., Machado, H. B., Strezoska, Ž., Lenger, S., McClelland, S., Birmingham, A., Vermeulen, A., & Smith, A. V. B. (2015). Systematic analysis of CRISPR-Cas9 mismatch tolerance reveals low levels of off-target activity. *Journal of Biotechnology*. https://doi.org/10.1016/j.jbiotec.2015.06.427

Arndt, D., Grant, J. R., Marcu, A., Sajed, T., Pon, A., Liang, Y., & Wishart, D. S. (2016). PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Research*. https://doi.org/10.1093/nar/gkw387

Artamonova, D., Karneyeva, K., Medvedeva, S., Klimuk, E., Kolesnik, M., Yasinskaya, A., Samolygo, A., & Severinov, K. (2020). Spacer acquisition by Type III CRISPR–Cas system during bacteriophage infection of Thermus thermophilus. *Nucleic Acids Research*. https://doi.org/10.1093/nar/gkaa685

Benson, D. A., Cavanaugh, M., Clark, K., Karsch-Mizrachi, I., Ostell, J., Pruitt, K. D., & Sayers, E. W. (2018). GenBank. *Nucleic Acids Research*. https://doi.org/10.1093/nar/gkx1094

Bernheim, A., Bikard, D., Touchon, M., & Rocha, E. P. C. (2020). Atypical organizations and epistatic interactions of CRISPRs and cas clusters in genomes and their mobile genetic elements. *Nucleic Acids Research*.

https://doi.org/10.1093/nar/gkz1091

Bolotin, A., Quinquis, B., Sorokin, A., & Dusko Ehrlich, S. (2005). Clustered regularly interspaced short

palindrome repeats (CRISPRs) have spacers of extrachromosomal origin. *Microbiology*.

https://doi.org/10.1099/mic.0.28048-0

Brouns, S. J. J., Jore, M. M., Lundgren, M., Westra, E. R., Slijkhuis, R. J. H., Snijders, A. P. L., Dickman, M. J.,

Makarova, K. S., Koonin, E. V, & van der Oost, J. (2008). Small CRISPR RNAs guide antiviral defense in

prokaryotes. *Science (New York, N.Y.)*, *321*(5891), 960–964. https://doi.org/10.1126/science.1159689

Callanan, J., Stockdale, S. R., Shkoporov, A., Draper, L. A., Ross, R. P., & Hill, C. (2020). Expansion of known

ssRNA phage genomes: From tens to over a thousand. *Science Advances*.

https://doi.org/10.1126/sciadv.aay5981

Chen, H., Mayer, A., & Balasubramanian, V. (2021). A scaling law in CRISPR repertoire sizes arises from

avoidance of autoimmunity. *BioRxiv*, 2021.01.04.425308. https://doi.org/10.1101/2021.01.04.425308

Chen, I. M. A., Markowitz, V. M., Chu, K., Palaniappan, K., Szeto, E., Pillay, M., Ratner, A., Huang, J., Andersen,

E., Huntemann, M., Varghese, N., Hadjithomas, M., Tennessen, K., Nielsen, T., Ivanova, N. N., & Kyrpides,

N. C. (2017). IMG/M: Integrated genome and metagenome comparative data analysis system. *Nucleic*

Acids Research. https://doi.org/10.1093/nar/gkw929

Clarke, R., Heler, R., MacDougall, M. S., Yeo, N. C., Chavez, A., Regan, M., Hanakahi, L., Church, G. M.,

Marraffini, L. A., & Merrill, B. J. (2018). Enhanced Bacterial Immunity and Mammalian Genome Editing via

RNA-Polymerase-Mediated Dislodging of Cas9 from Double-Strand DNA Breaks. *Molecular Cell*.

https://doi.org/10.1016/j.molcel.2018.06.005

Cock, P. J. A., Antao, T., Chang, J. T., Chapman, B. A., Cox, C. J., Dalke, A., Friedberg, I., Hamelryck, T., Kauff, F.,

Wilczynski, B., & De Hoon, M. J. L. (2009). Biopython: Freely available Python tools for computational

molecular biology and bioinformatics. *Bioinformatics*. https://doi.org/10.1093/bioinformatics/btp163

Collias, D., & Beisel, C. L. (2021). CRISPR technologies and the search for the PAM-free nuclease. *Nature*

Communications, *12*(1), 555. https://doi.org/10.1038/s41467-020-20633-y

Crawley, A. B., Henriksen, E. D., Stout, E., Brandt, K., & Barrangou, R. (2018). Characterizing the activity of

597  abundant, diverse and active CRISPR-Cas systems in lactobacilli. *Scientific Reports*.

598  https://doi.org/10.1038/s41598-018-29746-3

599 Datsenko, K. A., Pougach, K., Tikhonov, A., Wanner, B. L., Severinov, K., & Semenova, E. (2012). Molecular

600  memory of prior infections activates the CRISPR/Cas adaptive bacterial immunity system. *Nature*

601  *Communications*. https://doi.org/10.1038/ncomms1937

602 Deng, L., Garrett, R. A., Shah, S. A., Peng, X., & She, Q. (2013). A novel interference mechanism by a type IIIB

603  CRISPR-Cmr module in Sulfolobus. *Molecular Microbiology*. https://doi.org/10.1111/mmi.12152

604 Deveau, H., Barrangou, R., Garneau, J. E., Labonte, J., Fremaux, C., Boyaval, P., Romero, D. A., Horvath, P., &

605  Moineau, S. (2008). Phage Response to CRISPR-Encoded Resistance in Streptococcus thermophilus.

606  *Journal of Bacteriology*, *190*(4), 1390–1400. https://doi.org/10.1128/JB.01412-07

607 Elmore, J. R., Sheppard, N. F., Ramia, N., Deighan, T., Li, H., Terns, R. M., & Terns, M. P. (2016). Bipartite

608  recognition of target RNAs activates DNA cleavage by the Type III-B CRISPR–Cas system. *Genes and*

609  *Development*. https://doi.org/10.1101/gad.272153.115

610 Federhen, S. (2012). The NCBI Taxonomy database. *Nucleic Acids Research*.

611  https://doi.org/10.1093/nar/gkr1178

612 Fineran, P. C., Gerritzen, M. J. H., Suarez-Diez, M., Kunne, T., Boekhorst, J., van Hijum, S. a. F. T., Staals, R. H. J.,

613  & Brouns, S. J. J. (2014). Degenerate target sites mediate rapid primed CRISPR adaptation. *Proceedings of*

614  *the National Academy of Sciences*, *111*(16), 1629–1638. https://doi.org/10.1073/pnas.1400071111

615 Fischer, S., Maier, L. K., Stoll, B., Brendel, J., Fischer, E., Pfeiffer, F., Dyall-Smith, M., & Marchfelder, A. (2012).

616  An archaeal immune system can detect multiple protospacer adjacent motifs (PAMs) to target invader

617  DNA. *Journal of Biological Chemistry*. https://doi.org/10.1074/jbc.M112.377002

618 Fu, L., Niu, B., Zhu, Z., Wu, S., & Li, W. (2012). CD-HIT: Accelerated for clustering the next-generation

619  sequencing data. *Bioinformatics*. https://doi.org/10.1093/bioinformatics/bts565

620 Gasiunas, G., Young, J. K., Karvelis, T., Kazlauskas, D., Urbaitis, T., Jasnauskaite, M., Grusyte, M. M., Paulraj, S.,

621  Wang, P. H., Hou, Z., Dooley, S. K., Cigan, M., Alarcon, C., Chilcoat, N. D., Bigelyte, G., Curcuru, J. L.,

622  Mabuchi, M., Sun, Z., Fuchs, R. T., … Siksnys, V. (2020). A catalogue of biochemically diverse CRISPR-Cas9

orthologs. *Nature Communications*. https://doi.org/10.1038/s41467-020-19344-1

Gleditzsch, D., Pausch, P., Müller-Esparza, H., Özcan, A., Guo, X., Bange, G., & Randau, L. (2019). PAM

identification by CRISPR-Cas effector complexes: diversified mechanisms and structures. *RNA Biology*,

16(4), 504–517. https://doi.org/10.1080/15476286.2018.1504546

Hale, C. R., Duff, M. O., Graveley, B. R., Hale, C. R., Zhao, P., Olson, S., Duff, M. O., Graveley, B. R., Wells, L., &

Terns, R. M. (2009). RNA-guided RNA cleavage by a CRISPR RNA- Cas protein complex RNA-Guided RNA

Cleavage by a CRISPR RNA-Cas Protein Complex. *Cell*, *139*(5), 945–956.

https://doi.org/10.1016/j.cell.2009.07.040

Hoikkala, V., Ravantti, J., Díez-Villaseñor, C., Tiirola, M., Conrad, R. A., McBride, M. J., Moineau, S., & Sundberg,

L.-R. (2021). Cooperation between Different CRISPR-Cas Types Enables Adaptation in an RNA-Targeting

System. *MBio*, *12*(2), e03338-20. https://doi.org/10.1128/mBio.03338-20

Hyatt, D., Chen, G. L., LoCascio, P. F., Land, M. L., Larimer, F. W., & Hauser, L. J. (2010). Prodigal: Prokaryotic

gene recognition and translation initiation site identification. *BMC Bioinformatics*.

https://doi.org/10.1186/1471-2105-11-119

Jackson, S. A., McKenzie, R. E., Fagerlund, R. D., Kieper, S. N., Fineran, P. C., & Brouns, S. J. J. (2017). CRISPR-Cas:

Adapting to change. *Science*. https://doi.org/10.1126/science.aal5056

Johnson, K., Learn, B. A., Estrella, M. A., & Bailey, S. (2019). Target sequence requirements of a type III-B

CRISPR-Cas immune system. *Journal of Biological Chemistry*, *294*(26), 10290–10299.

https://doi.org/10.1074/jbc.RA119.008728

Jore, M. M., Lundgren, M., Van Duijn, E., Bultema, J. B., Westra, E. R., Waghmare, S. P., Wiedenheft, B., Pul, Ü.,

Wurm, R., Wagner, R., Beijer, M. R., Barendregt, A., Zhou, K., Snijders, A. P. L., Dickman, M. J., Doudna, J.

A., Boekema, E. J., Heck, A. J. R., Van Der Oost, J., & Brouns, S. J. J. (2011). Structural basis for CRISPR

RNA-guided DNA recognition by Cascade. *Nature Structural and Molecular Biology*, *18*(5), 529–536.

https://doi.org/10.1038/nsmb.2019

Kazlauskiene, M., Kostiuk, G., Venclovas, Č., Tamulaitis, G., & Siksnys, V. (2017). A cyclic oligonucleotide

signaling pathway in type III CRISPR-Cas systems. *Science*, *357*(6351), 605–609.

649  https://doi.org/10.1126/science.aao0100

650 Krupovic, M., Makarova, K. S., Forterre, P., Prangishvili, D., & Koonin, E. V. (2014). Casposons: A new

651  superfamily of self-synthesizing DNA transposons at the origin of prokaryotic CRISPR-Cas immunity. *BMC*

652  *Biology*. https://doi.org/10.1186/1741-7007-12-36

653 Lange, S. J., Alkhnbashi, O. S., Rose, D., Will, S., & Backofen, R. (2013). CRISPRmap: An automated classification

654  of repeat conservation in prokaryotic adaptive immune systems. *Nucleic Acids Research*.

655  https://doi.org/10.1093/nar/gkt606

656 Leenay, R. T., & Beisel, C. L. (2017). Deciphering, Communicating, and Engineering the CRISPR PAM. In *Journal*

657  *of Molecular Biology*. https://doi.org/10.1016/j.jmb.2016.11.024

658 Leenay, R. T., Maksimchuk, K. R., Slotkowski, R. A., Agrawal, R. N., Gomaa, A. A., Briner, A. E., Barrangou, R., &

659  Beisel, C. L. (2016). Identifying and Visualizing Functional PAM Diversity across CRISPR-Cas Systems.

660  *Molecular Cell*, *62*(1), 137–147. https://doi.org/10.1016/j.molcel.2016.02.031

661 Majumdar, S., Zhao, P., Pfister, N. T., Compton, M., Olson, S., Glover, C. V. C., Wells, L., Graveley, B. R., Terns, R.

662  M., & Terns, M. P. (2015). Three CRISPR-Cas immune effector complexes coexist in Pyrococcus furiosus.

663  *RNA (New York, N.Y.)*. https://doi.org/10.1261/rna.049130.114

664 Makarova, K. S., Haft, D. H., Barrangou, R., Brouns, S. J. J., Mojica, F. J. M., Wolf, Y. I., Yakunin, A. F., Oost, J. Van

665  Der, & Koonin, E. V. (2011). Evolution and classification of the CRISPR–Cas systems. *Nature Publishing*

666  *Group*, *9*(6), 467–477. https://doi.org/10.1038/nrmicro2577

667 Makarova, K. S., Wolf, Y. I., Alkhnbashi, O. S., Costa, F., Shah, S. A., Saunders, S. J., Barrangou, R., Brouns, S. J. J.,

668  Charpentier, E., Haft, D. H., Horvath, P., Moineau, S., Mojica, F. J. M., Terns, R. M., Terns, M. P., White, M.

669  F., Yakunin, A. F., Garrett, R. A., van der Oost, J., … Koonin, E. V. (2015). An updated evolutionary

670  classification of CRISPR–Cas systems. *Nature Reviews Microbiology*, *13*(11), 722–736.

671  https://doi.org/10.1038/nrmicro3569

672 Makarova, K. S., Wolf, Y. I., Iranzo, J., Shmakov, S. A., Alkhnbashi, O. S., Brouns, S. J. J., Charpentier, E., Cheng,

673  D., Haft, D. H., Horvath, P., Moineau, S., Mojica, F. J. M., Scott, D., Shah, S. A., Siksnys, V., Terns, M. P.,

674  Venclovas, Č., White, M. F., Yakunin, A. F., … Koonin, E. V. (2020). Evolutionary classification of CRISPR–

Cas systems: a burst of class 2 and derived variants. In *Nature Reviews Microbiology*.

https://doi.org/10.1038/s41579-019-0299-x

Malone, L. M., Warring, S. L., Jackson, S. A., Warnecke, C., Gardner, P. P., Gumy, L. F., & Fineran, P. C. (2020). A

jumbo phage that forms a nucleus-like structure evades CRISPR–Cas DNA targeting but is vulnerable to

type III RNA-based immunity. In *Nature Microbiology*. https://doi.org/10.1038/s41564-019-0612-5

Manica, A., Zebec, Z., Steinkellner, J., & Schleper, C. (2013). Unexpectedly broad target recognition of the

CRISPR-mediated virus defence system in the archaeon sulfolobus solfataricus. *Nucleic Acids Research*.

https://doi.org/10.1093/nar/gkt767

Marraffini, L. A. (2015). CRISPR-Cas immunity in prokaryotes. *Nature*, *526*(7571), 55–61.

https://doi.org/10.1038/nature15386

Marraffini, L. A., & Sontheimer, E. J. (2010). Self versus non-self discrimination during CRISPR RNA-directed

immunity. *Nature*, *463*(7280), 568–571. https://doi.org/10.1038/nature08703

Mendoza, B. J., & Trinh, C. T. (2018). In Silico Processing of the Complete CRISPR-Cas Spacer Space for

Identification of PAM Sequences. *Biotechnology Journal*. https://doi.org/10.1002/biot.201700595

Mendoza, S. D., Nieweglowska, E. S., Govindarajan, S., Leon, L. M., Berry, J. D., Tiwari, A., Chaikeeratisak, V.,

Pogliano, J., Agard, D. A., & Bondy-Denomy, J. (2020). A bacteriophage nucleus-like compartment shields

DNA from CRISPR nucleases. *Nature*, *577*(7789), 244–248. https://doi.org/10.1038/s41586-019-1786-y

Mitchell, A. L., Almeida, A., Beracochea, M., Boland, M., Burgin, J., Cochrane, G., Crusoe, M. R., Kale, V., Potter,

S. C., Richardson, L. J., Sakharova, E., Scheremetjew, M., Korobeynikov, A., Shlemov, A., Kunyavskaya, O.,

Lapidus, A., & Finn, R. D. (2020). MGnify: the microbiome analysis resource in 2020. *Nucleic Acids

Research*, *48*(D1), D570–D578. https://doi.org/10.1093/nar/gkz1035

Mojica, F. J. M., Díez-Villaseñor, C., García-Martínez, J., & Almendros, C. (2009). Short motif sequences

determine the targets of the prokaryotic CRISPR defence system. *Microbiology*, *155*(3), 733–740.

https://doi.org/10.1099/mic.0.023960-0

Musharova, O., Sitnik, V., Vlot, M., Savitskaya, E., Datsenko, K. A., Krivoy, A., Fedorov, I., Semenova, E., Brouns,

S. J. J., & Severinov, K. (2019). Systematic analysis of Type I-E Escherichia coli CRISPR-Cas PAM sequences

ability to promote interference and primed adaptation. *Molecular Microbiology*.

https://doi.org/10.1111/mmi.14237

Nicholson, T. J., Jackson, S. A., Croft, B. I., Staals, R. H. J., Fineran, P. C., & Brown, C. M. (2019). Bioinformatic

evidence of widespread priming in type I and II CRISPR-Cas systems. *RNA Biology*, *16*(4), 566–576.

https://doi.org/10.1080/15476286.2018.1509662

Niewoehner, O., Garcia-Doval, C., Rostøl, J. T., Berk, C., Schwede, F., Bigler, L., Hall, J., Marraffini, L. A., & Jinek,

M. (2017). Type III CRISPR-Cas systems produce cyclic oligoadenylate second messengers. *Nature*,

548(7669), 543–548. https://doi.org/10.1038/nature23467

Nobrega, F., Walinga, H., Dutilh, B., & Brouns, S. (2020). Prophages are associated with extensive, tolerated

CRISPR-Cas auto-immunity. *BioRxiv*, 2020.03.02.973784. https://doi.org/10.1101/2020.03.02.973784

Paez-Espino, D., Roux, S., Chen, I. M. A., Palaniappan, K., Ratner, A., Chu, K., Huntemann, M., Reddy, T. B. K.,

Pons, J. C., Llabrés, M., Eloe-Fadrosh, E. A., Ivanova, N. N., & Kyrpides, N. C. (2019). IMG/VR v.2.0: An

integrated data management and analysis system for cultivated and environmental viral genomes.

Nucleic Acids Research. https://doi.org/10.1093/nar/gky1127

Pasolli, E., Asnicar, F., Manara, S., Zolfo, M., Karcher, N., Armanini, F., Beghini, F., Manghi, P., Tett, A., Ghensi,

P., Collado, M. C., Rice, B. L., DuLong, C., Morgan, X. C., Golden, C. D., Quince, C., Huttenhower, C., &

Segata, N. (2019). Extensive Unexplored Human Microbiome Diversity Revealed by Over 150,000

Genomes from Metagenomes Spanning Age, Geography, and Lifestyle. *Cell*, *176*(3), 649-662.e20.

https://doi.org/10.1016/j.cell.2019.01.001

Pawluk, A., Davidson, A. R., & Maxwell, K. L. (2017). *Anti-CRISPR: discovery, mechanism and function*.

https://doi.org/10.1038/nrmicro.2017.120

Peterson, J., Garges, S., Giovanni, M., McInnes, P., Wang, L., Schloss, J. A., Bonazzi, V., McEwen, J. E.,

Wetterstrand, K. A., Deal, C., Baker, C. C., Di Francesco, V., Howcroft, T. K., Karp, R. W., Lunsford, R. D.,

Wellington, C. R., Belachew, T., Wright, M., Giblin, C., … Guyer, M. (2009). The NIH Human Microbiome

Project. *Genome Research*. https://doi.org/10.1101/gr.096651.109

Pinilla-Redondo, R., Mayo-Muñoz, D., Russel, J., Garrett, R. A., Randau, L., Sørensen, S. J., & Shah, S. A. (2020).

727 Type IV CRISPR-Cas systems are highly diverse and involved in competition between plasmids. *Nucleic*

728 *Acids Research*. https://doi.org/10.1093/nar/gkz1197

729 Pourcel, C., Touchon, M., Villeriot, N., Vernadet, J. P., Couvin, D., Toffano-Nioche, C., & Vergnaud, G. (2020).

730 CRISPRCasdb a successor of CRISPRdb containing CRISPR arrays and cas genes from complete genome

731 sequences, and tools to download and query lists of repeats and spacers. *Nucleic Acids Research*, *48*(D1),

732 D535–D544. https://doi.org/10.1093/nar/gkz915

733 Pruitt, K. D., Tatusova, T., & Maglott, D. R. (2005). NCBI Reference Sequence (RefSeq): A curated non-redundant

734 sequence database of genomes, transcripts and proteins. *Nucleic Acids Research*.

735 https://doi.org/10.1093/nar/gki025

736 Sayers, E. W., Barrett, T., Benson, D. A., Bryant, S. H., Canese, K., Chetvernin., V., Church, D. M., Dicuccio, M.,

737 Edgar, R., Federhen, S., Feolo, M., Geer, L. Y., Helmberg, W., Kapustin, Y., Landsman, D., Lipman, D. J.,

738 Madden, T. L., Maglott, D. R., Miller, V., … Ye, J. (2009). Database resources of the National Center for

739 Biotechnology Information. *Nucleic Acids Research*. https://doi.org/10.1093/nar/gkn741

740 Schneider, T. D., & Stephens, R. M. (1990). Sequence logos: A new way to display consensus sequences. *Nucleic*

741 *Acids Research*. https://doi.org/10.1093/nar/18.20.6097

742 Shah, S. A., Erdmann, S., Mojica, F. J. M., & Garrett, R. A. (2013). Protospacer recognition motifs: Mixed

743 identities and functional diversity. In *RNA Biology*. https://doi.org/10.4161/rna.23764

744 Shmakov, S. A., Sitnik, V., Makarova, K. S., Wolf, Y. I., Severinov, K. V., & Koonin, E. V. (2017). The CRISPR spacer

745 space is dominated by sequences from species-specific mobilomes. *MBio*.

746 https://doi.org/10.1128/mBio.01397-17

747 Silas, S., Lucas-Elio, P., Jackson, S. A., Aroca-Crevillén, A., Hansen, L. L., Fineran, P. C., Fire, A. Z., & Sánchez-

748 Amat, A. (2017). Type III CRISPR-Cas systems can provide redundancy to counteract viral escape from

749 type I systems. *ELife*. https://doi.org/10.7554/eLife.27601

750 Silas, S., Mohr, G., Sidote, D. J., Markham, L. M., Sanchez-Amat, A., Bhaya, D., Lambowitz, A. M., & Fire, A. Z.

751 (2016). Direct CRISPR spacer acquisition from RNA by a natural reverse transcriptase-Cas1 fusion protein.

752 *Science*, *351*(6276). https://doi.org/10.1126/science.aad4234

Soto-Perez, P., Bisanz, J. E., Berry, J. D., Lam, K. N., Bondy-Denomy, J., & Turnbaugh, P. J. (2019). CRISPR-Cas

System of a Prevalent Human Gut Bacterium Reveals Hyper-targeting against Phages in a Human Virome

Catalog. *Cell Host & Microbe*, *26*(3), 325-335.e5. https://doi.org/10.1016/j.chom.2019.08.008

Strutt, S. C., Torrez, R. M., Kaya, E., Negrete, O. A., & Doudna, J. A. (2018). RNA-dependent RNA targeting by

CRISPR-Cas9. *ELife*. https://doi.org/10.7554/eLife.32724

Swarts, D. C., Mosterd, C., van Passel, M. W. J., & Brouns, S. J. J. (2012). CRISPR Interference Directs Strand

Specific Spacer Acquisition. *PLOS ONE*, *7*(4), e35888. https://doi.org/10.1371/journal.pone.0035888

Vale, P. F., Lafforgue, G., Gatchitch, F., Gardan, R., Moineau, S., & Gandon, S. (2015). Costs of CRISPR-Cas-

mediated resistance in Streptococcus thermophilus. *Proceedings of the Royal Society B: Biological*

Sciences. https://doi.org/10.1098/rspb.2015.1270

Vink, J. N. A., Martens, K. J. A., Vlot, M., McKenzie, R. E., Almendros, C., Estrada Bonilla, B., Brocken, D. J. W.,

Hohlbein, J., & Brouns, S. J. J. (2020). Direct Visualization of Native CRISPR Target Search in Live Bacteria

Reveals Cascade DNA Surveillance Mechanism. *Molecular Cell*, *77*(1), 39-50.e10.

https://doi.org/10.1016/j.molcel.2019.10.021

Walton, R. T., Christie, K. A., Whittaker, M. N., & Kleinstiver, B. P. (2020). Unconstrained genome targeting with

near-PAMless engineered CRISPR-Cas9 variants. *Science*, *368*(6488), 290 LP – 296.

https://doi.org/10.1126/science.aba8853

Walton, R. T., Hsu, J. Y., Joung, J. K., & Kleinstiver, B. P. (2021). Scalable characterization of the PAM

requirements of CRISPR-Cas enzymes using HT-PAMDA. *Nature Protocols*.

https://doi.org/10.1038/s41596-020-00465-2

Westra, E. R., Buckling, A., & Fineran, P. C. (2014). CRISPR-Cas systems: Beyond adaptive immunity. *Nature*

Reviews Microbiology. https://doi.org/10.1038/nrmicro3241

Westra, E. R., Semenova, E., Datsenko, K. A., Jackson, R. N., Wiedenheft, B., Severinov, K., & Brouns, S. J. J.

(2013). Type I-E CRISPR-Cas Systems Discriminate Target from Non-Target DNA through Base Pairing-

Independent PAM Recognition. *PLoS Genetics*, *9*(9), e1003742.

https://doi.org/10.1371/journal.pgen.1003742

Wolf, Y. I., Silas, S., Wang, Y., Wu, S., Bocek, M., Kazlauskas, D., Krupovic, M., Fire, A., Dolja, V. V., & Koonin, E.

V. (2020). Doubling of the known set of RNA viruses by metagenomic analysis of an aquatic virome.

Nature Microbiology. https://doi.org/10.1038/s41564-020-0755-4

Xue, C., Zhu, Y., Zhang, X., Shin, Y. K., & Sashital, D. G. (2017). Real-Time Observation of Target Search by the

CRISPR Surveillance Complex Cascade. *Cell Reports*, *21*(13), 3717–3727.

https://doi.org/10.1016/j.celrep.2017.11.110

Figure 1. Spacer targets found with BLAST. (A) Computational pipeline for finding spacer targets. Targets of
72 099 spacers were found using blastn and filtered based on the fraction of spacer nucleotides matching a target
sequence (See methods). (B) Venn diagram of spacers with matches in the NCBI nucleotide database versus
metagenomic databases (C) Number of spacers per subtype. The subtype of a spacer was predicted based on
similarity of the repeat sequence to repeats with a known subtype (See methods). (D) Fraction of spacers with hits
per subtype. (E) Fraction of spacers with hits for the ten genera with the highest and ten genera with the lowest
fraction of hits. Only genera with at least 500 spacers are shown.

Figure 2. PAM determination of repeat clusters. (A) Sequence logos of upstream flank of hits to spacers from

Type I repeat clusters. Sequence logos of protospacer flanking regions per repeat cluster. Y-axes show information

content per nucleotide position. Label includes subtype of the repeat cluster and a representative genus in which

this repeat cluster is found. (B) Same as (A) but for downstream flanks of spacers from Type II repeat clusters. (C)

Same as (A) but for upstream flanks from Type III repeat clusters. (D) Frequency of PAM determined repeat

clusters with more than 25 hits. Nucleotide positions were considered part of PAM with a bitscore of at least 0.4

and 10 times above the median bitscore of the 23 nucleotides surrounding the hits. PAM size was at least 2

nucleotides. (E) Frequency of PAM determined repeat clusters for Type III systems that contain Cas1-2 vs Type

III systems that lack Cas1-2.

Figure 3. Relationship between repeat and PAM sequence. (A) Schematic of the analysis of PAM and repeat

sequence. The nucleotide identity of the PAM in each position is compared to the nucleotide of the repeat. (B).

PAM nucleotide frequency for Type I repeats. For each given repeat nucleotide position (indicated with coloured

boxes) the PAM nucleotide (pie chart) for each unique PAM-repeat combination of our database. Number of

occurrence is indicated above the pie chart (n). (C). The frequency of matches (red) and mismatches (grey) between

the PAM and the corresponding repeat nucleotide for each position in relationship to the spacer. For Type II, the

positions are compared on the other side of the spacer.

Figure 4. Template and coding strand targeting of spacers. (A) Schematic representation of a spacer targeting

the template strand and a spacer targeting the coding strand inside an ORF. Spacers targeting the coding strand are

also able to base pair with and target transcribed RNA. (B) Fraction of *Escherichia* spacers targeting template

(blue) and coding (orange) strand by subtype. (C) Fraction of *Moraxella* spacers targeting template and coding

strand by subtype. (D) Fraction of spacers targeting template and coding strand for Type I and Type IV subtypes.

(E) Fraction of spacers targeting template and coding strand for Type II and Type III subtypes. (F) Fraction of

spacers targeting template and coding strand for Type I. Spacers are grouped based on which other type of Cas

effector genes are present in the genome. (G) Same as (F) but for Type II spacers. Significance of strand bias is

calculated with a binomial test and a p-value<0.01 is indicated with an asterisk.

Figure 5. Different organisations of subtypes containing compatible spacer sequences. (A) Pie chart of

frequency of genomes each category of organisation, based on the subtype combination involved. Total number

of genomes for which this category was found (n) is noted in each chart (n).(B-D) Genome representations of

examples for the different organisation categories, (b) Type I-Type I compatibility, (c) Type I- Type III

compatibility (different repeat sequences), (d) Type I- Type III compatibility (same repeat sequences). Genes

involved in interference (blue) and adaptation (red) are shown for the different subtypes within the genome. PAM

logo and strand bias of each associated repeat cluster is depicted below the genomic representations.

 Supplementary figures

Supplementary figure 1. Taxonomy of spacer targets and number of found targets per database. (A) The

taxonomy of targeted sequences of the NCBI nucleotide database was obtained from the NCBI taxonomy database.

For hits in viral sequences, the taxonomy of known hosts was used to label the virus as a eukaryotic or prokaryotic

virus. (B) The contribution of each database to the total number of hits after filtering. All databases were accessed

in February 2020.

843

Supplementary figure 2 Average number and signal to noise ratio of clustered hits. Different clustering

methods were compared for their average number of unique hits (blue) and average signal to noise ratio (orange).

Signal to noise ratio was calculated by dividing the average information content of the two top positions in the

flank (potential PAM nucleotides) by the median information content in sequence logos generated from flanks of

hits. The clustering categories is based on whether spacers come from same species and subtype (species-subtype),

from same genus and subtype (genus-subtype), from clusters of repeat sequences with 90% identity (repeat-cluster)

or clusters of repeat sequences with 90% identity and additionally compensation for GC-content of spacers within

the cluster (see Materials and Methods, repeat-clusterGC).

a

Supplementary figure 3. Sequence logo Type III repeats. ClustalW alignment of Type III repeats for which

orientation was determined based on presence of PAM (n = 21 unique repeats). The 3'end of the repeat, which is

the 5' handle of the transcribed crRNA, has a conserved motif (ATTGAAAC).

Supplementary figure 4. Sequence logos of protospacer flanking regions per repeat cluster. (A) The sequence

logo of subtype IF repeat cluster is based on flanks of 1629 unique spacers. The reverse complement of the subtype

IF CC PAM is found, due to incorrect orientation of spacers from the repeat cluster. (B) The sequence logos of

subtype IIA and IIC repeat clusters. No positions with conserved nucleotides are visible, despite the high number

of unique spacers for each cluster (52, 155 respectively). (C) The sequence logo of a subtype IIIB repeat cluster

showing no positions with conserved nucleotides based on flanks of 31 unique spacers.

Relationship between repeat and PAM sequence of Type II and Type III systems. Same as Figure 3B except

for Type II (A) and Type III (B) systems.

Subtype	PAM	Genus	Subtype	PAM	Genus	Subtype	PAM	Genus
I-A	ATG	*Leptospira*	**I-B**	TTTA	*Petrimonas*	**I-E**	AAG	*Geobacter*
	CCN	*Acidianus*		TTG	*Thermobacillus*		AAN	*Klebsiella*
	TTA	*Thermodesulfobacterium*	**I-C**	CTN	*Anaerobutyricum*		AAT	*Lactobacillus*
	TCN	*Sulfurisphaera*		CCN	*Porphyromonas*		AAA	*Kosakonia*
	ATN	*Aminobacterium*		CTT	*Ruminococcus*		AC	*Corynebacterium*
I-B	CCA	*Moorella*		TTC	*Geobacillus*		AG	*Xenorhabdus*
	CCN	*Clostridium*		TTN	*Acidovorax*		AWG	*Escherichia*
	CCT	*Ureibacillus*		TTT	*Lachnoclostridium*	**I-F**	ACC	*Aeromonas*
	TAC	*Halorubrum*	**I-D**	GCN	*Haloquadratum*		CC	*Pseudomonas*
	TCA	*Campylobacter*		GGTG	*Halorubrum*		CCA	*Pseudomonas*
	TCN	*Campylobacter*		GTN	*Methanotrix*	**I-G**	TAC	*Rothia*
	TTA	*Methanosarcina*		GTT	*Microcystis*		TAN	*Propionibacterium*
	TTC	*Halobacterium*		GTG	*Methanospirillum*		TTN	*Pseudopropionibacterium*
	TTN	*Novibacillus*	**I-E**	AAC	*Bifidobacterium*		AAN	*Acidipropionibacterium*
							TTC	*Rhodothermus*

 Supplementary Table 1. Unique Type I PAM sequences. Table of all unique Type I PAMs found for the

 different subtypes and representative genera that contain the repeat cluster for which each PAM was determined.