The tiny, conserved zinc-finger protein GTSF1

helps PIWI proteins achieve their full catalytic potential

Amena Arif[1,2], Shannon Bailey[2], Natsuko Izumi[3], Todd A. Anzelon[4],

Deniz M. Ozata[2,6], Cecilia Andersson[2], Ildar Gainetdinov[2],

Ian J. MacRae[4], Yukihide Tomari[3,5] & Phillip D. Zamore[2,*]

[1]Department of Biochemistry and Molecular Pharmacology, University of Massachusetts
Chan Medical School, Worcester, MA 01605, USA

[2]Howard Hughes Medical Institute and RNA Therapeutics Institute, University of
Massachusetts Chan Medical School, 368 Plantation Street, Worcester, MA 01605,
USA

[3]Laboratory of RNA Function, Institute for Quantitative Biosciences, The University of
Tokyo, Tokyo, Japan

[4]Department of Integrative Structural and Computational Biology, The Scripps Research
Institute, La Jolla, CA 92037, USA

[5]Department of Computational Biology and Medical Sciences, Graduate School of
Frontier Sciences, The University of Tokyo, Kashiwa, Japan

[6]Current address: Department of Molecular Biosciences, Stockholm University,
SE-106 91 Stockholm, Sweden

*Correspondence: phillip.zamore@umassmed.edu (P.D.Z.)

Abstract

Argonaute proteins use nucleic acid guides to find and bind specific DNA or RNA target sequences. Argonaute proteins can be found in all kingdoms of life, and play diverse biological functions including genome defense, gene regulation, and chromosome partitioning. Many Argonautes retain their ancestral endoribonuclease activity, cleaving the phosphodiester bond between target nucleotides t10 and t11. In animals, a specialized class of Argonautes, the PIWI proteins, use 21–35 nt PIWI-interacting RNAs (piRNAs) to direct transposon silencing, protect the germline genome, and regulate gene expression during gametogenesis[1]. The piRNA pathway is required for fertility in one or both sexes of nearly all animals. Both piRNA production and function require RNA cleavage catalyzed by PIWI proteins. Spermatogenesis in mice and other placental mammals requires three distinct, developmentally regulated PIWI proteins: MIWI (PIWIL1), MILI (PIWIL2), and MIWI2 (PIWIL4)[2–4]. The piRNA-guided endoribonuclease activities of MIWI and MILI are essential to produce functional sperm[5,6]. piRNA-directed silencing in mice and insects also requires Gametocyte-Specific Factor 1 (GTSF1), a PIWI-associated protein of unknown function[7–12]. Here, we report that GTSF1 potentiates the weak, intrinsic, piRNA-directed RNA cleavage activities of PIWI proteins, transforming them into efficient endoribonucleases. GTSF1 represents the first example of an auxiliary protein that potentiates the catalytic activity of an Argonaute protein.

Main

In animals, 21–35 nt PIWI-interacting RNAs (piRNAs) direct PIWI proteins to silence

transposons and regulate gene expression[1,13–27]. Invertebrates produce piRNAs and

PIWI proteins in both the soma and the germline[28–35] but mammalian piRNAs act only in

the germline[36–43]. Mice lacking any of their three PIWI proteins—MIWI2 (PIWIL4), MILI

(PIWIL2), and MIWI (PIWIL1)—or other proteins required for piRNA production are

invariably male sterile[1–4]. As in other animals, mouse piRNA production requires

catalytically active PIWI proteins, MILI and MIWI[5,6]. Transposon silencing is the

ancestral function of piRNAs; uniquely, placental mammals also produce pachytene

piRNAs, which first appear shortly after the onset of meiosis I (refs. 40–42,44–46) and

reach a peak abundance in spermatocytes rivalling that of ribosomes[16]. Pachytene

piRNAs tune the abundance of mRNAs required for spermiogenesis, the process by

which round spermatids become sperm[47–50]. Pachytene piRNAs have been proposed to

direct MIWI and MILI to cleave specific mRNAs, ensuring appropriate levels of their

protein products[47,50–52]. Testing this idea has been thwarted by the absence of a cell-

free system in which MIWI or MILI can be loaded with synthetic piRNAs of defined

sequence.

Recombinant MIWI loaded with a piRNA of defined sequence

We used lentiviral transduction to engineer a stable HEK293T cell line over-expressing

epitope-tagged MIWI. Tagged MIWI was captured from cell lysate using anti-FLAG

antibody coupled to paramagnetic beads, incubated with a synthetic piRNA bearing a

monophosphorylated 5′ terminus and 2′-O-methylated 3′ end, and eluted from the

magnetic beads using 3XFLAG peptide. MIWI loaded with a synthetic piRNA (MIWI

piRISC; Fig. 1a), but not unloaded apo-MIWI (Extended Data Fig. 1a), cleaved a 5′ ^{32}P-

radiolabled target RNA fully complementary to the synthetic guide (Extended Data

Fig. 1b, c and Supplementary Data Fig. 1a).

Recombinant MIWI bound stably to an RNA guide bearing a 5′ monophosphate

but not to a guide with a 5′ hydroxyl group (Extended Data Fig. 1d). The MID domain of

Argonaute proteins contains a 5′ monophosphate-binding pocket that anchors the RNA

guide to the protein[53–59]. Mutations predicted to disrupt 5′ monophosphate-binding

perturb PIWI function[55,60–64]. We immobilized MIWI on paramagnetic beads via its

3XFLAG tag, incubated it with guide RNA, washed the beads, eluted the MIWI piRISC

with 3XFLAG tag peptide, and tested its ability to cleave a fully complementary target

RNA. Incubation with a 5′ monophosphorylated but not an otherwise identical 5′ hydroxy

guide yielded MIWI piRISC that cleaved target RNA (Extended Data Fig. 1d). In vivo,

the methyltransferase HENMT1 adds a 2′-O-methyl group to the 3′ ends of piRNAs.

Henmt1[−/−] male mice and *henn1*[−/−] worms are sterile[65–71]. Terminal 2′-O-methyl

modification likely stabilizes small RNAs against degradation by cellular ribonucleases

rather than secures the guide to the protein[72,73]. Consistent with this function for 2′-O-

methylation, piRNAs bearing a 3′ terminal 2′ hydroxyl or 2′-O-methyl were equally

functional in directing target cleavage, provided that the piRNA was 5′

monophosphorylated (Extended Data Fig. 1d).

Purified MIWI RISC is a sluggish enzyme

Although affinity-purified MIWI loaded with a piRNA bearing 5′ monophosphorylated and

3′, 2′-O-methylated termini specifically cleaved a fully complementary target RNA at the

phosphodiester bond that links target nucleotides t10 to t11, the rate of cleavage

$(0.01 \ \mathrm{min}^{-1})$ was >300 times slower than that catalyzed by mouse AGO2 RISC (≥

$3 \ \mathrm{min}^{-1})$[74,75]. At physiological temperature (37°C), mouse AGO2 RISC catalyzes

multiple rounds of target cleavage[76]. By contrast, 5 nM MIWI piRISC cleaved only ~15%

of the target RNA (5 nM) after 1 h (Fig. 1b, Extended Data Fig. 1e, and Supplementary

Data Fig. 1b). Inefficient target cleavage by MIWI piRISC was not caused by the

presence of the amino-terminal 3XFLAG-SNAP tandem tag: removing the tandem tags

using tobacco etch virus protease generated an untagged protein (Extended Data

Fig. 1f and Supplementary Data Fig. 1c) whose target cleavage kinetics were identical

to that of piRISC assembled with the tagged MIWI (Extended Data Fig. 1g).

The ubiquitously expressed arginine methyltransferase PRMT5 modifies the

amino terminus of PIWI proteins, allowing it to bind Tudor domain-containing proteins,

many of which are required for piRNA biogenesis, gametogenesis, and fertility[77–79]. A

potential explanation for the sluggish activity of recombinant MIWI piRISC is that it lacks

arginine methylation. We used mass spectrometry to map the positions of methyl

arginine in our affinity-purified MIWI. All of the arginine residues previously shown to be

methylated in endogenous MIWI immunoprecipitated from mouse testis[77] were

methylated in recombinant MIWI produced in HEK293T cells (Extended Data Fig. 1h).

Another possible explanation for the inefficiency of target cleavage by MIWI

piRISC is that the recombinant protein, although properly arginine methylated, lacks

other post-translational modifications. To test this idea, we immobilized apo-MIWI,

incubated it with wild-type (C57BL/6) mouse testis lysate in the presence of an ATP

regenerating system at 25°C for 15 min, removed the testis lysate by washing, and then

loaded MIWI with a synthetic piRNA and eluted the resulting piRISC. Pre-incubation of

MIWI with testis lysate either before or after loading with a piRNA failed to increase its

target cleaving activity (Extended Data Fig. 2a, b). We conclude that neither a missing

post-translational modification nor a tightly associated protein partner is likely to explain

the low activity of MIWI piRISC compared to AGO2 RISC.

MIWI piRISC requires an auxiliary factor to efficiently cleave its target

Adding testis lysate increased the rate of target cleavage by affinity-purified MIWI

piRISC ~20-fold (Fig. 1b; Extended Data Fig. 2b). This effect cannot be attributed to the

lysate contributing additional piRISC, because testis lysate alone failed to generate

detectable cleavage product (Fig. 1b).

MIWI is first produced as spermatogonia enter meiosis and differentiate into

spermatocytes[3,80]. To determine whether the MIWI-potentiating factor is differentially

expressed during spermatogenesis, we supplemented the target cleavage assay with

lysate prepared from FACS-purified germ cells: piRISC-potentiating activity was

greatest in lysate from secondary spermatocytes (Fig. 1c), a cell type in which

pachytene-piRNA-directed target cleavage is readily detected in vivo[47,50]. Moreover, the

potentiating activity was testis-specific: lysates from brain, liver, or kidney failed to

enhance piRNA-directed target cleavage by MIWI (Extended Data Fig. 3a). Finally, the

potentiating activity was specific for PIWI proteins and had no effect on the rate of target

cleavage by mouse AGO2 (Extended Data Fig. 3b).

Three lines of evidence suggest that the MIWI-potentiating activity contains one

or more structural Zn^{2+} ions. First, pre-treating testis lysate with the sulfhydryl alkylating

agent *N*-ethylmaleimide inactivated the potentiating activity (Extended Data Fig. 3c),

indicating reduced cysteine residues, which often bind divalent metal cations[81], are

essential. Second, the MIWI-potentiating activity was unaltered by EGTA, which

chelates Ca^{2+}, but was irreversibly inactivated by EDTA, which chelates many divalent

metals, and by 1,10-Phenanthroline, which specifically chelates Zn^{2+} (Extended Data

Fig. 3d). Adding additional metal ions failed to rescue the activity (Extended Data Fig.

3e–g), suggesting that loss of Zn^{2+} irreversibly denatures the MIWI-potentiating factor, a

characteristic of zinc-finger proteins[82,83]. Third, the MIWI-potentiating activity bound

more tightly to an immobilized metal-affinity resin charged with Zn^{2+} than to resin

charged with Ni^{2+} (Extended Data Fig. 3h).

To identify the MIWI-potentiating activity, we developed a chromatographic

purification scheme using cation-exchange, hydrophobic-interaction, and size-exclusion

chromatography (Fig. 2a). Notably, the activity eluted from a Superdex 200 size-

exclusion column as a single ~17 kDa peak (Fig. 2b). Taken together, our data suggest

that the MIWI-potentiating activity corresponds to a small, testis-specific, Zn^{2+}-binding

protein abundantly expressed in meiotic and post-meiotic male germ cells (Fig. 2c).

GTSF1 is necessary and sufficient for efficient target cleavage by MIWI

Gametocyte Specific Factor 1 (GTSF1) is a conserved, 19,083 Da, tandem CHHC-type

zinc-finger protein essential for piRNA function and fertility in flies[7,8,10], silk moths[11],

worms[84], and mice[9,12]. Like *Miwi* and *Mili*, the mRNA abundance of *Gtsf1*, as well as its

paralogs *Gtsf1L* and *Gtsf2*, rises sharply as male germ cells enter meiosis I, peaks in

secondary spermatocytes, and then declines in round spermatids (Extended Data

Fig. 4a). Immobilized mouse GTSF1 captures all three PIWI proteins from testis lysate,

but GTSF1 and PIWI proteins co-expressed in HEK293T cells do not co-

immunoprecipitate, leading to the suggestion that the GTSF1-PIWI interaction requires

additional protein or RNA components[9]. MIWI2-bound piRNAs are lost in $Gtsf1^{-/-}$ male

mice[9], which are sterile[12] likely because loss of GTSF1 causes loss of MIWI2-directed

retrotransposon promoter methylation[4,85–87]. Although RNA cleavage by MIWI2 is not

required for piRNA biogenesis or function, production of MIWI2 piRISC requires MILI-

dependent piRNA amplification, a process that requires the endonuclease activity of

MILI[6]. In flies, piRNA-directed transcriptional silencing of transposons by Piwi, but not

Piwi piRISC-assembly, requires the GTSF1 ortholog Asterix[7,8].

To test whether the MIWI-potentiating activity in the testis lysate corresponded to

GTSF1, we used CRISPR-Cas9 to engineer a mouse strain with a 3XFLAG-HA epitope

tag inserted into the endogenous *Gtsf1* coding sequence (*Gtsf1^{KI}*; Fig. 2d). Because the

MIWI potentiating activity was greatest in secondary spermatocytes, we prepared lysate

from FACS-purified secondary spermatocytes from homozygous *Gtsf1^{KI/KI}* male mice.

Incubation with anti-FLAG antibody coupled to paramagnetic beads depleted the lysate

of both epitope-tagged GTSF1 (Fig. 2e) and the MIWI-potentiating activity (Fig. 2f). The

activity was recovered from the beads by elution with 3XFLAG peptide (Fig. 2f). In

contrast, lysate from C57BL/6 secondary spermatocytes retained the MIWI-potentiating

factor after incubation with anti-FLAG antibody, and no activity was eluted from the

beads after incubation with FLAG peptide. Thus, GTSF1 is necessary to potentiate the

catalytic activity of MIWI.

GTSF1 is also sufficient to potentiate the catalytic activity of MIWI. Adding

purified recombinant GTSF1 (Extended Data Fig. 4b) to the MIWI-piRISC target

cleavage assay increased the rate of target cleavage by MIWI >100-fold (Fig. 3a, b).

The addition of 500 nM GTSF1 to MIWI piRISC (5 nM) caused the pre-steady-state rate

(k_{burst}) of target cleavage to increase from ~0.01 min^{-1} to 1.5 min^{-1} (Fig. 3b), a rate

similar to that of AGO2 (≥ 3 min^{-1})[74,75].

Our data suggest that GTSF1 does not promote product release or turnover: the

steady-state rate (k_{ss} <0.005 min^{-1}) of target cleavage under multiple-turnover

conditions (i.e., GTSF1 >> target RNA >> MIWI) was essentially unchanged from the

rate in the absence of GTSF1 (Fig. 1b and 3b). Perhaps MIWI remains bound to the

cleaved products, preventing it from catalyzing multiple rounds of target cleavage.

Supporting this idea, at incubation times >15 min, 3′-to-5′ exonucleases present in testis

lysate degrade the uncut, full-length target RNA, but the 5′ cleavage product remains

stable, consistent with its being protected by MIWI bound to its 3′ end (Fig. 1b). In vivo,

product release has been proposed to be facilitated in insects and mammals by the

RNA-stimulated ATPase Vasa (DDX4)[88–90] and the Vasa-like protein DDX43 (ref. 91).

RNA-binding is essential for GTSF1 function

Only three known eukaryotic proteins contain CHHC zinc-fingers: the spliceosomal

RNA-binding protein U11-48K, the TRM13 tRNA methyltransferase, and GTSF1 and its

paralogs[92]. In vitro, the first zinc-finger of GTSF1 directly binds RNA; RNA binding

requires four basic surface residues (R26, R29, K36 and K39)[93]. Potentiation of target

cleavage required RNA-binding by GTSF1: purified mutant GTSF1[R26A,R29A,K36A,K39A]

(Extended Data Fig. 4b) did not detectably increase the rate of catalysis by either MIWI

or MILI (Fig. 3c–d and Extended Data Table 2), suggesting that GTSF1 must interact

with RNA to function.

GTSF1 function requires PIWI-binding

In flies and mice, GTSF1 interacts with PIWI proteins through conserved aromatic

residues in its carboxy-terminus[7,9] (Fig. 2c). Mutating these residues in the recombinant

protein—W98A, W107A, and W112A for mouse GTSF1—reduced the stimulatory effect

of mouse GTSF1 on MIWI in a concentration-dependent manner, supporting the idea

that GTSF1 binds MIWI directly when potentiating target cleavage (Fig. 3e and

Extended Data Table 2) and suggesting that W98, W107, and W112 define a conserved

surface of GTSF1-that interacts with PIWI proteins. We measured the pre-steady-state

catalytic rate (k_{burst}) as a function of GTSF1 concentration. Fitting the data to a

hyperbolic function revealed that GTSF1 binds MIWI piRISC >60-fold more tightly than

GTSF1^{W98A,W107A,W112A} (GTSF1: K_D = 8 nM, 95% confidence interval [C.I.] = 6 – 9 nM;

GTSF1^{W98A,W107A,W112A}: K_D = 500 nM, 95% C.I. = 100 – 900 nM) (Fig. 3e). Moreover,

PIWI-binding appears to be the sole defect in GTSF1^{W98A,W107A,W112A}, since the first-

order rates of target cleavage by MIWI at saturating concentrations of either wild-type or

mutant GTSF1 were essentially indistinguishable. (wild-type: 1.2 min^{-1}, 95% C.I. = 0.7 –

2 min^{-1}; mutant GTSF1: 0.8 min^{-1}, 95% C.I. = 0.6 – 0.9 min^{-1}).

The PIWI-stimulatory function of GTSF1 orthologs is evolutionarily conserved

GTSF1 orthologs can be found in many metazoan genomes[92], suggesting that GTSF1

may potentiate target cleavage by PIWI proteins in many animals. Supporting this idea,

purified recombinant Gtsf1 from the arthropod *Bombyx mori* (Extended Data Fig. 4c)

potentiated the catalytic activity of the *B. mori* PIWI protein, BmSiwi (Fig. 3f).

The structure and kinetics of EfPiwi from the sponge *Ephydatia fluviatilis* were

recently described[64]. Like MIWI, EfPiwi possesses inherently weak catalytic activity, a

feature common to all PIWI proteins examined to date. Although the *E. fluviatilis*

genome is yet to be sequenced, the closely related, fully sequenced genome of

Ephydatia muelleri contains a readily identifiable GTSF1 ortholog. Purified recombinant

EmGtsf1 (Extended Data Fig. 4c) stimulated the single-turnover catalytic rate of EfPiwi

piRISC ~6 fold (Fig. 3g). Sponges (Porifera) are the sister group to all other animals, the

Eumetazoa, having separated ~900 Mya. Thus, the last common ancestor of all animals

likely required GTSF1 to potentiate target cleavage by PIWI proteins.

The GTSF1 tandem zinc-finger domains are conserved across phyla, whereas

the GTSF1 carboxy-terminal sequence diverges substantially between mammals and

arthropods (Extended Data Table 1). For example, the sequences of the mouse and

rhesus macaque first and second zinc-fingers are 100% identical, whereas their C-

terminal domains share 88.5% identity. The first zinc-finger of mouse GTSF1 is 37.5%

and 45.8% identical to its fly and moth orthologs, but the mouse protein shares just 8%

and 8.3% identity with the C-terminal domains of the fly and moth proteins, respectively.

Consistent with the evolutionary divergence of their carboxy terminal domains, testis

lysate from rat or rhesus macaque enhanced target cleavage by mouse MIWI piRISC,

whereas lysate from *Drosophila melanogaster* or *Trichoplusia ni* ovaries, *T. ni* Hi5 cells,

or purified recombinant *Bombyx mori* BmGtsf1 did not (Extended Data Fig. 4c, 5a-b and

Supplementary Data Fig. 1d).

241   GTSF1 paralogs can distinguish among PIWI proteins within a species

Animal genomes often encode more than one GTSF protein[92] (Extended Data Fig. 5c).

For example, *D. melanogaster* has four GTSF paralogs. The *D. melanogaster* OSC and

OSS cell lines, which derive from somatic follicle cells that support oogenesis, express

Piwi but lack the PIWI paralogs Aub or Ago3. Piwi-mediated, piRNA-guided transposon

silencing in these cells requires Asterix, a GTSF1 ortholog[7,8,10]. In vivo, *asterix* mutants

phenocopy *piwi* mutants and are female sterile, even though Piwi is successfully loaded

with piRNAs and transits to the nucleus in the absence of Asterix[7,8]. Whether the other

fly GTSF paralogs have a function in vivo, perhaps as auxiliary factors for Aub or Ago3,

remains to be tested. Like fly Asterix, mouse GTSF1 is essential for piRNA function and

fertility. In mice, two GTSF1 paralogs, GTSF1L and GTSF2 are also expressed during

spermatogenesis and interact with PIWI proteins[94]. Unlike GTSF1, single and double

Gtsf1l and *Gtsf2* knockout males are fertile[94]. Genes encoding the *Gtsf* paralogs are

syntenic in mammals, while *Gtsf2* is lost in primates (Extended Data Fig. 6).

The C-terminal domains of GTSF orthologs are more similar among closely

related species than among GTSF paralogs within the same species (Extended Data

Fig. 4d and 5c and Extended Data Table 1), further supporting the view that this domain

has evolved to bind specific PIWI proteins. Consistent with this idea, mouse GTSF1,

GTSF1L, and GTSF2 differ in their ability to potentiate target cleavage by MIWI and

MILI. While GTSF1 accelerated target cleavage by both MIWI and MILI, purified,

recombinant GTSF1L and GTSF2 (Extended Data Fig. 4b) efficiently potentiated target

cleavage by MIWI but not MILI (Fig. 3c–d and Extended Data Table 2). GTSF2 was

unable to detectably increase the rate of target cleavage by MILI. Although GTSF1L had

a modest effect on the rate of target cleavage by MILI piRISC, this enhancement was

less than one-sixteenth that provided by GTSF1 and half that of the PIWI-binding

mutant GTSF1^{W98A,W107A,W112A} (Fig. 3c–d and Extended Data Table 2). We conclude that

GTSF1L and GTSF2 are specialized to potentiate target cleavage by MIWI piRISC.

Silk moth BmGtsf1-like is more similar to mouse GTSF1 than to BmGtsf1

(Extended Data Fig. 5c and Extended Data Table 1); BmGtsf1-like more than doubled

the amount of target cleavage by MIWI piRISC, but had no detectable effect on MILI

(Extended Data Fig. 5b), further supporting the idea that the GTSF carboxy terminal

domain determines the affinity of the protein for specific PIWI proteins. In vivo, BmGtsf1

interacts with Siwi and is required for transposon silencing and sex determination[11].

BmGtsf1 increased the rate of target cleavage by affinity-purified BmSiwi but not that of

the other silk moth PIWI protein, BmAgo3 (Fig. 3f).

Target cleavage by MIWI or MILI requires extensive pairing between piRNA and

target

piRNA:target RNA complementarity from g2–g22 (i.e., 21 base-pairs) is required for

efficient target cleavage directed by endogenous piRNAs loaded into MIWI piRISC

immunoprecipitated from adult mouse testis[5]. However, GTSF1 does not detectably co-

immunoprecipitate with MIWI piRISC from mouse testis[5,77]. A requirement for 21 bp

complementarity between target and guide is unprecedented among Argonaute

proteins: fly Ago2 slices a target with as few as 11 contiguous base pairs[95]; mammalian

AGO2 requires only 11 contiguous base pairs for detectable cleavage[96]; and the

eubacterial DNA-guided DNA endonuclease TtAgo requires as few as 14 (ref. 97).

Affinity-purified MIWI, programmed with either of two different synthetic, 30 nt

piRNAs, readily cleaved target RNA complementary to guide nucleotides g2–21 but not

a target complementary to g2–g16 (Extended Data Fig. 7a, b). Under multiple-turnover

conditions with saturating amounts of purified, recombinant GTSF1 ([GTSF1] >> [target]

> [piRISC]), MIWI readily cleaved a target RNA with 19 nucleotides (g2–g20)

complementary to its synthetic piRNA guide (Fig. 4a, top). The lower background of

single-turnover experiments ([GTSF1] >> [piRISC] > [target]) allowed longer incubation

times. Using these conditions, we could detect GTSF1-stimulated cleavage of a target

RNA with as few as 15 complementary nucleotides (g2–g16; Fig. 4a, bottom). We note

that the pachytene stage of meiosis in mouse spermatogenesis lasts about 175 h, and

the pachytene piRNA pathway components are expressed until at least the round

spermatid stage, a time interval spanning >400 h.

Guide length limits the rate of target cleavage by MIWI

In vivo, piRNAs are trimmed to a length characteristic of the PIWI protein in which they reside: ~30 nt for MIWI and ~26–27 nt for MILI[41,42,87]. An attractive hypothesis is that these piRNA lengths are optimal for target cleavage catalyzed by the specific PIWI protein. Our data suggest a more complex relationship between piRNA length, PIWI protein identity, and target complementarity. In the presence of GTSF1, MIWI loaded with a 30 nt piRNA and MILI loaded with a 26 nt piRNA readily cleaved a fully complementary target RNA in a 60 min reaction (Fig. 4b). By contrast, neither piRISC cleaved a target complementary to piRNA positions g2–g16 (Fig. 4a, b and Extended Data Fig. 7a, b). Similarly, MIWI loaded with a 26 nt or MILI loaded with a 26 or 21 nt guide produced little cleavage for a target complementary to positions g2–g16 although both guide lengths supported cleavage of a fully complementary target (Fig. 4b). In fact, for MIWI, a 21mer was more active than a 30 nt guide (Fig. 4b). Without GTSF1, MIWI or MILI loaded with any of these guide lengths produced little cleaved target in 60 min (Fig. 4b). Remarkably, MIWI or MILI loaded with a 16 nt guide RNA, a piRNA length not present in vivo, readily cleaved the RNA target (Fig. 4b).

GTSF1 accelerates the rate of pre-steady state target cleavage (k_{burst}) by MIWI and MILI but has little effect on the nearly slow rate of steady-state cleavage (k_{ss}), suggesting that piRISC remains bound to its cleavage by-products (Fig. 3e). We incubated GTSF1 and MIWI—loaded with a 30, 26, 21, or 16 nt guide—with a target RNA fully complementary to each guide and measured k_{burst} and k_{ss} (Fig. 4c). As the guide length decreased, k_{burst} decreased and k_{ss} increased. Compared to its native 30 nt guide length, the 16 nt guide increased k_{ss} ~9-fold and decreased k_{burst} >50-fold. These data suggest that as the guide was shortened, the cleaved products were released more rapidly.

We estimated the binding affinity (ΔG) of the piRNA for its target using the

nearest-neighbor rules for base pairing at 37°C. As the strength of the piRNA base

pairing to the target increased, the rate of pre-steady-state cleavage increased (Fig. 5a),

consistent with base pairing serving to extract the 3′ end of the piRNA from the PAZ

domain, facilitating the transition to a more catalytically competent piRISC

conformation[98]. Conversely, decreased base-pairing strength increased the steady-state

rate of target cleavage, supporting the view that for biologically relevant piRNA lengths,

product release is the rate-determining step for MIWI-catalyzed target cleavage

(Fig. 5a).

Discussion

In nearly all animals, both piRNA biogenesis and piRNA function require the PIWI

endoribonuclease activity. Yet purified mouse MIWI and MILI are intrinsically slow to

cleave complementary target RNAs. Our data show that unlike AGO proteins, PIWI

proteins require an auxiliary factor, GTSF1, to efficiently cleave their RNA targets.

The ability of GTSF1 to potentiate PIWI-catalyzed target cleavage provides a

biochemical explanation for the genetic requirement for this small zinc-finger protein in

the piRNA pathway. GTSF1 function requires that it bind both RNA and the PIWI

protein, and differences among carboxy-terminal domains restrict individual GTSF1

paralogs to specific PIWI proteins.

We propose a testable kinetic scheme for target cleavage by MIWI (Fig. 5b, c

and Extended Data Fig. 8) that incorporates the requirement for GTSF1 and the

observation that a 16 nt guide changes the rate-determining step of target cleavage

catalyzed by MIWI. As originally proposed for fly Ago2 (ref. 98), piRISC bound to a

target is presumed to exist in two-states: one in which the piRNA 3′ end is secured in

the PAZ domain and a competing, pre-catalytic conformation in which the piRNA is fully

base paired to its target. Structures of Piwi-A from the freshwater sponge *Ephydatia*

fluviatilis show that extensive pairing between a piRNA and its target induces a

catalytically competent conformation in which the PAZ domain is rotated away from the

piRNA 3′ end[64]. We propose that the PAZ-bound state is more favorable for guides

bound to PIWI proteins than to AGOs, requiring a high degree of complementarity

between guide and target to extract the piRNA 3′ terminus from the PAZ domain. Our

data also suggest that a slow rate of product release after cleavage results directly from

the extensive piRNA-target RNA complementarity required for this conformational

change. A 16 nt piRNA allows MIWI more easily to assume a pre-catalytic state,

perhaps because the 3′ end of the short guide cannot reach the PAZ domain. Notably,

single-stranded siRNA guides as short as 14 nt allow mammalian AGO3, initially

believed to have lost its endonuclease activity, to efficiently cleave RNA targets[99]. In

golden hamsters, piRNAs bound to PIWIL1 are initially ~29 nt long, but a shorter

population of ~23 nt piRNAs appears at metaphase II and predominates in two-cell

embryos[100]. piRNAs bound to PIWIL3, a female-specific PIWI protein absent from mice,

are ~19 nt long in hamster and ~20 nt in human oocytes[100,101]. We speculate that these

short piRNAs allow PIWIL1 and PIWIL3 to function as multiple-turnover endonucleases.

Our model proposes that GTSF1 recognizes the pre-catalytic piRISC state,

facilitating a second conformational change in PIWI proteins that may occur

spontaneously in AGO clade proteins. GTSF1 binding likely stabilizes the catalytically

active conformation, facilitating target cleavage, but has no detectable effect on

subsequent release of the cleaved products. Slow product release may be a general

property of PIWI proteins: purified *Ephydatia fluviatilis* (freshwater sponge) PIWI

similarly catalyzes only a single round of target cleavage[64].

References

1. Ozata, D. M., Gainetdinov, I., Zoch, A., O'Carroll, D. & Zamore, P. D. PIWI-
interacting RNAs: small RNAs with big functions. *Nat. Rev. Genet.* **20**, 89-108
(2018).

2. Kuramochi-Miyagawa, S. et al. *Mili*, a mammalian member of *piwi* family gene, is
essential for spermatogenesis. *Development* **131**, 839-849 (2004).

3. Deng, W. & Lin, H. *miwi*, a murine homolog of *piwi*, encodes a cytoplasmic protein
essential for spermatogenesis. *Dev. Cell* **2**, 819-830 (2002).

4. Carmell, M. A. et al. MIWI2 is essential for spermatogenesis and repression of
transposons in the mouse male germline. *Dev. Cell* **12**, 503-514 (2007).

5. Reuter, M. et al. Miwi catalysis is required for piRNA amplification-independent
LINE1 transposon silencing. *Nature* **480**, 264-267 (2011).

6. De Fazio, S. et al. The endonuclease activity of Mili fuels piRNA amplification that
silences LINE1 elements. *Nature* **480**, 259-263 (2011).

7. Dönertas, D., Sienski, G. & Brennecke, J. *Drosophila* Gtsf1 is an essential
component of the Piwi-mediated transcriptional silencing complex. *Genes Dev.* **27**,
1693-1705 (2013).

8. Ohtani, H. et al. DmGTSF1 is necessary for Piwi-piRISC-mediated transcriptional
transposon silencing in the *Drosophila* ovary. *Genes Dev.* **27**, 1656-1661 (2013).

9. Yoshimura, T. et al. Mouse GTSF1 is an essential factor for secondary piRNA
biogenesis. *EMBO Rep.* **19**, e42054 (2018).

10. Muerdter, F. et al. A genome-wide RNAi screen draws a genetic framework for
transposon control and primary piRNA biogenesis in *Drosophila*. *Mol. Cell.* **50**,
736-748 (2013).

11. Chen, K. et al. Gtsf1 is essential for proper female sex determination and

transposon silencing in the silkworm, *Bombyx mori*. *PLoS Genet.* **16**, e1009194

(2020).

12. Yoshimura, T. et al. *Gtsf1/Cue110*, a gene encoding a protein with two copies of a

CHHC Zn-finger motif, is involved in spermatogenesis and retrotransposon

suppression in murine testes. *Dev. Biol.* **335**, 216-227 (2009).

13. Mohn, F., Handler, D. & Brennecke, J. Noncoding RNA. piRNA-guided slicing

specifies transcripts for Zucchini-dependent, phased piRNA biogenesis. *Science*

348, 812-817 (2015).

14. Han, B. W., Wang, W., Li, C., Weng, Z. & Zamore, P. D. Noncoding RNA. piRNA-

guided transposon cleavage initiates Zucchini-dependent, phased piRNA

production. *Science* **348**, 817-821 (2015).

15. Homolka, D. et al. PIWI Slicing and RNA Elements in Precursors Instruct

Directional Primary piRNA Biogenesis. *Cell Rep* **12**, 418-428 (2015).

16. Gainetdinov, I., Colpan, C., Arif, A., Cecchini, K. & Zamore, P. D. A Single

Mechanism of Biogenesis, Initiated and Directed by PIWI Proteins, Explains piRNA

Production in Most Animals. *Mol. Cell.* **71**, 775-790.e5 (2018).

17. Ipsaro, J. J., Haase, A. D., Knott, S. R., Joshua-Tor, L. & Hannon, G. J. The

structural biochemistry of Zucchini implicates it as a nuclease in piRNA biogenesis.

Nature **491**, 279-283 (2012).

18. Ding, D. et al. PNLDC1 is essential for piRNA 3′ end trimming and transposon

silencing during spermatogenesis in mice. *Nat Commun* **8**, 819 (2017).

19. Zhang, Y. et al. An essential role for PNLDC1 in piRNA 3′ end trimming and male

fertility in mice. *Cell Res* **27**, 1392-1396 (2017).

20. Nishimura, T. et al. PNLDC1, mouse pre-piRNA Trimmer, is required for meiotic

and post-meiotic male germ cell development. *EMBO Rep.* **19**, (2018).

21. Wang, H. et al. Antagonistic roles of Nibbler and Hen1 in modulating piRNA 3'

ends in *Drosophila*. *Development* **143**, 530-539 (2016).

22. Feltzin, V. L. et al. The exonuclease Nibbler regulates age-associated traits and

modulates piRNA length in *Drosophila*. *Aging Cell* **14**, 443-452 (2015).

23. Izumi, N., Shoji, K., Suzuki, Y., Katsuma, S. & Tomari, Y. Zucchini consensus

motifs determine the mechanism of pre-piRNA production. *Nature* **578**, 311-316

(2020).

24. Izumi, N. et al. Identification and Functional Analysis of the Pre-piRNA 3' Trimmer

in Silkworms. *Cell* **164**, 962-973 (2016).

25. Haase, A. D. et al. Probing the initiation and effector phases of the somatic piRNA

pathway in *Drosophila*. *Genes Dev.* **24**, 2499-2504 (2010).

26. Gunawardane, L. S. et al. A Slicer-Mediated Mechanism for Repeat-Associated

siRNA 5′ End Formation *in Drosophila*. *Science* **315**, 1587-1590 (2007).

27. Brennecke, J. et al. Discrete Small RNA-Generating Loci as Master Regulators of

Transposon Activity in *Drosophila*. *Cell* **128**, 1089-1103 (2007).

28. Lewis, S. H. et al. Pan-arthropod analysis reveals somatic piRNAs as an ancestral

defence against transposable elements. *Nat Ecol Evol* **2**, 174-181 (2018).

29. Fu, Y. et al. The genome of the Hi5 germ cell line from *Trichoplusia ni*, an

agricultural pest and novel model for small RNA biology. *eLife* **7**, e31628 (2018).

30. Jehn, J. et al. PIWI genes and piRNAs are ubiquitously expressed in mollusks and

show patterns of lineage-specific adaptation. *Commun Biol* **1**, 137 (2018).

31. Lim, R. S., Anand, A., Nishimiya-Fujisawa, C., Kobayashi, S. & Kai, T. Analysis of
Hydra PIWI proteins and piRNAs uncover early evolutionary origins of the piRNA
pathway. *Dev. Biol.* **386**, 237-251 (2014).

32. Juliano, C. E. et al. PIWI proteins and PIWI-interacting RNAs function in Hydra
somatic stem cells. *Proc. Natl. Acad. Sci. U S A* **111**, 337-342 (2014).

33. Praher, D. et al. Characterization of the piRNA pathway during development of the
sea anemone *Nematostella vectensis*. *RNA Biol* **14**, 1727-1741 (2017).

34. Grimson, A. et al. Early origins and evolution of microRNAs and Piwi-interacting
RNAs in animals. *Nature* **455**, 1193-1197 (2008).

35. Kerner, P., Degnan, S. M., Marchand, L., Degnan, B. M. & Vervoort, M. Evolution
of RNA-binding proteins in animals: insights from genome-wide analysis in the
sponge *Amphimedon queenslandica*. *Mol. Biol. Evol.* **28**, 2289-2303 (2011).

36. Murchison, E. P. et al. Conservation of small RNA pathways in platypus. *Genome*
Res. **18**, 995-1004 (2008).

37. Houwing, S. et al. A role for Piwi and piRNAs in germ cell maintenance and
transposon silencing in Zebrafish. *Cell* **129**, 69-82 (2007).

38. Toombs, J. A. et al. Xenopus Piwi proteins interact with a broad proportion of the
oocyte transcriptome. *RNA* **23**, 504-520 (2017).

39. Sun, Y. H. et al. Domestic chickens activate a piRNA defense against avian
leukosis virus. *eLife* **6**, e24695 (2017).

40. Grivna, S. T., Beyret, E., Wang, Z. & Lin, H. A novel class of small RNAs in mouse
spermatogenic cells. *Genes Dev.* **20**, 1709-1714 (2006).

41. Aravin, A. et al. A novel class of small RNAs bind to MILI protein in mouse testes.
Nature **442**, 203-207 (2006).

42.   Girard, A., Sachidanandam, R., Hannon, G. J. & Carmell, M. A. A germline-specific
469   class of small RNAs binds mammalian Piwi proteins. *Nature* **442**, 199-202 (2006).

470   43.   Lim, S. L. et al. Conservation and expression of PIWI-interacting RNA pathway
471   genes in male and female adult gonad of amniotes. *Biol. Reprod.* **89**, 136 (2013).

472   44.   Lau, N. C. et al. Characterization of the piRNA complex from rat testes. *Science*
473   **313**, 363-367 (2006).

474   45.   Özata, D. M. et al. Evolutionarily conserved pachytene piRNA loci are highly
475   divergent among modern humans. *Nat Ecol Evol* **4**, 156-168 (2020).

476   46.   Li, X. Z. et al. An Ancient Transcription Factor Initiates the Burst of piRNA
477   Production during Early Meiosis in Mouse Testes. *Mol. Cell.* **50**, 67-81 (2013).

478   47.   Wu, P. H. et al. The evolutionarily conserved piRNA-producing locus *pi6* is
479   required for male mouse fertility. *Nat. Genet.* **52**, 728-739 (2020).

480   48.   Choi, H., Wang, Z. & Dean, J. Sperm acrosome overgrowth and infertility in mice
481   lacking chromosome 18 pachytene piRNA. (2020).

482   49.   Gou, L. T. et al. Pachytene piRNAs instruct massive mRNA elimination during late
483   spermiogenesis. *Cell Res* 1-21 (2014).

484   50.   Goh, W. S. et al. piRNA-directed cleavage of meiotic transcripts regulates
485   spermatogenesis. *Genes Dev.* **29**, 1032-1044 (2015).

486   51.   Vourekas, A. et al. Mili and Miwi target RNA repertoire reveals piRNA biogenesis
487   and function of Miwi in spermiogenesis. *Nat. Struct. Mol. Biol.* **19**, 773-781 (2012).

488   52.   Zhang, P. et al. MIWI and piRNA-mediated cleavage of messenger RNAs in
489   mouse testes. *Cell Res* **25**, 193-207 (2015).

490   53.   Wang, Y., Sheng, G., Juranek, S., Tuschl, T. & Patel, D. J. Structure of the guide-
491   strand-containing argonaute silencing complex. *Nature* **456**, 209-213 (2008).

54. Schirle, N. T. & MacRae, I. J. The crystal structure of human Argonaute2. *Science*
336, 1037-1040 (2012).

55. Matsumoto, N. et al. Crystal Structure of Silkworm PIWI-Clade Argonaute Siwi
Bound to piRNA. *Cell* **167**, 484-497.e9 (2016).

56. Parker, J. S., Roe, S. M. & Barford, D. Crystal structure of a PIWI protein suggests
mechanisms for siRNA recognition and slicer activity. *EMBO J.* **23**, 4727-4737
(2004).

57. Ma, J. B. et al. Structural basis for 5′-end-specific recognition of guide RNA by the
A. fulgidus Piwi protein. *Nature* **434**, 663-666 (2005).

58. Yuan, Y. R. et al. Crystal structure of *A. aeolicus* Argonaute, a site-specific DNA-
guided endoribonuclease, provides insights into RISC-mediated mRNA cleavage.
Mol. Cell. **19**, 405-419 (2005).

59. Ma, J. B. et al. Structural basis for 5′-end-specific recognition of guide RNA by the
A. fulgidus Piwi protein. *Nature* **434**, 666-670 (2005).

60. Le Thomas, A., Marinov, G. & Aravin, A. A Transgenerational Process Defines
piRNA Biogenesis in *Drosophila virilis*. *Cell Reports* **8**, 1617-1623 (2014).

61. Yamaguchi, S. et al. Crystal structure of Drosophila Piwi. *Nat Commun* **11**, 858
(2020).

62. Kawaoka, S., Izumi, N., Katsuma, S. & Tomari, Y. 3′ end formation of PIWI-
interacting RNAs in vitro. *Mol. Cell.* **43**, 1015-1022 (2011).

63. Cora, E. et al. The MID-PIWI module of Piwi proteins specifies nucleotide- and
strand-biases of piRNAs. *RNA* **20**, 773-81 (2014).

64. Anzelon, T. A. et al. Structural basis for piRNA targeting. *Nature* **597**, 285-289
(2021).

65.   Kirino, Y. & Mourelatos, Z. The mouse homolog of HEN1 is a potential methylase

for Piwi-interacting RNAs. *RNA* **13**, 1397-1401 (2007).

66.   Kirino, Y. & Mourelatos, Z. 2′-*O*-methyl modification in mouse piRNAs and its

methylase. *Nucleic Acids Symp Ser (Oxf)* 417-418 (2007).

67.   Lim, S. L. et al. HENMT1 and piRNA Stability Are Required for Adult Male Germ

Cell Transposon Repression and to Define the Spermatogenic Program in the

Mouse. *PLoS Genet.* **11**, e1005620 (2015).

68.   Kamminga, L. M. et al. Hen1 is required for oocyte development and piRNA

stability in zebrafish. *EMBO J.* **29**, 3688-3700 (2010).

69.   Montgomery, T. A. et al. PIWI associated siRNAs and piRNAs specifically require

the *Caenorhabditis elegans* HEN1 ortholog henn-1. *PLoS Genet.* **8**, e1002616

(2012).

70.   Billi, A. C. et al. The *Caenorhabditis elegans* HEN1 ortholog, HENN-1, methylates

and stabilizes select subclasses of germline small RNAs. *PLoS Genet.* **8**,

e1002617 (2012).

71.   Kamminga, L. M. et al. Differential impact of the HEN1 homolog HENN-1 on 21U

and 26G RNAs in the germline of *Caenorhabditis elegans*. *PLoS Genet.* **8**,

e1002702 (2012).

72.   Pelisson, A., Sarot, E., Payen-Groschene, G. & Bucheton, A. A novel repeat-

associated small interfering RNA-mediated silencing pathway downregulates

complementary sense *gypsy* transcripts in somatic cells of the *Drosophila* ovary. *J.*

Virol. **81**, 1951-1960 (2007).

73.   Ameres, S. L. et al. Target RNA-directed trimming and tailing of small silencing

RNAs. *Science* **328**, 1534-1539 (2010).

74.   Salomon, W. E., Jolly, S. M., Moore, M. J., Zamore, P. D. & Serebrov, V. Single-
541   Molecule Imaging Reveals that Argonaute Reshapes the Binding Properties of Its
542   Nucleic Acid Guides. *Cell* **162**, 84-95 (2015).

543   75.   Becker, W. R. et al. High-Throughput Analysis Reveals Rules for Target RNA
544   Binding and Cleavage by AGO2. *Mol. Cell.* **75**, 741-755.e11 (2019).

545   76.   Hutvagner, G. & Zamore, P. D. A microRNA in a multiple-turnover RNAi enzyme
546   complex. *Science* **297**, 2056-2060 (2002).

547   77.   Vagin, V. V. et al. Proteomic analysis of murine Piwi proteins reveals a role for
548   arginine methylation in specifying interaction with Tudor family members. *Genes*
549   *Dev.* **23**, 1749-1762 (2009).

550   78.   Siomi, M. C., Mannen, T. & Siomi, H. How does the royal family of Tudor rule the
551   PIWI-interacting RNA pathway? *Genes Dev.* **24**, 636-646 (2010).

552   79.   Bedford, M. T. & Clarke, S. G. Protein arginine methylation in mammals: who,
553   what, and why. *Mol. Cell.* **33**, 1-13 (2009).

554   80.   Kuramochi-Miyagawa, S. et al. Two mouse *piwi*-related genes: *miwi* and *mili*. *Mech*
555   *Dev* **108**, 121-133 (2001).

556   81.   Michael, S. F., Kilfoil, V. J., Schmidt, M. H., Amann, B. T. & Berg, J. M. Metal
557   binding and folding properties of a minimalist Cys2His2 zinc finger peptide. *Proc.*
558   *Natl. Acad. Sci. U S A* **89**, 4796-4800 (1992).

559   82.   Matt, T., Martinez-Yamout, M. A., Dyson, H. J. & Wright, P. E. The CBP/p300
560   TAZ1 domain in its native state is not a binding partner of MDM2. *Biochem. J.* **381**,
561   685-691 (2004).

562   83.   Nyborg, J. K. & Peersen, O. B. That zincing feeling: the effects of EDTA on the
563   behaviour of zinc-binding transcriptional regulators. *Biochem. J.* **381**, e3-4 (2004).

84. Almeida, M. V. et al. GTSF-1 is required for formation of a functional RNA-

dependent RNA Polymerase complex in *Caenorhabditis elegans. EMBO J.* **37**,

e99325 (2018).

85. Kuramochi-Miyagawa, S. et al. DNA methylation of retrotransposon genes is

regulated by Piwi family members MILI and MIWI2 in murine fetal testes. *Genes*

Dev. **22**, 908-917 (2008).

86. Kojima-Kita, K. et al. MIWI2 as an Effector of DNA Methylation and Gene Silencing

in Embryonic Male Germ Cells. *Cell Rep* **16**, 2819-2828 (2016).

87. Aravin, A. A. et al. A piRNA pathway primed by individual transposons is linked to

de novo DNA methylation in mice. *Mol. Cell.* **31**, 785-799 (2008).

88. Wenda, J. M. et al. Distinct Roles of RNA Helicases MVH and TDRD9 in PIWI

Slicing-Triggered Mammalian piRNA Biogenesis and Function. *Dev. Cell* **41**, 623-

637.e9 (2017).

89. Xiol, J. et al. RNA Clamping by Vasa Assembles a piRNA Amplifier Complex on

Transposon Transcripts. *Cell* (2014).

90. Nishida, K. M. et al. Respective functions of two distinct Siwi complexes

assembled during PIWI-interacting RNA biogenesis in *Bombyx* germ cells. *Cell*

Rep **10**, 193-203 (2015).

91. Murakami, R., Sumiyoshi, T., Negishi, L. & Siomi, M. C. DEAD-box polypeptide 43

facilitates piRNA amplification by actively liberating RNA from Ago3-piRISC. *EMBO*

Rep. **22**, e51313 (2021).

92. Andreeva, A. & Tidow, H. A novel CHHC Zn-finger domain found in spliceosomal

proteins and tRNA modifying enzymes. *Bioinformatics* **24**, 2277-2280 (2008).

93. Ipsaro, J. J., O'Brien, P. A., Bhattacharya, S., Palmer III, A. G. & Joshua-Tor, L.

Asterix/Gtsf1 links tRNAs and piRNA silencing of retrotransposons. *Cell Reports*

34, 108914 (2021).

94. Takemoto, N., Yoshimura, T., Miyazaki, S., Tashiro, F. & Miyazaki, J. *Gtsf1l and*

Gtsf2 Are Specifically Expressed in Gonocytes and Spermatids but Are Not

Essential for Spermatogenesis. *PLoS One* **11**, e0150390 (2016).

95. Haley, B. & Zamore, P. D. Kinetic analysis of the RNAi enzyme complex. *Nat.*

Struct. Mol. Biol. **11**, 599-606 (2004).

96. Jackson, A. L. et al. Expression profiling reveals off-target gene regulation by

RNAi. *Nat. Biotechnol.* **21**, 635-637 (2003).

97. Wang, Y. et al. Nucleation, propagation and cleavage of target RNAs in Ago

silencing complexes. *Nature* **461**, 754-761 (2009).

98. Tomari, Y. & Zamore, P. D. Perspective: machines for RNAi. *Genes Dev.* **19**, 517-

529 (2005).

99. Park, M. S., Sim, G., Kehling, A. C. & Nakanishi, K. Human Argonaute2 and

Argonaute3 are catalytically activated by different lengths of guide RNA.

Proceedings of the National Academy of Sciences **117**, 28576-28578 (2020).

100. Ishino, K. et al. Hamster PIWI proteins bind to piRNAs with stage-specific size

variations during oocyte maturation. *bioRxiv* 2020.12.01.407411; doi:

https://doi.org/10.1101/2020.12.01.407411 (2020).

101. Yang, Q. et al. Single-cell CAS-seq reveals a class of short PIWI-interacting RNAs

in human oocytes. *Nature Comm.* **10**, 1-15 (2019).

Methods

Plasmids and cell lines

Supplementary Data Table 1 provides the sequences of all oligonucleotides used. To

create pScalps_Puro_EGFP, an IRES-driven EGFP was inserted downstream of the

multiple cloning site and upstream of the puromycin coding sequence of the lentivirus

transfer vector pScalps_Puro. MIWI cDNA was obtained from Mammalian Gene

Collection (https://genecollections.nci.nih.gov/MGC/). MILI cDNA was amplified by RT-

PCR from mouse testis total RNA. Gibson assembly and restriction cloning were used

to clone the MILI and MIWI coding sequences into pScalps_Puro_EGFP, fusing them

in-frame with N-terminal 3XFLAG and SNAP tags. Lentivirus transfer vectors were

packaged by co-transfection with psPAX2 and pMD2.G (4:3:1) using TransIT-2020

(Mirus Bio) in HEK293T cells. Supernatant containing lentivirus was used to transduce

HEK293T cells in the presence of 16 ug/ml polybrene (Sigma) to obtain stable PIWI-

expressing cell lines. Three sequential transductions were performed to maximize

recombinant protein production. The transduced cells were selected in the presence of

2 ug/ml Puromycin for two weeks, then the cells expressing the 5–10% highest EGFP

fluorescence were selected by FACS (UMASS Medical School Flow Cytometry Core).

Selected cells stably expressing the recombinant PIWI proteins were expanded,

harvested, and cell pellets flash-frozen and stored at −80°C.

Mouse GTSF1 cDNA was synthesized at Twist Biosciences and cloned into

pCold-GST (Takara Bio) bacterial expression vector by restriction cloning. GTSF1

mutants, GTSF1L, and GTSF2- expressing pCold-GST vectors were synthesized at

Twist Biosciences. pIZ-FLAG6His-Siwi was described previously[102]. FLAG-tagged MILI

and MIWI-expressing vectors were the kind gift of Shinpei Kawaoka (Kyoto University,

Kyoto, Japan). MmGtsf1 cDNA was amplified by RT-PCR from mouse spermatogonial

stem cell total RNA[103]. BmGtsf1 and BmGtsf1-like cDNAs were amplified by RT-PCR

from BmN4 cell total RNA. The amplified cDNA fragment and a DNA fragment coding

637   V5SBP were cloned into pcDNA5/FRT/TO vector (Thermo Fisher Scientific) by In-fusion

638   cloning (Takara). The *Ephydatia muelleri* Gtsf1 coding sequence was obtained from [104]

639   and cloned into bacterial expression vector pSV272 (IJM).

Mice

Generation of 3XFLAG-HA-tagged GTSF1 mice (*Gtsf1$^{KI/KI}$*) was performed at Cyagen.

The coding sequence for the tags was inserted into the endogenous locus by CRISPR-

Cas9. Briefly, fertilized mouse embryos were injected with the sgRNA targeting the

sequence GTCTTCCATGCTGATGGCAA<u>AGG</u> (<u>PAM</u>), a 3XFLAG-HA-tag cassette HDR

donor, and Cas9 mRNA. Supplementary Data Table 1 provides the sequences of the

HDR donor and oligonucleotide primers used for genotyping. Founder F0 animals were

genotyped and bred to generate F1 animals carrying the germline-transmitted knock-in

allele. Mice were maintained and used according to the guidelines of the Institutional

Animal Care and Use Committee of the University of Massachusetts Chan Medical

School (A201900331). C57BL/6J mice (RRID: IMSR_JAX:000664) were used as wild-

type control, where indicated.

Recombinant protein purification

PIWI proteins. PIWI-expressing stable cells were harvested by centrifugation and stored

at −80°C until lysed in 10 ml lysis buffer (30 mM HEPES-KOH, pH 7.5, 100 mM

potassium acetate, 3.5 mM magnesium acetate, 1 mM DTT, 0.1% v/v Triton X-100,

20% v/v glycerol, and 1× protease inhibitor cocktail [1 mM 4-(2-

aminoethyl)benzenesulfonyl fluoride hydrochloride [Sigma; A8456], 0.3 μM Aprotinin, 40

μM betanin hydrochloride, 10 μM. E-64 [Sigma; E3132], 10 μM leupeptin hemisulfate])

per g frozen cells. Cell lysis was monitored by staining with trypan blue. Crude

cytoplasmic lysate was clarified at 20,000 (S20) or 100,000 × *g* (S100) for 30 mins,

aliquoted, flash-frozen, and stored at −80°C. To capture PIWI proteins, clarified lysate

was incubated with 20 µl Anti-FLAG M2 paramagnetic beads (Sigma) per ml of lysate

for 2 h to overnight rotating at 4°C. Beads were washed five times with high salt wash

buffer (30 mM HEPES-KOH, pH 7.5, 2 M potassium acetate, 3.5 mM magnesium

acetate, 1 mM DTT), twice with low salt buffer (high salt wash buffer except containing

100 mM potassium acetate and 0.01% v/v Triton X-100). To assemble RISC, beads

were resuspended in low salt buffer containing 100 nM synthetic guide piRNA and

incubated with rotation at 37°C or room temperature for 30 min. MIWI RISC or unloaded

apo-MIWI was eluted from the beads with 200 ng/ul 3XFLAG peptide in lysis buffer

without 0.1% v/v Triton X-100 or protease inhibitors for 2 h at 4°C. Eluate containing

PIWI piRISC was aliquoted and stored at −80°C. EfPiwi piRISC purification has been

described[64]. AGO2 siRISC was purified using oligo-affinity purification as described[105].

GTSF1, GTSF1 mutants, and GTSF1 homologs. pCold-GST GTSF-expression vectors

were transformed into Rosetta-Gami 2 competent cells (Novagen). Cells were grown in

the presence of 1 µM $ZnSO_4$ at 37°C until OD_{600} ~0.6–0.8, then chilled on ice for 30 min

to initiate cold shock. Protein expression was induced with 0.5 mM IPTG for 18 h at

15°C. Cells were harvested by centrifugation, washed twice with PBS, and cell pellets

were flash frozen and stored at −80°C. Cell pellets were resuspended in lysis/GST

column buffer containing 20 mM Tris-HCl pH 7.5, 500 mM NaCl, 1 mM DTT, 5% v/v

glycerol, and complete EDTA-free protease inhibitor cocktail (Roche). Cells were lysed

by a single pass at 18,000 psi through a high-pressure microfluidizer (Microfluidics

M110P), and the resulting lysate clarified at 30,000 × *g* for 1 h at 4°C. Clarified lysate

was filtered through a 0.2 µm low-protein binding syringe filter (Millex Durapore; EMD

Millipore) and applied to glutathione Sepharose 4b resin (Cytiva) equilibrated with GST

column buffer. After draining the flowthrough, the resin was washed with 50 column-

volumes GST column buffer. To elute the bound protein and cleave the GST tag in a

single step, 50 U HRV3C Protease (Novagen) was added to the column, the column

was sealed and incubated for 3 h at 4°C, following which, the column was drained to

collect the cleaved protein. The protein was diluted to 50 mM NaCl and further purified

using a Hi-Trap Q (Cytiva) anion exchange column equilibrated with 20 mM Tris-HCl,

pH 7.5, 50 mM NaCl, 1 mM DTT, and 5% v/v glycerol. The bound protein was eluted

using a 100–500 mM NaCl gradient in the same buffer. Peak fractions were analyzed

for purity by SDS-PAGE and the purest were pooled and dialyzed into storage buffer

containing 30 mM HEPES-KOH, pH 7.5, 100 mM potassium acetate, 3.5 mM

magnesium acetate, 1 mM DTT, 20% v/v glycerol. To avoid precipitation of GTSF2,

which has a pI = 7.3, 20 mM Tris-HCl, pH 8.8, was substituted for HEPES-KOH during

purification and dialysis.

For Figure 3f and Extended Data Fig. 5b, HEK293T cells were transfected with

mouse and insect GTSF1 coding sequences cloned into pcDNA5 expression vectors

using Lipofectamine 3000 (Thermo Fisher Scientific) according to the manufacturer's

instructions. Cells were harvested 36 h later and homogenized in lysis buffer [20 mM

Tris-HCl (pH 7.4), 150 mM NaCl, 1.5 mM $MgCl_2$, 0.15% v/v Triton X-100, 100 µg/ml

RNase A (Qiagen), 0.5 mM DTT, 1× Complete EDTA-free protease inhibitor (Roche)].

After centrifugation at 17,000 × g for 20 min at 4°C, additional NaCl (final concentration:

0.65 M) and 1% v/v Triton X-100 (f.c.) were added to the supernatant and the lysate

was incubated with Streptavidin-Sepharose High Performance (Cytiva) beads at 4°C for

1 h. The beads were washed with wash buffer [20 mM Tris-HCl (pH 7.4), 1 M NaCl,

1.5 mM $MgCl_2$, 1% v/v Triton X-100, 0.5 mM DTT] five times and rinsed with lysis buffer

without RNase A. SBP-tagged recombinant proteins were eluted with elution buffer

(30 mM HEPES-KOH, pH 7.4, 100 mM potassium acetate, 2 mM magnesium acetate,

2.5 mM biotin, 0.5 mM DTT).

For Figure 3g, EmGtsf1 expression vector was transformed into BL21(DE3) cells

(New England Biolabs). Transformed cells were grown in LB supplemented with 1 µM

ZnSO4 at 37°C until OD600 ~0.6–0.8. The incubation temperature was lowered to 16°C

and protein expression was induced by addition of 1 mM IPTG for 4 h. Cells were

harvested by centrifugation and cell pellets were flash frozen in liquid nitrogen and

stored at −80°C for future use. Thawed cell pellets were resuspended in lysis buffer

(50 mM Tris, pH 8, 300 mM NaCl, 0.5 mM TCEP) and passed through a high-pressure

(18,000 psi) microfluidizer (Microfluidics M110P) to induce cell lysis. The resulting lysate

was clarified by centrifugation at 30,000 × g for 20 min at 4°C. Clarified lysate was

applied to Ni-NTA resin (Qiagen) and incubated for 1 h. The resin was extensively

washed with Nickel Wash Buffer (300 mM NaCl, 20 mM imidazole, 0.5 mM TCEP,

50 mM Tris, 8). Protein was eluted in four column volumes of Nickel Elution Buffer

(300 mM NaCl, 300 mM imidazole, 0.5 mM TCEP, 50 mM Tris, pH 8). TEV protease

was added to the eluted protein to induce cleavage and removal of the N-terminal His6

and MBP tags. The resulting mixture was dialyzed against Hi-Trap Dialysis Buffer

(300 mM NaCl, 20 mM imidazole, 0.5 mM TCEP, 50 mM Tris, pH 8) at 4°C overnight.

The dialyzed protein was then passed through a 5-ml Hi-Trap Chelating column (Cytiva)

and the unbound material collected. Unbound material was concentrated and further

purified by size exclusion chromatography using a Superdex 200 Increase 10/300

column (Cytiva) equilibrated in 50 mM Tris, pH 8, 300 mM NaCl, and 0.5 mM TCEP.

Peak fractions were analyzed for purity by SDS-PAGE, and the purest were pooled,

concentrated to 150 μM, aliquoted, and stored at −80°C.

Immunodepletion and Western Blotting

Approximately 300,000 FACS-purified Secondary Spermatocytes (Sp2) from wild-type

(C57BL/6) or 3XFLAG-HA-GTSF1-expressing (*Gtsf1*$^{KI/KI}$) mice were lysed in lysis buffer

(30 mM HEPES-KOH, pH 7.5, 100 mM potassium acetate, 3.5 mM magnesium acetate,

1 mM DTT, 0.1% v/v Triton X-100, 20% v/v glycerol, and 1× protease inhibitor cocktail

[1 mM 4-(2-Aminoethyl)benzenesulfonyl fluoride hydrochloride [Sigma; A8456], 0.3 μM

Aprotinin, 40 μM betanin hydrochloride, 10 μM. E-64 [Sigma; E3132], 10 μM leupeptin

hemisulfate]). Crude cytoplasmic lysate was clarified at 20,000 × g (S20) for 30 min at

4°C. The S20 protein concentration was measured using Bradford Assay. The S20 was

incubated with Anti-FLAG paramagnetic beads overnight at 4°C. The supernatant was

removed, and the beads washed thrice with lysis buffer but containing 1M potassium

acetate (high salt wash) and then twice with lysis buffer. Bound proteins were eluted by

incubation with 200 ng/ul 3XFLAG peptide for 2 h at 4°C. The extent of depletion was

assayed by western blotting. Briefly, the samples were resolved on SDS-PAGE and

transferred to nitrocellulose membrane (Amersham Protran 0.45 NC; Cytiva). The

membrane was probed using mouse anti-FLAG antibody (Sigma, F3165; 1:1,000 in

blocking buffer, Rockland Immunochemicals, MB-070) and rabbit anti-Tubulin (Cell

Signaling, 2144; 1:1000) for 2 hours at room temperature. After washing 5× with TBS

(+0.1% v/v Tween-20), the membrane was incubated with secondary antibodies -

donkey anti-rabbit IRDye 680RD (LI-COR, 926-68073; 1:15,000) and goat anti-mouse

IRDye 800CW (LI-COR, 926-32210; 1:15,000) for 30 minutes at room temperature. The

membrane was washed 5× with TBS (+0.1% v/v Tween-20) and imaged using Odyssey

Infrared Imaging System (LI-COR).

Northern Blotting

Northern blotting was performed as described[106]. Briefly, piRNA guide standards and

PIWI RISCs were first resolved on a denaturing 15% polyacrylamide gel, followed by

transfer to Hybond-NX (Cytiva) neutral nylon membrane by semi-dry transfer at 20 V for

1 h. Next, crosslinking was performed in the presence of 0.16 M EDC in 0.13 M 1-

methylimidazole, pH 8.0, at 60°C for 1 h. The crosslinked membrane was pre-hybridized

in Church's buffer (1% w/v BSA, 1 mM EDTA, 0.5 M phosphate buffer, and 7% w/v

SDS) at 45°C for 1 h. Radiolabeled, 5′ ^{32}P-DNA probe (25 pmol) in Church's buffer was

added to the membrane and allowed to hybridize overnight at 45°C, followed by five

washes with 1× SSC containing 0.1% w/v SDS. The membrane was air dried and

exposed to a storage phosphor screen.

Chromatographic fractionation of the MIWI-potentiating activity

Dissected animal tissues were homogenized in lysis buffer in a Dounce homogenizer

using 10 strokes of the loose-fitting pestle A, followed by 20 strokes of tight-fitting pestle

B. Lysate was clarified at 20,000 × g, followed by 0.2 μm filtration to yield an S20 for

further chromatographic purification. Lysates used without further purification were

directly prepared in 30 mM HEPES-KOH, pH 7.5, 100 mM potassium acetate, 3.5 mM

magnesium acetate, 1 mM DTT, and 20% v/v glycerol ("dialysis buffer") with 1×

protease inhibitor homemade cocktail; column fractions were dialyzed into this buffer

before assaying. Protein concentration was measured using the BCA assay.

Chromatography buffers were filtered prior to use.

For chromatography, lysate was prepared as described except using 30 mM

HEPES-KOH, pH 7.5, 50 mM NaCl, 1 mM DTT, 5% v/v glycerol, and protease

inhibitors. The lysate was applied to HiTrap SP column (Cytiva) equilibrated with the

lysis buffer. The column was washed, and the bound proteins eluted stepwise using

increasing NaCl concentrations. The NaCl content of the SP column fractions containing

the peak MIWI-potentiating activity was adjusted to 2 M and applied to HiTrap Phenyl

(Cytiva) equilibrated with column buffer containing 2 M NaCl. Bound proteins were

eluted stepwise using decreasing NaCl concentrations. The peak MIWI-potentiating

fractions elute from the HiTrap Phenyl column were pooled, concentrated (10 kDa

MWCO Amicon Ultra filter), and applied to Superdex 200 Increase 10-300 GL size-

exclusion chromatography column (bed volume ~24 ml) equilibrated with the dialysis

buffer but containing 5% v/v glycerol. The void volume (V_0) of the gel filtration column

was determined with blue dextran, and all fractions (0.5 ml each), starting from just

before V_0 to the end of the column (V_t) were assayed for MIWI potentiating activity. The

molecular weight of the potentiating activity was determined relative to molecular weight

markers (beta amylase, 200 kDa; alcohol dehydrogenase, 150 kDa; albumin, 66 kDa;

carbonic anhydrase, 29 kDa; and cytochrome C, 12.4 kDa).

Zn^{2+} and Ni^{2+} immobilized-metal affinity chromatography

HiTrap Chelating HP (Cytiva) columns were charged with 0.1 M $NiSO_4$ or $ZnSO_4$,

washed with water, and then equilibrated in column buffer (20 mM potassium phosphate

buffer, pH 7.5, 500 mM NaCl, 0.5 mM DTT, 5% v/v glycerol). S20 testis lysate was

applied to the column, the flow-through collected, and the column washed with the

column buffer until absorbance at 280 nm stabilized. Bound proteins were eluted in two

steps: first, with 20 mM potassium phosphate, pH 7.5, 2 M ammonium chloride, 0.5 mM

DTT, 5% v/v glycerol (elution buffer 1), and, second, with 20 mM potassium phosphate,

pH 7.5, 500 mM NaCl, 200 mM imidazole, pH 8.0, 0.5 mM DTT, 5% v/v glycerol (elution

buffer 2). The peak of each step was dialyzed into the dialysis buffer and assayed for

the ability to potentiate MIWI catalysis.

Target cleavage assays

Target RNA substrates for in vitro cleavage assays were prepared as described[74,107,108].

Briefly, piRNA target site-containing templates were amplified by PCR, in vitro

transcribed with T7 RNA polymerase, purified by urea-PAGE, and radiolabeled using α-

^{32}P GTP (3000 Ci/mmol; Perkin Elmer), *S*-adenosyl methionine, and vaccinia virus RNA

guanylyl transferase as described[108]. Unincorporated α-^{32}P GTP was removed using a

G-25 spin column (Cytiva), and target RNA gel purified. In Extended Data Fig. 7a, target

cleavage was monitored using synthetic RNA oligonucleotides radiolabeled by ligating

[5′-^{32}P] cytidine 3′,5′-bisphosphate to the 3′ end of the target with T4 RNA ligase I

(Ambion). The [5′-^{32}P] cytidine 3′,5′-bisphosphate was prepared by incubating 1 mM

cytidine 3′-monophosphate (Sigma) with 312.5 pmole [γ-^{32}P] ATP (6000 Ci/mmol;

Perkin Elmer) with 25 U T4 polynucleotide kinase (NEB) at 37°C for 1 h, followed by

70°C for 30 min to inactivate the kinase.

Radiolabeled target (3–100 nM f.c.) was added to purified PIWI piRISC (2–8 nM),

plus ~1 µg tissue or sorted germ cell lysate per 10 µl reaction volume or 0.5 µM (f.c.) of

purified GTSF protein. At the indicated times, an aliquot of a master reaction was

quenched in 4 volumes 50 mM Tris-HCl, pH 7.5, 100 mM NaCl, 25 mM EDTA, 1% w/v

SDS, then proteinase K (1 mg/ml f.c.) was added and incubated at 45°C for 15 min. An

equal volume of urea loading buffer (8 M urea, 25 mM EDTA) was added to the reaction

time points, heated at 95°C for 2 min, and resolved by 7–10% denaturing PAGE. Gels

were dried, exposed to a storage phosphor screen, and imaged on a Typhoon FLA

7000 (GE).

The raw image file was used to quantify the substrate and product bands,

corrected for background. Data were fit to the reaction scheme

$$E + S \underset{k_{-1}}{\overset{k_1}{\rightleftharpoons}} ES \overset{k_2}{\rightarrow} EP \overset{k_3}{\rightarrow} E + P$$

using the burst-and-steady-state equation, Equation 1.

$[P] = f(t) = [E]([k_2/(k_2 + k_3)]^2 \times (1 - e^{-[k2+k]t}) + [k_2k_3/(k_2 + k_3)]t)$　　　(Equation 1)

The time-dependence of product formation corresponded to a pre-steady-state

exponential burst ($k_{burst} = k_2 + k_3$) followed by a linear steady-state phase, described by

k_{cat}, where $k_{cat} = k_{ss} = k_2k_3/(k_2 + k_3)$.

The affinity (K_D) of wild-type or GTSF1[W98A,W107A,W112A] for MIWI piRISC-target

ternary complex and the maximum observable rate (k_{pot}) were estimated by measuring

the pre-steady-state rate of target cleavage (k_{burst}) of MIWI at increasing concentrations

of GTSF1 (1–5,000 nM) and fitting the data to Equation 2 (ref. 109:

$k_{burst} = k_{pot} [GTSF1]/(K_D + [GTSF1])$　　　(Equation 2).

FLAG-Siwi, MIWI, and MILI target-cleavage assays

FLAG-tagged BmSiwi, MIWI, or MILI-expressing vectors were transfected into BmN4

cells with X-tremeGENE HP DNA Transfection Reagent (Sigma). Preparation of loading

lysates and single-stranded RNA loading were as described[110]. After loading with

synthetic guide RNAs in the cell lysate, piRISC was immunoprecipitated with anti-FLAG

antibody (Sigma) conjugated to Dynabeads protein G superparamagnetic beads

(Thermo Fisher Scientific). The beads were washed five times with lysis buffer

containing 30 mM HEPES-KOH, pH 7.4, 100 mM potassium acetate, 2 mM magnesium

acetate, 0.5 mM DTT, and 0.1% v/v Empigen. Target cleavage assays were performed

at 25°C for 3.5 h in 8 µl reaction containing 2.4 µl "40×" reaction mix[108]. 100 nM

recombinant GTSF1 protein and 0.5 nM ^{32}P cap-radiolabeled 28 nt target RNA as

described[24].

BmAgo3 target-cleavage assays

Naive BmN4 cells were resuspended in lysis buffer (30 mM HEPES-KOH, pH 7.4,

100 mM potassium acetate, 2 mM magnesium acetate, 0.05% v/v Triton X-100, 0.5 mM

DTT, 1× Complete EDTA-free protease inhibitor [Roche]) and lysed using a Dounce

homogenizer. After centrifugation at 17,000 × g for at 4°C 20 min, the supernatant was

incubated at 4°C for 1 h with anti-BmAgo3 antibody[23] conjugated to Dynabeads

protein G (Thermo Fisher Scientific). The superparamagnetic beads were washed five

times with wash buffer (30 mM HEPES-KOH, pH 7.4, 100 mM potassium acetate, 2 mM

magnesium acetate, 0.5% v/v Empigen, 0.5% v/v Triton X-100, 0.5 mM DTT) and rinsed

with lysis buffer. Target cleavage assays were performed at 25°C for 2.5 h in 8 µl

reactions containing 2.4 µl "40×" reaction mix, 55 nM recombinant protein, and 0.5 nM

^{32}P cap-radiolabeled target 1 (complementary to endogenous piRNA #1) or target 2

(complementary to endogenous piRNA #2), prepared by in vitro transcription as

described[111].

EfPiwi target cleavage assays

EfPiwi piRISC was purified as described[64]. Purified EfPiwi piRISC (100 nM, final concentration) was incubated at 37°C with a 5′-^{32}P-labeled target RNA complementary to g2-g21 (10 nM, final concentration) in reaction buffer composed of 20 mM Tris pH 8.0, 100 mM NaCl, 2 mM $MgCl_2$, 2 mM $MnCl_2$, 0.5 mM TCEP, and 0.1 mg/mL baker's yeast tRNA. Where indicated, purified *Ephydatia muelleri* GTSF1 (500 nM, final concentration) was also included. Target cleavage was stopped at indicated times by mixing aliquots of each reaction with an equal volume of denaturing gel loading buffer (98% w/v formamide, 0.025% xylene cyanol, 0.025% w/v bromophenol blue, 10 mM EDTA pH 8.0). Intact and cleaved target RNAs were resolved by denaturing PAGE (15%) and visualized by phosphorimaging. Quantification of signal was performed using ImageQuant TL (GE Healthcare).

Mouse germ cell purification

Germ cells from mouse testes were sorted and purified as described[16,47]. Briefly, freshly dissected mouse testes were decapsulated with 0.4 mg/ml collagenase type IV (Worthington) in 1× Gey's balanced salt solution (GBSS) at 33°C for 15 min. The separated seminiferous tubules were treated with 0.5 mg/ml trypsin and 1 µg/ml DNase I in 1× GBSS at 33°C for 15 min. Trypsin was then inactivated by adding 7.5% v/v fetal bovine serum (FBS). The cell suspension was filtered through a 70-µm cell strainer, and cells pelleted at 300 × *g* at 4°C for 10 min. Cell staining was performed at 33°C for 15 min with 5 µg/ml Hoechst 33342 prepared in 1× GBSS, 5% v/v FBS, and 1 µg/ml DNase I, then the cells were treated with 0.2 µg/ml propidium iodide, followed by final pass through a 40-µm cell strainer before sorting. Sorted cells were pelleted at 100 × *g* for 5 min, the buffer removed, and the cell pellets flash frozen and stored at −80°C. Cell lysates were prepared as described for HEK293T cells. Protein

concentration was estimated using the BCA assay, and an equal amount of total protein

from each cell type was used to assay for the ability to potentiate MIWI catalysis.

Analysis of RNA-seq data

Publicly available datasets[45,112–114] were analyzed. rRNA reads were removed using

Bowtie 2.2.5 with default parameters[115]. After rRNA removal, the remaining reads were

mapped to corresponding genomes (mouse, mm10; rat, rn6; macaque, rheMac8;

human, hg19) using STAR. 2.3 with default parameters that allowed ≤ 2 mismatches

and 100 mapping locations (ref. 116). Mapped results were generated in SAM format,

duplicates removed and translated to BAM format using SAMtools 1.8 (ref. 117).

HTSeq 0.9.1 with default parameters was used to count uniquely mapping reads[118]

(steady-state transcript abundance was reported in reads per kilobase per million

uniquely mapped reads [RPKM])

Methylarginine analysis

Recombinant MIWI was immunopurified and resolved by electrophoresis on a 4–20%

gradient SDS-polyacrylamide gel. The gel was fixed, stained with Coomassie G-250

(Simply Blue, Invitrogen), and the recombinant MIWI band excised and analyzed at the

UMASS Mass Spectrometry Core. Gel slices were chopped into ~1 mm^2 pieces, 1 ml

water added, followed by 20 µl 45 mM DTT in 250 mM ammonium bicarbonate.

Samples were incubated at 50°C for 30 min, cooled to room temperature, then 20 µl

100 mM iodoacetamide (IAA) was added and incubated for 30 min. The solution was

removed, and the gel pieces were three times washed with 1 ml water, 1 ml 50 mM

ammonium bicarbonate:acetonitrile (1:1), quenched with 200 µl acetonitrile, and dried in

a SpeedVac. Gel pieces were then rehydrated in 50 ul 50 mM ammonium bicarbonate

containing 4 ng/µl trypsin (Promega, Madison, WI) and 0.01% proteaseMAX (Promega)

and incubated at 37°C for 18 h. Supernatants were collected and extracted with 200 µl

an 80:20 solution of acetonitrile: 1% (v/v) formic acid in water. Supernatants were then

combined, and the peptides lyophilized in a SpeedVac and re-suspended in 25 µl 5%

acetonitrile, 0.1% (v/v) formic acid and subject to mass spectrometry analysis. Data

were acquired using a NanoAcquity UPLC (Waters Corporation, Milford, MA) coupled to

an Orbitrap Fusion Lumos Tribrid (Thermo Fisher Scientific, Waltham, MA) mass

spectrometer. Peptides were trapped and separated using an in-house 100 µm I.D.

fused-silica pre-column (Kasil frit) packed with 2 cm ProntoSil (Bischoff

Chromatography, DE) C18 AQ (200 Å, 5 µm) media and configured to an in-house

packed 75 µm I.D. fused-silica analytical column (gravity-pulled tip) packed with 25 cm

ProntoSil (Bischoff; 100 Å, 3 µm) media. Mobile phase A was 0.1 % (v/v) formic acid in

water; mobile phase B was 0.1 % (v/v) formic acid in acetonitrile. Following a 3.8 µl

sample injection, peptides were trapped at flow rate of 4 µl/min with 5% B for 4 min,

followed by gradient elution at a flow rate of 300 nl/min from 5–35% B in 90 min (total

run time, 120 min). Electrospray voltage was delivered by liquid junction electrode

(1.5 kV) located between the columns and the transfer capillary to the mass

spectrometer was maintained at 275°C. Mass spectra were acquired over *m/z* 300–

1750 Da with a resolution of 120,000 (*m/z* 200), maximum injection time of 50 ms, and

an AGC target of 400,000. Tandem mass spectra were acquired using data-dependent

acquisition (3 s cycle) with an isolation width of 1.6 Da, HCD collision energy of 30%,

resolution of 15,000 (*m/z* 200), maximum injection time of 22 ms, and an AGC target of

50,000.

Raw data were processed using Proteome Discoverer 2.1.1.21 (Thermo Fisher

Scientific), and the database search performed by Mascot 2.6.2 (Matrix Science,

London, UK) using the Swiss-Prot human database (download 04/09/2019). Search

parameters were: semi-tryptic digestion with up to two missed cleavages; precursor

mass tolerance 10 ppm; fragment mass tolerance 0.05 Da; peptide N-terminal

acetylation, cysteine carbamidomethylation, methionine oxidation, N-terminal glutamine

to pyroglutamate conversion, arginine methylation and arginine demethylation were

specified as variable modifications. Peptide and protein validation and annotation was

done in Scaffold 4.8.9 (Proteome Software, Portland, OR) employing Peptide Prophet[119]

and Protein Prophet[120]. Peptides were filtered at 1% FDR, while protein identification

threshold was set to greater than 99% probability and with a minimum of two identified

peptides per protein. Only arginine modification sites detected in all three replicates

from separate immunoprecipitation experiments are reported in the figure.

Statistics and reproducibility

Figs. 1b, 3, 4 and Extended Data Figs. 4a, 5c show mean ± SD for three independent

trials. Figs. 2a, b, Extended Data Figs. 1a–c and 1h (three independent trials) and

Figs. 2e, f, and Extended Data Figs. 1f–g, 2b, 3a, 3c (two independent trials) and

Extended Data Figs. 4b, 4c show representative data. The experiments in Figs. 1c, 5a,

and Extended Data Figs. 1d–e, 2, 3b, 3d–h, 5a, 7a–b were performed once. Protein

sequences were aligned using Clustal Omega; unrooted tree was constructed using

Randomized Axelerated Maximum Likelihood with default parameters[121] and visualized

in Interactive Tree Of Life[122].

Reporting summary

Further information on research design is available in the Nature Research Reporting

Summary linked to this paper.

Data availability

All data are available from the authors upon request.

102. Kawaoka, S. et al. The *Bombyx* ovary-derived cell line endogenously expresses

PIWI/PIWI-interacting RNA complexes. *RNA* **15**, 1258-1264 (2009).

103. Aoshima, K., Baba, A., Makino, Y. & Okada, Y. Establishment of alternative culture

method for spermatogonial stem cells using knockout serum replacement. *PLoS*

One **8**, e77715 (2013).

104. Kenny, N. J. et al. Tracing animal genomic evolution with the chromosomal-level

assembly of the freshwater sponge Ephydatia muelleri. *Nat Commun* **11**, 3676

(2020).

105. Flores-Jasso, C. F., Salomon, W. E. & Zamore, P. D. Rapid and specific

purification of Argonaute-small RNA complexes from crude cell lysates. *RNA* **19**,

271-279 (2013).

106. Pall, G. S. & Hamilton, A. J. Improved northern blot method for enhanced detection

of small RNA. *Nat Protoc* **3**, 1077-1084 (2008).

107. Wee, L., Flores-Jasso, C., Salomon, W. & Zamore, P. Argonaute Divides Its RNA

Guide Into Domains with Distinct Functions and RNA-Binding Properties. *Cell* **151**,

1055-1067 (2012).

108. Haley, B., Tang, G. & Zamore, P. D. In vitro analysis of RNA interference in

Drosophila melanogaster. *Methods* **30**, 330-336 (2003).

109. Carlson, K. D., Johnson, R. E., Prakash, L., Prakash, S. & Washington, M. T.

Human DNA polymerase kappa forms nonproductive complexes with matched

primer termini but not with mismatched primer termini. *Proc. Natl. Acad. Sci. U S A*

103, 15776-15781 (2006).

110. Kawaoka, S. et al. Zygotic amplification of secondary piRNAs during silkworm

embryogenesis. *RNA* **17**, 1401-1407 (2011).

111. Yoda, M. et al. ATP-dependent human RISC assembly pathways. *Nat. Struct. Mol.*

Biol. **17**, 17-23 (2010).

112. Merkin, J., Russell, C., Chen, P. & Burge, C. B. Evolutionary dynamics of gene and
992   isoform regulation in Mammalian tissues. *Science* **338**, 1593-1599 (2012).

993   113. Yu, T. et al. Long first exons and epigenetic marks distinguish conserved
994   pachytene piRNA clusters from other mammalian genes. *Nat Commun* **12**, 73
995   (2021).

996   114. Gainetdinov, I. et al. Terminal Modification, Sequence, and Length Determine
997   Small RNA Stability in Animals. *bioRxiv* 2020.09.08.287979 (2020).

998   115. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat*
999   *Methods* **9**, 357-359 (2012).

116. Dobin, A. et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-
21 (2013).

117. Li, H. et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078-2079 (2009).

118. Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-
throughput sequencing data. *Bioinformatics* **31**, 166-169 (2015).

119. Keller, A., Nesvizhskii, A. I., Kolker, E. & Aebersold, R. Empirical statistical model
to estimate the accuracy of peptide identifications made by MS/MS and database
search. *Anal. Chem.* **74**, 5383-5392 (2002).

120. Nesvizhskii, A. I., Keller, A., Kolker, E. & Aebersold, R. A statistical model for
identifying proteins by tandem mass spectrometry. *Anal. Chem.* **75**, 4646-4658
(2003).

121. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis
of large phylogenies. *Bioinformatics* **30**, 1312-1313 (2014).

122. Letunic, I. & Bork, P. Interactive tree of life (iTOL) v3: an online tool for the display

and annotation of phylogenetic and other trees. *Nucl. Acids Res.* **44**, W242-W245

(2016).

Acknowledgements We are grateful to Ken-Edwin Aryee (Greiner/Bremh Lab) for

providing pScalps_Puro, psPAX2, pMD2.G plasmids and for guidance on lentiviral

transduction; Pei-Hsuan Wu for help with mouse testis germ cell FACS and for

providing sorted cells; Katharine Cecchini for mouse tissues; Elaine Norowski

(Greiner/Bremh labs) for rat testes; Kyle Orwig (University of Pittsburgh) for rhesus

macaque testes; Paul Albosta and Cindy Tipping for guidance with *T. ni* and *D.*

melanogaster dissection; Leemor Joshua-Tor (Cold Spring Harbor Laboratory) for

sharing data prior to publication; Tiffanie Gardner for help formatting the manuscript; the

UMASS Flow Cytometry and Mass Spectrometry cores; and members of the Zamore

lab for discussions and comments on the manuscript. This work was supported in part

by NIGMS grants R37 GM062862 and R35 GM136275 (P.D.Z.) and R35 GM127090

(I.J.M) and JSPS KAKENHI grants 18H05271 (Y.T.) and 19K06484 (N.I.).

Author contributions A.A., P.D.Z., I.J.M., and Y.T. conceived and designed the

experiments. A.A., S.M.B., N.I., T.A.A., D.M.O., C.A., performed the experiments.

D.M.O. and I.G. provided and analyzed the sequencing data. P.D.Z. supervised the

research. A.A. and P.D.Z. wrote the manuscript. All the authors discussed the results

and approved the manuscript.

Competing interests The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at .

Correspondence and requests for materials should be addressed to P.D.Z.

 Figure legends

 Figure 1 | A component of mouse testis lysate potentiates piRNA-directed target

 RNA cleavage by MIWI.

 a, Strategy for programming recombinant MIWI with synthetic piRNA. **b,** Top**:**

 representative denaturing polyacrylamide gel image for target RNA cleavage by MIWI

 piRISC with and without added testis lysate. Bottom: product formed as a function of

 time by MIWI piRISC (mean ± SD, n =3). Rate constants were determined by fitting the

 data to the burst-and-steady-state equation (Equation 1 in Methods). $[E]_{active}$: the

 apparent concentration of active MIWI piRISC estimated from data fitting; k_{burst}: the pre-

 steady-state rate; k_{ss}: the steady-state rate. **c,** Target RNA cleavage by MIWI piRISC in

 the presence of lysate from either whole testis or FACS-purified germ cells. Spg,

 spermatogonia; PS, pachytene spermatocytes; DS, diplotene spermatocytes; SpII,

 secondary spermatocytes; Sptd, spermatids.

 Figure 2 | Testis protein GTSF1 potentiates target RNA cleavage by MIWI piRISC.

 a, Scheme to purify the MIWI-potentiating factor from mouse testis lysate. **b,** Target

 cleavage assay to estimate the apparent molecular weight of the MIWI-potentiating

 activity by size-exclusion chromatography (SEC). Arrowheads indicate the peak

 concentration of the molecular weight standards, and their peak elution volumes. V_0:

 void volume. **c,** The properties of the MIWI-potentiating activity compared to those of

 GTSF1. **d,** Strategy for creating knock-in mouse expressing 3XFLAG-HA-GTSF1

 (*Gtsf1$^{KI/KI}$*). Arrows represent sequencing primers used to validate the insertion. The

 exon encodes an incomplete Isoleucine codon, which is completed upon splicing. **e,**

 Western Blotting to show immuno-depletion of 3XFLAG-HA-GTSF1 from secondary

 spermatocyte lysate using anti-FLAG paramagnetic beads. I, Input; S, Supernatant; E,

 3XFLAG peptide eluate. **f,** Product formed by MIWI piRISC in the presence of the

indicated components. Abbreviations as in **e**. Numbers below the gel indicate relative

product formed in each condition.

Figure 3 | GTSF1 paralogs can distinguish between MIWI and MILI.

a, Representative denaturing polyacrylamide gel image showing that purified GTSF1

recapitulates the effect of testis lysate on MIWI catalysis. **b,** Product generated as a

function of time (mean ± SD, n = 3). Data were fit to the burst-and-steady-state

equation. Data for MIWI RISC alone are from Fig. 1. **c,** Representative denaturing

polyacrylamide gel images of the assay to test GTSF1 mutants and paralogs in target

cleavage by MIWI or MILI. **d,** Product formed as a function of time (mean ± SD, n = 3).

e, Observed pre-steady-state first order rate constants of target cleavage by MIWI

piRISC in the presence of either wild-type or PIWI-interacting mutant

GTSF1^{W98A,W107A,W112A} were plotted as a function of GTSF1 concentration. Dissociation

constant (K_D) and maximum potentiation rate (k_{pot}) were determined by fitting each

replicate individually to Equation 2 (see Methods). The curve corresponds to the mean

K_D and k_{pot} of the three replicates (n = 3). **f,** Representative target cleavage assay using

the indicated PIWI protein in the presence of different GTSF1 orthologs (100 nM).

Numbers below the gel lanes report fraction target cleaved (mean ± SD; n = 3). **g,**

Single-turnover rate of target cleavage by EfPiwi piRISC in the presence or absence of

EmGtsf1 (mean ± SD, n = 3).

Figure 4 | Cleavage by MIWI piRISC is sensitive to the extent of complementarity

between the piRNA and target.

a, MIWI piRISC target cleavage in the presence of GTSF1 for targets with increasing

complementarity to synthetic piRNA guide. Top: multiple-turnover conditions; bottom:

single-turnover conditions. All reactions contained saturating amounts of GTSF1. **b,**

Target cleavage assay using targets complementary to piRNA guide nucleotides g2–

g16 or g2–g30 and MIWI or MILI loaded with piRNAs of the indicated lengths, with or

without GTSF1 (mean ± SD, n = 3). **c,** Absolute (mean ± SD, n = 3) and relative pre-

steady-state and steady-state rates of cleavage of the g2–g30 target by MIWI loaded

with piRNAs of the indicated lengths in the presence of GTSF1.

Figure 5 | A model for the function of guide length and GTSF1 in target cleavage

by PIWI proteins.

a, The energy of base pairing, estimated by standard nearest-neighbor methods vs. pre-

and steady-state rates of GTSF1-potentiated target cleavage (mean ± SD, n = 3),

directed by piRNAs of different lengths loaded into MIWI. The same target, which was

fully complementary to each piRNA, was used in all experiments. **b,** piRNA-directed,

MIWI-catalyzed target cleavage is envisioned to require two sequential conformational

changes: (1) a target-dependent conformational change in piRISC (E_{PAZ}) in which the

piRNA 3′ end leaves the PAZ domain, allowing extensive base pairing of the piRNA with

the target substrate (S); and (2) GTSF1-dependent conversion of this piRISC pre-

catalytic state (E_C) to the fully competent catalytic state ($E_{C'}$). P: cleaved target products.

c, Proposed effects of different piRNA lengths, extents of guide:target complementarity,

and GTSF1 on the forward and reverse rates of the two conformational rearrangements.

The wide, central cleft like that observed in the cryo-EM structure of *Ephydatia fluviatilis*

Piwi-A[64] is envisioned to allow the central region of a 30 nt piRNA to be mobile and

exposed to solvent when its 3′ end is secured to the PAZ domain (upper right).

Extended Data Table 1. Pairwise comparison of percent identity or similarity for domain sequence of GTSF1-related proteins.

Zinc finger 1

		Mm GTSF1	Mm GTSF1L	Mm GTSF2	Rn GTSF1	Rn GTSF1L	Rn GTSF2	Mmul GTSF1	Mmul GTSF1L	Tn TnGtsf1	Tn TnGtsf1l	Bm BmGtsf1	Bm BmGtsf1l	Dm Asterix
Mm	GTSF1	100	50	54.2	100	50	54.2	100	54.2	45.8	54.2	45.8	33.3	37.5
Mm	GTSF1L	62.5	100	73.9	50	100	73.9	50	95.7	41.7	52.2	37.5	56.5	39.1
Mm	GTSF2	66.7	87	100	54.2	73.9	100	54.2	78.3	33.3	43.5	33.3	56.5	34.8
Rn	GTSF1	100	62.5	66.7	100	50	54.2	100	54.2	45.8	54.2	45.8	33.3	37.5
Rn	GTSF1L	62.5	100	87	62.5	100	73.9	50	95.7	41.7	52.2	37.5	56.5	39.1
Rn	GTSF2	66.7	87	100	66.7	87	100	54.2	78.3	33.3	43.5	33.3	56.5	34.8
Mmul	GTSF1	100	62.5	66.7	100	62.5	66.7	100	54.2	45.8	54.2	45.8	33.3	37.5
Mmul	GTSF1L	66.7	95.7	91.3	66.7	95.7	91.3	66.7	100	37.5	47.8	37.5	52.2	34.8
Tn	Gtsf1	54.2	54.2	54.2	54.2	54.2	54.2	54.2	50	100	41.7	83.3	45.8	37.5
Tn	Gtsf1-like	66.7	65.2	52.2	66.7	65.2	52.2	66.7	60.9	62.5	100	41.7	43.5	39.1
Bm	Gtsf1	58.3	45.8	54.2	58.3	45.8	54.2	58.3	50	87.5	54.2	100	37.5	37.5
Bm	Gtsf1-like	54.2	65.2	65.2	54.2	65.2	65.2	54.2	60.9	66.7	73.9	58.3	100	52.2
Dm	Asterix	58.3	56.5	56.5	58.3	56.5	56.5	58.3	52.2	66.7	69.6	62.5	73.9	100

Zinc finger 2

		Mm GTSF1	Mm GTSF1L	Mm GTSF2	Rn GTSF1	Rn GTSF1L	Rn GTSF2	Mmul GTSF1	Mmul GTSF1L	Tn TnGtsf1	Tn TnGtsf1l	Bm BmGtsf1	Bm BmGtsf1l	Dm Asterix
Mm	GTSF1	100	39.1	39.1	100	39.1	39.1	100	39.1	43.5	43.5	39.1	30.4	39.1
Mm	GTSF1L	65.2	100	78.3	39.1	100	78.3	39.1	82.6	39.1	30.4	34.8	30.4	26.1
Mm	GTSF2	56.5	91.3	100	39.1	78.3	100	39.1	82.6	34.8	39.1	34.8	39.1	26.1
Rn	GTSF1	100	65.2	56.5	100	39.1	39.1	100	39.1	43.5	43.5	39.1	30.4	39.1
Rn	GTSF1L	65.2	100	91.3	65.2	100	78.3	39.1	82.6	39.1	30.4	34.8	30.4	26.1
Rn	GTSF2	56.5	91.3	100	56.5	91.3	100	39.1	82.6	34.8	39.1	34.8	39.1	26.1
Mmul	GTSF1	100	65.2	56.5	100	65.2	56.5	100	39.1	43.5	43.5	39.1	30.4	39.1
Mmul	GTSF1L	52.2	87	87	52.2	87	87	52.2	100	34.8	39.1	39.1	30.4	26.1
Tn	Gtsf1	56.5	52.2	52.2	56.5	52.2	52.2	56.5	52.2	100	43.5	82.6	34.8	26.1
Tn	Gtsf1-like	52.2	47.8	56.5	52.2	47.8	56.5	52.2	52.2	52.2	100	43.5	60.9	30.4
Bm	Gtsf1	52.2	47.8	52.2	52.2	47.8	52.2	52.2	56.5	91.3	52.2	100	39.1	26.1
Bm	Gtsf1-like	60.9	47.8	47.8	60.9	47.8	47.8	60.9	39.1	65.2	82.6	60.9	100	26.1
Dm	Asterix	56.5	60.9	56.5	56.5	60.9	56.5	56.5	60.9	60.9	65.2	56.5	47.8	100

Carboxy terminal domain

		Mm GTSF1	Mm GTSF1L	Mm GTSF2	Rn GTSF1	Rn GTSF1L	Rn GTSF2	Mmul GTSF1	Mmul GTSF1L	Tn TnGtsf1	Tn TnGtsf1l	Bm BmGtsf1	Bm BmGtsf1l	Dm Asterix
Mm	GTSF1	100	7.3	8.3	100	7.3	10.4	88.5	7.3	7.3	6.2	8.3	6.2	8
Mm	GTSF1L	20.8	100	32.6	7.3	93.2	31.5	9.4	60	4.3	4.4	5.4	4.4	5.5
Mm	GTSF2	28.1	54.3	100	8.3	32.6	89	9.4	34.8	2.2	4.4	4.3	5.5	10.1
Rn	GTSF1	100	20.8	28.1	100	7.3	10.4	88.5	7.3	7.3	6.2	8.3	6.2	8
Rn	GTSF1L	20.8	97.7	54.3	20.8	100	31.5	9.4	58.9	3.2	4.4	4.3	4.4	5.5
Rn	GTSF2	28.1	53.3	93.4	28.1	53.3	100	11.5	33.7	2.2	4.4	4.3	5.5	9.2
Mmul	GTSF1	92.7	19.8	28.1	92.7	19.8	28.1	100	8.3	8.3	8.3	9.4	7.3	7.1
Mmul	GTSF1L	24	66.7	53.3	24	68.9	51.1	24	100	6.5	1.1	9.7	2.2	4.6
Tn	Gtsf1	20.8	20.4	17.4	20.8	20.4	16.3	16.7	19.4	100	6.2	60	5	8.3
Tn	Gtsf1-like	11.5	8.9	8.8	11.5	7.8	8.8	12.5	6.7	7.5	100	5	44.4	4.6
Bm	Gtsf1	27.1	17.2	21.7	27.1	19.4	20.7	22.9	22.6	71.2	11.2	100	7.5	8.3
Bm	Gtsf1-like	11.5	12.2	14.3	11.5	11.1	13.2	12.5	7.8	6.2	60	10	100	1.8
Dm	Asterix	17.9	15.6	18.3	17.9	14.7	17.4	18.8	14.7	18.3	10.1	17.4	11	100

Extended Data Table 2.

		GTSF1	GTSF1 PIWI-interaction mutant	GTSF1 RNA-binding mutant	GTSF1L	GTSF2	Buffer
MIWI	k_{burst} (min⁻¹)	1.1 ± 0.1	0.67 ± 0.08	< LQ	1.1 ± 0.1	1.1 ± 0.1	< LQ
	k_{rel}	1	0.6	—	1	1	—
MILI	k_{burst} (min⁻¹)	0.17 ± 0.03	0.04 ± 0.01	< LQ	0.02 ± 0.02	< LQ	< LQ
	k_{rel}	1	0.2	—	0.06	—	—

Extended Data Figures and Legends

Extended Data Figure 1 | Recombinant MIWI programmed with guide piRNA cleaves complementary target RNA.

a, Purification of apo-MIWI and of MIWI loaded with a synthetic piRNA guide (piRISC). **b,** Coomassie-stained SDS-PAGE of purified apo-MIWI and MIWI piRISC. E1, E2: first and second eluates from sequential incubation of the paramagnetic beads with 3XFLAG peptide. E1 was used for the cleavage assay. **c,** Representative denaturing polyacrylamide gel of target cleavage assay using apo-MIWI or MIWI piRISC. A digital overexposure is shown below the gel. **d,** MIWI piRISC loaded with synthetic guides differing at their 5′ or 3′ termini was assayed for the ability to cleave a complementary target RNA. **e,** Northern blot to estimate the yield of purified MIWI piRISC. **f,** Silver stained gel showing purified MIWI piRISC with and without an amino-terminal epitope tag. **g,** Target cleavage assay comparing the activity of piRISC with and without an amino-terminal epitope tag. Data were fit to Equation 1 (see Methods); error of fit (SD) is reported. **h,** Mass spectrometry was used to determine arginine methylation status for recombinant, affinity-purified MIWI (n = 3); arginine methylation status of endogenous MIWI from mouse testis is provided for comparison[77]. Arginine position reports the amino acid number in endogenous mouse MIWI and in parentheses, the corresponding residue in the epitope-tagged recombinant protein. Only modified arginine residues detected in all three independent preparations are reported. N.D., not detected.

Extended Data Figure 2 | The target-cleavage potentiating component in testis lysate component transiently interacts with MIWI.

a, Immobilized, unloaded apo-MIWI was preincubated with testis lysate, purified as depicted, and then assayed for target cleavage activity. **b,** Immobilized MIWI piRISC was preincubated with testis lysate. purified as depicted, and then assayed for target cleavage activity. A digital overexposure is shown below the gel.

a

Pre-incubated with buffer

Time: 0 0.5 1 2 3 4 5 15 30 60 90 120

Substrate—

Product—

Pre-incubated with testis lysate

Time: 0 0.5 1 2 3 4 5 15 30 60 90 120 min

Substrate—

Product—

b

Affinity-purified MIWI piRISC pre-incubated with testis lysate then washed:

No wash | Low salt (120 mM KOAc) | High salt (2 M KOAc) | Affinity-purified piRISC alone

Time: 0 1 2 3 4 5 15 30 60 120 | 1 2 3 4 5 15 30 60 120 | 1 2 3 4 5 15 30 60 120 | 0 1 2 3 4 5 15 30 60 120 min

Substrate—

Product—

Product—

Extended Data Figure 3 | Biochemical properties of the MIWI potentiating component in testis lysate.

a, The MIWI-potentiating factor is specific to testis lysate. **b,** Testis lysate does not enhance target cleavage by purified mouse AGO2. **c,** The MIWI-potentiating factor is sensitive to alkylation by *N*-ethylmaleimide. **d,** Effect of divalent cation chelators on MIWI-potentiating activity. For 0 nM chelator, water was added for EDTA and EGTA and ethanol for 1,10-phenanthroline. **e,** Strategy to test metal rescue after incubation of testis lysate with divalent cation chelators. **f, g,** Metal rescue experiments were performed as in (**e**) using EDTA (**f**) or 1,10-phenanthroline (**g**). **h,** Fractionation of testis lysate using Zn^{2+} and Ni^{2+} immobilized metal affinity chromatography (IMAC). In **c** and **f-h**, a digital overexposure is shown below each gel. Numbers below the gel indicate relative product produced in each lane.

Extended Data Figure 4 | *Gtsf1, Gtsf1l*, and *Gtsf2* **mRNA abundance and recombinant protein expression.**

a, For each germ cell type, mRNA abundance (from previously published data; see methods) is reported as mean ± SD (*n* = 3) in molecules/cell. Spg: spermatogonia; SpI: primary spermatocytes; SpII: secondary spermatocytes; RS: round spermatids. **b,** Coomassie-stained SDS-PAGE of purified, recombinant wild-type GTSF1, GTSF1 mutants, and wild-type GTSF1Like and GTSF2. **c,** Oriole protein stain detection of purified, recombinant, C-terminal V5SBP-tagged GTSF1 orthologs. Mm: *Mus musculus* (mouse); Bm: *Bombyx mori* (silkmoth). Coomassie protein stain detection of purified, recombinant *Ephydatia muelleri* Gtsf1. **d**, Clustal Omega amino acid sequence alignment of the three mouse GTSF1 paralogs.

Extended Data Figure 5 | GTSF proteins are PIWI selective.

a, MIWI piRISC target cleavage assay in the presence of lysates from different animals. *T.ni: Trichoplusia ni.* Macaque: rhesus macaque. A digital overexposure is shown below the gel. **b,** Representative target cleavage assay using the indicated PIWI protein in the presence of different GTSF1 orthologs (100 nM). Numbers below the gel lanes report fraction target cleaved (mean ± SD; *n* = 3). **c,** Left: Percent identity of different GTSF1 orthologs Right: Unrooted tree of GTSF1 orthologs.

a

Affinity-purified MIWI piRISC plus lysate from:

b

Affinity-purified PIWI piRISC plus:

c

		% Identity (GTSF1)	
🪰	*Drosophila melanogaster* (Dm)	100.00	19.86
🦋	*Bombyx mori* (Bm)	20.00	22.38
🦋	*Trichoplusia ni* (Tn)	19.40	21.17
🐁	*Mus musculus* (Mm)	19.86	100.00
🐀	*Rattus norvegicus*	19.86	100.00
🐒	*Macaca mulatta* (Mmul)	20.55	91.62
🧍	*Homo sapiens* (Hs)	20.55	91.62
🦫	*Ephydatia muelleri* (Em)	14.97	21.47

Extended Data Figure 6 | Expression of GTSF1 paralogs in mouse, rat, rhesus macaque, and human whole testes.

Gtsf genes are syntenic in mice, rats, macaques and humans. Primates, including humans, do not express GTSF2. RNA-seq from total testis RNA of the indicated animals sourced from previously published datasets (see methods).

Extended Data Figure 7 | Complementarity and GTSF1 requirements for target cleavage by AGO2, MIWI, and MILI.

a, MIWI target cleavage ± GTSF1 using synthetic piRNA guide 1. **b,** Target cleavage by AGO2 siRISC compared with MIWI or MILI piRISC loaded with synthetic piRNA guide 1 or 2, for targets with complete (g2–g30) or partial (g2–g16) complementarity to the piRNA.

a

g2–g21   g2–g16

rGTSF1:

Time:

Substrate—

Product—

Affinity-purified MIWI piRISC

b

g2–g30   g2–g16

AGO2:
MILI:
MIWI:

Buffer AGO2 MIWI MILI

Substrate

Product

GTSF1:   − + +   − + +   − + +

Extended Data Figure 8 | Model predicting free-energy change during GTSF1-induced potentiation of target cleavage by PIWI.

Two conformational changes in piRISC (E) are proposed to be required for efficient target cleavage catalyzed by PIWI proteins. E_{PAZ}: piRISC with the 3′ end of the piRNA bound to the PAZ domain; E_C: piRISC with the piRNA fully paired to the target RNA, i.e., the pre-catalytic conformation; $E_{C'}$: E_C in the catalytically competent conformation; $S^{‡}$: transition-state. For piRNAs of biologically relevant length, extensive complementarity is proposed to promote the first conformational change; GTSF1 is proposed to promote the second. Blue: energy barrier to catalysis without GTSF1, i.e., the spontaneous conversion of E_C to $E_{C'}$.

Supplementary Table 1. Oligonucleotides and DNA used in this study

Name	Sequence (5′-to-3′)	Notes

AGO/PIWI RNA guides (m, 2′-O-methyl; phos, 5′ phosphate)

Name	Sequence (5′-to-3′)	Notes
p30.47	/phos/UGA GGU AGU AGG UUG UAU AGU AUC CAG AGG	First 21 nucleotides correspond to *let-7* miRNA sequence; 3′,2′ OH; orange label in figures
p30.50	/phos/UGA GGU AGU AGG UUG UAU AGU AUC CAG AGmG	
p30.50DNA	/phos/TGA GGT AGT AGG TTG TAT AGT ATC CAG AGmG	DNA guide
p21.46	/phos/UGA GGU AGU AGG UUG UAU AGU	
p30.53	/phos/UUA GGU AAC CCA GUA GAU CCA GAG GAA UUmC	Guide sequence targeting luciferase in pGL2 plasmid; red in figures
p26.53	/phos/UUA GGU AAC CCA GUA GAU CCA GAG GmA	
p26.53 (3'OH)	/phos/UUA GGU AAC CCA GUA GAU CCA GAG GA	
p21.1175	/phos/UUA GGU AAC CCA GUA GAU CCA	
p16.53	/phos/UUA GGU AAC CCA GUA G	
FLAG-Siwi/MIWI/MILI pre-piRNA	/phos/UCG AAG UAU UCC GCG UAC GUU AUG CUA GCU GAU CCU GAC CGC UGA GUC GU	

AGO passenger strand

Name	Sequence (5′-to-3′)	Notes
p21.443	/phos/UAU ACA ACC UAC UAC CUC AUU	PS for 21.46
p21.1176	/phos/UGG AUC UAC UGG GUU ACC UUC	PS for 21.1175

Target RNA oligos

Name	Sequence (5′-to-3′)	Notes
5′ CH₃O/Tar3	/5′ CH₃O/UAU AGA AAA AAU AAC UAU ACA ACC UAC UAC CUC AAU GCU	2-21 matched target against let-7 type guide sequences
5′ CH₃O/Tar4	/5′ CH₃O/UAU GAA GAG AUC GUA ACC ACA ACC UAC UAC CUC AAU GCU ACA	2-16 matched target against let-7 type guide sequences

Primers for IVT targets

Name	Sequence (5′-to-3′)	Notes
Pr8F_T7_pGl2	GCG TAA TAC GAC TCA CTA TAG GGT TTT AAT GAA TAC GAT TT	Forward Primer to add T7 promoter to template
Pr9R	CCC ATT TAG GTG ACA CTA TAG ATT TAT ACC TAG TTA AAC AGC GGA ACT GTG TAT AAA AGG TTG AGG TAG TAG GTT GTA TAG TAT CCA GAG GAA TTC ATT ATC AGT G	Reverse primer for fully complementary let-7 target
Pr_let7_(2-16)R	GCA CTG TTT TCA TGA AGA CAT GAA GAG ATC TGT AGC ATT GAG GTA GTA GGT TGT GGT TAC GAT CTC TTC ATG AGG AAT TCA TTA TCA GTG	Reverse primer for 2-16 let-7 target
Pr_Luc_Tar1R	TCC GGA ATG ATT TGA TTG CC	Reverse primer for fully complementary luciferase target

Pr_Luc_tar8R	GCG GAA GGG CCA CAC CCT TAG GTA ACC CAG TAG ATC CAC CTC TGG ATA TTA TCA GTG	Reverse primer for 2-21 match luciferase target
Pr_Luc_tar7R	TAT GCG GAA GGG CCA CAC CCT TAG GTA ACC CAG TAG ATC CTC CTC TGG ATA TTA TCA GTG	Reverse primer for 2-20 match luciferase target
Pr_Luc_Tar6R	TTC TAT GCG GAA GGG CCA CAC CCT TAG GTA ACC CAG TAG ATC TTC CTC TGG ATA TTA TCA GTG	Reverse primer for 2-19 match luciferase target
Pr_Luc_Tar5R	GCG GAA GGG CCA CAC CCT TAG GTA ACC CAG TAG ATA TTC CTC TGG ATA TTA TCA GTG	Reverse primer for 2-18 match luciferase target
Pr_Luc_Tar4R	TTC TAT GCG GAA GGG CCA CAC CCT TAG GTA ACC CAG TAG AAA TTC CTC TGG ATA TTA TCA GTG	Reverse primer for 2-17 match luciferase target
Pr_Luc_tar3R	TAT GCG GAA GGG CCA CAC CCT TAG GTA ACC CAG TAG GAA TTC CTC TGG ATA TTA TCA GTG	Reverse primer for 2-16 match luciferase target

Sequence-affinity purification (m, 2′-O-methyl)

luciferase capture oligo	/Biotin/mAmCmU mAmCmU mGmCmA mGmCmA mCmAmA mCmCmC mUmAmC mCmAmA mAmUmU mAmCmC mUmAmA mCmCmUmG mC	
luciferase competitor oligo	GCA GTT AGG TAA TTT GGT AGG GTT GTG CTG CAG TAG T	DNA oligo
let-7 capture oligo	/Biotin/mAmUmA mGmAmC mUmGmC mGmAmC mAmAmU mAmGmC mCmUmA mCmCmU mCmCmG mAmAmC mG	
let-7 competitor oligo	CGT TCG GAG GTA GGC TAT TGT CGC AGT CTA T	DNA oligo

Primers for BmAgo3 targets

Target-F	GCG TAA TAC GAC TCA CTA TAG TCA CAT CTC ATC TAC CTC C	Forward Primer to add T7 promoter to the template
Target #1 (pi848-perfect target-R)	AGG TGA CAC TAT AGA TTT ACA TCG CGT TGA GTG TAG AAC GGT TGT ATA AAA GGT CGA GTG ACA TGT TTG GAA GCA ACA GCA TCC AGA GGA GTT CAT G	Reverse primer for fully complementary target
Target #2 (pi2148ds-perfect target-R)	AGG TGA CAC TAT AGA TTT ACA TCG CGT TGA GTG TAG AAC GGT TGT ATA AAA GGT AGG GCC AAG ACC AAA TAT AGT ACG TTT CAT CCA GAG GAG TTC ATG	Reverse primer for fully complementary target

Construction of *Gtsf1*^{KI/KI} mouse

| HDR donor | ACTTTGTCCAATTCACCGACAGCCTGTTATTGGGGGAAAGGAATCAA
ATCACCTGTTAACAATTTGATGGGGAAGATAAGAGACAGCTGAATGA
TGGTGTTATGAGTTCCAGCGTCCTTGGCCTACTTGGAATCAGTCTGT
TTGCTAAGAAAGAATAGGTCAGCTCTCTAAGTCGCTGAATATCTTTA
CTCATCTTTTTCAAAATAATAATAATAATGTAATTCTTGGTTTTTCCTT
AAAACCTTTTTTTATTTTTAACAAAGTATGACATAACTGTAGTAGAAGT
TCCATACGGGTTAAGATAGTTTAACTAATTCTCAGATAGTTTCCCCAA
TGATTGCCATTAACAAATGGGCTCTGGAGCAAAGTCCGAAACTAGAG
TTTAGTTGATCACCCCTATTGCTTAACAAAGGTTCTAATGCTGCCTAT
GTCATAGAATGGGGGGGAAGGGGGTGTGGAATTTTTGTGAGAGGC
CAAAGGAGAGCTTACCTAGAGACAACAGCTGTATGTGCATCCAATTT
TCAGAAGATTCCTATGGGAGAGGTTATATACAAAATTAGTTTTCCCAT
AATACGTGAACTAAATCAGTGTTCAATAGAGATTCCAGTAAAAGAGA
ATTCAAACTCAGAGACATTTCCTCTTTTAATTTTTTTTCATGTTTGTTAT
ATTTGTTCTCTTAGATGTTCTCTATGAGCTCTTAATTTTTTTAACATTC
ACATATATTCCCTCAGTGCAGTTAAGTAATCAGTACTGACTAAAACAG
TATTTATCTTGTTATTATATAGAAATGAGAGGAACAAAATGAAAAATG
GCTAGAAAGAGAGAATATTGGTGTGACGTTTTCTGATCATGATTAAG
AAAAGTTACTAAGCTGGGTATGGTTGAAAATGCCTTTAATCCCCAGAA
CTCAGGAGGCAGAGGCAGGTGGATCTCTGTGTGTTCAAGGTCAGTC
TGGTCTGCATATTGAGTTCCAGGACAGCCAAAACTATACAGAGAAAC
CTTGTCTCTTAAAAAAGAAGAAAAAGAAAGGTCATTAAATGAGATGTA
GCTTTTATTTTACATGTGTTTGTGTGCATTTTGCCCACAGAAGCCAGA
AGAGGGCATCTGATCCCCTGAAACTGGAGTTATAGGCAGTTGTAAG
CCATGTGGGTGCTGAGAATCAAACCCAGATGTCACTAAGAACACCC
AGAGCTCTTAACCATTTAATCATCTTTCCATCTGGAAAGAAGATCTTT
AAAGTAGTTTATCCAGAAGTCTAAATACTGGTATTTTTATGACCTTGA
CTTGGCCCCCATTACTTATGTTAAATACCATAACCCCTAATAGCATGT
AAACAGTGGCAGGTGTTGTAATACCACTTAAAATAAATTGTTATTATT
GTCACCTGAATGTAGTTTGTCAGTTGTCATATTTGACCTAGCAATTTA
TTCCAAGAATGTGAACTATGCAGTGATGCTAAACTTAGTCATAGGAG
TAAAGAATGCCTTTAAGAAGAAACTGGAGAGAAGGTTGTATGCTAAG
GGTTTATTTTATTGAACTTGGGGCAGCTTCAGGTCCAGGAATGACTC
CACATCTTAGGGCTACTATTTGATGCGTTTCATTTCTGATTCTCTGAA
CCATGCTTTTTCCTCTAGCACTTGAACAGGACTCCTTTGCCATCAGC
<u>ATG</u>GACTACAAGGACCACGACGGTGACTACAAGGACCACGACATCG
ACTACAAGGACGACGACGACAAGGGCTCCGGATACCCATACGATGT
GCCAGATTACGCTGGATCAGGCGAAGACACTTACAGTATGTACTACT
TGCTGCTTTTAGAAAATAATACAATATTTATGTGTAGTTTGATGCCTG
ACTAGACTGTATGTTGTGAAAACTTCAGATTCCTCCCTGAAATGCTG
AGAAAGAAAGGTCTGTGTGATATTGAGGTATACAGATGCTCAAATTT
TAAATAAGTAACTTTAGTCTACAGCCATATTACCCTGAAAAAAAATGG
TCTGAAACAAGCACTTTAAACTTAAGTAATTAGACATATGAGAACCAA
CTGTAATGATACACAACTGTAATCCCAACTCATAGGAAGCTGAATCA
TGGGGATCACAACTTCTAGGCCATCATGGGCTTCACAGAAAGACCA
TCAAAAGATAAACCAGTAGTCCACTGAGATGGCTCAAGGGTGAAGT
GCTTATTACCATAAAAGCCTGACAACCTTAGTTTGAATTTTAGAACTC
TTGTAAAAGTAGGAAAGAAAACTGAATCTATAAAGTCGTCCTCTGAC
CTACACACAGACCACACACAATAATAAGTACAAAATTTTTACAATACT
GGGGATGTAACTCAGTGGTAGAGCATTTGTGTAGCATTCTCAAGACC
CATATTTTAATACCAAGATAATAGGTAGAGAGAGAGAGAGAGAGATATG
AATGATTGATTGGTGGTTAAATTGGTGACCAAGATAGATGGATGAAT
GGATGATTGATAAGTAGATGGATGGGTGGATGGATGTAGATAAATAG
ATGGATCATTGATAGATAGATAGATAGACAGATTCTTTTCAAGGTAGA
ACAGTAAGTTAGAGCTGACCTAAATTGCCAGACTAAGTAAAGCATTT
ATGGTAGTTCTCATTATCAACTTTGACACATGCTCTATATGAGCAGTT
ACAATATGTGTGAATCATGGGGTACCTCTGTTCTGTCAACTGCTGAC
ATTTGCTAGTGATAGAATGGATTAAATGAAGGAAAACTAATAAGAATT
TCATCTGTGACTGCTCCCTGGGTGTTCTGAAGCTTCAGAAATAAGAA
GAGATTGCTGGCTAAGAGCTCAGTGGCACAGTAGCATGCCTGTCCT
CTGGGGCCTTGCCTCCCTGTTCATTCTGTCCTTCAGATCACCTTTCA
AGCTGATTTGCATCATGACCGTGGTTAGCAATTTATTCCATGTTTGC
CACTCTTCCAGCGAAACTGTCAGTTTCAATTCCCCTCTGGTTTCTGA
TTTTCATTGCAGTCTCCTGGTTTATAAGCTTCTTATCAATGGTGAATT
GTTTGTCTTGCTTTTATCCTTACCTTCTGGGGTGGCAAGAGTGTTTTA
TTAGGTGCCTTTTGCTCTCAGAATGCACTATTTAACTTTCTTTCCTTA
ACCAGGCTTGTTAAGATTTGGGGCATAGCCTGAGGGCAGCTTTGAA
TAACATGGTATAGGTTCTGGGTTTTCTTGACACAGCAGTCTGTGTCT
ACAATTGTCTTAGCTGTAGGAGGTTATTTTTTATTTTAGCTGGCCAGT
TTGGTAGTCCTTAGATTAAGGAACTTTTTTTCTCATCTATGTTCCAAAA
CATGAAAAGTAAAATCACCTACCTGACAAAAGTAATGAGATCATTGT
CAGGGGAACGAGAGCAGCCAATACAGCATACAGGAAGGAGTAAAG
CAAAGAACAAATTGGGGGGAAACTCTAGTTATTAATACTTCATAGAA
CATTCAATTGAAGCTATCAACTTCTTAACTAGAAATAGACAAGGATGC
TAATGAGATTATTCTCAACTACGGCTAGCTAGATTGAGAGAGTGTTG
GAAAGTGCAGTGGGC | Left and right homology arms;
<u>ATG</u>: *Gtsf1* start codon;
3XFLAG-HA knock-in cassette |

sgRNA target sequence	GTCTTCCATGCTGATGGCAA<u>AGG</u>	<u>PAM</u> sequence
Primer F1	CTGTTTTACTGAGACCTGCAACTTT	Genotyping and validation primers
Primer R1	GATCCAGCGTAATCTGGCACAT	
Primer F2	ACTACAAGGACCACGACGGTG	
Primer R2	GACTGGATTGCTGGTGTAACAATA	

Arif et al., Supplementary Data Fig. 1

a

b

c

d