Local rather than global H3K27me3 dynamics associates with differential gene expression in *Verticillium dahliae*

H. Martin Kramer[1], Michael F. Seidl[1,2], Bart P.H.J. Thomma[1,3,#,*], David E. Cook[1,4,*]

[1]Laboratory of Phytopathology, Wageningen University and Research, Droevendaalsesteeg 1, 6708 PB Wageningen, the Netherlands

[2]Theoretical Biology & Bioinformatics Group, Department of Biology, Utrecht University, Utrecht, The Netherlands

[3]University of Cologne, Institute for Plant Sciences, Cluster of Excellence on Plant Sciences (CEPLAS), 50674 Cologne, Germany

[4]Department of Plant Pathology, Kansas State University, 1712 Claflin Road, Manhattan, Kansas 66506, USA

[*]These authors contributed equally

[#]For correspondence: bthomma@uni-koeln.de

Running title: Global histone dynamics in *Verticillium dahliae*

Word count Abstract: 237

Word count Importance: 150

Word count text: 6.466

ABSTRACT

Differential growth conditions typically trigger global transcriptional responses in

filamentous fungi. Such fungal responses to environmental cues involve epigenetic

regulation, including chemical histone modifications. It has been proposed that

conditionally expressed genes, such as those that encode secondary metabolites but also

effectors in pathogenic species, are often associated with a specific histone modification,

lysine27 methylation of H3 (H3K27me3). However, thus far no analyses on the global

H3K27me3 profiles have been reported under differential growth conditions in order to

assess if H3K27me3 dynamics governs differential transcriptional. Using ChIP- and RNA-

sequencing data from the plant pathogenic fungus *Verticillium dahliae* grown on three *in*

vitro cultivation media, we now show that a substantial number of the identified

H3K27me3 domains globally display stable profiles among these growth conditions.

However, we do observe local quantitative differences in H3K27me3 ChIP-seq signal that

associate with differentially transcribed genes between media. Comparing the *in vitro*

results to expression during plant infection suggests that *in planta* induced genes require

chromatin remodelling to achieve expression. Overall, our results demonstrate that

although some loci display H3K27me3 dynamics that can contribute to transcriptional

variation, other loci do not show such dependence. Thus, we conclude that while

H3K27me3 is required for transcriptional repression, it is not a conditionally responsive

global regulator of differential transcription. We propose that the H3K27me3 domains

that do not undergo dynamic methylation may contribute to transcription through other

mechanisms or may serve additional genomic regulatory functions.

IMPORTANCE

In many organisms, including filamentous fungi, epigenetic mechanisms that involve

chemical and physical modifications of DNA without changing the genetic sequence have

been implicated in transcriptional responses upon developmental or environmental cues.

In fungi, facultative heterochromatin that can de-condense to allow transcription in

response to developmental changes or to environmental stimuli is characterized by tri-

methylation of lysine 27 on histone H3 (H3K27me3), and H3K27me3 has been implicated

in transcriptional regulation although precise mechanisms and functions remain

enigmatic. Based on ChIP- and RNA-sequencing data, we show for the soil-borne broad

host range vascular wilt plant pathogenic fungus *Verticillium dahliae* that although some

loci display H3K27me3 dynamics that can contribute to transcriptional variation, other

loci do not show such dependence*.* Thus, although we recognize that H3K27me3 is

required for transcriptional repression, we also conclude that this mark is not a

conditionally responsive global regulator of differential transcription upon responses to

environmental cues.

INTRODUCTION

The fungal kingdom comprises a plethora of species occupying an enormously diverse

range of ecological niches (Schmit & Mueller, 2007). As environments are typically

dynamic, including the effects of daily and yearly cycles, fungi continuously need to

respond and adapt to survive (Gibson, 2008; Selvig & Alspaugh, 2011). To this end,

fungi have evolved various mechanisms to monitor their environment and to

transcriptionally respond to environmental cues (Bahn *et al.*, 2007). For instance, the

yeast *Saccharomyces cerevisiae* senses cold stress by increased membrane rigidity,

which leads to transcription of genes that, among others, encode cell damage-preventing

proteins (Aguilera *et al.*, 2007). Furthermore, *S. cerevisiae* senses the quantitative

availability of carbon and nitrogen sources in the environment to determine which

developmental program maximises the potential for survival (Schneper *et al.*, 2004). In

animal systems, epigenetic mechanisms (i.e. those affecting nuclear processes by

modifying DNA characteristics without changing the genetic sequence) are implicated in

transcriptional responses to changing environments (Heijmans *et al.*, 2008; Zannas *et

al.*, 2015; Martin & Fry, 2018). Such epigenetic mechanisms have similarly been

proposed to contribute to environmental response in filamentous fungi. For instance, the

saprotrophic fungus *Neurospora crassa* phenotypically reacts to environmental stimuli,

such as changes in temperature and pH, yet *N. crassa* mutants impaired in transcription-

associated epigenetic mechanisms display reduced growth in response to these stimuli

(Kronholm *et al.*, 2016). Similarly, the nectar-feeding yeast *Metschnikowia reukaufii* and

the ubiquitous fungus *Aureobasidium pullulans* fail to properly respond to changing

carbon sources when DNA methylation or histone acetylation are inhibited (Slepecky &

Starmer, 2009; Herrera *et al.*, 2012). These results suggest that epigenetic mechanisms

are important for transcriptional responses to changing environments in diverse fungi,

but many questions remain regarding the precise mechanisms and function of epigenetic

dynamics in fungi.

Epigenetic mechanisms, such as direct modifications of DNA and histone proteins

or physical changes to the chromatin architecture, can influence transcription by

regulating DNA accessibility (Slotkin & Martienssen, 2007). Chromatin that is accessible

and potentially active is termed euchromatin, while heterochromatin is condensed and

often transcriptionally silent (Bannister & Kouzarides, 2011). However, heterochromatic

regions are not always repressed. Heterochromatin is subcategorized into constitutive

heterochromatin that remains condensed throughout the cell cycle, and facultative

heterochromatin that can de-condense to allow transcription in response to

developmental changes or to environmental stimuli (Trojer & Reinberg, 2007; Kim *et al.*,

2015; Wang *et al.*, 2016). In fungi, constitutive heterochromatin is often associated with

repeat-rich genome regions and is typically characterized by tri-methylation of lysine 9

on histone H3 (H3K9me3), while facultative heterochromatin is characterized by tri-

methylation of lysine 27 on histone H3 (H3K27me3) (Lewis *et al.*, 2009, 2010; Jamieson

et al., 2013; Freitag, 2017). Empirically, both H3K9me3 and H3K27me3 have been

implicated in transcriptional regulation in various fungi (Reyes-Dominguez *et al.*, 2010,

2012; Connolly *et al.*, 2013; Wiemann *et al.*, 2013; Chujo & Scott, 2014; Soyer *et al.*,

2014). The majority of these studies rely on genetic perturbation of the enzymes that

deposit methylation at H3K9 and H3K27, and the results consistently show that depletion

of methylation at these lysine residues mainly results in transcriptional induction.

However, as global depletion of a histone modification can result in pleiotropic effects,

such as improper localization of other histone modifications or altered development (Tian

& Chen, 2001; Fan *et al.*, 2017), it is difficult to infer transcriptional control mechanisms

used for natural gene regulation from these genetic perturbation experiments. Therefore,

additional research is needed to directly test the hypothesis that heterochromatin-

associated histone modifications directly regulate transcription either through their

dynamics, or their action to form or recruit transcriptional complexes (Galazka & Freitag,

2014; Seidl *et al.*, 2016; Ridenour *et al.*, 2020).

The filamentous fungus *Verticillium dahliae* is a soil-borne broad host-range plant

pathogen that infects plants through the roots to invade the xylem vessels and cause

vascular wilt disease (Fradin & Thomma, 2006). Genomic and transcriptomic studies

have revealed that the *V. dahliae* genome harbours lineage-specific (LS) regions that are

variable between strains and enriched for genes that are *in planta* induced (Klosterman

et al., 2011; de Jonge *et al.*, 2013). These LS regions are generally considered genomic

hotspots for evolutionary adaptation to plant hosts (Klosterman *et al.*, 2011; de Jonge *et*

al., 2013; Faino *et al.*, 2016; Shi-Kunne *et al.*, 2018; Depotter *et al.*, 2019; Cook *et al.*,

2020). Recently we explored the epigenome of *V. dahliae* and distinguished LS regions

from the core genome based on particular chromatin signatures, including elevated levels

of H3K27me3 accompanied with accessible DNA and active transcription (Cook *et al.*,

2020). Using a machine learning approach and supported by orthogonal analyses, we

identified nearly twice as much LS DNA as previously considered, collectively referred to

as adaptive genomic regions (Cook *et al.*, 2020). Given the elevated levels of H3K27me3

at adaptive genomic regions in *V. dahliae*, and previous reports that removal of

H3K27me3 results in transcriptional induction (Connolly *et al.*, 2013; Chujo & Scott,

2014; Soyer *et al.*, 2014), we now tested if H3K27me3 dynamics are required for

transcriptional activation of genes in adaptive genomic regions. Ideally, the involvement

of H3K27me3 dynamics in differential gene expression of *V. dahliae* would be studied

between *in vitro* and *in planta* growth, as adaptive genomic regions are enriched for *in*

planta induced genes (de Jonge *et al.*, 2013; Cook *et al.*, 2020). However, *V. dahliae*

displays a low pathogen-to-plant biomass during infection (Faino *et al.*, 2012), which

impedes technical procedures to determine histone modification levels over the genome

(Soyer *et al.*, 2015). Nevertheless, as H3K27me3 is generally reported to regulate

transcription in response to environmental stimuli (Trojer & Reinberg, 2007; Wang *et al.*,

2016), we hypothesize that it may be involved in transcriptional regulation in *V. dahliae*

during differential growth conditions *ex planta* as well. Here, we analyse RNA-seq, ChIP-

seq and ATAC-seq data of *V. dahliae* cultured in various axenic growth media to

understand if transcriptional dynamics require concomitant changes in the H3K27me3

modification status.

RESULTS

Genes induced between growth conditions are associated with H3K27me3

We sought to test if genes in adaptive genomic regions, which we recently reported to

have distinct chromatin profiles (Cook *et al.*, 2020), are transcriptionally more responsive

to differential *in vitro* and *in planta* growth conditions. We found that genes that are

higher expressed in Czapec-Dox (CZA) or Murashige-Skoog (MS) media when compared

with potato dextrose broth (PDB) are significantly enriched in adaptive genomic regions

(two-sided two-proportions Z-test, $p<0.01$) (Fig. 1A). More specifically, 125 (14.6%) out

of 856, and 84 (11.8%) out of 711 genes that are higher expressed in CZA or MS than in

PDB, respectively, locate in adaptive genomic regions, while these regions only comprise

998 (8.7%) of the total 11,426 genes in the genome (two-sided two-proportions Z-test,

$p<0.01$) (Fig. 1A). Interestingly, genes with greater transcriptional differences between

the media (i.e. higher $\log_2$-fold changes) appear to be increasingly more overrepresented

in adaptive genomic regions than those with smaller or no differences (Fig. 1A).

We next compared expression during infection of *Arabidopsis thaliana* to

cultivation in PDB*.* We found that genes that are higher expressed during *A. thaliana*

colonization are also overrepresented in adaptive genomic regions, consistent with other

host-infection experiments for *V. dahliae* (de Jonge *et al.*, 2013; Cook *et al.*, 2020).

Similar to what we observed for *in vitro* growth conditions, genes with the greatest

transcriptional dynamics are increasingly more overrepresented in adaptive genomic

regions. That is, there are 145 (14.7%) out of 987 genes higher expressed *in planta*

($\log_2$-fold change >2) residing in adaptive genomic regions, while at $\log_2$-fold change >7

this increases to 26 (36.6%) out of 71 (Fig. 1A). These results show that genes higher

expressed in other *in vitro* growth media compared to PDB*,* as well as genes that are

higher expressed during plant infection compared to PDB, are overrepresented in

adaptive genomic regions. Thus, these data suggest that genes that respond to

environmental cues in general, rather than only genes that are induced *in planta*, are

overrepresented at adaptive genomic regions.

Given the elevated levels of H3K27me3 at adaptive genomic regions in *V. dahliae*

(Cook *et al.*, 2020), one potential explanation for the co-localization of genes that are

differentially expressed *in vitro* and those that are differentially expressed *in planta* is

that these responses to different growth conditions share an adaptive origin that is

epigenetically regulated by H3K27me3. However, given that H3K27me3 occurs not only

in adaptive genomic regions in *V. dahliae*, but also throughout the core genome (Cook *et*

al., 2020), we sought to determine if H3K27me3 is associated with *in vitro* and *in planta*

differentially expressed genes, not only in adaptive genomic regions, but also genome-

wide. Thus, we first performed peak calling on H3K27me3-ChIP data of *V. dahliae*

cultivated for 6 days in PDB and obtained 304 H3K27me3 domains that cover

approximately 10.5 Mb (28.9%) of the genome (Fig. S1), comprising 3,280 (28.7%)

genes and overlapping 87.4% of adaptive genomic regions. Interestingly, the observed

overrepresentation of differentially expressed genes in the generally H3K27me3-marked

adaptive genomic regions extends to H3K27me3 domains throughout the genome,

showing that H3K27me3 is genome-wide associated with differentially expressed genes

(Fig. 1B). Specifically, when assessing what fraction of higher expressed genes in CZA,

MS or Arabidopsis when compared with PDB locate in an H3K27me3 domain, we found

that 421 (49.2%) out of 856, 271 (38.1%) out of 711, and 453 (45.9%) out of 987

locate in H3K27me3 domains, respectively (Fig. 1B, $\log_2$-fold change >2), which is

considerably more than the 28.7% genome-wide average. Additionally, similar to our

observations for adaptive genomic regions, we also observed that genes displaying

higher $\log_2$-fold changes were increasingly overrepresented in H3K27me3 domains. Here

we found that 41 out of 51 (80.4%), 4 out of 7 (57.1%), and 54 out of 71 (76.0%)

genes higher expressed with a $\log_2$-fold change >7 in CZA, MS and Arabidopsis,

respectively, located in H3K27me3 domains (Fig. 1B). Collectively, these data reveal a

strong association between differential gene expression and the presence of H3K27me3

domains, both for genes that are differentially expressed between *in vitro* growth

conditions, as well as for genes that are higher expressed *in planta* than *in vitro.* As we

observed that the association with differentially expressed genes is not limited to

adaptive genomic regions, but is a feature for H3K27me3 domains in general, we infer

that H3K27me3 may be involved in regulating transcriptional changes in response to

environmental cues.

Figure 1. Genes that are higher expressed in CZA, MS and *in planta* than in PDB are overrepresented in adaptive genomic regions and in H3K27me3 domains. Fractions of total genes and genes that are higher expressed than in PDB that locate in (A) adaptive genomic regions and (B) H3K27me3-enriched regions. Increased expression is determined between six days of cultivation *in vitro* of *V. dahliae* in potato dextrose broth (PDB) and 6 days of cultivation in Czapec-Dox (CZA) and half-strength Murashige Skoog medium (MS), and infection of *Arabidopsis thaliana* (21 dpi) (*in planta*). Fractions of higher expressed genes that reside in adaptive genomic regions or H3K27me3-enriched regions are reported for log2-fold changes >1, 2, 3, 4, 5, 6 and 7, respectively. H3K27me3-enrichment was determined for *V. dahliae* cultivation during six days in PDB. Black numbers above the bars indicate numbers of genes per gene set. White numbers at bottom of the bars indicate the numbers of genes that locate in adaptive genomic or H3K27me3-domain per gene set. Significance of fraction differences was calculated for each fraction of higher expressed genes in adaptive genomic regions when compared with the fraction of total genes in adaptive genomic regions, by the two-sided two-proportions Z-test (*: p≤0.05).

Chromatin features correlate with gene expression levels

To generally assess how chromatin impacts transcriptional activity in *V. dahliae*, we determined the occurrence of the heterochromatic histone mark H3K9me3, the euchromatic histone mark H3K4me2, and chromatin accessibility determined by assay for transposase accessible chromatin followed by sequencing (ATAC-seq) (Buenrostro *et al.*, 2015). Grouping the *V. dahliae* genes into five expression quintiles, from quintile 1 containing the highest 20% expressed genes to quintile 5 with the 20% lowest expressed

genes, we are able to integrate histone modification profiles and transcriptional activity.

Expressed genes (quintiles 1-4) displayed low H3K4me2 coverage upstream of the

transcription start site (TSS), followed by a steep increase of coverage over the start of

the gene that decreases over the gene body (Fig. 2B). The strength of the association

corresponds to the level of transcription, also within quintiles (Fig. S2). The low

H3K4me2 coverage near the TSS coincides with increased chromatin accessibility, where

higher expressed quintiles have more open DNA (Fig. 2B, Fig. S2). There is little evidence

of H3K9me3 over gene bodies and *cis*-regulatory regions (Fig. 2B, Fig. S2), which

corroborates that H3K9me3 marks TEs in constitutive heterochromatic regions such as

the centromeres (Lewis *et al.*, 2009; Schotanus *et al.*, 2015; Cook *et al.*, 2020; Seidl *et*

al., 2020). During cultivation in PDB, H3K27me3 is mainly present on genes that are

lowly or not expressed (quintile 4 and 5) (Fig. 2B, Fig. S2). Collectively, this empirical

description of chromatin and transcriptional activity is consistent with reports for other

fungi (Reyes-Dominguez *et al.*, 2010; Connolly *et al.*, 2013; Möller *et al.*, 2019).

To further analyse the association between H3K27me3 and gene expression, we

generated expression quintiles for the genes located in H3K27me3 domains during PDB

growth (3,280 genes). Interestingly, we found that genes in this group with higher

expression values (e.g. quintile 1 and 2) had higher H3K4me2 over the gene body, more

accessible promoters, and lower H3K27me3 values (Fig. 2C). This suggests that genes in

H3K27me3 domains are not uniformly repressed or heterochromatic, and there appeared

to be a quantitative, rather than qualitative association between H3K27me3 association

and gene activity. Genes with a lower association to H3K27me3 may represent loci that

are not in a stable state under the tested conditions, where some cells have a more

euchromatic profile and other have a more heterochromatic profile. While we cannot infer

these details from the current data, it is clear that genes with lower expression in PDB

are generally marked with H3K27me3, have less H3K4me3 and have less accessible DNA

in the region of transcription initiation (i.e. their promoter).

Figure 2. Lowly and non-expressed genes associate with H3K27me3. A) Whole genome distribution of the euchromatin-associated histone modification H3K4me2, the constitutive heterochromatin-associated histone modification H3K9me3, the facultative heterochromatin-associated histone modification H3K27me3 and chromatin accessibility as determined by ATAC-seq, based on chromosome 5 as an example. Genes are indicated in light blue, transposons are indicated in red, centromere is indicated in dark blue and adaptive genomic regions are indicated in yellow. B,C) Relative coverage of chromatin accessibility and the histone marks H3K4me2, H3K9me3, H3K27me3 over gene bodies (between transcription start site (TSS) and transcription end site (TES)) ±800 bp of flanking sequence, grouped into quintiles based on gene expression levels upon cultivation for six days potato dextrose broth (PDB), for (B) all genes, and for (C) all H3K27me3-enriched genes.

Genetic perturbation of H3K27me3 induces transcription of many genes that are differentially expressed *in vitro* and *in planta*

To further characterize the influence of H3K27me3 on gene expression in *V. dahliae*, we deleted the histone methyltransferase component of the Polycomb Repressive Complex 2 (PRC2), termed *Set7* (Δ*Set7*), leading to loss of H3K27me3 (Fig. S3). As H3K27me3 is

generally associated with facultative heterochromatin, we anticipated that the loss of H3K27me3 would mainly lead to induction of genes that were located in H3K27me3 domains in wild-type *V. dahliae*. Out of the 839 genes that are induced in ΔSet7 (log$_2$-fold change >2, p<0.05), 615 (73.3%) locate in H3K27me3 domains, which is significantly higher than expected given that only 28.7% of genes locate in H3K27me3 domains (Fisher's exact test, p<0.00001). In contrast, we find that 214 (28.0%) of 765 repressed genes in ΔSet7 are in H3K27me3 domains (no association, Fisher's exact test, p=0.78) (Fig. 3A,B). Additionally, when comparing log$_2$-fold changes in expression between ΔSet7 and the wild type strain, we observed that genes located in a H3K27me3 domain in wild type are more significantly induced in ΔSet7 than genes not locating in H3K27me3 domains (Two-sample Student's T-test, p<0.001) (Fig. S4). These findings suggest that H3K27me3 represses transcription at marked regions.

We next sought to integrate the repressive transcriptional effects of H3K27me3 inferred from ΔSet7 RNA-seq with the differential transcription observed between growth conditions. For this, we compared genes induced in ΔSet7 versus wild-type PDB growth with genes higher expressed in CZA, MS or *in planta* versus PDB growth. There is a significant overrepresentation of genes induced in ΔSet7 (839), and those induced in CZA (254 genes, 30.3% overlap, Fisher's exact test p<0.00001), MS (197 genes, 23.5% overlap, Fisher's exact test p<0.00001) and *in planta* (208 genes, 24.8% overlap, Fisher's exact test p<0.00001). These associations are higher than the background levels of genes induced under CZA, MS or *in planta*, which is 7.5%, 6.2%, and 8.6% of genes, respectively (Fig. 3C-E). Together, these data show that H3K27me3 is required for transcriptional silencing of mainly H3K27me3 associated genes, and that many of these genes are also differentially expressed between the tested growth conditions. These results support the hypothesis that transcriptional dynamics between the tested growth conditions are regulated, at least partly, by levels of H3K27me3.

Figure 3. Genetic perturbation of H3K27me3 results in induction of genes that are transcriptionally regulated in different growth conditions. A) Fractions of induced (log2 fold change >1, p<0.05) and repressed (log2 fold change <-1, p<0.05) genes and genes that are not differentially expressed between wild-type and ΔSet7 that (B) locate in H3K27me3 domains and (C-E) are higher expressed in CZA, MS and *in planta* than in PDB.

H3K27me3 domains are globally stable between *in vitro* growth conditions

Under the hypothesis that H3K27me3 dynamics contribute to differential gene expression, we predict that H3K27me3 ChIP-seq domains will change when *V. dahliae* is grown under different conditions. In addition, the observed changes in H3K27me3 domains will be associated with transcriptional differences of the underlying genes. To test this hypothesis, we performed additional H3K27me3 ChIP-seq for *V. dahliae* cultured in MS and CZA, and identified H3K27me3 domains from replicate H3K27me3 ChIP-seq in each of the three growth media. We identified a total of 1,631 genes that were always present in H3K27me3 domains, regardless of the *in vitro* growth media (Fig. 4A). Not surprisingly, the 1,631 genes present in stable H3K27me3 domains were significantly lower expressed when compared with non-marked genes in all three growth conditions (Fig. 4B). To assess if changes in H3K27me3 domains between the media were associated with changes in transcription, we compared domains and gene expression between media. We first analysed RNA-seq output for genes present in H3K27me3

domains in two media but not in an H3K27me3 domain in the third medium. For

example, we identified 703 genes that were in a H3K27me3 domain in PDB and in MS but

not in CZA (Fig. 4A), and found that these genes were more highly expressed during CZA

growth when compared with growth in PDB or in MS (Fig. 4C). These result corroborated

the association between the lack of H3K27me3 in CZA and increased transcription of the

underlying genes. However, genes present in shared H3K27me3 domains for MS and CZA

growth (229), or CZA and PDB growth (44) did not display statistically significant

transcriptional differences between the growth conditions where the genes lack

H3K27me3 domains (Fig. 4A, C). Analysing H3K27me3 domains unique to a media, we

found that MS growth had the greatest number and proportion of unique genes locating

in H3K27me3 domains (18.3%), followed by 12.4% unique to CZA growth, and 4.7%

unique to PDB growth (Fig. 4A). The genes uniquely marked in PDB had a non-

significant, but lower median expression value when compared with the other two media.

Genes uniquely located in MS H3K27me3 domains were significantly lower expressed

when compared with growth in CZA, but not with growth in PDB (Fig. 4D). The genes

uniquely present in CZA H3K27me3 domains did not display transcriptional differences

when *V. dahliae* was grown in the other two media. Overall, these results show that

H3K27me3 contributes to transcriptional repression, but there is not a simple global

association between H3K27me3 presence/absence dynamics and transcriptional output.

We observed clear examples where the loss of H3K27me3 in one medium is associated

with increased transcription in that medium, but this is not ubiquitous. We note that the

majority of genes located in H3K27me3 domains were common to all three growth

conditions, accounting for 75%, 65.4% and 52% of the identified genes in H3K27me3

domains from CZA, PDB, and MS growth, respectively, indicating that the qualitative

presence of H3K27me3 domains is globally stable.

Figure 4. Differential H3K27 methylation partially explains differential expression. A) The number of genes located in H3K27me3 domains for *V. dahliae* cultivated for 6 days in Czapec-Dox (CZA), half-strength Murashige Skoog medium (MS) and Potato dextrose broth (PDB). B) Transcriptional values from RNA-seq, represented as TPM values in each growth medium. The genes were grouped between those consistently localized in H3K27me3 domains (1631 genes) and genes not present in H3K27me3 domains. C,D) Transcription TPM values in each growth medium for genes in H3K27me3 domains found in C) two or D) one of three tested *in vitro* growth media. Median values of violin plots are indicated with diamonds. Significant difference of TPM values between growth media are determined with the One-Sample Wilcoxon Signed Rank Test (*: $p <= 0.05$).

Local quantitative differences in H3K27me3 levels are associated with transcriptional differences

Our analysis on H3K27me3 presence/absence dynamics did not account for potential quantitative differences between growth media. Whole chromosome plots of H3K27me3 domains identified between media reflect their generally stable presence (Fig. S5A), but the quantitative level of ChIP-seq signal appeared to vary at some loci within called domains (Fig. S5B). To address this, we examined quantitative differences between H3K27me3 ChIP-seq signal and transcriptional output between pairs of growth conditions. Genes were grouped based on differential gene expression between media,

where genes higher expressed in medium A were one group and genes higher expressed

in medium B another group. The H3K27me3 ChIP-seq levels were compared between the

groups for the ChIP conducted in both the transcriptionally permissive and non-

permissive growth conditions. Comparing results for PDB versus MS, we see that genes

higher expressed in MS have a significantly lower H3K27me3 signal from MS-ChIP versus

the same genes from PDB-ChIP (Fig. 5A). For genes more highly expressed in PDB, there

is a higher, but not significantly different, H3K27me3 ChIP-seq level when grown in MS

when compared with PDB (Fig. 5A). These results support the idea that when a gene is

transcriptionally active in one media, the H3K27me3 levels are lower than H3K27me3

levels under less transcriptionally permissive growth. Integrating the transcriptional and

ChIP fold-changes for all genes between the media, we see that this association is true

for some, but not all genes that undergo transcriptional changes between the media (Fig.

5B). Globally, the distribution of H3K27me3 fold-changes per gene between media shows

that genes higher expressed in PDB have a significantly higher H3K27me3 fold-change

(i.e. less H3K27me3 in PDB growth), while MS higher expressed genes have a

significantly lower H3K27me3 fold-change (i.e. less H3K27me3 in MS growth) (Fig. 5B).

Results for PDB versus CZA growth showed that genes higher expressed in CZA

have higher H3K27me3 levels in PDB, consistent with the expected association (Fig. 5C).

For genes higher expressed in PDB however, we observed a higher H3K27me3 level in

PDB. Globally, the data indicated a lower H3K27me3 ChIP-signal for samples grown in

CZA (Fig. 5D, data shifted to left on x-axis). The integrated results indicated that genes

higher expressed in CZA had less H3K27me3 in CZA, but the PDB higher expressed

genes also had lower H3K27me3 in CZA. This difference, the overall lower H3K27me3

levels in CZA, may represent a biologically relevant difference between the growth

media, but may also reflect a technical bias encountered for ChIP using CZA grown

material. It is not possible to fully resolve these two possibilities with our current data.

Results for CZA versus MS growth showed that genes higher expressed in MS had

statistically similar levels of H3K27me3 ChIP-signal between the two media (Fig. 5E). For

genes higher expressed in CZA, there was significantly higher H3K27me3 ChIP-signal

during MS growth (Fig. 5E).   The genome-wide integrated analysis, combining fold-

change between CZA and MS for transcription and H3K27me3 ChIP, showed that genes

more highly expressed in CZA indeed have significantly higher levels of H3K27me3 when

grown in MS (Fig. 5F). We again see the effect of the lower H3K27me3 ChIP-signal from

CZA-grown material, where the global fold-change values indicate higher levels of

H3M27me3 in MS-grown material overall (Fig. 5F). However, there is a clear group of

genes that are more highly expressed in MS medium that have an H3K27me3 fold-

change indicating a shift toward higher levels in CZA. This can be seen by the

significantly different marginal density plot for H3K27me3 fold-change, where the genes

higher expressed in MS have a significantly greater change towards higher H3K27me3

levels in CZA (Fig. 5F). Overall, the results show that in most comparisons, there is an

association between quantitative transcriptional levels and H3K27me3 signal, where

genes more highly expressed in a transcriptionally permissive media have less

H3K27me3 when compared to H3K27me3 levels in the repressive media. These results

are consistent with the observations of the qualitative H3K27me3 presence/absence

comparisons and suggests that H3K27me3 levels are generally stable, but can vary for

some genes in a quantitative manner to impact their transcription.

Figure 5. Differentially expressed genes associate with local changes in H3K27me3 coverage. Pairwise comparisons of differentially expressed genes for *V. dahliae* cultivated for 6 days in A,B) half-strength Murashige Skoog medium (MS) and Potato dextrose broth (PDB), C,D) Czapec-Dox (CZA) and PDB, and E,F) MS and CZA. A,C,E) H3K27me3 coverage over differentially expressed genes in each corresponding growth medium. Median values of violin plots are indicated with diamonds. B,D,F) per gene comparison of log2-fold change in transcription and genic H3K27me3 coverage. Black horizontal lines indicate TPM $\log_2$-fold change of -2 and 2, black vertical lines indicate H3K27me3 $\log_2$-fold change of -1 and 1. Significant difference of TPM values between growth media are determined with the One-Sample Wilcoxon Signed Rank Test and indicated by letters or by asterisks (*: $p \leq 0.05$).

Genes induced *in planta* are largely H3K27me3 associated across all tested growth conditions

The presented analysis address histone modification and transcriptional dynamics between axenic growth medium. A significant question is how do H3K27me3 levels compare between more environmentally drastic growth conditions, such as *in vitro* versus *in planta* growth. As we have thus far been unable to perform ChIP on *V. dahliae* during host colonization due to a low pathogen-to-plant biomass, we assessed the *in vitro* chromatin profiles for genes that are significantly induced during infection. First, we grouped genes based on differential expression between PDB and *in planta* growth and analysed the PDB grown chromatin profiles. We saw that genes higher expressed *in planta* have significantly more H3K27me3 during PDB growth when compared with the genes more highly expressed in PDB (Fig. 6A). These *in planta*-induced genes lacked H3K4me2 and the DNA is less accessible during PDB growth, while genes highly expressed in PDB are open and in active euchromatin (Fig. 6A). This pattern of *in planta*-induced genes being highly H3K27me3-associated is also observed when analysed relative to growth in MS and CZA (Fig. 6B,C).

To understand what genes are driving these differences, we investigated whether *in planta*-induced genes that locate in H3K27me3 domains are overrepresented for genes potentially involved in infection. There were approximately 1,000 genes more highly expressed *in planta* when compared with any of the three media (Fig. 6D). We found that, depending on the medium, from 36.5% to 53.4% of the *in planta*-induced genes were located in H3K27me3 domains (Fig. 6D). We observed that the *in planta*-induced genes within H3K27me3 domains have a higher, yet not statistically significant, fraction of genes encoding secreted proteins or putative effectors when compared to all *in planta* induced genes. Specifically, for genes in H3K27me3 domains that were *in planta*-induced compared to PDB growth (376 genes total), 16.0% had a secretion signal and 5.1% were predicted effectors. This is compared to all *in planta* induced genes relative to PDB (987 genes total), where 12.3% had a secretion signal and 3.2% were predicted effectors. For CZA, 17.0% of the genes in H3K27me3 domains that were *in planta*-induced (418 genes

total) had a secretion signal and 5.7% were predicted effectors. This is compared to all *in*

planta induced genes relative to CZA (1,144 genes total), where 13.4% had a secretion

signal and 3.8% were predicted effectors. The results were similar for comparison to MS

grown samples. Thus, we consistently observed that *in planta*-induced genes in

H3K27me3 domains have a slightly higher fraction of genes that are potentially involved

in infection. Collectively, these results indicate that genes required for infection associate

with H3K27me3 and that chromatin dynamics are likely needed during *in planta* infection

to increase DNA accessibility concomitant with *in planta* transcription compared to the

measured state in PDB. Whether or not this activation is accompanied by loss or

dynamics of H3K27me3 levels will need to be directly assessed *in planta* in the future.

Figure 6. *In planta*-induced genes are H3K27me3-associated under transcriptionally
repressive conditions. Average coverage of chromatin features over genes differentially
expressed between in planta and A) 6 days of cultivation in Potato dextrose broth (PDB), B) half
strength Murashige Skoog medium (MS), and C) Czapec-Dox medium (CZA). D) overlap of genes
that are higher expressed *in planta* than in PDB, MS or CZA with genes located in an H3K27me3
domain in the corresponding growth medium. Significant differences of H3K27me3 coverage are
determined with the One-Sample Wilcoxon Signed Rank Test (*: p <= 0.05).

DISCUSSION

Chemical histone modifications play an essential role in transcriptional regulation, but questions remain regarding their mechanism, especially in filamentous fungi. In our study we show that many differentially transcribed genes between *in vitro* and *in planta* growth are located in previously defined adaptive genomic regions (Cook *et al.*, 2020). We report that a majority of the genes identified as being differentially transcribed between the tested conditions reside in H3K27me3 domains. This mark only occupies 28.7% of the genes in the genome, but the rate of H3K27me3 association significantly increases for differentially expressed genes by 35% to 70%.

Previous reports propose that H3K27me3 represses transcription of secondary metabolite clusters in *Fusarium spp.* and *Epichloe festucae*, and effectors in *Zymoseptoria tritici* and *Magnaporthe oryzae* (Connolly *et al.*, 2013; Chujo & Scott, 2014; Studt *et al.*, 2016; Möller *et al.*, 2019; Zhang *et al.*, 2021). However, these reports have mainly relied on genomic association with H3K27me3, genetic perturbation altering H3K27me3 deposition, and the analysis of only a few genes under natural conditions. Given these previous findings, our hypothesis was that H3K27me3 levels are dynamic during growth in different conditions and contribute to differential transcription. We directly addressed this using genome-wide H3K27me3 profiling and RNA-seq under different *in vitro* growth conditions. Consistent with reports in other fungi, we see that H3K27me3 is associated with transcriptional repression, and that deleting *Set7*, encoding the histone methyltransferase that is responsible for H3K27me3, led to induction of many genes normally present in H3K27me3 domains. Importantly, our direct observations of H3K27me3 levels and transcriptional output from three *in vitro* media showed that H3K27me3 domains are generally stable globally. We see that 50% to 75% of the identified H3K27me3 domains, depending on the medium analyzed, did not change between the tested conditions. Despite many H3K27me3 domains not undergoing presence/absence dynamics, numerous genes in these domains displayed differential transcription between the tested conditions, indicating that complete loss of H3K27me3 is not strictly required for transcriptional induction.

Even though our data suggest that qualitative H3K27me3 changes are not

sufficient to explain differential transcription under natural conditions alone, we do

provide empirical evidence that quantitative H3K27me3 dynamics are associated with

differentially expressed genes in *V. dahliae* grown under different *in vitro* conditions.

These results show that some genes grown in transcriptionally permissive media have a

lower H3K27me3 ChIP-signal when compared with growth in a transcriptionally

repressive media. We interpret these results as evidence for local, not global,

contribution of H3K27me3 quantitative dynamics to transcriptional regulation for these

genes. These results are consistent with data from *M. oryzae*, in which direct *in planta*

H3K27me3 ChIP-qPCR showed that some, but not all, analyzed regions displayed

differential H3K27me3 levels between *in planta* and *in vitro* conditions (Zhang *et al.*,

2021). We have not been able to directly assess histone modifications for *V. dahliae*

during plant infection, but we did analyze the *in vitro* growth H3K27me3 status of *in*

planta-induced genes. The identified *in planta*-induced genes have significantly higher

levels of H3K27me3 and the DNA is substantially less accessible compared to genes

highly expressed during *in vitro* growth. We conclude that dynamics for H3K27me3 can

contribute to differential gene expression under natural conditions, but it is not required,

and additional regulatory mechanisms are likely important to increase DNA accessibility

and enact transcription. An interesting finding in our research was the quantitative

association we observed between H3K27me3 levels and transcriptional output between

the growth conditions. This is interesting because it is not clear what mechanism explains

a quantitative rather than qualitative change in H3K27me3 levels. That is, a nucleosome

can have none, one, or two H3K27 tails with a tri-methyl modification, and quantitative

differences seen in our samples reflect either heterogeneity in H3K27me3 status for cells

in the population (e.g., 50% of cells are H3K27me3 and 50% are H3K27 non-modified)

or it represents bivalent chromatin domains (e.g., 100% of cells have 50% H3K27me3).

Bivalent domains are generally defined as DNA regions interacting with nucleosomes that

contain chemical modifications representative of both active and repressed chromatin,

with the prevailing example being the presence of both H3K4me3 and H3K27me3 (Blanco

et al., 2020). Bivalent domains have been identified and described in animal systems,

and are thought to exist in a poised transcriptional state, with low steady state

transcription capable of rapid transcriptional activation during cellular differentiation

(Bernstein *et al.*, 2006; Voigt *et al.*, 2013). To our knowledge, bivalent domains have not

been formally described in fungi. Whether our data are the result of a heterogenous

population of cells or represent bivalent domains remains to be resolved. Nevertheless,

our results show that many genes differentially expressed between growth conditions are

associated with quantitative differences in H3K27me3 levels, providing evidence that

chemical histone modifications dynamics can play a transcriptional regulatory mechanism

in *V. dahliae*. As our experiments were based on differences between growth conditions,

these results may represent a general regulatory mechanism for environmentally

response genes.

At present, our conclusions are limited to the tested conditions, and it is possible

that analyzing H3K27me3 levels in different cell-types or growth stages (e.g. spores,

microsclerotia, *in planta* infection) may yield a different picture of global H3K27me3

distribution. Additionally, our experiments focused on steady-state growth on different

media as these provide a clear and reproducible transcriptional difference. It is possible

that H3K27me3 dynamics occur rapidly in response to environmental changes, cues, or

developmental stages that we did not capture.

It appears evident that H3K27me3 contributes to additional genome functions

beyond strict transcriptional repression, at least in fungi (Ridenour *et al.*, 2020). This is

supported by our data showing the a majority of H3K27me3 domains were stable

between the tested growth conditions, indicating these H3K27me3 domains have

additional functions. These stable H3K27me3 domains may represent another form of

constitutive heterochromatin, but this fact seems unlikely given that in our data many

stable H3K27me3 domains harbored differentially expressed genes between growth

media. Additionally, results in *N. crassa* have shown that genetic perturbation leading to

the genome-wide loss of H3K9me3 causes a redistribution of H3K27me3 to previously

H3K9me3 marked sites, but interestingly, this re-distribution does not result in

transcriptional silencing (Basenko *et al.*, 2015). Rather, the redistribution of H3K27me3

in *N. crassa* lacking H3K9me3 appears to contribute to genomic instability (Basenko *et

al.*, 2015; Lewis, 2017). In *Z. tritici*, the enrichment of H3K27me3 at dispensable

chromosomes and empirical evidence that H3K27me3 somehow increases genomic

instability, additionally supports the hypothesis that H3K27me3 contributes to additional

genomic functions beyond transcriptional regulation (Schotanus *et al.*, 2015; Möller *et

al.*, 2019). Additional research is needed to fully understand the mechanisms of

H3K27me3 targeting, dynamics, and impact on genome stability in fungi. The results

presented here show that H3K27me3 domains are largely similar between *in vitro* growth

conditions, but that quantitative differences in H3K27me3 levels are associated with

concomitant transcriptional differences between the tested conditions.

MATERIALS AND METHODS

Fungal growth conditions

Verticillium dahliae strain JR2 (CBS 143773; Faino et al., 2015) was cultured on potato dextrose agar (PDA) (Oxoid, Thermo Scientific) at 22°C in the dark. Liquid cultures were obtained by collecting conidiospores from PDA plates after approximately 2 weeks of growth followed by inoculation at a final concentration of 1×10^4 spores per mL into liquid growth media. Media used in this study are potato dextrose broth (PDB) (Difco, Becton Dickinson, Franklin Lakes, NJ, USA), half strength Murashige & Skoog plus vitamins (MS) (Duchefa-Biochemie, Haarlem, The Netherlands) medium supplemented with 3% sucrose, and Czapec-Dox medium (CZA) (Oxoid, Thermo Scientific, Waltham, MA, USA). Liquid cultures were grown for 6 days in the dark at 22°C at 140 RPM. Mycelium was collected by straining cultures through miracloth (22 μm) (EMD Millipore, Darmstadt, Germany) and pressing to remove liquid after which the mycelium was flash frozen in liquid nitrogen and ground to powder with a mortar and pestle. If required, samples were stored at -20°C prior to nucleic acid extraction. All analyses were performed based on triplicate cultures that were processed individually.

RNA sequencing and analysis

RNA was isolated from ground mycelium using TRIzol (Thermo Fisher Science, Waltham, MA, USA) following manufacturer guidelines. Contaminating DNA was removed using the TURBO DNA-free kit (Ambion, Thermo Fisher Science, Waltham, MA, USA) and RNA integrity was assessed with gel-electrophoresis and quantified using a Nanodrop (Thermo Fisher Science, Waltham, MA, USA). Library were prepared and singleend 50 bp sequenced on the DNBseq platform at BGI (Hong Kong, China).

Sequencing reads were mapped to the reference annotation of *V. dahliae* strain JR2 (Faino *et al.*, 2015) using Kallisto quant (Bray *et al.*, 2016) to calculate per gene TPM values. Differential expression between cultivation in PDB and CZA, MS or during colonization of *Arabidopsis thaliana* was determined using DESeq2 (Love *et al.*, 2014).

H3K27me3-enriched regions were determined using epic2 with a binsize of 2,500 bp

(Stovner & Sætrom, 2019).

ChIP-sequencing and analysis

ChIP-seq was performed as described previously (Cook et al., 2020). Frozen ground

mycelium was added to ChIP lysis buffer, dounced 40 times and subsequently sonicated

for 5 rounds of 20 seconds with 40 second rest stages on ice. Supernatants were

collected after centrifugation and treated with MNase (New England Biolabs, Ipswich, MA,

United States) for 10 minutes in a 37°C waterbath. MNase activity was quenched by

addition of EGTA and samples were subsequently pre-cleared by addition of 40 µl Protein

A Magnetic Beads (New England Biolabs, Ipswich, MA, United States) and rotating at 4°C

for 60 min. Beads were captured and supernatant was divided over new tubes containing

antibodies against either H3K4me2, H3K9me3 or H3K27me3 (ActiveMotif; #39913,

#39765 and #39155) and incubated overnight with continuous rotation at 4°C.

Subsequently, the antibodies were captured, washed and nucleosomes were eluted from

beads, after which DNA was treated with Proteinase-K and cleaned-up using chloroform.

DNA was isolated by overnight precipitation in ethanol and DNA concentration was

determined with the Qubit™ dsDNA HS Assay Kit (Thermo Fisher Science, Waltham, MA,

USA). Sequencing libraries were generated using the TruSeq ChIP Library Preparation Kit

(Illumina, San Diego, CA, United States) according to instructions, but without gel

purification and with use of the Velocity DNA Polymerase (BioLine, Luckenwalde,

Germany) for 25 cycles of amplification. Single-end 125bp sequencing was performed on

the Illumina HiSeq2500 platform at KeyGene N.V. (Wageningen, the Netherlands).

Raw reads were trimmed using TrimGalore (Martin, 2011) and mapped to the *V.*

dahliae strain JR2 reference genome (Faino *et al.*, 2015) using BWA-mem with default

settings (Li, 2013). Three regions of the genome were masked due to aberrant mapping,

(chr1:1-45000, chr2:3466000-3475000, chr3:1-4200). ATAC-seq reads of *V. dahliae*

cultured in PDB (Cook *et al.*, 2020) were treated similarly as ChIP-seq reads, but the

paired end read pairs were trimmed and mapped simultaneously, and only read-pairs

<100 bp were considered for further analyses, as these represent open DNA. Mapped

reads were RPGC-normalized using deepTools bamCoverage (Ramírez *et al.*, 2014) with

binsize 1,000 and smoothlength 3,000 for plotting over the genome, and binsize 10 and

smoothlength 30 for further analysis. Average coverage over gene bodies per expression

quintile was calculated using deepTools computeMatrix in scale-regions mode (Ramírez *et*

al., 2014). Expression quintiles were generated by sorting all genes based on their

average TPM value from three replicates of *V. dahliae* cultured for 6 days in PDB. To

assess the presence of the histone marks over promoters (500 bp upstream of TSS) and

gene bodies (region between TSS and TES) of genes located in LS regions and the core

genome, we used deepTools multiBigwigSummary (Ramírez *et al.*, 2014) to calculate the

average coverage of the histone marks and chromatin accessibility for each gene.

Changes in H3K27me3 levels was determined by calculating the average coverage

over a gene using featureCounts (Liao *et al.*, 2014) on sorted bam files, and comparing

the coverage between duplicates of cultivation in PDB and CZA or MS with DESeq2 (Love

et al., 2014). Changes in H3K27me3 and differential expression per gene were analysed.

Generation of *Set7* deletion mutant

The *Set7* deletion mutant (Δ*Set7*) was constructed as previously described (Santhanam,

2012). Briefly, genomic DNA regions flanking the 5' and 3' ends of the coding sequences

were amplified with PCR using primers listed in Table S1 (primers 1-4) and cloned in to

the pRF-HU2 vector (Frandsen *et al.*, 2008) using USER enzyme following the

manufacturer's protocol (New England Biolabs, MA, USA). Sequence-verified vectors were

transformed into *Agrobacterium tumefaciens* strain AGL1 and used for *V. dahliae*

conidiospore transformation as described previously (Santhanam, 2012). *V. dahliae*

transformants that appeared on hygromycin B were transferred to fresh PDA

supplemented with hygromycin B after five days. Putative transformants were screened

using PCR to verify deletion of the target gene sequence and integration of the selection

marker at the designated locus using primers listed in Table S1 (primers 5-8). Analysis

and comparison of gene expression in the *Set7* deletion mutant to type *V. dahliae* was

performed as described above.

ACKNOWLEDGEMENTS

This work was supported by a PhD grant of the Research Council Earth and Life Sciences

(project 831.15.002) to HMK, and by Human Frontier Science Program Postdoctoral

Fellowship (HFSP, LT000627/2014-L), by USDA's National Institute of Food and

Agriculture (award no. 2018-67013-28492) through the Plant Biotic Interactions

Program, and by the National Science Foundation (award no. 1936800) Division of

Molecular and Cellular Biosciences to DEC. Work in the laboratories of M.F.S and

B.P.H.J.T. is supported by the Research Council Earth and Life Sciences (ALW) of the

Netherlands Organization of Scientific Research (NWO). B.P.H.J.T acknowledges funding

by the Alexander von Humboldt Foundation in the framework of an Alexander von

Humboldt Professorship endowed by the German Federal Ministry of Education and

Research is furthermore supported by the Deutsche Forschungsgemeinschaft (DFG,

German Research Foundation) under Germany´s Excellence Strategy – EXC 2048/1 –

Project ID: 390686111. The authors declare no conflict of interest.

REFERENCES

Aguilera J, Randez-Gil F, Prieto JA. **2007**. Cold response in *Saccharomyces cerevisiae*: new functions for old mechanisms. *FEMS Microbiology Reviews* **31**: 327–341.

Bahn Y-S, Xue C, Idnurm A, Rutherford JC, Heitman J, Cardenas ME. **2007**. Sensing the environment: lessons from fungi. *Nature Reviews Microbiology* **5**: 57–69.

Bannister AJ, Kouzarides T. **2011**. Regulation of chromatin by histone modifications. *Cell Research* **21**: 381–395.

Basenko EY, Sasaki T, Ji L, Prybol CJ, Burckhardt RM, Schmitz RJ, Lewis ZA. **2015**. Genome-wide redistribution of H3K27me3 is linked to genotoxic stress and defective growth. *Proceedings of the National Academy of Sciences* **112**: E6339–E6348.

Bernstein BE, Mikkelsen TS, Xie X, Kamal M, Huebert DJ, Cuff J, Fry B, Meissner A, Wernig M, Plath K. **2006**. A bivalent chromatin structure marks key developmental genes in embryonic stem cells. *Cell* **125**: 315–326.

Blanco E, González-Ramírez M, Alcaine-Colet A, Aranda S, Di Croce L. **2020**. The bivalent genome: characterization, structure, and regulation. *Trends in Genetics* **36**: 118–131.

Bray NL, Pimentel H, Melsted P, Pachter L. **2016**. Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology* **34**: 525–527.

Buenrostro JD, Wu B, Chang HY, Greenleaf WJ. **2015**. ATAC-seq: A method for assaying chromatin accessibility genome-wide. *Current Protocols in Molecular Biology* **2015**: doi: 10.1002/0471142727.mb2129s109.

Chujo T, Scott B. **2014**. Histone H3K9 and H3K27 methylation regulates fungal alkaloid biosynthesis in a fungal endophyte-plant symbiosis. *Molecular Microbiology* **92**: 413–434.

Connolly LR, Smith KM, Freitag M. **2013**. The *Fusarium graminearum* histone H3 K27 methyltransferase KMT6 regulates development and expression of secondary metabolite gene clusters. *PLoS Genetics* **9**: e1003916.

Cook DE, Kramer HM, Torres DE, Seidl MF, Thomma BPHJ. **2020**. A unique chromatin profile defines adaptive genomic regions in a fungal plant pathogen. *eLife* **9**: e62208.

Depotter JRL, Shi-Kunne X, Missonnier H, Liu T, Faino L, Berg GCM, Wood TA,

Zhang B, Jacques A, Seidl MF, _et al._ 2019. Dynamic virulence-related regions of the

plant pathogenic fungus _Verticillium dahliae_ display enhanced sequence conservation.

Molecular Ecology: doi: 10.1111/mec.15168.

Faino L, de Jonge R, Thomma BPHJ. **2012**. The transcriptome of _Verticillium dahliae-_

infected _Nicotiana benthamiana_ determined by deep RNA sequencing. _Plant Signaling &_

Behavior **7**: 1065–1069.

Faino L, Seidl M, Datema E, van den Berg GCM, Janssen A, Wittenberg AHJ,

Thomma BPHJ. **2015**. Single-molecule real-time sequencing combined with optical

mapping yields completely finished fungal genome. _mBio_ **6**: e00936–15.

Faino L, Seidl MF, Shi-Kunne X, Pauper M, Van Den Berg GCM, Wittenberg AHJ,

Thomma BPHJ. **2016**. Transposons passively and actively contribute to evolution of the

two-speed genome of a fungal pathogen. _Genome Research_ **26**: 1091–1100.

Fan A, Mi W, Liu Z, Zeng G, Zhang P, Hu Y, Fang W, Yin W-B. **2017**. Deletion of a

histone acetyltransferase leads to the pleiotropic activation of natural products in

Metarhizium robertsii. _Organic Letters_ **19**: 1686–1689.

Fradin EF, Thomma BPHJ. **2006**. Physiology and molecular aspects of Verticillium wilt

diseases caused by _V. dahliae_ and _V. albo-atrum_. _Molecular Plant Pathology_ **7**: 71–86.

Frandsen RJN, Andersson JA, Kristensen MB, Giese H. **2008**. Efficient four

fragment cloning for the construction of vectors for targeted gene replacement in

filamentous fungi. _BMC Molecular Biology_ **9**: 70.

Freitag M. **2017**. Histone Methylation by SET Domain Proteins in Fungi. _Annual Review_

of Microbiology **71**: 413–439.

Galazka JM, Freitag M. **2014**. Variability of chromosome structure in pathogenic fungi-

of 'ends and odds'. _Current Opinion in Microbiology_ **20**: 19–26.

Gibson G. **2008**. The environmental contribution to gene expression profiles. _Nature_

Reviews Genetics **9**: 575–581.

Heijmans BT, Tobi EW, Stein AD, Putter H, Blauw GJ, Susser ES, Slagboom PE,

Lumey LH. **2008**. Persistent epigenetic differences associated with prenatal exposure to

famine in humans. *Proceedings of the National Academy of Sciences* **105**: 17046–17049.

Herrera CM, Pozo MI, Bazaga P. **2012**. Jack of all nectars, master of most: DNA

methylation and the epigenetic basis of niche width in a flower-living yeast. *Molecular*

Ecology **21**: 2602–2616.

Jamieson K, Rountree MR, Lewis Z a, Stajich JE, Selker EU. **2013**. Regional control

of histone H3 lysine 27 methylation in *Neurospora*. *Proceedings of the National Academy*

of Sciences of the United States of America **110**: 6027–32.

De Jonge R, Bolton MD, Kombrink A, van den Berg GCM, Yadeta KA, Thomma

BPHJ. **2013**. Extensive chromosomal reshuffling drives evolution of virulence in an

asexual pathogen. *Genome Research* **23**: 1271–1282.

Kim J-M, Sasaki T, Ueda M, Sako K, Seki M. **2015**. Chromatin changes in response to

drought, salinity, heat, and cold stresses in plants. *Frontiers in Plant Science* **6**: doi:

10.3389/fpls.2015.00114.

Klosterman SJ, Subbarao K V., Kang S, Veronese P, Gold SE, Thomma BPHJ,

Chen Z, Henrissat B, Lee YH, Park J, *et al.* 2011. Comparative genomics yields

insights into niche adaptation of plant vascular wilt pathogens. *PLoS Pathogens* **7**:

e1002137.

Kronholm I, Johannesson H, Ketola T. **2016**. Epigenetic control of phenotypic

plasticity in the filamentous fungus *Neurospora crassa*. *G3: Genes, Genomes, Genetics* **6**:

4009–4022.

Lewis ZA. **2017**. Polycomb group systems in fungi: new models for understanding

polycomb repressive complex 2. *Trends in Genetics* **33**: 220–231.

Lewis ZA, Adhvaryu KK, Honda S, Shiver AL, Knip M, Sack R, Selker EU. **2010**.

DNA methylation and normal chromosome behavior in *Neurospora* depend on five

components of a histone methyltransferase complex, DCDC. *PLoS Genet* **6**: e1001196.

Lewis ZA, Honda S, Khlafallah TK, Jeffress JK, Freitag M, Mohn F, Schübeler D,

Selker EU. **2009**. Relics of repeat-induced point mutation direct heterochromatin

formation in *Neurospora crassa*. *Genome Research* **19**: 427–437.

Li H. **2013**. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. *arXiv:1303.3997*.

Liao Y, Smyth GK, Shi W. **2014**. featureCounts: an efficient general purpose program

for assigning sequence reads to genomic features. *Bioinformatics* **30**: 923–930.

Love MI, Huber W, Anders S. **2014**. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**: 550.

Martin M. **2011**. Cutadapt removes adapter sequences from high-throughput

sequencing reads. *EMBnet Journal* **17**: 10–12.

Martin EM, Fry RC. **2018**. Environmental influences on the epigenome: exposure-

associated DNA methylation in human populations. *Annual review of public health* **39**:

309–333.

Möller M, Schotanus K, Soyer JL, Haueisen J, Happ K, Stralucke M, Happel P,

Smith KM, Connolly LR, Freitag M, *et al.* **2019**. Destabilization of chromosome

structure by histone H3 lysine 27 methylation. *PLoS Genetics* **15**: e1008093.

Ramírez F, Dündar F, Diehl S, Grüning BA, Manke T. **2014**. deepTools: a flexible

platform for exploring deep-sequencing data. *Nucleic Acids Research* **42**: W187–W191.

Reyes-Dominguez Y, Boedi S, Sulyok M, Wiesenberger G, Stoppacher N, Krska R,

Strauss J. **2012**. Heterochromatin influences the secondary metabolite profile in the

plant pathogen *Fusarium graminearum*. *Fungal Genetics and Biology* **49**: 39–47.

Reyes-Dominguez Y, Bok JW, Berger H, Shwab EK, Basheer A, Gallmetzer A,

Scazzocchio C, Keller N, Strauss J. **2010**. Heterochromatic marks are associated with

the repression of secondary metabolism clusters in *Aspergillus nidulans*. *Molecular*

Microbiology **76**: 1376–1386.

Ridenour JB, Möller M, Freitag M. **2020**. Polycomb repression without bristles:

facultative heterochromatin and genome stability in fungi. *Genes* **11**: 638.

Santhanam P. **2012**. Random insertional mutagenesis in fungal genomes to identify

virulence factors. In: Plant Fungal Pathogens. Springer, 509–517.

Schmit JP, Mueller GM. **2007**. An estimate of the lower limit of global fungal diversity.

Biodiversity and Conservation **16**: 99–111.

Schneper L, Düvel K, Broach JR. **2004**. Sense and sensibility: nutritional response

and signal integration in yeast. *Current Opinion in Microbiology* **7**: 624–630.

Schotanus K, Soyer JL, Connolly LR, Grandaubert J, Happel P, Smith KM, Freitag

M, Stukenbrock EH. **2015**. Histone modifications rather than the novel regional

centromeres of Zymoseptoria tritici distinguish core and accessory chromosomes.

Epigenetics & Chromatin **8**: 41.

Seidl MF, Cook DE, Thomma BPHJ. **2016**. Chromatin biology impacts adaptive

evolution of filamentous plant pathogens. *PLoS Pathogens* **12**: e1005920.

Seidl MF, Kramer HM, Cook DE, Lorencini Fiorin G, van den Berg GCM, Faino L,

Thomma BPHJ. **2020**. Repetitive elements contribute to the diversity and evolution of

centromeres in the fungal genus *Verticillium*. *mBio*: doi: 10.1128/mBio.01714–20.

Selvig K, Alspaugh JA. **2011**. pH response pathways in fungi: adapting to host-derived

and environmental signals. *Mycobiology* **39**: 249–256.

Shi-Kunne X, Faino L, van den Berg GCM, Thomma BPHJ, Seidl MF. **2018**.

Evolution within the fungal genus *Verticillium* is characterized by chromosomal

rearrangement and gene loss. *Environmental Microbiology* **20**: 1362–1373.

Slepecky RA, Starmer WT. **2009**. Phenotypic plasticity in fungi: a review with

observations on *Aureobasidium pullulans*. *Mycologia* **101**: 823–832.

Slotkin RK, Martienssen R. **2007**. Transposable elements and the epigenetic

regulation of the genome. *Nature Reviews Genetics* **8**: 272–285.

Soyer JL, El Ghalid M, Glaser N, Ollivier B, Linglin J, Grandaubert J, Balesdent

MH, Connolly LR, Freitag M, Rouxel T, *et al.* **2014**. Epigenetic control of effector

gene expression in the plant pathogenic fungus *Leptosphaeria maculans*. *PLoS Genetics*

10: e1004227.

Soyer JL, Möller M, Schotanus K, Connolly LR, Galazka JM, Freitag M,

Stukenbrock EH. **2015**. Chromatin analyses of *Zymoseptoria tritici*: methods for

chromatin immunoprecipitation followed by high-throughput sequencing (ChIP-seq).

Fungal Genetics and Biology **79**: 63–70.

Stovner EB, Sætrom P. **2019**. epic2 efficiently finds diffuse domains in ChIP-seq data.

Bioinformatics **35**: 4392–4393.

Studt L, Rösler SM, Burkhardt I, Arndt B, Freitag M, Humpf HU, Dickschat JS,

Tudzynski B. **2016**. Knock-down of the methyltransferase Kmt6 relieves H3K27me3

and results in induction of cryptic and otherwise silent secondary metabolite gene

clusters in *Fusarium fujikuroi*. *Environmental Microbiology* **18**: 4037–4054.

Tian L, Chen ZJ. **2001**. Blocking histone deacetylation in *Arabidopsis* induces pleiotropic

effects on plant gene regulation and development. *Proceedings of the National Academy*

of Sciences **98**: 200–205.

Trojer P, Reinberg D. **2007**. Facultative heterochromatin: is there a distinctive

molecular signature? *Molecular Cell* **28**: 1–13.

Voigt P, Tee W-W, Reinberg D. **2013**. A double take on bivalent promoters. *Genes &*

development **27**: 1318–1338.

Wang J, Jia ST, Jia S. **2016**. New insights into the regulation of heterochromatin.

Trends in Genetics **32**: 284–294.

Wiemann P, Sieber CMK, von Bargen KW, Studt L, Niehaus EM, Espino JJ, Huß K,

Michielse CB, Albermann S, Wagner D, *et al.* 2013. Deciphering the cryptic genome:

genome-wide analyses of the rice pathogen *Fusarium fujikuroi* reveal complex regulation

of secondary metabolism and novel metabolites. *PLoS Pathogens* **9**: e1003475.

Zannas AS, Arloth J, Carrillo-Roa T, Iurato S, Röh S, Ressler KJ, Nemeroff CB,

Smith AK, Bradley B, Heim C. **2015**. Lifetime stress accelerates epigenetic aging in an

urban, African American cohort: relevance of glucocorticoid signaling. *Genome biology*

16: 1–12.

Zhang W, Huang J, Cook DE. **2021**. Histone modification dynamics at H3K27 are

associated with altered transcription of in planta induced genes in *Magnaporthe oryzae*.

PLoS genetics **17**: e1009376.

 FIGURE LEGENDS

Figure 1*.* **Genes that are higher expressed in CZA, MS and *in planta* than in PDB**

are overrepresented in adaptive genomic regions and in H3K27me3 domains.

Fractions of total genes and genes that are higher expressed than in PDB that locate in

(A) adaptive genomic regions and (B) H3K27me3-enriched regions. Increased expression

is determined between six days of cultivation *in vitro* of *V. dahliae* in potato dextrose

broth (PDB) and 6 days of cultivation in Czapec-Dox (CZA) and half-strength Murashige

Skoog medium (MS), and infection of *Arabidopsis thaliana* (21 dpi) (*in planta*). Fractions

of higher expressed genes that reside in adaptive genomic regions or H3K27me3-

enriched regions are reported for log2-fold changes >1, 2, 3, 4, 5, 6 and 7, respectively.

H3K27me3-enrichment was determined for *V. dahliae* cultivation during six days in PDB.

Black numbers above the bars indicate numbers of genes per gene set. White numbers at

bottom of the bars indicate the numbers of genes that locate in adaptive genomic or

H3K27me3-domain per gene set. Significance of fraction differences was calculated for

each fraction of higher expressed genes in adaptive genomic regions when compared

with the fraction of total genes in adaptive genomic regions, by the two-sided two-

proportions Z-test (*: $p \leq 0.05$).

Figure 2. Lowly and non-expressed genes associate with H3K27me3. A) Whole

genome distribution of the euchromatin-associated histone modification H3K4me2, the

constitutive heterochromatin-associated histone modification H3K9me3, the facultative

heterochromatin-associated histone modification H3K27me3 and chromatin accessibility

as determined by ATAC-seq, based on chromosome 5 as an example. Genes are

indicated in light blue, transposons are indicated in red, centromere is indicated in dark

blue and adaptive genomic regions are indicated in yellow. B,C) Relative coverage of

chromatin accessibility and the histone marks H3K4me2, H3K9me3, H3K27me3 over

gene bodies (between transcription start site (TSS) and transcription end site (TES))

±800 bp of flanking sequence, grouped into quintiles based on gene expression levels

upon cultivation for six days potato dextrose broth (PDB), for (B) all genes, and for (C)

all H3K27me3-enriched genes.

Figure 3. Genetic perturbation of H3K27me3 results in induction of genes that

are transcriptionally regulated in different growth conditions. A) Fractions of

induced (log2 fold change >1, p<0.05) and repressed (log2 fold change <-1, p<0.05)

genes and genes that are not differentially expressed between wild-type and ΔSet7 that

(B) locate in H3K27me3 domains and (C-E) are higher expressed in CZA, MS and *in*

planta than in PDB.

Figure 4. Differential H3K27 methylation partially explains differential

expression. A) The number of genes located in H3K27me3 domains for *V. dahliae*

cultivated for 6 days in Czapec-Dox (CZA), half-strength Murashige Skoog medium (MS)

and Potato dextrose broth (PDB). B) Transcriptional values from RNA-seq, represented as

TPM values in each growth medium. The genes were grouped between those consistently

localized in H3K27me3 domains (1631 genes) and genes not present in H3K27me3

domains. C,D) Transcription TPM values in each growth medium for genes in H3K27me3

domains found in C) two or D) one of three tested *in vitro* growth media. Median values

of violin plots are indicated with diamonds. Significant difference of TPM values between

growth media are determined with the One-Sample Wilcoxon Signed Rank Test (*: p <=

0.05).

Figure 5. Differentially expressed genes associate with local changes in

H3K27me3 coverage. Pairwise comparisons of differentially expressed genes for *V.*

dahliae cultivated for 6 days in A,B) half-strength Murashige Skoog medium (MS) and

Potato dextrose broth (PDB), C,D) Czapec-Dox (CZA) and PDB, and E,F) MS and CZA.

A,C,E) H3K27me3 coverage over differentially expressed genes in each corresponding

growth medium. Median values of violin plots are indicated with diamonds. B,D,F) per

gene comparison of log2-fold change in transcription and genic H3K27me3 coverage.

Black horizontal lines indicate TPM $\log_2$-fold change of -2 and 2, black vertical lines

indicate H3K27me3 $\log_2$-fold change of -1 and 1. Significant difference of TPM values

between growth media are determined with the One-Sample Wilcoxon Signed Rank Test

and indicated by letters or by asterisks (*: $p \leq 0.05$).

Figure 6. *In planta*-induced genes are H3K27me3-associated under

transcriptionally repressive conditions. Average coverage of chromatin features over

genes differentially expressed between in planta and A) 6 days of cultivation in Potato

dextrose broth (PDB), B) half strength Murashige Skoog medium (MS), and C) Czapec-

Dox medium (CZA). D) overlap of genes that are higher expressed *in planta* than in PDB,

MS or CZA with genes located in an H3K27me3 domain in the corresponding growth

medium. Significant differences of H3K27me3 coverage are determined with the One-

Sample Wilcoxon Signed Rank Test (*: $p <= 0.05$).

 ADDITIONAL FILES

Fig. S1. Distribution of chromatin features over all chromosomes. Whole genome

distribution of the euchromatin-associated histone modification H3K4me2 (green line),

the constitutive heterochromatin-associated histone modification H3K9me3 (red line), the

facultative heterochromatin-associated histone modification H3K27me3 (yellow line) and

chromatin accessibility (blue line) as determined by ATAC-seq. Genes are indicated in

light blue, transposons are indicated in red, centromere is indicated in dark blue and

adaptive genomic regions are indicated in yellow.

Fig. S2. Gene expression correlates with histone modification presence and

chromatin accessibility. RPGC normalized coverage of the histone marks H3K4me2,

H3K9me3, H3K27me3 and chromatin accessibility over gene bodies (between

transcription start site (TSS) and transcription end site (TES)) ±800 bp of flanking

sequence. Each row represents a single gene, and are sorted based on their TPM value

upon cultivation for six days potato dextrose broth (PDB), with the top gene being most

highly expressed. Genes are grouped in expression quintiles as in Figure 3A.

Fig. S3. The *V. dahliae ΔSet7* deletion mutant loses H3K27me3. Histone isolations

of wild-type, *ΔDim2*, *ΔDim5*, *ΔHP1* and *ΔSet7* were tested for presence of the H3K27me3

histone modification by Western blot. Antibody against H3 was used as loading control.

Fig. S4. Genes associated with H3K27me3 in wild type *V. dahliae* are stronger

transcriptionally induced in the *ΔSet7* mutant than genes that are not

associated with H3K27me3. Log2-fold change of expression between wild type and

ΔSet7 mutant for genes associated with H3K27me3 in wild type (green) and those not

associated with H3K27me3 in wild type (red). Significant difference of log2-fold change

between gene sets are determined with the One-Sample Wilcoxon Signed Rank Test (*:

$p <= 0.05$).

Fig. S5. Distribution of H3K27me3 for *V. dahliae* cultivated in *in vitro* growth

media. Replicates of *V. dahliae* cultivated for 6 days in Potato dextrose broth (PDB), half

strength Murashige Skoog medium (MS) and Czapec-Dox medium (CZA). A) Distribution

of H3K27me3 over the *V. dahliae* genome, predicted H3K27me3 domains are indicated

as blocks below each H3K27me3 track. Adaptive genomic regions are highlighted in

yellow. B) Distribution of H3K27me3 over chromosome 5 of *V. dahliae*. Genomic regions

within H3K27me3 domains that are visibly different between growth conditions are

highlighted in grey.

Table S1. Primers used to delete and analyse the *Set7* coding sequence in *V. dahliae*.