Bayesian genome-wide analysis of cattle traits using variants with

functional and evolutionary significance

Ruidong Xiang[1,2,*], Ed J. Breen[2], Claire P. Prowse-Wilkins[1,2], Amanda J. Chamberlain[2],

Michael E. Goddard[1,2]

[1] *Faculty of Veterinary & Agricultural Science, The University of Melbourne, Parkville 3052,*

Victoria, Australia

[2] *Agriculture Victoria, AgriBio, Centre for AgriBiosciences, Bundoora, Victoria 3083, Australia.*

* Corresponding author: ruidong.xiang@unimelb.edu.au

Abstract

Context. Functional genomics studies have revealed genomic regions with regulatory and

evolutionary significance. Such information independent of association analysis may benefit

fine-mapping and genomic selection of economically important traits. However, systematic

evaluation of the use of functional information in mapping, and genomic selection of cattle

traits is lacking. Also, Single Nucleotide Polymorphisms (SNPs) from the high-density (HD)

panel are known to tag informative variants, but the performance of genomic prediction using

HD SNPs together with variants supported by different functional genomics is unknown.

Aims. We selected six sets of functionally important variants and modelled each set together

with HD SNPs in Bayesian models to map and predict protein, fat, and milk yield as well as

mastitis, somatic cell count and temperament of dairy cattle.

Methods. Two models were used: 1) BayesR which includes priors of four distribution of

variant-effects, and 2) BayesRC which includes additional priors of different functional

classes of variants. Bayesian models were trained in 3 breeds of 28,000 cows of Holstein,

Jersey and Australian Red and predicted into 2,600 independent bulls.

Key results. Adding functionally important variants significantly increased the enrichment of

genetic variance explained for mapped variants, suggesting improved genome-wide mapping

precision. Such improvement was significantly higher when the same set of variants were

modelled by BayesRC than by BayesR. Combining functional variant sets with HD SNPs

improves genomic prediction accuracy in the majority of the cases and such improvement

was more common and stronger for non-Holstein breeds and traits like mastitis, somatic cell

count and temperament. In contrast, adding a large number of random sequence variants to

HD SNPs reduces mapping precision and has a worse or similar prediction accuracy,

compared to using HD SNPs alone to map or predict. While BayesRC tended to have better

genomic prediction accuracy than BayesR, the overall difference in prediction accuracy

between the two models was insignificant.

Conclusions. Our findings demonstrate the usefulness of functional data in genomic mapping

and prediction.

Implications. We highlight the need for effective tools exploiting complex functional
datasets to improve genomic prediction.

Key words: Functional genomics, Animal breeding, Genetic mapping, Quantitative genetics

Introduction

Emerging evidence shows that genomic variants with causal roles in biology can be used to improve genomic prediction of complex traits. The biological function of genomic variants provides information independent of genotype-trait associations which are usually confounded by linkage disequilibrium (LD). Such independent information can be exploited to identify informative variants. Once identified, informative variants can be used to improve genomic prediction (Xiang, MacLeod, Daetwyler, de Jong, O'Connor, Schrooten, Chamberlain & Goddard, 2021). While the use of functional data in improving genomic mapping and prediction has been reported in humans (Amariuta, Ishigaki, Sugishita, Ohta, Koido, Dey, Matsuda, Murakami, Price & Kawakami, 2020; Weissbrod, Hormozdiari, Benner, Cui, Ulirsch, Gazal, Schoech, Van De Geijn, Reshef & Márquez-Luna, 2020), using functional data in predicting the genetic merit of animal traits has not been comprehensively examined. However, there is evidence in cattle supporting the advantage of the use of functional information in genomic mapping and prediction with the linear mixed model (Fang, Sahana, Ma, Su, Yu, Zhang, Lund & Sørensen, 2017a; Fang, Sahana, Ma, Su, Yu, Zhang, Lund & Sørensen, 2017b; Liu, Fang, Zhou, Santos, Xiang, Daetwyler, Chamberlain, Cole, Li, Yu, Ma, Zhang & Liu, 2019; Xiang, Berg, MacLeod, Hayes, Prowse-Wilkins, Wang, Bolormaa, Liu, Rochfort, Reich, Mason, Vander Jagt, Daetwyler, Lund, Chamberlain & Goddard, 2019; Xu, Gao, Wang, Xu, Liu, Chen, Xu, Gao, Zhang & Gao, 2020).

The Functional Annotation of ANimal Genomes (FAANG) consortium (Clark, Archibald, Daetwyler, Groenen, Harrison, Houston, Kühn, Lien, Macqueen & Reecy, 2020) provides many types of sequencing data indicating the functionality of genome-wide sites (examples reviewed in (Clark *et al.*, 2020)). While these public datasets await exploitation, the structure and information content of different functional datasets vary significantly. For example, we recently showed that amongst all analysed functional datasets, a set of 300,000+ sequence

variants within sites highly conserved across 100 vertebrate species had the strongest

enrichment with cattle trait heritability (Xiang *et al.*, 2019), which primarily influences

genomic prediction accuracy. Additionally, a few thousand variants affecting the

concentration of milk fat metabolites, i.e., metabolic mQTLs, also had significantly higher

variance than SNPs in the 50K panel for cattle traits. Millions of variants that change gene

expression levels (geQTLs) or RNA splicing (sQTLs) are also enriched with complex trait

QTL (Fink, Lopdell, Tiplady, Handley, Johnson, Spelman, Davis, Snell & Littlejohn, 2020;

Li, van de Geijn, Raj, Knowles, Petti, Golan, Gilad & Pritchard, 2016; Lopdell, Tiplady,

Struchalin, Johnson, Keehan, Sherlock, Couldrey, Davis, Snell & Spelman, 2017; Silva,

Fonseca, Pinheiro, Magalhães, Muniz, Ferro, Baldi, Chardulo, Schnabel & Taylor, 2020;

Xiang, Hayes, Vander Jagt, MacLeod, Khansefid, Bowman, Yuan, Prowse-Wilkins, Reich,

Mason, Garner, Marett, Chen, Bolormaa, Daetwyler, Chamberlain & Goddard, 2018).

However, recent studies showed that variants close to genes with high or specific expression

patterns had limited improvement in prediction accuracy (de Las Heras-Saldana, Lopez,

Moghaddar, Park, Park, Chung, Lim, Lee, Shin & van der Werf, 2020; Fang, Cai, Liu,

Canela-Xandri, Gao, Jiang, Rawlik, Li, Schroeder & Rosen, 2020). Another common type of

functional data is peaks from ChIP-seq for histone modifications which are enriched with

promoters and/or enhancers regulating gene activities (Carey, Peterson & Smale, 2009). Our

work showed that hundreds of thousands of variants under ChIP-seq peaks are enriched for

complex trait QTL in cattle (Prowse-Wilkins, Wang, Xiang, Goddard & Chamberlain, 2021;

Xiang *et al.*, 2019). In addition, variants within the gene coding regions are expected to have

a high impact on complex traits. However, we and others previously found coding-related

variants (around 100,000) have limited contributions to cattle trait heritability (Koufariotis,

Chen, Stothard & Hayes, 2018; Xiang *et al.*, 2019), although their use in improving genomic

prediction has not been studied.

One way to assess the information content of functional data is to compare variants

prioritised by functional data with SNPs from standard genotyping panels. We have

previously performed such assessment using the standard 50K bovine SNP chip and showed

that functional information can improve genomic prediction accuracy compared to the 50K

chip SNPs (Xiang *et al.*, 2021). However, denser panels such as the high-density (HD) SNP

chip containing ~700,000 SNPs across the genome may be able to tag many functional

elements via LD, although it is not routinely used in animal genomic evaluation. With the

development of animal breeding, the HD panel may be intensively used in the future genomic

evaluation. Therefore, it is of interest to know if functional information can provide any

advantage in genomic mapping and prediction when HD SNPs are used. Also, since causal

variants are expected to have similar phenotypic effects across different breeds, we aim to

compare the use of functionally important variants in genomic prediction across different

breeds.

In the present study, we evaluate sequence variant sets prioritised by 6 types of functional and

evolutionary data in combination with the standard HD SNPs in genomic mapping and

prediction of 6 dairy cattle traits. We train the prediction equations using the BayesR method

(Erbe, Hayes, Matukumalli, Goswami, Bowman, Reich, Mason & Goddard, 2012) which fits

a mixture of 4 distributions of variant-effects and using the BayesRC method which fits

different distributions for each functional class of variant classifications (MacLeod, Bowman,

Vander Jagt, Haile-Mariam, Kemper, Chamberlain, Schrooten, Hayes & Goddard, 2016).

Genomic predictors were trained using 28,000 cows that included 3 breeds: Holstein, Jersey

and Australian Red. Genomic estimated breeding values (gEBVs) were predicted and

validated in 2,500 Holstein, Jersey and Australian Red bulls. We compare the results of

mapping and genomic prediction across the above-described scenarios, discuss these results

and provide suggestions for future studies.

Materials and Methods

The phenotype data analysed in this study were collected by DataGene Australia (http://www.datagene.com.au/) and no further live animal experimentation was required for our analyses. A set of 28,049 Australian cows were used as the discovery population and a set of 2,567 bulls were used as the validation population. The bull phenotypes were obtained as daughter trait deviations: i.e. the average trait deviations of a bull's daughters pre-corrected for known fixed effects by DataGene. The cow phenotypes were measured on themselves. Note that these bulls and cows were not included in those 44,000+ animals used to discover functional variants (Xiang *et al.*, 2019; Xiang *et al.*, 2021; Xiang, van den Berg, MacLeod, Daetwyler & Goddard, 2020). We also checked the pedigree to make sure that bulls used in the validation population were not the sires of cows from the discovery population. Cows in the discovery set included 24,305 Holstein, 2,486 Jersey, 1,258 Australian Red. Bulls in the validation datasets contained 2,091 Holstein, 385 Jersey, 91 Australian Red. Traits considered in the analysis included protein yield (Prot), fat yield (Fat), milk yield (Milk), Mastitis (Mas), somatic cell count (Scc) and temperament (Temp).

The genotypes used in the study were imputed sequence variants based on Run7 of the 1000 Bull Genomes Project (Daetwyler, Capitan, Pausch, Stothard, Van Binsbergen, Brøndum, Liao, Djari, Rodriguez & Grohs, 2014; Hayes & Daetwyler, 2018) based on the ARS-UCD1.2 reference bovine genome (https://www.ncbi.nlm.nih.gov/assembly/GCF_002263795.1/) (Rosen, Bickhart, Schnabel, Koren, Elsik, Tseng, Rowan, Low, Zimin & Couldrey, 2020). Variants with Minimac3 (Fuchsberger, Abecasis & Hinds, 2014; Howie, Fuchsberger, Stephens, Marchini & Abecasis, 2012) imputation accuracy $R^2 > 0.4$ and minor allele frequency (MAF) > 0.005 in bulls and cows. Most bulls were genotyped with a medium-density SNP array (50K) or a high-density

SNP array and most cows were genotyped with a low-density panel of approximately 6,900

SNPs overlapping with the standard-50K panel (BovineSNP50 beadchip, Ilumina Inc). The

low-density genotypes were first imputed to the Standard-50K panel and then all 50K

genotypes were imputed to the HD panel using Fimpute v3 (Sargolzaei, Chesnais &

Schenkel, 2014; Xiang *et al.*, 2019). Then, all HD genotypes were imputed to sequence using

Minimac3 with Eagle (v2) to pre-phase genotypes (Howie *et al.*, 2012; Loh, Danecek,

Palamara, Fuchsberger, Reshef, Finucane, Schoenherr, Forer, McCarthy & Abecasis, 2016).

We aimed to test whether variant sets selected from different functional and/or evolutionary

information, in addition to the standard HD SNP panel, can be useful for genomic prediction.

Therefore, we first included a baseline set, which is 610,764 SNPs from the standard bovine

high-density panel. There were six functional and/or evolutionary variant sets: 549,007

variants under multiple ChIP-seq peaks (Kern, Wang, Xu, Pan, Halstead, Chanthavixay,

Saelao, Waters, Xiang & Chamberlain, 2021; Prowse-Wilkins *et al.*, 2021) ('ChiPseq'),

106,538 variants annotated as related to coding activities by Ensembl Variant Effect Predictor

(McLaren, Gil, Hunt, Riat, Ritchie, Thormann, Flicek & Cunningham, 2016) ('Coding'),

943,315 variants affecting RNA splicing sQTLs from 4 cattle tissues (Chamberlain, Hayes,

Xiang, Vander Jagt, Reich, Macleod, Prowse-Wilkins, Mason, Daetwyler & Goddard, 2018;

Daetwyler, Xiang, Yuan, Bolormaa, Vander Jagt, Hayes, van der Werf, Pryce, Chamberlain

& Macleod, 2019; Xiang *et al.*, 2018) ('sQTL'), 65,394 finely mapped variants with

pleiotropic effects genome-wide (Xiang *et al.*, 2021) ('Finemap80k'), 4,871 variants affecting

milk fat metabolites mQTLs (Xiang *et al.*, 2019) ('mQTL') and 317,279 conserved sites

across 100 vertebrates (Xiang *et al.*, 2019) ('Cons100w'). Note that some of these functional

variant sets were initially determined on the UMD3.1 genome and were from different cattle

populations. These sets were lifted over from the older genome to ARS-UCD1.2 and filtered

with imputation accuracy and MAF in the new cattle populations.

The model training of the above-described data used BayesR (Erbe *et al.*, 2012) and

BayesRC (MacLeod *et al.*, 2016), which are now implemented via BayesR3, with improved

efficiency using blocks. BayesR jointly models all variants together with different effect

distribution priors. BayesRC follows the same approach but in addition allows a 'C' prior

which models classes of variants. Another aim is to see whether there are differences in

genomic prediction accuracy by modelling the same variants using BayesR and BayesRC. To

aid this comparison, we combined each functional variant set with the HD variants which led

to 6 combined variant sets: 1) ChIP-seq peak tagged variants + HD SNPs ('ChIPseq_HD'), 2)

coding variants + HD variants ('Coding_HD'), 3) sQTL variants + HD SNPs ('sQTL_HD'),

4) finely mapped variants + HD SNPs ('Finemap80k_HD'), 5) mQTL variants + HD SNPs

('mQTL_HD') and 6) conserved variants + HD SNPs ('Cons100w_HD'). The average minor

allele frequency of these sets of variants were 0.22 (±0.00014) for ChiPseq_HD,

0.25(±0.0002) for Coding_HD, 0.24 (±0.0001) for sQTL_HD, 0.27 (±0.0002) for

Finemap80k_HD, 0.27 (±0.0002) for mQTL_HD, 0.23 (±0.0002) for Cons100w_HD, and

0.27 (±0.0002) for HD alone.

In single-trait BayesR, we directly model these 6 variant sets one set at a time. To create a

reference baseline, we also used single-trait BayesR to fit the HD variant set ('HD') alone. In

single-trait BayesRC, for each of the same 6 combined variant sets, we specify 2 different

variant classes: 1) Variants appeared in the functional and/or evolutionary set and 2) variants

only appeared in the HD variant set.

Both BayesR and BayesRC modelled variant effects as a mixture distribution of four normal

distributions including a null distribution, $N(0, 0.0\sigma^2_g)$, and three others: $N(0, 0.0001\sigma^2_g)$,

$N(0, 0.001\sigma^2_g)$, $N(0, 0.01\sigma^2_g)$, where σ^2_g was the additive genetic variance for the trait.

The starting value of σ^2_g for each trait was estimated using GREML implemented in the

MTG2 (Lee & Van der Werf, 2016) with a single genomic relationship matrix made of all

sequence variants. The statistical model used in the single-trait BayesR and BayesRC in was:

$$\mathbf{y} = \mathbf{Wv} + \mathbf{Xb} + \mathbf{e} \text{ (equation 1)}$$

where $\mathbf{y}$ was a vector of phenotypic records; $\mathbf{W}$ was the design matrix of marker genotypes;

centred and standardised to have a unit variance; $\mathbf{v}$ was the vector of variant effects,

distributed as a mixture of the four distributions as described above; $\mathbf{X}$ was the design matrix

allocating phenotypes to fixed effects; $\mathbf{b}$ was the vector of fixed effects, including breeds;

$\mathbf{e}$ = vector of residual errors. As a result, the effect b for each variant jointly estimated with

other variants were obtained for further analysis.

BayesRC used the same linear model as BayesR. The C component of BayesRC had two

categories $c (c = 2)$ as described above. Within each category c, an uninformative Dirichlet

prior (α) was used for the proportion of effects in each of the four normal distributions of

variant effects: $P_c \sim Dir(\alpha_c)$, where $a_c = [1, 1, 1, 1]$. α_c was updated each iteration within

each category: $P_c \sim Dir(\alpha_c + \beta_c)$, where β_c was the current number of variants in each of the

four distributions within category c, as estimated from the data.

Two metrics were evaluated for mapping results. One is the mixing proportion, i.e., the

proportion of variants with small effect $N(0, 0.0001\sigma^2_g)$, medium effect $N(0, 0.001\sigma^2_g)$

and large effect $N(0, 0.01\sigma^2_g)$ for each BayesRC run across the functional variant class and

the HD SNP class. This metric shows the information content of the two classes. The other

metric was the percentage of 50kb segments needed by the model to explain 50% of the

cumulative sum of posterior probability (PP), which indicated the mapping precision. For

each variant, PP was calculated as $1 - P_0$ where P_0 was the probability for the variant to be

within the zero-effect distribution $N(0, 0.0\sigma^2_g)$. The sum of PP across all variants estimates

the number of variants causing genetic variance in the trait. The smaller amount of genomic

segments needed to explain a cumulative sum of PP, the higher the mapping precision. We

also compared genomic prediction accuracy, defined as the Pearson correlation r between

genomic estimated breeding value (gEBV) and phenotype in the validation populations.

gEBV of the validation animals was calculated as $gEBV = Z\hat{s}$ (equation 2), where Z was a

matrix of the standardisd genotypes of animals in the validation set, and $\hat{s}$ was the vector of

variant effects from the training model. In addition, to test if adding a large number of

random variants to the HD panel can increase mapping precision and prediction accuracy, a

random set of 944,616 variants matching the size of the largest set of functional variants

(sQTL, 943,315 variants) was also selected and added to the HD panel ('Random_HD'). This

random set was analysed for BayesR, mapping precision and prediction accuracy in the same

fashion as other variant sets described above.

Results

Information content in the functional variant sets

Averaged across mixing proportions from single-trait BayesRC, we show that compared to

HD SNPs, the finely mapped variants had consistently higher enrichment with variants

showing small, medium and large effects (Figure 1). Variants within coding regions showed

higher enrichment than HD SNPs for large- and medium-effect variants. Interestingly,

mQTLs, which were variants affecting the concentration of milk fat metabolites (Benedet,

Ho, Xiang, Bolormaa, De Marchi, Goddard & Pryce, 2019; Xiang *et al.*, 2019), had lower

enrichment of small-effect variants than HD SNPs, but had higher enrichment of medium and

large-effect variants than HD SNPs.

Mapping precision

Across traits, we show that all models using functional variants, except mQTL, needed a

smaller amount of genome-wide segments to explain 50% of the cumulative sum of PP,

compared to HD SNPs (Figure 2). This means that when adding to the HD SNPs, most

functional variants increased mapping precision. In contrast, adding randomly selected

944,000 variants to HD SNPs increased the amount of genome-wide segments (by 2.82%±

0.13%) across scenarios to explain 50% of the cumulative sum of PP, compared to only using

HD SNPs. This suggested that adding random variants to HD decreases mapping precision. It

is worth noting that when using 106,538 coding variants and 65,394 finely mapped variants,

BayesRC provided a further increase in mapping precision over HD SNPs than BayesR. On

the other hand, when using 549,007 ChIP-seq tagged variants and 943,315 sQTL variants,

BayesRC had less increase in mapping precision over HD SNPs than BayesR. This could be

due to the reduced signal-to-noise ratio in large variant sets of ChIP-seq tagged variants and

sQTLs.

Genomic prediction of traits

In total, we evaluated the genomic prediction accuracy in 216 scenarios, across 6 single-trait

analysis, 6 functional categories, 4 breeds in the validation population, and 2 Bayesian

methods. Out of these 216 scenarios, 142 (66%) times, HD SNPs combined with functional

variants increased genomic prediction accuracy, compared to the prediction only using the

HD SNPs (Figure 3 and 4). In 51 out of 216 times (24%), the increase in prediction accuracy

($[r_{functional} - r_{HD}] \times 100\%$) was greater than 1%. These 51 cases were almost all accounted

for by Jersey (15/51) and Australian Red (34/51), with only 2 cases in Holstein cattle. In 29

analyses (14%), the increase in prediction accuracy over HD SNPs was greater than 2%. All

these 29 cases were for non-Holstein breeds. Amongst tested functional sets, genomic

prediction accuracy was the best when the HD variants were combined with conserved

variants (Cons100w_HD). In contrast, averaged across tested scenarios, adding randomly

selected 944,000 variants to HD had a slightly worse or no improvement in prediction

accuracy (-0.5%±0.49%) compared to only using the HD panel to predict.

As shown in Figure 3, the genomic prediction accuracy of milk production traits using HD

SNPs in Holstein cattle was already high (around 0.7) and the increases in accuracy from

functional variants were very small. However, larger increases were evident in Jersey and

Australian Red. For milk production traits, 10 out of 18 times the genomic prediction

accuracy was the most improved by conserved variants and coding variants combined with

HD SNPs, followed by finely mapped variants combined with HD SNPs (4/18), ChIP-seq

tagged variants (3/18) combined with HD SNPs. sQTL combined with HD variants had the

highest accuracy when predicting protein yield in Holstein.

As shown in Figure 4, the greatest increases in prediction accuracy for traits mastitis, somatic

cell count and temperament were again seen in non-Holstein breeds. Chip-seq peak tagged

variants combined with HD SNPs (5/18 times) and conserved variants combined with HD

SNPs (5/18 times) had the best performances in predicting mastitis, somatic cell count and

temperament.

Across all scenarios, we did not see a clear distinction in prediction accuracy between

BayesR and BayesRC in the current study. There may be some tendencies where BayesRC

had a higher accuracy than BayesR for somatic cell count, mastitis and temperament.

However, none of these differences were significant.

Discussion

Our systematic evaluations show that functional information can improve genomic mapping

and prediction of cattle traits, even when HD SNPs are used, although there were times where

HD SNPs alone still had robust performances. It is usually the less represented breeds, such

as Jersey and Australian Red who benefited the most from the improvements using functional

data. This suggests functional information can well complement HD SNPs especially in

breeds with smaller training sets. Adding randomly selected variants to the HD panel reduces

mapping precision and provided no improvement in prediction accuracy compared to only

using the HD panel. This supports that the The benefit provided via selecting variants based

on functional importance can not simply be achieved by adding more sequence variants.

We show that the biological information content which can be used to benefit mapping and/or

prediction is different between functional datasets. One of the top-performing functional

variant sets in mapping large-effect variants was the finely mapped 80,000 variants (Xiang *et*

al., 2021). This result is somewhat expected as these variants combined information from

multiple functional datasets and also included variants affecting multiple dairy cattle traits.

These finely mapped 80,000 variants outperformed the SNPs from the 50K panel in previous

evaluations (Xiang *et al.*, 2021). Furthermore, finely mapped 80,000 variants showed

enhanced enrichment of large-effect variants and improvement in mapping precision when

modelled with BayesRC. This suggests that this much more refined set of variants (chosen

because they were more relevant to the traits of interest) are likely more enriched for variants

that are more strongly associated with the trait or are causal. BayesRC would only

outperform BayesR when there is strong enrichment for QTL in at least one of the defined

classes. The other functional groups tested are not trait specific (except mQTL for fat) so

likely less enriched relative to each trait.

Previous results showed that coding-related variants did not explain a significant amount of

heritability (Koufariotis *et al.*, 2018; Xiang *et al.*, 2019). In the current study, coding-related

variants combined with HD SNPs showed enhanced enrichment with large-effect variants

and improvement in mapping precision. This implies that variants affecting protein coding

may not necessarily be good at capturing all the genetic variance of polygenic traits. The

small set of mQTLs, derived from milk fat showed strong enrichment of large-effect variants

but did not show improvement in mapping precision over HD SNPs. This set of variants

needs future investigations.

317 Unlike the results in mapping large-effect variants, for genomic prediction, the top-

318 performing variant set is the conserved variants combined with HD SNPs. The advantage of

319 adding conserved variants to HD SNPs was particularly evident when predicting somatic cell

320 count, mastitis and temperament of non-Holstein breeds (Figure 4). In fact, in these scenarios

321 HD SNPs alone did not perform so well and this leaves more room for functional variants to

322 improve the prediction accuracy. Another variant set that performed well in genomic

323 prediction is the set of ChIP-seq peak tagged variants. Again, such an advantage was the most

324 evident when predicting somatic cell count, mastitis and temperament in non-Holstein breeds.

325 Interestingly, ChIP-seq variants combined with HD SNPs appear to show some particular

326 advantages in predicting temperament. There may be some large-effect variants for

327 temperament captured by ChIP-seq peaks.

328 We found that sQTL variants combined with HD SNPs had variable performances in

329 mapping and prediction. This set did not show good performance in detecting enrichment of

330 informative variants, but overall significantly increased mapping precision over HD SNPs. In

331 genomic prediction, its performance was not impressive. This is somewhat different from

332 previous studies which showed that sQTLs are enriched with complex trait QTL(Li *et al.*,

333 2016; Xiang *et al.*, 2019; Xiang *et al.*, 2018). One explanation is that sQTLs or any other

334 eQTLs were not trait specific and are plagued by LD, which is particularly strong for

335 Holstein breeds that dominated the discovery population. Another explanation is that the

336 sample size with which we used to discover sQTLs is still small (N~120) and we should re-

337 discover and re-evaluate this set of variants when there is a larger sample size.

338 As mentioned earlier, BayesRC would only outperform BayesR when there is strong

339 enrichment for QTL in at least one of the defined classes. It would also require functional

340 information to be trait-specific. We saw advantages in BayesRC over BayesR in detecting

341 enrichment with large-effect variants using finely mapped variants, coding variants and

mQTLs. BayesRC also had advantages over BayesR in mapping precision when used with

finely mapped variants and coding variants. While these functional data are expected to be

informative, they did not provide consistent advantages for BayesRC to predict traits over

BayesR. Across all tested cases, we did not see strong advantages in BayesRC over BayesR

in genomic prediction (Figure 4). BayesRC may have some tendencies to better predict

somatic cell count, mastitis and temperament than BayesR. However, the differences were

not statistically significant. The reason behind these observations may be complex.

We know that not all variants in the functional datasets are informative and many sequence

variants are in strong LD. BayesR and BayesRC both have limitations where variants are in

very strong LD. In addition, if most causal variants are quite well tagged by HD variants and

if validation animals are highly related to the discovery animals, the room to improve

prediction accuracy is limited. Also, there may be less common variants that are not tagged

by HD SNPs, but these variants are not well imputed. Further, the optimal tissues and/or

experimental conditions to generate functional data that can be better used for improving

genomic prediction are usually not known. Therefore, the marriage between functional data

and genomic prediction is still at its very early stage.

We therefore suggest two future research directions to improve on the current results. The

first is to increase the information content in functional datasets. This can be achieved by

either increasing the sample size (biological replicates, tissues and experimental conditions)

of functional datasets or by developing better bioinformatic tools to increase the signal-to-

noise ratio in functional datasets before they can be processed by genomic prediction models.

The second direction is to improve the current genomic prediction models. Because the type

and complexity of functional data will keep growing, it will be necessary to develop more

sophisticated and flexible methods to better extract information from complex functional

data. For example, an extended BayesRC that can model quantitative biological priors,

instead of qualitative classes will be needed. Similarly, in the future we will use larger sample

sizes and diverse breeds in the training model to reduce LD between sequence variants. This

will also increase the need for Bayesian methods to be more efficient.

In conclusion, our evaluation of Bayesian genomic prediction using functional and

evolutionary information with HD SNPs provides novel insights into this emerging area. We

show that functional datasets of conserved variants, coding variants, ChIP-seq peaks and

previously finely mapped variants can improve genomic mapping and/or genomic prediction,

even when HD SNPs are used. Such improvements usually benefit non-Holstein breeds,

given the current available functional datasets. We found that by using informative biological

priors, BayesRC has significant advantages over BayesR in detecting enrichment with large-

effect variants and in mapping precision. However, the advantage of BayesRC over BayesR

for genomic prediction was not consistent. Our results highlight the need to develop better

tools to extract information from complex functional datasets which will benefit genomic

prediction in large datasets. Fusing functional genomics with genomic selection presents

great opportunities to develop new technologies that improve animal breeding and genetics.

Acknowledgments

Australian Research Council's Discovery Projects (DP160101056 and DP200100499)

supported R.X. and M.E.G. DairyBio, a joint venture project between Agriculture Victoria

(Melbourne, Australia), Dairy Australia (Melbourne, Australia) and the Gardiner Foundation

(Melbourne, Australia), funded computing resources used in the analysis. The authors also

thank the University of Melbourne, Australia for supporting this research. No funding bodies

participated in the design of the study nor analysis, or interpretation of data nor in writing the

manuscript. DataGene and CRV provided access to the reference data used in this study. We

thank Gert Nieuwhof, Kon Konstantinov and Timothy P. Hancock (DataGene) and staff from

DairyNZ for the preparation and provision of data. We thank Dr. Sunduimijid Bolormaa for

the sequence variant data imputation. We thank Drs. Iona M. MacLeod and Hans D.

Daetwyler for critical reading of the manuscript.

Conflict of interest

397 The authors declare no conflicts of interest.

References:

Amariuta, T., Ishigaki, K., Sugishita, H., Ohta, T., Koido, M., Dey, K.K., Matsuda, K.,
Murakami, Y., Price, A.L. & Kawakami, E. 2020. Improving the trans-ancestry portability of
polygenic risk scores by prioritizing variants in predicted cell-type-specific regulatory
elements. *Nature Genetics,* 52(12):1346-1354.
Benedet, A., Ho, P., Xiang, R., Bolormaa, S., De Marchi, M., Goddard, M. & Pryce, J. 2019.
The use of mid-infrared spectra to map genes affecting milk composition. *Journal of dairy*
science, 102(8):7189-7203.
Carey, M.F., Peterson, C.L. & Smale, S.T. 2009. Chromatin immunoprecipitation (chip).
Cold Spring Harbor Protocols, 2009(9):pdb. prot5279.
Chamberlain, A., Hayes, B., Xiang, R., Vander Jagt, C., Reich, C., Macleod, I., Prowse-
Wilkins, C., Mason, B., Daetwyler, H. & Goddard, M. 2018. Identification of regulatory
variation in dairy cattle with RNA sequence data.254.
Clark, E.L., Archibald, A.L., Daetwyler, H.D., Groenen, M.A., Harrison, P.W., Houston,
R.D., Kühn, C., Lien, S., Macqueen, D.J. & Reecy, J.M. 2020. From FAANG to fork:
application of highly annotated genomes to improve farmed animal production. *Genome*
Biology, 21(1):1-9.
Daetwyler, H., Xiang, R., Yuan, Z., Bolormaa, S., Vander Jagt, C., Hayes, B., van der Werf,
J., Pryce, J., Chamberlain, A. & Macleod, I. 2019. Integration of functional genomics and
phenomics into genomic prediction raises its accuracy in sheep and dairy cattle. *Proceedings*
of the Association for the Advancement of Animal Breeding and Genetics, Armidale, NSW,
Australia:11-14.
Daetwyler, H.D., Capitan, A., Pausch, H., Stothard, P., Van Binsbergen, R., Brøndum, R.F.,
Liao, X., Djari, A., Rodriguez, S.C. & Grohs, C. 2014. Whole-genome sequencing of 234
bulls facilitates mapping of monogenic and complex traits in cattle. *Nature genetics,*
46(8):858.
de Las Heras-Saldana, S., Lopez, B.I., Moghaddar, N., Park, W., Park, J.-e., Chung, K.Y.,
Lim, D., Lee, S.H., Shin, D. & van der Werf, J.H. 2020. Use of gene expression and whole-
genome sequence information to improve the accuracy of genomic prediction for carcass
traits in Hanwoo cattle. *Genetics Selection Evolution,* 52(1):1-16.
Erbe, M., Hayes, B., Matukumalli, L., Goswami, S., Bowman, P., Reich, C., Mason, B. &
Goddard, M. 2012. Improving accuracy of genomic predictions within and between dairy
cattle breeds with imputed high-density single nucleotide polymorphism panels. *Journal of*
dairy science, 95(7):4114-4129.

Fang, L., Cai, W., Liu, S., Canela-Xandri, O., Gao, Y., Jiang, J., Rawlik, K., Li, B., Schroeder, S.G. & Rosen, B.D. 2020. Comprehensive analyses of 723 transcriptomes enhance genetic and biological interpretations for complex traits in cattle. *Genome research,* 30(5):790-801.

Fang, L., Sahana, G., Ma, P., Su, G., Yu, Y., Zhang, S., Lund, M.S. & Sørensen, P. 2017a. Exploring the genetic architecture and improving genomic prediction accuracy for mastitis and milk production traits in dairy cattle by mapping variants to hepatic transcriptomic regions responsive to intra-mammary infection. *Genetics Selection Evolution,* 49(1):1-18.

Fang, L., Sahana, G., Ma, P., Su, G., Yu, Y., Zhang, S., Lund, M.S. & Sørensen, P. 2017b. Use of biological priors enhances understanding of genetic architecture and genomic prediction of complex traits within and between dairy cattle breeds. *BMC genomics,* 18(1):604.

Fink, T., Lopdell, T.J., Tiplady, K., Handley, R., Johnson, T.J., Spelman, R.J., Davis, S.R., Snell, R.G. & Littlejohn, M.D. 2020. A new mechanism for a familiar mutation–bovine DGAT1 K232A modulates gene expression through multi-junction exon splice enhancement. *BMC genomics,* 21(1):1-13.

Fuchsberger, C., Abecasis, G.R. & Hinds, D.A. 2014. minimac2: faster genotype imputation. *Bioinformatics,* 31(5):782-784.

Hayes, B.J. & Daetwyler, H.D. 2018. 1000 Bull Genomes Project to Map Simple and Complex Genetic Traits in Cattle: Applications and Outcomes. *Annual review of animal biosciences*.

Howie, B., Fuchsberger, C., Stephens, M., Marchini, J. & Abecasis, G.R. 2012. Fast and accurate genotype imputation in genome-wide association studies through pre-phasing. *Nature genetics,* 44(8):955.

Kern, C., Wang, Y., Xu, X., Pan, Z., Halstead, M., Chanthavixay, G., Saelao, P., Waters, S., Xiang, R. & Chamberlain, A. 2021. Functional annotations of three domestic animal genomes provide vital resources for comparative and agricultural research. *Nature Communications,* 12(1):1-11.

Koufariotis, L.T., Chen, Y.-P.P., Stothard, P. & Hayes, B.J. 2018. Variance explained by whole genome sequence variants in coding and regulatory genome annotations for six dairy traits. *BMC genomics,* 19(1):237.

Lee, S.H. & Van der Werf, J.H. 2016. MTG2: an efficient algorithm for multivariate linear mixed model analysis based on genomic information. *Bioinformatics,* 32(9):1420-1422.

Li, Y.I., van de Geijn, B., Raj, A., Knowles, D.A., Petti, A.A., Golan, D., Gilad, Y. & Pritchard, J.K. 2016. RNA splicing is a primary link between genetic variation and disease. *Science,* 352(6285):600-604.

Liu, S., Fang, L., Zhou, Y., Santos, D.J.A., Xiang, R., Daetwyler, H.D., Chamberlain, A.J., Cole, J.B., Li, C.J., Yu, Y., Ma, L., Zhang, S. & Liu, G.E. 2019. Analyses of inter-individual variations of sperm DNA methylation and their potential implications in cattle. *BMC Genomics,* 20(1):888.

Loh, P.-R., Danecek, P., Palamara, P.F., Fuchsberger, C., Reshef, Y.A., Finucane, H.K., Schoenherr, S., Forer, L., McCarthy, S. & Abecasis, G.R. 2016. Reference-based phasing using the Haplotype Reference Consortium panel. *Nature genetics,* 48(11):1443.

Lopdell, T.J., Tiplady, K., Struchalin, M., Johnson, T.J., Keehan, M., Sherlock, R., Couldrey, C., Davis, S.R., Snell, R.G. & Spelman, R.J. 2017. DNA and RNA-sequence based GWAS highlights membrane-transport genes as key modulators of milk lactose content. *BMC genomics,* 18(1):1-18.

MacLeod, I., Bowman, P., Vander Jagt, C., Haile-Mariam, M., Kemper, K., Chamberlain, A., Schrooten, C., Hayes, B. & Goddard, M. 2016. Exploiting biological priors and sequence

variants enhances QTL discovery and genomic prediction of complex traits. *BMC genomics,*
17(1):144.

McLaren, W., Gil, L., Hunt, S.E., Riat, H.S., Ritchie, G.R.S., Thormann, A., Flicek, P. &
Cunningham, F. 2016. The Ensembl Variant Effect Predictor. *Genome Biology,* 17(1):122.

Prowse-Wilkins, C., Wang, J., Xiang, R., Goddard, M. & Chamberlain, A. 2021. Putative
causal variants are enriched in annotated functional regions from 6 bovine tissues. *Submitted.*

Rosen, B.D., Bickhart, D.M., Schnabel, R.D., Koren, S., Elsik, C.G., Tseng, E., Rowan, T.N.,
Low, W.Y., Zimin, A. & Couldrey, C. 2020. De novo assembly of the cattle reference
genome with single-molecule sequencing. *GigaScience,* 9(3):giaa021.

Sargolzaei, M., Chesnais, J.P. & Schenkel, F.S. 2014. A new approach for efficient genotype
imputation using information from relatives. *BMC Genomics,* 15(1):478.

Silva, D.B., Fonseca, L.F., Pinheiro, D.G., Magalhães, A.F., Muniz, M.M., Ferro, J.A., Baldi,
F., Chardulo, L.A., Schnabel, R.D. & Taylor, J.F. 2020. Spliced genes in muscle from Nelore
Cattle and their association with carcass and meat quality. *Scientific reports,* 10(1):1-13.

Weissbrod, O., Hormozdiari, F., Benner, C., Cui, R., Ulirsch, J., Gazal, S., Schoech, A.P.,
Van De Geijn, B., Reshef, Y. & Márquez-Luna, C. 2020. Functionally informed fine-
mapping and polygenic localization of complex trait heritability. *Nature Genetics,*
52(12):1355-1363.

Xiang, R., Berg, I.v.d., MacLeod, I.M., Hayes, B.J., Prowse-Wilkins, C.P., Wang, M.,
Bolormaa, S., Liu, Z., Rochfort, S.J., Reich, C.M., Mason, B.A., Vander Jagt, C.J.,
Daetwyler, H.D., Lund, M.S., Chamberlain, A.J. & Goddard, M.E. 2019. Quantifying the
contribution of sequence variants with regulatory and evolutionary significance to 34 bovine
complex traits. *Proceedings of the National Academy of Sciences,* 116(39):19398-19408.

Xiang, R., Hayes, B.J., Vander Jagt, C.J., MacLeod, I.M., Khansefid, M., Bowman, P.J.,
Yuan, Z., Prowse-Wilkins, C.P., Reich, C.M., Mason, B.A., Garner, J.B., Marett, L.C., Chen,
Y., Bolormaa, S., Daetwyler, H.D., Chamberlain, A.J. & Goddard, M.E. 2018. Genome
variants associated with RNA splicing variations in bovine are extensively shared between
tissues. *BMC Genomics,* 19(1):521.

Xiang, R., MacLeod, I.M., Daetwyler, H.D., de Jong, G., O'Connor, E., Schrooten, C.,
Chamberlain, A.J. & Goddard, M.E. 2021. Genome-wide fine-mapping identifies pleiotropic
and functional variants that predict many traits across global cattle populations. *Nature
Communications,* 12(1):860.

Xiang, R., van den Berg, I., MacLeod, I.M., Daetwyler, H.D. & Goddard, M.E. 2020. Effect
direction meta-analysis of GWAS identifies extreme, prevalent and shared pleiotropy in a
large mammal. *Commun Biol,* 3(1):88.

Xu, L., Gao, N., Wang, Z., Xu, L., Liu, Y., Chen, Y., Xu, L., Gao, X., Zhang, L. & Gao, H.
2020. Incorporating Genome Annotation Into Genomic Prediction for Carcass Traits in
Chinese Simmental Beef Cattle. *Frontiers in Genetics,* 11.

Figure 1. The proportion of small-effect, medium effect and large effect variants in
functional variants and HD SNPs. The mean and standard error bars are averaged across 6
traits. ChiPseq_HD: ChIP-seq peaks + HD SNPs. Coding_HD: coding variants + HD SNPs.
mQTL_HD: mQTLs + HD SNPs. sQTL_HD: sQTL variants + HD SNPs. Cons100w_HD:
conserved variants across 100 vertebrates + HD SNPs. Finemap80k_HD: finely mapped
variants + HD SNPs.

Figure 2. Mapping precision of different models. The Y-axes represent the percentage of 50kb segments needed by the model to explain 50% of the cumulative sum of posterior probability (PP) of variants. A shorter bar means less amount of segments the model needs to explain the same amount of genetic variance, indicating higher mapping precision. Black dashed line indicates the Y value for the HD SNPs, fitted along in BayesR. ChIPseq_HD: ChIP-seq peaks + HD SNPs. Coding_HD: coding variants + HD SNPs. mQTL_HD: mQTLs + HD SNPs. sQTL_HD: sQTL variants + HD SNPs. Cons100w_HD: conserved variants across 100 vertebrates + HD SNPs. Finemap80k_HD: finely mapped variants + HD SNPs.

Figure 3. Genomic prediction accuracy (Pearson correlation coefficient, Y-axis) for
production traits, across different functional/evolutionary variant sets, breeds and Bayesian
methods. A black border and a dashed line of a bar indicate that it has the highest genomic
prediction accuracy in the panel. HOL: Holstein breed. JER: Jersey breed. RED: Prot: milk
protein yield. Fat: milk fat yield. Milk: milk yield. Australian Red. ChiPseq_HD: ChIP-seq
peaks + HD SNPs. Coding_HD: coding variants + HD SNPs. mQTL_HD: mQTLs + HD
SNPs. sQTL_HD: sQTL variants + HD SNPs. Cons100w_HD: conserved variants across 100
vertebrates + HD SNPs. Finemap80k_HD: finely mapped variants + HD SNPs.

Figure 4. Genomic prediction accuracy (Pearson correlation coefficient, Y-axis) for mastitis,
somatic cell count and temperament across different functional/evolutionary variant sets,
breeds and Bayesian methods. A black border and a dashed line of a bar indicate that it has
the highest genomic prediction accuracy in the panel. HOL: Holstein breed. JER: Jersey
breed. RED: Australian Red. Mas: mastitis. Scc: somatic cell count. Temp: temperament.
ChiPseq_HD: ChIP-seq peaks + HD SNPs. Coding_HD: coding variants + HD SNPs.
mQTL_HD: mQTLs + HD SNPs. sQTL_HD: sQTL variants + HD SNPs. Cons100w_HD:
conserved variants across 100 vertebrates + HD SNPs. Finemap80k_HD: finely mapped
variants + HD SNPs.