Title

Visualization of lake succession as indicated by macrophyte species

composition and trophic levels in lakes subdivided by volcanic eruption at

Akan Caldera, Hokkaido, Japan

Short title

Visualization of lake succession in caldera lakes

Author list

Isamu Wakana[1*], Yasuro Kadono[2], Jotaro Urabe[3], Yuki Tamura[4], Yoshifusa

Suzuki[5], Hiroyuki Yamada[6], Yoichi Oyama[7], Keiji Wada[8], Takeshi Hasegawa[9],

and Masashi Ohara[10]

[1]Kushiro International Wetland Center, Kushiro, 085-8505 Japan

[2] Graduate School of Science, Kobe University, Kobe, 657-8501 Japan

[3]Graduate School of Life Science, Tohoku University, Sendai, 980-8577 Japan

[4]Kankyo Consultant Co., Ltd., Kushiro-cho, 088-0606 Japan

[5]Kaiyoutansa Co., Ltd., Otaru, 047-0036 Japan

[6]Graduate School of Agriculture, Hokkaido University, Sapporo, 060-8589

Japan

[7] Faculty of Life and Environmental Sciences, University of Tsukuba,

Tsukuba, 305-0006 Japan (Present address: Kushiro City Board of

Education, Kushiro, 085-0467 Japan)

[8]Hokkaido University of Education, Asahikawa Campus, Asahikawa,

070-0825 Japan

[9]Department of Earth Science, Ibaraki University, Mito, 310-0056 Japan

[10]Graduate School of Environmental Science, Hokkaido University, Sapporo,

060-0810 Japan

* Corresponding author

E-mail: akantbe1@seagreen.ocn.ne.jp

Abstract

It is not feasible to continuously observe ecological succession in lakes
because of the long time-scales generally involved. Thus, the process has
been inductively deduced by comparing many lakes with different succession
states, or indirectly simulated by tracking studies of smaller water bodies,
experiments using microcosms or mesocosms, and reconstruction of lake
history by sediment analysis. However, the reality of succession processes in
large lakes with slow succession is not well understood, and new approaches
are needed. Theoretically, in a group of large and small lakes of similar ages
and with similar initial and watershed environments, the rate of nutrient
accumulation in each lake depends on the ratio of watershed area to lake size,
and the lakes are predicted to evolve to different trophic levels over time.
Here, we tested this hypothesis on the 10 lakes of varying sizes in Akan
Caldera, Japan, which were formed thousands of years ago by fragmentation
due to volcanic eruptions within the caldera. Topographic and water quality
studies showed that the ratio of accumulated watershed area to lake size
(area and volume), expressed logarithmically, had a positive linear
regression with the total phosphorus concentration, an indicator of trophic

level. The trophic levels of the lakes were diverse, including oligotrophic,

mesotrophic, and eutrophic types in the traditional "lake type" classification

based on total phosphorus and chlorophyll-a concentrations. Furthermore,

21 species of aquatic macrophytes were observed by a diving survey, and the

plant species composition was classified into five groups corresponding to the

trophic status of the lakes, indicating a conventional "hydrarch succession".

The diversity of water quality and aquatic vegetation in a group of lakes with

similar origins paves the way for new comparative studies of lakes, including

large lakes.

Introduction

Ecological succession within lakes is not only a major component of

freshwater ecology and limnology, but also a platform for diagnosis and

management of worldwide deteriorating aquatic environments resulting

from human activity since the 20th century [1-7]. In general, lake succession

proceeds thusly: when a lake is first formed the water is oligotrophic with a

paucity of biota, but the trophic level subsequently increases, as trophic

substances inflow from the watershed, and the lake basin shallows from

accumulation of sediments. The diversity of biota and biomass increase or

fluctuate throughout this process, and eventually the lake attains the state

of bog or marsh [1-4, 8]. Historically, Forel, the "father" of limnology [4],

explained this transition as analogous to human ageing [9], and this concept

has been widely accepted due to both intuitive and experiential evidence [1, 2,

4, 10]. Forel also pointed out that it is impossible to follow a normal lake's

evolution, because of the immense time-scales required for a lake to

disappear by sedimentation [9]. Therefore, this general picture of long-term

succession has been indirectly formed by comparing many lakes with

different trophic conditions [10-16], follow-up survey of small-scale dams or

reservoirs [2, 17], experimental microcosms and mesocosms [4, 18-22], and

historical reconstruction of sedimentation processes [5, 8, 23]. Despite these

efforts, however, the perception of ecological succession in large lakes still

relies on many assumptions, due to their long ageing process [2, 18, 24-26].

Is there any other approach to visualize lake succession processes,

including within large lakes? In principle, the rate of eutrophication, a major

factor in ecological succession, is determined by lake size, watershed area

and trophic condition of the watershed [4]. Therefore, if trophic conditions

are equal among different watersheds, the eutrophication rates, i.e., the

rates of nutrient increase, should vary depending on the ratio of the

watershed area to the lake size in each lake. In addition, given a group of

lakes with similar formation time and initial environment, the

concentrations of nutrients are expected to change with a positive linear

regression between the ratios of watershed area to lake size. As described,

such lakes can be considered as a series that evolved into different trophic

levels, providing a new approach to visualizing lake succession. In practice,

however, no study has yet demonstrated this relationship, because nutrient

loading from the watershed is generally influenced by indigenous variables,

such as land form, soil, erosion, local climate (temperature and rainfall),

vegetation, and land use (farmland, factory and urbanization) in addition to

the time effect [1-4, 10]. We therefore offer insight into this problem by

comparing a group of lakes within a large caldera. The lakes of Akan Caldera,

Hokkaido, Japan, were formed by volcanic eruptions thousands of years ago,

which divided a huge caldera lake into several lakes of varying sizes [27-29].

These lakes share a similar terrestrial environment within the caldera. In

contrast, the trophic levels in these lakes are known to be diverse, ranging

from oligotrophic to eutrophic [30-33], and may show a series of "lake

successions" or "lake types" in the natural ecological succession of lakes [2, 5,

10]. In this paper, by studying the topography and water quality of Akan

Caldera Lakes, we tested the hypothesis that lakes with a lower ratio of

watershed area to lake size are oligotrophic, while lakes with a higher ratio

are eutrophic.

Furthermore, the lake ecological succession is also displayed as

"hydrarch succession" according to changes in aquatic vegetation [2, 5, 10,

34]. The macrophytes, which are major players in this process, are known to

change their species composition in response to changes in trophic levels,

especially in eutrophic lakes affected by anthropogenic activity [35-38], but

the natural hydrarch succession in large lakes is not fully understood, as is

the case with lake succession or lake type. Through preliminary surveys

across the Akan Caldera Lakes, we have observed that each lake has a

unique macrophyte species composition, so we also examined the prediction

that the species composition may correspond to the trophic level.

Materials and methods

Study area

Akan Caldera is situated at the southern end of the Akan-Shiretoko Volcanic

Chain, a volcanic region in eastern Hokkaido, Japan [29]. The outer shape of

the caldera is oblong (24 km east-west × 13 km north-south) [29], and a

"central cone", Mt. O-akan, rises at the center of its inner basin (Fig 1).

Within the caldera there are lakes and marshes of various sizes surrounding

Mt. O-akan (Fig 1). Lake watersheds are isolated from external input by the

caldera wall [39, 40], and the water systems, connected by rivers or

underground flow [27, 40, 41], are roughly divided north-south and join at

the southern foot of Mt. O-akan, where they then discharge through Akan

River, the notch in the south caldera wall (Fig 1A).

Fig 1. Map and landscape of Akan Caldera Lakes. (A) The watershed of Akan

Caldera is isolated by a caldera wall shown by red dotted lines, but the western

caldera wall is obscured by volcanos erupting after the formation of the caldera.

Ten lakes surround Mt. O-akan within the caldera, and are connected by rivers

and underground flow from water systems to the north and south. The water

flows converge south of Mt. O-akan and discharge is through the Akan River.

The map was generated by Kashmir 3C [42], a map software. (B) Land area is

mostly covered with subarctic forests except for a town south of Lake Akan,

Akanko-onsen.

Large lakes are distributed from west to northeast of Mt. O-akan and

smaller lakes are localized from south to east (Fig 1A). This peculiar lake

arrangement is a result of the formation history of the caldera and Mt.

O-akan. The Akan region has witnessed more than ten large volcanic

eruptions in the last 1.5 million years, and the present oblong-shaped

caldera was formed by the largest eruption of 200,000 years ago [27, 29].

After the last large eruption (150,000 years ago), a huge lake, "Ko-akanko"

(ancient Lake Akan), was generated in the caldera [27-29]. The lake was

narrowed by post-caldera volcanic activity in the southwest part of the

caldera, and by 110,000 years ago landform of the inside caldera wall was

almost completed [27-29]. A minor eruption of Mt. O-akan occurred 13,000

years ago slightly southeast of the center of the caldera, and it stopped when

the lava flow reached the caldera wall, so that Ko-akanko was separated into

large and small basins [27-29].

The developmental history of Mt. O-akan also shows that Pond

Hyotan and Pond Junsai of the southern water system were first divided

from Ko-akanko, and other lakes of the northern water system were formed

5,000 to 2,500 years ago [27-29]. The depth charts of the large lakes, Akan,

Panke and Penke, show the remains of valleys on the bottom of the inside

caldera wall extending as far as the base of Mt. O-akan [27, 39, 41]. This

suggests that the water level of Ko-akanko was extremely low or the basin

was exposed by discharge through the notch before the eruption of Mt.

O-akan [27]. Therefore, the lakes of the northern water system are thought

to have formed by re-flooding after damming by Mt. O-akan, but the timing

of lake formation and the developmental processes are not fully understood.

Akan Caldera occupies a part of Akan-Mashu National Park

(designated in 1934), and is mostly covered with subarctic forests except for a

town on the south side of Lake Akan, Akanko-onsen (Fig 1B) [30]. Only Lake

Akan has been developed as a sightseeing area due to the presence of

Marimo, *Aegagropila linnaei*, a ball-shaped green alga designated a

Japanese natural treasure [30, 43]. Since the 1950s, the increase in tourism

has resulted in eutrophication from sewage discharge. This continued until

the 1980s when public sewage treatment service was provided [44].

Topography of lakes and watersheds

Lake area and boundary length were calculated using ARCGIS10 (Esri

Japan Co.) based on the 1/25,000 numerical map data of the Geographical

Survey Institute, Japan. Land area includes island areas.

Land watershed area was computed using the DEM10m data of the

Geographical Survey Institute. After altitude data were changed into raster

(altitude grid), subtle undulations were removed by the Fill tool, and bearing

azimuth of flow was computed by the Flow Direction tool (north; 64,

northeast; 128, east; 1, southeast; 2, south; 4, southwest; 8, west; 16 and

northwest; 32). Accumulation value (number of cells accumulated toward the

direction of flow) computed by the Flow Accumulation tool was extracted at

accumulation values more than 30000 (sl30000) and more than 200 (sl200)

by the Reclass command, and each watershed (ws30000, ws200) was

computed by being grouped by every feeder of sl30000 and sl200 by the

Stream Link tool. Finally, the Ws raster was converted into the polygon, and

Ws30000 and ws200 were manually divided as a watershed for each lake,

observing the DEM.

Lake volume and mean depth were calculated based on lake charts.

Depth sounding for chart drawing was performed in autumn of 2014 in seven

lakes, excepting Lakes Akan, Panke and Penke which already have lake

charts. The depth of the entire lake was uniformly measured by a GPS fish

finder (Lowrance HDS-8, Navico) on a motor boat, and the depth-sounding

data were converted into contour drawings by chart drawing software,

Reefmaster (ReefMaster Software Ltd.). Elevation of the lake surface was

obtained by the GNSS survey. Residence time was calculated by dividing

lake volume by inflow obtained through multiplying annual rainfall at 1,200

mm [30] by the accumulated watershed area (see Statistical analysis, below).

Water quality

Measurements of physical and chemical variables and collection of lake

water were performed twice in ten lakes of Akan Caldera from October to

November 2013 and in July 2014. Secchi depth, water temperature (Temp),

electrical conductivity (EC) and pH were directly recorded using a Secchi

disk and portable sensors at the center of each lake. Water was collected at

the same point using 2 l polycarbonate bottles and taken immediately to the

laboratory. Dissolved oxygen (DO) and chemical oxygen demand (COD) were

measured by titration with standard sodium thiosulfate solution and

potassium permanganate solution, respectively [45]. Total nitrogen (TN) and

total phosphorus (TP) were measured by an auto analyzer (AACS-II,

Bran+Luebbe Ltd.) [46]. Additionally, an aliquot of the water sample was

filtered onto Whatman GF/F glass fiber filters, and suspended solids (SS)

measured gravimetrically after drying at 110 °C for two hours [45].

Chlorophyll-a (Chl-a), concentrated onto a Whatman GF/F glass filter, was

quantified with a spectrophotometer (UV-1600, Shimadzu Co. Ltd.) after

extraction using methanol (100%) [47]. The filtrate was used for measuring

dissolved organic carbon (DOC) with a TOC meter (TOC VC, Shimadzu Co.

Ltd.) [48].

Macrophyte survey

Macrophytes were surveyed in 10 lakes of Akan Caldera by SCUBA diving or

snorkeling from October to November 2013 and in July 2014. The maximum

depth surveyed was 18 m in Lake Panke.

Statistical analysis

A total of 31 variables were analyzed (S1 Table). The lake topographic factors

were 11 variables: elevation, boundary length, lake area (LA), shore line

development, maximum depth, mean depth, lake volume (LV), residence

time, land watershed area (LWA), total watershed area (TWA) which is the

sum of LA and LWA, and accumulated watershed area (AWA) which is the

sum of the TWA above a given lake. LA and LWA correspond to the aquatic

and terrestrial watershed areas specific to individual lakes, respectively, and

AWA indicates the total watershed area including the upstream portion. Six

variables reflected inflow and outflow of materials [3]: LWA-LA ratio,

LWA-LV ratio, TWA-LA ratio, TWA-LV ratio, AWA-LA ratio and AWA-LV

ratio. Water chemistry parameters consisted of ten variables: Temp, pH, DO,

EC, SS, Chl-a, DOC, COD, TN and TP, and the macrophyte communities

were defined by four variables: number of submerged species, floating-leaved

species, free-floating species and total species. Secchi depth was excluded

from the analysis due to missing data (S1 Table).

To explore the relationship among topography, water quality and

macrophyte community, a correlation matrix of these standardized data was

made (S2 Table) and variables with strong correlation coefficients ($|r| \geq 0.7$)

were investigated in detail. In lakes where macrophytes occurred, the

presence/absence of individual species was converted into 1/0 data, and a

cluster analysis was conducted by the Ward method using the Euclid

distance. The BellCurbe (Social Survey Research Information), an add-in

program for Excel, was used for the analysis.

Results and discussion

Water quality and lake types

The water quality of the 10 lakes was diverse (S1 Table), and classified as

oligotrophic (2 lakes), mesotrophic (5 lakes) and eutrophic (3 lakes) according

to Chl-a and TP concentrations (Fig 2A) [7]. Among these lakes, Pond Junsai

(eutrophic) had brownish water with high TN (0.438 mg l^{-1}), DOC (7.1 mg l^{-1})

and Chl-a concentrations (12.5 µg l^{-1}) (S1 Table), indicating dystrophic

properties [5, 21].

Fig 2. Aspects of trophic status in Akan Caldera Lakes. (A) A variety of lake

types categorized by Chlorophyll-a and total phosphorous (TP) according to the

Organization for Economic Co-Operation and Development (OECD) [7]. Even

though the lakes share the same origin and most of them were formed at the

same time, the lake types are diversified into oligotrophic, mesotrophic and

eutrophic. Additionally, Pond Junsai (red square) has dystrophic characteristics

(S1 Table). (B) Relationship between ratio of accumulated watershed area to

lake volume (AWA/LV) and TP. The AWA/LV shows significant correlation with

TP thought to be a parameter of eutrophication rate, but only Lake Akan (red

square), subjected to anthropogenic eutrophication in the past, shows an

unusually high TP.

Among ten water quality variables and 17 topographic characters,

strong ($|r| \geq 0.7$) and significant ($p < 0.05$, by test of no correlation)

correlation coefficients were found between: TP and AWA, TP and AWA-LV

ratio (AWA/LV), EC and AWA, EC and AWA-LA ratio (AWA/LA), EC and

AWA/LV, pH and shoreline development, DO and AWA/LV, and Temp and

elevation (Table 1). Although correlation coefficients between TP and

AWA/LA, and DO and AWA/LA did not reach 0.7 and -0.7, respectively, both

were statistically significant (Table 1).

Table 1. Combinations indicating strong correlation coefficients (r)

between water quality variables and topographic characters.

Water quality	Topography			
	Shoreline development	AWA	AWA/LA	AWA/LV
TP	$-0.052^{n.s}$	0.820^{**}	0.665^{*}▯	0.751^{*}▯
EC	$0.236^{n.s}$	0.991^{***}	0.720^{*}▯	0.726^{*}▯
pH	0.885^{***}	$0.283^{n.s}$	$-0.149^{n.s.}$	$-0.205^{n.s.}$
DO	$0.517^{n.s}$	$-0.395^{n.s}$	-0.692^{*}	-0.763^{*}
Temp	-0.924^{***}	$0.610^{n.s}$	$0.338^{n.s}$	$0.386^{n.s}$

The original correlation matrix is available in S2 Table. TP: total phosphorus, EC:

electrical conductivity, DO: dissolved oxygen, Temp: temperature, AWA:

accumulated watershed area, AWA/LA: AWA-lake area ratio, AWA/LV: AWA-lake

volume ratio, ***: $p < 0.001$, **: $p < 0.01$, *: $p < 0.05$, n. s.: not significant, ▯: Lake

Akan as an outlier.

When two-dimensional plots were drawn on these 10 combinations,

Lake Akan was discriminated as an outlier in TP and AWA/LA, TP and

AWA/LV, EC and AWA/LA, and EC and AWA/LV (Table 1, Fig 2B). As

previously noted, Lake Akan was subject to anthropogenic eutrophication in

the second half of the 20th century [44]. We thus estimated phosphorus

concentration before eutrophication. The oldest P_2O_5 data, 0.010 mg l^{-1},

measured in Lake Akan in 1931 [49] was calculated at 0.004 mg l^{-1}

phosphorus by the conversion formula which divides the value of P_2O_5 by

2.29 [10]. This result was close to the regression lines of TP and AWA/LA (r =

0.787, $p < 0.05$), and TP and AWA/LV (r = 0.881, $p < 0.01$) drawn for nine

lakes except for Lake Akan (Fig 3A). The difference (0.022 mg l^{-1}) from our

observed data (0.026 mg l^{-1}, S1 Table) is likely part of the increase from

eutrophication, suggesting that before eutrophication, AWA/LA and AWA/LV

of the lakes including Lake Akan had a tighter linear regression relationship

with TP. Conversely, without including the outlier in the case of TP and AWA,

the resulting plot when Lake Akan TP data were replaced by the above 0.004

mg l^{-1} drew away from the regression line (r = 0.795, $p < 0.05$, Fig 3B).

Fig 3. Estimation of phosphorus concentration in Lake Akan before

eutrophication. Because Lake Akan was subject to anthropogenic

eutrophication in the second half of the 20th century [44], phosphorus

concentration before eutrophication was estimated based on data from 1931 [49].

Each regression line is drawn for nine lakes (blue circles) except for Lake Akan.

In (A) relationship between ratio of accumulated watershed area (AWA) to lake

volume and total phosphorus (TP), the 1931 estimate of phosphorus

concentration for Lake Akan (light red square) is significantly lower than the

2013–14 TP data (dark red square), and is close to the regression line.

Conversely, in (B) the relationship between AWA and TP, the same 1931

estimate is distant from the regression line.

A strong correlation was observed between EC and TP ($r = 0.821$, $p <$

0.01, S2 Table). The regression formula ($y = 819.023x - 2.343$) estimated the

EC of Lake Akan at 0.933 mS m^{-1}, near the regression lines of EC and

AWA/LA ($r = 0.973$, $p < 0.001$,) and EC and AWA/LV ($r = 0.979$, $p < 0.001$) for

the nine lakes when TP was 0.004 mg l^{-1} (Fig 4A). By contrast, the calculated

result of Lake Akan EC drew away from the regression between EC and AWA

without the outlier ($r = 0.988$, $p < 0.001$, Fig 4B). This suggests that the

involvement of AWA may be invalid, along with the results of similar

substitution of that between TP and AWA, shown in Fig 3B.

Fig 4. Estimation of electrical conductivity (EC) in Lake Akan before

eutrophication. Because phosphorus concentration in Lake Akan was

estimated to be significantly lower before eutrophication (Fig. 3), EC showing a

high correlation with total phosphorus (TP) was also examined. The EC for 1931

was calculated at 0.933 mS m^{-1} by substituting the estimated phosphorus

concentration for 1931 (0.004 mg l^{-1}) into the regression formula for EC and TP.

Each regression line in the graphs is drawn for nine lakes (blue circles) except

for Lake Akan. In (A) relationship between ratio of accumulated watershed area

(AWA) to lake volume and EC, the 1931 estimate of EC for Lake Akan (light red

square) is significantly lower than the 2013–14 EC data (dark red square), and is

close to the regression line, similar to the case of phosphorus in Fig. 3.

Conversely, in (B) the relationship between AWA and EC, the same 1931

estimate is distant from the regression line, as in the case of phosphorus.

The regression line of pH and shoreline development was likewise

drawn without including the outlier (Fig 5). Because shoreline development

is proportional to length of the horizontal littoral zone, the correlation with

pH was presumed to relate to consumption of CO_2 through hydrophytic

photosynthesis [2, 4, 50].

Fig 5. Relationship between shoreline development and pH in Akan

Caldera Lakes. Despite the lakes having a rich diversity of lake basin sizes and

water quality, including trophic levels (S1 Table), shoreline development, the

degree of shoreline flexion, is strongly correlated with lake water pH.

The regression lines of DO and AWA/LA, and DO and AWA/LV were

drawn with a slight negative slope due to the specific low DO data for Lake

Jiro (Fig 6), and may not indicate an environmental gradient. Lake Jiro has

no inflow and outflow rivers (Fig 1A), and the water appears to be supplied

through underground flow from the upstream Lake Akan [51]. However, in

addition to DO, Chl-a, DOC, and COD in Lake Jiro were the lowest

concentrations among the Akan Caldera Lakes, and TP was the highest (S1

Table). Furthermore, a portion of the lake surface of Lake Jiro does not

freeze in winter [51]. These results suggest that other water sources, such as

groundwater, may be involved in water formation in Lake Jiro.

Fig 6. Relationship between ratios of accumulated watershed area to lake

volume and dissolved oxygen (DO). Despite high correlation coefficients

(Table 1), the regression line may not indicate an environmental gradient of DO,

because its slight negative slope results from the uniquely low DO data for Lake

Jiro (red square) with a relatively heterogeneous water quality (S1 Table).

Finally, regardless of the lake's size, water temperature decreases

with elevation upstream, and the difference in water temperature was 3.5 °C

for an elevation difference of about 150 m (Fig 7).

Fig 7. Relationship between elevation and water temperature in Akan

Caldera Lakes. Water temperature is lower in lakes located upstream at higher

elevations, regardless of lake size.

Distribution and species composition of macrophytes

We recorded 21 species of macrophytes in total (excluding emergent plants

and macro algae) in 7 lakes, while no macrophyte species were observed in 3

lakes without inflow rivers (S1 Table). The correlation matrix of the number

of macrophyte species was strong and significant with the following

variables among the above-mentioned lake topographic characters (S2

Table): boundary length (r = 0.720, p < 0.05), shoreline development (r =

0.703, p < 0.05), maximum depth (r = 0.924 < 0.001), mean depth (Fig 8, r =

0.928, p < 0.001) and residence time (r = 0.921, p <0.001). Several previous

studies on relatively shallow or small lakes and ponds reported that the

number of macrophyte species is correlated with lake area and that

MacArthur and Wilson's "theory of island biogeography", which theorizes

that larger islands are more biodiverse than smaller islands, is applicable

here [52-59]. However, in our study no significant correlation was found

between lake area and the number of macrophyte species (S2 Table). The

boundary length and the shoreline development are parameters affecting

horizontal length of the littoral zone, and the magnitude of the maximum

and mean depths and the residence time related to depth and volume of lake

basin contribute to expansion of the vertical littoral zone under conditions of

greater water clarity. Therefore, the number of macrophyte species in the

large lakes of Akan Caldera is presumed to be more closely related to the

littoral zone area than the lake area.

Fig 8. Effect of lake size on number of aquatic macrophyte species in Akan

Caldera Lakes. In total, 21 species were observed in 7 lakes. The number of

species in each lake shows high correlation with mean depth as well as with

maximum depth, boundary length, shoreline development and residence time

(S2 Table). Pond Junsai (red square) has dystrophic water quality (S1 Table) and

is dominated by floating-leaved and free-floating plants, whereas most other

lakes have a submerged plant community (S1 Table).

The two-dimensional plots of the number of macrophyte species and

the above five topographic characters showed that Pond Junsai, a dystrophic

lake, was designated an outlier (Fig 8). Furthermore, when the individual

species was classified as submerged, floating-leaved and free-floating, the

correlation coefficient was significantly greater with the number of species of

submerged plants (S2 Table). The number of species of floating-leaved and

free-floating plants showed significant and strong correlation with the water

quality parameters Chl-a, DOC, COD and TN (S2 Table), with most of these

species localized in Pond Junsai. Thus, we conducted a cluster analysis to

understand species composition of macrophytes in each lake (Fig 9). The

cluster was firstly divided into two groups: 6 species of floating-leaved and

free-floating plants of Pond Junsai, and 14 submerged and 1 floating-leaved

plants in the other lakes. Pond Junsai contained some indicator species of

the dystrophic water: *Brasenia schreberi* (Fig 10), *Nuphar pumila* var.

pumila, Nymphaea tetragona var. *tetragona* and *Utricularia macrorhiza* [60].

The remaining species recorded from the other lakes were classified as

oligotrophic, mesotrophic, oligo-mesotrophic and oligo-meso-eutrophic,

following the lake type based on the trophic status shown in Fig 2A.

Ranunculus nipponicus var. *submersus* (Fig 10), *Potamogeton alpinus* and

Isoetes asiatica, typically found in more pristine water [60-62], were

designated oligotrophic. *Myriophyllum spicatum* (Fig 10), *Hydrilla*

verticillata, Potamogeton compressus, Potamogeton pectinatus and

Ceratophyllum demersum, occurring in more eutrophic settings [38, 60, 61],

were designated oligo-mesotrophic. *Potamogeton crispus* (Fig 10)*,*

distributed in a variety of water environments [60], was designated

mesotrophic.

Fig 9. Cluster analysis of distribution and species composition of aquatic

macrophytes in Akan Caldera Lakes. The tree diagram was drawn by the

Ward method using Euclid distance based on the 1/0 data converted from the

presence/absence data for the seven lakes where macrophytes occurred.

According to the lake habitat types (Fig. 2A), clusters are classified into five

vegetation types: oligotrophic, mesotrophic, oligo-mesotrophic,

oligo-meso-eutrophic, and dystrophic.

Fig 10. Determination model for macrophyte species composition

corresponding to trophic types in Akan Caldera Lakes. Individual

macrophyte species belong to any one of the five occurrence types of

macrophytes, accompanied by inherent trophic type(s) and range. Macrophytes

of oligotrophic-, mesotrophic- and dystrophic-occurrence types are only

distributed within oligotrophic-, mesotrophic- and dystrophic-type lakes,

respectively. In contrast, within oligo-mesotrophic- and

oligo-meso-eutrophic-occurrence types, macrophytes are able to occur widely in

oligotrophic/mesotrophic- and oligotrophic/mesotrophic/eutrophic-type lakes.

Accordingly, species composition of macrophytes in oligotrophic- and

mesotrophic-type lakes is determined by combination of the occurrence types,

oligotrophic/oligo-mesotrophic/oligo-meso-eutrophic and

mesotrophic/oligo-mesotrophic/oligo-meso-eutrophic. In eutrophic and

dystrophic lakes, oligo-meso-eutrophic- and dystrophic-occurrence types are

distributed, respectively.

As mentioned above, prior to anthropogenic eutrophication in the

first half of the 20th century, the phosphorus level in Lake Akan appears to

have been much lower than currently (Fig 3). Thus, the current aquatic flora

was also compared with results from the oldest known vegetation survey

conducted in 1897 [63]. Of the ten species of macrophytes in Lake Akan

observed in our study, nine species were also found in 1897, except for *P.*

crispus, classified as mesotrophic. However, the two oligotrophic species *R.*

nipponicus var. *submersus* and *I. asiatica* in the old list were not found.

These results suggest a shift to a more eutrophic vegetation type.

Uniqueness of Akan Caldera Lakes

To summarize, although the lakes of Akan Caldera have the same origin,

after being divided by a volcanic eruption they have developed into a series of

oligotrophic, mesotrophic, eutrophic and dystrophic lakes (Fig 2A). The

trophic level of each lake indicated by TP was closely related to the ratio of

watershed area (AWA) to lake size (LA and LV) (Fig 2B). These results

strongly suggest the possibility that the rate of eutrophication was different

among the lakes, and we see the various stages of lake succession in progress

in this system. However, the observed TP of Lake Akan and its downstream

neighbors, Lake Jiro and Lake Taro, are assumed to have been impacted by

anthropogenic eutrophication in the past (Fig 3A). Furthermore, the

formation history of the Akan Caldera Lakes suggests that Ponds Hyotan

and Junsai are significantly older than the other lakes [27-29]. As shown in

Fig 2A, Chl-a levels in these two ponds are specifically higher than those of

other lakes, which might be due to differences in time of formation. To

understand the long-term changes in nutrient loading, including TP, and in

primary production, the correct time of formation and subsequent

eutrophication history of each lake should be clarified through research on

lake sediment, etc. In addition, the linear regression of the relationship

between TP and AWA/LA and AWA/LV (Fig 2B) suggests that the indigenous

environmental variables of the watersheds may vary little. However, in

reality, the geology and vegetation within the Akan Caldera are not uniform

[27, 30], and further research is needed to determine the actual phosphorus

loading from the watersheds.

Macrophyte species composition varied among lakes, driven by lake

trophic conditions (Fig 9), and is indicative of aquatic plant succession, or

"hydrarch succession". Many species of macrophytes are classified into a

variety of "types" according to environmental characteristics of habitats,

including trophic level [56, 61, 64-68]. Schneider and Melzer [68], for

example, proposed seven categories ranging from oligotrophic to eutrophic

and even polytrophic types. Meanwhile, Lacoul and Freedman [61]

simplified into three categories: oligotrophic, eutrophic, and general types,

based on the opinion that many macrophytes generally have a broad

ecological range, occurring over wide trophic levels, while other species have

a narrower distribution. In our study, however, all of the observed species

belonged to only one of the five occurrence types, with a species-specific

range of trophic level (Fig 10). Thirteen species were distributed among

oligotrophic, mesotrophic and dystrophic lake types, respectively, and eight

species occurred in the oligotrophic and mesotrophic, and the oligotrophic,

mesotrophic and eutrophic lakes with wider trophic ranges. These results

explain how the species composition of macrophytes in the Akan Caldera

Lakes with different trophic types is determined by the combination of

species with different trophic requirements. Importantly, we noted

vegetation changes in Lake Akan due to anthropogenic eutrophication: this

was characterized by disappearance of oligotrophic-type species, appearance

of mesotrophic-type species, and survival of oligo-mesotrophic and

oligo-meso-eutrophic-type species, when the trophic level of Lake Akan

shifted from oligotrophic to mesotrophic as shown in Fig 10. This might be

the first example to clearly and simply illustrate the replacement of species

in an aquatic plant community caused by trophic level change. Further

investigation is necessary to test whether this is a universal phenomenon in

limnology.

Trophic levels are not the only factor known to affect macrophyte

distribution and species composition. At a smaller scale, the following

physical factors have been important in producing environmental gradients

between or within lakes: topography, geological qualities, inflow waters as

physical factors in the watershed, lake basin morphology (mainly depth and

area), water temperature, light conditions, turbidity, current flow, substrate

(sediment). And, chemical factors in play include inorganic ions, salinity,

organic matter, conductivity, alkalinity, pH and nutrients [11, 13, 21, 54, 56,

59, 61, 62, 64, 65, 69-73]. As this itemization indicates, a comprehensive

understanding of the mechanisms that determine the distribution and

species composition of macrophytes is difficult, due to multiple influencing

factors. In the Akan Caldera Lakes, species distribution was related to

trophic conditions (Fig 9 and 10), while number of species was closely related

to lake size (Fig 8). In general, species diversity at smaller scales increased

depending upon spatial environmental heterogeneity [74, 75]. Lakes Akan,

Panke and Penke, all large and oligo- and mesotrophic lakes with high

species count, appear to offer a large variety of the physical and chemical

factors noted above. The topography of capes, bays and islands in these lakes

diversifies wave action and substrate via varying wind-wave parameters,

and "fetch", i.e. the length of the lake surface over which wind blows [4, 61].

Inflow rivers locally alter substrate, current and water quality, greater water

depth lowers water temperature and reduces substrate grain size, and oligo-

or mesotrophic water allows sunlight to penetrate deeper into the littoral

zone, leading to a gradual gradient in the light environment [4, 5, 21, 55, 56,

59, 61, 62, 69]. This environmental variability may offer habitat for a

number of macrophytes in these large lakes. On the other hand, Pond Junsai,

dominated by floating-leaved and free-floating plants, is the only example of

dystrophic water quality in the Akan Caldera Lakes. Its brownish lake water,

derived from high DOC containing abundant humic substances, suppresses

submerged plant growth due to high light absorption [5, 21, 56, 61, 62].

Humic substances are thought to originate from decomposing terrestrial

and/or littoral plant material [5, 22, 50, 61]. Thus, eutrophication in Pond

Junsai may have undergone a different process than in the other lakes

because of differing surrounding vegetation, even if the initial process was

the same. To understand the factors affecting plant distribution the

relationships between detailed macrophyte distribution and habitat

micro-environments in each lake, including impacts of the surrounding

vegetation, must be elucidated.

Conclusions

In this study, we demonstrated that the ratio of Akan Caldera watershed

area to lake size was positively correlated with TP concentration, an

indicator of trophic level in lakes. Furthermore, the variety of trophic levels

in the lakes, ranging from oligotrophic to eutrophic, coincided with the "lake

succession" or "lake type" series of ecological succession in lakes. Further,

"hydrarch succession", a change in the species composition of macrophytes,

also indicates a regular series corresponding to the trophic levels. These facts

indicate that the unique eutrophication rates, defined by the ratio of

watershed area to lake size, may have diversified the water quality and the

aquatic flora of the lakes. Uniquely, this possibly illustrates lake ecological

succession in "real time", previously thought impossible to see [2, 9, 18, 24].

In other words, the Akan Caldera Lakes can be seen as a massive

experiment conducted by nature in an "Akan Caldera Laboratory", or seen as

a showcase visualizing a freshwater ecosystem, focusing on lake and

hydrarch succession. The essence of ecological succession is, simply put,

nothing more than a developing and changing process of biodiversity [75]. In

this context, the Akan Caldera Lakes will enable a new approach to the

comparative study of spatio-temporal fluctuations in biodiversity and

aquatic environments, including responses to changes in the surrounding

environment, especially climate change, now seen as an issue of paramount

importance [24, 26].

591   Acknowledgements

A part of this study is "Research of the aquatic plants in Lake Akan and

circumference lakes" and "Lake charts drawing of Lake Akan and

circumference lakes" conducted in 2013 and 2014 by the Ministry of

Environment of Japan. The City of Kushiro provided facilities and funding

for compilation of the study. Árni Einarrson, Mývatn Research Station,

Iceland, provided advice on research design. Takehiko Fukushima,

University of Tsukuba and Koichi Kamiya, Ibaraki Kasumigaura

Environmental Science Center supported the water analyses. The Hokkaido

Shimbun Press contributed to the aerial photography. Satoshi Kofuku and

Kokoro kikuchi, IDEA Consultants, Inc., assisted with literature research.

Shuji Hino, Yamagata University and Yasushi Ishikawa, Hokkaido Research

Organization, advised regarding caldera lakes and calculation of water

quality. Mami Yamazaki, Sapporo Museum Activity Center, provided advice

on macrophytes.

607 References

1. Whittaker RH. Communities and ecosystems. 2nd ed. New York:

Macmillan; 1975.

2. Sakamoto M. Ecological succession, II. Tokyo: Kyoritsu-Shuppan; 1976.

(in Japanese)

3. Owen OS. Natural resource conservation: an ecological approach, 4th ed.

New York: Macmillan; 1985.

4. Horne AJ, Goldman CR. Limnology, 2nd ed. New York: McGraw-Hill;

1994.

5. Wetzel RG. Limnology: lake and river ecosystems, 3rd ed. San Diego:

Academic Press; 2001.

6. Schindler DW, Vallentyne JR. The algal bowl, overfertilization of the

world's freshwaters and estuaries. Edmonton: University of Alberta

Press; 2008.

7. Vollenweider R, Kerekes J. Eutrophication of waters: Monitoring,

assessment and control. Paris: Organization for Economic Co-Operation

and Development (OECD); 1982.

8.   Lindeman RL. The trophic-dynamic aspect of ecology. Ecology. 1942;23:

399-417. https://doi.org/10.2307/1930126.

9.   Forel FA. Handbuch der Seenkunde: allgemeine Limnologie. Stuttgart: J.

Engelhorn; 1901. (in Germany)

10. Yoshimura S. Limnology. Tokyo: Sanseido; 1937. (in Japanese)

11. Pearsall WH. The development of vegetation in the English lakes,

considered in relation to the general evolution of glacial lakes and rock

basins. Proc R Soc B, Biol Sci. 1921;92: 259-284.

https://doi.org/10.1098/rspb.1921.0024.

12. Dachnowski A. Peat deposits of Ohio: their origin, formation and uses.

Columbus: Division of Geological Survey, Ohio; 1912.

http://hdl.handle.net/1811/78409

13. Pearsall WH. The aquatic vegetation of the English lakes. J. Ecol.

1920;8: 163-201. doi: 10.2307/2255612.

14. Dachnowski AP. Factors and problems in the selection of peat lands for

different uses. Washington, D. C.: United States Department of

Agriculture; 1926. https://doi.org/10.5962/bhl.title.108906

15. Kormondy EJ. Concepts of ecology. 2nd ed. Englewood Cliffs:

Prentice-Hall; 1969. http://ci.nii.ac.jp/ncid/BA1127514X

16. Odum EP. Fundamentals of ecology, 3rd ed. Philadelphia: Saunders;

1971.

17. Morishita I. Dam lake ecology. Tokyo: Sankaidou; 1983. (in Japanese)

18. Margalef R. Perspectives in ecological theory. Chicago: University of

Chicago Press; 1968.

19. Odum EP. The strategy of ecosystem development. Science. 1969;164:

262-270. http://habitat.aq.upm.es/boletin/n26/aeodu.en.html

20. Wetzel RG, Likens GE. Limnological analyses, 3ed ed. New York:

Springer; 2000.

21. Dodson SI. Introduction to limnology. New York: McGraw-Hill; 2005.

22. Dodds WK, Whiles MR. Freshwater ecology, 2nd ed. Burlington:

Academic Press; 2010.

23. Smith J, Appleby P, Battarbee R, Dearing J, Flower R, Haworth E, et al.

Environmental history and palaeolimnology. Springer Science &

Business Media; 1991.

24. Kling GW. A lake's life is not its own. Nature. 2000;408: 149-150. doi:

https://doi.org/10.1038/35041659.

25. Johnson EA, Miyanishi K. Testing the assumptions of chronosequences

in succession. Ecol Lett. 2008;11: 419-431. doi:

https://doi.org/10.1111/j.1461-0248.2008.01173.x

26. Sayer C, Roberts N, Sadler J, David C, Wade P. Biodiversity changes in a

shallow lake ecosystem: a multi - proxy palaeolimnological analysis. J

Biogeogr. 1999;26: 97-114. doi:

https://doi.org/10.1111/j.1365-2699.1999.00298.x

27. Satoh H. Akanko, explanatory text of the geological map of Japan,

Kushiro, No. 7. Tokyo: Geological Survey of Japan; 1965. (in Japanese

with English summary)

28. Tamada J, Nakagawa M. Eruption history of Oakan volcano, eastern

Hokkaido, Japan. Kazan. 2009;54: 147-162. (in Japanese with English

summary)

29. Hasegawa T, Nakagawa M. Large scale explosive eruptions of Akan

volcano, eastern Hokkaido, Japan: a geological and petrological case

study for establishing tephro-stratigraphy and-chronology around a

caldera cluster. Quat Int. 2016;397: 39-51. doi:

https://doi.org/10.1016/j.quaint.2015.07.058.

30. Maeda Ippoen Foundation. The nature of Akan National Park, 1993.

Akan: Maeda Ippoen Foundation; 1994. (in Japanese)

31. Hino S, Mikami H, Arisue J, Ishikawa Y, Imada K, Yasutomi R, et al.

Limnological characteristics and vertical distribution of phytoplankton

in oligotrophic lake Akan-Panke. Jpn J Limnol. 1998;59: 263-279. doi:

https://doi.org/10.3739/rikusui.59.263

32. Tanaka M. The lakes in Japan. Nagoya: Nagoya University Press; 1992.

(in Japanese)

33. Hokkaido Institute of Environmental Science. Lakes and marshes in

Hokkaido, revised edition. Sapporo: Hokkaido Institute of

Environmental Science; 2005. (in Japanese)

34. Clements FE. Plant succession: an analysis of the development of

vegetation. Washington: Carnegie Institution of Washington; 1916.

http://ci.nii.ac.jp/ncid/BA39490309

35. Solińska-Górnicka B, Symonides E. Long-term changes in the flora and

vegetation of Lake Mikołajskie (Poland) as a result of its eutrophication.

Acta Soc Bot Pol. 2001;70: 323-334. doi:

https://doi.org/10.5586/asbp.2001.040.

36. Kowalczewski A, Ozimek T. Further long-term changes in the submerged

macrophyte vegetation of the eutrophic Lake Mikolajskie (North Poland).

Aquat Bot. 1993;46: 341-345. doi:

https://doi.org/10.1016/0304-3770(93)90013-M.

37. Coops H, Doef RW. Submerged vegetation development in two shallow,

eutrophic lakes. Hydrobiol. 1996;340:115-120. doi:

https://doi.org/10.1007/BF00012742.

38. Egertson CJ, Kopaska JA, Downing JA. A century of change in

macrophyte abundance and composition in response to agricultural

eutrophication. Hydrobiol. 2004;524:145-156. doi:

https://doi.org/10.1023/B:HYDR.0000036129.40386.ce.

39. Tanakadate S. Research report on the volcano lakes in Hokkaido.

Sapporo: Hokkaido Government; 1925. (in Japanese)

40. Takayasu S, Igarashi H, Sawa K. Lake survey, Lake Akan. Sci Rep

Hokkaido Fish Exp Stn. 1930;21: 67-92. (in Japanese)

41. Takayasu S, Kondo K. River and lake survey, Lake Panke and Lake

Penke. Sci Rep Hokkaido Fish Exp Stn. 1936;39: 1-24. (in Japanese)

42. Sugimoto T. Introduction to Kashmir C3. Tokyo: Jitsugyo no Nihon Sha;

2010. (in Japanese)

43. Umekawa T, Wakana I, Ohara M. Reproductive behavior and role in

maintaining an aggregative form of the freshwater green alga Marimo,

Aegagropila linnaei, in Lake Akan, Hokkaido, Japan. Aquat Bot.

2021;168: 103309. doi: https://doi.org/10.1016/j.aquabot.2020.103309.

44. Igarashi S, Ishikawa Y, Mikami H. Long-term changes in limnological

characteristics in Lake Akan, Hokkaido. Res Rep NIES, Japan. 2000;153:

34-54. (in Japanese)

45. K0102 J. Testing Methods for Industrial Wastewater. Tokyo: Japanese

Standards Association; 2013. (in Japanese)

46. Grasshoff K, Kremling K, Ehrhardt M. Methods of seawater analysis:

Weinheim: John Wiley & Sons; 2009.

47. SCOR-UNESCO. Determination of photosynthetic pigments in seawater.

Monographs on oceanographic methodology 1. New York: UNESCO

Publications Center; 1966.

48. Fukushima T, Imai A, Matsushige K, Aizaki M, Otsuki A. Freshwater

DOC measurements by high-temperature combustion: comparison of

differential (DTC-DIC) and DIC purging methods. Water Res. 1996;30:

2717-2722. doi: https://doi.org/10.1016/S0043-1354(96)00198-4.

49. Miyadi D. Studies on the bottom fauna of Japanese lakes, VII. Lakes of

Hokkaido. Jpn J Zool. 1932;4: 223-252.

50. Brönmark C, Hansson L-A. The biology of lakes and ponds, 2nd ed.

Oxford: Oxford University Press; 2005.

51. Motoda S. The lakes in Hokkaido Sci Rep Hokkaido Fish Hatchery.

1950;5: 1-96. (in Japanese)

52. Wilson EO, MacArthur RH. The theory of island biogeography.

Princeton: Princeton University Press; 1967.

53. Møller TR, Rørdam CP. Species numbers of vascular plants in relation to

area, isolation and age of ponds in Denmark. Oikos. 1985;45: 8-16. doi:

https://doi.org/10.2307/3565216.

54. Rørslett B. Principal determinants of aquatic macrophyte richness in

northern European lakes. Aquat Bot. 1991;39: 173-193. doi:

https://doi.org/10.1016/0304-3770(91)90031-Y.

55. Weiher E, Boylen CW. Patterns and prediction of α and β diversity of

aquatic plants in Adirondack (New York) lakes. Can. J. Bot. 1994;72:

1797-1804. doi: doi.org/10.1139/b94-221.

56. Toivonen H, Huttunen P. Aquatic macrophytes and ecological gradients

in 57 small lakes in southern Finland. Aquat Bot. 1995;51: 197-221. doi:

https://doi.org/10.1016/0304-3770(95)00458-C.

57. Linton S, Goulder R. Botanical conservation value related to origin and

management of ponds. Aquat Conserv Mar Freshw Ecosyst. 2000;10:

77-91. doi:

https://doi.org/10.1002/(SICI)1099-0755(200003/04)10:2<77::AID-AQC39

1>3.0.CO;2-Y.

58. Mäkelä S, Huitu E, Arvola L. Spatial patterns in aquatic vegetation

composition and environmental covariates along chains of lakes in the

Kokemäenjoki watershed (S. Finland). Aquat Bot. 2004;80: 253-269. doi:

https://doi.org/10.1016/j.aquabot.2004.08.006.

59. Søndergaard M, Jeppesen E, Jensen JP. Pond or lake: does it make any

difference? Arch Hydrobiol. 2005;162: 143-165. doi:

10.1127/0003-9136/2005/0162-0143.

60. Kadono Y. A field guide to aquatic plants of Japan. Tokyo: Bun-ichi Sogo

Shuppan; 2014. (in Japanese)

61. Lacoul P, Freedman B. Environmental influences on aquatic plants in

freshwater ecosystems. Environ Rev. 2006;14: 89-136. doi:

https://doi.org/10.1139/a06-001.

62. Søndergaard M, Johansson LS, Lauridsen TL, Jørgensen TB,

Liboriussen L, Jeppesen E. Submerged macrophytes as indicators of the

ecological quality of lakes. Freshw Biol. 2010;55: 893-908. doi:

https://doi.org/10.1111/j.1365-2427.2009.02331.x.

63. Kawakami T. Collection record in Akan, Kushiro Province. Bot Mag

Tokyo. 1898;12: 220-225. (in Japanese)

64. Seddon B. Aquatic macrophytes as limnological indicators. Freshw Biol.

1972;2: 107-130. doi: doi.org/10.1111/j.1365-2427.1972.tb00365.x.

65. Wiegleb G. A study of habitat conditions of the macrophytic vegetation in

selected river systems in Western Lower Saxony (Federal Republic of

Germany). Aquat Bot. 1984;18: 313-352. doi:

https://doi.org/10.1016/0304-3770(84)90055-X.

66. Mäkirinta U. Classification of South Swedish Isoetid vegetation with the

help of numerical methods. Vegetatio. 1989;81: 145-157.

https://link.springer.com/chapter/10.1007/978-94-009-2432-1_12.

67. Heegaard E, Birks HH, Gibson CE, Smith SJ, Wolfe-Murphy S.

    Species–environmental relationships of aquatic macrophytes in

    Northern Ireland. Aquat Bot. 2001;70: 175-223. doi:

    https://doi.org/10.1016/S0304-3770(01)00161-9.

68. Schneider S, Melzer A. The trophic index of macrophytes (TIM)–a new

    tool for indicating the trophic state of running waters. Internat Rev.

    Hydrobiol. 2003;88: 49-67. doi: https://doi.org/10.1002/iroh.200390005.

69. Hutchinson GE. A treatise on limnology, Vol III, Limnological Botany.

    New York: John Wiley; 1975.

70. Spence DHN. Factors controlling the distribution of freshwater

    macrophytes with particular reference to the lochs of Scotland. J Ecol.

    1967;55: 147-170. doi: 10.2307/2257723.

71. Kadono Y. Occurrence of aquatic macrophytes in relation to pH,

    alkalinity, Ca^{++}, Cl^- and conductivity. Jap J Ecoi. 1982;32: 39-44. doi:

    10.18960/seitai.32.1_39.

72. Vestergaard O, Sand-Jensen K. Alkalinity and trophic state regulate

    aquatic plant distribution in Danish lakes. Aquat Bot. 2000;67: 85-107.

doi: https://doi.org/10.1016/S0304-3770(00)00086-3.

73. Lukács BA, Tóthmérész B, Borics G, Várbíró G, Juhász P, Kiss B, et al.

Macrophyte diversity of lakes in the Pannon Ecoregion (Hungary).

Limnologica. 2015;53: 74-83. doi

74. Wright DH, Currie DJ, Maurer BA. Energy supply and patterns of

species richness on local and regional scales. In: Ricklefs RE, Schluter D,

editors. Species diversity in ecological communities: historical and

geographical perspectives. 7. Chicago: University of Chicago Press; 1993.

75. Miyashita T, Noda T. Community ecology. Tokyo: University of Tokyo

Press; 2003. (in Japanese)

Supporting information

814   S1 Table. Data set of topography, water quality and number of species of macrophytes in

815   Akan Caldera Lakes.

816   S2 Table. Correlation matrix of topography, water quality and number of species of

817   macrophytes in Akan Caldera Lakes.

Lake / Species	L. Panke	L. Penke	L. Akan	P. Kyodai-naka	P. Hyotan	L. Taro	P. Junsai	Type of occurrence	Euclidean distance
Ranunculus nipponicus var. *submersus*	1	0	0	0	0	0	0	Oligotrophic	
Potamogeton alpinus	1	0	0	0	0	0	0	Oligotrophic	
Najas yezoensis	1	1	0	0	0	0	0	Oligotrophic	
Potamogeton gramineus	1	1	0	0	0	0	0	Oligotrophic	
Isoetes asiatica	1	1	0	0	0	0	0	Oligotrophic	
Potamogeton berchotoldii	1	1	1	1	0	1	0	Oligo-meso-eutrophic	
Potamogeton perfoliatus	1	1	1	1	1	0	0	Oligo-meso-eutrophic	
Potamogeton maackianus	1	1	1	0	0	0	0	Oligo-mesotrophic	
Myriophyllum spicatum	1	1	1	0	0	0	0	Oligo-mesotrophic	
Hydrilla verticillata	1	0	1	0	0	0	0	Oligo-mesotrophic	
Potamogeton compressus	1	0	1	0	0	0	0	Oligo-mesotrophic	
Stuckenia pectinata	1	0	1	0	0	0	0	Oligo-mesotrophic	
Ceratophyllum demersum	1	0	1	0	0	0	0	Oligo-mesotrophic	
Potamogeton crispus	0	0	1	0	0	0	0	Mesotrophic	
Persicaria amphibia	0	0	1	0	0	0	0	Mesotrophic	
Brasenia schreberi	0	0	0	0	0	0	1	Dystrophic	
Nuphar pumila var. *pumila*	0	0	0	0	0	0	1	Dystrophic	
Nymphaea tetragona var. *tetragona*	0	0	0	0	0	0	1	Dystrophic	
Potamogeton natans	0	0	0	0	0	0	1	Dystrophic	
Utricularia macrorhiza	0	0	0	0	0	0	1	Dystrophic	
Utricularia minor	0	0	0	0	0	0	1	Dystrophic	

1: presence, 0: absence.

Occurrence type of macrophyte	Species of macrophyte	Trophic type of lake			
		Oligotrophic	Mesotrophic	Eutrophic	Dystrophic
Oligotrophic	*Ranunculus nipponicus* var. *submersus* *Potamogeton alpinus* *Najas yezoensis* *Potamogeton gramineus* *Isoetes asiatica*	*R. nipponicus* var. *submersus* — L. Panke			
Mesotrophic	*Potamogeton crispus* *Persicaria amphibia*		*P. crispus* — L. Akan		
Oligo-mesotrophic	*Potamogeton maackianus* *Myriophyllum spicatum* *Hydrilla verticillata* *Potamogeton compressus* *Stuckenia pectinata* *Ceratophyllum demersum*	*M. spicatum* — L. Panke	*M. spicatum* — L. Akan		
Oligo-meso-eutrophic	*Potamogeton berchotoldii* *Potamogeton perfoliatus*	*P. perfoliatus* — L. Penke	*P. perfoliatus* — P. Kyodai-naka	*P. perfoliatus* — P. Hyotan	
Dystrophic	*Brasenia schreberi* *Nuphar pumila* var. *pumila* *Nymphaea tetragona* var. *tetragona* *Potamogeton natans* *Utricularia macrorhiza* *Utricularia minor*				*B. schreberi* — P. Junsai