July 1, 2023

Detecting natural selection in trait-trait coevolution

Daohan Jiang[1,2*] and Jianzhi Zhang[1]

[1]Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, Michigan 48109, USA
[2]Present address: Department of Quantitative and Computational Biology, University of Southern California, Los Angeles, California 90007, USA

*Correspondence to
Daohan Jiang
Department of Quantitative and Computational Biology
University of Southern California
416F Ray R. Irani Hall
1050 Childs Way
Los Angeles, CA 9008, USA
Phone: 734-545-0764
Email: daohanji@usc.edu

Running title: Selection in trait-trait coevolution

Keywords: fly, morphology, mutation, modularity, pleiotropy, yeast

ABSTRACT

No phenotypic trait evolves independently of all other traits, but the cause of trait-trait

coevolution is poorly understood. While the coevolution could arise simply from pleiotropic

mutations that simultaneously affect the traits concerned, it could also result from multivariate

natural selection favoring certain trait relationships. To gain a general mechanistic

understanding of trait-trait coevolution, we examine the evolution of 220 cell morphology traits

across 16 natural strains of the yeast *Saccharomyces cerevisiae* and the evolution of 24 wing

morphology traits across 110 fly species of the family Drosophilidae, along with the variations of

these traits among gene deletion or mutation accumulation lines (a.k.a. mutants). For numerous

trait pairs, the phenotypic correlation among evolutionary lineages differs significantly from that

among mutants. Specifically, we find hundreds of cases where the evolutionary correlation

between traits is strengthened or reversed relative to the mutational correlation, which, according

to our population genetic simulation, is likely caused by multivariate selection. Furthermore, we

detect selection for enhanced modularity of the yeast traits analyzed. Together, these results

demonstrate that trait-trait coevolution is shaped by natural selection and suggest that the

pleiotropic structure of mutation is not optimal. Because the morphological traits analyzed here

are chosen largely because of their measurability and thereby are not expected to be biased with

regard to natural selection, our conclusion is likely general.

BACKGROUND

Many phenotypic traits covary during evolution. For example, the logarithm of brain weight and that of body weight show a nearly perfect linear relationship across mammals [1, 2]. In theory, four processes may explain such trait-trait coevolution. First, it could arise simply from pleiotropic mutations that simultaneously influence these traits with a more or less constant ratio of effects [3-5], as has been previously shown empirically [6-10]. Second, trait covariation could arise from the linkage disequilibrium between genes controlling these traits [5, 11-13], but such trait covariation is expected to be restricted to closely related individuals due to the deterioration of linkage disequilibrium as a result of recombination. If the linkage disequilibrium is stably maintained due to, for example, chromosomal inversion, the involved linked genes can be regarded as a supergene with mutational pleiotropy [13]. For this reason, linkage disequilibrium is negligible except for trait covariation among closely related individuals. Third, shared ancestry can also create apparent trait correlations across lineages, which, however, can be explained away when the phylogenetic relationships are taken into account in correlation analysis [14]. Finally, trait covariation could be a result of natural selection for particular trait relationships that are advantageous, a phenomenon known as correlational selection or multivariate selection [2, 15-20].

Despite a long-standing interest in trait correlation in evolution [2, 13, 21], which is also referred to as phenotypic integration in the literature [22, 23], our understanding of the roles of mutation and selection in trait-trait coevolution remains limited. Most studies on the subject focused on a small number of traits that are physiologically or ecologically important [24], such as skull anatomy characters [25-30], behavioral syndrome (i.e., sets of correlated behavioral traits) [31, 32], and ecological or organismal traits correlated with the metabolic rate [33-37]; hence, they may not provide a general, unbiased picture of trait-trait coevolution. Additionally, it is the trait correlation resulting from standing genetic variation and its effect on adaptation that have received the most attention [38-44]. But, because standing genetic variation could have been influenced by selection [40], the resulting trait correlation may not inform the correlation produced by mutation. Not knowing the mutational correlation hinders a full understanding of the contribution of selection.

Related to trait-trait correlation is the concept of modularity. It has been hypothesized that it is beneficial for organisms to have a modular organization such that functionally related

traits belonging to the same module covary and genotypes and/or phenotypes that lead to low

fitness are less likely to occur [21, 25, 45-47]. Although modularity is a well-recognized feature

of many trait correlation networks, the relative contribution of selection and mutational

pleiotropy to modularity has not been assessed at the phenome scale [46-48].

To gain a general mechanistic understanding of trait-trait coevolution, we study the

phenotypic correlations for a large number of trait pairs at the levels of mutation and long-term

evolution; natural selection is inferred when the evolutionary correlation between traits cannot be

fully explained by the mutational correlation. We also ask if the overall pattern of trait

correlation (i.e., phenotypic integration) differ at the two levels. Our primary data include 220

cell morphology traits of the budding yeast *Saccharomyces cerevisiae* that have been measured

in 4817 single-gene deletion lines [49], 89 mutation accumulation (MA) lines (for a subset of

187 traits) [50], and 16 natural strains with clear phylogenetic relationships [49, 51]. These traits

were quantified from fluorescent microscopic images of triple-stained cells and were originally

chosen for study because of their measurability regardless of potential roles in evolution and

adaptation [49]. Subsequent studies found that these cell morphological traits are correlated with

the yeast mitotic growth rate (i.e., a proxy for fitness) to varying degrees [7]. Hence, these traits

may be considered representatives of phenotypic traits that have different contributions to

fitness. Previous analyses of these traits among natural strains unveiled signals of positive

selection on individual traits [52], but their potential coevolution has not been studied. While

studying these trait pairs can offer a general picture of trait-trait coevolution, we recognize that

the selective agent would be hard to identify should selection be detected, because the biological

functions of these traits (other than correlations with the growth rate) are generally unknown

[52]. To verify the generality of the findings from the yeast traits, we analyze another dataset

that includes 12 landmark vein intersections on the fly wings that have been measured in 150

MA lines of *Drosophila melanogaster* [9] and 110 Drosophilid species [53]. At last, using

computer simulations, we demonstrate how certain regimes of selection could explain the

observed differences between mutational and evolutionary correlations.

RESULTS

Evolutionary correlations differ from mutational correlations for many trait pairs

To investigate if trait correlations in evolution can be fully accounted for by the

correlations generated by mutation, we examined all pairs of the 220 yeast cell morphology traits

previously measured. For each pair of traits, we computed the mutational correlation COR_M,

defined as Pearson's correlation coefficient across 4,817 gene deletion lines (upper triangle in

Fig. 1A, **Data S1**), and evolutionary correlation COR_E, defined as Pearson's correlation

coefficient across 16 natural strains (lower triangle in **Fig. 1A**, **Data S1**) with their phylogenetic

relationships (**Fig. S1**) taken into account (see Materials and Methods). Note that the original

data contained 37 natural strains [51], of which 21 belong to the "mosaic" group [54, 55]—their

phylogenetic relationships with other *S. cereviase* strains vary among genomic regions—so

cannot be included in our analysis that requires considering phylogenetic relationships.

We found that the frequency distribution of COR_E across all trait pairs differs

significantly from that of COR_M (**Fig. 1B**), suggesting the action of selection. For each pair of

traits, we transformed the COR_M and COR_E to Z-scores using Fisher's r-to-Z transformation and

conducted Z-test to determine whether the two correlations are significantly different. Of the

24,090 trait pairs examined, 6743 pairs (or 28.0%) have a COR_E that deviates significantly from

COR_M at the false discovery rate (FDR) of 5% (**Table 1**, **Data S1**), suggesting that natural

selection has shaped the coevolution of many trait pairs. To investigate whether the above result

is biased because of the use of each trait in many trait pairs, we randomly arranged the 220 traits

into 110 non-overlapping pairs and counted the number of pairs with COR_E significantly

different from COR_M. This was repeated 1,000 times to yield 1,000 estimates of the proportion

of trait pairs with significantly different COR_E and COR_M. The middle 95% of these estimates

ranged from 14.5% to 40.1%, with the median estimate being 28.2%, almost identical to the

result (28.0%) from all pairwise comparisons. Hence, there is no indication that using

overlapping trait pairs has biased the estimate of the fraction of trait pairs with significantly

different COR_E and COR_M.

To further test selection, we simulated neutral evolution along the yeast tree 1000 times

under a Brownian motion model with the observed mutational covariance matrix M used as the

mutational input, generating 1,000 simulated datasets. Before the simulation, we confirmed that

the sampling error of our estimated M is negligible, likely because of the large number of

mutants used in M estimation (**Table S1**; see Materials and Methods). From each simulated

dataset, we calculated the number of trait pairs with COR_E significantly different from COR_M.

Only in 0.7% of the simulated data did we find this number equal to or greater than that from the

actual data (**Table 1**), indicating that the observed evolutionary correlations between traits

cannot be explained by the neutral Brownian motion model. The distribution of mutational

effects can be asymmetric and skewed [56] while it is assumed normal in the Brownian motion

model. Nevertheless, simulations showed that mutational bias will not render COR_E deviate

from COR_M in the absence of selection and will not enlarge the variance of COR_E (**Table S2**; see

Materials and Methods).

We divided the 6743 cases of significantly different COR_E and COR_M into three

categories. In the first category, the trait correlation generated by mutation is strengthened by

natural selection during evolution. A total of 2,727 trait pairs are considered to belong to this

"strengthened" category (**Table 1**) because they satisfy the following criteria: COR_E and COR_M

have the same sign and $|COR_E| > |COR_M|$, or COR_E and COR_M have different signs but only

COR_E is significantly different from 0 (at the nominal P-value of 0.05) (**Fig. 2A**). In the second

category, the trait correlation generated by mutation is weakened by natural selection during

evolution. A total of 1,221 trait pairs satisfying the following criteria are classified into this

"weakened" category (**Table 1**): COR_E and COR_M have the same sign and $|COR_E| < |COR_M|$, or

COR_E and COR_M have different signs but only COR_M is significantly different from 0 (**Fig. 2B**).

In the last category, the trait correlation generated by mutation is reversed in sign by natural

selection during evolution. A total of 2,795 trait pairs satisfying the following criteria are in this

"reversed" category (**Table 1**): COR_E and COR_M have different signs and are both significantly

different from 0 (**Fig. 2C**).

To assess the robustness of the selection signals detected, we repeated the above analysis

using COR_M estimated from 89 mutation accumulation (MA) lines [43] (**Fig. S2A**, **Data S1**).

Again, the overall frequency distribution across all trait pairs differs significantly between COR_E

and COR_M (**Fig. S2B**). We found that 5,146 trait pairs exhibit a significantly different COR_E

from the corresponding COR_M (**Table 1**, **Data S1**), supporting a role of selection in the

coevolution of many trait pairs. When comparing the analysis using COR_M from gene deletion

lines and that using COR_M from MA lines, we found 990 trait pairs to exhibit selection signals

and fall into the same category in both analyses, including 275 pairs in the "strengthened"

category, 223 pairs in the "weakened" category, and 574 pairs in the "reversed" category. All of

these numbers substantially exceed the corresponding expected random overlaps ($P < 0.001$

based on 1,000 random draws in each case; the medians across the 1,000 draws are 271, 68 and

163, respectively), suggesting the reliability of both analyses. Although mutations in MA lines

are more natural than those in gene deletion lines, the number of MA lines is much smaller than

the number of gene deletion lines and only 187 of the original 220 traits were measured in the

MA lines. For these reasons, we focused on the COR_M estimated from the gene deletion lines in

subsequent analyses.

To examine the generality of the above yeast-based findings, we analyzed the 24 wing

morphology traits of Drosophilid flies. The COR_M and COR_E have been previously estimated

from 150 MA lines [9] and 110 Drosophilid species, respectively (**Fig. S3A**, **Data S1**). The

overall frequency distribution across all trait pairs differs significantly between COR_E and COR_M

(**Fig. S3B**). Of the 276 pairs of traits, 144 (52.2%) showed a significant difference between

COR_E and COR_M (**Table 1**, **Data S1**), indicating widespread actions of selection in the

coevolution of fly wing morphology traits.

Together, these results demonstrate that, for many trait pairs, mutational and evolutionary

correlations between morphological traits are more different than expected under neutrality.

This observation suggests an important role of selection in shaping the strength and/or direction

of trait correlation in evolution.

Effects of different selection regimes on trait-trait coevolution

The strengthened, weakened, and reversed trait correlations in evolution may have

resulted from different selection regimes. Below we consider various selection regimes that

could potentially explain these types of difference between COR_M and COR_E (**Fig. 3**). First,

when a specific allometric relationship between two traits is selectively favored, the population

mean trait values are expected to be concentrated near the fitness ridge or the optimal allometric

line, resulting in a strong evolutionary correlation between the traits (i.e., a high $|COR_E|$) (**Fig.**

3A). Unless COR_M is already similar to COR_E, we expect to see strengthened or reversed COR_E

depending on COR_M. Second, if there is a single fitness peak for an optimal combination of trait

values and if there is sufficiently strong stabilizing selection on the optimal phenotype, the

population mean phenotype should be restricted within a small range of the optimal phenotype in

all directions in the phenotypic space regardless of the mutational variance. Consequently,

COR_E is expected to be close to 0, which could account for a weakened evolutionary correlation

relative to the mutational correlation (**Fig. 3B**). Finally, if the fitness optimum varies across

lineages in a random fashion, the steady-state COR_E will be close to zero, potentially leading to

the weakening of the evolutionary correlation relative to the mutational correlation (**Fig. 3C**).

To verify these predictions, we simulated the evolution of two traits. Under each

parameter set, we simulated 50 independent replicate lineages and computed the correlation

coefficient, or COR_E, between the traits across the replicate lineages at the end of the simulated

evolution. This was repeated 200 times to obtain an empirical distribution of COR_E. To evaluate

the difference between COR_M and COR_E, we examined the location of COR_M in the distribution

of COR_E; a significant ($P < 0.05$) difference is inferred if COR_M is in the left or right 2.5% tail of

the COR_E distribution.

As expected, in the absence of selection, the distribution of COR_E is centered around

COR_M (first block in **Table 2**). When a specific allometric relationship is selectively favored, a

high $|COR_E|$ always emerges regardless of the COR_M used, resulting in either strengthened or

reversed evolutionary correlations ($P < 0.005$ for all parameter sets examined; the second to fifth

blocks in **Table 2**). By contrast, stabilizing selection of an optimal phenotype leads to weakened

correlation across replicate lineages when $|COR_M|$ is not small (sixth block in **Table 2**). Finally,

when different lineages have different phenotypic optima that are randomly picked from the

standard bivariate normal distribution, weakened evolutionary correlations are generally

observed except when COR_M is close to zero (bottom block in **Table 2**). These results suggest

that the strengthened and reversed evolutionary correlations of yeast and fly morphological traits

are likely caused by selections of allometric relationships, while the weakened correlations are

likely caused by selections of individual traits either when there is a single optimal phenotype or

when the optimal phenotype randomly varies among lineages.

Selection for enhanced modularity of yeast morphological traits

While all of the above analyses focused on individual trait pairs, here we ask whether the

overall trait correlation across divergent lineages is stronger or weaker than that created by

mutation. As a measure of the overall level of trait correlation (i.e., overall integration), we

calculated the variance of eigenvalues (V_{eigen}) of the correlation matrix from divergent lineages

and mutants, respectively. A greater V_{eigen} corresponds to a stronger overall correlation between

traits because the eigenvalues become less evenly distributed as the absolute values of the

correlation coefficients become larger [57]. However, the sample size (i.e., the number of

strains) in the estimation of the correlation matrix also influences V_{eigen}; a matrix estimated from

a smaller sample naturally tends to have fewer positive eigenvalues and greater V_{eigen}. To

exclude the influence of this factor, we randomly sampled the mutant strains to obtain 5000

control datasets. Because the rank number of the evolutionary correlation matrix is 15 for the

yeast data (i.e., 15 positive eigenvalues), each control dataset also consists of 15 randomly drawn

strains such that the corresponding mutational correlation matrix also has 15 positive

eigenvalues. We examined the location of the observed V_{eigen} in this distribution and computed a

P-value based on this location (see Materials and Methods). For the yeast traits, V_{eigen} of the

observed evolutionary correlation matrix exceeds that in 96% of control datasets ($P = 0.08$ in a

two-tailed test; **Table 3**). Furthermore, only two of the 5000 control datasets have V_{eigen}

significantly different from that of the observed evolutionary correlation matrix (Fligner-Kileen

test). Hence, there is little evidence for a difference between the overall evolutionary correlation

and the overall mutational correlation in yeast. For the fly data, the number of positive

eigenvalues is unlimited by the sample size for both the evolutionary and mutational correlation

matrices, hence we directly compared V_{eigen} between the two matrices, but found them to be

similar ($P = 0.459$, Figner-Kileen test; **Table 3**). We also compared the overall integration

between yeast and flies using $V_{eigen}/(n-1)$, where n is the number of traits examined. $V_{eigen}/(n-1)$

equals 0.157 and 0.268 for the yeast mutational and evolutionary matrices, respectively, whereas

the corresponding values in flies are 0.153 and 0.190, respectively.

In addition to the overall level of trait correlation, we asked whether the correlational

structure of traits exhibits different levels of modularity among divergent lineages when

compared with that among mutants. To this end, we used a covariance ratio (*CR*) test [58] that

compares covariance within and between pre-defined modules (see Materials and Methods).

Specifically, we calculated *CR* for the evolutionary covariance matrix and compared it to the *CR*

distribution based on 5000 mutational covariance matrices estimated from the randomly drawn

subsets of mutants aforementioned. We treated the three non-overlapping categories of the yeast

morphological traits—actin traits, nucleus traits, and cell wall traits [49]—as three modules

(**Data S1**). We found that the *CR* of the evolutionary covariance matrix exceeded that of every

control dataset ($P < 0.001$; **Table 3**), suggesting natural selection for increased modularity in

evolution.

DISCUSSION

By comparing the trait-trait correlation across mutants (COR_M) with that across divergent lineages (COR_E) for 24,090 pairs of yeast cell morphology traits and 276 pairs of fly wing morphology traits, we detected the action of natural selection in trait-trait coevolution. The fraction of trait pairs showing evidence for selection is substantially higher in the fly (52%) than yeast (28%) data ($P < 10^{-4}$, chi-squared test). This is at least in part caused by a difference in statistical power, because the number of strains/species used for estimating COR_E is much greater for the fly (110) than yeast (16) data. It is likely that a higher fraction than 28% of the yeast trait pairs are subject to selection in their coevolution. Furthermore, our comparison between COR_E and COR_M intends to test selection on trait correlations common among the evolutionary lineages considered. If different evolutionary lineages have different trait correlations, the COR_E estimated from all lineages may not be significantly different from COR_M even when selection occurs in some or all of the lineages. In other words, our test is expected to underestimate the proportion of trait pairs subject to selection.

One potential biological explanation of the yeast-fly disparity in the prevalence of correlational selection is divergence time: the fly species represent a group that is tens of millions of years old while the yeast strains diverged from each other much more recently [53-55]. It is known that genetic correlations predict evolutionary correlations better over shorter timescales [38]. Similarly, selection might have had more time to decouple the pattern of evolutionary divergence from the mutational input in the flies but not yet in the yeast strains.

While we assumed that the mutants used carry all designed or natural mutations, extremely deleterious mutations such as lethal mutations are not represented. However, because such mutations are quickly selectively purged in natural populations, they should only be present transiently and are presumably unlikely to contribute to long-term evolution. Hence, their absence from our mutant data should not qualitatively alter our results.

We demonstrated by simulations that various selection regimes can explain differences between COR_M and COR_E. In particular, strengthened or reversed COR_E relative to COR_M can occur when a specific allometric relationship is preferred, while weakened COR_E can occur under directional or stabilizing selection of individual traits. A notable difference between the simulation results and empirical observations is that the simulations tend to end up with extreme

values of $|COR_E|$ (i.e., close to either 1 or 0) except in the case of neutrality, whereas the

empirically observed $|COR_E|$ is usually less extreme even when COR_M and COR_E are

significantly different. This is due to the fact that the simulation results usually represent steady-

state correlations across lineages. That is, the mean phenotype of each lineage is at or near the

corresponding optimum (if any); consequently, $|COR_E|$ is close to 1 when the optimum is a line

and close to 0 when the optimum is a single combination of two trait values. However, the

population mean phenotypes may not be close to their optima in some strains because of recent

changes of the optima or the sparsity of mutations toward the optima, the latter of which is well

known as a potential hindrance to adaptation [38, 42, 43, 59]. Another possibility is the

existence of a wide range of preferred allometry such that there is no strong selection for extreme

$|COR_E|$. Finally, selection may not result in the preferred allometry between two traits because

of the constraints from unconsidered traits [60].

It is worth noting that the yeast natural strains had been cultured in synthetic media

before phenotyping [51] while the mutant strains were all grown in the rich medium YPD [49,

50]. Hence, it remains a possibility that the difference between COR_E and COR_M reported here

contains a component caused by the environmental difference in phenotyping. Notwithstanding,

our analysis suggests that this component is small (see Materials and Methods), which is

expected because both media are meant to provide an ideal, stress-free environment for yeast

growth. This said, future phenotyping in the same medium will be needed to validate our

findings.

While selection was detected for many trait pairs, a large fraction of trait pairs, especially

in the yeast data, do not show a significant difference between COR_E and COR_M. These trait

pairs may be divided into two groups. In the first group, COR_E and COR_M are actually different,

but the difference is not found significant due to the limited statistical power. As mentioned, we

believe that a substantial fraction of yeast trait pairs belong to this category due to the relatively

low statistical power in detecting the difference between COR_E and COR_M in the yeast data. In

the second group, COR_E truly equals COR_M, which could result from one of the following three

scenarios. First, the specific trait-trait correlation does not impact fitness so evolves neutrally.

Second, the two traits have an intrinsic, immutable relationship (such as the hypothetical traits of

body size and twice the body size), so will yield equal COR_E and COR_M; this possibility can be

tested by examining the correlation of the two traits across isogenic individuals that show non-

heritable phenotypic variations [61]. The last and perhaps the most interesting scenario is that
the trait-trait correlation impacts fitness and hence has driven the optimization of COR_M via a
second-order selection [52, 59, 62, 63] such that the first-order selection of mutations that affect
the two traits is no longer needed. However, the relative frequencies of these three scenarios are
unknown.
In addition to pairwise trait correlations, we tested hypotheses regarding the evolution of
overall phenotypic integration and modularity. In the yeast data, we observed a higher
modularity across natural strains than across mutants but did not find evidence for a change of
overall phenotypic integration in evolution. These results support the view of increasing
modularity during evolution [21, 25, 45, 46, 64] but also suggest that modularity is enhanced by
both strengthening trait-trait correlations within modules and weakening trait-trait correlations
across modules. We found the overall integration lower for the fly than yeast traits, but whether
this observation indicates a difference between different types of traits (i.e., cellular traits and
multicellular organisms' morphological traits) or between multicellular and unicellular
organisms requires analyzing more species and traits.
Our analysis compared COR_M estimated from one yeast strain (BY) with COR_E estimated
from 16 different strains, under the assumption of a constant COR_M across different strains.
While it is a common practice to assume that the mutational architecture is more or less constant
during evolution and to study phenotypic evolution by comparing mutational or genetic
(co)variances in one species with those among different species [53, 65, 66], genetic variations
affecting the genetic (co)variances of phenotypic traits have been reported [67-69]. As discussed
earlier, such genetic variations may allow second-order selection of COR_M. For instance, it has
been hypothesized that the optimization of mutational (co)variances driven by selection for
mutational robustness and/or adaptability can lead to modularity [21, 46]. It has indeed been
found in the study of *Drosophila* gene expression traits that variational modules identified from
mutants can be predicted to some extent by functional grouping of genes (i.e., Gene Ontology
terms), although there is still much difference between functional modules and modules resulting
from mutational pleiotropy, suggesting that optimization of the mutational architecture is far
from complete even if it did take place [47]. Even without second-order selection, COR_M could
still vary across strains because the pleiotropic effects of a mutation can vary by the environment
and genetic background [19, 70, 71]. Regardless, in the future, it would be desirable to measure

mutant phenotypes from multiple lineages to investigate whether COR_M evolves, how rapidly it

evolves, and whether its evolution is largely neutral or adaptive.

Our analysis of the yeast dataset is subject to a major limitation resulting from the

structure of the dataset. As many yeast strains are mosaic, only a small number of strains (16)

were used in our study. Most of the remaining strains fall in one clade (**Fig. S1**), which is the

Wine/European clade [54, 55]. That is, a substantial fraction of evolution along the yeast tree

took place on internal branch(es), which would further reduce the effective sample size [72]. As

a result, the COR_E estimate may not be very accurate, and the selection test suffers from low

statistical power. It would be desirable if more non-mosaic strains from non-Wine/European

clades are included. Another caveat regarding the calculation of COR_E is that correction

methods like independent contrast do not always sufficiently account for the tree structure and

can be susceptible to singular evolutionary events (e.g., shift of evolutionary rate in a clade) [73];

in our case, such a singular event could have taken place in the Wine/European clade after it had

split from other yeast strains.

In summary, we detected the action of natural selection in shaping trait-trait coevolution.

Because the traits analyzed here, especially the yeast traits, were chosen almost exclusively due

to their measurability, our results likely reflect a general picture of trait-trait coevolution.

Measuring these yeast traits in additional divergent natural strains with clear phylogenetic

positions could improve the statistical power and clarify whether the fraction of trait pairs whose

coevolution is shaped by selection is much greater than detected here. Finally, the detection of

selection for enhanced modularity of the yeast traits analyzed supports the hypothesis that

modularity is beneficial [21, 25]. The detection of selection in trait-trait coevolution and

selection for enhanced modularity suggests that the current pleiotropic structure of mutation is

not optimal. This nonoptimality could be due to the weakness of the second-order selection on

mutational structure and/or a high dependence of the optimal mutational structure on the

environment, which presumably changes frequently. Future studies on how the mutational

structure evolves will likely further enlighten the mechanism of trait-trait coevolution.

CONCLUSION

In this study, we analyzed morphological traits of yeast and flies and compared patterns

of trait-trait correlation at the levels of mutation and long-term evolution. In both datasets, we

discover that the evolutionary correlation differs significantly from the mutational correlation for

numerous trait pairs, revealing a role of natural selection in trait-trait coevolution. We also

provide evidence for selection for enhanced modularity of the yeast traits. Insights gained in this

study can be summarized as follows:

1) Can trait-trait correlations in long-term evolution be explained by mutations? Our

analyses showed that some correlations observed across divergent lineages differ

significantly from correlations created by mutations. In addition, the pattern of

phenotypic covariance among natural yeast strains has stronger modularity (i.e.,

stronger within-module correlations and/or weaker between-module correlations)

than among mutants. These observations together indicate that selection likely played

a role in shaping trait correlations in long-term evolution.

2) What evolutionary forces drive trait-trait correlation during evolution? Our

simulations show how various selection regimes render the pattern of correlation

during evolution different from that caused by mutation. Some types of differences,

including strengthening and reversal of correlations, are explained by selection for an

optimal allometric relationship, but not selection on individual traits.

MATERIALS AND METHODS

Phenotypic data

The *S. cerevisiae* cell morphology traits were previously measured by analyzing

fluorescent microscopic images. Three phenotypic datasets were compiled and analyzed in this

study, including (i) 220 traits measured in 4,718 gene deletion lines that each lack an

nonessential gene [49], (ii) the same 220 traits measured in 37 natural strains [51], and (iii) 187

of the 220 traits measured in 89 mutation accumulation (MA) lines [50]. When comparing

patterns of trait correlation between two datasets, we used traits available in both datasets. For

each deletion strain, many cells (95 on average) were phenotyped, and the average trait value of

all these cells were used to represent the strain in our analyses.

Three types of traits were measured in the deletion strains and the natural strains,

including actin traits (i.e., measurements based on dyed actin cytoskeleton), cell wall traits (i.e.,

measurements based on dyed mannoprotein and cell wall markers), and nucleus traits (i.e.,

measurements based on dyed nuclear DNA) [49, 51]. These three categories were treated as

three modules in our analysis of modularity. Only the cell membrane traits and nucleus traits

were measured in the MA lines [50].

Before the analyses, we first standardized all trait values by converting each trait value to

the natural log of the ratio of the original trait value to a reference such that the distributions

become approximately normal and suitable for the Z-test. The standardized value of the ith trait

in the jth strain is $\tilde{X}_{i,j} = ln\frac{X_{i,j}}{X_{i,r}}$, where $X_{i,j}$ is the original trait value and $X_{i,r}$ is the trait value of

the reference. For the gene deletion lines, the reference is the wild-type BY strain. For the MA

lines, the reference is the progenitor strain used in MA. For natural strains, the reference is the

same as the reference of the mutant strains to be compared with (i.e., wild-type BY or progenitor

of the MA lines).

The locations of 12 vein intersections on the fly wing were previously measured in 150

MA lines of *Drosophila melanogaster* and a mutational covariance matrix was estimated [9].

Because each intersection is described by two coordinates, which are counted as two traits, there

are 24 traits in this dataset. These traits were also measured in 110 Drosophilid species and an

evolutionary covariance matrix was estimated with species phylogeny taken into account [53].

Both matrices are based on log-scale trait values.

Influence of the sampling error on the correlational structure

To evaluate the influence of sampling error on the estimated mutational covariance

matrix (i.e., the M matrix) of yeast or fly, we took samples (vectors of phenotypes) from the

multivariate distribution of M (4,817 samples for yeast gene deletion data and 150 samples for

fly MA data), estimated a covariance matrix ($\widetilde{M}$) from these samples, and calculated Pearson's

correlation coefficient between the eigenvalues of M and $\widetilde{M}$. For instance, for the yeast data, M

and $\widetilde{M}$ each has 220 eigenvalues, and we calculated the correlation between these two sets of

eigenvalues as a measurement of similarity between M and $\widetilde{M}$. This was repeated 1,000 times

and the distribution of the correlation coefficient was used to evaluate the potential impact of

sampling error on M.

Impact of the environmental difference on the correlational structure of the yeast traits

Because the natural strains of yeast had been grown in synthetic media before

phenotyping [51] while the mutant strains were all grown in the rich medium YPD [49, 50], we

tested whether this environmental difference affected the correlational structure of the yeast

morphological traits under consideration. Specifically, we examined whether the phenotype of

the BY strain grown in synthetic media (referred to as "synthetic phenotype" for short) falls in

the distribution of 123 biological replicates of BY grown in YPD (referred to as "YPD

phenotypes" for short). The phenotypes were normalized in the way described earlier with the

mean phenotype of the YPD replicates used as the reference. We decomposed YPD phenotypes

into principal components (PCs) and focused on the first three PCs, which together explained

67.5% of the variance among the 123 YPD phenotypes. We then calculated the values of the

three PC traits of the synthetic phenotype. The synthetic phenotype is in the central 95% of the

distribution of the YPD phenotypes for each of the three PC traits, indicating a lack of major

effect of the difference between synthetic and YPD media on the correlational structure of the

yeast traits concerned.

Comparison between mutational and evolutionary correlations

To take into account the phylogenetic relationships among yeast strains in estimating

COR_E, we utilized a distance-based tree previously inferred [55] (**Fig. S1**). Strains with mosaic

origins inferred in the same study [55] were removed before analysis, resulting in 16 remaining

natural strains. Because the BY strain was not included in the data file in that study [55], W303,

a laboratory strain closely related to BY, was chosen to represent BY. We obtained the

evolutionary covariance matrix using the *ratematrix* function from the R package *geiger* [74,

75], which calculates evolutionary covariances using the independent contrast method [14]. The

evolutionary covariance matrix was then converted to the corresponding correlation matrix.

To test whether the observed pairwise trait correlation at the level of evolutionary

divergence is significantly different from that expected by mutation alone for each pair of traits,

we first converted both correlations to Z-scores by $Z = \frac{1}{2}\left[ln\,(1 + r)\ -ln\,(1 - r)\right]$, where r is

the correlation coefficient. The testing statistic was then computed by $Z = \frac{Z_E - Z_M}{\sqrt{\frac{1}{n_E-3}+\frac{1}{n_M-3}}}$, where Z_E

and Z_M are Z-scores converted from COR_E and COR_M, respectively, n_E is the number of

independent contrasts, which equals the number of natural strains minus one, and n_M is the

number of mutant strains. Two-sided P-value was calculated from each Z and converted to

adjusted P-value following the Benjamini-Hochberg procedure [76]. An adjusted P-value below

0.05 indicates selection.

To see how many trait pairs would show a significant difference between COR_E and

COR_M under neutrality, we simulated neutral evolution along the phylogenetic tree that had been

used in estimating COR_E. A Brownian motion model was used to simulate neutral phenotypic

evolution such that the amount of evolution in branch i is $M_i l$, where M_i is a vector sampled

from the multivariate normal distribution of the mutational covariance matrix M and l is the

branch length. Sampling was performed using the *rmvnorm* function in the R package *mvtnorm*

[77]. The starting value of each trait is 0 in all simulations. The phenotypic value of each strain

was obtained by adding up the amount of evolution on all branches ancestral to the strain. This

was repeated 1,000 times to generate 1,000 datasets.

To account for the difference in V_eigen caused by different sample sizes in estimating the

correlation matrices, we randomly sampled subsets of the gene deletion strains. Because the

evolutionary correlation matrix has a rank number of 15 and has 15 positive eigenvalues, each

subset consists of 15 strains randomly drawn from the 4718 gene deletion strains such that the

mutational correlation matrix computed from each subset of mutants also has 15 positive

eigenvalues. From each subset of strains, we computed V_eigen, leading to a null distribution of

V_eigen. The observed V_eigen from the evolutionary correlation matrix is then compared with the

null distribution; a significant difference is inferred if the observed value falls in either the left or

right 2.5% tail.

To test whether there exists a significant modular structure among traits, we performed

the covariance ratio (CR) test. For each pair of predefined modules, traits were first re-ordered

such that traits belonging to each module were located in the upper-left and lower-right corners

of the covariance matrix, respectively, and $CR = \sqrt{\dfrac{trace(M_{12}M_{21})}{\sqrt{trace(M_{11}^*M_{11}^*)+trace(M_{22}^*M_{22}^*)}}}$, where M_{12} and

M_{21} are the upper-right and lower-left sections of the original covariance matrix, respectively,

containing all between-module covariances, M_{11}^* is the upper-left section with diagonal elements

replaced by zeros, M_{22}^* is the lower-right section with diagonal elements replaced by zeros, and

$trace(M)$ denotes the trace, or the sum of diagonal elements, of matrix M [58]. Because three

modules were defined in the yeast data, the average of all pairwise CR values was used to

represent the overall modularity. A test for selection on *CR* was performed following the test of

selection on V_{eigen}.

Computer simulation of trait-trait coevolution under selection

In each simulation, we considered a pair of traits with equal amounts of mutational

variance V_M, which is set to be 0.01. The mutational covariance matrix is thus $M =$

$\begin{bmatrix} V_M & COV_M \\ COV_M & V_M \end{bmatrix} \begin{bmatrix} V_M & V_M COR_M \\ V_M COR_M & V_M \end{bmatrix}$, where COV_M is the mutational covariance. The number

of mutations is a random Poisson variable with the mean equal to 1. The phenotypic effect of a

mutation is drawn from the multivariate normal distribution of *M* using the *rmvnorm* function in

the R package *mvtnorm* [77]. The starting phenotype is (0, 0) in all simulations.

We considered a Gaussian fitness function of $f = exp\left(-\frac{D^2}{2}\right)$, where *f* is the fitness and

D is the distance between the current phenotype and the optimal phenotype. When there is a

single fitness peak (i.e., the fitness optimum is a single point), *D* is the Euclidean distance

defined by $\sqrt{d_1{}^2 + d_2{}^2}$, where d_1 and d_2 are the distances between the current phenotypic values

of the two traits and their corresponding optima, respectively. When there is a fitness ridge (i.e.,

the fitness optimum is a line), *D* is the shortest distance from the current phenotype to the fitness

ridge. The selection coefficient *s* equals $\frac{f}{f_{WT}} - 1$, where *f* and f_{WT} are the fitness values of the

mutant and wild type, respectively. The fixation probability of a newly arisen mutant is $P_f =$

$\frac{1 - exp\,(-2s)}{1 - exp\,(-2N_e s)}$ in a haploid population [78], where the effective population size N_e was set at 10^4.

After each unit time, the phenotypic effect of each mutation is added to the population mean with

a probability of $N_e P_f$; this probability is treated as 1 when $N_e P_f > 1$ or when there is no

selection as in the latter case $P_f = \frac{1}{N_e}$. Combinations of parameters used in the simulations are

listed in Table 2.

In simulations where different lineages were assigned different optima, each lineage's

optimum was obtained by independently drawing the optimal values of the two traits from the

standard normal distribution. Before conducting simulations, we confirmed that the optima of

the two traits are not correlated (correlation coefficient = 0.0882, *P* = 0.54, *t*-test).

Computer simulation of trait-trait coevolution under mutational bias

To investigate the effect of mutational bias on trait correlation, we introduced the bias

coefficient B. Each mutation, after being sampled from a multivariate normal distribution

described above, was rescaled using B. Let the mutational effect be $m = (m_1, m_2)$, where m_1 and

m_2 are the effects on trait 1 and trait 2, respectively. The rescaled mutational effect, $\tilde{m}$, is

obtained by

553
$$\tilde{m} = \begin{cases} mB & (m_1 > 0) \\ \dfrac{m}{B} & (m_1 < 0) \end{cases}$$

Because mutational effects are first drawn from a pre-set multivariate normal distribution

and then rescaled, we examined if COR_M estimated from the rescaled effects ($\widehat{COR_M}$) is different

from the pre-set value of COR_M. For each pre-set value of COR_M, we obtained

$\widehat{COR_M}$ from 5,000 rescaled mutations. This was repeated 200 times with different random

mutations, yielding 200 $\widehat{COR_M}$ estimates. A series of different B values were used in the

simulation (**Table S2**). For comparison, we also estimated B from yeast gene deletion lines and

found the maximal B of any trait to be 1.503. To estimate B for a trait from the yeast gene

deletion lines, we respectively calculated the mean trait value of all deletion lines with positive

trait values and mean trait value of all deletion lines with negative values. We then computed the

ratio of their absolute values with the greater absolute value used as the numerator. The square

root of the ratio is B. We found that COR_M is always near the center of the distribution of these

200 $\widehat{COR_M}$ estimates (**Table S2**). Hence, mutational bias will not bias our test.

All analyses in this study were conducted in R [79].

LIST OF ABBREVIATIONS

MA: mutation accumulation.

CR: covariance ratio.

DECLARATION

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

580   The datasets generated and/or analysed during the current study are available at

581   https://github.com/RexJiangEvoBio/Trait-Correlation.

Competing interest

584   The authors declare that they have no competing interests.

Funding

587   This work was supported by U.S. National Institutes of Health grant R35GM139484 to

588   J.Z. D.J. was supported by Predoctoral Fellowship of Rackham Graduate School, University of

589   Michigan while working on the project.

Authors' contributions

592   D.J. and J.Z. designed the study. D.J. performed the analyses and prepared all figures.

593   D.J. and J.Z. wrote the paper.

Acknowledgements

596   We thank members of the Zhang lab and Dr. Matthew Pennell for valuable comments.

REFERENCES

1.   Gould SJ: **Allometry and Size in Ontogeny and Phylogeny**. *Biol Rev* 1966, **41**(4):587-640.
2.   Lande R: **Quantitative Genetic Analysis of Multivariate Evolution, Applied to Brain - Body Size Allometry**. *Evolution* 1979, **33**(1):402-416.
3.   Lande R: **The Genetic Covariance between Characters Maintained by Pleiotropic Mutations**. *Genetics* 1980, **94**(1):203-215.
4.   Wagner GP: **Multivariate Mutation-Selection Balance with Constrained Pleiotropic Effects**. *Genetics* 1989, **122**(1):223-234.
5.   Wagner GP, Zhang J: **The pleiotropic structure of the genotype-phenotype map: the evolvability of complex organisms**. *Nature Reviews Genetics* 2011, **12**(3):204-213.

6.   Dugand RJ, Aguirre JD, Hine E, Blows MW, McGuigan K: **The contribution of**
mutation and selection to multivariate quantitative genetic variance in an outbred
population of Drosophila serrata. *Proc Natl Acad Sci U S A* 2021, **118**(31).

7.   Ho WC, Zhang J: **The genotype-phenotype map of yeast complex traits: basic**
parameters and the role of natural selection. *Mol Biol Evol* 2014, **31**(6):1568-1580.

8.   McGuigan K, Collet JM, McGraw EA, Ye YH, Allen SL, Chenoweth SF, Blows MW:
The nature and extent of mutational pleiotropy in gene expression of male
Drosophila serrata. *Genetics* 2014, **196**(3):911-921.

9.   Houle D, Fierst J: **Properties of Spontaneous Mutational Variance and Covariance**
for Wing Size and Shape in Drosophila Melanogaster. *Evolution* 2013, **67**(4):1116-
1130.

10.   McGuigan K, Collet JM, Allen SL, Chenoweth SF, Blows MW: **Pleiotropic mutations**
are subject to strong stabilizing selection. *Genetics* 2014, **197**(3):1051-1062.

11.   Lande R: **The maintenance of genetic variability by mutation in a polygenic**
character with linked loci (Reprinted). *Genetics Research* 2007, **89**(5-6):373-387.

12.   Gardner KM, Latta RG: **Shared quantitative trait loci underlying the genetic**
correlation between continuous traits. *Mol Ecol* 2007, **16**(20):4195-4209.

13.   Saltz JB, Hessel FC, Kelly MW: **Trait Correlations in the Genomics Era**. *Trends Ecol*
Evol 2017, **32**(4):279-290.

14.   Felsenstein J: **Phylogenies and the Comparative Method**. *Am Nat* 1985, **125**(1):1-15.

15.   Roff DA, Mostowy S, Fairbairn DJ: **The evolution of trade-offs: Testing predictions**
on response to selection and environmental variation. *Evolution* 2002, **56**(1):84-95.

16.   Sinervo B, Svensson E: **Correlational selection and the evolution of genomic**
architecture. *Heredity (Edinb)* 2002, **89**(5):329-338.

17.   Shoval O, Sheftel H, Shinar G, Hart Y, Ramote O, Mayo A, Dekel E, Kavanagh K, Alon
U: **Evolutionary trade-offs, Pareto optimality, and the geometry of phenotype space**.
Science 2012, **336**(6085):1157-1160.

18.   Bolstad GH, Cassara JA, Marquez E, Hansen TF, van der Linde K, Houle D, Pelabon C:
Complex constraints on allometry revealed by artificial selection on the wing of
Drosophila melanogaster. *P Natl Acad Sci USA* 2015, **112**(43):13284-13289.

19.   Svensson EI, Arnold SJ, Burger R, Csillery K, Draghi J, Henshaw JM, Jones AG, De
Lisle S, Marques DA, McGuigan K *et al*: **Correlational selection in the age of**
genomics. *Nat Ecol Evol* 2021.

20.   Svensson EI: **Multivariate selection and the making and breaking of mutational**
pleiotropy. *Evol Ecol* 2022, **36**:807-828.

21.   Wagner GP, Altenberg L: **Perspective: Complex Adaptations and the Evolution of**
Evolvability. *Evolution* 1996, **50**(3):967-976.

22.   Olson EC, Miller RL: **Morphological integration**, Pbk. edn. Chicago: University of
Chicago Press; 1999.

23.   Pigliucci M: **Phenotypic integration: studying the ecology and evolution of complex**
phenotypes. *Ecol Lett* 2003, **6**(3):265-272.

24.   Kingsolver JG, Hoekstra HE, Hoekstra JM, Berrigan D, Vignieri SN, Hill CE, Hoang A,
Gibert P, Beerli P: **The strength of phenotypic selection in natural populations**. *Am*
Nat 2001, **157**(3):245-261.

25.   Goswami A, Smaers JB, Soligo C, Polly PD: **The macroevolutionary consequences of**
655   **phenotypic integration: from development to deep time**. *Philos Trans R Soc Lond B*
656   *Biol Sci* 2014, **369**(1649):20130254.
657   26.   Porto A, Sebastiao H, Pavan SE, VandeBerg JL, Marroig G, Cheverud JM: **Rate of**
658   **evolutionary change in cranial morphology of the marsupial genus Monodelphis is**
659   **constrained by the availability of additive genetic variation**. *J Evol Biol* 2015,
660   **28**(4):973-985.
661   27.   Simon MN, Machado FA, Marroig G: **High evolutionary constraints limited adaptive**
662   **responses to past climate changes in toad skulls**. *Proc Biol Sci* 2016, **283**(1841).
663   28.   Watanabe A, Fabre AC, Felice RN, Maisano JA, Muller J, Herrel A, Goswami A:
664   **Ecomorphological diversification in squamates from conserved pattern of cranial**
665   **integration**. *Proc Natl Acad Sci U S A* 2019, **116**(29):14688-14697.
666   29.   Fabre AC, Bardua C, Bon M, Clavel J, Felice RN, Streicher JW, Bonnel J, Stanley EL,
667   Blackburn DC, Goswami A: **Metamorphosis shapes cranial diversity and rate of**
668   **evolution in salamanders**. *Nat Ecol Evol* 2020, **4**(8):1129-1140.
669   30.   Navalon G, Marugan-Lobon J, Bright JA, Cooney CR, Rayfield EJ: **The consequences**
670   **of craniofacial integration for the adaptive radiations of Darwin's finches and**
671   **Hawaiian honeycreepers**. *Nature Ecology & Evolution* 2020, **4**(2):270-+.
672   31.   Sih A, Bell A, Johnson JC: **Behavioral syndromes: an ecological and evolutionary**
673   **overview**. *Trends Ecol Evol* 2004, **19**(7):372-378.
674   32.   Dochtermann NA, Dingemanse NJ: **Behavioral syndromes as evolutionary**
675   **constraints**. *Behav Ecol* 2013, **24**(4):806-811.
676   33.   Martin RD: **Relative brain size and basal metabolic rate in terrestrial vertebrates**.
677   *Nature* 1981, **293**(5827):57-60.
678   34.   Brown JH, Gillooly JF, Allen AP, Savage VM, West GB: **Toward a metabolic theory of**
679   **ecology**. *Ecology* 2004, **85**(7):1771-1789.
680   35.   Glazier DS: **A unifying explanation for diverse metabolic scaling in animals and**
681   **plants**. *Biol Rev* 2010, **85**(1):111-138.
682   36.   Pettersen AK, White CR, Marshall DJ: **Metabolic rate covaries with fitness and the**
683   **pace of the life history in the field**. *P Roy Soc B-Biol Sci* 2016, **283**(1831).
684   37.   White CR, Marshall DJ, Alton LA, Arnold PA, Beaman JE, Bywater CL, Condon C,
685   Crispin TS, Janetzki A, Pirtle E *et al*: **The origin and maintenance of metabolic**
686   **allometry in animals**. *Nat Ecol Evol* 2019, **3**(4):598-603.
687   38.   Schluter D: **Adaptive radiation along genetic lines of least resistance**. *Evolution* 1996,
688   **50**(5):1766-1774.
689   39.   Steppan SJ, Phillips PC, Houle D: **Comparative quantitative genetics: evolution of the**
690   **G matrix**. *Trends Ecol Evol* 2002, **17**(7):320-327.
691   40.   Arnold SJ, Burger R, Hohenlohe PA, Ajie BC, Jones AG: **Understanding the Evolution**
692   **and Stability of the G-Matrix**. *Evolution* 2008, **62**(10):2451-2461.
693   41.   Walsh B, Blows MW: **Abundant Genetic Variation plus Strong Selection =**
694   **Multivariate Genetic Constraints: A Geometric View of Adaptation**. *Annu Rev Ecol*
695   *Evol S* 2009, **40**:41-59.
696   42.   Agrawal AF, Stinchcombe JR: **How much do genetic covariances alter the rate of**
697   **adaptation?** *P Roy Soc B-Biol Sci* 2009, **276**(1659):1183-1191.
698   43.   Blows MW, Mcguigan K: **The distribution of genetic variance across phenotypic**
699   **space and the response to selection**. *Mol Ecol* 2015, **24**(9):2056-2072.

44.   Walter GM, Aguirre JD, Blows MW, Ortiz-Barrientos D: **Evolution of Genetic**
Variance during Adaptive Radiation. *Am Nat* 2018, **191**(4):E108-E128.

45.   Wagner GP: **A research programme for testing the biological homology concept**.
Novartis Found Symp 1999, **222**:125-134; discussion 134-140.

46.   Wagner GP, Pavlicev M, Cheverud JM: **The road to modularity**. *Nature Reviews*
Genetics 2007, **8**(12):921-931.

47.   Collet JM, McGuigan K, Allen SL, Chenoweth SF, Blows MW: **Mutational Pleiotropy**
and the Strength of Stabilizing Selection Within and Between Functional Modules of
Gene Expression. *Genetics* 2018, **208**(4):1601-1616.

48.   Wang Z, Liao BY, Zhang JZ: **Genomic patterns of pleiotropy and the evolution of**
complexity. *P Natl Acad Sci USA* 2010, **107**(42):18034-18039.

49.   Ohya Y, Sese J, Yukawa M, Sano F, Nakatani Y, Saito TL, Saka A, Fukuda T, Ishihara
S, Oka S *et al*: **High-dimensional and large-scale phenotyping of yeast mutants**. *P*
Natl Acad Sci USA 2005, **102**(52):19015-19020.

50.   Geiler-Samerotte KA, Zhu YO, Goulet BE, Hall DW, Siegal ML: **Selection Transforms**
the Landscape of Genetic Variation Interacting with Hsp90. *Plos Biol* 2016, **14**(10).

51.   Yvert G, Ohnuki S, Nogami S, Imanaga Y, Fehrmann S, Schacherer J, Ohya Y: **Single-**
cell phenomics reveals intra-species variation of phenotypic noise in yeast. *BMC Syst*
Biol 2013, **7**:54.

52.   Ho WC, Ohya Y, Zhang JZ: **Testing the neutral hypothesis of phenotypic evolution**. *P*
Natl Acad Sci USA 2017, **114**(46):12219-12224.

53.   Houle D, Bolstad GH, van der Linde K, Hansen TF: **Mutation predicts 40 million years**
of fly wing evolution. *Nature* 2017, **548**(7668):447-+.

54.   Liti G, Carter DM, Moses AM, Warringer J, Parts L, James SA, Davey RP, Roberts IN,
Burt A, Koufopanou V *et al*: **Population genomics of domestic and wild yeasts**. *Nature*
2009, **458**(7236):337-341.

55.   Peter J, De Chiara M, Friedrich A, Yue JX, Pflieger D, Bergstrom A, Sigwalt A, Barre B,
Freel K, Llored A *et al*: **Genome evolution across 1,011 Saccharomyces cerevisiae**
isolates. *Nature* 2018, **556**(7701):339-344.

56.   Hodgins-Davis A, Duveau F, Walker EA, Wittkopp PJ: **Empirical measures of**
mutational effects define neutral models of regulatory evolution in Saccharomyces
cerevisiae. *Proc Natl Acad Sci U S A* 2019, **116**(42):21085-21093.

57.   Pavlicev M, Cheverud JM, Wagner GP: **Measuring Morphological Integration Using**
Eigenvalue Variance. *Evol Biol* 2009, **36**(1):157-170.

58.   Adams DC: **Evaluating modularity in morphometric data: challenges with the RV**
coefficient and a new test measure. *Methods Ecol Evol* 2016, **7**(5):565-572.

59.   Hansen TF, Houle D: **Measuring and comparing evolvability and constraint in**
multivariate characters. *J Evolution Biol* 2008, **21**(5):1201-1219.

60.   Houle D, Jones LT, Fortune R, Sztepanacz JL: **Why does allometry evolve so slowly?**
Integr Comp Biol 2019, **59**(5):1429-1440.

61.   Geiler-Samerotte KA, Li S, Lazaris C, Taylor A, Ziv N, Ramjeawan C, Paaby AB, Siegal
ML: **Extent and context dependence of pleiotropy revealed by high-throughput**
single-cell phenotyping. *Plos Biol* 2020, **18**(8).

62.   Wagner A: **Robustness and evolvability in living systems**. Princeton, NJ: Princeton
University Press; 2005.

63. Jones AG, Arnold SJ, Burger R: **Evolution and stability of the G-matrix on a**
landscape with a moving optimum. *Evolution* 2004, **58**(8):1639-1654.
64. Clune J, Mouret JB, Lipson H: **The evolutionary origins of modularity**. *Proc Biol Sci*
2013, **280**(1755):20122863.
65. Ackermann RR, Cheverud JM: **Detecting genetic drift versus selection in human**
evolution. *Proc Natl Acad Sci U S A* 2004, **101**(52):17946-17951.
66. Lynch M: **The Rate of Morphological Evolution in Mammals from the Standpoint of**
the Neutral Expectation. *Am Nat* 1990, **136**(6):727-741.
67. Jerison ER, Kryazhimskiy S, Mitchell JK, Bloom JS, Kruglyak L, Desai MM: **Genetic**
variation in adaptability and pleiotropy in budding yeast. *Elife* 2017, **6**.
68. Jones AG, Burger R, Arnold SJ: **Epistasis and natural selection shape the mutational**
architecture of complex traits. *Nat Commun* 2014, **5**:3709.
69. Pavlicev M, Kenney-Hunt JP, Norgard EA, Roseman CC, Wolf JB, Cheverud JM:
Genetic variation in pleiotropy: Differential epistasis as a source of variation in the
allometric relationship between long bone lengths and body weight. *Evolution* 2008,
62(1):199-213.
70. Pavlicev M, Cheverud JM: **Constraints Evolve: Context Dependency of Gene Effects**
Allows Evolution of Pleiotropy. *Annual Review of Ecology, Evolution, and Systematics,*
Vol 46 2015, **46**:413-434.
71. Wei X, Zhang J: **Environment-dependent pleiotropic effects of mutations on the**
maximum growth rate r and carrying capacity K of population growth. *Plos Biol*
2019, **17**(1):e3000121.
72. Ané C: **Analysis of comparative data with hierarchical autocorrelation**. *Annals of*
Applied Statistics 2008, **2**(3):1078-1102.
73. Uyeda JC, Zenil-Ferguson R, Pennell MW: **Rethinking phylogenetic comparative**
methods. *Syst Biol* 2018, **67**(6):1091-1109.
74. Pennell MW, Eastman JM, Slater GJ, Brown JW, Uyeda JC, FitzJohn RG, Alfaro ME,
Harmon LJ: **geiger v2.0: an expanded suite of methods for fitting macroevolutionary**
models to phylogenetic trees. *Bioinformatics* 2014, **30**(15):2216-2218.
75. Revell LJ, Harmon LJ, Langerhans RB, Kolbe JJ: **A phylogenetic approach to**
determining the importance of constraint on phenotypic evolution in the neotropical
lizard Anolis cristatellus. *Evol Ecol Res* 2007, **9**(2):261-282.
76. Benjamini Y, Hochberg Y: **Controlling the False Discovery Rate - a Practical and**
Powerful Approach to Multiple Testing. *J R Stat Soc B* 1995, **57**(1):289-300.
77. Genz AB, F.; Miwa, T.; Mi, X.; Leisch, F.; Scheipl, F.; Hothorn, T.: **mvtnorm:**
Multivariate Normal and t Distributions. R package version 1.1-0, https://CRAN.R-
project.org/package=mvtnorm. 2020.
78. Kimura M: **On the probability of fixation of mutant genes in a population**. *Genetics*
1962, **47**:713-719.
79. R Core Development Team: **R: A language and environment for statistical**
computing. 2010.
786

788
789

Table 1. Numbers of trait pairs with significantly different COR_E and COR_M in the yeast and fly data.

	Yeast						Fly (276 trait pairs)		
	COR_M from gene deletion lines (24,090 trait pairs)			COR_M from MA lines (17,391 trait pairs)					
	Observed[a]	Expected[b]	Fraction[c]	Observed	Expected	Fraction	Observed	Expected	Fraction
Strengthened	2727	145	3%	2395	17	3.5%	68	0	0%
Weakened	1221	78	1.4%	1348	6	0.1%	48	0	0%
Reversed	2795	51	0%	1403	2	0%	28	0	0%
Total	6743	302.5	0.7%	5146	31	0.7%	144	0	0%

[a]Number of trait pairs with significantly different COR_M and COR_E inferred by a Z-test.

[b]Number of trait pairs that show a significant difference between COR_M and COR_E derived from neutral Brownian motion simulations. The median from 1000 simulations is shown.

[c]Fraction of the 1000 Brownian motion simulations where the number of trait pairs with significantly different COR_M and COR_E exceeds the number under the "observed" column.

798   Table 2. Parameters and results of simulations of trait-trait coevolution.

Optimum	COR_M	Median COR_E at the end of simulation	Fraction of simulations with $COR_E > COR_M$	COR_E compared with COR_M
No optimum	0.9	0.900	49.5%	No difference
	0.5	0.495	47.5%	No difference
	0.1	0.113	55.5%	No difference
$y = x$ *	0.9	1.000	100%	Strengthened
	0.5	1.000	100%	Strengthened
	0.1	1.000	100%	Strengthened
$y = 0.5x$	0.9	1.000	100%	Strengthened
	0.5	1.000	100%	Strengthened
	0.1	1.000	100%	Strengthened
$y = -0.5x$	0.9	-0.995	0%	Reversed
	0.5	-0.999	0%	Reversed
	0.1	-1.000	0%	Reversed
$y = -x$	0.9	-0.997	0%	Reversed
	0.5	-0.999	0%	Reversed
	0.1	-1.000	0%	Reversed
(0, 0)	0.9	0.0213	0%	Weakened
	0.5	0.00142	0.5%	Weakened
	0.1	-0.0109	24%	No difference
Drawn from $N(0, 1)$	0.9	0.0895	0%	Weakened
	0.5	0.0874	0%	Weakened
	0.1	0.0866	6%	No difference

*x and y respectively represent the values of the two traits considered.

Table 3. Overall phenotypic integration (V_{eigen}) and modularity (CR) at the levels of mutation
and evolutionary divergence. Values at the level of mutation for yeast are medians of 1,000
control datasets. P-values for yeast are computed from locations of the observed values in the
corresponding distributions of 5000 control datasets, while the P-value for fly is from a
Fligner-Killeen test.

Statistic	Taxon	Mutation	Divergence	P-value
V_{eigen}	Yeast	34.414	58.656	0.0788
	Fly	3.530	4.359	0.459
CR	Yeast	0.649	0.997	< 0.001

808
809
810

Figure 1. Mutational (COR_M) and evolutionary (COR_E) correlations for all pairs of the 220 yeast

morphological traits. COR_M is based on yeast gene deletion lines. (A) COR_M (upper triangle)

and COR_E (lower triangle) for all pairs of traits ordered according to their IDs. (B) Frequency

distributions of COR_M and COR_E across all trait pairs. The two distributions are significantly

different ($P < 10^{-10}$, Kolmogorov–Smirnov test).

Figure 2. Examples of yeast trait pairs with COR_E significantly different from the corresponding

COR_M. (A) An example of evolutionarily strengthened correlation. (B) An example of

evolutionarily weakened correlation. (C) An example of evolutionarily reversed correlation.

Each blue dot represents a gene deletion line (a.k.a. mutant) while each red dot represents an

independent contrast derived from natural strains. Blue and red lines are linear regressions

between the standardized values of the two traits in mutants and independent contrasts,

respectively, while the dotted blackline shows the diagonal ($y = x$). Trait IDs are shown along

the axes. All COR_M and COR_E values shown are significantly different from 0 except when

indicated by "NS" in the parentheses.

Figure 3. Schematic illustration of predictions made by models of trait-trait coevolution. Each

circle represents the equilibrium mean phenotype of two hypothetical traits (trait 1 and trait 2) of

a diverging lineage. (A) When a specific allometric relationship is selectively favored, the

population mean phenotypes are distributed along the fitness ridge (i.e., the optimal allometric

line shown in red), resulting in a strong trait correlation across lineages. (B) When a specific

value is selectively favored for each trait, the population mean phenotypes are concentrated near

the optimal phenotype (marked by the red cross) and the trait correlation across lineages is weak.

(C) When different lineages have different optimal phenotypes (marked by red crosses) that are

randomly distributed, the trait correlation across lineages is weak.

837
838

839 **Figure 1**

841 A

843 B

844
845
846
847

Figure 2

A

B

C

Figure 3

A

859
860 B

861
862 C