Title: Flexible foraging behaviour increases predator vulnerability to climate change

Author list: Benoit Gauzens[1,2,*], Benjamin Rosenbaum[1,2], Gregor Kalinkat[3], Thomas Boy[1,2],

Malte Jochum[4,5,6], Susanne Kortsch[7], Eoin J. O'Gorman[8], Ulrich Brose[1,2].

Affiliations:

[1] EcoNetLab, German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-

Leipzig, Leipzig, Germany

[2] Institute of Biodiversity, Friedrich Schiller University Jena, Jena, Germany

[3] Department of Community and Ecosystem Ecology, Leibniz Institute of Freshwater

Ecology and Inland Fisheries (IGB), Berlin, Germany

[4] Experimental Interaction Ecology, German Centre for Integrative Biodiversity Research

(iDiv) Halle-Jena-Leipzig, Leipzig, Germany

[5] Leipzig University, Institute of Biology, Leipzig, Germany

[6] Department of Global Change Ecology, Biocenter, University of Würzburg, Würzburg,

Germany

Tvärminne Zoological Station, University of Helsinki, Hanko, Finland

[8] School of Life Sciences, University of Essex, Wivenhoe Park, Colchester CO4 3SQ, UK

* Corresponding author: benoit.gauzens@idiv.de

Abstract:

Higher temperatures are expected to reduce species coexistence by increasing energetic demands.

However, flexible foraging behaviour could balance this effect by allowing predators to target specific

prey species to maximize their energy intake, according to principles of optimal foraging theory. We

test these assumptions using a large dataset comprising 2,487 stomach contents from six fish species

with different feeding strategies, sampled across environments with varying prey availability over 12

years in Kiel Bay (Baltic Sea). Our results show that foraging shifts from trait- to density-dependent

prey selectivity in warmer and more productive environments. This behavioural change leads to lower

consumption efficiency at higher temperature as fish select more abundant but less energetically

rewarding prey, thereby undermining species persistence and biodiversity. By integrating this

behaviour into dynamic food web models, our study reveals that flexible foraging leads to lower

species coexistence and biodiversity in communities under global warming.

Main text:

Ecosystems are experiencing abrupt changes in climatic conditions, making it ever more important to

predict and understand how they will respond in the future. Global warming will affect various levels

of biological organization; from physiological processes occurring at the individual level[1] to patterns

at macroecological scales[2]. Warming impacts will cascade through these different organizational

levels, changing species composition[3], as well as community and food web structure[4]. By scaling up

temperature effects from species physiology to food webs[5], trophic interactions play a key role in an

ecosystem's response to global warming[6]. Therefore, food web models have become important tools

for predicting the future of communities amidst rising temperatures, and understanding the underlying

mechanisms[7]. However, current food web models are mostly based on networks composed of static

feeding interactions, thereby disregarding flexibility in consumer behaviour in response to varying

environmental conditions.

Food web models building on biological processes observed at the level of individual organisms can

be used to test mechanisms and generate predictions at the ecosystem level and therefore offer an

integrative solution to study temperature effects on communities. For example, Allometric Trophic

Networks[8] (ATNs) quantify effects of body mass and temperature on the biological rates of

consumers and resources to predict species biomass changes over time and across environmental

conditions[8–11]. Thus, ATNs facilitate our understanding of how physiological responses to warming

translate into species coexistence and biodiversity[7]. However, the ability of ATNs to derive sound

predictions for large communities under changing environmental conditions has been challenged,

stressing the need for more biological realism[6,12]. Indeed, a strong limitation of these models is that

species are solely characterized by a set of biological rates that respond to temperature, such as

metabolic or attack rates. Therefore, species are limited to physiological responses to warming,

whereas the behavioural component is largely ignored. However, it is well established that species

also respond to warming by changing their behaviour[13,14]. From a food web perspective, flexibility in

species' foraging behaviour is usually modelled using the principles of optimal foraging behaviour,

which posit that consumers will modify their diet to maximize their energy intake. Under this

hypothesis, models predict that behavioural changes help to support species coexistence in

communities. However, it remains uncertain whether the premise of optimal foraging holds true in the

context of rising temperatures and how behavioural changes prompted by warming translate into

altered species coexistence. Therefore, integrating behavioural flexibility into food web models is

critical to improve their accuracy in predicting the consequences of global warming[15–19].

Energetic demands increase with temperature, but species can offset them by adopting various

strategies to increase their energy intake. Species can actively forage on energetically more rewarding

resources[13,20], typically prey that are close to the maximum body mass that consumers can feed on[21].

Therefore, we expect that predators favour larger prey individuals (trait-based selectivity) at higher

temperatures, reducing predator-prey body mass ratios (hypothesis H1a). Alternatively, individuals

under high energetic stress may be driven by their increased demand for food and accept less

rewarding (smaller), but more abundant prey upon random encounter (hypothesis H2a) leading to a

reduced trait-based selectivity, and a trophic niche driven more by neutral processes (random

encounter probability). These two hypotheses would lead to contrasting effects on communities[22].

Trait-based selectivity may increase the interaction strengths between predators and larger prey,

which could increase their extinction risk due to the lower abundance of large organisms[23], therefore

having negative consequences on species coexistence (hypothesis H1b). Alternatively, if neutral

processes are found to be responsible for driving selectivity, predators would primarily forage on the

most abundant species, which could result in a more even distribution of biomass across all prey size

classes, enhancing the likelihood of species coexistence[18,24] (hypothesis H2b). To test these

hypotheses, we used a dataset of 2,487 stomach contents from 6 demersal fish species[25,26] collected in

the Baltic Sea between 1968 and 1978. This data set compiles body mass and information for the fish

species, as well as body mass and biomass for the resource species observed in fish stomachs and in

environment. We used this dataset to analyse the behavioural responses of fish to changes in

temperature and resource availability (i.e., the total biomass of their prey). Subsequently, we

addressed the consequences of these empirical relationships by integrating them into a food web

model to predict how species coexistence changes with warming and the related foraging choices of

consumers.

Response to temperature and resource availability gradients

We used our database to document how consumer foraging behaviour responds to temperature and

resource availability. The six fish species considered belong to two functional groups differing in

body shape and foraging behaviour (flat, sit-and-wait predators versus fusiform, more active hunters).

To describe the prey body mass distributions observed in fish stomachs (hereafter called the *realized*

distribution) and in the environment (hereafter called the *environmental distribution*), we used

empirical medians and standard deviations (Fig. 1). The environmental distribution defines what is

expected if neutral processes drive fish diets: it represents the expected body mass distribution of the

consumer diets if consumption was driven by density-based encounter rates only. However, the size

distributions of prey in the environment and in consumer diets are usually not identical because

consumers actively select prey individuals with specific body masses. We used the ratio of the

realized and environmental distributions to calculate fish selectivity with respect to these different

prey body masses to obtain a *preference distribution* (see conceptual explanation in Fig. SI 1), which

describes consumer selectivity based on traits and consumer behavioural decisions (i.e. foraging

behaviour). While traits define the fundamental trophic niche of a species (what a consumer can eat),

behavioural decisions define the realised part of the fundamental niche. Therefore, a shift in behaviour

does not necessarily imply a shift in the identity of prey species, but can simply lead to a shift in the

individual traits that are selected, within or across different prey species. A more detailed description

of the body mass relationships observed in consumer stomachs can be found in SI II.

We tested how the median of the fish's preference distributions was affected by the local resource

availability and temperature conditions using a Bayesian linear model (note that we did not detect any

strong correlation between resource availability and temperature; Spearman correlation = -0.167). The

model included main effects for resource availability, temperature, fish functional group, and fish

body mass, and the interactions between resource availability and temperature and between fish

functional group and temperature. Including fish body mass as a covariate in the model ensures that

our results are independent of predator size. We compared the full model to versions without the

interaction between fish functional group and temperature and without fish functional group as a

covariate using a "Leave One Out Cross Validation" approach (LOO-CV)[27]. The most parsimonious

model used for the subsequent analyses was the one that did not include fish functional group or its

interaction with temperature (results presented in SI III), indicating that the behavioural responses to

temperature and resource availability were similar for fish species with different body shapes and

foraging strategies. By contrast, we observed a significant interaction between resource availability

and temperature on the median of the preference distribution (Table 1, Fig. 2) suggesting that

behavioural responses of fish to temperature depend on environmental resource availability levels.

By examining the effect size and significance of the temperature effect along the resource availability

gradient in our dataset, we found that the temperature effect is only significant at high levels of

resource availability (Fig. 3). Here, fish tend to forage on smaller prey as temperature increases. This

indicates that fish only change their feeding behaviour to temperature by foraging on smaller prey in

warmer conditions (so supporting H2a against H1a) when resources are plentiful

The energetic stress that warming imposes on individuals through increased metabolic rates should be

mitigated by higher feeding rates at higher prey availability in more productive environments. Thus,

because the effects of temperature and resource availability should cancel each other out, we expected

a stronger behavioural response at low resource levels, where consumers must cope with maximum

energetic stress (regardless of temperature). Under such stressful conditions, there may be no scope

for predators to change their feeding behaviour as temperature increases, especially in the Baltic Sea

where growth rates of fish species tend to be limited by resource availability in general[28,29].

In more productive environments (See SI IV for a description of resource availability and of prey

body mass structure), feeding behaviour may be less constrained, increasing the behavioural

flexibility of the fish. We also observed that the width of the fish preference distribution decreases

with temperature in the most productive environments (Table 2, Extended data 1), which means that

this reduction in selected prey body masses is not explained by neutral processes but instead comes

from fish actively foraging for smaller prey when temperature increases. This result challenges our

hypothesis H2a, which suggested that foraging on smaller prey would be associated with a decrease in

selectivity, but could be explained by the commonly observed type III functional response[23,30]. Here,

when temperature increases consumers gradually shift their focus on more abundant resources

(corresponding to smaller species) which comes at the cost of ignoring the less abundant, bigger ones,

leading to a reduction in the width of the consumer trophic niche. With this behaviour, fish can satisfy

their immediate demand for energy by targeting abundant species, but at the cost of missing out on

larger and thus energetically more rewarding prey individuals, which can be critical to satisfy their

energetic demands in the long term[31]. Indeed, fish metabolic rates increase with warming over large

temperature gradients[32] and do so faster than their feeding rates[33], which can lead to the extinction of

top predators due to starvation[34]. Combining this physiological starvation effect with our observed

behavioural response indicates that consumption outside of the most efficient predator-prey body

mass ratio should reduce energy flux through food webs, limiting the coexistence of consumer

species[31,35]. The combination of physiological and behavioural warming effects could thus increase

the likelihood of top predator extinction in food webs, which are usually considered key species for

maintaining biodiversity and ecosystem functioning[36].

Consequences for species coexistence under global warming

Flexible foraging in response to varying local conditions is considered key to promote species

coexistence[18,19,37]. The general assumption behind this notion is that consumer species will adapt their

foraging strategies in order to maximize their energetic gains[38]. However, our results suggest that

consumers tend to depart from this optimal foraging behaviour under stressful conditions. We

explored the consequences of this behaviour using a dynamic population model, which predicts the

temporal dynamics and coexistence of species in food webs (see Methods). We ran two versions of

this model: one including changes of species diets based on local temperature and resource

availability conditions as informed by our empirical results, and one model with static feeding links

representing the classical food-web modelling approach. We simulated the dynamics for synthetic

food webs of 50 species (30 consumers and 20 basal species) over a temperature gradient spanning

from 0°C to 18°C to predict the number of species extinctions at different temperatures. Overall, we

observed that models incorporating flexible foraging were more sensitive to warming, with more

consumer extinctions over the temperature gradient (Fig. 4) and relatively few extinctions of basal

resources (results presented in Fig. SI 6). These results were not affected by the functional response

type, which is a free parameter in our modelling approach, but tended to weaken at very low levels of

nutrient availability (i.e. productivity), consistent with our empirical results (model predictions for the

different values are available in SI VI).

Generally, food web models incorporating foraging behaviour are based on optimal foraging theory

and thus lack a data-driven description of how the selectivity of consumer diets changes in a natural

context. To address this gap, we developed a trait-based framework to document the response of

foraging behaviour to temperature, which can be incorporated into predictive models of food web

structure and species coexistence. Our approach can be generalized to other ecological variables that

affect food webs and foraging behaviour, such as fear of predators or habitat complexity[39]. Finally, the

effects documented here come from data sampled at rather low levels of temperature and resource

availability. Therefore, it will be crucial to extend our regression models to warmer and more

productive ecosystems to assess whether very high levels of productivity could balance the energetic

stress related to warming, thus limiting behavioural responses in eutrophic environments.

A potential limitation of our approach is that the variations in temperature we employed were

primarily driven by seasonal changes, which may not fully reflect the long-term dynamics associated

with climate change as well as indirect effects through modification of habitat structure potentially

interacting with the foraging behaviour of consumers or lowering oxygen concentrations. Similarly,

response of social behaviour, like aggregation of predator or prey could influence the shape of the

functional response but are rarely considered in the literature. Nevertheless, our hypotheses and

interpretations are rooted in energetic budget calculations that stem from fundamental mechanistic

principles concerning the impact of temperature and prey body masses on population growth rates.

This mechanistic framework enabled us to formulate generalisable predictions regarding temperature

effects across different ecological contexts. Consequently, we propose a testable explanation for the

observed decline in biomass among the fish populations in the Baltic Sea[40] . However, it remains

possible that fish could exhibit increased fitness by consuming smaller, more abundant organisms

under certain conditions. Hence, these predictions should now be confronted with a comprehensive

examination of how the fitness of fish changes depending on temperature and behavioural choices, for

example in more controlled experimental settings (e.g., mesocosms). Such experimental approaches

could also consider other mechanisms leading to shifts in species diets and interactions in the context

of temperature increase, e.g., ontogenetic shits[41].

The effects of warming on the trait structure of communities and the distribution of trophic

interactions[42] are well documented, but a framework for integrating changes in feeding behaviour with

a general modelling approach has been lacking. Closing this gap, our results stress the importance of

accounting for foraging behaviour, to better understand and predict community responses to climate

change, and challenge previous conclusions on this topic. Indeed, the discrepancies between the

models with and without flexible foraging suggest that the classical approach, which only accounts for

changes in species physiology[6,7] completely ignoring behavioural aspects, may have overlooked a

significant portion of community responses to warming. Importantly, our results show that, contrary

to common expectation, behavioural responses to climatic stress can reduce the likelihood of species

coexistence and thus decrease community biodiversity. The similarity in the responses of the two fish

feeding strategies (sit-and-wait and active foraging) indicates some generality in our results, but it is

important to investigate these patterns for a wider range of species and ecosystem types in future

studies. For instance, consumer metabolic type has an important effect on the response of species to

temperature[43] and endotherms could respond differently than ectotherms such as fish.

Conclusion

It is generally assumed that consumers respond to environmental conditions by making choices that maximize their energy intake[19,44]. This assumption has been used to derive several predictions in ecology about community structure and species coexistence, and is often considered a solution to May's paradox[45] that highly diverse communities are mostly mathematically infeasible, despite their widespread occurrence in nature. It is therefore commonly assumed that behaviour not included in these simple mathematical models is a strong driver of community organization and supports species coexistence[19]. We challenge this optimistic view of nature by demonstrating how consumer species can shift to less efficient foraging behaviour under stressful conditions, for instance when resources are scarce and when they face additional energetic stress due to warming. Therefore, the ecological conclusions built into the assumptions that flexible behaviour favours coexistence do not necessarily hold in the context of global warming. Our mechanistic modelling approach demonstrates the consequences of this observation, with more species extinctions in response to warming when flexible foraging is considered. This indicates that global warming may lead to a greater reduction in species coexistence than predicted by classical ecological models. Our findings thus challenge the general paradigm that flexible foraging should mitigate the consequences of global warming for natural ecosystems and call for a general data-driven theory approach to forecasting biodiversity and functioning in future ecosystems.

Acknowledgements: We are profoundly grateful that Wolf E. Arntz collected and provided the
valuable data set from his early work in Kiel Bay that we used in this study. We are also thankful to
Astrid Jarre who digitized the stomach content data, Ute Jacob for her help in the early phase of this
project and Carlos Melian for his friendly review of the manuscript. We gratefully acknowledge the
support of:
iDiv funded by the German Research Foundation (DFG–FZT 118, 202548816), BG, UB, BR, TB,
MJ.
German Academic Exchange Service (DAAD, 57070483), GK.
European Research Council (ERC) under the European Union's Horizon 2020 research and
innovation programme (grant agreement no. 677232), MJ.
273

274 **Author contributions:**

275 BG, BR, UB conceptualised the study. BG, GK, BR, TB where involved in data compilation and code
276 development and BG and BR analysed the data. BG wrote the first draft of the manuscript ad all authors
277 contributed to its revision.

279 **Competing interest:** Authors declare that they have no competing interests

Tables

Table 1: Response of the median prey body mass to predator body mass and environmental gradients. Presentation of mean estimates and uncertainties (CI95), as well as effective sample size for the different predictors of the model. Rhat values were all lower than 1.003 for all estimated parameters. Total number of samples was 4000.

Response of the median prey body mass of the preference distribution to

Predictors	Estimates	CI (95%)	Effective samples
Intercept	-1.06	-3.14 – 1.09	1646
Predator body mass	0.55	0.41 – 0.70	2090
Temperature	0.18	-0.04 – 0.38	2000
Resource availability	0.19	-0.59 – 0.90	2017
Temperature: Resource availability	-0.07	-0.14 – 0.01	1961
Observations	290		
R^2 Bayes	0.279		

Table 2: Response of the standard deviation of the body mass preference distribution to predator body mass and environmental gradients. Presentation of mean estimates and uncertainties (CI95), as well as effective sample size for the different predictors of the model. Rhat values were all lower than 1.002. Total number of samples was 4000.

Response of standard deviation of the preference body mass distribution to

Predictors	Estimates	CI (95%)	Effective samples
Intercept	-1.05	-1.92 – -0.17	1046
Predator body mass	0.19	0.12 – 0.25	2059
Temperature	0.13	0.04 – 0.21	1053
Resource availability	0.39	0.09 – 0.69	1053
Temperature: Resource availability	-0.05	-0.08 – -0.02	1050
Observations	290		
R^2 Bayes	0.157		

Figures Legends

Figure 1: Conceptual representation of temperature impacts on the preference distribution. The preference distribution of prey body masses (turquoise) is estimated from the environmental (distribution of prey body masses in the environment, purple) and realised (distribution of prey body masses in consumers' stomachs, yellow) body mass distributions (top panel). It represents how different prey body masses are selected by the consumers. Based on our two hypotheses, temperature increase can either: (H1, bottom left panel) lead consumers to preferentially select for larger species (trait-based selectivity), which can create an imbalance in the trophic fluxes (blue arrows): some species are preferentially selected even when their abundance is low, increasing their extinction risks and initiating large dynamical oscillations in species densities. Alternatively, temperature increase (H2, bottom right panel) leads consumers to have their diet more driven by encounter rates (density-based selectivity), which creates a stronger control of species with high biomass in comparison to less abundant, smaller ones, favouring coexistence of resource species and thus community species richness.

Figure. 2: Response of the median prey body mass of the preference distribution. Effect of (a) predator body mass and (b) temperature and resource availability. Points represent log-transformed data across all resource availability levels and lines represent model predictions. Regression lines represent model predictions on the median of the preferred distribution when all other covariates are considered. The shaded areas show the 95% confidence interval on the predicted values. Low and high resource availability values correspond to the two modes of the bimodal distribution of resource availability values (presented in SI IV).

Figure. 3: Effect of resource availability on the effect size of temperature on the median body mass of the preference distribution. The solid black line represents the effect size of the temperature effect calculated from the original model coefficients and the grey area shows the associated uncertainty (95% confidence interval). The vertical dashed line represents the resource availability threshold above which the temperature effect becomes significant (effect of 0 outside of the 95% confidence interval).

Figure. 4: Number of consumer species extinctions predicted by the model at different temperatures (out of an initial richness of 30 consumers). Points represent the number of observed extinctions for each simulation. The blue line represents the model prediction on the average number of extinctions without the response of species' foraging behaviour to local temperature and resource availability conditions considered, whilst the red line shows average predictions of extinctions when allowing for this flexibility. The shaded areas show the 95% confidence interval on the predicted values. Predictions were estimated using a GAM with a binomial link function.

References

1. Dell, A. I., Pawar, S. & Savage, V. M. Systematic variation in the temperature dependence of physiological and ecological traits. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 10591–10596 (2011).

2. Free, C. M. *et al.* Impacts of historical warming on marine fisheries production. *Science* **363**, 979–983 (2019).

3. Lenoir, J. *et al.* Species better track climate warming in the oceans than on land. *Nature Ecology and Evolution* **4**, 1044–1059 (2020).

4. Brose, U. *et al.* Predator traits determine food-web architecture across ecosystems. *Nature Ecology and Evolution* **3**, (2019).

5. Petchey, O. L., Brose, U. & Rall, B. C. Predicting the effects of temperature on food web connectance. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 2081–2091 (2010).

6. Gauzens, B., Rall, B. C., Mendonça, V., Vinagre, C. & Brose, U. Biodiversity of intertidal food webs in response to warming across latitudes. *Nature Climate Change* **10**, 264–269 (2020).

7. Binzer, A., Guill, C., Rall, B. C. & Brose, U. Interactive effects of warming, eutrophication and size structure: Impacts on biodiversity and food-web structure. *Global Change Biology* **22**, 220–227 (2016).

8. Martinez, N. D. Allometric Trophic Networks From Individuals to Socio-Ecosystems: Consumer–Resource Theory of the Ecological Elephant in the Room. *Frontiers in Ecology and Evolution* vol. 8 92 Preprint at https://doi.org/10.3389/fevo.2020.00092 (2020).

9. Boit, A., Martinez, N. D., Williams, R. J. & Gaedke, U. Mechanistic theory and modelling of complex food-web dynamics in Lake Constance. *Ecology Letters* **15**, 594–602 (2012).

10. Curtsdotter, A. *et al.* Ecosystem function in predator–prey food webs—confronting dynamic models with empirical data. *Journal of Animal Ecology* **88**, 196–210 (2019).

11. Hudson, L. N. & Reuman, D. C. A cure for the plague of parameters: constraining models of complex population dynamics with allometries. *Proceedings of the Royal Society B: Biological Sciences* **280**, 20131901 (2013).

12. Sauve, A. M. C. & Barraquand, F. From winter to summer and back: Lessons from the parameterization of a seasonal food web model for the Białowieża forest. *Journal of Animal Ecology* **89**, 1628–1644 (2020).

13. Sentis, A., Hemptinne, J. L. & Brodeur, J. Towards a mechanistic understanding of temperature and enrichment effects on species interaction strength, omnivory and food-web structure. *Ecology Letters* **17**, 785–793 (2014).

14. Abram, P. K., Boivin, G., Moiroux, J. & Brodeur, J. Behavioural effects of temperature on ectothermic

animals: unifying thermal physiology and behavioural plasticity. *Biological Reviews* **92**, 1859–1876

(2017).

15. Abrams, P. A. Dynamics and Interactions in Food Webs with Adaptive Foragers. in *Food Webs* 113–121

(Springer US, 1996). doi:10.1007/978-1-4615-7007-3_11.

16. Abrams, P. & Matsuda, H. Effects of adaptive predatory and anti-predator behaviour in a two-prey-one-

predator system. *Evolutionary Ecology* **7**, 312–326 (1993).

17. Valdovinos, F., Ramos-Jiliberto, R., Garay-Narva´ez, L., Pasquinell, U. & Dunne, J. A. Consequences of

adaptive behaviour for the structure and dynamics of food webs. *Ecology Letters* **13**, 1546–1559 (2010).

18. Kondoh, M. Does foraging adaptation create the positive complexity-stability relationship in realistic food-

web structure? *Journal of theoretical biology* **238**, 646–51 (2006).

19. Kondoh, M. Foraging adaptation and the relationship between food-web complexity and stability. *Science*

299, 1388–1391 (2003).

20. Lemoine, N. P., Drews, W. A., Burkepile, D. E. & Parker, J. D. Increased temperature alters feeding

behavior of a generalist herbivore. *Oikos* **122**, 1669–1678 (2013).

21. Portalier, S. M. J., Fussmann, G. F., Loreau, M. & Cherif, M. The mechanics of predator–prey interactions:

First principles of physics predict predator–prey size ratios. *Funct Ecol* **33**, 323–334 (2019).

22. Lazzaro, X., Lacroix, G., Gauzens, B., Gignoux, J. & Legendre, S. Predator foraging behaviour drives

food-web topological structure. *Journal of Animal Ecology* **78**, 1307–1317 (2009).

23. Kalinkat, G., Rall, B. C., Uiterwaal, S. F. & Uszko, W. Empirical evidence of type III functional responses

and why it remains rare. *Frontiers in Ecology and Evolution* **11**, 125 (2023).

24. Gauzens, B., Legendre, S., Lazzaro, X. & Lacroix, G. Intermediate predation pressure leads to maximal

complexity in food webs. *Oikos* **125**, 595–603 (2016).

25. Gauzens, B. & Kalinkat, G. Dietary information from the benthic community of the Western Baltic (1968-

1978). German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig

https://doi.org/10.25829/IDIV.3547-RTGQ13 (2023).

26. Gauzens, B. *et al.* Quantitative description of six fish species' gut contents and abundances of their prey in

the benthic food web of the Baltic Sea (1968-1978). 2023.12.08.570787 Preprint at

https://doi.org/10.1101/2023.12.08.570787 (2023).

27. Vehtari, A., Gelman, A. & Gabry, J. Practical Bayesian model evaluation using leave-one-out cross-

validation and WAIC. *Statistics and Computing* **27**, 1413–1432 (2017).

28. Elmgren, R. Man's Impact on the Ecosystem of the Baltic Sea: Energy Flows Today and at the Turn of the

Century. *Ambio* **18**, 326–332 (1989).

29. Elmgren, Ragnar. & Hill, Cathy. Ecosystem function at low biodiversity – the Baltic example. in *Marine*

Biodiversity: Patterns and Processes (eds. Gage, J. D., Angel, M. V. & Ormond, R. F. G.) 319–336

(Cambridge University Press, Cambridge, 1997). doi:10.1017/CBO9780511752360.015.

30. Uiterwaal, S. F. & DeLong, J. P. Functional responses are maximized at intermediate temperatures.

Ecology **101**, e02975 (2020).

31. Brose, U. Body-mass constraints on foraging behaviour determine population and food-web dynamics.

Functional Ecology **24**, 28–34 (2010).

32. Clarke, A. & Johnston, N. M. Scaling of metabolic rate with body mass and temperature in teleost fish. *J*

Anim Ecology **68**, 893–905 (1999).

33. Lindmark, M., Ohlberger, J. & Gårdmark, A. Optimum growth temperature declines with body size within

fish species. *Global Change Biology* **28**, 2259–2271 (2022).

34. Lindmark, M., Ohlberger, J., Huss, M. & Gårdmark, A. Size-based ecological interactions drive food web

responses to climate warming. *Ecol Lett* **22**, 778–786 (2019).

35. Guzman, L. M. & Srivastava, D. S. Prey body mass and richness underlie the persistence of a top predator.

Proc. R. Soc. B. **286**, 20190622 (2019).

36. Birkeland, C. & Dayton, P. K. The importance in fishery management of leaving the big ones. *Trends in*

Ecology & Evolution **20**, 356–358 (2005).

37. Stump, S. M. & Chesson, P. How optimally foraging predators promote prey coexistence in a variable

environment. *Theoretical Population Biology* **114**, 40–58 (2017).

38. Pyke, G. H., Pulliam, H. R. & Charnov, E. L. Optimal Foraging: A Selective Review of Theory and Tests.

The Quarterly Review of Biology **52**, 137–154 (1977).

39. Jinks, K. I. *et al.* Habitat complexity influences the structure of food webs in Great Barrier Reef seagrass

meadows. *Ecosphere* **10**, (2019).

40. Möllmann, C. *et al.* Tipping point realized in cod fishery. *Sci Rep* **11**, 14259 (2021).

41. Thunell, V., Lindmark, M., Huss, M. & Gårdmark, A. Effects of Warming on Intraguild Predator

Communities with Ontogenetic Diet Shifts. *The American Naturalist* **198**, 706–718 (2021).

42. O'Gorman, E. J. *et al.* A simple model predicts how warming simplifies wild food webs. *Nat. Clim. Chang.*

9, 611–616 (2019).

43. Pinsky, M. L., Eikeset, A. M., McCauley, D. J., Payne, J. L. & Sunday, J. M. Greater vulnerability to

warming of marine versus terrestrial ectotherms. *Nature* **569**, 108–111 (2019).

44. Petchey, O. L., Beckerman, A. P., Riede, J. O. & Warren, P. H. Size, foraging, and food web structure.

Proceedings of the National Academy of Sciences of the United States of America **105**, 4191–6 (2008).

45. May, R. M. Qualitative Stability in Model Ecosystems. *Ecology* **54**, 638–641 (1973).

443
444
445
Methods

The Kiel Bay database

To estimate how species body mass preferences change depending on local environmental conditions

we used the Kiel Bay database[26]. This publicly available database[25] compiles the stomach contents of

seven fish species (adult and sub-adult individuals) classified into two functional groups, based on

their body shape and habitat use: fusiform and benthopelagic species (*Gadus morhua* and *Merlangius*

merlangus) versus flat and demersal species (*Limanda*, *Pleuronectes platessa*, *Platichthys flesus*, and

Hippoglossoides platessoides). This shape characteristic also corresponds to specific foraging

behaviour[46], where flat fish tend to be "sit-and-wait" predators, while fusiform fish are more actively

foraging feeders. Together with taxonomic information, the database contains the body length of the

fish species (rounded to the nearest integer in cm) as well as the abundance and biomass of the

different prey species in fish stomachs and in their environments. Species in the environment were

sampled with van Veen grabs directly along the trawling routes used for fish sampling. In total, 6

sampling sites (3 for each trawling route) were used for the bottom grab sampling, each site being

sampled several times during the period covered by our dataset. To reduce biases associated with site

specificity of benthos communities (i.e., sampled with the van Veen grabs), data from sampling sites

were pooled together if they were harvested on the same date and corresponded to the same trawling

route.

The dataset also contains sea temperature data (representing monthly average taken at 20m depth).

For our analysis, we used a subset of the database for which it was possible to associate the content of

fish stomachs to the environmental abundances of their prey (see the method section "*Filtering data*

and association between fish and environments" for more details).

Estimation of fish body masses

To allow for correspondences between our allometric food web model (that relies on species body

masses for its parametrisation), the energetic approach discussed, and the empirical part of our study,

we converted the body lengths of fish to body masses. We used a power law[47]: $BM_i = aBL_i^b$, were BM_i

and BL_i represent the body mass and body length of fish i, respectively. a and b are constants that are

species- and location-specific. Values for our parameters a and b obtained from fishBase[48] are:

Gadus morhua: a = 0.00708, b = 3.08; *Merlangius merlangus*: a = 0.00631, b = 3.05; *Limanda*: a =

0.00776, b = 3.08; *Pleuronectes platessa*; a = 0.00776, b = 3.06; *Platichthys flesus*: a = 0.00776   , b

= 3.07; *Hippoglossoides platessoides*: a = 0.00562, b = 3.09; *Enchelyopus cimbrius*: a = 0.00389, b =

3.08.

Filtering data and association between fish and environments

To make comparisons between the distributions of prey observed in fish stomachs and the ones

observed in the environment, we only used a subset of the Kiel Bay database for which we were able

to associate information about a fish's stomach contents to information about the related prey in the

environment. We considered this association between fish and environment possible when they were

sampled in the same area (i.e. bottom grabs carried out directly on the route of a bottom trawl) and

within less than 31 days (see SI VII for a distribution of the differences). Then, associations were

made between fish and environments from the same location that minimised the time difference

between the fish and environmental sampling. This first filter reduced the number of fish used in our

analysis to 2,487. From this subset, we pooled all individuals from the same fish species occurring at

the same place on the same date (i.e., when they were harvested during the same sampling event) with

the same body mass (in our case, body mass is a discrete variable as it was estimated from body

length, rounded to the nearest integer in cm) into a unique entity for statistical analysis, which we

hereafter call "statistical fish". This choice is led by the allometric approach used in our analysis,

where all individuals from the same species and with the same body mass are considered identical.

This aggregation increases the quality of the estimation of the prey body mass distribution in

stomachs at the cost of a lower statistical power for the analyses done on the shape of these

distributions. For instance, with a high aggregation level, fewer data points are available to consider

the effect of temperature on the average body mass of prey. This approach is therefore conservative as

it reduces the probability of type 1 error. Lastly, to make sure that the sampling of the prey in the

environment was representative of the statistical fish's diet, we checked if the species composition in

the environment matched that of the fish stomachs (<future reference to the data paper>). This led to

the removal of 26 statistical fish where less than 90% of the biomass found in the diet corresponded to

species also found in the environment. This resulted in a final dataset of 290 statistical fish,

underpinned by 2,487 individuals. For our statistical analysis, we used fish body functional group as a

covariate instead of fish species, as models based on fish functional group were always found to be

more parsimonious (based on AIC).

Fitting of gut content and environmental distributions

We used empirical medians and standard deviations to describe all environmental distributions of

$\log_{10}$ body masses and realised distributions of each predator identity. Taxon-specific characteristics

of the prey, such as body toughness, could bias the dietary distributions towards prey containing shells

or skeletons. We assumed that prey with hard body parts are more likely to be detected in stomach

contents than species composed of soft tissues (due to their higher digestion time) and weighted their

occurrence by a correction factor of 0.8, (according to[49]) Overall, the trends and effects observed

when including this correction were similar to those observed without correction, thus suggesting an

absence of systematic biases (see SI VIII for an analysis without correction factor).

Determining allometric species' preferences

We assumed that a feeding event is defined by two independent probabilities: the probability for a

consumer to encounter a prey of a certain body size x (defined by the environmental distribution $E(x)$)

and the probability for a consumer to consume the prey when encountered (given by the preference

distribution $P(x)$). Then, the realised distribution is proportional to their product:

$R(x) \sim E(x) \cdot P(x)$

The preference distribution can therefore be expressed by the departure of the realised niche from the

environmental distribution, or by filtering out the effect of species environmental availability from the

realised distribution:

$P(x) \frac{R(x)}{E(x)}$

As such, the preference distribution is independent of seasonal changes in prey communities or any

other factors that could lead to an altered size distribution of prey in the environment.

Theoretically, it is possible to compute continuous distributions R and E from observed body masses

for the realised distribution, r_i ($i=1\ldots n$) and for the environmental distribution e_i ($i=1\ldots m$),

respectively, with e.g. kernel density estimation, and compute:

$P(x) = \frac{R(x)}{E(x)} / \int \frac{R(x)}{E(x)} dx$

We chose, however, a more conservative approach that requires just a kernel density estimate for the

environmental distribution $E(x)$: Moments (i.e., mean, standard deviation, and skewness) of $P(x)$ can

be computed as weighted moments of the observed realised body masses r_i with weights (i.e., the

contribution of the datapoint to the estimation of the moments) $w_i=1/E(r_i)$ being the inverse of

environmental abundances. Thus, realised body masses that are highly abundant in the environment

contribute less to the preference distribution, while those that are rare contribute more. Following[50]

and assuming $W = \sum_i w_i$, the mean μ, variance σ^2 and skewness γ of the preference distribution $P(x)$

are:

543
$$\mu = \frac{1}{W} \sum_i w_i r_i,$$

544
$$\sigma^2 = \frac{1}{W} \sum_i w_i (r|i - \mu)^2$$

$$\gamma = \frac{1}{W\sigma^3} \sum_i w_i (r|i - \mu)^3$$

To assess changes in the distributions and how they depart from each other, we used variations in the

point estimates (median and standard deviation).

Statistical analyses

To fit the preference distributions, we used a Bayesian linear model to explicitly test our hypotheses.

We started by predicting the median and standard deviation of the preference distributions using

temperature, resource availability, fish functional group, and fish body mass as covariates, as well as

interactions between temperature and functional group and temperature and resource availability.

Body masses used for the different distributions, as well as resource availability, were log-

transformed for visualisation purposes and to meet homoscedasticity assumptions from linear models.

The inclusion of fish body mass as a fixed effect in our model ensures that the other covariates are

corrected by fish body mass and therefore independent of it. We first checked if fish functional group

was an important predictor in our model using a "Leave-one-out" cross validation[27], and finally

simplified our model by removing fish functional group from the covariates (see SI III for a more

comparison of the different models and for the presentation of posterior distributions). The log-

transformation of the body mass and biomass variables were done to fulfil the assumptions of linear

models.

Dynamic model

To simulate the population dynamics, we used a previously published model[51], based on the Yodzis

and Innes framework[52]. The growth of consumer species B_i is determined by the balance between its

energetic income (predation) and its energetic losses (predation metabolism)

$\frac{dB_i}{dt} = e_P B_i \sum_j F_{ij} + e_A B_i \sum_j F_{ij} - \sum_j B_l F_{ji} - x_i B_i,$

where $e_p = 0.545$ and $e_a = 0.906$ represent the assimilation efficiency of a consumer foraging on plants

and animals, respectively[53]. x_i defines the metabolic rate of species i, which scales allometrically with

body mass:

$\quad x_i = x_0 m_i^{-0.25} e^{E_x \frac{T_0-T}{kT_0T}},$

$\quad$ where $x_0 = 0.314$ is the normalisation constant[51], $E_x = -0.69$ is the activation energy of metabolic rate[7],

$\quad$ k the Boltzmann constant, $T_0 = 293.15$ the reference temperature in Kelvin and T the temperature at

$\quad$ which the simulation is performed. The trophic interactions are determined using a functional

$\quad$ response F_{ij} that describes the feeding rate of consumer i over resource j:

$$F_{ij} = \frac{\omega_{ij} b_{ij} B_j^{1+q}}{1 + cB_i + \omega_{ij} \sum_k h_i b_{ik} B_k^{1+q}} \cdot \frac{1}{m_i}.$$

$\quad$ b_{ij} represent the species-specific capture and is determined by predator and prey body masses:

$\quad$ $b_{ij} = P_{ij} L_{ij}.$

$\quad$ It corresponds to the product of encounter probabilities P_{ij} by the probability that an encounter leads

$\quad$ to a realised predation event L_{ij}. As such, the parameters encode neutral processes (encounter

$\quad$ probabilities) and trait-based selectivity, as the distribution L_{ij} represents the fundamental trophic

$\quad$ niche of consumer i, i.e. the set of prey it can consume based on its traits. Both quantities are

$\quad$ determined by species body masses. We assume that encounter probability is more likely for species

$\quad$ with higher movement speeds of both consumer and resource species:

$$P_{ij} = p_0 m_i^{\beta_i} m_j^{\beta_j} e^{E_p \frac{T_0-T}{kT_0T}}.$$

$\quad$ Since movement speed scales allometrically with consumer and resource masses[54], we drew β_i and β_j

$\quad$ from normal distributions (consumer: $\mu_\beta = 0.47$, $\sigma_\beta = 0.04$, resource: $\mu_\beta = 0.15$, $\sigma_\beta = 0.03$), following[51].

$\quad$ Activation energy E_p is equal to -0.38, from[7]. L_{ij} is assumed to follow a Ricker curve[51], defined as:

$\quad$ $L_{ij} = \left(\frac{m_i}{m_j R_{opt}} e^{1 - \frac{m_i}{m_j R_{opt}}} \right)^\gamma,$

$\quad$ where the optimal consumer-resource body mass ratio $R_{opt} = 71.68$ was calculated from the observed

$\quad$ realised interactions in our dataset. We used a threshold $L_{ij} < 0.01$ under which values were set to 0,

$\quad$ assuming that consumers do not consider prey which are too small or too large. This fundamental

$\quad$ niche described by L_{ij} was used to generate our food webs, as we considered that species i consumes

$\quad$ species j when $L_{ij} > 0.01$. The handling time h_{ij} of i on j is defined as:

$\quad$ $h_{ij} = h_0 m_i^{\eta_i} m_j^{\eta_j} e^{E_h \frac{T_0-T}{kT_0T}},$

where the scaling constant h_0 was set to 0.4 and the allometric coefficients for η_i and η_j were drawn

from a normal distribution with mean and standard deviation of -0.48 and 0.03 for η_i and of -0.66 and

0.02 for η_j. E_h is equal to 0.26. c_i represents the species-specific interference competition defined as:

$c_i = c_0 e^{E_c \frac{T_0 - T}{k T_0 T}}$,

Where the scaling constant c_0 was drawn from a normal distribution of mean 0.8 and standard

deviation 0.2. The activation energy E_c is equal to -0.65. The hill exponent q is a free parameter

coding for the type of functional response and is set to 1.2 by default. The term w_{ij} informs on species

selectivity[55], describing the foraging effort of a given consumer on part of its fundamental niche

(based on traits only, as described by the L_{ij}). For the models without behavioural expectations, we

defined w_{ij} for every resource j as 1 over the number of prey species of consumer i. This

parametrisation, which corresponds to what is usually done in modelling studies, means that a

consumer will split its foraging effort equally among its prey, independent of the local environmental

conditions. When flexible behaviour was included in the model, the values of w_{ij} were determined by

the empirical preference distributions associated with fish. Our empirical distributions were

characterised by the 3 first moments (mean, standard deviation, and skewness) which can be used to

estimate the parameters of a skewed normal distribution, named location (ξ), scale (ω), and shape (α),

respectively. As for the mean and standard deviations, we used linear models to relate these

parameters to consumer body mass, as well as the interaction between resource availability and

temperature. The scale parameter (ω) is constrained to positive values, so we used a generalised linear

model with a log link function for this specific parameter. For each consumer, based on its body mass,

as well as local temperature and resource availability conditions, we predicted the complete

parametrisation of a skewed normal distribution. The preference of consumer i on resource j was

estimated with the values returned by the probability density functions associated to each

parameterisation for the different resource body masses.

To maintain the comparability with the model without flexible behaviour, the w_{ij} values were

transformed so that their sum was equal to 1 for each consumer. The biomass dynamic of the basal

species i is defined as:

$\frac{dB_i}{dt} = r_i G_i B_i - \sum_j B_j F_{ji} - x_i B_i,$

where $r_i = m_i^{-0.25}$ defines the species growth rate (E_r is the associated activation energy, set to 0.25).

G_i is the species-specific growth factor, determined by the concentration of two nutrients N_1 and N_2:

$G_i = min\left(\frac{N_1}{K_{i1}+N_1}, \frac{N_2}{K_{i2}+N_1}\right),$

Where K_{il} determines the half saturation density of plant i nutrient uptake rate, determined randomly

from a uniform distribution in [0.1, 0.2]. The dynamic of the nutrient concentrations is defined by:

$\frac{dN_l}{d_t} = D(S_l - N_l) - v_l \sum_i r_i G_i P_i,$

Where $D = 0.25$ determines the nutrient turnover rate and $S_l = 5$ determines the maximal nutrient

level. The loss of a specific nutrient N_l is limited by its relative content in plant species biomass ($v_1=1$,

$v_2=0.5$). We ran our model on food webs of 50 species, composed of 30 consumers and 20 basal

species. A link was drawn between two species i and j when $L_{ij} > 0$. For each temperature we ran 50

replicates of the two versions of the model (with and without flexible behaviour) using R 4.0.0-2 and

with an updated version of the ATNr package[56],using the "Unscaled ATN with nutrients" version. The

parameters used correspond to the one provided by the package and summarised in SI IX. We fitted a

GAM model on this number of extinctions. The code used for our analysis is available at[57]

https://zenodo.org/doi/10.5281/zenodo.10554928

642
643
644
645 **Data Availability**

646
647 The data used for this study can be accessed at:
648 https://doi.org/10.25829/idiv.3547-rtgq13 (see[25])

649
650 **Code Availability**

651
652 The code used for this study is available at:
653 https://zenodo.org/doi/10.5281/zenodo.10554928 (see[57])

Methods-only references

46. Russo, T., Costa, C. & Cataudella, S. Correspondence between shape and feeding habit changes throughout ontogeny of gilthead sea bream Sparus aurata L., 1758. *Journal of Fish Biology* **71**, 629–656 (2007).

47. Froese, R., Thorson, J. T. & Reyes, R. B. A Bayesian approach for estimating length-weight relationships in fishes. *Journal of Applied Ichthyology* **30**, 78–85 (2014).

48. Jobling, M. Mythical models of gastric emptying and implications for food consumption studies. *Environmental Biology of Fishes* **16**, 35–50 (1986).

49. Rimoldini, L. Weighted skewness and kurtosis unbiased by sample size and Gaussian uncertainties. *Astronomy and Computing* **5**, 1–8 (2014).

50. Schneider, F. D., Brose, U., Rall, B. C. & Guill, C. Animal diversity and ecosystem functioning in dynamic food webs. *Nature Communications* **7**, 1–8 (2016).

51. Yodzis, P. & Innes, S. Body Size and Consumer-Resource Dynamics. *The American Naturalist* **139**, 1151–1175 (1992).

52. Lang, B., Ehnes, R. B., Brose, U. & Rall, B. C. Temperature and consumer type dependencies of energy flows in natural communities. *Oikos* **126**, 1717–1725 (2017).

53. Hirt, M. R., Lauermann, T., Brose, U., Noldus, L. P. J. J. & Dell, A. I. The little things that run: a general scaling of invertebrate exploratory speed with body mass. *Ecology* **98**, 2751–2757 (2017).

54. Delmas, E., Brose, U., Gravel, D., Stouffer, D. B. & Poisot, T. Simulations of biomass dynamics in community food webs. *Methods in Ecology and Evolution* **8**, 881–886 (2017).

55. Gauzens, B., Brose, U., Delmas, E. & Berti, E. ATNr: Allometric Trophic Network models in R. *Methods in Ecology and Evolution* **14**, 2766–2773 (2023).

685   56. Gauzens, B., Berti, E., Delmas, E. & Brose, U. ATNr : Allometric trophic models in R

686   Unscaled version. *bioRxiv preprint* **2**, 1–12 (2022).

57. *g*auzens. gauzens/Allometric_preferences: v1.0.1. Zenodo
    https://doi.org/10.5281/zenodo.10554929 (2024).