Metamorphosis imposes variable constraints on genome expansion

Article type: Major Article

Keywords: genome size, constraint, development, comparative methods, life history

Word Count: 7,729

Figures: 2

Tables: 4

Abstract

Genome size varies ~ 100,000-fold across eukaryotes. Genome size is heavily shaped

by transposable element accumulation, the dynamics of which are increasingly well un-

derstood. However, given that traits like cell size and rate of development co-vary

strongly with genome size, organism-level trait evolution likely shapes genome size di-

versity as well. Metamorphosis — a radical transformation of morphology — has been

hypothesized to impact genome size because it can be a vulnerable part of the life cy-

cle. Thus, selection may act to limit metamorphic duration, indirectly constraining the

rate of development as well as genome and cell sizes. Salamanders have large and

variable genomes — 3 to 40 times that of humans — and species exhibit a range of

metamorphic and non-metamorphic life histories. Using salamanders, we test the hy-

pothesis that different types of metamorphic repatterning during the life cycle impose

different constraints on genome expansion. We show that metamorphosis during which

animals are unable to feed imposes the most severe constraint against genome expan-

sion. Other types of metamorphosis that differ in energetic provisioning impose less se-

vere constraints. More generally, our work demonstrates the utility of phylogenetic com-

parative methods in testing the role of constraint in shaping phenotypic evolution.

Introduction

Across the tree of life, few characters exhibit the tremendous scale of variation of genome size, encompassing a ~100,000-fold range across eukaryotes (Gregory 2021). Decades of research have revealed the consistent covariation of two organismal features with genome size: a negative correlation with cell division rate, and a positive correlation with cell size (Gregory 2001, 2005). Genome size has also been associated with a variety of organismal or ecological factors including: developmental rate or complexity (Gregory 2002b), temperature (Hessen, et al. 2013), metabolic rate (Waltari and Edwards 2002; Roddy, et al. 2019), invasiveness (Pandit, et al. 2014), or speciation and extinction rates (Vinogradov 2004; Jeffery, et al. 2016), but these associations vary across studies.

Comparative biologists often think of adaptive explanations for character associations, which would suggest that genomes evolve toward an "optimum" size with respect to one or more of these correlated traits. However, the context-sensitive nature of associations with these factors belies a strongly adaptive explanation for genome size. For example, metabolic rate and genome size are correlated in some vertebrate clades, but not in others, and genome size provides no overall explanatory power for basal metabolic rate across vertebrates (Licht and Lowcock 1991; Gregory 2002a; Smith, et al. 2013; Uyeda, et al. 2017; Gardner, et al. 2020). Genome size evolution may instead evolve nearly neutrally until some threshold value is reached, beyond which fitness is

impacted (Gregory 2002b). This process is more aptly described as governed by con-

straints. As far as we are aware, the constraint model has never been formally tested

within a phylogenetic comparative framework for a univariate trait.

Recent years have seen tremendous improvement in our understanding of the

mechanistic processes by which genome size evolves. Variation in genome size can re-

flect the accumulation of many types of sequences, from simple repeats to increases in

ploidy (Elliott and Gregory 2015; Pasquesi, et al. 2018; Carta, et al. 2020), such that the

majority of the genome is non-coding or "junk" DNA. In vertebrate animals, genome size

is strongly determined by the accumulation of transposable elements (TEs), sequences

that replicate and spread throughout host genomes (Sotero-Caio, et al. 2017; Shao, et

al. 2019). TEs are also deleted by mutations introduced during replication, recombina-

tion, and DNA repair (Michael 2014; Vu, et al. 2017). In the absence of selection or con-

straint on genome size, the background process for genome evolution is stochastic, with

genome size increasing if TE insertions outpace deletions. TE activity is often neutral at

the cellular and organismal levels, with most individual insertions and deletions missing

functional regions of the genome and resulting in negligible fitness consequences

(Arkhipova 2018). Non-coding DNA can therefore accumulate until genome size

crosses a threshold where it begins to impact fitness through a correlated trait (e.g. cell

size or developmental rate).

While the notion of constraint on genome size is conceptually appealing, there

are few comparative methods that can detect constraint and possibly distinguish it from

adaptation. Whether by correlation of phylogenetically independent contrasts

(Felsenstein 1985) or by phylogenetic autocorrelation (Cheverud, et al. 1985), phyloge-

netic comparative methods that focus primarily on the mean or "location" of the pheno-

type are primed to detect adaptive evolution, as selection is expected to move the phe-

notype toward an optimum (Simpson 1953; Lande 1980). For example, testing for a cor-

relation between traits after independent contrasts (Felsenstein 1985) may reveal

whether, on average, a phenotype covaries across species with some other trait, with

the relationship maintained presumably as a result of adaptive evolution. Phylogenetic

autocorrelation (Cheverud, et al. 1985) and phylogenetic regression (Grafen and

Hamilton 1989) use regression approaches to separate phylogenetic and environmental

(adaptive) effects on variation of the mean phenotype. Ancestral character state recon-

struction methods can infer the mean phenotype of interest along nodes of the tree and

are used to explore evolution in relation to some other factor, but vary in the underlying

models employed (Huey and Bennett 1987; Maddison 1991; Schluter, et al. 1997). How

mean-focused methods enable the detection of constraint, however, is unclear. Meth-

ods that model both the mean and variance of a stochastically evolving phenotype, such

as Ornstein-Uhlenbeck-based methods (Hansen 1997; Butler and King 2004; O'Meara,

et al. 2006; Beaulieu, et al. 2012), may open the door to exploring constraint by reveal-

ing whether a stochastically spreading phenotype evolves by selection, or by weak se-

lection bounding the phenotype consistent with the presence of constraint, or by no se-

lection at all.

Salamanders are an ideal group to test hypotheses of selection and constraint on

genome size. They have both the largest genomes and the largest range of variation in

genome size among vertebrates. Genome sizes range from 9 Gb – 120 Gb across the

763 extant species (Decena-Segarra, et al. 2020; AmphibiaWeb 2021; Gregory 2021),

reflecting varying levels of TE accumulation. Salamanders also have exceptional life-

history diversity, in particular, with metamorphosis lost, modified, and regained through-

out their evolutionary history (Wiens, et al. 2005; Bonett, et al. 2014). Metamorphosis

has been hypothesized to shape genome size evolution because of the effects that ge-

nome (and therefore cell) size has on the rate of development (Wake and Marks 1993;

Gregory 2002b). Because metamorphosis can be a vulnerable part of the life cycle, its

duration has been proposed as a target of natural selection. Selection to limit the time in

metamorphosis would indirectly select for faster development, constraining genomes

(and cells) to smaller sizes (Gregory 2002b; Bonett, et al. 2020).

Several studies have linked life history to genome size in salamanders, with

smaller (albeit still enormous relative to other taxa) genome sizes associated with meta-

morphosis (Wake and Marks 1993; Gregory 2002b; Sessions 2008; Bonett, et al. 2020).

Bonnett (2020) used phylogenetic comparative methods to demonstrate that salaman-

der genome size is better explained by models that account for differences in life history

— including the presence or absence of metamorphosis — as opposed to variation in

ecology or habitat stability. However, this study was largely focused on the effects of

facultative loss of metamorphosis on genome size evolution; thus, it did not explore the

concept of evolutionary constraint versus adaptation. In addition, it did not examine the

connection between energetic vulnerability and metamorphic duration, which we pro-

pose as a key conceptual link between metamorphosis and genome size.

In this study, we build on this previous body of work, revisiting the hypothesis that

the radical morphological repatterning associated with metamorphosis imposes evolu-

tionary constraints on genome size in salamanders. In particular, we model characteris-

tics that are likely to place energetic limits on the duration of metamorphosis: Does re-

patterning happen inside the egg, fueled solely by yolk stores? Does repatterning hap-

pen in an organism that is free-living, but unable to feed? We use life history and ge-

nome size data to inform stochastic models of trait evolution that explore how different

metamorphic regimes interact with genome expansion. More generally, we demonstrate

the ability of OU-based stochastic models to identify trait evolution governed by con-

straint, expanding their use beyond classic scenarios of adaptive evolution.

Life History Regimes Constraining Genome Size Evolution

In the absence of any selection or constraint, genome size in salamanders is expected

to expand by biased stochastic evolution. Overall mutation pressure in the clade leads

to TE accumulation, as TE deletion rates are low and TE silencing is incomplete (Sun,

Arriaza, et al. 2012; Frahry, et al. 2015; Madison-Villar, et al. 2016; Mohlhenrich and

Mueller 2016). Salamanders' enormous genomes are the cumulative result of unusually

high levels of TE activity and retention (Sun, Shepard, et al. 2012; Sun and Mueller

2014; Keinath, et al. 2015; Liedtke, et al. 2018; Nowoshilow, et al. 2018).

The ancestral salamander life history included a larval growth stage followed by

metamorphosis. Metamorphosis is a radical transformation of morphology during the

life cycle that produces strikingly different, largely decoupled larval and adult forms

(Moran 1994). During metamorphosis, as during embryogenesis, salamanders undergo

rapid cell division, differentiation, migration, and apoptosis (Alberch 1989). The dynam-

ics of these cellular processes are affected by genome size and cell size; as ge-

nome/cell sizes increase, developmental rates throughout ontogeny — from embryo-

genesis through metamorphosis — slow down (Horner and Macgregor 1983; Jockusch

1997). Metamorphosis has been lost and regained from the life cycle numerous times

by different mechanisms (Chippindale, et al. 2004; Mueller, et al. 2004; Wiens, et al.

2005; Bonett, et al. 2014), producing multiple life history regimes that could shape ge-

nome size evolution. We outline these regimes, and their predicted effects on genome

size evolution, below and in Figure 1:

Paedomorphosis: In paedomorphic species, some or all stages of metamorphic repat-

terning are lost, and organisms reach sexual maturity retaining largely larval traits

(Gould 1977). Because there is no selection to limit time in metamorphosis, there is no

associated constraint on genome size. Thus, paedomorphs are expected to most

closely reflect the background condition of stochastic genome expansion.

Direct development: In direct-developing species, the larval growth stage is eliminated,

and embryogenesis and metamorphosis are integrated into a single sequence of devel-

opmental events that takes place inside the egg (Alberch 1989; Rose 2014). This entire

sequence must be fueled by yolk provisioned by the mother. Thus, we might expect se-

lection to limit time in metamorphosis, imposing a constraint on genome size.

Non-feeding metamorphosis: Within the salamander family Plethodontidae, metamor-

phic repatterning events happen relatively synchronously in a free-living organism. Dur-

ing this metamorphosis, the organisms are unable to feed because of the replacement

of cartilaginous elements associated with the change from suction to projectile feeding

(Rose 1995a; Deban and Marks 2002). The entire process must be fueled by energy re-

serves built up during the larval stage. Thus, we might expect selection to limit time in

metamorphosis, imposing a constraint on genome size.

Feeding metamorphosis: In non-plethodontid salamanders, metamorphic repatterning

events happen in a free-living organism that is able to feed throughout the transfor-

mation. Thus, we would not expect selection based on energy demands to limit time

spent in metamorphosis. There are, however, other ways in which metamorphosis can

increase vulnerability that might translate into selection on metamorphic duration, con-

straining genome size.

173
174 Figure 1. Alternative hypotheses for constraints imposed by development on genome
175 size evolution in salamanders. On each phylogeny, alternative life history regimes are
176 painted in different colors as indicated in each legend (see text). Genome sizes are
177 shown on the right in pg (1 pg = 978 Mb).
178

Methods

Taxon Sampling, Genome Size, and Phylogeny

Genome size data were available for 106 species of salamanders (out of a total of 763 currently named extant species), encompassing all ten salamander families and 35 of 68 genera (Supplemental Table 1) (AmphibiaWeb 2021; Gregory 2021). We transformed the data with natural logarithms prior to analysis. Our sampling includes representatives with diverse life histories: direct development, paedomorphosis, feeding metamorphosis, and non-feeding metamorphosis. Hereafter, we distinguish between "feeding" and "non-feeding metamorphosis", and use "metamorphosis" to indicate all species which undergo metamorphosis, whether feeding or non-feeding. Several lineages are facultative paedomorphs, which retain the ability to undergo metamorphosis under certain conditions. We coded these taxa as metamorphic, reflecting the experience of constraint, even if only occasionally, and supported by recent work examining the evolutionary impacts of facultative paedomorphosis (Bonett, et al. 2020).

We used a previously published phylogeny (Mueller, et al. 2008; Pyron and Wiens 2011; Vieites, et al. 2011; Zheng, et al. 2011) and estimated branch lengths on this topology using sequence data for two mitochondrial genes – cytochrome-b and 16S obtained from GenBank. Additional Sanger sequences were collected as necessary at the Colorado State University sequencing core facility or the Joint Genome Institute following (Mueller, et al. 2004) (Supplemental Table 1). Loci were aligned using MUSCLE

(Edgar 2004). Branch lengths were estimated for each locus independently and then av-

eraged, weighted by gene length. Branch lengths were estimated using RAxML

(Stamatakis 2006) using the best-fit (AIC) model of nucleotide substitution in ModelTest

3.6 (Posada and Crandall 1998), with Cyt-b partitioned by codon. The resulting tree was

ultrametricized using penalized likelihood implemented in r8s with the truncated Newton

algorithm and cross validation to select the optimal smoothing parameter value

(Sanderson 2003).

Models of Genome Size Evolution

We modeled genome size evolution using both Brownian motion (BM) and Ornstein-Uh-

lenbeck (OU) models of evolution (Hansen 1997; Butler and King 2004; O'Meara, et al.

2006; Beaulieu, et al. 2012). As these models have been described previously, we

cover them only briefly here. The BM model is the simplest stochastic model used in

comparative analysis and has a single rate parameter σ, which can be thought of as a

stochastic noise intensity parameter shared by all taxa and describing the magnitude of

the independent random walks of the trait evolving along the branches of the phylogeny.

This model predicts that across lineages, there will be no change in mean phenotype,

but variance will grow through time as lineages wander over trait space, and the ex-

pected variance between two lineages will be proportional to the time since their diver-

gence. Conceptually, the BM model may be a good candidate for purely stochastic ge-

nome size evolution if increases occur as frequently as decreases.

The multiple-rate BM model, introduced by O'Meara, et al. (2006), allows differ-

ent σ values across different portions of a phylogenetic tree. We use this model to rep-

resent changes in the rate of stochastic evolution accompanying changes in constraint

resulting from shifts in life history regime. Under this model, lineages evolving under dif-

ferent regimes may differ in variance.

OU models generalize the BM model by including terms that allow the mean to

shift as well as allowing variance to narrow. They include a deterministic component of

trait evolution that models the tendency of the trait to move toward an equilibrium value.

Mathematically, the model for trait evolution expressed as a differential equation is

$$dX(t) = \alpha(\theta(t) - X(t)) + \sigma dB(t),$$

where $\theta(t)$ is the deterministic equilibrium for the trait at time *t* and α is an evolutionary

rate describing the *strength of the deterministic pull (e.g. selection)* towards that equilib-

rium along a branch of a tree. Hypotheses regarding trait evolution are specified by

painting "regimes" on the tree to indicate where these parameters are expected to shift.

The simplest OU models allow multiple equilibria leading to the evolution of differences

in mean phenotype across regimes (Hansen 1997; Butler and King 2004); for example,

in the current study, separate equilibria might exist for paedomorphs, metamorphosers,

and direct developers. Further model extensions also allow the strength of the determin-

istic pull and the stochastic noise intensity to vary across regimes (Beaulieu, et al.

2012), as might be consistent with the lifting or imposition of constraints on genome size

evolution.

We formalized five hypotheses for how life history regime could constrain ge-

nome size evolution in salamanders: (1) Brownian motion: Genome size evolves by

purely stochastic evolutionary processes with neither constraint nor bias. (2) *metamor-*

phosis-other: Metamorphosis imposes a constraint on genome expansion that is dis-

tinct from the other life histories. (3) *meta-paed-dd*: Metamorphosis, direct develop-

ment, and paedomorphosis each impose distinct constraints on genome expansion. (4)

meta$_f$-meta$_{nf}$-other: Feeding metamorphosis (*meta$_f$*) and non-feeding metamorphosis

(*meta$_{nf}$*) each impose constraints that are distinct from those experienced by salaman-

der species with other life histories. (5) *meta$_f$-meta$_{nf}$-paed-dd*: non-feeding metamor-

phic species, feeding metamorphic species, paedomorphic species, and direct-devel-

oping species each experience different levels of constraint on genome expansion.

These five biologically-inspired hypotheses are specified by fitting evolutionary

models that allow their parameters to vary with life history regime. The simplest form al-

lows the equilibrium to vary with life history regime while assuming the rates of evolution

are constant across the phylogeny. Additional subhypotheses fit evolutionary models

that vary stochastic noise intensity values (σ), and/or deterministic pull values (α) ac-

cording to life history regime (Table 1). In all, we tested 21 models to explore life history

constraints on genome size and model parameters (Table 2). We fit all of the models

with the character state of the node at the base of the plethodontid lineage defined as

metamorphosing as well as direct-developing (Bonett, et al. 2014), and found that the

choice of the basal character state made no qualitative difference to our conclusions. In

addition, we made a series of pairwise comparisons to further assess the degree of in-

fluence of feeding metamorphosis, paedomorphosis, direct development, and non-feed-

ing metamorphosis on genome size evolution in combinations of increasing complexity

(Table 3).

We fit models of evolution using the software package OUwie (Beaulieu, et al.

2012) and compared the multiple optimum OU models with results fitted in OUCH

(Butler and King 2004; King and Butler 2009). We modified OUwie to fit models with re-

gime changes that occur at the nodes, as in OUCH, rather than the default behavior of

OUwie which places regime changes mid-way along the branch between a clade of in-

terest and its ancestor. We recognize that while either choice can be argued to be arbi-

trary, assuming that the regime change occurred simultaneously with the origin of the

clade that bears the phenotype as indicated in Figure 1 seems reasonable. All analyses

were conducted in the R statistical computing environment (R Core Team 2020). We

note here that OUwie has different options for dealing with the root state, $X(0)$. Absent

any information about the phenotype deep in the tree, this parameter is often very diffi-

cult to estimate in OU models (Cressler, et al. 2015). One alternative is to assume that

the value of $X(0)$ is distributed according to the stationary distribution of an OU process,

which eliminates this parameter by absorbing the variance into the covariance matrix

implied by the phylogeny itself (Ho and Ané 2013). However, OUwie does not currently

support this approach for OU models with multiple α or σ parameters, and as far as we

know, the mathematical modifications for these models have not been worked out. Be-

cause the option to estimate the value of the root node as in the earlier implementations

(Hansen 1997; Butler and King 2004); and specified in OUwie by setting root.sta-

tion=FALSE) is available for all of the models of interest, we used it for all model fitting.

However, as we show in the Supplementary Materials, the parameter estimates of a sin-

gle-α, single-σ OU model fit assuming the root is in the stationary distribution are very

different from the parameter estimates obtained by fitting the root state, although model

selection conclusions hold. This large difference in the parameter values reinforces a

general point in phylogenetic comparative hypothesis testing, which is that parameter

estimation is often more fraught than hypothesis testing, and as such, parameter esti-

mates should be interpreted with caution (Beaulieu, et al. 2012; Ho and Ané 2013; Ho

and Ané 2014; Cressler, et al. 2015; Cooper, et al. 2016).

When fitting OU models, there are datasets for which it is possible to recover an

extremely low value of α (weak deterministic pull), with equally extremely large θ (equi-

librium values). This combination can reasonably be interpreted as weak selection, or a

biased nearly-random walk. Rather than look only at these parameters in isolation, we

considered whether it may be more informative to determine their combined effect on

trait evolution. To this end, we calculated the value of $\alpha(X - \theta)$ for each life history re-

gime to estimate any "deterministic trend" imposed by the best-fitting model, where X is

the average genome size of species in that regime at the end of the evolutionary pro-

cess.

Table 1. BM and OU models with single or multiple parameters used to fit the data. Numbers in parentheses specify (1) model parameters and notation, (2) parameters that remain constant across the phylogeny, (3) parameters that vary with shifts in life history regime, (4) O'Meara (2006) model notation, and (5) notes for the model implementations and citations.

Models	Uniform	Variable with Regime	O'Meara Notation	Notes
(1)	(2)	(3)	(4)	(5)
Single noise intensity σ	σ		BM1	Classic BM model of (Felsenstein 1985)
Multiple noise intensities σ_i		σ_i	BMS	Multiple-rate BM model of O'Meara et al. (2006)
Multiple equilibria $\theta i, \sigma, \alpha$	σ α	θ_i	OUM	OU model of Butler and King (2004)
Multiple equilibria and deterministic pull strengths $\theta i, \sigma, \alpha i$	σ	α_i θ_I	OUMA	Multiple-α model of Beaulieu et al. (2012).
Multiple equilibria and noise intensities $\theta i, \sigma i, \alpha$	α	σ_i θ_i	OUMV	Multiple-σ model of Beaulieu et al. (2012)

Multiple equilibria, noise intensities, and deterministic pull strengths $\theta i, \sigma i, \alpha i$		σ_i α_i θ_i	OUMVA	Full model of Beaulieu et al. (2012).

317
318

Model Comparison and Parameter Estimation

We compared the fit of each of the models using the Akaike Information Criterion cor-

rected for small sample size (AIC$_c$). We further performed model selection bootstrap

analysis (phylogenetic Monte Carlo; (Boettiger, et al. 2012) for targeted comparisons of

the models in Table 3 to assess the strength of evidence for different hypotheses as

well as the power to detect differences in model support. Model selection bootstrap

analysis is necessary because AIC$_c$ differences can favor more complicated models,

even when a simpler model is correct (Boettiger, et al. 2012). For each comparison, we

computed the observed likelihood difference,

$$\delta_{obs} = -2 (\log L_0 - \log L_1),$$

where L_0 is the likelihood of the simpler model and L_1 is the likelihood of the more com-

plex model.

Determining whether δ_{obs} is significantly different from a null expectation requires

an approximate *p*-value — the probability of observing δ_{obs} *if the simpler model were*

true. That is, we need to compare the value δ_{obs} to the distribution of δ values under the

simpler model. To create this distribution, we generated 500 datasets by simulating the

simpler model at its MLE parameter estimates; we then fit both the simpler and more

complex models to each simulated dataset and computed the values of δ, producing a

null distribution of δ under the simpler model. We compared the observed value of δ to

this null distribution to calculate an approximate *p*-value.

Power conveys the (desirable) probability of rejecting the simpler model when the

more complex model is true. To estimate power, we generated 500 datasets by simulat-

ing the more complex model at its MLE parameter estimates; we then fit the two models

and computed the values of δ. The fraction of these δ values that are greater than the

95% quantile of the distribution generated under the simpler model (described above)

gives an estimate of power. All data and code necessary to carry out the analysis in this

manuscript can be found at https://github.com/claycressler/genomesize and in Supple-

mental Material.

Results

The best-fitting model for salamander genome size evolution accounted for four

regimes: both non-feeding and feeding metamorphosis, paedomorphosis, and direct de-

velopment (*meta$_f$-meta$_{nf}$-paed-dd*; Table 2) under an OU model that allowed both equi-

librium genome size and noise intensity to vary across these regimes (θ_i, σ_i, α, Table 2).

An identical 4-regime model with only a single noise intensity fit nearly as well (Table 2).

Additionally, the three-regime *meta$_f$-meta$_{nf}$-other* (θ_i, σ_i, α) hypothesis provided some

explanatory power. These three models were far superior to the remaining models.

Overall, the addition of multiple α values resulted in worse model fit relative to a

uniform α value, whereas the addition of multiple σ values relative to a single σ both im-

proved and worsened model fit, depending on the hypothesis (Table 2). We present the

results with the character state of the node at the base of the plethodontid lineage de-

fined as metamorphosing. In the Supplementary Materials, we show that defining this

node as direct-developing has no effect on the evolutionary conclusions we draw here.

The model selection bootstrap analysis allows us to quantify the degree of im-

provement in explanatory power provided by moving between specific hypotheses (Fig-

ure 2). Note that for each comparison, we used the version of the OU model that had

the lowest AICc (Table 2), so comparisons involving *metamorphosis-other* and *meta-*

paed-dd used the single-σ model, whereas comparisons involving *meta$_f$-meta$_{nf}$-paed-*

dd and *meta$_f$-meta$_{nf}$-other* used the multiple-σ model.

Based on these results, we can reject any purely stochastic hypothesis for ge-

nome size evolution, as a model that allows for separate equilibrium values for meta-

morphosers was far superior to any purely neutral model (Table 2; Figure 2A). Specify-

ing distinct equilibrium values for non-feeding and feeding metamorphosis substantially

improves the explanatory power of any model (Figure 2C, D). In particular, the separa-

tion of metamorphosis into feeding and non-feeding categories provided far greater im-

provement of the model than subdividing the "other" category into direct development

and paedomorphosis (compare the results in Figure 2C to those in Figure 2B, and Fig-

ure 2D to 2E). Allowing distinct noise parameters for each regime slightly improves the

fit of the OU models with separate equilibria for non-feeding and feeding metamorphosis

(Table 2), although the improvement over the single-σ model is not significant (Compari-

son [6], Table 3; Figure 2F). Therefore, the best model included distinct equilibrium val-

ues for direct development, paedomorphosis, feeding metamorphosis, and non-feeding

metamorphosis.

Table 2. Model comparison statistics. Best models (interrogated by bootstrap, Table 3) indicated in bold. Model parameterizations are indicated by: σ = Brownian motion; σ_i = Brownian motion with multiple noise intensities; θ_i, σ, α = OU model with multiple equilibria; $\theta_i, \sigma_i, \alpha$ = OU model with multiple equilibria and multiple noise intensities; $\theta_i, \sigma, \alpha_i$ = OU model with multiple equilibria and multiple deterministic pull strengths; $\theta_i, \sigma_i, \alpha_i$ = OU model with multiple equilibria, noise intensities, and deterministic pull strengths.

ΔAICc	Model					
	OU Models				BM Models	
Hypotheses	$\theta i, \sigma i, \alpha$	$\theta i, \sigma, \alpha$	$\theta i, \sigma i, \alpha i$	$\theta i, \sigma, \alpha i$	σi	σ
metaf -metanf -paed-dd	**0**	**1.1**[a]	7.5	8.1	12.9	
metaf -metanf -other	1.4[b]	3.6	6.1	8.1	11.1	
meta-paed-dd	6.0	3.4	10.7	7.9	11.4	
metamorphosis-other	5.8	4.1	8.0	6.3	9.5	
Brownian motion						31.8

[a] *metaf -metanf -dd-paed* with single or multiple noise parameters σ are compared in Figure 2F.

[b] *metaf -metanf -other* is compared against the best model via bootstrap in Figure 2E.

Figure 2. Bootstrap distributions of the likelihood difference (δ) calculated by generating 500 datasets under each of two competing models at their MLE parameter estimates, refitting the two models, and computing δ. The light gray region shows the probability density of the parameter when the data is generated by the simpler model; the dark gray region shows the density when the data is generated by the more complex model. The dashed line gives the observed value (δ_{obs}) from fitting the actual genome size data. The reported p-value is the fraction of the light gray distribution that lies to the right of δ_{obs}; the power is the fraction of the dark gray distribution that lies to the right of the 95th percentile of the light gray distribution. Each panel evaluates the support for a different hypothesis: (A) metamorphosis imposes a constraint on genome expansion; (B) there are distinct constraints imposed by the different non-metamorphosing strategies, direct development and paedomorphosis; (C) non-feeding metamorphosis imposes a distinct constraint from feeding metamorphosis; (D) non-feeding metamorphosis imposes a distinct constraint from feeding metamorphosis, after accounting for differences in non-metamorphosing strategies; (E) non-metamorphosing strategies impose unique constraints, after accounting for differences between feeding and non-feeding metamorphosis; (F) after identifying *meta$_f$-meta$_{nf}$-paed-dd* as best-fitting, this comparison tested whether fitting each regime with a distinct stochastic noise intensity improved fit to the data.

Parameter values for each regime estimated under the best-fitting model are presented

in Table 3. Direct-developers, non-feeding metamorphosers, paedomorphs, and feeding

metamorphosers have broadly overlapping stochastic noise intensity (σ) values. Deter-

ministic pull strength (α) takes an extremely small value, and the equilibrium values (θ)

take extreme values both large and small. Although the deterministic pull strength pa-

rameter is vanishingly small, nevertheless, the models that account for metamorphosis

are far superior to a purely stochastic model (Table 2). We note that the extreme nature

of the parameter estimates is a consequence of the handling of the root state (see Sup-

plementary Materials).

430
431

432 Table 3: Maximum likelihood parameter estimates and parametric bootstrap confidence
433 intervals for the best-fitting model (*meta_f* -*meta_nf* -*dd-paed* $\theta i, \sigma i, \alpha$: separate equilibrium
434 values and noise intensities for lineages in the four life history regimes: dd = direct de-
435 velopment, meta$_{nf}$ = non-feeding metamorphosis, meta$_f$ = feeding metamorphosis, paed
436 = paedomorphosis).
437

Parameter		MLE	95% CI
Deterministic pull strength	α	1.05e-7	(1.01e-7, 1.7)
Stochastic noise intensity	σ_{dd}	0.365	(0.308, 0.545)
	σ_{meta_nf}	0.355	(0.252, 0.525)
	σ_{meta_f}	0.187	(0.108, 0.283)
	σ_{paed}	0.249	(0.112, 0.464)
Equilibrium value	θ_{dd}	-3.77e5	(-2.67e6, 1.99e6)
	θ_{meta_nf}	3.62	(3.32, 3.90)
	θ_{meta_f}	-5.55e6	(-7.62e6, 3.28)
	θ_{paed}	4.41e6	(3.96, 6.29e6)

438
439

There is evidence of a deterministic trend $\alpha(X - \theta)$ towards small genome size in sal-

amanders that undergo non-feeding metamorphosis as well as a trend towards large

genome size in paedomorphic salamanders (Table 4). In contrast, genome sizes in di-

rect developing and feeding metamorphosing salamanders show no such trends. The

range of values for the strength of the overall deterministic trend (Table 4) is much

greater than the range of values for stochastic noise intensity (Table 3).

Table 4: Estimates of the deterministic trend in the best-fitting model for lineages evolv-
ing in each life history regime, based on the average genome size (log-scale) of sala-
manders in each regime. Genome sizes (in pg) are also shown in parentheses.
450

Regime	Average log genome size X	Deterministic trend $\alpha(\theta - X)$
Direct development	3.66 (38.9 pg)	-0.0396
Feeding metamorphosis	3.46 (31.8 pg)	1.68e-8
Non-feeding metamorphosis	3.07 (21.5 pg)	-0.583
Paedomorphosis	3.89 (48.9 pg)	0.463

451
452

Discussion

Biased stochastic expansion and life history constraints shape genome size.—
Overall, evolution of genome size in salamanders is well-described by a weakly deter-
ministic model with separate equilibria for each life history regime and moderate sto-
chasticity. An increased level of constraint in lineages that evolved non-feeding meta-
morphosis, as well as a decreased level of constraint in lineages that have lost meta-
morphosis, exert the greatest deterministic influences on the evolution of genome size.
The strength of the deterministic trend towards genome expansion for paedomorphs is
roughly the same as the strength of the deterministic trend towards genome reduction
for non-feeding metamorphosers; both are an order of magnitude stronger than the
trend for direct-developers. Feeding metamorphosers have no deterministic trend (Ta-
ble 4). Taken together, these results, and the observed spread of genome sizes within
regimes, suggest that genome size evolution is driven by stochasticity with a bias to-
ward increase — likely representing TE insertions that outpace deletions — constrained
to varying degrees by metamorphic repatterning, when selection acts to limit metamor-
phic duration.

Metamorphosis as a vulnerable stage of the life cycle in amphibians
Metamorphosis has been posited to exert evolutionary pressure for reduced genome
size in amphibians to speed development during a vulnerable life stage. This hypothesis
comes primarily from evidence in frogs. Metamorphosing frogs experience higher pre-
dation levels because they can neither swim nor hop effectively (Wassersug and Sperry

1977; Arnold and Wassersug 1978). In addition, frogs are unable to feed during meta-

morphosis, at a time when their energetic requirements can nearly double as compared

to the period immediately preceding metamorphosis (Orlofske and Hopkins 2009;

Wright, et al. 2011). Despite being homologous and retaining broad similarities at the

transcriptomic, hormonal, and organismal levels (Sanchez, et al. 2018), salamander

metamorphosis is different — and less dramatic — than frog metamorphosis. The pro-

cess takes much longer in salamanders; timescales are on the order of weeks to

months rather than days (Norman 1985; Downie, et al. 2004; Vladimirova, et al. 2012;

Sanchez, et al. 2018), suggesting little, if any, time pressure. Metamorphosing salaman-

ders do not experience compromised locomotion and are thus not more vulnerable to

predation as are frogs (Landberg and Azizi 2010). In addition, metamorphosing sala-

manders do not have higher energetic requirements compared to non-metamorphosing

individuals of the same species (Vladimirova, et al. 2012). Some salamanders, how-

ever, are unable to feed during metamorphosis, requiring that they undergo the transfor-

mation using only stored energy reserves (i.e., non-feeding metamorphosers) (Deban

and Marks 2002)). Direct developers undergo the transformation inside the egg, fueled

only by yolk stores. Genome size constraints for these two life history regimes likely re-

flect energetic vulnerability that is not relevant for the other two life histories considered

here: feeding metamorphosis and paedomorphosis.

Genome size evolution in feeding metamorphosers.—Lineages that undergo metamor-

phosis, but are able to feed throughout the process, show no deterministic trend in ge-

nome size evolution. Rather, trait evolution is described by moderate stochastic noise

around an equilibrium value that we interpret as a balance between TE accumulation

and a constraint imposed by metamorphosis. Although the animals are able to feed,

there are other ways in which fitness can be lowered during salamander metamorpho-

sis; for example, metamorphosing individuals are less able to exploit stream habitat re-

fugia than either larvae or adults, which increases their mortality (Lowe, et al. 2019).

Our results indicate that feeding metamorphosis imposes a less severe constraint on

genome size than non-feeding metamorphosis, and we infer that the constraint is medi-

ated by vulnerabilities other than depletion of energetic stores.

Genome size evolution in non-feeding metamorphosers.— Although other analyses

have demonstrated a link between metamorphosis and genome size in salamanders

(Wake and Marks 1993; Gregory 2002b; Sessions 2008; Bonett, et al. 2020), here we

show that non-feeding metamorphosis imposes a substantial and distinct constraint, rel-

ative to feeding metamorphosis, as predicted if energetic vulnerability shapes the dura-

tion of metamorphosis. The deterministic trend toward smaller genome sizes within this

regime is consistent with the imposition of a more severe constraint against genome ex-

pansion — or, put another way, selection towards genome size reduction —— to

shorten the duration of metamorphosis.

The evolution of non-feeding metamorphosis in plethodontids has been an im-

portant target of research because the phylogeny suggests that it evolved from a direct-

developing ancestor(s), which necessitates the evolutionary reappearance of the "lost"

larval stage (Chippindale, et al. 2004; Mueller, et al. 2004). This evolutionary transfor-

mation series was historically considered unlikely, although more recent work has re-

vealed potential scenarios for regain of the larval stage (Bonett, et al. 2005; Bonett, et

al. 2014). Under this scenario, in the direct-developing ancestral lineage(s), metamor-

phic repatterning steps were retained as part of the longer sequence of developmental

events that occurred inside the egg (Kerney, et al. 2012). These changes were likely

mediated by evolutionary changes in the timing of thyroid hormone activity and re-

sponse (Rose 1995a, b; Bonett 2016). The re-evolution of (now non-feeding) metamor-

phosis reflected the insertion of a long, slow-growing larval growth phase back into on-

togeny, followed by the synchronous occurrence of more drastic metamorphic repattern-

ing events in the free-living organism (Rose 1995b, c; Beachy, et al. 2017). Metamor-

phic repatterning is more extreme in plethodontids than in other salamanders. Im-

portantly, it involves a complete remodeling of the feeding apparatus, with the cerato-

branchials (cartilaginous components of the tongue skeleton) replaced by new struc-

tures in the adult rather than remodeled from existing larval structures; this results in the

inability to feed during metamorphosis (Alberch 1989; Rose 1995c; Deban and Marks

2002). Under the classical scenario, in contrast, this synchronization of metamorphosis

and drastic remodeling of feeding structures would have evolved in a metamorphosing

ancestor to produce non-feeding metamorphosis (Wake and Hanken 2004). Under ei-

ther scenario, our results illustrate how phylogenetic comparative methods can reveal

the evolutionary forces that have acted on genome size as lineages moved through the

different life history regimes.

Genome size evolution in direct developers.—In direct-developing lineages, some or all

of the developmental steps of metamorphic repatterning occur inside the egg at the end

of embryogenesis (Alberch 1989; Kerney, et al. 2012); these lineages show a weak de-

terministic trend towards genome size reduction. Because they are occurring in an em-

bryo rather than a free-living organism that has undergone a growth period, the repat-

terning happens to a smaller number of cells in a smaller overall mass of tissue com-

pared with metamorphosing lineages (Downie, et al. 2004). Thus, the energetic require-

ments for comparable developmental steps are lower in direct developers than in meta-

morphosers. On the other hand, the energy to fuel these steps comes from yolk stores

which, although plentiful in direct developers, are still finite (Wake and Hanken 2004).

Thus, we infer that direct development imposes a less severe constraint on genome

size than does non-feeding metamorphosis, mediated by the potential for depletion of

energy stores if the duration of metamorphic repatterning during embryogenesis is too

long. We note that there is greater variation across direct developers in metamorphic re-

patterning than is modeled here. In some cases, the sequence of developmental events

is shortened because the formation of larval structures is lost from ontogeny. In other

cases, most or all events of embryogenesis and metamorphosis are retained but occur

inside the egg (which allows for the possibility of re-evolution of metamorphosis; Alberch

1989). We would predict more severe constraints in these latter lineages. Although we

treated both scenarios as a single category for simplicity, these two types of direct de-

velopment may be different in their effects on genome size evolution and warrant more

detailed study.

Genome size evolution in paedomorphs.—Paedomorphic salamanders retain a larval

body form throughout life and show a deterministic trend towards genome expansion.

This trend is consistent with TE accumulation proceeding to higher overall levels, un-

checked by any constraints imposed by metamorphic repatterning. However, we do not

suggest that genome size is free from all constraints. The impacts of decreased sur-

face-area-to-volume ratio that accompany increased cell size likely impose an upper

limit on cell function that salamanders may well have reached (Chan and Marshall

2010); their cells are among the largest found in animals (Horner and Macgregor 1983).

In addition, the duration of embryogenesis may well have an upper bound that con-

strains genome expansion at the extremely high end. In the past, huge cells have been

proposed as adaptive because they coincide, at broad taxonomic levels, with low meta-

bolic rates; salamanders and lungfishes have the lowest metabolic rates and the largest

genomes/cells within vertebrates. This correlation led to the proposal that selection

shaped an adaptive "frugal metabolic strategy" in these taxa (Szarski 1983; Olmo, et al.

1989). More recent analyses of the relationship between genome/cell size and meta-

bolic rate, however, have failed to find a clear relationship (Licht and Lowcock 1991;

Uyeda, et al. 2017; Gardner, et al. 2020). Thus, empirical evidence that huge genomes

are a product of directional selection is currently lacking. Our results are more con-

sistent with the relaxation of a constraint against genome expansion because of the ex-

tremely weak deterministic pull strength and strong stochastic noise parameters.

Model complexity to capture the evolutionary process.—While the best model includes a

deterministic pull parameter, its magnitude is miniscule. Yet our model selection results

demonstrate that models with deterministic pull provide a huge improvement over any

purely stochastic model (Table 2). One of the challenges of an OU model with a weak

deterministic component is that the model parameters will be poorly defined (Cressler,

et al. 2015). Intuitively, if the pull parameter (α) is important but close to zero, the equi-

libria (θi) can take on a wide range of values in combination with a range of values for

the stochastic parameter σ and explain the phenotypic distribution equally as well. The

difference between a purely stochastic (BM) model and one that has any degree of de-

terministic pull is that the variance of a BM model will grow unbounded over time (as

variance of a trait undergoing a Brownian motion process is proportional to time),

whereas the variance in a model with deterministic pull will not (variance $\sim \sigma^2 / 2\alpha$)

(Hansen and Martins 1996; Butler and King 2004). Thus, while the phenotype may ex-

plore a wide range of values in an OU model with very small α, it will remain bounded.

For paedomorphs, a weak deterministic pull allows the mean to wander, while a far-

away equilibrium value captures a deterministic trend toward increase. In OU models,

increasing deterministic pull strength influences the approach to the equilibrium, but

also will tend to dampen stochastic effects (apart from the influence of σ), so it is possi-

ble to have both weak deterministic pull and substantial noise in the stochastic process.

This analysis demonstrates that deterministic pull can exert an important evolutionary

influence, even if the magnitude of alpha is weak.

But how complex a model is necessary? One might suppose that stronger deter-

ministic pull on one portion of the tree would support a rate shift in α. However, all multi-

ple α models performed poorly. This is consistent with extensive simulation results

showing that, among the three basic parameters of the OU model, α is most poorly de-

fined (Boettiger, et al. 2012; Ho and Ané 2013; Cressler, et al. 2015). Thus, even if a

rate shift in α existed, there is probably little power to detect it. In this study, we find that

with multiple free parameters, we can readily capture shifts in the evolutionary process

with variable θ and perhaps σ over the tree, with stronger deterministic trends accom-

plished by moving θ to more extreme values. A shift in α is superlative, as changes in α

and θ are not independently identifiable. Beyond cases with weak deterministic pull, dif-

ficulty in identifiability of α may be a general problem for all comparative studies, as we

do not know of a case as of yet where a multiple α model was superior.

What this analysis clearly illustrates is the importance of including relevant bio-

logically-informed hypotheses without over-parameterizing the evolutionary model. Iso-

lating the evolutionary "signal" from the "noise" in this dataset was most strongly aided

by including relevant hypotheses for variation in constraint, particularly by specifying

shifts in metamorphosis and non-feeding metamorphosis along the evolutionary history

of salamanders. When looking at adding parameters to the evolutionary model, the ef-

fects were variable with strongly positive (α), neutral (multiple σ), or strongly detri-

mental (multiple α) effects on explanatory power. We note that the harm from overpa-

rameterizing is not only significantly poorer model scores, but at least in this dataset, the

order of the hypotheses changed. It is possible that these significantly worse-fit models

leak variation from some of the regime categories to the extraneous parameters. It is

critical for model testing, therefore, to include a test of all of the meaningful biological

hypotheses with a range of parameterizations including models with fewer parameters.

Exploring evolutionary constraints with the comparative method.—Constraint has long

been argued as a necessary component of the evolutionary toolbox, as workers realized

that selection alone is insufficient to explain macroevolutionary shifts in phenotype

(Alberch 1980; Gould 1980; Cheverud 1984). The concept of constraint is paradoxically

both simple: "evolutionary constraints are restrictions or limitations on the course or out-

come of evolution" and wide-reaching, including genetic, selective, developmental, and

functional constraints (Arnold 1992). Many fields have attempted to quantify constraints

at various levels of biological organization using the notion of limitation. Genetic con-

straints have been identified as limitations on genetic variation (Kirkpatrick and Lofsvold

1992), constrained responses to selection (Cheverud 1984), or multivariate correlations

that are antagonistic to the direction of selection (Etterson and Shaw 2001). All of these

mechanisms allow the phenotype to change in ways that are not directly adaptive. De-

velopmental canalization, decisions in the developmental program which narrow the

phenotypic possibilities in some traits later in ontogeny, illustrate the potentially creative

force of constraint when coupled with modularity by opening opportunities for the larger

jumps in phenotype that become possible by evolutionary changes in modules (Wagner

1988; Wagner and Altenberg 1996; Jones, et al. 2018). Here, bounds to evolution

clearly extend the range of possible phenotypic evolution. In terms of broader compara-

tive study, modularity in multivariate shape evolution has been studied via patterns of

multivariate covariation, especially for skeletal features (Parsons, et al. 2012; Jones, et

al. 2018). However, progress in applying concepts of constraint on single characters

has been hampered by a lack of precision in the concept of constraint and how it could

be detected.

Adaptation is typically envisioned as an evolutionary process whereby pheno-

types evolve in response to the strong pull toward an optimum, which tends to concen-

trate phenotypes about one or more optima after sufficient time has proceeded. On the

other hand, an evolutionary process bounded by constraint would be dominated by sto-

chastic evolution for much of the range of a phenotype, appearing as random evolution

with a substantially larger — but bounded — variance. Only when the phenotype

crosses a threshold would selection come into play, and this may occur due to corre-

lated selection on another linked trait. Thus, the evolutionary "driving force" can simply

be stochastic change with limits imposed by constraints. Our results support this theo-

retical model as an explanation for genome size evolution in salamanders. More gener-

ally, our results show that OU models, which incorporate both the change in the mean

as well as variance of the phenotype, can be used to distinguish between trait evolution

explained by a strong pull toward an optimum versus weak selection with bounded vari-

ance. Phylogenetic comparative methods have by and large focused on explaining

shifts in mean phenotype, and have thus lent themselves well to studying adaptation,

convergent evolution, and parallelism. By diving deeper into the exploration of stochas-

tic models with opportunities for varying selection and noise, we can make great pro-

gress in understanding the action of constraint in shaping evolution.

Salamanders as a model for linking organismal traits to genome biology.—Much re-

mains to be learned about how selection on organismal traits translates into changes in

genome size, and this represents a gap in our knowledge of genome biology and evolu-

tion that studies of salamanders can help to fill. For example, our results suggest that

salamanders that undergo non-feeding metamorphosis can be leveraged as a model

system to distinguish between possible mechanisms of genome reduction. Variation in

genome size is introduced into a population by TE insertions and deletions. Selection

could, in principle, sort among these genome size variants, even as the fitness conse-

quences of individual TE loci are typically miniscule and effectively neutral (Lynch 2007;

Arkhipova 2018). In addition, selection could sort among differences in TE control ma-

chinery including the pathways that underlie TE silencing and deletion, which could yield

variants with greater differences in TE composition and fitness (Mueller 2015; Parhad

and Theurkauf 2019; Parhad, et al. 2020). Comparing TE control machinery in non-

feeding metamorphosers versus other life history regimes (e.g. paedomorphs) could

identify the mechanisms underlying genome reduction.

Conclusion.— Overall, our study shows that the evolution of a life history that includes

non-feeding metamorphosis exerts a unique influence on the evolution of genome size,

imposing the most severe constraints of any salamander life history strategy. This result

suggests that selection to shorten the duration of metamorphosis because of energetic

vulnerability has indirectly selected for faster development, constraining cells and ge-

nomes to smaller sizes. We show that genome size evolution is shaped by strong sto-

chastic forces that are not widely variable across the salamander clade, but that deter-

ministic forces — which vary enormously across life histories — have also played an im-

portant role. We broaden the application of stochastic models of trait evolution to in-

clude constraint, and we highlight the challenges inherent in connecting model parame-

ters and parameter values to complex biological phenomena.

705   Literature Cited
706

Alberch P. 1989. Development and the evolution of amphibian metamorphosis. In:

Splechtna/Hilgers, editor. Trends in Vertebrate Morphology. Stuttgart: Gustav Fischer

Verlag. p. 163-173.

Alberch P. 1980. Ontogenesis and morphological diversification. Am Zool 20:653-667.

AmphibiaWeb: Information on amphibian biology and conservation [Internet]. 2021.

Berkeley, California. Available from http://amphibiaweb.org/.

Arkhipova IR. 2018. Neutral theory, transposable elements, and eukaryotic genome

evolution. Mol Biol Evol 35:1332-1337.

Arnold SJ. 1992. Constraints on phenotypic evolution. Am Nat 140:S85-S107.

Arnold SJ, Wassersug RJ. 1978. Differential predation on metamorphic anurans by

garter snakes (Thamnophis): Social behavior as a possible defense. Ecology 59:1014-

1022.

Beachy CK, Ryan TJ, Bonett RM. 2017. How metamorphosis is different in

plethodontids: Larval life history perspectives on life-cycle evolution. Herpetologica

73:252-258.

Beaulieu JM, Jhwueng D-C, Boettiger C, O'Meara BC. 2012. Modeling stabilizing

selection: Expanding the Ornstein–Uhlenbeck model of adaptive evolution. Evolution

66:2369-2383.

Boettiger C, Coop G, Ralph P. 2012. Is your phylogeny informative? Measuring the

power of comparative methods. Evolution 66:2240-2251.

Bonett RM. 2016. An integrative endocrine model for the evolution of developmental

timing and life history of plethodontids and other salamanders. Copeia 104:209-221.

Bonett RM, Hess AJ, Ledbetter NM. 2020. Facultative transitions have trouble

committing, but stable life cycles predict salamander genome size evolution. Evol Biol

47:111-122.

Bonett RM, Mueller RL, Wake DB. 2005. Why should reaquisition of larval stages by

Desmognathine salamanders surprise us? Herpetol Rev 36:112.

Bonett RM, Steffen MA, Robison GA. 2014. Heterochrony repolarized: a phylogenetic

analysis of developmental timing in plethodontid salamanders. Evo Devo 5:27.

Butler Marguerite A, King Aaron A. 2004. Phylogenetic comparative analysis: A

modeling approach for adaptive evolution. Am Nat 164:683-695.

Carta A, Bedini G, Peruzzi L. 2020. A deep dive into the ancestral chromosome number

and genome size of flowering plants. New Phytol 228:1097-1106.

Chan Y-HM, Marshall WF. 2010. Scaling properties of cell and organelle size.

Organogenesis 6:88-96.

Cheverud JM. 1984. Quantitative genetics and developmental constraints on evolution

by selection. J Theor Biol 110:155-171.

Cheverud JM, Dow MM, Leutenegger W. 1985. The quantitative assessment of

phylogenetic constraints in comparative analyses: Sexual dimorphism in body weight

among primates. Evolution 39:1335-1351.

Chippindale PT, Bonett RM, Baldwin AS, Wiens JJ. 2004. Phylogenetic evidence for a

major reversal of life-history evolution in plethodontid salamanders. Evolution 58:2809-

2822.

Cooper N, Thomas GH, Venditti C, Meade A, Freckleton RP. 2016. A cautionary note

on the use of Ornstein Uhlenbeck models in macroevolutionary studies. Biol J Linn Soc

118:64-77.

Cressler CE, Butler MA, King AA. 2015. Detecting adaptive evolution in phylogenetic

comparative analysis using the Ornstein–Uhlenbeck model. Syst Biol 64:953-968.

Deban SM, Marks SB. 2002. Metamorphosis and evolution of feeding behaviour in

salamanders of the family Plethodontidae. Zool J Linn Soc 134:375-400.

Decena-Segarra LP, Bizjak-Mali L, Kladnik A, Sessions SK, Rovito SM. 2020.

Miniaturization, genome size, and biological size in a diverse clade of salamanders. Am

Nat 196:634-648.

Downie JR, Bryce R, Smith J. 2004. Metamorphic duration: an under-studied variable in

frog life histories. Biol J Linn Soc 83:261-272.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res 32:1792-1797.

Elliott TA, Gregory TR. 2015. What's in a genome? The C-value enigma and the

evolution of eukaryotic genome content. Philos Trans R Soc B: Biol Sci 370:20140331.

Etterson JR, Shaw RG. 2001. Constraint to adaptive evolution in response to global

warming. Science 294:151.

Felsenstein J. 1985. Phylogenies and the comparative method. Am Nat 125:1-15.

Frahry MB, Sun C, Chong R, Mueller RL. 2015. Low levels of LTR retrotransposon

deletion by ectopic recombination in the gigantic genomes of salamanders. J Mol Evol

80:120-129.

Gardner JD, Laurin M, Organ CL. 2020. The relationship between genome size and

metabolic rate in extant vertebrates. Philos Trans R Soc B 375:20190146.

Gould SJ. 1980. Is a new and general theory of evolution emerging? Paleobiology

6:119-130.

Gould SJ. 1977. Ontogeny and Phylogeny. Cambridge, MA: Harvard University Press.

Grafen A, Hamilton WD. 1989. The phylogenetic regression. Philosophical Transactions

of the Royal Society of London. B, Biological Sciences 326:119-157.

Gregory TR. 2002a. A bird's eye view of the c-value enigma: genome size, cell size, and

metabolic rate in the class Aves. Evolution 56:121-130.

Gregory TR. 2001. Coincidence, coevolution, or causation? DNA content, cell size, and

the C-value enigma. Biol Rev Camb Philos Soc 76:65-101.

Gregory TR. 2005. The Evolution of the Genome. San Diego, CA: Academic Press.

Gregory TR. 2002b. Genome size and developmental complexity. Genetica 115:131-
146.

Gregory, T. R. Animal Genome Size Database (http://www.genomesize.com) [Internet].
2021.

Hansen TF. 1997. Stabilizing selection and the comparative analysis of adaptation.
Evolution 51:1341-1351.

Hansen TF, Martins EP. 1996. Translating between microevolutionary process and
macroevolutionary patterns: The correlation structure of interspecific data. Evolution
50:1404-1417.

Hessen DO, Daufresne M, Leinaas HP. 2013. Temperature-size relations from the
cellular-genomic perspective. Biol Rev 88:476-489.

Ho LST, Ané C. 2013. Asymptotic theory with hierarchical autocorrelation: Ornstein-
Uhlenbeck tree models. Ann Statist 41:957-981.

Ho LST, Ané C. 2014. Intrinsic inference difficulties for trait evolution with Ornstein-
Uhlenbeck models. Methods Ecol Evol 5:1133-1146.

Horner HA, Macgregor HC. 1983. C value and cell volume: their significance in the
evolution and development of amphibians. J Cell Sci 63:135.

Huey RB, Bennett AF. 1987. Phylogenetic studies of coadaptation: preferred
temperatures versus optimal performance temperatures of lizards. Evolution 41:1098-
1115.

Jeffery NW, Yampolsky L, Gregory TR. 2016. Nuclear DNA content correlates with

depth, body size, and diversification rate in amphipod crustaceans from ancient Lake

Baikal, Russia. Genome 60:303-309.

Jockusch EL. 1997. An evolutionary correlate of genome size change in plethodontid

salamanders. Proc Royal Soc B: Biol Sci 264:597.

Jones KE, Benitez L, Angielczyk KD, Pierce SE. 2018. Adaptation and constraint in the

evolution of the mammalian backbone. BMC Evol Biol 18:172.

Keinath MC, Timoshevskiy VA, Timoshevskaya NY, Tsonis PA, Voss SR, Smith JJ.

2015. Initial characterization of the large genome of the salamander Ambystoma

mexicanum using shotgun and laser capture chromosome sequencing. Sci Rep

5:16413.

Kerney RR, Blackburn DC, Müller H, Hanken J. 2012. Do larval traits re-evolve?

Evidence from the embryogenesis of a direct-developing salamander, *Plethodon*

cinereus. Evolution 66:252-262.

King Aaron A, Butler Marguerite A. 2009. OUCH: Ornstein-Uhlenbeck models for

phylogenetic comparative hypotheses (R package).

Kirkpatrick M, Lofsvold D. 1992. Measuring selection and constraint in the evolution of

growth. Evolution 46:954-971.

Landberg T, Azizi E. 2010. Ontogeny of escape swimming performance in the spotted

salamander. Funct Ecol 24:576-587.

Lande R. 1980. Genetic variation and phenotypic evolution during allopatric speciation.

Am Nat 116:463-479.

Licht LE, Lowcock LA. 1991. Genome size and metabolic rate in salamanders. Comp
Biochem Physiol B: Comp Biochem 100:83-92.

Liedtke HC, Gower DJ, Wilkinson M, Gomez-Mestre I. 2018. Macroevolutionary shift in
the size of amphibian genomes and the role of life history and climate. Nat Ecol Evol
2:1792-1799.

Lowe WH, Swartz LK, Addis BR, Likens GE. 2019. Hydrologic variability contributes to
reduced survival through metamorphosis in a stream salamander. Proc Natl Acad Sci
USA 116:19563.

Lynch M. 2007. The Origins of Genome Architecture. Sunderland, MA: Sinauer
Associates, Inc.

Maddison WP. 1991. Squared-change parsimony reconstructions of ancestral states for
continuous-valued characters on a phylogenetic tree. Syst Biol 40:304-314.

Madison-Villar MJ, Sun C, Lau N, Settles M, Mueller RL. 2016. Small RNAs from a big
genome: the piRNA pathway and transposable elements in the salamander species
Desmognathus fuscus. J Mol Evol 83:126-136.

Michael TP. 2014. Plant genome size variation: bloating and purging DNA. Brief Funct
Genom 13:308-317.

Mohlhenrich E, Mueller RL. 2016. Genetic drift, mutational hazard, and the evolution of
genomic gigantism in salamanders. Evolution 70:2865-2878.

Moran NA. 1994. Adaptation and constraint in the complex life cycles of animals. Ann
Rev Ecol Syst 25:573-600.

Mueller RL. 2015. Genome biology and the evolution of cell size diversity. In: Heald R,

Hariharan I, Wake DB, editors. Size Control in Biology, from Organelles to Organisms:

Cold Spring Harbor Press.

Mueller RL, Gregory TR, Gregory SM, Hsieh A, Boore JL. 2008. Genome size, cell size,

and the evolution of enucleated erythrocytes in attenuate salamanders. Zoology

111:218-230.

Mueller RL, Macey JR, Jaekel M, Wake DB, Boore JL. 2004. Morphological homoplasy,

life history evolution, and historical biogeography of plethodontid salamanders inferred

from complete mitochondrial genomes. Proc Natl Acad Sci USA 101:13820-13825.

Norman MF. 1985. A practical method for staging metamorphosis in the tiger

salamander *Ambystoma tigrinum*. Anatom Rec 211:102-109.

Nowoshilow S, Schloissnig S, Fei J-F, Dahl A, Pang AWC, Pippel M, Winkler S, Hastie

AR, Young G, Roscito JG, et al. 2018. The axolotl genome and the evolution of key

tissue formation regulators. Nature 554:50-55.

O'Meara BC, Ané C, Sanderson MJ, Wainwright PC. 2006. Testing for different rates of

continuous trait evolution using likelihood. Evolution 60:922-933.

Olmo E, Capriglione T, Odierna G. 1989. Genome size evolution in vertebrates: Trends

and constraints. Comp Biochem Phys B: Comp Biochem 92:447-453.

Orlofske SA, Hopkins WA. 2009. Energetics of metamorphic climax in the pickerel frog

(*Lithobates palustris*). Comp Biochem Phys A: Molec Integ Phys 154:191-196.

Pandit MK, White SM, Pocock MJO. 2014. The contrasting effects of genome size,

chromosome number and ploidy level on plant invasiveness: a global analysis. New

Phytol 203:697-703.

Parhad SS, Theurkauf WE. 2019. Rapid evolution and conserved function of the piRNA

pathway. Roy Soc Open Biol 9:180181.

Parhad SS, Yu T, Zhang G, Rice NP, Weng Z, Theurkauf WE. 2020. Adaptive evolution

targets a piRNA precursor transcription network. Cell Rep 30:2672-2685.e2675.

Parsons KJ, Márquez E, Albertson RC. 2012. Constraint and opportunity: The genetic

basis and evolution of modularity in the cichlid mandible. Am Nat 179:64-78.

Pasquesi GIM, Adams RH, Card DC, Schield DR, Corbin AB, Perry BW, Reyes-Velasco

J, Ruggiero RP, Vandewege MW, Shortt JA, et al. 2018. Squamate reptiles challenge

paradigms of genomic repeat element evolution set by birds and mammals. Nat

Commun 9:2774.

Posada D, Crandall KA. 1998. MODELTEST: testing the model of DNA substitution.

Bioinformatics 14:817-818.

Pyron AR, Wiens JJ. 2011. A large-scale phylogeny of Amphibia including over 2800

species, and a revised classification of extant frogs, salamanders, and caecilians. Mol

Phylogenet Evol 61:543-583.

R Core Team. 2020. R: A language and environment for statistical computing. Vienna,

Austria: R Foundation for Statistical Computing.

Roddy AB, Théroux-Rancourt G, Abbo T, Benedetti JW, Brodersen CR, Castro M,

Castro S, Gilbride AB, Jensen B, Jiang G-F, et al. 2019. The scaling of genome size

and cell size limits maximum rates of photosynthesis with implications for ecological

strategies. Int J Plant Sci 181:75-87.

Rose CS. 2014. The importance of cartilage to amphibian development and evolution.

Int J Dev Biol 58:917-927.

Rose CS. 1995a. Skeletal morphogenesis in the urodele skull: II. Effect of

developmental stage in thyroid hormone-induced remodeling. J Morph 223:149-166.

Rose CS. 1995b. Skeletal morphogenesis in the urodele skull: III. Effect of hormone

dosage in th-induced remodeling. J Morph 223:243-261.

Rose CS. 1995c. Skeletal morphogenesis in the urodele skull:I. Postembryonic

development in the hemidactyliini (amphibia: Plethodontidae). J Morph 223:125-148.

Sanchez E, Küpfer E, Goedbloed DJ, Nolte AW, Lüddecke T, Schulz S, Vences M,

Steinfartz S. 2018. Morphological and transcriptomic analyses reveal three discrete

primary stages of postembryonic development in the common fire salamander,

Salamandra salamandra. J Exp Zool B: 330:96-108.

Sanderson MJ. 2003. r8s: inferring absolute rates of molecular evolution and

divergence times in the absence of a molecular clock. Bioinformatics 19:301-302.

Schluter D, Price T, Mooers AØ, Ludwig D. 1997. Likelihood of ancestor states in

adaptive radiation. Evolution 51:1699-1711.

Sessions SK. 2008. Evolutionary cytogenetics in salamanders. Chromosome Res

16:183-201.

Shao F, Han M, Peng Z. 2019. Evolution and diversity of transposable elements in fish
genomes. Sci Rep 9:15399.

Simpson GG. 1953. The Major Features of Evolution. New York: Columbia University
Press.

Smith JDL, Bickham JW, Gregory TR. 2013. Patterns of genome size diversity in bats
(order Chiroptera). Genome 56:457-472.

Sotero-Caio CG, Platt RN, II, Suh A, Ray DA. 2017. Evolution and diversity of
transposable elements in vertebrate genomes. Genome Biol Evol 9:161-177.

Stamatakis A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses
with thousands of taxa and mixed models. Bioinformatics 22:2688-2690.

Sun C, Arriaza JRL, Mueller RL. 2012. Slow DNA loss in the gigantic genomes of
salamanders. Genome Biol Evol 4:1340-1348.

Sun C, Mueller RL. 2014. Hellbender genome sequences shed light on genome
expansion at the base of crown salamanders. Genome Biol Evol 6:1818-1829.

Sun C, Shepard DB, Chong RA, Arriaza JL, Hall K, Castoe TA, Feschotte C, Pollock
DD, Mueller RL. 2012. LTR retrotransposons contribute to genomic gigantism in
plethodontid salamanders. Genome Biol Evol 4:168-183.

Szarski H. 1983. Cell size and the concept of wasteful and frugal evolutionary
strategies. J Theor Biol 105:201-209.

Uyeda JC, Pennell MW, Miller ET, Maia R, McClain CR. 2017. The evolution of
energetic scaling across the vertebrate tree of life. Am Nat 190:185-199.

Vieites DR, Rom·n SN, Wake MH, Wake DB. 2011. A multigenic perspective on
phylogenetic relationships in the largest family of salamanders, the Plethodontidae. Mol
Phylogenet Evol 59:623-635.

Vinogradov AE. 2004. Genome size and extinction risk in vertebrates. Proc Roy Soc B
271:1701-1705.

Vladimirova IG, Kleimenov SY, Alekseeva TA. 2012. Dynamics of body mass and
oxygen consumption in the ontogeny of the Spanish ribbed newt (*Pleurodeles waltl*): 2.
Larval stage. Biol Bull 39:10-14.

Vu GTH, Cao HX, Reiss B, Schubert I. 2017. Deletion-bias in DNA double-strand break
repair differentially contributes to plant genome shrinkage. New Phytol 214:1712-1721.

Wagner GP. 1988. The significance of developmental constraints for phenotypic
evolution by natural selection. In: de Jong G, editor. Population Genetics and Evolution.
Berlin, Heidelberg: Springer.

Wagner GP, Altenberg L. 1996. Perspective: Complex adaptations and the evolution of
evolvability. Evolution 50:967-976.

Wake D, Hanken J. 2004. Direct development in the lungless salamanders: what are
the consequences for developmental biology, evolution and phylogenesis? Int J Dev
Biol 40:859-869.

Wake DB, Marks SB. 1993. Development and evolution of plethodontid salamanders: a
review of prior studies and a prospectus for future research. Herpetologica 49:194-203.

Waltari E, Edwards SV. 2002. Evolutionary dynamics of intron size, genome size, and
physiological correlates in archosaurs. Am Nat 160:539-552.

Wassersug RJ, Sperry DG. 1977. The relationships of locomotion to differential

predation on *Pseudacris triseriata* (anura: Hylidae). Ecology 58:830-839.

Wiens JJ, Bonett RM, Chippindale PT. 2005. Ontogeny discombobulates phylogeny:

paedomorphosis and higher-level salamander relationships. Syst Biol 54:91-110.

Wright ML, Richardson SE, Bigos JM. 2011. The fat body of bullfrog (*Lithobates*

catesbeianus) tadpoles during metamorphosis: Changes in mass, histology, and

melatonin content and effect of food deprivation. Comp Biochem Phys A 160:498-503.

Zheng Y, Peng R, Kuro-o M, Zeng X. 2011. Exploring patterns and extent of bias in

estimating divergence time from mitochondrial DNA sequence data in a particular

lineage: a case study of salamanders (Order Caudata). Mol Biol Evol 28:2521-2535.