1

3 **Metamorphosis imposes variable constraints on genome expansion**

4 **through effects on development**

6 **Article type:** Original Research

8 **Keywords**: genome size, metamorphic vulnerability, developmental system, compara-

9 tive methods, life history, salamander

Abstract

Genome size varies ~ 100,000-fold across eukaryotes and has long been hypothesized to be influenced by metamorphosis in animals. Transposable element accumulation has been identified as a major driver of increase, but the nature of constraints limiting the size of genomes has remained unclear, even as traits such as cell size and rate of development co-vary strongly with genome size. Salamanders, which possess diverse metamorphic and non-metamorphic life histories, have the largest vertebrate genomes ⸺ 3 to 40 times that of humans ⸺ as well as the largest range of variation in genome size. We tested 13 biologically-inspired hypotheses exploring how the form of metamorphosis imposes varying constraints on genome expansion in a broadly representative phylogeny containing 118 species of salamanders. We show that metamorphosis during which animals undergo the most extensive and synchronous remodeling imposes the most severe constraint against genome expansion, with the severity of constraint decreasing with reduced extent and synchronicity of remodeling. More generally, our work demonstrates the potential for broader interpretation of phylogenetic comparative analysis in exploring the balance of multiple evolutionary pressures shaping phenotypic evolution.

Introduction

Across the tree of life, few traits exhibit the tremendous scale of variation shown by genome size, which encompasses a ~100,000-fold range across eukaryotes alone (Gregory 2022). Decades of research have explored the question of whether this variation has a cohesive evolutionary explanation, revealing two traits that consistently covary with large genome size: cell division rate slows down and cell size increases (Gregory 2001, 2005). However, whether genome size evolves by adaptation or constraint, and what drives these processes, have been challenging questions to answer, in part because the organismal features involved have been unclear. For example, genome size co-varies in a context-dependent manner with metabolic rate, showing an association within some vertebrate clades, but not others, and shows no correlation across vertebrates as a whole (Licht and Lowcock 1991; Gregory 2002a; Smith, et al. 2013; Wright, et al. 2014; Kapusta, et al. 2017; Uyeda, et al. 2017; Gardner, et al. 2020). Genome size has also been associated with developmental rate or complexity (Gregory 2002b), temperature (Hessen, et al. 2013), invasiveness (Pandit, et al. 2014), and speciation and extinction rates (Vinogradov 2004; Jeffery, et al. 2016). As these associations vary in their generality, the explanation may not lie in a single overarching factor. Rather, genome size variation may reflect the balance of multiple evolutionary pressures. Thus, to disentangle the forces affecting genome size requires close consideration of the interaction of ecology and organismal biology (Gregory 2004; Roddy, et al. 2019).

The association between genome size and metamorphosis has been repeatedly noted across ectotherms (salamanders: Larson 1984; Wake and Marks 1993; Gregory

2002b; Sessions 2008; fish and insects: Gregory 2002b). It has long been proposed that

natural selection acts to shorten the duration of metamorphosis (Szarski 1957) to limit

exposure to potentially lethal stresses during transformation, termed "metamorphic vul-

nerabilities" (Gregory 2002b; Lowe, et al. 2021). Salamanders have received special fo-

cus because they have the largest genomes among vertebrates (9 Gb – 120 Gb per

haploid genome; the vast majority are diploid), as well as exceptional diversity in life his-

tory across the 773 extant species (Decena-Segarra, et al. 2020; AmphibiaWeb 2022;

Gregory 2022). Metamorphosis has been lost, modified, and regained throughout the

clade's evolutionary history leading to metamorphosers, paedomorphs, and direct de-

velopers among extant salamanders. These three life history types vary widely in the

degree and duration of transformation, as well as many other organismal features that

have important consequences for natural selection. Metamorphosers undergo a mor-

phological transformation from an aquatic larval to terrestrial adult form, paedomorphs

retain the aquatic larval morphology throughout life, and direct developers hatch from

terrestrial eggs as miniature versions of the adults (Rose 1996; Rose 1999;

Chippindale, et al. 2004; Mueller, et al. 2004; Wiens, et al. 2005; Bonett, et al. 2014).

The observation that metamorphosis is linked to smaller genome size (Wake and

Marks 1993; Gregory 2002b; Sessions 2008; Bonett, et al. 2020), and larger genome

size is linked to an overall slow-down of developmental rates, led to the hypothesis of

time-limited metamorphosis as a constraint on genome expansion (Gregory 2002b). An

evolutionary constraint is broadly defined as a limit to phenotypic variation (Arnold 1992)

that can arise from non-adaptive sources or from indirect selection on a correlated trait

(e.g., Savell, et al. 2016). For example, because the rate of development is an emergent

property of complex cellular processes that are slowed down by large genome/cell sizes

(Horner and Macgregor 1983; Jockusch 1997), selection to shorten metamorphosis

should limit the sizes that genomes can attain, resulting in a constraint on genome size.

There has been some ecological confirmation of the time-limited metamorphosis hy-

pothesis in salamanders, with species in ephemeral habitats possessing smaller ge-

nomes (Lertzman-Lepofsky, et al. 2019). However, large-scale phylogenetic compara-

tive analysis has both rejected (Liedtke, et al. 2018) and confirmed the classical associ-

ation between genome size and life history (Bonett, et al. 2020). A key difference in the

latter was partitioning life history diversity to better reflect how developmental complex-

ity influences genome size. Combining lineages with the ability to metamorphose at all

— even if only occasionally, as seen in facultative paedomorphs — and separating

them from lineages that are obligate paedomorphs was key to finding any association

between genome size and life history (Bonett, et al. 2020).

Although this was a big step forward, there was no clear best model; a model

that separated direct developers from paedomorphs and a model that grouped them as

non-metamorphosers explained the data nearly equally well. This result is surprising if

developmental remodeling is a primary factor influencing genome size because direct

developers undergo a substantial amount of metamorphic remodeling inside the egg.

We hypothesize that this result may be explained by the models not yet considering nat-

ural diversity across salamander lineages in the synchronicity and extent of develop-

mental remodeling during metamorphosis, as well as the associated vulnerabilities.

Thus, the relevant features of metamorphosing organisms, and the evolutionary pres-

sures they exert on genome size, likely remain incompletely understood.

In this study, we explore how metamorphosis shapes genome size, expanding

the consideration of life history diversity to include natural variation in developmental

complexity during metamorphosis and associated metamorphic vulnerabilities. We in-

form stochastic models of genome size evolution with life history data and the molecular

mechanisms of genome expansion. We detail how OU model-based comparative meth-

ods can be used to explore adaptation and constraint in genome size and, by extension,

in other traits that also may not be shaped exclusively by adaptive evolution. Whereas

OU models have been used widely to study adaptive significance (Hansen 1997; Butler

and King 2004), some traits such as extremely large genome size have no known fit-

ness benefit, which is at odds with an interpretation of adaptive evolution. Indeed, these

general models can be interpreted in multiple ways, and rather than evolutionary "op-

tima," the phenotypic locations predicted by the models may be better envisioned as

"equilibria" to reflect a balance of forces such as upwardly-biased mutation pressure op-

posed by a constraint set by other aspects of organismal biology. The sigma term, de-

scribing the intensity of random fluctuations of the evolutionary process, has received

much less attention, but in combination with the other parameters may be helpful in di-

agnosing release from constraint.

Methods

Genome Size and Life History Regimes

Stochastic Genome Expansion In vertebrates, genome size is strongly shaped by

transposable elements (TEs), sequences that replicate and spread throughout host ge-

nomes, increasing genome size (Sotero-Caio, et al. 2017; Shao, et al. 2019). TEs can

be deleted by errors in replication, recombination, and DNA repair (Michael 2014; Vu, et

al. 2017). TE activity can be nearly neutral, largely missing functional genomic regions

and therefore resulting in negligible fitness consequences (Arkhipova 2018). Salaman-

ders are particularly prone to stochastic increases in genome size through TE accumu-

lation (Sun, et al. 2012; Keinath, et al. 2015; Nowoshilow, et al. 2018) because they

possess low TE deletion rates and incomplete TE silencing (Sun and Mueller 2014;

Frahry, et al. 2015; Madison-Villar, et al. 2016). Against this clade-wide background of

biased stochastic genome expansion, we hypothesize that different life history regimes

exert varying levels of constraint on genome size mediated through exposure to meta-

morphic vulnerabilities as explained below.

Metamorphic Vulnerabilities Metamorphosis is a radical morphological transformation

often coordinated in a short window of time, and thus subject to multiple risks. During

metamorphosis, animals undergo rapid cell division, differentiation, migration, and apop-

tosis (Alberch 1989), as they remodel their skin and glands, blood, gut, teeth, and mus-

culoskeletal, excretory, and immune systems (Rose 1999). Accordingly, metamorphos-

ing animals can experience three classes of vulnerabilities: 1) <u>Performance and physio-

logical handicap.</u> Animals can suffer reductions in performance in escaping predation or

accessing food and shelter; or reductions in physiological tolerance to environmental

swings during transformation. 2) <u>Energetic limitations</u> imposed when resorbing or re-

modeling larval structures and forming adult structures (Wassersug and Sperry 1977;

Orlofske and Hopkins 2009; Enriquez-Urzelai, et al. 2019; Lowe, et al. 2019; Lowe, et

al. 2021). 3) Random developmental errors resulting from developmental system pertur-

bation, which can become more likely with both increased developmental complexity

and reduced cell numbers when remodeling occurs at earlier embryonic stages (Hanken

1984; Gregory 2002b; Rose 2003).

In addition to the ancestral form of metamorphosis, salamanders have evolved a

derived mode that differs markedly in timing and extent of remodeling (Alberch, et al.

1985; Rose 1999, 2003; Beachy, et al. 2017) In contrast to previous studies, we there-

fore differentiate four life history regimes for the evolution of genome size: Paedomor-

phosis, Direct Development, and two forms of metamorphosis called Gradual Progres-

sive Metamorphosis and Abrupt Synchronous Metamorphosis. Because they account

for < 2% of salamander species, we did not include viviparous and ovoviviparous life

histories.

Paedomorphosis: Individuals do not undergo metamorphosis, instead reaching sexual

maturity retaining largely larval traits without any radical developmental remodeling

(Gould 1977; Rose 1996). Paedomorphosis has evolved at least eight times within sala-

manders (Rose 1999). Paedomorphs experience no metamorphic vulnerabilities and

are thus predicted to be free from any related constraints on genome expansion.

Direct development: The larval growth stage is eliminated, and metamorphosis is inte-

grated with embryogenesis into a single sequence of developmental events that takes

place much earlier in ontogeny, inside the egg (Alberch 1989; Rose 2014). Thus, meta-

morphic remodeling occurs, but the events involve small amounts of tissue and few

cells. Direct development has likely evolved at least twice within salamanders (Bonett,

et al. 2014). Direct developers experience the metamorphic vulnerabilities of energetic

limitation and developmental error due to decreased cell numbers. Thus, direct develop-

ers are predicted to have intermediate levels of constraint on genome size associated

with metamorphosis.

Gradual, progressive metamorphosis: This type of metamorphosis is likely ancestral for

salamanders (Rose 1999), and involves remodeling events happening sequentially and

over relatively long timeframes in a free-living, aquatic organism in preparation for tran-

sition to a terrestrial habitat (Rose 1996). In contrast to frogs, these salamander larvae

are able to feed throughout the process, as well as use larval energy stores (Semlitsch,

et al. 1988; Deban and Marks 2002), alleviating energetic limitations. Furthermore, the

timing of metamorphic onset is flexible and can be delayed when larval food resources

are limited (Beachy, et al. 2017). Gradual metamorphosers experience the metamorphic

vulnerabilities of decreased performance and physiological tolerance and are thus pre-

dicted to have intermediate levels of constraint on genome size associated with meta-

morphosis.

Abrupt, synchronous metamorphosis: This form of metamorphosis differs markedly from

gradual, progressive metamorphosis in that remodeling events are more radical, occur

simultaneously, and begin at a fixed time point in larval development, irrespective of lar-

val size (Alberch, et al. 1985; Rose 1999, 2003; Beachy, et al. 2017). Organisms un-

dergo a more extensive remodeling of the feeding apparatus and consequently trans-

form with reduced feeding performance (Deban and Marks 2002). This type of meta-

morphosis has evolved approximately three times within salamanders (Bonett, et al.

2014). Abrupt metamorphosers experience the metamorphic vulnerabilities of de-

creased performance and physiological tolerance, energetic demand, and developmen-

tal error due to increased complexity and are thus predicted to have the most extreme

levels of constraint on genome expansion associated with metamorphosis.

200
201
202 Figure 1. Alternative hypotheses for constraints imposed by development on genome
203 size evolution in salamanders. On each phylogeny (www.vertlife.org), alternative life his-
204 tory regimes are painted in different colors as indicated in each legend (see text). Hap-
205 loid genome sizes are shown on the right in pg (1 pg = 978 Mb).
206

Taxon Sampling, Genome Size, and Phylogeny

We analyzed haploid genome size data for 118 species of salamanders (all of which are diploid), including all ten families and 33 of 68 genera, and representing all four life-history regimes (Supplemental Table 1) (AmphibiaWeb 2022). We excluded miniaturized taxa, as miniaturization is associated with decreased genome size independent from any selection imposed by metamorphosis (Decena-Segarra, et al. 2020). Several lineages are facultative paedomorphs; we coded these taxa as metamorphic, as they retain the need to successfully (albeit occasionally) metamorphose (Bonett, et al. 2020).

The dataset includes all species that are represented in the Animal Genome Size Database (www.genomesize.com) and the VertLife database for phylogeny subsampling (www.vertlife.org) (Jetz and Pyron 2018; Gregory 2022). We took the mean genome size where estimates were reported from multiple studies, and natural-log transformed the data to better conform with assumptions of Gaussian errors. We excluded one study that reported consistently higher values than all others (Bachmann 1970). We obtained the 118-species phylogeny by sampling 1,000 ultrametric trees from the pseudo-posterior distribution of the VertLife database (Jetz and Pyron 2018) and computing mean branch lengths using the consensus.edges function in the R package phytools v 0.7-90 (Revell 2012) (Fig. 1). All analyses were conducted in the R statistical computing environment (R Core Team 2020).

Models of Genome Size Evolution

We modeled genome size evolution using both Brownian motion (BM) and OU models of evolution (Hansen 1997; Butler and King 2004; O'Meara, et al. 2006; Beaulieu, et al.

2012). The BM model is the simplest stochastic model with a single rate parameter σ for

stochastic noise intensity describing the magnitude of the independent random walks of

the trait evolving along the branches of the phylogeny. The multiple-rate BM model al-

lows σ to vary across a phylogeny (O'Meara, et al. 2006).

OU models generalize the BM model by allowing the mean to shift and the vari-

ance to narrow. They include a deterministic component of trait evolution that models

the tendency to move toward an equilibrium. In contrast to previous work, and to more

closely reflect our hypothesis that salamander genome size evolves in response to a

balance of mutation pressure and deterministic forces such as selection or constraint,

we generalize the notion of "selective optima" to "*deterministic equilibria*", which encom-

passes both adaptive and non-adaptive forces affecting the equilibrium ($\theta(t)$). Similarly,

we generalize "selection" to "*deterministic pull*" to encompass non-selective, but direc-

tional, evolutionary forces (α; e.g., biased mutation pressure and constraint in addition

to selection). Mathematically, the model for trait evolution expressed as a differential

equation is

$$dX(t) = \alpha(\theta(t) - X(t)) + \sigma dB(t),$$

where $\theta(t)$ is the *deterministic equilibrium* for the trait at time *t* and α is an evolutionary

rate describing the *strength of the deterministic pull* towards that equilibrium. The sim-

plest OU models allow equilibria to vary across the tree, reflecting the evolution of differ-

ences in mean phenotype across regimes (Hansen 1997; Butler and King 2004). Fur-

ther model extensions also allow the strength of the deterministic pull and the stochastic

noise intensity to vary across regimes (Beaulieu, et al. 2012).

We formalized five hypotheses for the influence of metamorphosis on genome

size evolution: (1) *random evolution*: With no influence of metamorphosis, stochastic

evolutionary processes may be sufficient to explain genome size evolution, as repre-

sented by BM. The remaining four hypotheses propose different groupings of con-

straint fit with OU models. (2) *metamorphosis-other*: Metamorphosis imposes a con-

straint on genome expansion distinct from all remaining life histories, grouped as

"other". (3) *meta-paed-dd*: This hypothesis refines (2) by dividing the "other" category

into direct developers and paedomorphs, allowing each to impose distinct constraints

on genome size evolution. (4) *meta$_{abrupt}$-meta$_{gradual}$-other*: Alternatively, we may refine (2)

by keeping the "other" category of non-metamorphosers, but differentiating the two

groups of metamorphosers, recognizing that abrupt (*meta$_{abrupt}$*) and gradual (*meta$_{gradual}$*)

metamorphosis each impose distinct constraints on genome size evolution. (5)

meta$_{abript}$-meta$_{gradual}$-paed-dd: Each category imposes distinct constraints on genome

expansion (Fig. 1).

We tested 21 models that varied in the number of parameters used to explore

these five biologically-inspired hypotheses. The simplest model allows the equilibria to

vary with life history regime while modeling a single stochastic noise intensity (σ) and

deterministic pull strength (α) across all species. Additional sub-hypotheses fit evolu-

tionary models with separate stochastic noise intensities (σ_i) for each regime (Table 1).

More complex models with multiple deterministic pull strengths (α_i) produced fitting er-

rors or nonsensical parameter estimates or likelihoods, so we do not consider them fur-

ther (see Supplementary Information; Table 2). We fit the remaining 13 models with the

ancestral plethodontid node assigned to be either metamorphosing or direct-developing

(Bonett, et al. 2014) and found that the choice made no qualitative difference (Supple-

mentary Information). Model fitting and parameter estimation were carried out using

OUwie (Beaulieu, et al. 2012).

Table 1. BM and OU models with single or multiple parameters used to fit the data.
Numbers in parentheses specify (1) model parameters and notation, (2) parameters that
remain constant across the phylogeny, (3) parameters that vary with shifts in life history
regime, (4) OUwie model notation, and (5) notes for the model implementations and ci-
tations.
285

Models	Uniform	Multiple with Regime	OUwie Notation	Notes
(1)	(2)	(3)	(4)	(5)
Single noise intensity σ	σ		BM1	Classic BM model of (Felsenstein 1985)
Multiple noise intensities σ_i		σ_i	BMS	Multiple-rate BM model of O'Meara et al. (2006)
Multiple equilibria θ_i, σ, α	σ α	θ_i	OUM	OU model of Hansen (1997) and Butler and King (2004)
Multiple equilibria and deterministic pull strengths $\theta_i, \sigma, \alpha_i$	σ	α_i θ_I	OUMA	Multiple-α model of Beaulieu et al. (2012)
Multiple equilibria and noise intensities $\theta_i, \sigma_i, \alpha$	α	σ_i θ_i	OUMV	Multiple-σ model of Beaulieu et al. (2012)

Multiple equilibria, noise intensities, and deterministic pull strengths $\theta_i, \sigma_i, \alpha_i$		σ_i α_i θ_i	OUMVA	Full model of Beaulieu et al. (2012)

Model Comparison

We compared the fit of each of the models using the Akaike Information Criterion corrected for small sample size (AIC_c). Because AIC_c differences can favor more complex models even when a simpler one is correct, we performed model selection bootstrap analysis (phylogenetic Monte Carlo; (Boettiger, et al. 2012). We additionally evaluated the support for hypotheses in six pairwise comparisons (Fig. 2) which assess and progressively refine the strength of evidence for successive levels of increased model complexity as well as the power to detect differences in model support. Support for the more complex model in each pair implies:

(A) *BM vs. metamorphosis-other:* metamorphosis imposes a constraint on genome expansion;

(B) *meta-other vs. meta-dd-paed:* distinct constraints are imposed by the different non-metamorphosing strategies, direct development and paedomorphosis;

(C) *meta-other vs. abrupt-gradual-other:* abrupt metamorphosis imposes a distinct constraint from gradual metamorphosis;

(D) *meta-dd-paed vs. abrupt-gradual-dd-paed:* abrupt metamorphosis imposes a distinct constraint from gradual metamorphosis, after accounting for differences in non-metamorphosing strategies;

(E) *abrupt-gradual-other vs. abrupt-gradual-dd-paed:* non-metamorphosing strategies impose unique constraints, after accounting for differences between abrupt and gradual metamorphosis;

(F) after identifying *meta$_{abrupt}$-meta$_{gradual}$-paed-dd* as the best-fitting hypothesis,

we further explored the power to discriminate between multiple versus single stochastic

noise intensity parameters.

For each comparison, we computed the observed likelihood difference,

$$\delta_{obs} = -2 (\log L_0 - \log L_1),$$

where L_0 is the likelihood of the simpler model and L_1 is the likelihood of the more com-

plex model. We used these parameters and stochastic simulation to compute approxi-

mate *p*-values and power.

Determining whether δ_{obs} is significantly different from a null expectation requires

an approximate *p*-value — the probability of observing δ_{obs} *if the simpler model were*

true. That is, we need to compare the value δ_{obs} to the distribution of δ values under the

simpler model. To create this distribution, we generated 500 datasets by simulating the

simpler model at its MLE parameter estimates; we then fit both the simpler and more

complex models to each simulated dataset and computed the values of δ, producing a

null distribution of δ assuming the simpler model. We compared the observed value of δ

to this null distribution to calculate an approximate *p*-value.

Power conveys the (desirable) probability of rejecting the simpler model when the

more complex model is true. To estimate power, we generated 500 datasets by simulat-

ing the more complex model at its MLE parameter estimates; we then fit the two models

and computed the values of δ. The fraction of these δ values that are greater than the

95% quantile of the distribution generated under the simpler model (described above)

gives an estimate of power. All data and code necessary to carry out the analysis in this

manuscript can be found at https://github.com/claycressler/genomesize and in Supple-

mental Material.

Results

The best-fitting model for salamander genome size evolution accounted for four re-

gimes: both abrupt and gradual metamorphosis, paedomorphosis, and direct develop-

ment (*meta_abrupt -meta_gradual -paed-dd*; Table 2) under an OU model that allowed both

equilibrium genome size and noise intensity to vary across these regimes (θ_i, σ_i, α, Ta-

ble 2). The three-regime *meta_abrupt -meta_gradual -other* (θ_i, σ_i, α) and *meta-dd-paed* (θ_i, σ_i,

α) hypotheses also had marginal support ($\Delta AIC_c \leq 3$).

Several authors have recommended parametric and model selection bootstraps

to assess OU models, as model selection tends to be robust while parameters can be

difficult to estimate accurately (Boettiger, et al. 2012; Ho and Ané 2013; Ho and Ané

2014; Cressler, et al. 2015). We found no evidence of identifiability issues; AIC_c values

correspond with parametric bootstrap results, and parameter estimates and confidence

intervals are well-behaved (Table 3, Fig. 2,3). We furthermore avoid interpreting point

estimates of parameters, limiting our discussion to features of the best models, relative

rankings of parameter estimates across regimes bolstered by AIC_c-derived confidence

intervals, and comparison of variation across models.

We reject a purely stochastic hypothesis for genome size evolution based on

both the results of model fitting and interrogation by parametric bootstrap (Table 2, Fig.

2). The BM model had the highest AIC_c, and a model that accounts for metamorphosis

vs. others with separate equilibrium and noise intensity values was far superior to a

purely neutral model (Table 2; Fig. 2a). Models that separate non-metamorphosing

strategies into direct development and paedomorphosis (comparing *metamorphosis-*

other to *meta-dd-paed*; Fig. 2b) or separate metamorphosing strategies into gradual and

abrupt metamorphosis (comparing *metamorphosis-other* to *meta$_{abrupt}$-meta$_{gradual}$-other*;

Fig. 2c) had only moderate power to reject the simpler *metamorphosis-other* hypothesis.

However, models that included all four selective regimes (*meta$_{abrupt}$-meta$_{gradual}$-dd-paed*)

had high power to reject any three-regime hypothesis (Fig. 2d,e). These model selection

bootstrapping results support our conclusion, based on the AIC$_c$ values obtained from

fitting the real data, that these life history regimes have different deterministic equilibria

(Table 3), and moreover, allowing noise intensity to vary among regimes is strongly sup-

ported (Fig. 2f). Therefore, we have compelling evidence that observed differences in

genome size between abrupt metamorphosers, gradual metamorphosers, direct devel-

opers, and paedomorphs (Fig. 3) reflect differences in the balance of evolutionary

forces shaping genome size in each regime.

Table 2. Model comparison statistics. Best model (interrogated by bootstrap, Fig. 2) in-
dicated in bold. Model parameterizations are indicated by: σ = Brownian motion; σ_i =
Brownian motion with multiple noise intensities; θ_i, σ, α = OU model with multiple equi-
libria; θ_i, σ_i, α = OU model with multiple equilibria and multiple noise intensities.
376

Hypotheses	ΔAIC_c			
	OU Models		BM Models	
	$\theta i, \sigma i, \alpha$	$\theta i, \sigma, \alpha$	σi	σ
meta$_{abrupt}$ -meta$_{gradual}$ - paed-dd	**0**	7.1	7.5	
meta$_{abrupt}$ -meta$_{gradual}$ - other	2.5	6.9	9.5	
meta-paed-dd	3.0	8.8	5.3	
metamorphosis-other	4.2	7.5	7.6	
Brownian motion				15.5

Figure 2. The power to discriminate among competing life-history regime models. Each
panel evaluates the support for a different, more complex, hypothesis relative to a sim-
pler one (panel keys). Pairwise bootstrap distributions of the likelihood difference (δ)
calculated by generating 500 datasets under each of two competing life history regime
models at their MLE parameter estimates, refitting the two models, and computing δ. All
comparisons are between multiple equilibria, multiple noise intensity models. Assuming
the simpler model is the truth, the probability density of δ is in light gray, while the den-
sity of the likelihood difference assuming the complex model is in dark gray. The dashed
line gives the observed value (δ_{obs}) from fitting the actual genome size data. The re-
ported *p*-value is the fraction of the light gray distribution that lies to the right of δ_{obs}; the
power is the fraction of the dark gray distribution that lies to the right of the 95[th] percen-
tile of the light gray distribution.
400

401
Parameter estimates for the best-fitting model are presented in Table 3. Abrupt meta-

morphosers have the smallest genome size equilibrium, gradual metamorphosers and

direct developing salamanders are intermediate, and paedomorphic salamanders have

the largest (Table 3), with multiple categories of stochastic noise intensity. To help visu-

alize what this best-fitting model tells us about genome size evolution, we compared the

distribution of observed genome sizes for species in each selective regime to the ex-

pected stationary distribution, given the model estimates of θ, α, and σ for each regime

(Fig. 3). In particular, (Ho and Ané 2013) showed that the stationary distribution of an

OU process will be a normal distribution with mean θ and variance $\sigma^2/(2\alpha)$, providing a

useful way to visualize the model fit that captures aspects of all parameters. In the di-

rect-developing regime (which is the most common across the tree), the expected distri-

bution completely overlaps the observed (Fig. 3c). The observed and expected distribu-

tions of genome size for gradual metamorphosers are broadly overlapping, although the

mean of the expected distribution is slightly higher than the observed (Fig. 3b); similar

patterns are observed for paedomorphs (Fig. 3d). In contrast, the expected mean is

smaller than almost all abrupt metamorphosing salamander genome sizes (Fig. 3a).

Furthermore, this best-fit model predicts that the variance of the trait distribution for

each regime is well-predicted by $\sigma^2/(2\alpha)$, which varies by regime because of differ-

ences in the noise parameter σ_i. Direct developers have larger stochastic noise in ge-

nome size than abrupt metamorphosers or paedomorphs, with gradual metamorphosers

intermediate (Table 3, Fig. 3).

424
425

426 Table 3: Maximum likelihood parameter estimates and parametric bootstrap confidence
427 intervals for the best-fitting model (*meta*$_{abrupt}$-*meta*$_{gradual}$-dd-paed $\theta i, \sigma i, \alpha$: separate
428 equilibrium values and noise intensities for lineages in the four life history regimes: dd =
429 direct development, meta$_{abrupt}$ = abrupt metamorphosis, meta$_{gradual}$ = gradual metamor-
430 phosis, paed = paedomorphosis)
431

Parameter	Abrupt Metamorpho­sis	Gradual Metamorpho­sis	Direct Develop­ment	Paedomorpho­sis
Deterministic pull (α)		1.27		
95% CI		(0.43, 4.24)		
Equilibrium value (θ_i)	2.65	3.66	3.72	4.31
95% CI	(0.71, 3.39)	(3.39, 3.93)	(2.87, 4.58)	(3.87, 5.03)
Stochastic noise intensity (σ_i)	0.38	0.59	0.84	0.36
95% CI	(0.22, 0.56)	(0.41, 0.88)	(0.72, 1.08)	(0.14, 0.68)

450
451
452 Figure 3. Comparison of the observed distribution of genome sizes for salamanders in
453 each regime (bars) against the stationary distribution predicted by the best-fitting pa-
454 rameter set (dotted lines, Table 3). The stationary distribution is normal with mean θ
455 and variance $\sigma^2/(2\alpha)$, where the values of θ and σ vary across regimes. The x-axis is
456 natural-log transformed genome size.

462 **Discussion**
463

Genome Size Evolution Across Different Life Histories

Genome size differs among metamorphic life history strategies in salamanders. Sala-

manders that undergo abrupt metamorphosis are experiencing deterministic pull down-

ward, suggesting that abrupt metamorphosis is driving the evolution of smaller genome

size (smaller equilibrium value than observed, Figure 3a). Gradual metamorphosers and

direct developers reach larger genome sizes than abrupt metamorphosers, but we see

little evidence from the best-fit model of any tendency towards either genome size con-

traction or expansion (Fig. 3b,c). When metamorphic remodeling is largely removed

from life history, genomes are expected to be unconstrained and permissive to TE accu-

mulation, consistent with our finding that paedomorphic lineages are experiencing deter-

ministic pull toward genome expansion (Fig. 3d). Although metamorphosis has been

suggested to influence the evolution of genome size, this hypothesis has rarely been

demonstrated in a broadly comparative manner, and the relevant features of metamor-

phosing organisms that exert evolutionary pressure on genome size have been under-

studied (but see Gregory 2002b; Bonett, et al. 2020). We discuss how three classes of

metamorphic vulnerabilities or their interaction may constrain biased TE accumulation

and, thus, shape genome size diversity across life history regimes in salamanders: (1)

performance and physiological handicap, (2) energetic limitation, and (3) developmental

error during metamorphosis. We track the influence of these evolutionary pressures in

qualitative form in Table 4.

Performance and Physiological Handicap

486
Performance handicaps are expected based on the classic and widely-cited work

on frogs; metamorphosing individuals can neither hop nor swim effectively and are sub-

ject to increased predation (Wassersug and Sperry 1977; Arnold and Wassersug 1978).

In contrast, metamorphosing salamanders do not suffer locomotor handicaps (Landberg

and Azizi 2010). However, some metamorphosing salamanders (*Gyrinophilus porphyrit-*

icus) do suffer a performance handicap in accessing shelter during critical environmen-

tal fluctuations; their decreased ability to exploit stream habitat refugia during extremely

high or low flow rates increases mortality (Lowe, et al. 2019). In addition, several spe-

cies of salamanders have decreased critical thermal maxima during metamorphosis

(*Ambystoma tigrinum*, *Notophthalmus viridescens*) (Hutchison 1961; Delson and Whit-

ford 1973). Thus, decreased performance and physiological tolerance likely contribute

to the increased mortality experienced by salamanders during metamorphosis (Peter-

son, et al. 1991), selecting for rapid metamorphosis and constraining genomes and cells

to smaller sizes. However, these vulnerabilities apply to all metamorphosers, accounting

for their smaller genome sizes relative to direct developers and paedomorphs, but not

for the differences between abrupt and gradual metamorphosers (Table 4).

Energetic Limitation

Vulnerability of metamorphosing individuals to energetic limitation is important

despite low overall metabolic rates across salamanders (Gatten, et al. 1992), and fur-

thermore affects the metamorphic strategies differently. The most strongly affected are

abrupt, synchronous metamorphosers. There is evidence that these larvae are food-lim-

ited in nature; individuals grow more slowly in the field than those fed *ad libitum* in the

lab (Beachy 1995). In addition, the timing of the onset of abrupt metamorphosis is fixed,

irrespective of food availability, larval growth rate, or body size (Beachy, et al. 2017;

Beachy 2018). Thus, these animals lack the flexibility to delay metamorphosis until they

reach an optimal size — unlike metamorphosing frogs, arthropods, and salamanders

that undergo gradual metamorphosis — resulting in the possibility that transformation is

forced at a smaller body size with lower energy reserves, yielding reduced fitness (Wilbur and Collins 1973; Beachy, et al. 2017). Once abrupt metamorphosis begins, feeding ceases, although there are mixed anecdotal reports of transforming organisms feeding or attempting to feed in the lab (Deban and Marks 2002). Metamorphosis itself does not increase energetic demand, as evidenced by gradually metamorphosing salamanders, in which metabolic rate remains unchanged during the transformation (Vladimirova, et al. 2012). However, the transformation is more extensive in abrupt metamorphosing salamanders, and frogs experience elevated metabolic rates during their extensive metamorphosis (Orlofske and Hopkins 2009; Wright, et al. 2011). Thus, abrupt metamorphosers are at risk of initiating a costly sequence of developmental events with sub-optimal energy reserves and impaired capacity to feed (Deban and Marks 2002; Vladimirova, et al. 2012; Beachy, et al. 2017). In contrast, gradual metamorphosers are not affected energetically, as they can feed throughout metamorphosis as well as delay its onset. In direct-developing lineages, some or all of the developmental steps of metamorphic remodeling occur inside the egg at the end of embryogenesis, in sequences that deviate from those of metamorphosers to varying degrees (Alberch 1989; Wake and Hanken 1996; Marks 2000; Rose 2003; Kerney, et al. 2012). The energy to fuel metamorphic repatterning comes from yolk stores which, although relatively large in direct developers, are still finite (Wake and Hanken 1996; Gregory 2002b), suggesting the potential for some energetic vulnerability. However, direct developers do not have smaller genome size equilibria than gradual metamorphosers, suggesting that any constraint imposed by energetic limitation in direct developers is weak. Abrupt metamorphosers have the smallest genome size equilibrium, consistent with energetic limitation

exerting the strongest selection for rapid metamorphosis and therefore the strongest

constraint on genome size (Table 4).

Developmental Error

Abrupt and gradual metamorphosers differ in both metamorphic synchronicity

and extent of transformation, leading to differential vulnerability to developmental errors.

Abrupt metamorphosis entails the synchronous execution of a large number of develop-

mental events, e.g.: the chondrification and ossification of the nasal capsule, dermal

bones of the skull, upper jaw, and palate; remodeling of the skin; and gill resorption. The

internal organs are likely also abruptly remodeled because they are under the same

hormonal control (Rose 1996; Rose 1999, 2003). Some systems undergo a more radi-

cal transformation altogether; larval elements undergo cell death, and adult structures

form *de novo* rather than being remodeled from larval elements. For example, the hy-

obranchial apparatus (i.e. tongue skeleton) in abrupt metamorphosers transforms this

way, enabling the formation of projectile and, in some species, ballistic tongues

(Alberch, et al. 1985; Alberch and Gale 1986). These transformations require precise

spatial coordination of the cell-autonomous programs and cell-cell interactions that drive

cell death, growth, and differentiation, often in very close proximity (Alberch, et al. 1985;

Schreiber, et al. 2009; Ishizuya-Oka 2011). Temporal organization is also critical; induc-

tion events (e.g. epithelial-mesenchymal interactions) that occur out of sequence result

in developmental anomalies (Hanken and Hall 1993). The evolution of abrupt metamor-

phosis increased the complexity of spatiotemporal coordination of cell-cell interactions

relative to gradual metamorphosers. We found that abrupt metamorphosers have dis-

tinctly smaller genome sizes, consistent with more intense vulnerability to developmen-

tal error (Table 4). This result is broadly consistent with a similar pattern in insects,

where holometabolous lineages (i.e. those that undergo a complete metamorphosis)

have smaller genome sizes than hemimetabolous lineages (i.e. those that undergo in-

complete metamorphosis), which has been interpreted as evidence for stronger con-

straint on genome size with greater developmental complexity (Gregory 2002b; Alfsnes,

et al. 2017).

Direct developers undergo repatterning at a developmental stage involving much

less tissue and fewer cells, therefore requiring less spatiotemporal cell-cell coordination.

Although this simplifies the overall developmental process, it also introduces the poten-

tial for error because stochastic noise in developmental processes (e.g. cell migration)

can have larger phenotypic effects when each cell represents a larger proportion of an

incipient structure. One of the best illustrations of the mechanistic consequences of low

cell numbers on development is the salamander genus *Thorius,* characterized by large

genomes, small bodies, and thus low cell numbers. Skeletal and cartilaginous elements

in the limbs and skull form embryonically from precartilage condensations, which are

tight aggregates of mesenchymal cells. Up to 70% of *Thorius* individuals show left-right

asymmetry in the arrangement of carpal or tarsal elements in the limbs and/or anterior

elements in the skull, and similar variation exists among individuals (Hanken 1982;

Hanken 1984). This degree of variability demonstrates that the outcome of precartilage

condensation is subject to stochastic noise in the cellular processes involved in cell ag-

gregation (i.e. cell movement, cell-cell adhesion, cell-extracellular matrix interaction)

(Chatterjee, et al. 2020; Glimm, et al. 2020).

More generally, vulnerability to stochastic noise is expected to impose stronger

constraints on genome expansion when developmental sequences are more complex.

Limited data from direct developers support this hypothesis. For example, in the genus

Bolitoglossa, the formation of larval hyobranchial apparatus components has been lost

from ontogeny, leading to a simpler developmental sequence in which adult structures

are formed without larval precursors (Alberch 1989; Wake and Hanken 1996). Other

taxa (e.g. *Desmognathus* and the eastern *Plethodon*) retain a more complex develop-

mental sequence with more of the larval and metamorphorphic stages of hyobranchial

development (Wake and Hanken 1996; Marks 2000; Kerney, et al. 2012); as predicted,

they have smaller genome sizes (~15 Gb and ~25 Gb for *Desmognathus* and the east-

ern *Plethodon*, respectively, versus ~50 Gb for *Bolitoglossa*).

Both direct developers and abrupt metamorphosers experience risks for develop-

mental error, but direct developers have a larger genome size equilibrium (Table 3),

supporting weaker developmental error constraints (Table 4). In addition, direct devel-

opers have the largest expected variance in genome size, which presents a new hy-

pothesis — that diverse developmental sequences within this life history regime also dif-

fer in vulnerability to error and, thus, constraint on genome expansion.

Paedomorphosis and the Loss of Metamorphic Constraints

Despite their freedom from metamorphosis-induced constraints, paedomorphs

may still face other evolutionary pressures that prevent their genomes from reaching

ever-larger sizes. The decreased surface-area-to-volume ratio that accompanies in-

creased cell size imposes a functional limit that salamanders may well have reached

(Chan and Marshall 2010), as their cells are among the largest in animals (Horner and

Macgregor 1983). In addition, the ability of adult salamanders to regenerate limbs and

organs declines dramatically in paedomorphs with the largest genomes, suggesting an-

other fitness consequence of extreme genome expansion (Scadding 1977; Sessions

and Wake 2021).

In the past, huge cells have been proposed as adaptive because salamanders

and lungfishes have the lowest metabolic rates and the largest genomes/cells among

vertebrates. This correlation led to the "frugal metabolic strategy" hypothesis (Szarski

1983; Olmo, et al. 1989), under which paedomorphs would show the greatest degree of

adaptation for this trait. However, more recent studies failed to find a clear relationship

between genome or cell size and metabolic rate, both within salamanders (Licht and

Lowcock 1991) and more generally (Uyeda, et al. 2017; Gardner, et al. 2020). Rather

than adaptation towards larger genome size, our results support the idea that loss of

metamorphosis has released constraints against genome expansion in paedomorphs.

Table 4. Mutation pressure and metamorphic vulnerabilities shaping genome size in different life history regimes

Evolutionary Pressure	Effect on Genome Size	Abrupt Metamor­phosers	Gradual Metamor­phosers	Direct Developers	Paedomorphs
TE accumulation bias	↑	✓[a]	✓	✓	✓
Performance or Physiologi­cal Handicap	↓	✓	✓		
Energetic Limitation	↓	✓✓		✓	
Developmen­tal Error	↓	✓✓	✓	✓	

625 [a]A single checkmark vs. two checkmarks indicates a difference in the intensity of a met-

626 amorphic vulnerability across life history regimes.

627
628

Summary of Metamorphic Vulnerabilities and Genome Size Evolution

Abrupt metamorphosers are challenged with every metamorphic vulnerability: de-

creased performance and physiological tolerance, energetic limitation, and developmen-

tal error, providing the explanation for their smallest genome sizes (Table 4). In contrast,

paedomorphs are free from all of these vulnerabilities and have the largest genome

sizes (Table 4). Direct developers have a similar genome size equilibrium to gradual

metamorphosers, but with larger variation (Table 3), which we propose reflects diversifi-

cation of direct developmental pathways. Gradual metamorphosers avoid developmen-

tal error at the cost of decreased performance and physiological tolerance whereas di-

rect developers avoid performance handicap at the cost of some susceptibility to sto-

chastic noise in development. More generally, these results illustrate how genome size

may be shaped by different constraints across taxa that are important to assess in order

to understand the major drivers of genome size evolution across the tree of life (Knight,

et al. 2005; Carta and Peruzzi 2016; Alfsnes, et al. 2017; Roddy, et al. 2019).

OU Models Evaluate the Balance of Evolutionary Forces

While we have no model tailored to biased evolutionary increase opposed by

constraining forces, the OU class of models (of which BM is a subset) is a powerful tool

that can differentiate among biologically-motivated, nuanced hypotheses. For example,

because TE accumulation is stochastic, one might expect the BM model to be a good

candidate for purely stochastic genome size evolution. However, this is not the case for

salamanders because there is bias for TE gains over losses, rendering BM a poor fit.

While it is widely recognized that OU models provide deterministic components that al-

low modeling evolutionary shifts in the means of phenotypes, it is less appreciated that

they also provide a way to model the information content in the variances. The differ-

ence between a purely stochastic (BM) model and one that has any degree of determin-

istic pull (OU) is that the variance of a BM model will grow unbounded over time,

whereas the variance in a model with deterministic pull will not (Hansen and Martins

1996; Butler and King 2004). Along these lines, our finding that allowing both stochastic

noise intensities and equilibria to vary across regimes produces a model with substan-

tially better fit than either set of parameters alone is very biologically informative. While

we have three broad categories of genome size equilibria, with abrupt metamorphosers

distinctly smaller and paedomorphs larger (Table 3), the metamorphic categories also

differ substantially in stochastic noise intensity. The smaller equilibrium value and noise

intensity of abrupt metamorphosers suggests that this regime is highly constrained with

deterministic pull downward, even in comparison to gradual metamorphosers (Table 3,

Fig. 3a vs. b). Direct developers have the highest estimated noise intensity, suggesting

that the evolution of diverse developmental sequences resulted in a range of vulnerabili-

ties to error and, thus, variable constraints on genome expansion (compare confidence

intervals in Table 3, Fig. 3c vs. d). Thus, both the mean and variance of the evolutionary

process shed light on the balance of forces acting on genome size across life histories.

670   In general, OU models are conceptualized as models for stabilizing selection,

with the θ parameters interpreted as trait optima towards which individual species are

evolving. However, this interpretation does not fit well with the biology of genome size.

As the variability in genome size is strongly determined by the quantity of nearly-neutral

noncoding DNA, the notion of an "optimal" genome size has little meaning (but see

Cavalier-Smith 2005). More realistically, there is a range of permissible variation within

which species can vary.

677   We propose that the strong support for OU models in our analysis reflects a bal-

ance between the biased stochastic forces driving genome size upward and evolution-

ary constraint acting to limit genome size. That is, rather than interpreting Fig. 3b as evi-

dence that gradual metamorphosing salamanders have genomes near their "optimal"

sizes, we interpret it as showing that these salamanders have settled on a genome size

distribution that is balanced between stochastic TE dynamics tending to bias genome

size upward and selection on metamorphosis imposing evolutionary constraint against

further size increase. Paedomorphs have the largest genome size equilibria of all, and

our model indicates that they are evolving with a deterministic pull toward even larger

size. There is no compelling adaptive interpretation for this genome size "optimum"

larger than all other vertebrates, although some have tried (see above); rather, we inter-

pret it as a balance between upwardly-biased TE accumulation and functional con-

straints not considered here (e.g. an upper limit on cell size). This is a novel way of in-

terpreting the results of comparative analysis, but one that is supported by our under-

standing of the biology of the system. Our results thus suggest that OU models can po-

tentially be used to detect other evolutionary processes beyond adaptation towards an

optimum, which broadens their applicability to the study of traits that do not evolve in re-

sponse to strong selection by a single factor.

698   Literature Cited
699

Alberch P. 1989. Development and the evolution of amphibian metamorphosis. In:

Splechtna/Hilgers, editor. Trends in Vertebrate Morphology. Stuttgart: Gustav Fischer

Verlag. p. 163-173.

Alberch P, Gale EA. 1986. Pathways of cytodifferentiation during the metamorphosis of

the epibranchial cartilage in the salamander *Eurycea bislineata*. Dev Biol 117:233-244.

Alberch P, Lewbart G, Gale EA. 1985. The fate of larval chondrocytes during the

metamorphosis of the epibranchial in the salamander, *Eurycea bislineata*. J Embryol

Exp Morph 88:71-83.

Alfsnes K, Leinaas HP, Hessen DO. 2017. Genome size in arthropods; different roles of

phylogeny, habitat and life history in insects and crustaceans. Ecol Evol 7:5939-5947.

AmphibiaWeb: Information on amphibian biology and conservation [Internet]. 2022.

Berkeley, California. Available from http://amphibiaweb.org/.

Arkhipova IR. 2018. Neutral theory, transposable elements, and eukaryotic genome

evolution. Mol Biol Evol 35:1332-1337.

Arnold SJ. 1992. Constraints on phenotypic evolution. Am Nat 140:S85-S107.

Arnold SJ, Wassersug RJ. 1978. Differential predation on metamorphic anurans by

garter snakes (*Thamnophis*): Social behavior as a possible defense. Ecology 59:1014-

1022.

Bachmann K. 1970. Feulgen slope determinations of urodele nuclear DNA amounts.

Histochemie 22:289-293.

Beachy CK. 2018. Effects of growth rate and temperature on metamorphosis in *Eurycea*

wilderae (Caudata, Plethodontidae, Hemidactyliinae, Spelerpini; blue ridge two-lined

salamander). Southeast Nat 17:423-432.

Beachy CK. 1995. Effects of larval growth history on metamorphosis in a stream-

dwelling salamander (*Desmognathus ochrophaeus*). J Herp:375-382.

Beachy CK, Ryan TJ, Bonett RM. 2017. How metamorphosis is different in
plethodontids: Larval life history perspectives on life-cycle evolution. Herpetologica
73:252-258.

Beaulieu JM, Jhwueng D-C, Boettiger C, O'Meara BC. 2012. Modeling stabilizing
selection: Expanding the Ornstein–Uhlenbeck model of adaptive evolution. Evolution
66:2369-2383.

Boettiger C, Coop G, Ralph P. 2012. Is your phylogeny informative? Measuring the
power of comparative methods. Evolution 66:2240-2251.

Bonett RM, Hess AJ, Ledbetter NM. 2020. Facultative transitions have trouble
committing, but stable life cycles predict salamander genome size evolution. Evol Biol
47:111-122.

Bonett RM, Steffen MA, Robison GA. 2014. Heterochrony repolarized: a phylogenetic
analysis of developmental timing in plethodontid salamanders. Evo Devo 5:27.

Butler Marguerite A, King Aaron A. 2004. Phylogenetic comparative analysis: A
modeling approach for adaptive evolution. Am Nat 164:683-695.

Carta A, Peruzzi L. 2016. Testing the large genome constraint hypothesis: plant traits,
habitat and climate seasonality in Liliaceae. New Phytol. 210:709-716.

Cavalier-Smith T. 2005. Economy, speed and size matter: evolutionary forces driving
nuclear genome miniaturization and expansion. Ann Bot 95:147-175.

Chan Y-HM, Marshall WF. 2010. Scaling properties of cell and organelle size.
Organogenesis 6:88-96.

Chatterjee P, Glimm T, Kaźmierczak B. 2020. Mathematical modeling of chondrogenic

pattern formation during limb development: Recent advances in continuous models.

Math Biosci 322:108319.

Chippindale PT, Bonett RM, Baldwin AS, Wiens JJ. 2004. Phylogenetic evidence for a

major reversal of life-history evolution in plethodontid salamanders. Evolution 58:2809-

2822.

Cressler CE, Butler MA, King AA. 2015. Detecting adaptive evolution in phylogenetic

comparative analysis using the Ornstein–Uhlenbeck model. Syst Biol 64:953-968.

Deban SM, Marks SB. 2002. Metamorphosis and evolution of feeding behaviour in

salamanders of the family Plethodontidae. Zool J Linn Soc 134:375-400.

Decena-Segarra LP, Bizjak-Mali L, Kladnik A, Sessions SK, Rovito SM. 2020.

Miniaturization, genome size, and biological size in a diverse clade of salamanders. Am

Nat 196:634-648.

Delson J, Whitford WG. 1973. Critical thermal maxima in several life history stages in

desert and montane populations of *Ambystoma tigrinum*. Herpetologica:352-355.

Enriquez-Urzelai U, Sacco M, Palacio AS, Pintanel P, Tejedo M, Nicieza AG. 2019.

Ontogenetic reduction in thermal tolerance is not alleviated by earlier developmental

acclimation in *Rana temporaria*. Oecologia 189:385-394.

Felsenstein J. 1985. Phylogenies and the comparative method. Am Nat 125:1-15.

Frahry MB, Sun C, Chong R, Mueller RL. 2015. Low levels of LTR retrotransposon

deletion by ectopic recombination in the gigantic genomes of salamanders. J Mol Evol

80:120-129.

Gardner JD, Laurin M, Organ CL. 2020. The relationship between genome size and
metabolic rate in extant vertebrates. Philos Trans R Soc B 375:20190146.

Gatten REJ, Miller K, Full RJ. 1992. Energetics at rest and during locomotion. In: Feder
ME, editor. Environmental Physiology of the Amphibians. Chicago: University of
Chicago Press. p. 314-377.

Glimm T, Bhat R, Newman SA. 2020. Multiscale modeling of vertebrate limb
development. Wiley Interdiscip Rev: Syst Biol Med 12:e1485.

Gould SJ. 1977. Ontogeny and Phylogeny. Cambridge, MA: Harvard University Press.

Gregory TR. 2002a. A bird's eye view of the c-value enigma: genome size, cell size, and
metabolic rate in the class Aves. Evolution 56:121-130.

Gregory TR. 2001. Coincidence, coevolution, or causation? DNA content, cell size, and
the C-value enigma. Biol Rev Camb Philos Soc 76:65-101.

Gregory TR. 2005. The Evolution of the Genome. San Diego, CA: Academic Press.

Gregory TR. 2002b. Genome size and developmental complexity. Genetica 115:131-
146.

Gregory, T. R. Animal Genome Size Database (http://www.genomesize.com) [Internet].
2022.

Gregory TR. 2004. Macroevolution, hierarchy theory, and the C-value enigma.
Paleobiology 30:179-202.

Hanken J. 1982. Appendicular skeletal morphology in minute salamanders, genus

Thorius (Amphibia: Plethodontidae): growth regulation, adult size determination, and

natural variation. J Morph 174:57-77.

Hanken J. 1984. Miniaturization and its effects on cranial morphology in plethodontid

salamanders, genus Thorius (Amphibia: Plethodontidae). I. Osteological variation. Biol J

Linn Soc 23:55-75.

Hanken J, Hall BK. 1993. Mechanisms of skull diversity and evolution. In: Hanken J,

Hall BK, editors. The Skull, Volume 3: Functional and Evolutionary Mechanisms.

London: University of Chicago Press.

Hansen TF. 1997. Stabilizing selection and the comparative analysis of adaptation.

Evolution 51:1341-1351.

Hansen TF, Martins EP. 1996. Translating between microevolutionary process and

macroevolutionary patterns: The correlation structure of interspecific data. Evolution

50:1404-1417.

Hessen DO, Daufresne M, Leinaas HP. 2013. Temperature-size relations from the

cellular-genomic perspective. Biol Rev 88:476-489.

Ho LST, Ané C. 2013. Asymptotic theory with hierarchical autocorrelation: Ornstein-

Uhlenbeck tree models. Ann Statist 41:957-981.

Ho LST, Ané C. 2014. Intrinsic inference difficulties for trait evolution with Ornstein-

Uhlenbeck models. Methods Ecol Evol 5:1133-1146.

Horner HA, Macgregor HC. 1983. C value and cell volume: their significance in the

evolution and development of amphibians. J Cell Sci 63:135.

Hutchison VH. 1961. Critical thermal maxima in salamanders. Physiol Zool 34:92-125.

Ishizuya-Oka A. 2011. Amphibian organ remodeling during metamorphosis: Insight into
thyroid hormone-induced apoptosis. Develop Growth Diff 53:202-212.

Jeffery NW, Yampolsky L, Gregory TR. 2016. Nuclear DNA content correlates with
depth, body size, and diversification rate in amphipod crustaceans from ancient Lake
Baikal, Russia. Genome 60:303-309.

Jetz W, Pyron RA. 2018. The interplay of past diversification and evolutionary isolation
with present imperilment across the amphibian tree of life. Nature Ecol Evol 2:850-858.

Jockusch EL. 1997. An evolutionary correlate of genome size change in plethodontid
salamanders. Proc Royal Soc B: Biol Sci 264:597.

Kapusta A, Suh A, Feschotte C. 2017. Dynamics of genome size evolution in birds and
mammals. Proc Nat Acad Sci 114:E1460-E1469.

Keinath MC, Timoshevskiy VA, Timoshevskaya NY, Tsonis PA, Voss SR, Smith JJ.
2015. Initial characterization of the large genome of the salamander *Ambystoma*
mexicanum using shotgun and laser capture chromosome sequencing. Sci Rep
5:16413.

Kerney RR, Blackburn DC, Müller H, Hanken J. 2012. Do larval traits re-evolve?
Evidence from the embryogenesis of a direct-developing salamander, *Plethodon*
cinereus. Evolution 66:252-262.

Knight CA, Molinari NA, Petrov DA. 2005. The large genome constraint hypothesis:
evolution, ecology and phenotype. Ann Bot 95:177-190.

Landberg T, Azizi E. 2010. Ontogeny of escape swimming performance in the spotted

salamander. Funct Ecol 24:576-587.

Larson A. 1984. Neontological inferences of evolutionary pattern and process in the

salamander family Plethodontidae. Evol Biol 17:119-217.

Lertzman-Lepofsky G, Mooers AØ, Greenberg DA. 2019. Ecological constraints

associated with genome size across salamander lineages. Proc Roy Soc B

286:20191780.

Licht LE, Lowcock LA. 1991. Genome size and metabolic rate in salamanders. Comp

Biochem Physiol B: Comp Biochem 100:83-92.

Liedtke HC, Gower DJ, Wilkinson M, Gomez-Mestre I. 2018. Macroevolutionary shift in

the size of amphibian genomes and the role of life history and climate. Nat Ecol Evol

2:1792-1799.

Lowe WH, Martin TE, Skelly DK, Woods HA. 2021. Metamorphosis in an era of

increasing climate variability. Trends Ecol Evol.

Lowe WH, Swartz LK, Addis BR, Likens GE. 2019. Hydrologic variability contributes to

reduced survival through metamorphosis in a stream salamander. Proc Natl Acad Sci

USA 116:19563.

Madison-Villar MJ, Sun C, Lau N, Settles M, Mueller RL. 2016. Small RNAs from a big

genome: the piRNA pathway and transposable elements in the salamander species

Desmognathus fuscus. J Mol Evol 83:126-136.

Marks SB. 2000. Skull development in two plethodontid salamanders (genus

Desmognathus) with different life histories. In. The Biology Of Plethodontid

Salamanders: Springer. p. 261-276.

Michael TP. 2014. Plant genome size variation: bloating and purging DNA. Brief Funct

Genom 13:308-317.

Mueller RL, Macey JR, Jaekel M, Wake DB, Boore JL. 2004. Morphological homoplasy,

life history evolution, and historical biogeography of plethodontid salamanders inferred

from complete mitochondrial genomes. Proc Natl Acad Sci USA 101:13820-13825.

Nowoshilow S, Schloissnig S, Fei J-F, Dahl A, Pang AWC, Pippel M, Winkler S, Hastie

AR, Young G, Roscito JG, et al. 2018. The axolotl genome and the evolution of key

tissue formation regulators. Nature 554:50-55.

O'Meara BC, Ané C, Sanderson MJ, Wainwright PC. 2006. Testing for different rates of

continuous trait evolution using likelihood. Evolution 60:922-933.

Olmo E, Capriglione T, Odierna G. 1989. Genome size evolution in vertebrates: Trends

and constraints. Comp Biochem Phys B: Comp Biochem 92:447-453.

Orlofske SA, Hopkins WA. 2009. Energetics of metamorphic climax in the pickerel frog

(*Lithobates palustris*). Comp Biochem Phys A: Molec Integ Phys 154:191-196.

Pandit MK, White SM, Pocock MJO. 2014. The contrasting effects of genome size,

chromosome number and ploidy level on plant invasiveness: a global analysis. New

Phytol 203:697-703.

Peterson CL, Wilkinson RF, Moll D, Holder T. 1991. Premetamorphic survival of

Ambystoma annulatum. Herpetologica:96-100.

R Core Team. 2020. R: A language and environment for statistical computing. Vienna,
Austria: R Foundation for Statistical Computing.

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and
other things). Methods Ecol Evol 3:217-223.

Roddy AB, Théroux-Rancourt G, Abbo T, Benedetti JW, Brodersen CR, Castro M,
Castro S, Gilbride AB, Jensen B, Jiang G-F, et al. 2019. The scaling of genome size
and cell size limits maximum rates of photosynthesis with implications for ecological
strategies. Int J Plant Sci 181:75-87.

Rose CS. 2003. The developmental morphology of salamander skulls. In: Heatwole H,
editor. Amphibian Biology: Osteology. Chipping Norton: Surrey Beatty & Sons. p. 1684-
1781.

Rose CS. 1996. An endocrine-based model for developmental and morphogenetic
diversification in metamorphic and paedomorphic urodeles. J Zool 239:253-284.

Rose CS. 1999. Hormonal control in larval development and evolution—amphibians. In.
The Origin And Evolution Of Larval Forms: Elsevier. p. 167-VI.

Rose CS. 2014. The importance of cartilage to amphibian development and evolution.
Int J Dev Biol 58:917-927.

Savell KRR, Auerbach BM, Roseman CC. 2016. Constraint, natural selection, and the
evolution of human body form. Proc Nat Acad Sci 113:9492-9497.

Scadding SR. 1977. Phylogenic distribution of limb regeneration capacity in adult
Amphibia. J Exp Zool 202:57-67.

Schreiber AM, Mukhi S, Brown DD. 2009. Cell–cell interactions during remodeling of the
intestine at metamorphosis in *Xenopus laevis*. Dev Biol 331:89-98.

Semlitsch RD, Scott DE, Pechmann JH. 1988. Time and size at metamorphosis related
to adult fitness in *Ambystoma talpoideum*. Ecology 69:184-192.

Sessions SK. 2008. Evolutionary cytogenetics in salamanders. Chromosome Res
16:183-201.

Sessions SK, Wake DB. 2021. Forever young: Linking regeneration and genome size in
salamanders. Dev Dyn 250:768-778.

Shao F, Han M, Peng Z. 2019. Evolution and diversity of transposable elements in fish
genomes. Sci Rep 9:15399.

Smith JDL, Bickham JW, Gregory TR. 2013. Patterns of genome size diversity in bats
(order Chiroptera). Genome 56:457-472.

Sotero-Caio CG, Platt RN, II, Suh A, Ray DA. 2017. Evolution and diversity of
transposable elements in vertebrate genomes. Genome Biol Evol 9:161-177.

Sun C, Mueller RL. 2014. Hellbender genome sequences shed light on genome
expansion at the base of crown salamanders. Genome Biol Evol 6:1818-1829.

Sun C, Shepard DB, Chong RA, Arriaza JL, Hall K, Castoe TA, Feschotte C, Pollock
DD, Mueller RL. 2012. LTR retrotransposons contribute to genomic gigantism in
plethodontid salamanders. Genome Biol Evol 4:168-183.

Szarski H. 1983. Cell size and the concept of wasteful and frugal evolutionary
strategies. J Theor Biol 105:201-209.

Szarski H. 1957. The origin of larva and metamorphosis in amphibia. Am Nat 91:283-
301.

Uyeda JC, Pennell MW, Miller ET, Maia R, McClain CR. 2017. The evolution of
energetic scaling across the vertebrate tree of life. Am Nat 190:185-199.

Vinogradov AE. 2004. Genome size and extinction risk in vertebrates. Proc Roy Soc B
271:1701-1705.

Vladimirova IG, Kleimenov SY, Alekseeva TA. 2012. Dynamics of body mass and
oxygen consumption in the ontogeny of the Spanish ribbed newt (*Pleurodeles waltl*): 2.
Larval stage. Biol Bull 39:10-14.

Vu GTH, Cao HX, Reiss B, Schubert I. 2017. Deletion-bias in DNA double-strand break
repair differentially contributes to plant genome shrinkage. New Phytol 214:1712-1721.

Wake D, Hanken J. 1996. Direct development in the lungless salamanders: what are
the consequences for developmental biology, evolution and phylogenesis? Int J Dev
Biol 40:859-869.

Wake DB, Marks SB. 1993. Development and evolution of plethodontid salamanders: a
review of prior studies and a prospectus for future research. Herpetologica 49:194-203.

Wassersug RJ, Sperry DG. 1977. The relationships of locomotion to differential
predation on *Pseudacris triseriata* (anura: Hylidae). Ecology 58:830-839.

Wiens JJ, Bonett RM, Chippindale PT. 2005. Ontogeny discombobulates phylogeny:
paedomorphosis and higher-level salamander relationships. Syst Biol 54:91-110.

Wilbur HM, Collins JP. 1973. Ecological aspects of amphibian metamorphosis:

nonnormal distributions of competitive ability reflect selection for facultative

metamorphosis. Science 182:1305-1314.

Wright ML, Richardson SE, Bigos JM. 2011. The fat body of bullfrog (*Lithobates*

catesbeianus) tadpoles during metamorphosis: Changes in mass, histology, and

melatonin content and effect of food deprivation. Comp Biochem Phys A 160:498-503.

Wright NA, Gregory TR, Witt CC. 2014. Metabolic 'engines' of flight drive genome size

reduction in birds. Proc Roy Soc B: Biol Sci 281:20132780.

942