An optimized protocol for assay for transposase-accessible chromatin by sequencing

(ATAC-seq) library preparation from adult *Drosophila melanogaster* neurons

Collin B. Merrill[1,*], Miguel A. Pabon[2], Austin B. Montgomery[2], Aylin R. Rodan[2,3,4,5], and

Adrian Rothenfluh[1,2,4,6,*]

[1]Huntsman Mental Health Institute, Department of Psychiatry, University of Utah, Salt

Lake City, UT 84108, USA.

[2]Molecular Medicine Program, University of Utah, Salt Lake City, UT 84112, USA.

[3]Division of Nephrology and Hypertension, Department of Internal Medicine, University

of Utah, Salt Lake City, UT 84112, USA.

[4]Department of Human Genetics, University of Utah, Salt Lake City, UT 84112, USA.

[5]Medical Service, Veterans Affairs Salt Lake City Health Care System, Salt Lake City,

UT 84148, USA

[6]Department of Neurobiology, University of Utah, Salt Lake City, UT 84112, USA.

*Corresponding authors:

Collin Merrill, collin.merrill@utah.edu

Adrian Rothenfluh, adrian.rothenfluh@hsc.utah.edu

ABSTRACT (200 words)

Assay for transposase-accessible chromatin by sequencing (ATAC-seq) is rapidly

becoming the assay of choice to investigate chromatin-mediated gene regulation,

largely because of low input requirements, a fast workflow, and the ability to interrogate

the entire genome in an unbiased manner. Many studies using ATAC-seq use

mammalian or human-derived tissues, and established protocols work well in these

systems. However, ATAC-seq is not yet widely used in *Drosophila*. Vinegar flies present

several advantages over mammalian systems that make them an excellent model for

ATAC-seq studies, including abundant genetic tools that allow straightforward targeting,

transgene expression, and genetic manipulation, and are not available in mammalian

models. Because current ATAC-seq protocols are not optimized to use flies, we

developed an optimized workflow that accounts for several complicating factors present

in *Drosophila*. We examined parameters affecting nuclei isolation, including input size,

freezing time, washing, and possible confounds from retinal pigments. Then, we

optimized the enzymatic steps of library construction to account for the smaller

Drosophila genome size. Finally, we used our optimized protocol to generate ATAC-seq

libraries that meet ENCODE quality metrics. Our optimized protocol enables extensive

ATAC-seq experiments in *Drosophila*, thereby leveraging the advantages of this

powerful model system to understand chromatin-mediated gene regulation.

41
42   Keywords: Drosophila, neurons, Tn5, tagmentation, ATAC-seq
43

INTRODUCTION

Assay for transposase-accessible chromatin by sequencing (ATAC-seq) is a relatively new technique that has advanced our understanding of chromatin-mediated gene regulation. The technique offers several advantages over other sequencing techniques that examine chromatin, such as micrococcal nuclease (MNase-seq) or formaldehyde-assisted isolation of regulatory elements (FAIRE-seq). These advantages include low input requirements, a fast workflow that does not require harsh chemicals, and genome-wide interrogation in an untargeted manner (Buenrostro et al., 2013). ATAC-seq is also robust to small sample sizes and has been adapted for single-cell applications (Cusanovich et al., 2015; Buenrostro et al., 2018).

The established ATAC-seq protocol was developed using peripheral blood samples (Buenrostro et al., 2013), and many studies using ATAC-seq focus on mammalian or human-derived tissues (Davie et al., 2015; Bysani et al., 2019; Liu et al., 2019). Recent studies used ATAC-seq to investigate other tissue types, including tissues from plants (Lu et al., 2017), zebrafish (Quillien et al., 2017), and livestock (Halstead et al., 2020). Some ATAC-seq studies using *Drosophila melanogaster* have been published, though many of these studies focus on embryogenesis and larvae (Cusanovich et al., 2018; Haines and Eisen, 2018; Bozek et al., 2019). One recent report performed ATAC-seq from central brains after fluorescent sorting of GFP-positive cell bodies (Brovkina et al., 2021). However, the physical dissection and dissociation of many brains can be labor-intensive and yield relatively few cells, in that case 6,000-10,000 cells (Brovkina et al., 2021). We were interested in developing a streamlined protocol that would allow

efficient ATAC-seq library preparation, while taking advantage of the many genetic tools

available for labeling specific *Drosophila* neurons (Jenett et al., 2012).

ATAC-seq library preparation has several parameters that can be optimized based on

the input tissue. Many of these parameters are addressed in the original ATAC-seq

protocol (Buenrostro et al., 2013). Additional studies optimized ATAC-seq library

preparation from various tissue types in several storage conditions (Corces et al., 2017;

Ho et al., 2018; Wang et al., 2018; Fujiwara et al., 2019). Most studies in *Drosophila*

larvae treat the tissue similarly to mammalian tissues, only adding an additional washing

step to clean the embryos or larvae before homogenization (Koenecke et al., 2016;

Haines and Eisen, 2018). ATAC-seq library preparation using adult *Drosophila* poses

challenges that are not present in other tissues. For example, the insect cuticle is a

chitin structure that is difficult to process by simple homogenization. Another challenge

is eye color, which is commonly used to identify the presence of transgenic insertions,

and many transgenic fly lines contain the *mini-white* (w^{+mC}) gene. In a w^- background,

mini-white confers a yellow to red eye color, so eye color can range from white (w^-) to

red, depending on the genotype. Some *Drosophila* retinal pigments emit fluorescence

(Franceschini et al., 1981; Miller et al., 1984) that can complicate fluorescent-activated

nuclei sorting (FANS). Finally, the *Drosophila* genome is considerably smaller than

mammalian genomes (Vinogradov, 2004; Canapa et al., 2016), so established ATAC-

seq parameters may use higher-than-optimal enzyme concentrations and long

incubation times that may lead to low-quality ATAC-seq libraries when prepared from fly

tissues. Thus, optimizing each parameter for ATAC-seq library preparation using adult

flies should be performed prior to sequencing. However, these experiments add

considerable sequencing costs, putting the approach out-of-reach for many labs.

Identifying cell- and tissue-specific roles in organism function is a major goal of many

current studies, which require specific and sensitive detection of target cells, especially

for rare cell types. To facilitate these studies, we developed a protocol to generate

ATAC-seq libraries from adult *Drosophila* neurons. Our goal was to optimize each

variable in the library preparation process, with particular attention to challenges specific

to *Drosophila* model systems. Our protocol provides guidelines for ATAC-seq library

preparation using *Drosophila* and outlines the optimal parameters for each library

preparation step. This protocol will accelerate studies of chromatin-mediated gene

regulation in flies, thereby coupling the powerful genetic tools in *Drosophila* with high-

throughput, in-depth examination of chromatin structure.

MATERIALS AND METHODS

Fly strains

The following fly lines were obtained from the Bloomington Drosophila Stock Center

(Indiana University, Bloomington, IN, USA) and used in this study: *Canton S* (BL64349),

tub-Gal4 (BL5138), *elav-Gal4* (BL458), *nSyb-Gal4* (BL51635) *UAS-GFP-nls* (nuclear

localization signal; BL4775), *UAS-Stinger* (a super-bright GFP-nls variant; BL84277),

TH-Gal4 (BL8848), *Cha-Gal4.19B* (BL6798), and *vGAT-Gal4* (BL58980). *w* Berlin* flies

have been in the lab for many generations (Wolf et al., 2002) and were originally a gift from

Martin Heisenberg. Flies were reared in bottles containing standard cornmeal agar and

grown at 25 °C with 70% relative humidity and a 12-h light/dark cycle.

Nuclei isolation

All equipment and buffers were pre-chilled to be ice-cold. All plastic and glassware used

for nuclei isolation was pre-treated with 1% bovine serum albumin (BSA) in phosphate-

buffered saline (PBS). Flies were collected into empty bottles and frozen at -80 °C.

Then, the flies were placed in pre-chilled (in a -80 °C freezer) sieves and agitated for 1

min to separate the heads from the bodies. The fly heads were added to lysis buffer [10

mM Tris-HCl pH 7.5, 10 mM NaCl, 3 mM $MgCl_2$, 0.1% Tween-20, 0.1% Nonidet P40

substitute, 0.01% digitonin (ThermoFisher Scientific, Waltham, MA, USA), and 1%

bovine serum albumin] in a Dounce homogenizer. The tissue was homogenized with the

A pestle until the resistance disappeared. The crude homogenate was passed through a

40 µm filter and homogenized with 15 slow strokes with the B pestle. The crude nuclei

were washed with wash buffer (10 mM Tris-HCl, 10 mM NaCl, 3 mM $MgCl_2$, 1% BSA,

and 0.1% Tween-20) and centrifuged at 500 *g* for 10 minutes per wash. The washed

nuclei were resuspended in 1 mL wash buffer with 3 mM DAPI for evaluation and

sorting.

Fluorescence-activated nuclei sorting (FANS)

Nuclei were evaluated and sorted with a BD FACS Aria flow cytometer (BD

Biosciences, San Jose, CA, USA) operated by the University of Utah Flow Cytometry

Core facility. Nuclei collected from *w* Berlin* flies (no GFP) were stained with 3 µM DAPI

and used as the GFP-negative control to set the sorting gates. GFP-positive nuclei were

collected into 500 µL wash buffer and stored on ice until use for ATAC-seq library prep.

Nuclei counts were determined using an internal control bead population (Spherotech,

Lake Forest, IL, USA) at 10^6 beads/mL. Briefly, 20 uL of well-mixed beads were added

to 200 uL of nuclei suspension and gently mixed. Data was recorded for the bead and

cell mixture until at least 10,000 singlet beads were collected. Nuclei were identified

based on the forward scatter signal and DAPI intensity. The number of nuclei was

calculated as follows: number of nuclei recorded × 10^5/number of beads recorded =

number of nuclei/mL.

DNA tagmentation, amplification, and purification

Purified nuclei were centrifuged for 10 min at 500 *g* at 4 °C. The pellet was mixed with

22.5 µL 2X Tn5 tagmentation buffer [20 mM Tris-HCl pH 7.6, 10 mM $MgCl_2$, and 20%

dimethyl formamide (Sigma Aldrich, St. Louis, MO, USA) in sterile water], 16.5 µL 1X

PBS, 0.5 µL Tween-20, 5.5 µL water, and Tn5 enzyme (Illumina, Inc., San Diego, CA,

USA). The mixture was incubated at 37 °C for various times and the DNA was purified

using a MinElute PCR purification kit (Qiagen, Germantown, MD, USA). Then, the

purified DNA was mixed with CD index primers (Illumina) and Phusion High Fidelity

PCR MasterMix (New England Biolabs, Ipswich, MA, USA) and amplified by 1 cycle of

72 °C for 5 min and 98 °C for 30 sec, followed by 5 cycles of 98 °C for 10 sec, 63 °C for

30 sec, and 72°C for 1 min in a C100 thermocycler (BioRad, Hercules, CA, USA). 5 µL

of this reaction was removed and used for a qPCR side reaction. The aliquot was mixed

with the same CD index primers and SsoFast EvaGreen Supermix (BioRad) and

amplified for 1 cycle of 98 °C for 30 sec and 40 cycles of 98 °C for 10 sec, 63 °C for 30

sec, and 72 °C for 1 min in an Applied Biosystems 7900HT qPCR instrument

(ThermoFisher Scientific). Total fluorescence was calculated by subtracting the

fluorescence from first five PCR cycles from the maximum fluorescence of each sample.

The cycle number corresponding to a portion of the total fluorescence was determined

(~8-10 cycles) and the DNA was amplified for this number of additional cycles. After

amplification, the DNA was purified with 0.5X and 1.1X AMPure XP beads (Beckman

Coulter, Indianapolis, IN, USA) to remove primer-dimers and high-molecular weight

DNA (> 5000 bp). The purified DNA was stored at -20 °C until further analysis.

ATAC-seq library quality assessment

ATAC-seq library quality was assessed using an Agilent 2200 TapeStation and High

Sensitivity D5000 ScreenTape assays (Agilent Technologies, Inc., Santa Clara, CA,

USA) at the Huntsman Cancer Institute High Throughput Genomics core facility or the

Genome Technology Access Center of the McDonnell Genome Institute at the

Washington University School of Medicine (St. Louis, MO, USA).

Next generation sequencing and analysis

ATAC-seq libraries constructed using nuclei isolated from dopaminergic and GABAergic

neurons were sequenced on a Novaseq 6000 instrument (Illumina) using 50-bp paired-

end reads. Fastq files were quality checked using FastQC (v 0.11.9). Adapter

sequences were removed with CutAdapt (v3.4). The sequenced reads were aligned to

the dmel_r6.26 genome assembly using Bowtie2 (v2.4.2; Langmead and Salzberg,

2012). Aligned reads were sorted with Samtools (v1.12; Li et al., 2009) and

deduplicated with Picard (v2.23) using the MarkDuplicates command. Aligned, filtered,

and deduplicated reads were used to call peaks using MACS2 software (v2.2.5; Zhang

et al., 2008). Then, featureCounts (v2.6.4; Liao et al., 2019) was used to generate the

count matrix. Differential accessibility analysis was performed using DESeq2 (v1.32.0;

Love et al., 2014). Insert size distributions were calculated using Picard. Peaks were

visualized using the IGV browser (v2.7.2).

Statistical analysis

The data are represented as means ± standard deviation. Comparisons between

homogenizer size and fly volume were analyzed by two-way ANOVA with Bonferroni 's

multiple comparison tests. Freezing time was analyzed by one-way ANOVA using

Tukey's multiple comparison tests. Eye color experiments, % read alignment, and

fractions of reads in peaks (FRiP) were analyzed using Student's t-tests. Gal4-UAS and

washing experiments were analyzed by simple linear regression, and analysis of

covariance (ANCOVA). All statistical analyses were performed using GraphPad Prism 9

software (San Diego, CA, USA).

RESULTS

Sample volumes, homogenization, and freezing time

Our goal was to identify the optimal conditions for nuclei isolation, purification, and

ATAC-seq library preparation using adult *Drosophila*. We first asked if the number of

flies used for head isolation and the size of the Dounce homogenizer play a role in the

number of extracted nuclei that are labeled with DAPI. Using a larger volume of flies

increased the number of extracted nuclei (2-way ANOVA, $F_{(1,8)} = 15.61$, $p = 0.0042$).

The size of the Dounce homogenizer used for extraction did not significantly affect the

total extracted nuclei if analyzed by 2-way ANOVA ($F_{(1,8)} = 1.984$, $p = 0.1966$; Fig. 1A),

however, with 0.5 mL of flies, more nuclei were isolated with the 7 mL homogenizer

(significant by isolated t-test). Thus, we recommend using the larger, 7 mL Dounce
homogenizer to maximize nuclei extraction.

One advantage to using *Drosophila* is the availability of sophisticated genetic tools that
allows the easy identification of target cells, which can then be isolated by flow
cytometry. In our nuclei isolation procedure, *nSyb-Gal4* flies expressing pan-neuronal
GFP-nls are frozen at -80 °C for 5 min and passed through a fly sieve to isolate the
tissue of interest (bodies vs. heads). We hypothesized that extended freezing at -80 °C
could reduce GFP fluorescence, so we performed a time-course experiment to examine
whether freezing time influences GFP fluorescence. Five min is about the minimum time
per sample in which flies can be frozen at -80 °C and homogenized when handling
multiple samples. We observed that GFP fluorescence decreased quickly with
additional time on dry ice but reached a plateau at ~60% fluorescence in the time
analyzed (Fig. 1B, one-phase exponential decay, K = 0.51 min^{-1}, goodness of fit R^2 =
0.64). Therefore, we suggest that flies should only be frozen briefly (5 min or less) at -80
°C when used for nuclei isolation and ATAC-seq library preparation.

Flow cytometer specificity and sensitivity

Sorting GFP+ nuclei using a flow cytometer is dependent on the gating scheme for the
detection of green fluorescent signal. Detection schemes for stains usually come at a
low gate (less signal required) and a high gate (more signal required). To determine the
sorting sensitivity at these gates, we ubiquitously expressed a brighter version of
nuclear GFP, *UAS-Stinger*, using a strong *Tubulin-Gal4* driver. At the low gate, we
recovered ~80% of nuclei from ~100,000 DAPI+ nuclei (Fig. 2A), while we recovered

~12% of nuclei with the high gate (Fig. 2B). An independent experiment showed similar

percentages. The data from these experiments suggest that the low gate has 6-7 fold

better sensitivity than the high gate.

To examine the sorting specificity, we evaluated the number of sorted 'GFP+ nuclei' in

w* Berlin flies, which are all GFP-negative. Sorting at the low gate yielded 0.093%

'GFP+' nuclei (Fig. 3A), while at the high gate, 0.026% of nuclei were 'GFP+' (Fig. 3B).

An independent experiment gave the same results. Together, these data suggest that

the high gate has 4-fold higher specificity and is better at calling true positives. As

expected, there is a tradeoff between sensitivity and specificity when using low and high

GFP+ sorting gates.

Drosophila-specific parameters affect nuclei isolation

Using the high sorting gate, 1 in ~4000 nuclei are selected as false-positive. At face

value, this is good specificity. However, assuming that a *Drosophila* head contains

>200,000 nuclei, >50 of those will be sorted as false positives at the high gate (and

>200 at the low gate). These numbers would significantly contaminate the nuclei sorted

from a genotype where Gal4 is expressed in a subset of 100s of neurons. The high gate

will produce fewer false positives, thus increasing the specificity for the Gal4-expression

pattern, but the high gate will also reduce the sensitivity and recovery of those Gal4+

neurons. Therefore, we asked whether we could increase the recovery via genetic

means by increasing the number of *Gal4* or *UAS-GFP-nls* transgenic constructs to

boost GFP expression. We used *ics-Gal4 UAS-GFP-nls* flies and crossed them to

themselves, *ics-Gal4*, *UAS-GFP-nls*, or *w* Berlin* to generate flies with 4, 3, 3, or 2

copies of *Gal4/UAS* insertions, respectively. *ics-Gal4* expression includes many

neurons and the expression pattern is not different between *ics-Gal4/+* hetero- and *ics-*

Gal4 homozygotes. (Ojelade et al., 2015). Expanding the copies of *Gal4/UAS* insertions

increased the number of detected GFP+ nuclei (linear regression, slope ≠ 0, *p* = 0.0008,

low gate; *p* = 0.001, high gate, Fig. 4A). At both gates, the recovery with 4 transgenes

was about 20-fold higher than recovery with 2 transgenes. We therefore recommend

that for Gal4-lines with sparse expression patterns, the amount of GFP produced should

be increased genetically.

Most *Drosophila* transgenic insertions are marked with a miniature copy of the *white$^+$*

gene, *w^{+mC}*. The presence of such a transgene will confer anywhere from pale yellow to

red-colored eyes. In addition, the allele *w^{+mC}* is dosage-sensitive, *i.e.* the more

transgenes in the fly labeled with *w^{+mC}*, the closer to wild-type red the eye color will be.

Since we suggested above that increasing the copy of *Gal4* and/or *UAS* transgenes will

increase the recovery of GFP+ nuclei, we next wanted to determine if red-colored eye

pigment interferes with sorting GFP+ nuclei. We crossed *w$^-$;Cha-Gal4 UAS-GFP-nls*

males to white-eyed (*w$^-$*), or to red-eyed (*w$^+$*) virgins and then collected the male

progeny. Both of these male genotypes contain 2 *w^{+mC}* alleles from the transgenes, but

one group is red-eyed (*w$^+$*) and the other is orange-eyed (in a *w$^-$* background). At the

low gate, we recovered almost twice as many 'GFP+ nuclei' (Fig. 4B), suggesting that

red eye pigment from *w$^+$* contributes to the false positive rate. We hypothesized that

increasing the number of washes during nuclei isolation would mitigate the false positive

rate caused by red eye pigments. Washing the crude nuclei extracts one, three-, or five-

times dose-dependently reduced the GFP+/DAPI+ ratio in red-eyed and orange-eyed

flies. For the red-eyed flies, we observed the steepest decrease between one and three

washes, and with three washes, there was no longer a difference in the GFP+/DAPI+

ratio between red- and orange-eye flies (Fig. 4C,D). We therefore recommend at least

three washes, especially when sorting with the low-stringency gate and as more w^{+mC}-

marked transgenes are used, since the eye color will approach wild-type red.

Reaction conditions affect library quality

Published ATAC-seq guidelines are based on mammalian genomes, which are

approximately 10-fold larger than the *Drosophila* genome (Vinogradov, 2004; Canapa et

al., 2016). Therefore, the parameters used for the Tn5 transposition reaction (incubation

time and enzyme concentration) in mammalian cells may not be appropriate for fly

nuclei. To determine the optimal Tn5 reaction conditions, we sorted GFP+ nuclei from

fly heads and constructed ATAC-seq libraries. We evaluated the library quality using the

guidelines recommended by Buenrostro et al. (2015). We first examined how reaction

time affects the fragment distribution by sorting pan-neuronal GFP+ nuclei from *nSyb-*

Gal4 UAS-GFP-nls fly heads. Then, we prepared ATAC-seq libraries using 1X Tn5

enzyme and sample incubation for 5, 23, and 60 min (Fig. 5A). A low reaction time

decreased the nucleosome-free peak and increased the proportion of larger fragments

up to 1000 bp, suggesting that di-, tri- and multi-nucleosome fragments are over-

represented. Incubation for 23 min provided a fragment distribution with more

nucleosome-free and fewer mono- and di-nucleosome fragments, while increasing the

reaction time to 60 min biased the library toward smaller fragment sizes with relatively

little evidence of larger fragments. Thus, for library preparation using *Drosophila* nuclei,

23 minutes is the optimal reaction time. We next examined how the Tn5 concentration

affects libraries by collecting GFP+ nuclei from *tubulin-Gal4 UAS-stinger-nls* flies and

tagmenting the DNA using 0.1X, 0.3X, 1X, and 3X Tn5 (compared to manufacturer

recommendation; Fig. 5A). 1X Tn5 showed the largest peak around 200 bp, with

additional skewing towards larger fragments. Reacting DNA using 3X Tn5 reduced both

skewness toward larger fragments and the number of nucleosome-free fragments due

to over-digestion of DNA [which was size-filtered (150~2000 bp) before analysis, Fig.

5B]. Reducing the Tn5 concentration to 0.1X showed skewness to higher fragments, but

overall fewer amplified fragments and reduced overall recovery. We therefore

recommend 1X Tn5 as the best balance between recovery of amplified fragments and

avoiding over-digestion. Finally, we examined how library amplification affects quality.

Like other sequencing library types, ATAC-seq libraries must be amplified to obtain

sufficient DNA concentrations for sequencing. Buenrostro et al. (2013, 2015) suggested

that ATAC-seq libraries should be amplified to 33% of maximal fluorescence obtained in

a qPCR pre-reaction to ensure that sufficient material is present for sequencing, without

the introduction of GC and size bias. We hypothesized that the smaller genome size in

Drosophila would require more PCR cycles than libraries derived from mammalian

nuclei. We tagmented DNA from pan-neuronal GFP+ nuclei from *nSyb-Gal4 UAS-GFP-*

nls fly heads with 1X Tn5 and amplified the tagmented DNA to 25% (8 cycles), 33% (9

cycles), and 50% (10 cycles) of total fluorescence in a qPCR side reaction (40 cycles) to

determine the number of PCR cycles needed for library amplification before sequencing

(Fig. 5C). The library that was amplified to 33% total fluorescence had the smallest

amount of nucleosome-free fragments and a large proportion of ~800-1200 bp

fragments, which may correspond to a PCR bubble that arose from depleted PCR

reagents, most likely primers (Kanagawa, 2003). In contrast, the library amplified to

25% total fluorescence showed a broad nucleosome-free peak and an even distribution

of higher molecular weight fragments, while the library amplified to 50% total

fluorescence showed a tall nucleosome-free peak that suggests a bias toward low

molecular weight amplification. These results suggest that ATAC-seq libraries derived

from *Drosophila* nuclei should be amplified to 25% of total qPCR fluorescence, though

33%-amplified libraries may also be suitable for sequencing after reconditioning PCR

(Thompson et al., 2002).

Optimized reaction conditions generate ATAC-seq libraries conforming to ENCODE

standards

Finally, we combined our optimized parameters to generate and sequence ATAC-seq

libraries. We drove nuclear GFP expression in dopaminergic and GABAergic neurons

and isolated the nuclei after freezing approximately 1 mL (300-400) flies at -80 °C for 5

min. The crude nuclei extracts were washed three times and sorted by flow cytometry

using a high-stringency gate. We collected 65,000 nuclei per sample for each neuron

type, which were tagmented for 23 min using 1X Tn5 (from an Illumina Nextera kit). The

resulting libraries were amplified to 25% qPCR fluorescence (8 cycles) and analyzed

using 50-bp paired end sequencing. We then examined the percent alignment to the

genome, the fraction of reads in peaks (FRiP), and fragment length distribution of the

sequenced libraries. Both libraries showed high alignment rates, with 79% and 94%

alignment in dopaminergic and GABAergic libraries, respectively (Fig. 6A). Dopamine

neuron libraries had 0.37 FRiP, while GABA neuron libraries had 0.40 FRiP (Fig. 6B),

which were both higher than the FRiP values (>0.3) recommended by the ENCODE

consortium for ATAC-seq libraries (Davis et al., 2018). Finally, each of these libraries

showed the expected nucleosomal banding pattern (Fig. 6C). Together, these metrics

indicate that we generated high-quality libraries and that our protocol returns high-

quality sequencing data. To test this hypothesis, we examined ATAC-seq peaks in

regions associated with classical and neuron markers identified by single-cell RNA-seq

(Davie et al., 2018). We observed differential peaks, indicating increased accessibility,

in *ple*, *DAT*, and *hth*, which encode tyrosine hydroxylase, the rate-limiting dopamine-

synthesis enzyme, the dopamine transporter, and homothorax, a homeobox

transcription factor, respectively. These peaks were significantly more open in

dopamine neurons than in GABA neurons (peaks indicated by asterisks in Fig. 7A, from

left to right; *ple*: p = 3.32e-65 and 5.01e-53; *DAT*: p = 1.03e-05 and 1.04e-05; *hth*: p =

0.009, 1.69e-36, 1.14e-21, and 0.0008). Peaks associated with dopamine-neuron

specific genes were much smaller or absent in GABA neurons (Fig. 7). The peaks

corresponding to the GABAergic markers include *Gad1*, which encodes glutamic acid

decarboxylase, which synthesizes GABA, *Lim3*, and *CG14989*. Peaks associated with

GABA marker genes were significantly more open in GABA neurons than in dopamine

neurons (peaks indicated by asterisks in Fig. 7B, from left to right; *GAD1*: p = 0.0005,

4.42e-08, 0.04, 0.01, and 6.6e-08; *Lim3*: p = 1.39e-05 and 0.001; *CG14989*; p = 0.1).

These peaks were much smaller or absent in dopaminergic neurons. These results

confirm that our nuclei isolation and library generation protocol yields high quality data,

including differential chromatin accessibility in known cell-type markers (Davie et al.,

2018).

DISCUSSION

ATAC-seq is a powerful way to interrogate the chromatin state within tissues of interest.

Critically, the technique provides key advantages over previous methods, such as ChIP-

seq, MNase-seq, or FAIRE-seq, including low input requirements (50,000-60,000 nuclei

vs. 1-10 million nuclei for MNase- or FAIRE-seq) (Kidder et al., 2011; Simon et al.,

2012; McKay and Lieb, 2013), less hands-on time, no harsh chemicals such as

paraformaldehyde, and no antibody optimization or complicated pull-down methods.

Thus, ATAC-seq is quickly becoming the assay of choice for chromatin studies.

Most published studies using ATAC-seq focus on either tumor tissue or white blood

cells, though recent studies have investigated the brain and other tissues in mice and

humans (Liu et al., 2019; Rocks et al., 2021). The initial study describing ATAC-seq was

performed in mammalian tissue (Buenrostro et al., 2013), and published protocols have

been optimized for use in these tissues (Corces et al., 2017; Buenrostro et al., 2018).

Drosophila melanogaster is a powerful model organism that is routinely used to study

development and human disease models. Because we are interested in *Drosophila*

neurons from the adult brain, we first determined some basic parameters for nuclei

isolation. We found that freezing flies for ≥ 5 min and shaking off the heads in a sieve to

enrich for brain tissue did not impede sorting GFP+ nuclei, but freezing flies for a longer

duration did reduce GFP+ nuclei recovery. Using adult *Drosophila* – heads or whole

flies – for ATAC-seq experiments presents some complicating factors, such as the

exoskeleton. While brains can be manually dissected, for more efficient throughput and

to be able to sort enough GFP+ nuclei from populations of few neurons, we used

undissected heads for nuclei isolation. We homogenized flies with a Dounce

homogenizer, with a filtration step between homogenization with the "A" and "B" pestles.

The first step disrupts the fly exoskeleton, as shown by empty fly heads in the crude

homogenate, indicating that the tissue inside the head is pushed out of the exoskeleton

during homogenization. Passing the crude homogenate through a cell strainer

eliminates much of the exoskeleton debris. Subsequent homogenization lyses the

plasma membrane without disrupting the nuclear membrane (Corces et al., 2017).

These freezing and homogenizing steps allowed us to recover up to 80% of GFP+

nuclei when ubiquitously expressing nuclear GFP. Thus, our physical extraction protocol

allows for efficient and sensitive recovery of GFP-labelled nuclei from adult *Drosophila*

heads.

An advantage of using *Drosophila* for ATAC-seq studies is the ability to genetically label

nuclei of interest using the *Gal4/UAS* system. Thousands of Gal4 lines labeling distinct

neurons are available (Jenett et al., 2012), including ones specific to neurotransmitters

(Deng et al., 2019), allowing for the specific, reproducible interrogation of only a defined

subset of neurons. As mentioned above, we were able to recover up to 80% of

ubiquitous GFP+ nuclei at the low stringency gate, indicating good sensitivity. To

determine the specificity, we sorted nuclei from GFP-negative flies and recovered 0.1%

and 0.025% false positive nuclei at the low- and high-stringency gates. While these

seem to be acceptable numbers for GFP+ specificity, they may present a problem when

sorting from Gal4 lines with sparse expression. Sorting at the high-stringency gate will

produce fewer false positives but will also reduce the sensitivity for collecting true

positives. Possible solutions to the issue of insufficient GFP+ nuclei recovered include

increasing the numbers of fly heads and increasing the GFP signal. We found that

increasing the number of *Gal4* and *UAS-GFP-nls* transgenes increased the number of

recovered GFP+ nuclei. The versatile *Gal4/UAS* system therefore allows for some

flexibility to be able to sort ≥50,000 nuclei for ATAC-seq library preparation. We

recommend that specific *Gal4* and *UAS-GFP-nls* combinations be tested by sorting

nuclei to get information about the recovery of GFP+ nuclei prior to performing actual

experiments consisting of library preparation and deep sequencing of experimental and

control flies.

The *Drosophila* genome is ~10-fold smaller than mammalian genomes, so enzyme-

mediated steps during ATAC-seq library preparation may be affected by the amount of

available DNA. Tn5 tagmentation in mammalian samples is generally performed using

undiluted Tn5 enzyme included in a library preparation kit (1X) for 30 min at 37 °C

(Buenrostro et al., 2015). Because the amount of accessible chromatin in *Drosophila*

nuclei may be lower than in mammalian nuclei, we investigated the effects of changing

the parameters of the enzymatic reactions (Tn5 tagmentation and PCR amplification) on

fragment distribution and library quality. We observed that increasing or decreasing the

Tn5 concentration reduced library complexity. Previous studies showed that decreasing

the Tn5 concentration in mouse embryonic stem cells only decreases tagmentation

efficiency when diluted to 10 nM or lower (Corces et al., 2017). Notably, this experiment

used homemade Tn5 (Picelli et al., 2014), which may have different kinetics or

concentrations than commercially-available Tn5 enzyme. Increasing the tagmentation

time decreased library complexity, while decreasing the tagmentation time increased

the number of mid-length fragments but did not improve nucleosomal periodicity. We

also examined the number of PCR cycles needed to amplify ATAC-seq libraries before

sequencing. Protocols for generating ATAC-seq libraries from mammalian tissues call

for amplification to one-third the total fluorescence in a qPCR side reaction (Buenrostro

et al., 2015). Because of the smaller genome size in *Drosophila*, amplifying ATAC-seq

libraries to one-third of the total qPCR fluorescence may introduce GC and size bias.

Indeed, amplifying our libraries to one-third qPCR fluorescence introduced a PCR

bubble, indicating that the available primers were depleted during amplification

(Kanagawa, 2003), causing amplification to occur using annealed adaptor sequences

and increasing the mid-size fragments in the library. While these libraries may be

sequenceable after one cycle of reconditioning PCR (Thompson et al., 2002), libraries

with PCR bubbles should be quality-checked to verify that the PCR bubble was resolved

before sequencing. Amplifying ATAC-seq libraries to more than 33% total qPCR

fluorescence biases the library toward smaller fragment sizes and reduces library

complexity. Based on these results, we recommend transposition with 1X Tn5 for 23

min, followed by amplification of the ATAC-seq libraries to 25% qPCR fluorescence,

which preserves library complexity and does not introduce GC or size bias.

Using these optimized conditions, we generated ATAC-seq libraries from 65,000 nuclei

isolated from *Drosophila* dopaminergic and GABAergic neurons. These libraries were of

high quality, with the expected nucleosomal banding patterns and fragment size

distributions. Additionally, these libraries had high alignment rates to the genome (92.4-

96.5%), further confirming their quality. Therefore, we propose that ATAC-seq libraries

from *Drosophila* tissues should use the following parameters (Fig. 8): freezing at -80 °C

for no more than 5 min, at least three washes, 23-min tagmentation time with 1X Tn5,

and amplification to 25% qPCR fluorescence. Our protocol harnesses the powerful

genetic tools available in *Drosophila* to achieve excellent cell-type specificity followed by ATAC-seq, which is used to interrogate the chromatin landscape with unprecedented resolution. Thus, our protocol will enable detailed investigations of the role of chromatin-mediated gene regulation in normal and pathological states.

ACKNOWLEDGMENTS

This work was supported by the University of Utah Flow Cytometry Facility, the University of Utah Genomics Core Facility, the High Throughput Sequencing Core at the Huntsman Cancer Institute, and the National Cancer Institute through Award Number 5P30CA042014-24. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

FUNDING

This study was supported by grants from the National Institute on Drug Abuse (Grant R21DA049635 to A.R.), the National Institute on Alcohol Abuse and Alcoholism (Grant R01AA026818 to A.R.), and the National Institute of Diabetes and Digestive and Kidney Diseases (R01DK110358 to A.R.R.).

AUTHOR CONTRIBUTIONS

CBM performed ATAC-seq experiments, analyzed data; and wrote the manuscript; MAP performed nuclei isolation, sorting, and optimization experiments; ABM analyzed data; ARR reviewed and edited the manuscript; AR supervised the project and revised and edited the manuscript. All authors reviewed the manuscript.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

DATA AVAILABILITY

The datasets generated by this study are available upon reasonable request.

REFERENCES

Bozek M, Cortini R, Storti AE, Unnerstall U, Gaul U, Gompel N (2019) ATAC-seq

reveals regional differences in enhancer accessibility during the establishment of spatial

coordinates in the Drosophila blastoderm. Genome Res 29:771–783.

Brovkina M V., Duffié R, Burtis AEC, Clowney EJ (2021) Fruitless decommissions

regulatory elements to implement cell-type-specific neuronal masculinization. PLOS

Genet 17:e1009338.

Buenrostro JD, Corces MR, Lareau CA, Wu B, Schep AN, Aryee MJ, Majeti R, Chang

HY, Greenleaf WJ (2018) Integrated Single-Cell Analysis Maps the Continuous

Regulatory Landscape of Human Hematopoietic Differentiation. Cell 173:1535-

1548.e16.

Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ (2013) Transposition of

native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-

binding proteins and nucleosome position. Nat Methods 10:1213–1218.

Buenrostro JD, Wu B, Chang HY, Greenleaf WJ (2015) ATAC-seq: A Method for

Assaying Chromatin Accessibility Genome-Wide. In: Current Protocols in Molecular

Biology, pp 21.29.1-21.29.9. Hoboken, NJ, USA: John Wiley & Sons, Inc.

Bysani M, Agren R, Davegårdh C, Volkov P, Rönn T, Unneberg P, Bacos K, Ling C

(2019) ATAC-seq reveals alterations in open chromatin in pancreatic islets from

subjects with type 2 diabetes. Sci Rep 9.

Canapa A, Barucca M, Biscotti MA, Forconi M, Olmo E (2016) Transposons, genome

size, and evolutionary insights in animals. Cytogenet Genome Res 147:217–239.

Corces MR, Trevino AE, Hamilton EG, Greenside PG, Sinnott-Armstrong NA, Vesuna

S, Satpathy AT, Rubin AJ, Montine KS, Wu B, Kathiria A, Cho SW, Mumbach MR,

Carter AC, Kasowski M, Orloff LA, Risca VI, Kundaje A, Khavari PA, Montine TJ,

Greenleaf WJ, Chang HY (2017) An improved ATAC-seq protocol reduces background

and enables interrogation of frozen tissues. Nat Methods 14:959–962.

Cusanovich DA, Daza R, Adey A, Pliner HA, Christiansen L, Gunderson KL, Steemers

FJ, Trapnell C, Shendure J (2015) Multiplex single cell profiling of chromatin

accessibility by combinatorial cellular indexing. Science 348:910–914.

Cusanovich DA, Reddington JP, Garfield DA, Daza RM, Aghamirzaie D, Marco-

Ferreres R, Pliner HA, Christiansen L, Qiu X, Steemers FJ, Trapnell C, Shendure J,

Furlong EEM (2018) The cis-regulatory dynamics of embryonic development at single-

cell resolution. Nature 555:538–542.

Davie K, Jacobs J, Atkins M, Potier D, Christiaens V, Halder G, Aerts S (2015)

Discovery of Transcription Factors and Regulatory Regions Driving In Vivo Tumor

Development by ATAC-seq and FAIRE-seq Open Chromatin Profiling. PLoS Genet

11:1–24.

Davie K, Janssens J, Koldere D, De Waegeneer M, Pech U, Kreft Ł, Aibar S, Makhzami

S, Christiaens V, Bravo González-Blas C, Poovathingal S, Hulselmans G, Spanier KI,

Moerman T, Vanspauwen B, Geurs S, Voet T, Lammertyn J, Thienpont B, Liu S,

Konstantinides N, Fiers M, Verstreken P, Aerts S (2018) A Single-Cell Transcriptome

Atlas of the Aging Drosophila Brain. Cell 174:982-998.e20.

Davis CA, Hitz BC, Sloan CA, Chan ET, Davidson JM, Gabdank I, Hilton JA, Jain K,

Baymuradov UK, Narayanan AK, Onate KC, Graham K, Miyasato SR, Dreszer TR,

Strattan JS, Jolanki O, Tanaka FY, Cherry JM (2018) The Encyclopedia of DNA

elements (ENCODE): data portal update. Nucleic Acids Res 46:D794–D801.

Deng B, Li Q, Liu X, Cao Y, Li B, Qian Y, Xu R, Mao R, Zhou E, Zhang W, Huang J,

Rao Y (2019) Chemoconnectomics: Mapping Chemical Transmission in Drosophila.

Neuron 101:876-893.e4.

Franceschini N, Kirschfeld K, Minke B (1981) Fluorescence of photoreceptor cells

observed in vivo. Science (80-) 213:1264–1267.

Fujiwara S, Baek S, Varticovski L, Kim S, Hager GL (2019) High Quality ATAC-Seq

Data Recovered from Cryopreserved Breast Cell Lines and Tissue. Sci Rep 9:1–11.

Haines JE, Eisen MB (2018) Patterns of chromatin accessibility along the anterior-

posterior axis in the early Drosophila embryo. PLoS Genet 14.

Halstead MM, Kern C, Saelao P, Chanthavixay G, Wang Y, Delany ME, Zhou H, Ross

PJ (2020) Systematic alteration of ATAC-seq for profiling open chromatin in

cryopreserved nuclei preparations from livestock tissues. Sci Rep 10.

Ho YT, Shimbo T, Wijaya E, Ouchi Y, Takaki E, Yamamoto R, Kikuchi Y, Kaneda Y,

Tamai K (2018) Chromatin accessibility identifies diversity in mesenchymal stem cells

from different tissue origins. Sci Rep 8:1–11.

Jenett A, Rubin GM, Ngo TTB, Shepherd D, Murphy C, Dionne H, Pfeiffer BD, Cavallaro

A, Hall D, Jeter J, Iyer N, Fetter D, Hausenfluck JH, Peng H, Trautman ET, Svirskas

RR, Myers EW, Iwinski ZR, Aso Y, DePasquale GM, Enos A, Hulamm P, Lam SCB, Li

HH, Laverty TR, Long F, Qu L, Murphy SD, Rokicki K, Safford T, Shaw K, Simpson JH,

Sowell A, Tae S, Yu Y, Zugates CT (2012) A GAL4-Driver Line Resource for Drosophila

Neurobiology. Cell Rep 2:991–1001.

Kanagawa T (2003) Bias and artifacts in multitemplate polymerase chain reactions

(PCR). J Biosci Bioeng 96:317–323.

Kidder BL, Hu G, Zhao K (2011) ChIP-Seq: Technical considerations for obtaining high-

quality data. Nat Immunol 12:918–922.

Koenecke N, Johnston J, Gaertner B, Natarajan M, Zeitlinger J (2016) Genome-wide

identification of Drosophila dorso-ventral enhancers by differential histone acetylation

analysis. Genome Biol 17:1–19.

Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. Nat

Methods 2012 94 9:357–359.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,

Durbin R (2009) The Sequence Alignment/Map format and SAMtools. Bioinformatics

25:2078–2079.

Liao Y, Smyth GK, Shi W (2019) The R package Rsubread is easier, faster, cheaper

and better for alignment and quantification of RNA sequencing reads. Nucleic Acids Res

47.

Liu C, Wang M, Wei X, Wu L, Xu J, Dai X, Xia J, Cheng M, Yuan Y, Zhang P, Li J, Feng

T, Chen A, Zhang W, Chen F, Shang Z, Zhang X, Peters BA, Liu L (2019) An ATAC-seq

atlas of chromatin accessibility in mouse tissues. Sci Data 6.

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol 15:1–21.

Lu Z, Hofmeister BT, Vollmers C, DuBois RM, Schmitz RJ (2017) Combining ATAC-seq

with nuclei sorting for discovery of cis-regulatory regions in plant genomes. Nucleic

Acids Res 45.

McKay DJ, Lieb JD (2013) A Common Set of DNA Regulatory Elements Shapes

Drosophila Appendages. Dev Cell 27:306–318.

Miller G V, Itoku KA, Fleischer AB, Stark WS (1984) Studies of fluorescence in

Drosophila compound eyes: changes induced by intense light and vitamin A deprivation.

Springer-Verlag.

Ojelade SA, Jia T, Rodan AR, Chenyang T, Kadrmas JL, Cattrell A, Ruggeri B, Charoen

P, Lemaitre H, Banaschewski T, Büchel C, Bokde ALW, Carvalho F, Conrod PJ, Flor H,

Frouin V, Gallinat J, Garavan H, Gowland PA, Heinz A, Ittermann B, Lathrop M, Lubbe

S, Martinot JL, Pausu T, Smolka MN, Spanagel R, O'Reilly PF, Laitinen J, Veijola JM,

Feng J, Desrivières S, Jarvelin MR, Schumann G, Rothenfluh A (2015) Rsu1 regulates

ethanol consumption in Drosophila and humans. Proc Natl Acad Sci U S A 112:E4085–

E4093.

Picelli S, Björklund AK, Reinius B, Sagasser S, Winberg G, Sandberg R (2014) Tn5

transposase and tagmentation procedures for massively scaled sequencing projects.

Genome Res 24:2033–2040.

Quillien A, Abdalla M, Yu J, Ou J, Zhu LJ, Lawson ND (2017) Robust Identification of

Developmentally Active Endothelial Enhancers in Zebrafish Using FANS-Assisted

ATAC-Seq. Cell Rep 20:709–720.

Rocks D, Jaric I, Tesfa L, Greally JM, Suzuki M, Kundakovic M (2021) Cell type-specific

chromatin accessibility analysis in the mouse and human brain. Epigenetics:1–18.

Simon JM, Giresi PG, Davis IJ, Lieb JD (2012) Using formaldehyde-assisted isolation of

regulatory elements (FAIRE) to isolate active regulatory DNA. Nat Protoc 7:256–267.

Thompson JR, Marcelino LA, Polz MF (2002) Heteroduplexes in mixed-template

amplifications: Formation, consequence and elimination by "reconditioning PCR."

Nucleic Acids Res 30:2083–2088.

Vinogradov AE (2004) Evolution of genome size: Multilevel selection, mutation bias or

dynamical chaos? Curr Opin Genet Dev 14:620–626.

Wang J, Zibetti C, Shang P, Sripathi SR, Zhang P, Cano M, Hoang T, Xia S, Ji H,

Merbs SL, Zack DJ, Handa JT, Sinha D, Blackshaw S, Qian J (2018) ATAC-Seq

analysis reveals a widespread decrease of chromatin accessibility in age-related

macular degeneration. Nat Commun 9:1–13.

Wolf FW, Rodan AR, Tsai LTY, Heberlein U (2002) High-resolution analysis of ethanol-

induced locomotor stimulation in Drosophila. J Neurosci 22:11035–11044.

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nussbaum C,

Myers RM, Brown M, Li W, Liu XS (2008) Model-based Analysis of ChIP-Seq (MACS).

Genome Biol 9:R137.

FIGURE CAPTIONS

Figure 1. Sample volume and freezing time affect nuclei yield.

A) The number of flies used to collect heads prior to nuclei isolation increases the

number of extracted nuclei, but the homogenizer size does not. The indicated volumes

of *w* Berlin* flies were frozen, the heads shaken off in a sieve, the nuclei were extracted

and stained with DAPI, and were then counted in a flow cytometer. Data represent the

mean ± SD of three independent experiments. B) Freezing time longer than 5 min

decreases GFP detection by flow cytometry. Flies with pan-neuronal GFP, *nSyb-Gal4*

UAS-GFP-nls were counted at the low gate. Data represent three biological replicates.

The data in A) were analyzed by two-way ANOVA and the data in B) were analyzed by

nonlinear regression using one-phase exponential decay, $K = 0.51$ min^{-1}, goodness of fit

$R^2 = 0.64$. **$p < 0.01$.

Figure 2. Flow cytometry shows high specificity and few false positives.

Nuclei were isolated from *w* Berlin* heads (GFP-negative) and analyzed by flow

cytometry. A) Representative dot plots of nuclei analyzed using the low gate. Nuclei

were identified by DAPI staining (Y-axis) and forward light scatter (X-axis, left). The box

indicates the sorted DAPI+ singlet nuclei, which were then analyzed for green

fluorescence (Y-axis) and the red PE-A channel (X-axis, right), which allowed the best

separation of GFP populations, and sorted at the low gate (solid trapezoid box). 0.093%

of input singlets were GFP false-positives (and 0.024% when applying the high gate in

this sort). B) Representative dot plots of nuclei analyzed using the high gate. Nuclei

were isolated as in A), shown on the left, and then sorted for green fluorescence at the

high gate (stippled trapezoid box, right). 0.026% of input singlets were GFP false-

positives (and 0.10% when applying the low gate in this sort).

Figure 3. Differential sensitivity at the low and high sorting gate.

UAS-Stinger, encoding a super-bright nuclear GFP, expression was driven using

ubiquitous *Tubulin-Gal4* (all nuclei are GFP+). Nuclei from heads were isolated as

described in Fig. 2 (left sides) and analyzed by flow cytometry using low- and high-

stringency gating (right). A) 79% of nuclei were called as GFP+ at the low gate (solid

trapezoid). B) 12% of nuclei were called as GFP+ at the high gate (stippled trapezoid,

right).

Figure 4. Physical parameters affecting ATAC-seq library preparation from

***Drosophila* nuclei. ChangesToFig**

A) GFP+ nuclei from flies expressing multiple copies of Gal4 + UAS (*ics-Gal4 UAS-*

GFP-nls) were isolated and analyzed by flow cytometry using low- and high-stringency

gating. B) Nuclei from cholinergic neurons (heads of *Cha-Gal4 UAS-GFP-nls*) red-eyed

(w^+ background) and orange-eyed flies (w^- background) were isolated and analyzed by

flow cytometry using low- and high-stringency gating. Error bars represent the mean ±

SD of six biological replicates and were analyzed by Student's t-tests. C-D) Nuclei from

the same flies used in B) were washed multiple times after isolation and before analysis

using low- (C) and high- (D) stringency gating. The data was analyzed using linear

regression. All regression lines have slopes ≠ 0, $p < 0.02$.

Figure 5. Optimal library construction parameters

GFP expression was driven using various fly genotypes. Then, nuclei were collected by

flow cytometry using high-stringency gating. The isolated nuclei were used in ATAC-seq

library construction, with variable preparation conditions as indicated. After

amplification, each library was purified using AMPure XP beads, which remove low- and

high-molecular weight fragments. The fragment sizes and relative fluorescence from

Tapestation data are shown. A) Tn5 reaction time; B) Tn5 concentration; and C) post-

Tn5 PCR amplification. GFP expression in A) was driven using *nSyb-Gal4 UAS-GFP-*

nls; GFP expression in B) was driven using *Tubulin-Gal4 UAS-Stinger*; and GFP

expression in C) was driven using *nSyb-Gal4 UAS-GFP-nls.*

Figure 6. Optimized ATAC-seq parameters generate consistent libraries.

Representative traces from ATAC-seq libraries prepared from dopaminergic and

GABAergic nuclei isolated from adult *Drosophila* heads. The libraries were prepared

using nuclear GFP+ expression driven by *TH-Gal4* (dopamine) and *vGAT-Gal4* (GABA).

Flies were frozen at -80 °C for 5 min per replicate (300-400 flies per replicate; 3

replicates per genotype), the heads were collected, and the nuclei were isolated by flow

cytometry using high-stringency gating. The target DNA was incubated for 23 minutes

with 1X Tn5, and the libraries were PCR-amplified for 8 cycles, which corresponded to

25% total fluorescence from the qPCR side reaction. Amplified libraries were purified

using AMPure XP beads and sequenced using a Novaseq 6000 instrument using 50 bp

paired-end reads. A) Fraction of reads aligned to the nuclear genome. B) Fraction of

reads in peaks (FRiP) after alignment, filtering, and peak calling. C) Insert size

distribution showing the expected nucleosomal periodicity. Inset: Log-scaled insert size

distribution. In A) and B), the data represent the mean ± SD of three replicates per

neuron type. Statistical differences were analyzed using Mann-Whitney U tests.

Figure 7. Optimized ATAC-seq libraries return expected cellular markers.

ATAC-seq peaks in marker genes for dopaminergic and GABAergic neurons. ATAC-seq

reads from dopamine neurons (tyrosine hydroxylase: TH+) are shown in orange and

reads from GABA (vesicular GABA transporter: vGAT+) neurons are shown in blue. The

genes and gene structures are shown below the ATAC-seq peaks. A) Peaks in

chromatin regions that are more open in dopamine neurons. *ple* (encoding tyrosine

hydroxylase; $p = 2.43e-68$) and *DAT* (dopamine transporter; $p = 9.64e-7$) are classical

dopamine neuron markers. *hth* (homothorax; $p = 1.16e-52$) is a newly described

dopamine neuron marker. B) Chromatin regions that are more open in GABA neurons.

Gad1 (glutamic acid decarboxylase 1; $p = 6.64e-8$) is a classical GABAergic neuron

marker, while *Lim3* (Lim3; $p < 0.0002$) and *CG14989* (uncharacterized protein; $p = 0.1$)

were identified as additional GABA neuron markers. *CG14989*, *Lim3*, and *hth* were

described as specific markers using single-cell RNA sequencing (Davie et al., 2018).

The boxes in the gene structures indicate exons and the arrows indicate the

transcription direction.

Figure 8: Flow chart of optimized ATAC-seq protocol

Suggested workflow, which we routinely use to generate ATAC-seq libraries from adult

Drosophila. The libraries prepared using the workflow conform to ENCODE guidelines,

with typical markers indicating a high-quality library, including nucleosomal banding

pattern, high genome alignment rates, and high fraction of reads in peaks (FRiP). In the

workflow, blue boxes indicate procedural steps, green diamonds indicate decision

points, and yellow diamonds indicate decision points based on QC metrics. On the right

of each procedural box is shown the approximate time required for each step.

Figure 1.

a

b

Figure 2.

a Low gate

Total nuclei: 375931
Singlets: 30.4%

Singlets: 114351
GFP+ singlets:
Low gate, 0.093%

b High gate

Total nuclei: 332585
Singlets: 32.7%

Singlets: 108782
GFP+ nuclei
High gate, 0.026%

Figure 3.

a
b

Total nuclei: 225812
Percent singlets: 44.2%

Total singlets: 99807
GFP+ singlets: 79.1%

c
d

Total nuclei: 231184
Percent singlets: 43.0%

Total singlets: 99329
GFP+ singlets: 12.2%

Figure 4.

Figure 5.

Figure 6.

Figure 7.

a Dopaminergic markers

Sorted for: TH+ vGAT+

b GABAergic markers

Figure 8.