Dynamics of pollination and seed dispersal mutualisms at low density

Kayla R. S. Hale[1*], Daniel P. Maes[1,2], & Fernanda S. Valdovinos[1,3*]

[1] Department of Ecology and Evolutionary Biology, University of Michigan, Biological

Sciences Building, 1105 North University Ave, Ann Arbor, MI 48109, USA.

[2] Department of Mathematics, University of Michigan, East Hall, 530 Church St, Ann

Arbor, MI 48109, USA.

[3] Department of Environmental Science and Policy, University of California, Davis,

Wickson Hall, One Shields Avenue, Davis, CA 95616.

*Corresponding authors; email: kaylasal@umich.edu, fvaldovinos@ucdavis.edu

Manuscript elements: Figure 1, Figure 2, Figure 3, Table 1, Online Appendices A and B

(including Table A1 and Figures A1-A3). Figures 1-3 are to print in color.

Keywords: transportation mutualism, reproductive services, population dynamics,

thresholds, Allee effects, germination, negative density-dependence

Manuscript type: Major Article.

Running title: Dynamics of mutualisms at low density

Word count: 5545.

Acknowledgements: This research was supported by National Science Foundation Graduate

Research Fellowship DGE-1143953 to K.R.S.H., National Science Foundation Graduate

Research Fellowship DGE-1256260 to D.P.M., and National Science Foundation grant

DEB-1834497 to F.S.V.

Competing interests: The authors declare no competing interests.

Statement of authorship: F.S.V. conceived the study; K.R.S.H. and D.P.M. analyzed the

models; K.R.S.H. and F.S.V. developed the biological interpretation of the model results;

K.R.S.H. wrote the manuscript with important contributions from F.S.V.; all authors

developed the models.

Code availability: Mathematica notebooks used to analyze the models are available by

request and will be available on GitHub upon acceptance.

Abstract

Pollination and seed dispersal mutualisms are critical for biodiversity and ecosystem

services yet face mounting threats from anthropogenic perturbations that cause their

populations to decline. Characterizing the dynamics of these mutualisms when populations

are at low density is important to anticipate consequences of these perturbations. We

developed simple population dynamic models detailed enough to distinguish different

mechanisms by which plant populations benefit from animal pollination or seed dispersal.

We modeled benefits as functions of foraging rate by animals on plant rewards and

specified whether they affected plant seed set, germination, or negative density dependence

during recruitment. We found that pollination and seed dispersal mutualisms are stable at

high density but exhibit different dynamics at low density, depending on plant carrying

capacity, animal foraging efficiency, and whether populations are obligate upon their

partners for persistence. Under certain conditions, all mutualisms experience destabilizing

thresholds in which one population declines because its partner is too rare. Plants

additionally experience Allee effects when obligate upon pollinators. Finally, both

pollination and seed dispersal mutualisms that reduce mortality during recruitment can

exhibit bistable coexistence at low or high density. Insights from our models can inform

conservation efforts, as mutualist populations continue to decline globally.

Introduction

Pollination and seed dispersal mutualisms support vast amounts of biodiversity and

productivity in terrestrial ecosystems. Pollinators aid the reproduction of 78% and 94% of

flowering plants in temperate and tropical regions, respectively (Ollerton *et al.* 2011), while

seed dispersers aid the reproduction of 56% of plant species worldwide (Aslan *et al.* 2013).

In addition, the production of 75% of the world's major crops partially or completely

depend on pollinators (Klein *et al.* 2007). Unfortunately, these important mutualisms face

threats globally. A recent meta-analysis of plant regeneration in forests (Neuschultz *et al.*

2016) found that pollination and seed dispersal are the processes most vulnerable to human

disturbance including climate change, nutrient runoff, pesticide use, and invasive species

(Stachowicz 2001, Traveset & Richardson 2006, Tylianakis *et al.* 2008, Zhou *et al.* 2013).

In fact, there is clear evidence showing the abundance of pollinators and key seed disperser

taxa such as frugivorous birds and mammals are declining globally (Potts *et al.* 2010, 2016,

Wotton & Kelly 2011). Therefore, understanding the population dynamics of species

involved in these mutualisms, especially at low abundances, can inform predictions of the

consequences of those declines on the mutualisms, plant populations and crop production.

Historically, theoretical research (Gause & Witt 1935, Whittaker 1975, Vandermeer

& Boucher 1978, Goh 1979, Addicott 1981, Wolin 1985, Bascompte *et al.* 2006, Okuyama

& Holland 2008, Bastolla *et al.* 2009) mostly used Lotka-Volterra type models (*sensu*

Valdovinos 2019) to investigate the dynamics of populations interacting mutualistically.

This phenomenological representation of mutualistic interactions as net positive effects

between interacting species provided insight for characterizing the effects of facultative,

obligate, linear, and saturating mutualisms on the long-term stability of mutualistic

systems. However, representing mutualisms simply as positive effects between species

overlooks important dynamics that emerge from the specific mechanisms by which species

positively affect each other. More recent research advanced upon these phenomenological

representations of mutualism by developing models that account for consumer-resource

mechanisms (Wright 1989, Holland & DeAngelis 2010, Valdovinos *et al.* 2013, Revilla

2015). Accounting for consumer-resource mechanisms enabled the discovery of important

dynamics, such Allee effects, alternative states, transitions between mutualism and

parasitism, competition among species sharing mutualistic partners, and niche partitioning

(May 1976, Soberón & Martinez del Rio 1981, Wells 1983, Wright 1989, Pierce & Young

1986, Holland & DeAngelis 2010, Valdovinos *et al.* 2013, 2016, Revilla 2015, Valdovinos

& Marsland 2021) as well as the integration of these mutualisms into food web dynamics

(Hale *et al.* 2020). This research, however, mostly focused on animal dynamics (i.e., the

consumer) and the dynamics of plant resources available to the animals (but see Wells

1983). To investigate the consequences of pollinator and seed disperser declines on plant

populations, more focus is needed on the mechanisms by which these animals affect plant

reproduction (Beckman *et al.* 2020). Here, we improve mechanistic understanding of the

ecological dynamics of these mutualisms by developing and analyzing consumer-resource

models that also account for simple mechanisms of reproductive benefits to plants.

Pollination and seed dispersal mutualisms share similarities as "transportation

mutualisms" (Bronstein 2015). In both cases, animals visit plants to feed upon rewards

(such as nectar and fruit) and provide reproductive services (such as transport of pollen and

seeds) incidentally during foraging. However, these interactions differ in the mechanism of

reproductive benefit to plants. Pollinators increase plant seed set by facilitating cross-

fertilization, by moving pollen between conspecific plant individuals (Willmer 2011). Seed

dispersers increase plant recruitment by lessening density-dependent seed(ling) mortality

caused by predators, pathogens, and competitors (Wotton & Kelly 2011, Moore & Dittel

2020). Seed dispersal can also increase germination by improving seed condition during

passage through dispersers' guts (Fricke *et al.* 2013). Here, we investigate the dynamical

consequences of these various mechanisms of reproductive benefits to plants. We find

distinct dynamical behaviors at low population densities, including thresholds, Allee

effects, and bistable coexistence, depending on species obligacy (i.e., their ability to persist

without their partner), plant carrying capacity, and animal foraging efficiency. We also

show that pollination and seed dispersal mutualisms have similar dynamics and stability at

high density but differ in the ecological conditions under which the populations are

vulnerable to collapse.

Methods

We begin by deriving models for animal mutualists, which closely follow previous

consumer-resource approaches (e.g., Wright 1989, Holland & DeAngelis 2010). Then, we

derive novel models for plant mutualists, balancing simplicity with the necessary

mechanistic detail to distinguish between the potential benefits of pollination versus seed

dispersal interactions. We assume that the growth rate of animal and plant populations are

functions of the density-independent per-capita birth (b_A, b_P) and death (b_A, b_P) rates in

absence of their mutualist, where subscripts A and P indicate animal and plant populations,

respectively. These population growth rates are also function of per-capita self-limitation

and other density-dependent processes (s_A, s_P), and the benefits provided by mutualists (see

below). Our models are continuous, which accommodates species with overlapping

generations. They are also deterministic and ignore migration, which allows us to focus on

the dynamics that emerge from the mutualistic interactions, unobscured by stochasticity and

the dynamics of other patches. Table 1 summarizes our parameter definitions for all

models.

121   *Population dynamics of animals*

122   Pollinators and seed dispersers benefit from visiting plants by foraging for rewards

123   that offer primarily nutritional benefit (Willmer 2011, Jordano 2014). We therefore model

124   the change in animal population density ($\boldsymbol{A}$) over time t as

125                         $$\frac{d\boldsymbol{A}}{dt} = b_A\boldsymbol{A} + \varepsilon C_R - s_A\boldsymbol{A} - d_A\boldsymbol{A}$$ Eqn. 1

where the growth rate of the animal population increases proportionally to its consumption

rate of plant rewards, C_R. Parameter ε is the efficiency of converting rewards to new animal

individuals via birth and maturation. We choose total consumption rate $C_R = \frac{a\boldsymbol{A}\boldsymbol{P}}{1+ah\boldsymbol{P}}$,

following a Holling Type II functional response with attack rate a and handling time h.

This encodes the reasonable assumption that animal consumption rate on plant rewards

saturates with increasing density of the plant population. Reproduction fueled by resources

other than plant rewards is included in b_A. We define an animal population ($\boldsymbol{A}$) as an

"obligate" mutualist of the plant population ($\boldsymbol{P}$) when $r_A = b_A - d_A \leq 0$, that is, when $\boldsymbol{A}$

cannot persist in the absence of its mutualist plant. Otherwise ($r_A > 0$), $\boldsymbol{A}$ is "facultative"

and its population is self-sustaining.

Population dynamics of plants

Plants receive reproductive services (S) from animals which increases the density of

mature (reproductive) plant individuals. Reproductive services are functions of animal

visitation to plants, which we assume to be well approximated by the foraging rate on

rewards (C_R, Vázquez *et al*. 2005). Our pollination and seed-dispersal models both account

for germination, g, and negative density-dependence, s_P, that limit the seeds that survive

and mature to reproductive adults. This density-dependence represents, for example, natural

enemies near a parent plant making the habitat inhospitable for the survival of seedlings

(Janzen-Connell effect) or seeds competing during recruitment. Below, we derive separate

models for pollination and seed dispersal based on the different mechanisms by which each

affects the population dynamics of plants.

Pollination: Increase in seed set

We model animal-pollinated plant dynamics ($\boldsymbol{P}$) as:

149
$$\frac{d\boldsymbol{P}}{dt} = \boldsymbol{P}[g(b_P + \beta S)(1 - s_P\boldsymbol{P}) - d_P] \qquad \text{Eqn. 2}$$

where β is the conversion efficiency of pollination services (S) to seed production.

Pollination requires the transfer of pollen from one individual's flower to the stigma of a

conspecific plant individual, which occurs when animals visit flowers to forage. We

therefore assume $S = C_R$, where the contribution of the animal population to per-plant seed

set is the total foraging rate of animals on plant rewards (C_R). This expression accounts for

the repeated interactions between plant and animal individuals required for conspecific

pollen transfer (Vázquez *et al.* 2005, Schupp *et al.* 2017). Using the Holling Type II

functional response for C_R from above yields our specific model for animal-pollination

mutualisms:

159
$$\begin{cases} \frac{dP}{dt} = P\left[\left(b_P + \beta\frac{aAP}{1+ahP}\right)g(1-s_PP) - d_P\right] \\ \frac{dA}{dt} = A\left[b_A + \varepsilon\frac{aP}{1+ahP} - s_AA - d_A\right] \end{cases}$$
Eqn. 3

Following our previous definitions, plant population (P) is an obligate mutualist of A when

it cannot persist in its absence ($r_P = g_Pb_P - d_P \le 0$); otherwise, P is facultative ($r_P > 0$).

Seed dispersal: Reduction in negative density-dependence or increase in germination

Seed dispersers visit plants to forage on fruit, elaiosomes, or even seeds directly,

later depositing the seed away from the parent plant. This process increases recruitment by

reducing density-dependent mortality caused by predators or pathogens, which are most

abundant near adult plants (Wotton & Kelly 2011, Fricke *et al.* 2013, Moore & Dittel

2020). Seed dispersal may also increase germination by improving seed condition during

passage through dispersers' guts, which removes pathogens or provides "chemical

camouflage" from natural enemies (Fricke *et al.* 2013). We consider both cases below.

First, animals rescue seeds from predators, pathogens, and intraspecific competition

by transporting them away from parent plants. This can increase successful recruitment:

172
$$\frac{dP}{dt} = P[b_Pg(1-(s_P-\sigma S)P) - d_P]$$
Eqn. 4

where σ converts dispersal services into reductions in negative density-dependence (s_P).

Eqn. 4 implies that plants must be potentially persistent (have positive demographic

rates, $r_P = b_P g - d_P > 0$) to benefit from reductions to negative density-dependence.

Therefore, we only consider facultative plants ($r_P > 0$) for this case.

Second, animals may increase seed condition by masticating or digesting fruits.

This can improve germinability or germination rate:

$$\frac{dP}{dt} = P[b_P(g + \gamma S)(1 - s_P P) - d_P]. \qquad \text{Eqn. 5}$$

where γ converts dispersal services into increased germination g. Plants may be obligate

($r_P \leq 0$) or facultative ($r_P > 0$) upon their animal partner for population persistence.

In both seed dispersal cases, services are directly related to consumption rate of

animals on plant rewards. We use $S = C_R/P$, where the per-plant benefit of seed dispersal

is proportional to the per-capita (i.e., per-plant) visitation rate of animals on plants. This is

because repeated interactions between plant and animal individuals are possible but not

required for effective seed dispersal, as it is for effective pollination (Schupp *et al.* 2017).

Using the Holling Type II functional response from above yields our specific models for

animal seed-dispersal mutualisms in which plants benefit via escape from negative density-

dependence:

$$\begin{cases} \frac{dP}{dt} = P\left[b_P g \left(1 - \left(s_P - \sigma \frac{aA}{1+ahP}\right)P\right) - d_P\right] \\ \quad \frac{dA}{dt} = A\left[b_A + \varepsilon \frac{aP}{1+ahP} - s_A A - d_A\right] \end{cases} \qquad \text{Eqn. 6}$$

191   or increased germination:

$$\begin{cases} \frac{dP}{dt} = P\left[b_P\left(g + \gamma\frac{aA}{1+ahP}\right)(1 - s_P P) - d_P\right] \\ \quad\frac{dA}{dt} = A\left[b_A + \varepsilon\frac{aP}{1+ahP} - s_A A - d_A\right] \end{cases}$$

Eqn. 7

193   Notice that Eqn. 7 is only biologically reasonable when $s_P - \sigma S \geq 0$, which is not

194 guaranteed when the ratio of animal to plant density is high. We restrict our analysis to the

195 space in which this condition holds, though our results are qualitatively the same outside of

196 this regime as well.

197 **Costs of mutualism**

198   The potential costs of pollination and seed dispersal mutualisms on plant

199 populations are less clear than their benefits. Plants tend to provide nutritional rewards such

200 as nectar or fruits to attract their animal transporters of pollen or seeds. The available

201 evidence suggests that the costs of providing these rewards tend to suppress seed set or

202 other aspects of an individual plant's reproduction (Edwards *et al.* 2006, Willmer 2011).

203 However, these individual-level costs are likely to be buffered at the population level for

204 several reasons. First, costs associated with rewards can simply act to reduce benefit

205 provided by mutualism (e.g., reduced seed set due to nectar production, Brandenburg *et al.*

206 2012; seed predation by scatterhoarders, Gómez *et al.* 2019). Second, rewards and the

207 organs that produce them are typically ephemeral in comparison to individuals in the

208 population. These organs can be damaged or decay, but these costs are unlikely to

209 instantaneously result in death or damage to individuals (Revilla 2015). More generally,

210 costs associated with rewards may be "fixed" in that they are independent of partner density

211 (e.g., sunk costs of molecular machinery used to produce rewards, Holland *et al.* 2002).

Finally, changes in metrics of individual fitness like fecundity do not necessarily manifest

in changes in the per-capita growth rate of the population (Ford *et al.* 2015). In summary,

rewards costs, even if highly impactful to individuals or a certain vital rate, may not have a

measurable impact on population growth. Therefore, we assume that costs associated with

rewards do not substantially impact population density or, if they do, the impact could be

represented as small, fixed changes to plants' per-capita birth and death rates approximated

as constant with respect to the producing species' density.

Analysis

We performed phase plane analyses on our pollination (Eqn. 3) and seed dispersal

models (Eqns. 5, 6) using Wolfram Mathematica 11. Mathematical analysis and simulation

code are presented in Appendix A and online, respectively. In the main text, we only

present results for the ecologically relevant cases where populations have positive densities

and can potentially coexist (i.e., are feasible). We investigated the behavior of populations

at low density, assessing our models for conditions under which Allee effects, thresholds,

and other alternative stable states occur. We use the term "Allee effects" for strong,

demographic Allee effects, in which a population declines under a certain threshold of its

own density (Kramer *et al.* 2009). We define "threshold effects" as population declines

caused by the decline of its partner's density under a certain threshold (Vandermeer &

Boucher 1978, Revilla 2015). Alternative stable states, such as single-species persistence or

bistability, occur when a system can settle stably into more than one equilibrium depending

only upon initial conditions ("historical accidents," May 1977).

Results

We found that pollination and seed dispersal mutualisms have similar dynamics and

stability at high density but differ in the ecological conditions under which the populations

are vulnerable to collapse. Different parameter regimes, in combination with whether each

partner is obligate or facultative, lead to different dynamics at low population densities.

Below, we describe these different dynamics for each of our models, assuming that

coexistence is feasible (but see Appendix A for results on feasibility). Then, we describe

some ecological scenarios under which each outcome is likely to occur. We provide these

scenarios rather than providing formal mathematical conditions, because the latter are

complex and difficult to interpret due to the nonlinearities in our models.

Pollination

The shape and intersections of the plant and animal "nullclines" (curves of zero

change in population density) determine the dynamics of the system. The animal nullclines

(black curves) are concave down, increasing functions that saturate with respect to plant

density (Figs. 1-3). The animal-pollinated plant nullclines (green curves) differ according to

whether the plant is obligate or facultative (Fig. 1). When obligate (Fig. 1A-B), the plant

nullcline is U-shaped bounded between vertical asymptotes at zero and at $1/s_P$. When

facultative (Fig. 1C-G), the plant nullcline is increasing and concave up at high densities.

Depending on the parameter regime, the plant nullcline may exhibit an inflection point at

low densities in which it transitions from concave down to up (e.g., Fig. 1D). When

facultative, animals or plants may persist in the absence of their partners at density $K_A =$

r_A/s_A or $K_P = r_P/b_P g s_P$, respectively. These densities are "carrying capacities" in the

original sense of single-species equilibrium densities (Vandermeer & Boucher 1978).

When plants are obligate (Fig. 1A-B), the system exhibits two equilibria: (1) A

stable coexistence equilibrium (off-axes filled circle). Density past this equilibrium in either

species causes its population to decrease due to negative density-dependence. Therefore,

unbounded population growth cannot occur. (2) A half-stable saddle point that attracts in

one dimension but repels in the other (off-axes hollow circle), bisecting the plane into two

portions, as marked by the "separatrix" (dashed line). This separatrix marks out a threshold

under which obligate partners go extinct even if initially highly abundant ("threshold

effects," regions shaded light red). For example, following a trajectory from the lower-right

region in Fig. 1B, initially highly abundant obligate plants go extinct while initially rare

pollinators persist at K_A. These threshold effects occur because one species is too low in

density to provide sufficient benefits to its partner, causing the partner's population to

decline. The low-density species continues to benefit from mutualism but its increase in

density cannot occur fast enough to save the system from extinction. Above the separatrix,

one or both species are of high enough density that benefits from mutualism cause positive

population growth in their partners and the system will achieve stable coexistence.

Obligate animal-pollinated plants are additionally susceptible to Allee effects

(regions shaded dark red), where their population declines under a threshold of their own

density regardless of the density of their partner (Fig. 1A-B). This occurs because benefit is

proportional to the total consumption rate by animals, i.e., plants require obligate

outcrossing for successful pollination behavior. At very low plant density, even highly

abundant pollinators are limited by the time required to harvest and (incidentally) transfer

pollen between two plant individuals, and thus cannot provide pollination services at a rate

sufficient to allow plant population growth. Unsurprisingly, animals cannot acquire enough

food and decline either to extinction (Fig. 1A) or to K_A once plants are fully extirpated (Fig.

1B).

When animal-pollinated plants are facultative (Fig. 1C-G), more diverse dynamics

are possible, depending on the relative growth rates and efficiency of plants and animals as

mutualists. High-density stable coexistence always occurs if partners are originally at high

enough density. At lower density, threshold effects can also occur when animals are

obligate mutualists (Fig.1 C) or even population growth from low densities (Fig. 1D-G)

with the potential for bistable coexistence (Fig. 1D, 1F).

When both partners exhibit population growth from low densities (Fig. 1D-G),

stable coexistence at higher densities than either species could achieve alone will always

occur. However, in a relatively small parameter range, the system exhibits three equilibria:

low-density stable, half-stable saddle, and high-density stable (Fig. 1D, 1F). Thus, there are

two possible points of stable coexistence – one at lower density and one at higher – to

which the system will be attracted, based upon initial conditions ("bistable coexistence"). A

separatrix (dashed line) running through the saddle point (hollow circle) marks out a

threshold under which the system will persist at the lower-density coexistence equilibrium.

Growth from low density (Fig. 1D-G) can occur when $-\frac{r_A}{a(\varepsilon+hr_A)} < K_P$. That is,

when the intercept of the animal nullcline in the x-axis (representing plant density) is higher

than the intercept of the plant nullcline. In this case, bistable coexistence (Fig. 1D, 1F) is

most likely when plants have low carrying capacity, K_P. That is, when the environment is

hostile to plants due to, for example, high density-dependence (s_P) or density-independent

mortality (d_P). In addition, bistable coexistence is most likely when pollinators are

inefficient at accessing nutrients from rewards when foraging upon plants (low a, low ε,

high h) and have near-zero population growth rate (r_A) in the absence of plants, regardless

of whether they are obligate or facultative. The single stable coexistence equilibrium at

high density is most likely under the opposite conditions (Fig. 1E, 1G). That is, when plants

have high carrying capacities and pollinators are efficient foragers. Finally, threshold

effects can occur instead (Fig. 1C) when $-\frac{r_A}{a(\varepsilon+hr_A)} > K_P$, that is, when the intercept of the

animal nullcline is lower than the intercept of the plant nullcline in the x-axis. This happens

when pollinators are highly obligate upon plants ($r_A \ll 0$). Threshold effects are also more

likely when pollinators are efficient foragers (high a, high ε, low h) but the environment is

hostile to plants (low K_P).

311   *Seed Dispersal*

Our seed dispersal model with benefit to plants as reduced negative density-

dependence displays a range of low-density outcomes (Fig. 2) like those described for

pollination mutualisms. Seed dispersers follow the same dynamics as pollinators. Plants are

always facultative (see Methods) and can persist in the absence of animal mutualists at

density K_P. The plant nullcline is a concave down, increasing function with linear slope of

$\frac{hs_P}{\sigma} \geq h$ at high densities. Therefore, based on the plant nullcline alone, benefits to plants

would increase indefinitely with increasing animal density. However, animal visitation rate

saturates due to satiation (encoded in the Holling type II functional response) and this

animal saturation bounds the benefits to plants by intersecting the plant nullcline at high

densities. At low densities, the plant nullcline can exhibit an inflection point in which it

transitions from concave down to linear growth. The same qualitative conditions hold as

described for the pollination model as to whether the system exhibits bistable coexistence

(Fig. 2B, 2D), a single high-density stable coexistence equilibrium (Fig. 2C, 2E), or

threshold effects when animals are obligate mutualists (Fig. 2A). Note that as negative

density dependence increases due to the Janzen-Connell effect or intraspecific competition

(high s_P), plant carrying capacity shifts to lower density, making the cases in which

threshold effects or bistable coexistence can occur more likely. Conversely, increasing the

effectiveness of escape from negative density-dependence due to seed dispersal (high σ)

makes the single high-density coexistence case more likely.

Our seed dispersal model with benefit to plants as increased seed germination displays

simpler dynamics than the other two models. The plant nullcline for both obligate and

facultative plants is a concave up, increasing function, bounded by a vertical asymptote at

$1/s_P$. Animal-seed dispersers follow the same dynamics as animal-pollinators, described

above. Like the pollination model, stable coexistence at higher densities than either can

achieve alone is always possible (Fig. 3A-F). When at least one partner is an obligate

mutualist, threshold effects can always occur, leading to extinction if at least one partner is

at low enough density (Fig. 3A, B, D). These effects are most likely when one partner is

highly obligate ($r_i \ll 0$) and when plants are difficult to find on the landscape (search time

$(1/a) \to \infty$). Specifically, they can occur in feasible systems when $K_A < -\frac{r_P}{b_P \gamma a}$ or

$-\frac{r_A}{a(\varepsilon + h r_A)} > K_P$. Alternatively, growth from low density and ultimate stable coexistence is

the only outcome when both partners are facultative (Fig. 3F).

Discussion

Pollination and seed dispersal mutualisms support the reproduction of a vast number

of plant species globally, including crops (Klein *et al.* 2007, Aslan *et al.* 2013, Garibaldi *et*

al. 2013, Neuschultz *et al.* 2016). Anthropogenic stressors threaten these mutualisms by

causing population declines that can disrupt the benefits provided by the mutualists

(Tylianakis *et al.* 2008, Traveset & Richardson 2006, 2014, Potts *et al.* 2010, 2016,

Beckman *et al.* 2020). These potential disruptions increase the need to study the dynamics

of mutualisms when the interacting populations exhibit low density. Historically, however,

theoretical studies have focused on the dynamics of mutualisms at high density. Authors

grappled with the unbounded growth predicted by the simplest models of mutualism

(Robert May's (1976) "orgy of mutual benefaction") and searched for mechanisms that

would lead to stable coexistence (Wolin & Lawlor 1984). This research found that the

saturation of mutualistic benefit stabilizes the population dynamics of mutualistic

interactions at high density (Holland *et al.* 2006, Okuyama & Holland 2008). Theoretical

research should now focus on the dynamics of mutualisms at low density, especially given

the global population declines of pollinators and seed dispersers.

We investigated the dynamics, especially at low density, of novel consumer-

resource models that incorporate the mechanisms by which pollinators and seed-dispersers

benefit plant populations. We found thresholds, Allee effects, and bistable coexistence, and

characterized when each of those dynamics occur. Thresholds occur when mutualists are

obligate because low-density populations cannot provide enough benefit to rescue their

higher density partner (Figs. 1A-C, 2A, 3A-B, D). Allee effects occur in obligate animal-

pollinated plants because the plants can become too low in density to attract sufficient

pollinator visits to save their own population (Figs. 1A-B). Bistable coexistence occurs in

pollination (Figs. 1D, F) and seed dispersal mutualisms when plants benefit through a

reduction in negative density-dependence during recruitment (Figs. 2B, D). In both

mutualisms, bistable coexistence occurs when plants are facultative but have low carrying

capacity while animals are inefficient foragers with near-zero per-capita growth rate.

Conversely, when facultative plants have high carrying capacity and animals are efficient

foragers, plant and animal populations grow from low density with subsequent stable

coexistence at a single, high-density equilibrium (Figs. 1E, 1G, 2C, 2E, 3C, 3E). This high-

density equilibrium is guaranteed for facultative seed dispersal mutualisms when animals

benefit plants via increased seed germination (Fig. 3F).

Threshold effects in mutualisms were first described in the late 1970s, specifically

in cases of obligate mutualists that strongly benefit each other (May 1976, Vandermeer and

Boucher 1978). The strong mutualistic interaction allowed the species to persist at high

densities, but was not sufficient for coexistence at densities below the threshold. The

authors also found that obligate mutualists interacting weakly always go extinct, that is,

thresholds effects do not occur in those mutualisms. This research used the Lotka-Volterra

model for mutualism with positive interacting coefficients and linear functional responses,

but the same qualitative results at low density are found when saturation is added to the

functional responses (Vandermeer & Goldberg 2013). Indeed, models incorporating

consumer-resource mechanisms also find these thresholds (May 1976, Soberón & Martinez

del Rio 1981, Wells 1983, Wright 1989, Fishman & Hadany 2010). Most recently, Revilla

(2015) predicts threshold effects in obligate plants, in which plants at low density cannot

recover fast enough to save their animal partners from decline, even if the animal

population was originally at very high density. Finally, Valdovinos and Marsland (2021)

identify the quality of visits needed from pollinators for plants to persist. Below such

threshold, the plant species and the animals depending on those plants go extinct. Here, we

also find that threshold effects occur when at least one mutualist is obligate and when the

benefits among species (in terms of efficient foraging) are strong enough to sustain

populations at high densities.

Allee effects may also cause irrecoverable declines in populations at low density.

We find Allee effects in plant populations that are obligate mutualists of pollinators, due to,

e.g., high self-incompatibility. This result is consistent with pollination ecology because

pollinators' benefits to plants tend to decrease at low plant density due to increased search

time, leading to a decrease in visitation rate and pollen limitation (Forsyth 2003). Low plant

density may also decrease visitation rate due to insufficient floral displays or reduce visit

quality due to pollen dilution (Forsyth 2003). Indeed, pollinator-mediated Allee effects

have been observed empirically in highly self-incompatible plant species (Kramer *et al.*

2009). We show that reductions in visit quantity alone are sufficient to induce an Allee

effect, whereas natural populations likely experience both a decrease in visit quality and

quantity at low plant density. In contrast, we did not find Allee effects in our seed dispersal

models. This is consistent with the mechanisms by which seed dispersers benefit plants.

Seed dispersers benefit plants by increasing the seed germination or their survival to

adulthood through refuge from intraspecific competition and natural enemies (seed

predators, herbivores) that are attracted to regions of high food densities (i.e., the Janzen-

Connell effect, Janzen 1970, Connell 1971, Wotton & Kelly 2011, Fricke *et al.* 2013,

Moore & Dittel 2020). These plants are thus less likely to experience Allee effects, because

at low density the primary threats are similarly reduced. However, plants that are obligate

upon animal dispersers for germination can experience Allee effects in our models when

animals follow a Type III functional response (Appendix A); this is because plants become

too rare to attract animals that are less active at low food density (also see Courchamp *et al.*

2018). Finally, Holland and DeAngelis (2010) predict an Allee effect in "unidirectional"

consumer-resource mutualisms, which include both animal-pollination and seed dispersal

interactions. However, their Allee effect occurs due to animal overexploitation of plant

rewards, which induces more costs to plants from supplying rewards than benefits, though

they comment that this is unlikely to lead to the extinction of seed dispersal systems.

Bistable coexistence, where partners coexist stably either at low or high density,

depending upon if at least one population dips below a certain threshold of density, is, to

our knowledge, a novel result. Ecologically, bistable coexistence results from the

dependence on plant density caused by the required transfer of pollen between conspecific

plants in pollination or refuge from threats at high conspecific density in seed dispersal

mutualism. Mathematically, it results from the change in concavity in the plant nullclines of

both models. As a result, bistable coexistence can also occur in our germination model

when seed dispersers follow a Holling Type III functional response. In such case, the

animal nullcline inflects because animals consume faster at medium, compared to low or

high, plant densities (Appendix A). Bistable coexistence occurs when plants have low

carrying capacity (K_P) and animals have near-zero growth rate and low foraging efficiency,

particularly due to low "attack rate" (a) on plants' rewards. High search times for plants

($1/a$) can cause low attack rate, potentially because the plant is not very visible to the

pollinator on the landscape, or because pollinators have low preference for that plants'

rewards (Valdovinos *et al.* 2013, 2016). Bistable coexistence is probably less likely to

occur in natural systems than our stable coexistence or threshold effects scenarios, in which

pollinators have higher a and plants have high or low K_P, respectively. However,

anthropogenic perturbations leading to population decline, as well as spatial and temporal

mismatches between compatible partners, may increase the occurrence of bistable

coexistence.

Overall, we find that obligate mutualisms are most at risk for experiencing

destabilizing thresholds, which may have cascading effects on the rest of the ecological

community (Latty & Dakos 2019). The existence of thresholds suggests that environmental

change could shift a pollination system from above a separatrix to below it. This will lead

to collapse of the system under certain conditions, but in very special cases, the system may

instead shift to a lower stable coexistence equilibrium, and still exhibit positive growth at

low density. Though this is a preferable outcome to deterministic extinction, these

dynamics still allow the system to reach very low population densities, and thus to be more

vulnerable to environmental or demographic stochasticity. However, the highly stable

outcomes for facultative mutualists, and particularly for facultative plants with high

carrying capacity, suggest that these species may buffer communities from perturbations to

low density in a network setting (also see van der Heide *et al.* 2020). This is because

mutualistic networks are typically highly nested, which means that generalist species tend

to interact with both generalist and specialist species but specialist species with only

generalists (Bascompte *et al.* 2003). Then, a nested network with generalist plants (by

definition facultative to each of their animal mutualists), that also have sufficiently high

carrying capacity, may not suffer threshold effects caused by rare, obligate animal

populations. In other words, those generalist plants with high carrying capacity may make

the network robust to potential threshold effects.

A unique aspect of our modelling approach is that we specify whether plants benefit

according to animal total or per-capita visitation rate. Our choice for pollination benefits to

plants to scale with animal total visitation rate accounts for outcrossing (Vázquez *et al.*

2005). This, to the best of our knowledge, makes our pollination model unique in the

literature of two-species models (but see Valdovinos *et al.* 2013, 2016, Hale *et al.* 2020 for

network models). In addition, visitation in our models differs from previous work (e.g.,

Revilla 2015) because we use nonlinear (specifically, Holling Type II) saturating functional

responses for consumption rate. This allows both plants and animals to experience intra-

specific competition for benefits, set by the maximum time animals can spend foraging on

rewards. We could follow previous work (Revilla 2015) and impose a direct limitation to

plant benefits, so that they saturate due to limited ovules or seeds (Appendix B). This does

not qualitatively change our results, except for the negative density-dependence seed

dispersal model, which displays the simpler results of our germination model (i.e., no

bistable coexistence).

Our qualitative results (Figs. 1-3) are generally robust to varying functional forms

from Type I (linear) to Type III (Appendix A). The exceptions are our negative density-

dependence seed dispersal model, in which unbounded population growth can occur when

dispersers follow a linear functional response, and our germination seed dispersal model

which has similar dynamics to our pollination model when dispersers follow a Type III

functional response. Our models can also accommodate functional responses interpreted as

"net benefit" curves (e.g., Holland *et al.* 2002, Morris *et al.* 2010) if costs and benefits

affect the same vital rate. That is, if "costs" simply reduce the benefits that accrue to a

given vital rate (e.g., Brandenburg *et al.* 2012). Under this interpretation, unimodal

functional forms may arise (e.g., Morris *et al.* 2010) which could lead to substantially

different dynamical predictions than those presented here. Another of our unique

assumptions is that negative density-dependence specifically reduces the potential plant

"births" that mature to reproductive adults. Our models exhibit qualitatively similar

dynamics if we instead assume that negative density-dependence limits overall per-capita

growth (using an additive term) as in previous works (e.g., Holland & DeAngelis 2010,

Revilla 2015). Finally, our models, as well as previous ones, assume that mutualisms have

population-level impacts. Most empirical studies, however, quantify the benefits and costs

of mutualisms at the individual level in terms of fitness or even by using a single proxy for

fitness (Bronstein 2001, Ford *et al.* 2015). Those effects do not necessarily imply

population- or community-level impacts of mutualism (Flatt & Weisser 2000, Ford *et al.*

2015). Therefore, empirical work on population dynamics of mutualisms is of foremost

importance to evaluate historical and current ecological theory on mutualisms.

Conclusion

This research increased mechanistic understanding of the dynamics of plant-animal

mutualisms by analyzing consumer-resource models that incorporate simple mechanisms of

reproductive benefits to plants. We found that these mutualisms may be vulnerable to

declines at low density due to Allee and threshold effects and identified bistable

coexistence in some cases. We also characterized the ecological scenarios under which

those dynamical behaviors are most likely to occur, which has been recognized a

conservation priority (Latty & Dakos 2019). Future work should continue to investigate

how different mechanisms of mutualism manifest in different population dynamics at low

density, both to advance ecological understanding and to aid in conservation objectives.

References

Addicott, J. F. 1981. Stability properties of 2-species models of mutualism: Simulation

studies. Oecologia 49:42–49.

Aslan, C. E., E. S. Zavaleta, B. Tershy, and D. Croll. 2013. Mutualism disruption Threatens

global plant biodiversity: A Systematic Review. PloS ONE 8.

Bascompte, J., Jordano, P., Melian, C.J. and Olesen, J.M. 2003. The nested assembly of

plant-animal mutualistic networks. Proceedings of the National Academy of Sciences USA

100:9383–9387.

Bascompte, J., P. Jordano, and J. M. Olesen. 2006. Asymmetric coevolutionary networks

facilitate biodiversity maintenance. Science 1:3-5.

Bastolla, U., M. A. Fortuna, A. Pascual-García, A. Ferrera, B. Luque, and J. Bascompte.

2009. The architecture of mutualistic networks minimizes competition and increases

biodiversity. Nature 458:1018-1020.

Beckman, N. G., C. E. Aslan, H. S. Rogers, O. Kogan, J. L. Bronstein, J. M. Bullock, F.

Hartig, et al. 2020. Advancing an interdisciplinary framework to study seed dispersal

ecology. AoB PLANTS 12:1–18.

Bronstein, J. L. 2001. The costs of mutualism. American Zoologist 41:825–839.

Bronstein, J. L. 2015. The study of mutualism. Pages 3-19 *in* J. L. Bronstein, ed.

Mutualism. Oxford University Press, Oxford.

Courchamp, F., L. Berec, and J. Gascoigne 2008. Population dynamics: modelling

demographic Allee effects. *In* Allee Effects in Ecology and Conservation, Oxford

University Press, Oxford.

Edwards, W., M. Dunlop, and L. Rodgerson. 2006. The evolution of rewards: Seed

dispersal, seed size and elaiosome size. Journal of Ecology 94:687–694.

Fishman, M. A., and L. Hadany. 2010. Plant-pollinator population dynamics. Theoretical

Population Biology 78:270–277.

Flatt, T., and W. W. Weisser. 2000. The effects of mutualistic ants on aphid life. Ecology

81:3522–3529.

Ford, K. R., J. H. Ness, J. L. Bronstein, and W. F. Morris. 2015. The demographic

consequences of mutualism: ants increase host-plant fruit production but not population

growth. Oecologia 179:435–446.

Forsyth, S. A. 2003. Density-dependent seed set in the Haleakala silversword: Evidence for

an Allee effect. Oecologia 136:551–557.

Fricke, E. C., M. J. Simon, K. M. Reagan, D. J. Levey, J. A. Riffell, T. A. Carlo, and J. J.

Tewksbury. 2013. When condition trumps location: Seed consumption by fruit-eating birds

removes pathogens and predator attractants. Ecology Letters 16:1031–1036.

Gause, G. F., and A. A. Witt. 1935. Behavior of mixed populations and the problem of

natural selection. American Naturalist 69:596–609.

Goh, B. S. 1979. Stability in models of mutualism. American Naturalist 113:261–275.

Gómez, J. M., E. W. Schupp, and P. Jordano. 2019. Synzoochory: the ecological and

evolutionary relevance of a dual interaction. Biological Reviews 94:874–902.

Hale, K. R. S., F. S. Valdovinos, and N. D. Martinez. 2020. Mutualism increases diversity,

stability, and function of multiplex networks that integrate pollinators into food webs.

Nature Communications 11.

Holland, J. N., and D. L. DeAngelis. 2010. A consumer-resource approach to the density-

dependent population dynamics of mutualism. Ecology 91:1286–1295.

Holland, J. N., D. L. DeAngelis, and J. L. Bronstein. 2002. Population dynamics and

mutualism: Functional responses of benefits and costs. American Naturalist 159:231–244.

Holland, J. N., Okuyama, T., & DeAngelis, D. L. 2006. Comment on "Asymmetric

coevolutionary networks facilitate biodiversity maintenance." Science, 313:1887–1887.

Janzen, D. H. 1970. Herbivores and number of tree species in tropical forests. American

Naturalist 104:501–508.

Jordano P. 2014. Fruits and frugivory. Pages 125–166 *in* R. S. Gallagher, ed. Seeds: The

ecology of regeneration in plant communities. CABI, Boston.

Klein, A. M., B. E. Vaissière, J. H. Cane, I. Steffan-Dewenter, S. A. Cunningham, C.

Kremen, and T. Tscharntke. 2007. Importance of pollinators in changing landscapes for

world crops. Proceedings of the Royal Society B: Biological Sciences 274:303–313.

Kramer, A. M., B. Dennis, A. M. Liebhold, and J. M. Drake. 2009. The evidence for Allee

effects. Population Ecology 51:341–354.

Latty, T., and V. Dakos. 2019. The risk of threshold responses, tipping points, and

cascading failures in pollination systems. Biodiversity and Conservation 28:3389–3406.

May, R. M. 1976. Theoretical Ecology, Principles and Applications. Blackwell Scientific,

Oxford.

May, R. M. 1977. Thresholds and breakpoints in ecosystems with a multiplicity of stable

states. Nature 269:471–477.

Moore, C. M., and J. W. Dittel. 2020. On mutualism, models, and masting: The effects of

seed-dispersing animals on the plants they disperse. Journal of Ecology 108:1775–1783.

Morris, W. F., D. P. Vázquez, and N. P. Chacoff. 2010. Benefit and cost curves for typical

pollination mutualisms. Ecology 91:1276–1285.

Neuschulz, E. L., T. Mueller, M. Schleuning, and K. Böhning-Gaese. 2016. Pollination and

seed dispersal are the most threatened processes of plant regeneration. Scientific Reports

6:6–11.

Okuyama, T., and J. N. Holland. 2008. Network structural properties mediate the stability

of mutualistic communities. Ecology Letters 11:208–216.

Ollerton, J., R. Winfree, and S. Tarrant. 2011. How many flowering plants are pollinated by

animals? Oikos 120:321–326.

Pierce, N. E., and W. R. Young. 1986. Lycaenid butterflies and ants: Two-species stable

equilibria in mutualistic, commensal, and parasitic Interactions. American Naturalist

128:216–227.

Potts, S. G., J. C. Biesmeijer, C. Kremen, P. Neumann, O. Schweiger, and W. E. Kunin.

2010. Global pollinator declines: Trends, impacts and drivers. Trends in Ecology and

Evolution 25:345–353.

Potts, S. G., V. Imperatriz-Fonseca, H. T. Ngo, M. A. Aizen, J. C. Biesmeijer, T. D.

Breeze, L. v. Dicks, et al. 2016. Safeguarding pollinators and their values to human well-

being. Nature 540:220–229.

Revilla, T. A. 2015. Numerical responses in resource-based mutualisms: A time scale

approach. Journal of Theoretical Biology 378:39–46.

Schupp, E. W., P. Jordano, and J. M. Gómez. 2017. A general framework for effectiveness

concepts in mutualisms. Ecology Letters 20:577–590.

Soberón, J. M., and C. Martinez del Rio. 1981. The dynamics of a plant-pollinator

interaction. Journal of Theoretical Biology 91:363–378.

Stachowicz, J. J. 2001. The structure of ecological communities. BioScience 51:235–246.

Traveset, A. and Richardson, D. M. 2006. Biological invasions as disruptors of plant

reproductive mutualisms. Trends in Ecology and Evolution 21:208–216.

Traveset, A. and Richardson, D. M. 2014. Mutualistic interactions and biological

invasions. Annual Review of Ecology, Evolution, and Systematics 45:89–113.

Tylianakis, J. M., R. K. Didham, J. Bascompte, and D. A. Wardle. 2008. Global change and

species interactions in terrestrial ecosystems. Ecology Letters 11:1351–1363.

Valdovinos, F. S. 2019. Mutualistic networks: moving closer to a predictive theory.

Ecology Letters 22:1517–1534.

Valdovinos, F.S., Moisset de Espanes, P., Flores, J.D. & Ramos-Jiliberto, R. 2013.

Adaptive foraging allows the maintenance of biodiversity of pollination networks. Oikos

122:907–917.

Valdovinos, F. S., B. J. Brosi, H. M. Briggs, P. Moisset de Espanés, R. Ramos-Jiliberto,

and N. D. Martinez. 2016. Niche partitioning due to adaptive foraging reverses effects of

nestedness and connectance on pollination network stability. Ecology Letters 19:1277–

1286.

Valdovinos, F.S., and Marsland, R. 2021. Niche theory for mutualism: A graphical

approach to plant-pollinator network dynamics. American Naturalist 197:393–404.

Van der Heide, T., C. Angelini, J. de Fouw, and J. S. Eklöf. 2021. Facultative mutualisms:

A double-edged sword for foundation species in the face of anthropogenic global change.

Ecology and Evolution 11:29–44.

Vandermeer, J. H., and D. H. Boucher. 1978. Varieties of mutualistic interaction in

population models. Journal of Theoretical Biology 74:549–558.

Vandermeer, J. H., and Goldberg, D. E. 2013. Pages 225-238 *in* Population ecology: First

principles. Princeton University Press, Princeton.

Vázquez, D. P., W. F. Morris, and P. Jordano. 2005. Interaction frequency as a surrogate

for the total effect of animal mutualists on plants. Ecology Letters 8:1088-1094.

Wells, H. 1983. Population equilibria and stability in plant-animal pollination systems.

Journal of Theoretical Biology 100:685–699.

Whittaker, R. H. 1975. Communities and Ecosystems. 2d ed. Macmillan, New York.

Willmer, P. 2011. Pollination and Floral Ecology. Princeton University Press, Princeton.

Wolin, C. L., and L. R. Lawlor. 1984. Models of facultative mutualism: Density effects.

American Naturalist 124:843–862.

Wolin, C.L. 1985. The population dynamics of mutualistic systems. Pages *in* 248–269. D.

H. Boucher, ed. The Biology of Mutualism. Oxford University Press, Oxford.

Wotton, D. M., and D. Kelly. 2011. Frugivore loss limits recruitment of large-seeded trees.

Proceedings of the Royal Society B: Biological Sciences 278:3345–3354.

Wright, D. H. 1989. A simple, stable model of mutualism incorporating handling time.

American Naturalist 134:664–667.

Zhou, Y., C. Newman, J. Chen, Z. Xie, and D. W. Macdonald. 2013. Anomalous, extreme

weather disrupts obligate seed dispersal mutualism: Snow in a subtropical forest ecosystem.

Global Change Biology 19:2867–2877.

Legends

Figure 1.

Phase plane diagrams for plant-pollinator mutualisms. Nullclines are curves of zero growth for plant (green) or animal pollinator (black) populations. Equilibria (dots) occur at the intersection of plant and animal nullclines. Filled black equilibria are stable attractors, filled red equilibria are unstable repellers, and hollow equilibria are half-stable saddle points. Arrows show the directions of population change for plants (x-axis) and animals (y-axis), with lighter colors indicating a faster rate of change. Dynamics depend upon the parameterization of the model and whether plants and animals are obligate (O) or facultative (F) mutualists (rows). When plants are obligate (**A, B**), they experience Allee effects at low density (dark red shaded region). Plants cannot attract sufficient pollinator visitation to support their own population growth, leading to collapse. When either plants or animals are obligate, both partners experience threshold effects (light red shaded region). Below the threshold marked by the separatrix (dashed line), one population's density is too low to support its partner's growth, resulting in system collapse to extinction (**A**), animal-only persistence at $K_A = (b_A - d_A)/s_A$ (**B**), or plant-only persistence at $K_P = (b_P g - d_P)/b_P g s_P$ (**D**). Thus, depending upon the initial densities of both partners, the mutualism may persist stably or collapse. When one partner is facultative, stable coexistence may be the only outcome (**C, E, G**). However, in a small parameter space, bistable coexistence may occur (**D, F**). Here, the separatrix divides the region in which the system will be attracted to the low- or high-density stable coexistence equilibrium, depending upon whether initial densities are below or above the separatrix, respectively. Parameter values are fixed to the

following: $a = 0.6, h = 0.75$; plants: $\beta = 2, g = 0.8, s_P = 0.15$; obligate: $b_P = 0.3, d_P =$

2.2; facultative: $b_P = 2, d_P = 1$; animals: $\varepsilon = 1, d_A = 1.5$; obligate: $b_A = 1.2, s_A = 0.11$;

facultative animals: $b_A = 2, s_A = 0.2$; except for: (**B**) $a = 0.4$, (**C**) $h = 0.5, \beta = 1, d_P =$

$1.4, b_A = 0.7, s_A = 0.07$, (**D**) $a = 0.05, b_P = 0.1, d_P = 0.05, g = 0.55, \beta = 0.55, s_P =$

$0.25, b_A = 0.19, d_A = 0.2, s_A = 0.05$, (**F**) $a = 0.014, b_P = 0.19, d_P = 0.11, g = 0.6, \beta =$

$0.55, s_P = 0.1, b_A = 0.18, \varepsilon = 2.5, d_A = 0.15, s_A = 0.04$, (**G**) $a = 0.4$.

668   *Figure 2.*

Phase plane diagrams for plant seed-disperser mutualisms, where animals reduce

negative density-dependence in plants. Nullclines are curves of zero growth for plant

(green) and animal seed-disperser (black) populations. Here, plants must be able to persist

at low density ($K_P = (b_P g - d_P)/b_P g s_P > 0$) to benefit from reductions in negative

density-dependence; therefore, we consider only facultative plant populations. All other

formatting and terminology follow Fig. 1. When animals are obligate mutualists (**A**), both

partners may experience threshold effects. Otherwise, both partners experience growth

from low density (**B, C, D, E**), resulting in either a single (**C, E**) or two (bi)stable (**B, D**)

coexistence equilibria. Parameter values are fixed to the following: $a = 1, h = 0.5$; plants:

$b_P = 1, g = 1, s_P = 0.2, \sigma = 0.15, d_P = 0.4$; animals: $\varepsilon = 1$; obligate: $b_A = 1, d_A =$

$2, s_A = 0.15$; facultative: $b_A = 2$; except for: (**A**) $g = 0.6, \sigma = 0.2$, (**B**) $g = 0.5, a =$

$0.36, b_A = 1.75, s_A = 0.12$; (**D**) $g = 0.45, \sigma = 0.175, s_A = 0.12, d_A = 1.75$; (**E**) $\varepsilon = 0.5$,

$s_A = 0.2, d_A = 1.5$.

682   *Figure 3.*

Phase plane diagrams for plant seed-disperser mutualisms, where animals increase

seed germination in plants. Nullclines are curves of zero growth for plant (green) and

animal seed-disperser (black) populations. Dynamics depend upon the parameterization of

the model and whether plants and animals are obligate (O) or facultative (F) mutualists

(rows). When either partner is an obligate mutualist, both partners may experience

threshold effects (**A, B, D**). When either partner is a facultative mutualist, the system may

instead exhibit a single coexistence equilibrium with growth from low density for both

partners (**C, E, F**). Parameter values are fixed to the following: $a = 1, h = 0.5$; plants:

$b_P = 1, g = 0.5, \gamma = 0.5, s_P = 0.1$; obligate: $d_P = 1$; animals: $b_A = 1, \varepsilon = 1$; obligate:

$s_A = 0.1, d_A = 1.4$; facultative: $s_A = 0.2, d_A = 0.2$; except for: (**A**) $s_A = 0.15$; (**B**) $s_P =$

$0.2, d_P = 1.6, d_A = 0.9$; (**C**) $a = 0.5, g = 1$; (**D**) $a = 0.5, s_P = 0.17, d_P = 0.4$; (**E**) $h =$

$1, d_P = 0.2$; (**F**) $h = 1, d_P = 0.3, \varepsilon = 0.8$.

 Figures and Tables

 Figure 1.

698 *Figure 2.*

Figure 3.

Table 1. Table of parameters.

Parameter	Interpretation	Units
a	Per-plant attack rate on rewards	$[P^{-1} t^{-1}]$
b_i	Per-capita birth or seed production rate	$[t^{-1}]$
β	Per-animal seed set	$[A^{-1}]$
d_i	Per-capita death rate	$[t^{-1}]$
ε	Efficiency of converting food rewards to animal offspring	*Unitless*
g	Per-plant seed germination	*Unitless*
γ	Seed germination per animal visitation rate	$[A^{-1} P\, t]$
h	Handling time on rewards	$[t]$
s_i	Negative density-dependence due to, e.g., intraspecific competition, Janzen-Connell effect	s_P: $[P^{-1}]$ or s_A: $[A^{-1} t^{-1}]$
σ	Reduced negative density-dependence per animal visitation rate	$[A^{-1} t]$

All parameters are assumed to be positive (> 0). The units A, P, and t refer to animal density, plant density, and time, respectively. Subscript i is used to specify the parameter for the plant (P) or animal (A) population. We use Greek letters (β, ε, γ, σ) to indicate conversion coefficients for benefits due to mutualism, while Roman letters (a, b, d, g, h, s) indicate plant and animal demographic and feeding parameters.

Appendix A: Mathematical Analysis

In all three of our models (Eqns. 3, 6, 7), plant and animal populations have "trivial"

nullclines: $\bar{P} = 0$, a vertical line along the y-axis, and $\bar{A} = 0$, a horizontal line along the x-

axis. The intersection of both species' trivial nullclines gives a trivial extinction equilibrium

($\bar{P} = 0,\ \bar{A} = 0$) in every model. The animal population, A, follows the same dynamic

equation (Eqn. 1) in all of our models and thus has the same nontrivial nullclines (hereafter,

simply "nullclines").

The animal nullcline (Figs. 1-3, black) is a concave down, increasing function that

saturates with respect to its partner's abundance:

720
$$\bar{A} = \frac{r_A}{s_A} + \frac{\varepsilon}{s_A} \frac{aP}{(1+ahP)}.$$
Eqn. A1

Eqn. A1 describes the balance between negative density-dependence, benefit from

mutualism, and intrinsic growth or decay for facultative or obligate animal populations,

respectively. When $A > \bar{A}$, negative density-dependence is stronger than benefit from

mutualism and/or intrinsic growth, causing A to decrease. When $A < \bar{A}$, negative density-

dependence is weaker and A increases.

The benefit of mutualism to A is $\frac{\varepsilon}{s_A} \frac{aP}{(1+ahP)}$, which saturates with respect to plant

density. As plant density becomes extremely high, A saturates to a higher value than it

could achieve in the absence of its partner:

729
$$\lim_{P \to \infty} \bar{A} = \frac{r_A}{s_A} + \frac{\varepsilon}{s_A h}.$$
Eqn. A2

730   This horizontal asymptote is an upper bound on the density of A. When A is facultative

731   ($r_A > 0$), its nullcline intersects the y-axis in the ecological region (positive quadrant) at

$$\overline{A}_{0,y} = \frac{r_A}{s_A} = K_A. \qquad\qquad \text{Eqn. A3}$$

733   This is animal density in the absence of plants; we call it the carrying capacity (K_A). All

734   else being equal, decreasing r_A pushes the visible part of the animal nullcline down so that

735   when the animal population is obligate ($r_A \leq 0$), its nullcline intersects the y-axis at zero or

736   negative values (not shown). The animal nullcline intersects the x-axis in the positive

737   quadrant at

$$\overline{A}_{x,0} = \frac{-r_A}{a(\varepsilon + h r_A)}. \qquad\qquad \text{Eqn. A4}$$

739   When the animal population is obligate, this intersection occurs in the positive quadrant,

740   whereas when the animal population is facultative, the intersection occurs at negative

741   values of the plant population.

742   The equation for the nullcline of animal-pollinated plants (Fig. 1, green) is:

$$A = \frac{(b_P g s_P \overline{P} - r_P)(1 + a h \overline{P})}{\beta a g (1 - s_P \overline{P})\overline{P}}. \qquad\qquad \text{Eqn. A5}$$

When plants are obligate ($r_P = b_P g - d_P \leq 0$, Fig. 1a-b), the nullcline is a U-shape bounded

by vertical asymptotes at $P = 0$ and $P = 1/s_P$. Note that the asymptote $P = 1/s_P$

represents the plant's maximum population density. When plants are facultative ($r_P > 0$,

Fig. 1c-g), the nullcline is concave up at high density and keeps increasing asymptotically

towards its maximum, $P = 1/s_P$. At low density, the nullcline is concave down, though

this may not be visible in the ecological region of the plane (positive quadrant). Its x-

intercept is

$$\overline{\boldsymbol{P}}_{x,0} = \frac{r_P}{b_P g s_P} = K_P > 0. \qquad \text{Eqn. A6}$$

752   This is plant density in the absence of animals; we call it the carrying capacity (K_P). Inside

753   the U or to the left of the nullcline, plant density increases; to the right or under, plant

754   density decreases due to strong negative density-dependence.

755   The equation for the nullcline of animal-dispersed plants that benefit through

756   reduced negative density-dependence (Fig. 2, green) is:

$$\boldsymbol{A} = \frac{(b_P g s_P \overline{\boldsymbol{P}} - r_P)(1 + a h \overline{\boldsymbol{P}})}{b_P g \sigma a \overline{\boldsymbol{P}}}. \qquad \text{Eqn. A7}$$

758   At high plant density, the nullcline has a linear slope of $\frac{h s_P}{\sigma} \geq h$. At low plant density, the

759   nullcline is concave down, bounded on the left side by a vertical asymptote at $\boldsymbol{P} = 0$.

760   Because we consider only facultative plants for this mutualism ($r_P > 0$), the nullcline

761   always has an x-intercept visible in the ecological region at

$$\overline{P}_{x,0} = \frac{r_P}{b_P g s_P} = K_P > 0. \qquad \text{Eqn. A8}$$

763   The equation for the nullcline of animal-dispersed plants that benefit through

764   germination (Fig. 3, green) is:

$$\boldsymbol{A} = \frac{(b_P g s_P \overline{\boldsymbol{P}} - r_P)(1 + a h \overline{\boldsymbol{P}})}{b_P \gamma a (1 - s_P \overline{\boldsymbol{P}})}. \qquad \text{Eqn. A9}$$

766   In the ecological region, the nullcline is an increasing, concave up curve. At high plant

767   density, the nullcline has a vertical asymptote at $P = 1/s_P$, the maximum plant density in

768   the presence of animal mutualists. At low plant density, the nullcline is concave down,

769   bounded on the left side by a vertical asymptote at $P = 0$. The nullcline intersects the x-

770   axis at

771
$$\overline{P}_{x,0} = \frac{r_P}{b_P g s_P} = K_P \qquad\qquad \text{Eqn. A10}$$

772   which is visible in the ecological region when plants are facultative ($r_P > 0$). It intersects

773   the y-axis at

774
$$\overline{P}_{0,y} = \frac{-r_P}{b_P \gamma a} \qquad\qquad \text{Eqn. A11}$$

which is visible in the ecological region when plants are obligate ($r_P \leq 0$).

<div align="center">*Feasibility*</div>

All our models are structurally unstable (Rohr *et al*. 2014), such that smooth

transitions in parameter values shift the nontrivial nullclines (hereafter, "nullclines") so that

they may intersect in various ways or even fail to intersect, with different dynamical

outcomes for each case (Fig. A1). In the main text, we focused on "feasible" systems, that

is, where coexistence of the partners with non-negative density (i.e. in the ecological

region) was possible. There are many parameterizations that lead to feasibility, but it is not

easy to provide an ecological interpretation for the conditions to achieve them. Here, we

enumerate the geometric conditions that lead to feasibility by describing the required

positioning of the nullclines.

There are four cases for how the plant and animal nullclines intersect:

(1) If the nullclines never intersect (e.g., Fig. A1A, A1E), then coexistence is not possible.

(2) If the nullclines intersect exactly once (e.g., Fig. A1B, A1F), this is an ecologically

unlikely case that we neglect.

(3) If the nullclines intersect twice with at least one intersection in the ecological region

(e.g., Fig. A1C-D, A1G-J), then coexistence is feasible. The low-density equilibrium is a

saddle point while the high-density equilibrium is stable (e.g., Fig. A1C, A1G, A1H).

When only one intersection occurs in the ecological region, it is the stable coexistence

equilibrium (e.g., Fig. A1D, A1J).

(4) If the nullclines intersect three times in the ecological region (e.g., Fig. B1I), then

coexistence is feasible. The low- and high-density equilibria are stable ("bistable

coexistence") and the intermediate equilibrium is a saddle point. Bistable coexistence

appears to only occur when at least one nullcline changes concavity in the ecological

region. Our main-text models display bistable coexistence because their nullclines change

from concave up to concave down (pollination, Figs. 1D, 1F) or from concave down to

linear growth (seed dispersal, Figs. 2B, 2D). However, bistability can more generally result

from Type III functional responses in facultative mutualists.

Functional Forms

To understand the robustness of our results, we analyzed our models using Holling

Type I and III functional forms in addition to the saturating Holling Type II forms

presented in the main text (Table B1, Figs. B1-B3).

References

Rohr, R. P., S. Saavedra, and J. Bascompte. 2014. On the structural stability of mutualistic

systems. Science 345(6195):1–9.

Table A1. Functional forms

Reproductive Services	Type I	Type II	Type III
Pollination $S = C_R$	AP	$\dfrac{aAP}{1 + ahP}$	$\dfrac{aAP^2}{1 + ahP^2}$
Seed Dispersal $S = C_R/P$	A	$\dfrac{aA}{1 + ahP}$	$\dfrac{aAP}{1 + ahP^2}$

Expressions for Holling Type I-III functional forms used to model reproductive services (S)

and consumption rate (C_R) in our pollination (Eqns. 1 & 2) and seed dispersal models:

reduced negative density-dependence (Eqns. 1 & 4), increased germination (Eqns. 1 & 5).

Because reproductive services (S) provided animals are a function of their consumption rate

on plant rewards (C_R), only one functional form needs to be specified. Pollination services

are assumed to be equal to total consumption rate, while seed dispersal services (both

germination and negative density-dependence) are assumed to be equal to per-capita

consumption rate.

Figure A1.

Examples of structural instability. Phase plane diagrams illustrating smooth transitions from infeasible to feasible coexistence (from
left to right panels within a row). Labels for each row identify the relevant models and Plants / Animals as obligate (O) or facultative
(F). (**A-D**) The *A* nullcline shifts smoothly to the left, due to, for example, increasing conversion efficiency (ε). This moves the system

from (**A**) infeasible, to (**B**) ecologically unlikely, through a "blue sky" bifurcation in which (**C**) coexistence is feasible and stable but
will not occur if species' initial densities fall below a threshold, to (**D**) guaranteed stable coexistence. This type of structural instability
occurs in both our pollination model when plants are facultative and in our seed dispersal: germination benefits model. (**E-G**) The
vertex of the *P* nullcline shifts smoothly down, due to, for example, increasing partner quality (β). This moves the system from (**E**)
infeasible, to (**F**) ecologically unlikely, through a blue-sky bifurcation in which (**G**) coexistence is feasible and stable but threshold and
Allee effects are present. (**H-J**) The *P* nullcline shifts smoothly to the right, due to, for example, decreasing negative density-
dependence (s_P). This moves the system from infeasible (not shown), to ecologically unlikely (not shown), through a blue-sky
bifurcation in which (**H**) coexistence is feasible and stable but threshold effects are present, to (**I**) bistability in which stable
coexistence is guaranteed but occurs either at a low- or high-density equilibrium, to (**J**) guaranteed stable coexistence. This type of
structural instability occurs in both our seed dispersal model (shown) and in our pollination model when plants are facultative (not
shown).

Figure A2.

Effect of functional form on pollination mutualisms. Panels show phase plane diagrams with

unique dynamical outcomes when each partner population (***P***, green; ***A***, black) is an obligate or

facultative mutualist following a Holling Type I, II, or III functional response (Table A1).

Equilibria are stable (filled, black) or unstable or saddle points (hollow). Saddle points are

bisected by a separatrix (dashed black line) that divides the plane into two regions.

Figure A3.

Effect of functional form on seed dispersal mutualisms that reduce negative density-

dependence in plants. Formatting follows figure A2.

Figure A3.

Effect of functional form on seed dispersal mutualisms that increase germination in plants.

Formatting follows figure A2.

Appendix B: Saturating Services

In our main text, benefits to plants saturate indirectly, due to the limited time available

for animals to forage upon rewards. In natural systems, benefits also saturate directly in terms of

the number of flowers available to be fertilized, seeds available to germinate, etc. We can ensure

this by writing benefit as a saturating function of reproductive services $S/(1 + S)$. However,

simpler models are desirable both for analytical tractability and because they can perform

sufficiently or even better than more complicated models when confronted with messy or

ecologically constrained data reflective of natural systems (Novak & Stouffer 2020). Indeed, our

pollination and germination models retain identical qualitative dynamics when modeling plant

benefit with saturating services, though our negative density-dependence model exhibits simpler

dynamics that are qualitatively like the germination model. For this reason, we presented the

simpler models in the main text, but we derive the more detailed models below. The more

detailed approach allows more specific parameter defintions that may be easier to measure

empirically.

For plant populations (P), we define b_P as per-capita seed set, d_P as the per-capita death

rate, and s_P as negative density-dependence in the absence of mutualism. We assume that

negative density-dependence limits the seeds that survive and mature to reproductive adults, due

to, e.g., the Janzen-Connell effect and seed competition for recruitment. Additionally, we assume

that plant reproduction is limited by the fraction of flowers that can be fertilized through wind

pollination or selfing ($0 \leq f \leq 1$) and the fraction of seeds that can germinate ($0 \leq g \leq 1$) even

when subjected to low negative density-dependence. Reproductive services (S) from animals

increase the density of mature (reproductive) plant individuals by pollinating flowers (φ),

improving germinability (γ), or providing refuge from seed predation and other sources of

density-dependent mortality during recruitment (σ).

When animals pollinate flowers by transferring pollen between them (reproductive

services, S), the fraction of flowers that are pollinated (f) can increase in the presence of animals

to a maximum of $f + \varphi \leq 1$. We can then model animal-pollinated plant population dynamics

($\mathbf{P}$) as

$$\frac{d\mathbf{P}}{dt} = \mathbf{P}\left[b_P(f + \varphi\frac{S}{1+S})g(1 - s_P\mathbf{P}) - d_P\right] \qquad \text{Eqn. B1}$$

where benefit from pollination services (S) saturates as the fraction of flowers that are pollinated

approaches its maximum. As in the main text, we assume $S = C_R$, total foraging rate of animals

on plant rewards. Using a Holling Type II functional response for C_R and simplifying the algebra

yields:

$$\begin{cases} \frac{d\mathbf{P}}{dt} = \mathbf{P}\left[b_P(f + \varphi\frac{aA\mathbf{P}}{1+ah\mathbf{P}+aA\mathbf{P}})g(1 - s_P\mathbf{P}) - d_P\right] \\ \frac{dA}{dt} = A\left[b_A + \varepsilon\frac{a\mathbf{P}}{1+ah\mathbf{P}} - s_AA - d_A\right] \end{cases} \qquad \text{Eqn. B2}$$

In this specific form of the model, plant benefits are limited both by the fraction of flowers

available to pollinate (φ) and by pollinators' handling time on flowers (h).

Similarly, when animals rescue seeds from predators, pathogens, and intraspecific

competition by transporting them away from parent trees dispersal services, negative density-

dependence (s) can be reduced to a minimum of $s - \sigma \geq 0$ in the presence of animals. Modeling

this yields:

$$\frac{d\mathbf{P}}{dt} = \mathbf{P}\left[b_Pfg\left(1 - \left(s - \sigma\frac{S}{1+S}\right)\mathbf{P}\right) - d_P\right] \qquad \text{Eqn. B3}$$

When animals increase germinability of seeds by masticating and/or digesting fruits

(reproductive services, S), the fraction of seeds that germinate (g) can increase in the presence of

animals to a maximum of $g + \gamma \leq 1$. Modeling this yields:

$$\frac{dP}{dt} = P\left[b_P f(g + \gamma \frac{S}{1+S})(1 - s_P P) - d_P\right]$$ Eqn. B4

As in the main text, we assume $S = C_R/P$, per-plant foraging rate of animals on plant rewards,

for both seed dispersal models. Using a Holling Type II functional response for C_R and

simplifying the algebra yields our specific model for escape from negative density-dependence in

animal seed-dispersal mutualisms:

$$\begin{cases} \frac{dP}{dt} = P\left[b_P f g \left(1 - \left(s_P - \sigma \frac{aA}{1+ahP+aA}\right)P\right) - d_P\right] \\ \frac{dA}{dt} = A\left[b_A + \varepsilon \frac{aP}{1+ahP} - s_A A - d_A\right] \end{cases}$$ Eqn. B5

and for increased germinability in animal seed-dispersal mutualisms:

$$\begin{cases} \frac{dP}{dt} = P\left[b_P f(g + \gamma \frac{aA}{1+ahP+aA})(1 - s_P P) - d_P\right] \\ \frac{dA}{dt} = A\left[b_A + \varepsilon \frac{aP}{1+ahP} - s_A A - d_A\right] \end{cases}$$ Eqn. B6

In these specific forms, plant benefits are limited both by the fraction of seeds available to

germinate (γ) or the negative density-dependent effects to reduce (σ) and by dispersers' handling

time on rewards (h).