Title

Quantitative assessment of multiple fish species around artificial reefs using environmental DNA

metabarcoding

Authors

Masaaki Sato[1*], Nariaki Inoue[1], Ryogen Nambu[1,2], Naoki Furuich[1], Tomohito Imaizumi[1], and

Masayuki Ushio[3,4]

[1]Fisheries Engineering Division, Fisheries Technology Institute, Japan Fisheries Research and

Education Agency (FRA), Kamisu, Ibaraki, Japan

* Correspondence (E-mail: msato22@affrc.go.jp)

[2]Present address: Fisheries Division, Japan International Research Center for Agricultural

Sciences (JIRCAS), Tsukuba, Ibaraki, Japan

[3]Hakubi Center, Kyoto University, Kyoto 606-8501 Japan

[4]Center for Ecological Research, Kyoto University, Otsu, Shiga 520-2113 Japan

Abstract

Since the early 1970s, many artificial reefs (ARs) have been deployed in Japanese coastal waters

to create fisheries grounds. Recently, researchers began to use environmental DNA (eDNA)

methods for biodiversity monitoring of aquatic species. A metabarcoding approach using internal

standard DNAs (i.e., quantitative MiSeq sequencing) makes it possible to monitor eDNA

concentrations of multiple species simultaneously. This method can improve the efficiency of

monitoring AR effects on fishes. Our study investigated distributions of marine fishes at ARs and

surrounding stations in the open oceanographic environment of Tateyama Bay, central Japan,

using quantitative MiSeq sequencing and echo sounder survey. Using the quantitative

metabarcoding method, we found higher quantities of fish eDNAs at the ARs than at surrounding

stations and different fish species compositions between them. Comparisons with eco sounder

survey also showed positive correlations between fish eDNA concentration and echo intensity,

which indicates a highly localized signal of eDNA at each sampling station. These results suggest

that quantitative MiSeq sequencing is a promising technique to complement conventional

methods to monitor distributions of multiple fish species.

Introduction

Artificial reefs (ARs) have been deployed worldwide to create fisheries grounds. Since the early 1970s, restrictions on overseas fishing have influenced high seas fisheries outside the exclusive economic zones (EEZ), and this situation led the Japanese government to implement a new fishery policy to develop and maintain coastal fisheries. In 1971, the Japanese government started the first national project of maintenance and development of coastal fisheries and has continued to implement similar projects, spending $ 5.4 billion to deploy ARs throughout coastal waters. Yamane[1] indicated that the total number of Japanese AR projects and proportion of shallow fishing grounds with ARs in 1986 reached 6,443 and 9.3%, respectively. After establishing a large number of ARs in Japanese waters, underwater visual census, surveys using fishery gear, and echo sounder surveys have been used to examine fish aggregation and stock enhancement effects of ARs in Japanese waters[2–5]. These survey methods have advantages and disadvantages. An underwater visual census is effective for species identification, fish counts, and size estimation, but it requires identification skills in situ and/or a long investigation time for large areas. Surveys with bottom trawls and gillnets can obtain similar information on the fish to the underwater visual census, but these surveys exploit fish individuals. Echo sounders can survey large geographical areas quickly without fish collection, but with this method, it is difficult to identify fish species.

Recently, researchers have begun to use environmental DNA (eDNA), DNAs derived from environmental samples such as water, to investigate the distribution of aquatic organisms, including marine fishes[6,7]. In the case of macro-organisms, eDNA originates from metabolic waste, damaged tissue, dead individuals, or spawning events[8], and the eDNA contains information about the species of the organisms that produced it[7]. This method does not need to exploit a body or body tissues of target species and can detect target organisms noninvasively. Also, this method requires less field time and is more likely to detect the presence of target species than the other survey methods[6,9]. Recently, the metabarcoding approach, using high-throughput sequencing technologies (e.g., Illumina MiSeq) and universal primer sets, has been utilized to reveal the fish diversity in a given area[6,10]. MiFish, a universal polymerase chain reaction (PCR) primer set, is especially popular for fish eDNA metabarcoding and has detected more fish species than other competing primers[11].

Although the eDNA metabarcoding approach has become a cost- and labor-effective approach for estimating aquatic biodiversity, it has one limitation, i.e., the quantity of eDNA cannot be estimated from the number of eDNA sequence reads obtained by high-throughput sequencing partly due to several experimental problems such as PCR inhibitions[7]. However, some studies have proposed an approach using an internal standard DNA to overcome this limitation. Ushio et al.[7] tested whether additions of internal standard DNAs (i.e., known amounts of short DNA

fragments from fish species that have never been observed in a sampling area) to eDNA samples
can create sample-specific regression lines between the numbers of sequence reads and DNA
copies. Based on regression lines for each sample, they converted the sequence reads of DNAs
from unknown fish species (i.e., non-standard fish species) to the number of fish DNA copies in
each sample. This study showed that the estimated numbers of total and two dominant fish
(Japanese anchovy and jack mackerel) DNA copies were significantly positively correlated with
the numbers of DNA copies quantified by qPCR. Therefore, this method (i.e., quantitative MiSeq
sequencing) can contribute to the quantitative assessment of multispecies fish eDNA. Some
studies have shown that the eDNA quantity could be a proxy for the abundance or biomass of
macro-organisms, not only in closed systems like tank and mesocosm[12], but also in marine
ecosystems[9,13]. The quantitative MiSeq sequencing with MiFish primer could be one of the most
cost- and labor-effective approaches for the quantitative monitoring of multi species fish in
marine and coastal ecosystems, including ARs.

Here, we investigated marine fish distributions around ARs in Tateyama Bay, central Japan,
using eDNA metabarcoding, catch record of fishery, and eco sounder method. The quantitative
MiSeq sequencing using internal standard DNAs and MiFish primers was applied to assess
whether the distribution of multi fish species around ARs could be monitored based on the spatial
variation in the estimated quantities of eDNA (i.e., identification of fish species and
quantification of the number of fish eDNA copies simultaneously). Specifically, the following
were tested: 1) whether fish communities detected by eDNA metabarcoding match with the catch
record of the coastal fishery, 2) whether the DNA quantities of multiple fish species calculated by
the quantitative MiSeq sequencing decrease with increasing distance from an AR, 3) whether the
spatial variation of fish DNA quantities correlates to the relative distribution estimated by the
acoustic survey.

Materials and methods

Study site, field survey, and in situ filtration

The field survey was performed in Tateyama Bay (34°60'N, 139°48'E), central Japan, in the
proximity of the Kuroshio warm current facing the Pacific Ocean (Fig. 1). This area has many
artificial reefs (ARs) created to improve fishing efficiency for fishers. Among the ARs, we
focused on one high-rise steel AR (AR1), with a height of 30 m, where fish tended to aggregate
(Fig. S1). Sampling stations were set up at the AR1 and at six linear distant points extending
northeast and southwest. These stations were named E150, E500, E750, W150, W500, and W750,
where "W" or "E" and the number of each station name represented northeast or southwest and
distance in meters from the AR1, respectively (Table 1 and Fig. 1). Another station was set up at
a second AR (AR2: 25 m height) 220 m from AR1 because we found AR2 by chance during the

survey (Table 1 and Fig. 1), and it might affect the eDNA concentration at other stations.

We conducted water sampling for eDNA analysis and performed an acoustic survey for

estimating relative fish density using research vessel *Taka-maru* (Japan Fisheries Research and

Education Agency: FRA) on May 23, 2018. We started the echo sounder survey at the southern

part of the bay and continued it during the water sampling. Although the echo sounder survey

could not differentiate between fish species, we collected this data to assess the association

between the estimated concentration fish of eDNA and the echo intensity measured by the echo

sounder. Water sampling began at E750, then continued along the transect line to E150, AR1,

W150, W500, W750, before going back to AR2. At each sampling station, we collected 10 L of

seawater from both the middle and bottom layers by one cast of two Niskin water samplers and

measured vertical profiles of water temperature and salinity with a

conductivity-temperature-depth sensor (RINKO profiler, JFE Advantech Co., Ltd.). Then, two

2L samples were subsampled from the seawater samples and immediately filtered using a

combination of Sterivex filter cartridges (nominal pore size = 0.45 μm; Merck Millipore) through

an aspirator (i.e., the two filters were subsets of a single water collection) in a laboratory on the

research vessel. After filtration (approximately 15 min), an outlet port of the filter cartridge was

sealed with an outlet luer cap, 1.5 ml RNAlater (ThermoFisher Scientific, Waltham, MA) was

injected into the cartridge using a filtered pipette tip to prevent eDNA degradation, and an inlet

port was also sealed with an inlet luer cap[14]. The Niskin water samplers were bleached before

each water collection using a commercial bleach solution while filtering devices (i.e., filter

funnels and measuring cups used for filtration) were bleached after every filtration. We filtered

2L MilliQ water with a filter funnel and measuring cup for every two or three sites as a field

negative control to test for possible contamination. The filter cartridges were placed in a freezer

immediately after filtration until eDNA extraction. Disposable latex or nitrile gloves were worn

during sampling and replaced between each sampling station.

DNA extraction and purification

Workspaces were sterilized prior to DNA extraction using 10% commercial bleach, and filter tip

pipettes were used to safeguard against cross-contamination. Following the method developed by

Miya et al.[14], the eDNA was extracted and purified. Briefly, after removing RNAlater inside the

cartridge using a centrifuge, proteinase-K solution was injected into the cartridge from the inlet

port, and the port was re-capped with the inlet lure cap. The eDNA captured on the filter

membrane was extracted by constant stirring of the cartridge at a speed of 20 rpm using a roller

shaker placed in an incubator heated at 56°C for 20 min. The eDNA extracts were transferred to a

2-ml tube from the inlet of the filter cartridges by centrifugation. The collected DNA was purified

using a DNeasy Blood & Tissue Kit (Qiagen) following the manufacturer's protocol. After the

purification, DNA was eluted using 100 µl of the elution buffer (buffer AE). All DNA extracts

were frozen at −20°C until paired-end library preparation.

Preparation of internal standard DNAs

Five artificially designed and synthetic internal standard DNAs, which were similar but not

identical to the region of any existing fish mitochondrial 12S rRNA, were included in the library

preparation process to estimate the number of fish DNA copies (i.e., quantitative MiSeq

sequencing)[7,15]. They were designed to have the MiFish primer□binding regions as those of

known existing fishes and to have the conserved regions in the insert region. Variable regions in

the insert region were replaced with random bases so that no known existing fish sequences had

the same sequences as the standard sequences. The standard DNA size distribution of the library

was estimated using an Agilent 2100 BioAnalyzer (Agilent, Santa Clara, CA, USA), and the

concentration of double-stranded DNA of the library was quantified using a Qubit dsDNA HS

assay kit and a Qubit fluorometer (ThermoFisher Scientific, Waltham, MA, USA). Based on the

quantification values obtained using the Qubit fluorometer, the copy number of the standard

DNAs was adjusted as follows: Std. A (100 copies/µl), Std. B (50 copies/µl), Std. C (25

copies/µl), Std. D (12.5 copies/µl) and Std. E (2.5 copies/µl). Then, these standard DNAs were

mixed.

Paired-end library preparation

Four PCR‐level negative controls (i.e., each two with and without internal standard DNAs) were

employed for MiSeq run to monitor contamination during the experiments. The first-round PCR

(1st PCR) was carried out with a 12-µl reaction volume containing 6.0 µl of 2 × KAPA HiFi

HotStart ReadyMix (KAPA Biosystems, Wilmington, WA, USA), 0.7 µl of each primer (5 µM),

2.6 µl of sterilized distilled H$_2$O, 1.0 µl of standard DNA mix and 1.0 µl of template. Note that the

standard DNA mix was included for each sample. The final concentration of each primer was 0.3

µM. We used a mixture of the following four PCR primers modified from original MiFish

primers[16]: MiFish-U-forward (5´-ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT

NNN NNG TCG GTA AAA CTC GTG CCA GC) and MiFish-U-reverse (5´-GTG ACT GGA

GTT CAG ACG TGT GCT CTT CCG ATC TNN NNN CAT AGT GGG GTA TCT AAT CCC

AGT TTG-3´), MiFish-E-forward (5´-ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA

TCT NNN NNG TTG GTA AAT CTC GTG CCA GC-3´), and MiFish-E-reverse (5´-GTG ACT

GGA GTT CAG ACG TGT GCT CTT CCG ATC TNN NNN CAT AGT GGG GTA TCT AAT

CCT AGT TTG-3´). These primer pairs co-amplify a hypervariable region of the fish

mitochondrial 12S rRNA gene (around 172 bp; hereafter called MiFish sequence) and append

primer-binding sites (5´ ends of the sequences before six Ns) for sequencing at both ends of the

amplicon. The five random bases were used to enhance cluster separation on the flow cells during

initial base call calibrations on the MiSeq platform. The thermal cycle profile after an initial 3 min

denaturation at 95°C was as follows (35 cycles): denaturation at 98°C for 20 s; annealing at 65°C

for 15 s; and extension at 72°C for 15 s, with a final extension at the same temperature for 5 min.

Eight replications were performed for the 1st PCR, and the replicates were pooled to minimize the

PCR dropouts. The 1st PCR products from the eight tubes were pooled in a single 1.5-ml tube.

Then, we sent the 1st PCR products to IDEA consultants, Inc. to outsource the following MiSeq

sequencing processes. The pooled products were purified and size-selected for 200−400bp using a

SPRIselect (Beckman Coulter inc.) to remove dimers and monomers following the

manufacturer's protocol.

The second-round PCR (2nd PCR) was carried out with a 24 µl reaction volume containing 12

µl of 2 × KAPA HiFi HotStart ReadyMix, 2.8 µl of each primer (5 µM), 4.4 µl of sterilized

distilled H2O, and 2.0 µl of template. We used the following two primers to append the

dual-index sequences (8 nucleotides indicated by Xs) and flowcell-binding sites for the MiSeq

platform (5´ ends of the sequences before eight Xs): 2nd-PCR-forward (5´-AAT GAT ACG GCG

ACC ACC GAG ATC TAC ACX XXX XXX XAC ACT CTT TCC CTA CAC GAC GCT CTT

CCG ATC T-3´); and 2nd- PCR-reverse (5´-CAA GCA GAA GAC GGC ATA CGA GAT XXX

XXX XXG TGA CTG GAG TTC AGA CGT GTG CTC TTC CGA TCT-3´). The thermal cycle

profile after an initial 3 min denaturation at 95°C was as follows (12 cycles): denaturation at 98°C

for 20 s; combined annealing and extension at 72°C for 15 s, with a final extension at 72°C for 5

min. The concentration of each second PCR product was measured by quantitative PCR using TB

Green Fast qPCR Mix (Takara inc.). Each sample was diluted to a fixed concentration and

combined (i.e., one pooled 2nd PCR product that included all samples). The pooled 2nd PCR

product was size-selected to approximately 370 bp using BluePippin (SAGE SCIENCE). The

size-selected library was purified using the Agencourt AMPure XP beads, adjusted to 4 nM by

quantitative PCR using TB Green Fast qPCR Mix (Takara inc.), and sequenced on the MiSeq

platform using a MiSeq v2 Reagent Kit (2 × 150 bp).

Data preprocessing and taxonomic assignment

The raw MiSeq data were converted into FASTQ files using the bcl2fastq program provided by

Illumina (bcl2fastq v2.18). The FASTQ files were then demultiplexed using the command

implemented in Claident[17]. We adopted this process rather than using FASTQ files demultiplexed

by the Illumina MiSeq default program in order to remove sequences whose 8☐mer index

positions included nucleotides with low-quality scores (i.e., Q-score < 30).

The processed reads were subjected to a BLASTN search against the full NCBI database. If the

sequence similarity between queries and the top BLASTN hit was < 98.5% and the sequence

length was less than ≤ 150 bp, the unique sequence was not subjected to the following analyses.

After BLASTN searches, assembled sequences assigned to the same species were clustered, and

we considered the clustered sequences as operational taxonomic units (OTUs). To remove

possible contaminants, we removed the OTUs whose sequence reads were < 0.05 % of the total

reads as a noise for each sample.

Catch record of coastal fishery

To confirm the feasibility of MiFish metabarcoding, we compared the fish community of MiFish

metabarcoding with a catch record of a net set near the study site (Fig.1). The detection rate was

defined as the proportion of species detected by metabarcoding compared to the species collected

by the net. Because our eDNA survey was conducted in May 2018, we used catch-record data

from the same period (May 2018).

Estimation of DNA copy numbers

According to Ushio et al.[7] and Ushio[15], the procedure to estimate DNA copy numbers consisted of

two parts: (i) linear regression analysis to examine the relationship between sequence reads and

the copy numbers of the internal standard DNAs for each sample and (ii) the conversion of

sequence reads of non-standard fish DNAs to estimate the copy numbers using the result of the

linear regression for each sample.

Linear regressions were used to examine how many sequence reads were generated from one

fish DNA copy through the library preparation process. Note that a linear regression analysis

between sequence reads and standard DNAs was performed for each sample, and the intercept

was set as zero. The regression equation was: MiSeq sequence reads = regression slope × the

number of standard DNA copies [/µl]. The number of linear regressions performed was

thirty-three (=the number of fish DNA samples + field negative controls with standard DNAs),

and thus thirty-three regression slopes were estimated in total.

The sequence reads of non-standard fish DNAs were converted to copy numbers using

sample-specific regression slopes estimated by the above regression analysis. The number of

non-standard fish DNA copies was estimated by dividing the number of MiSeq sequence reads by

a sample-specific regression slope (i.e., the number of DNA copies = MiSeq sequence

reads/regression slope). A previous study demonstrated that these procedures provide a

reasonable estimation of DNA copy numbers using high-throughput sequencing[7].

Acoustic survey of fish school using echo sounder

We estimated relative fish density using a quantitative echo sounder following the acoustic survey

of Yamamoto et al.[9]. We used the echo sounder, KFC-6000 (Sonic Co. Tokyo, Japan), with a

transducer (frequency, 38 kHz; beam type, split-beam; beam width, approximately 7°; pulse
duration, 0.3 ms; ping rate, 0.4 s). The transducer was set at the bottom of the research vessel at a
depth of 2.1 m to avoid cavitation bubbles generated by the research vessel. The acoustic devices
were operated during the entire survey cruise, and all signals were recorded. The average ship
speed was ~10 knots between sampling stations, although the ship slowed down when
approaching a sampling station and completely stopped when water samples were being
collected.

We eliminated noise from the obtained echo intensity data using Echoview ver. 11.0
(Echo-view Software Pty. Ltd., Tasmania, Australia). Signals between the sea bottom and 1.0 m
above the sea bottom were eliminated to avoid possible confounding with the acoustic dead zone.
In addition, the signas of Niskin water samplers and RINKO profiler were also eliminated.
Finally, we eliminated the signals of bubbles generated by the movement of the screw propeller.
After eliminating these noises, we re-obtained echo intensity data to assess the association with
eDNA concentration. Note that the obtained echo intensity data would include echo signals from
a variety of fish species.

Data analyses
We performed three types of univariate analysis using generalized linear models (GLMs). First,
we used likelihood ratio tests to assess variation in the eDNA copy numbers of total fish, demersal
fish, pelagic fish, and five dominant OTUs, the number of OTUs, as well as echo intensity among
sampling stations and between vertical positions of water sampling. When the effects of sampling
stations were significant, the Tukey-HSD test was used to determine the significant differences
between sampling stations. The second analysis assessed the number of OTUs and eDNA copy
numbers of total, demersal, pelagic, and five dominant OTUs in relation to distance from the
AR1, vertical position of water samples (i.e., middle or bottom layers, represented as 'depth'),
and the interaction between the distance and vertical position. Although our sampling stations
contained two ARs, we focused on AR1 in the second analysis because more fish aggregated
around AR1 and transport of eDNAs from AR1 may have more impact than those from AR2. The
third analysis examined the relationships between eDNA copy numbers (each eDNA copy
number of total fish, demersal fish, pelagic fish, or five dominant OTUs described below) and
echo intensity (S_V). Because echo intensity can correlate to distance from the AR1, the second and
third analyses were separately performed. Based on Fishbase[18], each fish OTU was classified into
demersal and pelagic species. For the analyses, we used the Poisson distribution with a log-link
function for OTU number because it was count data, while a gamma distribution with a log-link
was used for eDNA copy numbers of total fish, pelagic fish, demersal fish, and *Scomber* spp. (one
dominant OTU) because these values are continuous and greater than zero. A mixture of binomial

and gamma distributions, which is often called a "gamma hurdle model," was also used as in
previous studies[19,20] for eDNA copy numbers of remaining dominant OTUs because these values
are continuous and contain zero values. Bayesian information criterion (BIC) was used to select
the correct model among candidate models for the second and third analyses[21].

We also performed multivariate analyses to determine differences in the fish community
structures between sampling stations/vertical positions. Community structures of each sampling
were visualized using non-metric dimensional scaling (NMDS) based on the Bray–Curtis
dissimilarity index of the copy numbers of each OTUs. Similarity tests between two variables
(sampling stations and vertical positions) were conducted using nonparametric multivariate
analysis of variance (NPMANOVA). All statistical analyses were performed using RStudio
version 1.1.423 (R version 3.6.1) with several packages. The packages used were: (a)
glmmTMB[22] for regression models; (b) multcomp[23] for Tukey-HSD test; (c) MuMIn[24] for BIC
model selection, (d) vegan[25] for multivariate analysis; (e) ggplot2[26] and (f) ggpubr[27] for figure
creation.

Results

Sequence reads

The MiSeq paired☐end sequencing (2 × 150 bp) of the 34 libraries for this study (containing 32
PCR replicates, one negative field control, and one PCR-level control), together with 142 libraries
from another study (total number of libraries = 176), yielded a total of 15.98 million reads, with
95.3% base calls containing Phred quality scores of $\geq$ 30.0 (Q30; error rate = 0.1% or base call
accuracy = 99.9%).

For this study, we obtained 4,862,616 MiSeq reads, of which 3,528,543 were high-quality,
merged reads. Number of non-standard fish Miseq reads were 2,153,777 out of the 3,528,543
(61.0%). In the MiSeq run, 18.8%−98.6% of sequence reads were from non-standard fish DNAs
of field samples, and 0% was from those of a field negative control. The final list included 95
OTUs distributed across 49 families and 83 genera (Table S1).

Comparison of eDNA metabarcoding with fish catch record

The fish catch record of a set net indicated that 289 thousand kg of total fish biomass belonging to
41 classifications (39 species and two classifications of closely related species) was caught in
May 2018. The catch biomass was highly variable between the classifications. For example, 255
thousands kg of *Scomber japonicus* was recorded, whereas *Fistularia petimba* was 0.2 kg only
(Table S2). Of the 41 classifications caught by the set net, eDNA metabarcoding detected 25
species. Three classifications of the 41 classifications were not recorded at the species level in the
list of the fish catch of the set net, while the metabarcoding cannot assign OTUs at the species

level for the *Scomber japonicus* and *S. australasicus* because they are so closely related. Thus,

excluding these five species, our metabarcoding eventually detected 58.3% (= 21/36) of the fishes

caught by the set net. Among the five dominant OTUs detected by MiFish metabarcoding

described below (*B. splendens*, *P. trilineatum*, *Scomber* spp., *P. major*, and *T. japonicus*), four

OTUs except *B. splendens* appeared in the list of the fish catch.

Relationship between the copy numbers and sequence reads of the standard DNA

The relationships between the sequence reads and copy numbers of standard DNAs were

examined by linear regression analysis (Fig. 2). Within each sample, the sequence reads linearly

and positively correlated with the copy numbers of standard DNAs (Fig. 2a). R^2 values of the

regression lines ranged from 0.71 to 0.99, and more than 75% of the samples had values over 0.9

(Fig. 2b). These results suggested that the number of sequence reads was generally proportional to

the number of copies of DNA initially added to the first PCR reaction within a single sample. The

slopes of the linear regressions varied depending on the sample, suggesting that the amplification

efficiency or removal amounts of DNAs during the purification of PCR products were dependent

on the individual sample. These sample-specific slopes were used to convert sequence reads to the

copy numbers of fish OTUs. Based on the estimated copy numbers of each OTU, five dominant

OTUs were *Beryx splendens*, *Parapristipoma trilineatum*, *Scomber* spp. (*S. japonicus* or *S.*

australasicus), *Pargus major*, and *Trachurus japonicus* (Fig. 1b–f).

Levels of contaminations were assessed using the field negative control. DNA copy numbers

detected in the field negative control (n = 1) were 0% of mean DNA copy numbers of

field-positive samples, suggesting that there was no amount of contamination during the library

preparation process.

Site and vertical variation in fish eDNA quantities, the number of OTUs, and echo intensity

The estimated copy numbers of fish OTUs were summed to estimate the total, demersal, and

pelagic fish eDNA quantities. The average eDNA copy numbers of total fish, demersal fish, and

pelagic fish were 64.3, 44.6, and 19.7 copies/ml water, respectively, while those of five dominant

OTUs, *B. splendens*, *P. trilineatum*, *Scomber* spp., *P. major*, and *T. japonicus,* were 20.3, 13.8,

12.9, 5.5, and 3.2 copies/ml water, respectively. Likelihood ratio tests showed that the eDNA

copy numbers significantly varied among sampling stations for total, demersal, pelagic fishes, *P.*

trilineatum, *Scomber* spp., *P. major*, and *T. japonicus*, whereas no significant variation was

detected for *B. splendens* (Table S3). Tukey-HSD tests indicated that significantly higher total

and demersal fish eDNA copy numbers were observed at AR1 than at W150–750 and E 500–750,

while the pelagic fish eDNA copy number was higher at AR1 than at E750 (Fig. S1). Also, the

total, demersal, and pelagic fish eDNA quantities were significantly higher at AR2 than at E750.

For dominant fish OTUs, *P. trilineatum*, *P. major*, and *T. japonicus* eDNA copies were

significantly higher at AR1 than at other stations, while *Scomber* spp. eDNA quantities were

higher at AR1 and AR2 than at E750. Echo intensity (S_V) ranged from 4.53×10^{-8} to 1.38×10^{-6}, and

a likelihood ratio test indicated significant variation in echo intensity among the stations. Like the

fish eDNA quantities, the echo intensity was higher at AR1 than at the surrounding stations.

Meanwhile, the number of OTUs detected at each station ranged from 9 to 27, with a mean of 16

OTUs, and there was no significant variation among the stations (Table S3).

Aside from echo intensity, likelihood ratio tests indicated that all the types of fish eDNA

quantities and number of OTUs were significantly higher in the bottom samples than in the

middle ones (Table S3 and Fig. S2).

Effects of distance from an artificial reef on fish eDNA quantities and the number of OTUs

Distance from the AR1 and vertical position were included in the Gamma component of the

minimum BIC model for total fish, demersal fish, pelagic fish, and *Scomber* spp. eDNA quantities

(Table 2). Estimated parameters indicated that these eDNA quantities decreased with increasing

distance from the AR1 and the eDNA quantities were lower in the middle samples than in the

bottom ones. For *Parapristipoma trilineatum*, *Pargus major*, and *Trachurus japonicus* eDNA

quantities, distance from the AR1, vertical position, and their interaction were included in the

Gamma component of the minimum BIC model. Estimated parameters indicated that these eDNA

quantities decreased with increasing distance from AR1, but the intercepts and slopes of the

relationship with distance differed between the layers. The eDNA quantities of *P. major* were

lower in the middle layer at 0 m from AR1, while those of *P. trilineatum* and *T. japonicus* were

higher in the middle samples at 0 m from AR1 (Fig. 3). Slopes for *P. trilineatum*, and *T. japonicus*

eDNA quantities were steeper in the middle layer, while that for *P. major* eDNA was steeper in

the bottom layer. Only the vertical position was included in the Gamma component of the

minimum BIC model for the number of OTUs and *Beryx splendens* eDNA quantity, and these

values were higher in the bottom layer than in the middle layer (Table 2 and Fig. 3).

Relationships between fish eDNA quantities and echo intensity

Based on the BIC values, echo intensity was selected as an explanatory variable for total fish,

demersal fish, pelagic fish, *B. splendens, P. trilineatum, Scomber* spp., and *T. japonicus* eDNA,

but not for *P. major*. The former eDNA quantities had positive relationships with echo intensity,

and echo intensity had high predictive power for these eDNA quantities (Fig. 4 and Table 3).

Fish community structure

There was dissimilarity in the NMDS in fish community structure between the AR and other

surrounding stations (Fig. 5, NMDS stress = 0.096). PERMANOVA detected significant

differences in community structure between the ARs (AR1 and AR2) and surrounding stations (E

and W) as well as between the middle and bottom sampling layers (ARs vs surrounding stations:

F = 3.040, p = 0.005; Middle vs bottom: F = 4.542, p = 0.002).

Discussion

Field studies have begun to use an eDNA approach for monitoring fish density or diversity in

coastal ecosystems[6,7,9,10], but, as far as we know, there have been no studies using this method to

assess artificial reef (AR) effects on marine fishes. We found that eDNA metabarcoding

efficiently detected the fish assemblages around ARs of Tateyama Bay. MiFish metabarcoding

detected 95 OTUs from 32 eDNA samples collected within 8 h on May 23, 2018 (Table S1),

while 41 classifications (39 species and two classifications of closely related species) were

recorded in the fish catch of the net set near the study site in May 2018 (Table S2). Of the 36

species recorded in the fish catch over a month, MiFish metabarcoding detected 58.3% of species

(i.e., 21 species/36 species). This detection rate of metabarcoding relative to the conventional

approach was within the range of previous MiFish metabarcoding studies (45−70%)[6,28,29]. Four

dominant OTUs in terms of eDNA quantities (*P. trilineatum Scomber* spp., *P. major*, and *T.*

japonicus) were also a higher proportion of fish catch of the set net (Table S1). Meanwhile,

MiFish metabarcoding detected 70 fish OTUs that were not collected by the set net. This

difference in the detected fish community between the MiFish metabarcoding and set net may be

partly because the set net could not collect small-sized fish, and non-targeted fishes (e.g.,

Myctophidae and Gobiidae) were not recorded in the fish catch. These results indicated that

eDNA metabarcoding approach had a high and reasonable resolution for examining a fish

community in the study site.

We observed linear relationships between the number of sequence reads and the quantities of

internal standard DNAs. This result suggested that the sequence reads were proportional to the

copy numbers of DNA within a single sample, indicating that the conversion of sequence reads

using a sample□specific regression slope would provide an estimation of DNA copy numbers in

the DNA extract (i.e., estimated DNA copy number = sequence reads/sample□specific regression

slope). This finding is consistent with previous studies showing that this procedure provides a

reasonable estimate of copy numbers of multispecies eDNAs[7,15]. Meanwhile, the quantitative

Miseq also has limitations; i.e., this method does not take account of the variation in amplification

efficiency among species. Therefore, we have to caution against comparisons of DNA copy

numbers between different species.

This study efficiently estimated the eDNA quantities of multiple fish species around an AR.

Quantitative MiSeq sequencing found that the eDNA quantities of total, demersal, pelagic fishes,

and four dominant OTUs, except *B. splendens*, were higher at AR1 than adjacent sites. In

addition, all eDNA quantities, except *B. Splendens*, sharply decreased with increasing distance

from the AR. In particular, those of *P. trilineatum, P. major*, and *T. japonicus* clearly decreased

even 150 m from AR1 (e.g., W150 and E150). These results are consistent with previous studies

showing a higher density of fishes in ARs than reference sites[2,30,31]. In fact, four dominant OTUs,

P. trilineatum, *Scomber* spp., *P. major*, and *T. japonicus*, were known to appear and aggregate in

ARs[2,31–33]. Only *B. splendens* was not reported to be dependent on ARs, although it is highly

targeted by fisheries and usually caught by vertical longline at midnight in a deeper zone of this

region (depth > 200 m, personal communication, M. Sato). A review paper of the biology of *B.*

splendens indicated that the fisheries grounds of this species around the mouth of Tokyo Bay are

100−500 m in depth, but their juveniles are sometimes collected in a shallower area (<100 m)[34].

Although we cannot exclude the possibility that eDNA from adult *B. splendens* could be detected

in this study, its higher eDNA quantities in the stations of 70☐80 m depth (e.g., AR2, AR1 and

E150) would be likely to reflect the presence of its juveniles. In future studies, combining eDNA

analysis with other methods such as observation by underwater drones or GoPro, which can

determine their body length, would clarify the details of ontogenetic habitat shift for *B. splendens*.

We expected the number of OTUs would be higher at ARs than the surrounding stations, but

did not observe such a pattern. In the following year's survey, we found a larger number of fishes

around AR1 than surrounding stations in a GoPro video (personal observation, N. Inoue, M. Sato,

N, Furuichi), which contradicted our eDNA results. We suspect that higher eDNA concentrations

of dominant OTUs at ARs consumed a large portion of the sequence reads, and species with low

eDNA concentrations were not detected by MiFish metabarcoding in this study.

MiFish metabarcoding also showed different fish species composition between the AR and

surrounding stations based on NMDS and PERMANOVA. The eDNA quantities of fishes,

including dominant OTUs (pelagic fishes: *Scomber* spp., *Etrumeus teres*, *Sardinops*

melanostictus; demersal fishes: *B. splendens*, *P. trilineatum*, *P. major*, *Sardinops melanostictus*,

Sacura margaritacea, *Thamnaconus modestus*) were more abundant in ARs, while those of some

demersal fishes such as flathead (*Platycephalus* sp.2), Asian sheephead wrasse (*Semicossyphus*

reticulatus), barred knifejaw (*Oplegnathus fasciatus*), and seabream (*Dentex tumifrons* or *D.*

cardinalis) were higher in the surrounding stations. Different fish responses to AR structures may

result in differences in the fish community compositions between the AR and surrounding

stations.

Previous studies have found positive relationships between the density of target species and

eDNA concentrations in semi-closed marine systems like estuaries[9,13]; our results found such

associations in an open marine system. We found eDNA quantities of total, demersal, pelagic

fishes, and four dominant species, except *P. major*, positively correlated with the echo intensity

measured by echo sounder. Our study site faces toward the open sea (i.e., Pacific ocean), and the
flow was stronger than in a previous study (5.0–25.0 cm/s in our study and 0.4–13.5 cm/s in a
semi-closed bay at Maizuru, Japan)[35]. Therefore, the flow is likely to transport more eDNAs from
sampling stations in this study, but we could also obtain a strongly localized eDNA signal at
sampling stations. One exception is the eDNA quantity of *P. major*, which did not correlate with
echo intensity. This finding was probably due to elimination of the echo signals around the sea
bottom, where this species is mainly found[18]. Overall, our result indicates that eDNA quantities
reflect spatial distributions of marine fish in an open coastal environment.

We found the eDNA quantities and species number were higher in the bottom sampling layers
than in the middle layers. Multivariate analyses, including NMDS and PERMANOVA, also
showed different fish compositions between the bottom and middle layers. Previous eDNA
studies have also indicated variations in eDNA quantities, the number of species, and community
compositions between different sampling depths[6,9,36,37]. Some of these studies using eDNA
methods suggested that water column stratification and the different species' responses to it cause
the observed different composition between sampling depths[36,37]. Murakami et al.[35] also discussed
that the vertical distribution of eDNA depends on both the depth of the source organisms and the
behavior of eDNA, with the latter dependent on its chemical condition and physical factors in the
environment. Although water column stratification between the middle and bottom layers was not
observed in our study (Fig. S3), we consider that the preference of detected fish species for
bottom habitats or the behavior of eDNA contributed to the higher eDNA quantities and OTU
number in the bottom layer in this study.

Overall, our results demonstrate the effectiveness of quantitative MiSeq sequencing to assess
AR effects on multiple fish species. We found higher quantities of fish eDNA at AR stations than
at surrounding stations and positive correlations between these quantities and eco intensity, which
indicates a highly localized eDNA signal in an oceanographically dynamic environment. This
spatial resolution indicates that quantitative MiSeq sequencing can improve techniques to
monitor the distribution of multiple fish species. Another method, called 'quantitative sequencing
(qSeq).' has also been developed for simultaneous quantifications of eDNA concentrations of
multiple species[38]. This method adopts a stochastic labeling approach with random-tag sequences
and can also quantify eDNA concentrations of fish species using the MiFish primer set[39]. The
quantitative metabarcoding methods (i.e., the Miseq sequencing and qSeq) can complement
conventional methods, especially in an environment with areas where the water is too deep to
adopt underwater visual census (>40m depth) or in no-take marine reserves where exploiting fish
is prohibited.

Data availability

DDBJ accession numbers of the DNA sequences analyzed in the present study are DRA011864
(Submission ID), PRJDB11389 (BioProject ID) and SAMD00298249–SAMD00298281
(BioSample ID).
509

Acknowledgements
We are grateful to the staff of the research vessel *Takamaru* for their support in the field, Tomoe
Ishibashi of Fisheries Technology Institute, and Tomohiro Shirako of IDEA consultants, Inc. for
support in molecular experiments. We also thank Yutaro Kawano of Mie University and Fumie
Yamaguchi of Fisheries Technology Institute for providing fish photos. This work was supported
by the research grants from the National Research Institute of Fisheries Engineering (NRIFE) in
2018□2019 fiscal year.
517

Author Contributions
M. S., R. N., and N. F. designed the study and conducted field surveys. T. I. and N. I. analyzed the
data of echo sounder survey. M. S. and N. I. performed molecular experiments, and M. U.
prepared the internal standards for eDNA quantification. M. S. and N. I. analyzed fish community
dataset extracted from results of MiSeq sequencing. M. S. wrote the first draft of the manuscript
and other authors commented on the manuscript.
524

525   **Additional information**
526   **Supplementary information** accompanies this paper.
527   **Competing Interests**: The authors declare no competing interests.
528

References

1.   Yamane, T. Status and future plans of artificial reef projects in Japan. *Bull. Mar. Sci.* **44**,
1038–1040 (1989)

2.   Inoue, N. *et al.* The present situation of utilization and the effects of artificial reefs on the
resource density of Japanese butterfish *Hyperoglyphe japonica* and red sea bream *Pagrus*
major in the waters of the Iki Islands, Nagasaki, Japan. *Nippon Suisan Gakkaishi* **84**(6),
1010–1016 (Japanese with English abstract) (2018). 10.2331/suisan.17-00081

3.   Polovina, J. J. & Sakai, I. Impacts of artificial reefs on fishery production in Shimamaki,
Japan. *Bull. Mar. Sci.* **44**, 997–1003 (1989)

4.   Kakimoto, H. Control of fish behavior by the artificial reef. *Fish. Eng.* **30**, 59–68 (1993)

5.   Kang, M., Nakamura, T. & Hamano, A. A methodology for acoustic and geospatial analysis of
diverse artificial-reef datasets. *ICES J. Mar. Sci.* **68**(10), 2210–2221 (2011).
10.1093/icesjms/fsr141

6.   Yamamoto, S. *et al.* Environmental DNA metabarcoding reveals local fish communities in a
species-rich coastal sea. *Sci. Rep.* **7**, 40368 (2017). 10.1038/srep40368, Pubmed:28079122

7.   Ushio, M. *et al.* Quantitative monitoring of multispecies fish environmental DNA using
high-throughput sequencing. *Metabarcoding Metagenomics*, e23297 (2018). 10.1101/113472

8.   Barnes, M. A. & Turner, C. R. The ecology of environmental DNA and implications for
conservation genetics. *Conserv. Genet.* **17**(1), 1–17 (2016). 10.1007/s10592-015-0775-4

9.   Yamamoto, S. *et al.* Environmental DNA as a 'snapshot' of fish distribution: A case study of
Japanese jack mackerel in Maizuru Bay, Sea of Japan. *PLOS ONE* **11**(3), e0149786 (2016).
10.1371/journal.pone.0149786, Pubmed:26933889

10.   Port, J. A. *et al.* Assessing vertebrate biodiversity in a kelp forest ecosystem using
environmental DNA. *Mol. Ecol.* **25**(2), 527–541 (2016). 10.1111/mec.13481,
Pubmed:26586544

11.   Miya, M., Gotoh, R. O. & Sado, T. MiFish metabarcoding: A high-throughput approach for
simultaneous detection of multiple fish species from environmental DNA and other samples.
Fish. Sci. 86(6), 939–970. 10.1007/s12562-020-01461-x

12.   Takahara, T., Minamoto, T., Yamanaka, H., Doi, H. & Kawabata, Z. Estimation of fish
biomass using environmental DNA. *PLOS ONE* **7**(4), e35868 (2012).
10.1371/journal.pone.0035868, Pubmed:22563411

13.   Stoeckle, M. Y., Soboleva, L. & Charlop-Powers, Z. Aquatic environmental DNA detects
seasonal fish abundance and habitat preference in an urban estuary. *PLOS ONE* **12**(4),
e0175186 (2017). 10.1371/journal.pone.0175186, Pubmed:28403183

14.   Miya, M. *et al.* Use of a filter cartridge for filtration of water samples and extraction of
environmental DNA. *J. Vis. Exp.* (117) (2016). 10.3791/54741, Pubmed:27911387

15.   Ushio, M. Use of a filter cartridge combined with intra-cartridge bead-beating improves
detection of microbial DNA from water samples. *Methods Ecol. Evol.* **10**(8), 1142–1156
(2019). 10.1111/2041-210X.13204

16.   Miya, M. *et al.* MiFish, a set of universal PCR primers for metabarcoding environmental DNA
from fishes: Detection of more than 230 subtropical marine species. *R. Soc. Open Sci.* **2**(7),
150088 (2015). 10.1098/rsos.150088

17.   Tanabe, A. S. & Toju, H. Two new computational methods for universal DNA barcoding: A
benchmark using barcode sequences of bacteria, Archaea, animals, fungi, and land plants.
PLOS ONE **8**(10), e76910 (2013). 10.1371/journal.pone.0076910, Pubmed:24204702

18.   Froese, R. & Pauly, D. FishBase. *World Wide Web Electronic Publication.* Available from:
www.fishbase.org (2019).

19.   Sato, M., Nakamura, Y. & Hori, M. Potential stocks of reef fish-based ecosystem services in
the Kuroshio Current region: Their relationship with latitude and biodiversity. *Popul. Ecol.*
63(1), 75–91 (2021). 10.1002/1438-390X.12061

20.   Cordeiro, T. C., Barrella, W., Butturi-Gomes, D. & Petrere Júnior, M. A modeling approach
for reposition dynamics of litter composition in coastal areas of the city of Santos, Sao Paulo,
Brazil. *Mar. Pollut. Bull.* **128**, 333–339 (2018). 10.1016/j.marpolbul.2018.01.054,
Pubmed:29571381

21.   Aho, K., Derryberry, D. & Peterson, T. Model selection for ecologists: the worldviews of AIC
and BIC. *Ecology.* **95**(3), 631–636 (2010). 10.1890/13-1452.1

22.   Nielsen, A. *et al.* glmmTMB:Generalized Linear Mixed Models Using Template Model
Builder, 2017

23.   Hothorn, T., Bretz, F., Westfall, P., Heiberger, R. M. & Schuetzenmeister, A. multcomp:
Simultaneous inference in general parametric models. R package version 1.4-15. (2019)

24.   Bartoń, K. MuMIn: Multi-Model Inference. R package version 1.43.17. (2013)

25.   Oksanen, Jari *et al.* vegan：Community ecology package. R package version 2.5□7. (2013)

26.   Wickham, H. ggplot2: An implementation of the Grammar of Graphics. R package version
3.3.2. https://github.com/hadley/ggplot2 (2014).

27.   Kassambara, A. ggpubr: 'ggplot2' Based Publication Ready Plots. R package version 0.4.0.
https://CRAN.R-project.org/package=ggpubr (2018).

28.   Fujii, K. *et al.* Environmental DNA metabarcoding for fish community analysis in backwater
lakes: A comparison of capture methods. *PLOS ONE* **14**(1), e0210357 (2019).
10.1371/journal.pone.0210357, Pubmed:30703107

29.   Oka, S. *et al.* Environmental DNA metabarcoding for biodiversity monitoring of a highly
diverse tropical fish community in a coral reef lagoon: Estimation of species richness and
detection of habitat segregation. *Environ. DNA* 3(1), 55–69 (2021). 10.1002/edn3.132

30.   dos Santos, L. N., Brotto, D. S. & Zalmon, I. R. Fish responses to increasing distance from

602   artificial reefs on the Southeastern Brazilian Coast. *J. Exp. Mar. Biol. Ecol.* **386**(1–2), 54–60

603   (2010). 10.1016/j.jembe.2010.01.018

604   31.   Inoue, N. *et al.* Using standarfized CPUE to estimate the effect of artificial reefs on fish

605   abundance in *Modern Fisheries Engineerng: Realizing a Healthy and Sustainable Marine*

606   *Ecocystem* (eds Bortone, A. S. & Otake, S.) 75–84 (CRC Press, 2020)

607   32.   Noh, J., Ryu, J., Lee, D. & Khim, J. S. Distribution characteristics of the fish assemblages to

608   varying environmental conditions in artificial reefs of the Jeju Island, Korea. *Mar. Pollut. Bull.*

609   **118**(1–2), 388–396 (2017). 10.1016/j.marpolbul.2017.02.066, Pubmed:28258725

610   33.   Nakamura, T. & Hamano, A. Seasonal differences in the vertical distribution pattern of

611   Japanese jack mackerel, Trachurus japonicus: Changes according to age? *ICES J. Mar. Sci.*

612   **66**(6), 1289–1295 (2009). 10.1093/icesjms/fsp114

613   34.   Watari, S. *et al.* Fisheries biology and resource management of splendid alfonsino Beryx

614   splendens. *Bull. Jpn Fish. Res. Educ. Agency*, 1-46 (Japanese with English abstract) (2017)

615   35.   Murakami, H. *et al.* Dispersion and degradation of environmental DNA from caged fish in a

616   marine environment. *Fish. Sci.* **85**(2), 327–337 (2019). 10.1007/s12562-018-1282-6

617   36.   Andruszkiewicz, E. A. *et al.* Biomonitoring of marine vertebrates in Monterey Bay using

618   eDNA metabarcoding. *PLOS ONE* **12**(4), e0176343 (2017). 10.1371/journal.pone.0176343,

619   Pubmed:28441466

620   37.   Jeunen, G. *et al.* Water stratification in the marine biome restricts vertical environmental DNA

621   (eDNA) signal dispersal. *Environ. DNA* **2**(1), 99–111 (2020). 10.1002/edn3.49

622   38.   Hoshino, T. & Inagaki, F. Application of stochastic labeling with random-sequence barcodes

623   for simultaneous quantification and sequencing of environmental 16S rRNA genes. *PLOS*

624   *ONE* **12**(1), e0169431 (2017). 10.1371/journal.pone.0169431, Pubmed:28052139

625   39.   Hoshino, T., Nakao, R., Doi, H. & Minamoto, T. Simultaneous absolute quantification and

626   sequencing of fish environmental DNA in a mesocosm by quantitative sequencing technique.

627   *Sci. Rep.* 11(1), 4372 (2021). 10.1038/s41598-021-83318-6, Pubmed:33623060

630   Figures and Tables

631
632

Figure 1 (a) Location of sampling stations, cruise track, and a set net in Tateyama Bay. Gray areas
indicate landmasses, a gray bold line indicates cruise track, and gray thin lines indicate depth
contours with an interval of 20 m. Pictures of the dominant species, (b) splendid alfonsino
(*Beryx splendens*), (c) chicken grunt (*Parapristipoma trilineatum*), (d) chub mackerel (*Scomber
japonicus*), (e) red seabream (*Pagrus major*), and (f) jack mackerel (*Trachurus japonicus*).
Photograph credits: (b) Fumie Yamaguchi, (c, d, f) Yutaro Kawano, and (e) Masaaki Sato.

642   Figure 2 The relationships between sequence reads and the copy numbers of standard DNAs

643   according to Ushio et al.[7]. (a) The relationships between sequence reads and the copy numbers of

644   standard DNAs with the maximum, median, and minimum slopes. (b) Distribution of R^2 values of

645   the regression lines.

Figure 3 The relationships between distance from AR1 as well as number of OTUs and eDNA copies of total fish, demersal fish, pelagic fish, *Beryx splendens*, *Parapristipoma trilineatum*, *Scomber* spp., *Pagrus major*, and *Trachurus japonicus*. The black line and the gray shaded area represent a median value and 95% confidence interval, respectively, as determined from the generalized linear model. Red and turquoise blue circles indicate values in the bottom and middle sampling layers.

Figure 4 The relationships between eco intensity and eDNA copies of total fish, demersal fish,

pelagic fish, *Beryx splendens*, *Parapristipoma trilineatum*, *Scomber* spp., *Pagrus major*, and

Trachurus japonicus. The black line and the gray shaded area represent a median value and the

95% confidence interval, respectively, as determined from the generalized linear model. Red and

turquoise blue circles indicate values in the bottom and middle sampling layers.

Figure 5 Nonmetric multidimensional scaling (NMDS) for fish communities detected at sampling

stations. NMDS stress was 0.096.

671   Table 1 Details of the eight sampling stations for fish eDNA survey on May 23, 2018

Station	Vertical position	Latitude (N)	Longitude (E)	Depth (m)	eDNA sampling time	eDNA sampling depth (m)	Water temperature (°C)	Salinity
E750	Middle	35.0020	139.7949	94	9:23	47	17.90	34.51
	Bottom				9:34	92	17.26	34.51
E500	Middle	34.9999	139.7937	86	10:08	43	18.50	34.49
	Bottom				10:13	84	17.45	34.49
E150	Middle	34.9971	139.7920	76	10:58	38	18.61	34.49
	Bottom				11:03	74	17.60	34.50
AR1	Middle	34.9957	139.7917	73	11:46	35	18.65	34.49
	Bottom				11:40	71	18.24	34.52
W150	Middle	34.9947	139.7906	71	12:35	35	19.43	34.46
	Bottom				12:39	69	18.67	34.50
W500	Middle	34.9928	139.7876	71	13:13	35	19.41	34.47
	Bottom				13:17	69	19.20	34.48
W750	Middle	34.9916	139.7853	67	13:52	32	19.93	34.44
	Bottom				13:56	65	19.43	34.48
AR2	Middle	34.9938	139.7911	65	14:33	30	19.96	34.48
	Bottom				14:26	63	19.13	34.47

673 Table 2. Parameters estimated from an BIC minimum model based on generalized liner modelling.

	Estimate	SE	Five dominant species	Estimate[b]	SE[b]	Estimate[c]	SE[c]
Number of OTUs			*Beryx splendens* eDNA[a]				
Intercept	2.87	0.06	Intercept	3.02	0.36	-3.30	1.02
Distance			log(Distance+0.1)				
Vertical position (middle)	-0.25	0.10	Vertical position (middle)	-1.64	0.50		
Distance : Vertical position			log(Distance+0.1) : Vertical position (middle)				
Total fish eDNA			*Parapristipoma trilineatum* eDNA[a]				
Intercept	5.42	0.41	Intercept	3.01	0.57	-3.30	1.02
log(Distance+0.1)	-0.30	0.06	log(Distance+0.1)	-0.23	0.10		
Vertical position (middle)	-1.25	0.36	Vertical position (middle)	0.55	0.84		
log(Distance+0.1) : Vertical position (middle)			log(Distance+0.1) : Vertical position (middle)	-0.41	0.15		
Demersal fish eDNA			*Scomber* spp. eDNA				
Intercept	5.08	0.41	Intercept	-1.86	0.41		
log(Distance+0.1)	-0.34	0.06	log(Distance+0.1)	-0.15	0.07		
Vertical position (middle)	-1.17	0.35	Vertical position (middle)	-1.86	0.41		
log(Distance+0.1) : Vertical position (middle)			log(Distance+0.1) : Vertical position (middle)				
Pelagic fish eDNA			*Pargus major* eDNA[a]				
Intercept	4.10	0.45	Intercept	3.21	0.47	-6.16	3.60
log(Distance+0.1)	-0.23	0.06	log(Distance+0.1)	-0.39	0.08	0.62	0.56
Vertical position (middle)	-1.34	0.39	Vertical position (middle)	-5.19	0.69	3.18	1.24
log(Distance+0.1) : Vertical position (middle)			log(Distance+0.1) : Vertical position (middle)	0.46	0.13		
			Trachurus japonicus eDNA[a]				
			Intercept	1.83	0.52	-3.30	1.02
			log(Distance+0.1)	-0.24	0.09		
			Vertical position (middle)	0.14	0.72		
			log(Distance+0.1) : Vertical position (middle)	-0.31	0.13		

[a]Gamma hurdle model was used. [b]An estimate and error of Gamma component when using Gamma or Gamma hurdle model. [c]An estimate and error of binomial component when using Gamma hurdle model.

676 Table 3. Summary results of Gamma or Gamma hurdle models (Null model: model with intercept
677 only; Model with eco intensity: minimum BIC model which includes eco intensity as an
678 explanatory variable) for each response variable of eDNA copy numbers based on automated
679 model selection.

	Intercept[b]	log(eco intensity)[b]	Intercept[c]	log(eco intensity)[c]	BIC	R^2
Total fish eDNA						
Null model	4.16				325.60	0.00
Model with eco intensity	18.97	0.94			315.40	0.41
Demersal fish eDNA						
Null model	3.80				299.20	0.00
Model with eco intensity	19.88	1.02			288.20	0.45
Pelagic fish eDNA						
Null model	2.98				251.90	0.00
Model with eco intensity	15.50	0.79			244.70	0.33
Beryx splendens eDNA[a]						
Null model	3.04		-3.43		235.70	0.00
Model with eco intensity	18.84	1.00	-3.43		230.70	0.44
Parapristipoma trilineatum eDNA[a]						
Null model	2.66		-3.43		202.20	0.00
Model with eco intensity	20.60	1.17	-3.43		184.80	0.52
Scomber spp. eDNA						
Null model	2.56				212.90	0.00
Model with eco intensity	16.32	0.87			207.60	0.37
Pargus major eDNA[a]						
Null model	2.04		-0.94		150.60	0.00
Model with eco intensity	13.99	0.75	-0.94		153.60	0.30
Trachurus japonicus eDNA[a]						
Null model	1.22		-2.71		118.50	0.00
Model with eco intensity	15.62	0.93	-2.71		130.70	0.40

[a]Gamma hurdle model was used. [b]An estimate and error of Gamma component when using Gamma hurdle model. [c]An estimate and error of binomial component when using Gamma hurdle model.