Title: Intracellular and intercellular gene regulatory networks inference from time-course

individual RNA-Seq.

Authors: Makoto Kashima[1*], Yuki Shida[2], Takashi Yamashiro[2], Hiromi Hirata[1], and

Hiroshi Kurosaka[2]

[1]College of Science and Engineering, Aoyama Gakuin University, 5-10-1, Fuchinobe,

Sagamihara Chuo Ku, Kanagawa 252-5258, Japan;

[2]Department of Orthodontics and Dentofacial Orthopedics, Osaka University, Yamadaoka,

Suita, Osaka, 565-0871, Japan.

* Corresponding author

Structured Abstract

Motivation: Recent technical advances in bulk RNA-Seq have enabled time-course

RNA-Seq of whole individual embryos to understand the underlying molecular

mechanisms. Thus, we hypothesized that gene regulatory networks (GRNs) inferred from

time-course individual RNA-Seq during embryonic development reveal intercellular

regulatory relationships involved in signaling pathways.

Results: Time-course bulk RNA-Seq of individual mouse embryos in early development,

followed by pseudo-time analysis and GRN inference, demonstrated that GRN inference

from RNA-Seq with pseudo-time can be applied for individual bulk RNA-Seq similar to

scRNA-Seq. Validation using an experimental-source-based database showed that our

approach could significantly infer GRN for all transcription factors in the database.

Furthermore, the inferred ligand-related and receptor-related downstream genes were

significantly overlapped. Overall, the inferred GRN include intracellular as well as

intercellular regulatory relationships, which cannot be inferred from scRNA-Seq. Thus,

inferring GRN from time-course bulk RNA-Seq is a novel approach for understanding

the regulatory relationships underlying cellular events.

Availability and Implementation: The R source code used in the present study is

available at https://github.com/Makoto-Kashima/TIODE

Contact: kashima@chem.aoyama.ac.jp

Supplementary information: Supplementary data are available at Bioinformatics online.

30 1   Introduction

Regulation of gene expression is a fundamental factor that controls cellular events

such as proliferation and differentiation. Understanding gene regulatory

interconnections is an important step to understand the molecular mechanisms

underlying cellular events. Recently, gene regulatory network (GRN) inference based

on time-course data has attracted much attention in single-cell RNA-Seq (scRNA-

Seq). State-of-the-art scRNA-Seq analysis techniques can generate transcriptome

information from thousands of cells. (Sasagawa *et al.*, 2018, 2013; Gao *et al.*, 2020;

Hayashi *et al.*, 2018; Klein *et al.*, 2015). Dimensional reduction of scRNA-Seq data

provides a trajectory of cellular events and enables assignment of pseudo-time to

each cell during differentiation and proliferation (Treutlein *et al.*, 2014; Haghverdi *et*

al., 2016). Because scRNA-Seq with pseudo-time is a dense time-course observation

of cellular events, gene regulatory networks can be inferred by comparing the timing

of gene upregulation and downregulation along pseudo-time (Matsumoto *et al.*,

2017; Aalto *et al.*, 2020).

Originally, GRN inference was applied for gene expression data from tissue and

pooled cells (bulk samples), generated using DNA microarray and RNA-Seq

[reviewed in (Ko and Brandizzi, 2020; Fernandez-Valverde *et al.*, 2018)]. Compared

to steady-state data, time-course data allows GRN inference by comparing the timing

of gene upregulation and downregulation (Iglesias-Martinez *et al.*, 2016; Zhang *et al.*, 2019; Krouk *et al.*, 2010; Ogami *et al.*, 2012). However, GRN inference from time-course data of bulk samples has not been popular due to the following drawbacks: 1) Transcriptome analysis of a large number of bulk samples is expensive (Spies and Ciaudo, 2015), and 2) biological variance results in inconsistency between real sampling time and transcriptome status (Kilfoil *et al.*, 2009). Fortunately, recent technical advances related to bulk RNA-Seq have overcome these difficulties. Advances in sequencing platforms and bulk 3′ RNA-Seq library preparation methods have enabled cost-effective time-course bulk RNA-Seq (Muir *et al.*, 2016; Alpern *et al.*, 2019; Kamitani *et al.*, 2019; Li *et al.*, 2020). Because these methods were developed based on the methods for scRNA-Seq, they can also be applied for individual RNA-Seq (Kashima *et al.*, 2020) (for instance, RNA-Seq of a single embryo or tissue of individuals with several ng of total RNA). Pseudo-time analysis for individual RNA-Seq might capture individual differences of gene expression in the time-course of biological events such as development. After assignment of pseudo-time to each individual RNA-Seq data, GRN can be inferred similar to an inference based on scRNA-Seq.

Theoretically, GRN inferred from whole-body and tissue RNA-Seq are different from those inferred from scRNA-Seq. scRNA-Seq provides transcriptomic information at the cellular level that enables inference of intracellular GRN involved in the progression of proliferation and differentiation. In contrast, bulk RNA-Seq of whole body and tissues could contain transcriptomic information at the level of cell populations. Time course individual RNA-Seq during development would enable inference of both intracellular and intercellular GRN (cell-cell communications)

involved in the developmental process. For instance, during embryonic development,

upregulation of ligand-related genes can be followed by upregulation of downstream

genes in cells expressing receptor genes.

We thus hypothesized that GRNs inferred from time-course individual RNA-Seq

during embryonic development would include intercellular regulatory relationships

between ligand genes and downstream genes of related signaling pathways. To test

this hypothesis in the present study, we conducted time-course bulk RNA-Seq of

individual mouse embryos in early development, followed by pseudo-time analysis

and GRN inference.

2 Materials and methods

2.1 Maintenance of mice

Embryos were collected from pregnant female Institute of Cancer Research (ICR) mice

(CLEA, Tokyo, Japan) at each stage. All sacrificed female mice were housed under a 12

h dark-light cycle in which the light phase started from 8 am. All animal experiments were

performed in accordance with the guidelines of the Animal Care and Use Committee of

Osaka University Graduate School of Dentistry, Osaka, Japan.

2.2 RNA extraction

Total RNA extraction was performed using the RNeasy® kit (QIAGEN, Hilden,

Germany) according to manufacturer's protocol. The total RNA concentration

was measured using Qubit™ RNA HS Assay Kit (Thermo Fisher Scientific,

Waltham, MA, USA) and was adjusted to 5 ng/ul, and stored at -80 °C until

subsequent analysis.

2.3 RNA-Seq library preparation and sequencing

Non targeted RNA-Seq were conducted according to the Lasy-Seq ver. 1.1

protocol (https://sites.google.com/view/lasy-seq/) (Kamitani *et al.*, 2019; Kashima *et al.*, 2020). Briefly, 50 ng of total RNA were reverse-transcribed using an RT primer with index and SuperScript IV reverse transcriptase (Thermo Fisher Scientific, Waltham, MA, USA). Then, all RT mixtures of the samples were pooled and purified using an equal volume of AMpure XP beads (Beckman Coulter, Brea, CA, USA) according to the manufacturer's instructions. Second strand synthesis was conducted on the pooled samples using RNaseH (5 U/µL, Enzymatics, Beverly, MA, USA), and DNA polymerase I (10 U/µL, Enzymatics, Beverly, MA, USA). To avoid the carryover of large amounts of rRNAs, the mixture was subjected to RNase treatment using RNase T1 (Thermo Fisher Scientific, Waltham, MA, USA). Then, purification was conducted with a 0.8× volume of AMpure XP beads. Fragmentation, end-repair, and A-tailing were conducted using 5× WGS Fragmentation Mix (Enzymatics, Beverly, MA, USA). The Adapter for Lasy-Seq was ligated using 5× Ligation Mix (Enzymatics, Beverly, MA, USA), and the adapter-ligated DNA was purified twice with a 0.8× volume of AMpure XP beads. After optimisation of PCR cycles for library amplification by qPCR using Evagreen, 20× in water (Biotium, Fremont, CA, USA) and the QuantStudio5 Real-Time PCR System (Applied Biosystems, Waltham, MA, USA), the library was amplified using KAPA HiFi HotStart ReadyMix (KAPA BIOSYSTEMS, Wilmington, MA, USA) on the ProFlex PCR System (Applied Biosystems, Waltham, MA, USA). The amplified library was purified with an equal volume of AMpure XP beads. One microliter of the library was then used for electrophoresis using a Bioanalyzer 2100 with the Agilent High Sensitivity DNA kit (Agilent Technologies, Santa Clara, CA, USA)

| 121 | to check for quality. Then, sequencing of 150-bp paired-end reads was performed |
| 122 | using HiSeq X Ten (Illumina, San Diego, CA, USA). |

2.4 Mapping and gene quantification

Read 1 reads were processed with fastp (version 0.21.0) (Chen *et al.*, 2018) using the following parameters: --trim_poly_x -w 20 -- adapter_sequence=AGATCGGAAGAGCACACGTCTGAACTCCAGTCA -- adapter_sequence_r2=AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTG T -l 31. The trimmed reads were then mapped to the mouse reference sequences of Mus_musculus.GRCm38.cdna.all.fa, using BWA mem (version 0.7.17-r1188) (Li and Durbin, 2009) with the default parameters. The read count for each gene was calculated with salmon using -l IU, which specifies the library type (version v0.12.0) (Patro *et al.*, 2017). Then, using R (version 4.0.1) (R Core Team, 2017), the sums of read counts per gene were calculated. Genes with read counts greater than zero were used in the subsequent analysis.

2.5 Pseudo-time analysis

Read counts were normalized using the "NormalizeData" function with the default parameters in Seurat (version 4.0.0) (Hao *et al.*, 2020). For principal component analysis (PCA), the normalized read counts were scaled using the "ScaleData" function with the default parameters except for do.scale = F. Then, PCA was conducted using the "RunPCA" function for genes with high dispersion, which were selected using the "FindVariableFeatures" function with default parameters except for selection.method = "mvp". Finally, SingleCellExperiment (version 1.10.1) (Amezquita *et al.*, 2020) and slingshot (version 1.6.1) (Street *et al.*, 2018) were used to calculate the pseudo-time for

145	each sample.

146	**2.6**	**Gene regulatory network inference**

To infer GRN, A was optimized 20 times using the SCODE algorithm

(Matsumoto *et al.*, 2017) with one hundred iterations and $D = 4$. Pearson's

correlation coefficients between values of each A from the 20 optimizations

and the mean$A$ (the average of each value of A) were calculated. In the

following analysis, we used the average values of the top 10 A showing higher

correlations with the mean$A$ from 20 optimizations. To define the thresholds

for downstream gene selection for each gene, the linear function was regressed

using the "nls" function in R for the scatter plot of absolute values of A for

downstream genes in the decreasing order, and the X and Y axes showed the

integers from one to 28117, and the absolute values of A, respectively. Genes

with larger absolute values of A compared to the Y values of the regressed line

were defined as the inferred gene downstream of each gene.

159	**2.7**	**Evaluation of inferred GRN by comparing with a TF-downstream gene**
160		**database**

For validating the inferred GRN, the information regarding the binding motifs

and targets for 438 mouse TFs in the TF2DNA database was used (Pujato *et al.*,

2014; Badis *et al.*, 2009; Berger *et al.*, 2006; Mathelier *et al.*, 2014; Matys *et*

al., 2006; Weirauch *et al.*, 2014; Berger *et al.*, 2008; Chen *et al.*, 2011; Jolma

et al., 2013; Sebé-Pedrós *et al.*, 2013; Wei *et al.*, 2010; Weirauch *et al.*, 2013).

Area under the curve (AUC) values for downstream prediction based on the

absolute values of A were calculated using the "performance" function in

ROCR (version 1.0-11) (Sing *et al.*, 2005). Statistical analysis for the

169 enrichment of validated target genes among the inferred genes was conducted

170 using the "fisher.test" function in R. Statistical analysis for the overlap of

171 inferred downstream genes for ligand-related and receptor-related genes was

172 conducted using the "enrichment_test" function in Rvenn (version 1.1.0)

173 (Akyol, 2019). For all statistical tests, BH correction was performed using the

174 "p.adjust" function. The upset plots were drawn using the "upset" function in

175 UpSetR (version 1.4.0) (Conway *et al.*, 2017).

176 **2.8** **Data availability**

177 All RNA-Seq sequence data is deposited in PRJNA725414 of SRA.

178 **3** **Results**

179 **3.1** **Pseudo-time analysis of the time-course bulk RNA-Seq for mouse embryos**

180 To examine whether gene regulatory network (GRN) inference based on time-course

181 bulk RNA-Seq works well, we conducted time-course bulk RNA-Seq for individual

182 mouse embryos. RNA was extracted from each individual whole embryo (n = 10 or

183 11) at seven time points: E7.5, E8.5, E9.5, E10.5, E11.5, E12.5, and E.13.5, followed

184 by 3′ RNA-Seq using the 'Lasy-Seq' method (Kamitani *et al.*, 2019). As a result, we

185 obtained 76 RNA-Seq datasets with an average of 8.5 million reads per sample. The

186 reads were mapped onto the mouse reference sequence, followed by calculation of

187 the read counts of each gene in each sample. We then used 'Seurat' (Stuart *et al.*,

188 2019), an R package used for single cell omics analysis, normalization of read counts,

189 detection of highly variable genes, dimension reduction of omics date, and so on. On

190 the PC1 and PC2 plane obtained with Seurat, each sample at the same stage was close

191 to each other (Fig. 1A). As expected, the clusters of each stage were ordered

192 according to the developmental process (Fig. 1A). Using an R package 'slingshot'

(Street *et al.*, 2018), we inferred the developmental trajectory of mouse embryos and

calculated the pseudo-time for each sample (Fig. 1 A and B). Pseudo-time analysis

revealed individual differences among embryos in the speed of their developmental

processes. For example, the pseudo-time of a sample at E10.5 and of E11.5 were

almost the same (Fig. 1 A and B). Using the pseudo-time analysis, the time-course

RNA-Seq data of seven time points could be converted into those of 76 time points.

3.2   Gene regulatory network inference based on individual RNA-Seq of whole

mouse embryos

Next, we inferred a GRN from the dataset of time-course individual RNA-Seq of

whole mouse embryos. We used SOCDE, which solves linear ordinary differential

equations to infer GRN (Matsumoto *et al.*, 2017). To avoid incorrect emphasis on the

technical noise of RNA-Seq, we used a non-scaled normalized gene expression matrix

as the input for SCODE. Selection of the size of *D,* a parameter of SCODE, is

important for robust GRN inference, because a needlessly large *D* results in an

unstable inferred GRN (Matsumoto *et al.*, 2017). In this study, we used *D* = 4 similar

to the previous study (Matsumoto *et al.*, 2017), with which the residual sum of squares

(RSS) was relatively small (Fig. 2A). For 28117 genes whose read counts were greater

than zero, SCODE produced *A* (28117 × 28117 matrix) corresponding to the inferred

gene regulatory network, in which the value of $A_{i,j}$ indicates regulatory effects on the

downstream gene *i* from the regulator *j*. $A_{i,j} > 0$ indicates that the regulator *j* positively

regulates gene *i,* whereas $A_{i,j} < 0$ indicates the opposite. Because SCODE optimizes

A by random sampling, we optimized *A* 20 times to check for reproducibility. Then,

Pearson's correlation coefficients between the values of each *A* from the 20

optimizations and the mean*A* were calculated (Fig. 2B). Almost all optimizations

produced similar A with high correlation coefficients (Fig. 2B). In the subsequent

analysis, we used the average values of the top 10 A showing higher correlations with

the mean A from 20 optimizations. We then tried to define the thresholds for significant

regulatory relationships between regulators and downstream genes. Because the

average expression level of downstream genes showed higher correlation with the

absolute values of A compared to the regulators (Fig. 2 C and D), we independently

defined a threshold for each regulator as follows. For example, absolute values of A

indicating regulatory relationships between *Sox8* and its regulators showed a positive

correlation (Pearson's correlation coefficient = 0.76) (Fig. 2E), whereas absolute

values of A showing regulatory relationships between *Sox8* and its downstream genes

showed smaller correlations (Pearson's correlation coefficient = 0.54) (Fig. 2F).

Although most of the inferred values of A for regulatory relationships between *Sox8*

and its downstream genes were around zero, some values were outliers (Fig. 2G). To

define the threshold for the downstream genes of *Sox8*, we regressed the linear

function for the scatter plot of absolute A values for downstream genes in the

decreasing order, and defined the threshold (See Material and methods) (Fig. 2G).

Finally, in the subsequent analysis, genes with larger absolute values of A compared

to the threshold were defined as the inferred genes downstream of *Sox8* (Fig. 2G).

3.3   Validation of the inferred network

Next, we evaluated the inferred GRN by comparing the inferred genes downstream

of the TFs with the information in the TF2DNA database, which is an experimental-

source-based database of the binding motifs and downstream genes for 438 mouse

TFs (Pujato *et al.*, 2014; Badis *et al.*, 2009; Berger *et al.*, 2006; Mathelier *et al.*, 2014;

Matys *et al.*, 2006; Weirauch *et al.*, 2014; Berger *et al.*, 2008; Chen *et al.*, 2011; Jolma

et al., 2013; Sebé-Pedrós *et al.*, 2013; Wei *et al.*, 2010; Weirauch *et al.*, 2013). First,

we evaluated the effectiveness of target prediction based on the absolute values of A

by calculating the area under the curve (AUC) (Fig. 3A). The average AUC was 0.704,

indicating that the inferred regulatory relationships could reflect the true regulatory

network. Second, we examined the validity of the defined thresholds by checking the

differences between the validated target gene rate (number of validated target genes /

number of all inferred downstream genes) above the defined thresholds and the best

validated rate (Fig. 3B). Below the thresholds, validated target gene rates for the 438

TFs were 0.69% smaller than the best validated target gene rates on average. The

validated target genes were 67.5% of the inferred downstream genes on average

(Table S1, Fig. 3C and D). Compared with the background rate (number of known

target genes / number of all genes), the validated target gene rates of the inferred

downstream genes of all TFs were statistically higher (adjusted p-value < 0.01) (Fig.

3D). These results suggest that our approach could infer the GRN underlying early

development in mice.

3.4 The inferred network contained regulatory relationships involved in cell-

cell interaction

Our GRN inference was based on bulk RNA-Seq containing the information of all

cells in the body. We thus hypothesized that the inferred GRN also included the

intercellular regulatory network. To examine this possibility, we checked the overlaps

of inferred downstream genes for genes related to the ligands and receptors of nine

major signaling pathways (Fig. 4A): Wnt/βcatenin, TNF, TGF-β, Hedgehog, FGF,

EGF, Delta/Notch, BMP, and Retinoic acid (RA) signaling pathways (Table S2). As

expected, the inferred downstream genes of all pairs of ligand and receptor genes were

significantly overlapped (adjusted p-value < 0.01) (Fig. 4A). On average, 94.2% of

the inferred downstream genes for the ligand-related and receptor-related genes were

overlapped (Fig. 4A). For example, 4510 genes were inferred as the downstream of

Wnt genes, and 4545 genes were downstream of *Fzd* genes. However, 97.5% of the

inferred downstream of *Wnt* genes were also inferred as the downstream of *Fzd* genes

(Fig. 4A). Because SCODE can infer whether each downstream gene is positively or

negatively regulated (Matsumoto *et al.*, 2017), we checked the overlaps of positively

and negatively regulated gene downstream of the representative ligand-related and

receptor-related genes of each signaling pathway, i.e., with the largest number of

inferred downstream genes among each gene family (Fig. 4 B-D and Supplementary

Figure 1,2). For six signaling pathways (TNFβ, TGF, FGF, EGF, Delta/Notch, and

BMP signaling pathways), which activate the downstream genes only when ligands

bind to the receptors (Gilbert and Barresi, 2017), the regulatory directions of ligand-

related and receptor-related genes for most inferred downstream genes were the same

(Fig. 4B and Supplementary Figure 1). In contrast, in case of the RA signaling

pathway, wherein RA receptors function as transcriptional repressors without RA

binding (Supplementary Figure 2A) (Glass and Rosenfeld, 2000), the regulatory

directions of a ligand-related gene, *Aldh1a3,* which encodes a protein involved in RA

synthesis, and a receptor-related gene, *Rara,* for the downstream genes were opposite

(Fig. 4C). In case of the Hedgehog signaling pathway, the regulatory directions of

Ssh2 and *Ptchd4* tended to be opposite (Fig. 4D). The regulatory directions of *Ssh2*

and *Gli3* for the downstream genes were also opposite (Fig. 4D). In the absence of

Hedgehog ligands, the full-length Gli family proteins are ubiquitinated and function

as repressors (Supplementary Figure 2B) (Skoda *et al.*, 2018). With the binding of

Hedgehog ligands, PATCHED inhibits the degradation of Gli family proteins and

allows them to function as transcriptional activators (Supplementary Figure 2A)

(Skoda *et al.*, 2018). Thus, the overlap of downstream genes that are positively and

negatively regulated by *Ssh2, Ptchd4,* and *Gli3* is reasonable. In case of the

Wnt/βcatenin signaling pathway, the regulatory directions of *Wnt7a* and *Fzd5* for the

downstream were opposite (Supplementary Figure 3A). Our time-course RNA-Seq

revealed that among the *Fzd* and *Wnt* genes, *Fzd5* and *Wnt8a,* but not *Wnt7a,* were

the only genes expressed dominantly in the early developmental stage among the

protein families. (Supplementary Figures 4 and 5). Similar regulatory directions were

found for most of the inferred downstream genes of *Wnt8a-Fzd5*, which could be a

functional pair in the early developmental stages. Furthermore, the regulatory

directions for most of the inferred downstream genes of *Wnt7a-Fzd6* (*Fzd* genes with

the second most inferred downstream genes) were also the same (Supplementary

Figure 3). In conclusion, our approach allowed successful inference the intercellular

regulatory relationships related to the major signaling pathways as well as the

intracellular pathways related to TFs.

4   **Discussion**

Recently, GRN inference based on the combination of scRNA-Seq and pseudo-time

analysis has attracted much attention (Dai *et al.*, 2020). However, to our knowledge,

this study is the first to report GRN inference based on the combination of individual

bulk RNA-Seq and pseudo-time analysis. Unlike scRNA-Seq, which can provide

transcriptomic dynamics in a certain cellular event such as proliferation and

differentiation, bulk RNA-Seq could provide a mixture of various transcriptomic

dynamics regarding the cellular events occurring in an embryo (Chasman and Roy,

2017). Theoretically, GRN inference based on bulk RNA-Seq cannot provide

multiple lineage-specific regulatory relationships. This might be the reason why

GRN inference based on bulk RNA-Seq has not been attempted as with scRNA-Seq.

In this study, we succeeded in inferring the known regulatory relationships of the TFs

in the TF2DNA database with high AUC compared to the GRN inferred from

scRNA-Seq (Chen and Mar, 2018) (Fig. 3). This suggests that GRN inference from

bulk RNA-Seq can provide insights into the regulatory interconnections of genes,

even though GRN inferred from bulk RNA-Seq would have limitations in the

comprehensiveness of inferred regulatory relationships compared to those from

scRNA-Seq. Furthermore, drastic changes in the cell population occur during early

developments. It is possible that our GRN inferred from bulk RNA-Seq merely

reflect the changes in cell populations but not the interconnections of genes.

Considering the results of the inference for *Wnt* and *Fzd* genes (Supplementary

Figure 2), these points should be considered when interpreting the inferred GRN

based on the time-course bulk RNA-Seq.

Assignment of pseudo-time is an important step in GRN inference from time-

course data. In case of scRNA-Seq, the accuracy of pseudo-time assignment is

controversial (Tritschler *et al.*, 2019). In contrast, in the case of time-course

individual bulk RNA-Seq, the accuracy of pseudo-time assignment could be assured

by the actual sampling time, which can be an advantage compared to GRN inference

from scRNA-Seq.

GRN indicate the intracellular interconnections of genes in a narrow sense;

intercellular regulation of genes via cell-cell communication is also a key factor to

understand the regulatory mechanisms underlying multicellular organisms. Several

studies have tried to systematically identify cell-cell communications based on single

cell gene expression profiles and information regarding ligand-receptor pairs (Kumar

et al., 2018; Jin *et al.*, 2021; Cabello-Aguilar *et al.*, 2021; Wang *et al.*, 2019). Because

these approaches required prior knowledge, they could only be applied for the major

model organisms and could not reveal novel signaling pathways. We demonstrated

that GRN inference based on time-course individual RNA-Seq could infer

intercellular regulatory relationships related to cell-cell communication via cell

signaling pathways. This approach only requires time-course RNA-Seq results and

is applicable for non-model organisms without ligand-receptor information.

Theoretically, the GRN inferred based on scRNA-Seq cannot include intercellular

interconnections of genes. Comparison of GRN from scRNA-Seq and bulk RNA-

Seq would be useful for systematic identification of genes involved in cell-cell

communications during cellular events of interest. Taken together, our approach is a

powerful tool for understanding intracellular and intercellular regulatory

relationships of genes, which cannot be achieved using existing GRN inferences

based on scRNA-Seq alone. As discussed above, bulk RNA-Seq has limitations in

the comprehensiveness of inferred regulatory relationship because multiple lineage-

specific regulatory relationships cannot be deduced from GRN inference based on

bulk RNA-Seq. A future novel bioinformatic approach that can deconvolute gene

expression in each tissue and cell lineage from bulk RNA-Seq will help overcoming

this limitation.

Finding: This work was supported by Japan Society for the Promotion of Science [grant

number JP19H03858 to H.K.].

Disclosures: Conflicts of interest: No conflicts of interest declared.

Acknowledgments. We would like to thank Editage (www.editage.com) for English
language editing.

References

Aalto,A. *et al.* (2020) Gene regulatory network inference from sparsely sampled noisy
data. *Nat. Commun.*, **11**, 3493.

Akyol,T.Y. (2019) RVenn: An R package for set operations on multiple sets.

Alpern,D. *et al.* (2019) BRB-seq: ultra-affordable high-throughput transcriptomics
enabled by bulk RNA barcoding and sequencing. *Genome Biol.*, **20**, 71.

Amezquita,R.A. *et al.* (2020) Orchestrating single-cell analysis with Bioconductor. *Nat.
Methods*, **17**, 137–145.

Badis,G. *et al.* (2009) Diversity and complexity in DNA recognition by transcription
factors. *Science (80-.).*, **324**, 1720–1723.

Berger,M.F. *et al.* (2006) Compact, universal DNA microarrays to comprehensively
determine transcription-factor binding site specificities. *Nat. Biotechnol.*, **24**,
1429–1435.

Berger,M.F. *et al.* (2008) Variation in Homeodomain DNA Binding Revealed by High-
Resolution Analysis of Sequence Preferences. *Cell*, **133**, 1266–1276.

Cabello-Aguilar,S. *et al.* (2021) SingleCellSignalR: Inference of intercellular networks
from single-cell transcriptomics. *Nucleic Acids Res.*, **48**, 55.

Chasman,D. and Roy,S. (2017) Inference of cell type specific regulatory networks on
mammalian lineages. *Curr. Opin. Syst. Biol.*, **2**, 130–139.

Chen,L. *et al.* (2011) hmChIP: a database and web server for exploring publicly
available human and mouse ChIP-seq and ChIP-chip data. *Bioinformatics*, **27**,

1447–1448.

Chen,S. *et al.* (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics, **34**, i884–i890.

Chen,S. and Mar,J.C. (2018) Evaluating methods of inferring gene regulatory networks

highlights their lack of performance for single cell gene expression data. *BMC*

Bioinformatics, **19**, 232.

Conway,J.R. *et al.* (2017) UpSetR: an R package for the visualization of intersecting

sets and their properties. *Bioinformatics*, **33**, 2938–2940.

Dai,H. *et al.* (2020) Reconstructing gene regulatory networks in single-cell

transcriptomic data analysis. *Zool. Res.*, **41**, 599–604.

Fernandez-Valverde,S.L. *et al.* (2018) Inference of developmental gene regulatory

networks beyond classical model systems: New approaches in the post-genomic

era. *Integr. Comp. Biol.*, **58**, 640–653.

Gao,C. *et al.* (2020) The Comparison of Two Single-cell Sequencing Platforms: BD

Rhapsody and 10x Genomics Chromium. *Curr. Genomics*, **21**, 602–609.

Gilbert,S.F. and Barresi,M.J.F. (2017) DEVELOPMENTAL BIOLOGY, 11TH

EDITION 2016. *Am. J. Med. Genet. Part A*, **173**, 1430–1430.

Glass,C.K. and Rosenfeld,M.G. (2000) The coregulator exchange in transcriptional

functions of nuclear receptors. *Genes Dev.*, **14**, 121–141.

Haghverdi,L. *et al.* (2016) Diffusion pseudotime robustly reconstructs lineage

branching. *Nat. Methods*, **13**, 845–848.

Hao,Y. *et al.* (2020) Integrated analysis of multimodal single-cell data. *bioRxiv*,

2020.10.12.335331.

Hayashi,T. *et al.* (2018) Single-cell full-length total RNA sequencing uncovers

dynamics of recursive splicing and enhancer RNAs. *Nat. Commun.*, **9**, 1–16.

Iglesias-Martinez,L.F. *et al.* (2016) BGRMI: A method for inferring gene regulatory

networks from time-course gene expression data and its application in breast

cancer research. *Sci. Rep.*, **6**, 1–12.

Jin,S. *et al.* (2021) Inference and analysis of cell-cell communication using CellChat.

Nat. Commun., **12**, 1–20.

Jolma,A. *et al.* (2013) DNA-binding specificities of human transcription factors. *Cell*,

152, 327–339.

Kamitani,M. *et al.* (2019) Lasy-Seq: a high-throughput library preparation method for

RNA-Seq and its application in the analysis of plant responses to fluctuating

temperatures. *Sci. Rep.*, **9**, 7091.

Kashima,M. *et al.* (2020) DeLTa-Seq: direct-lysate targeted RNA-Seq from crude tissue

lysate. (8/15 words) Running Title: Development of direct-lysate targeted RNA-

Seq method Corresponding Author. *bioRxiv*, 2020.09.15.299180.

Kilfoil,M.L. *et al.* (2009) Stochastic variation: From single cells to superorganisms.

HFSP J., **3**, 379–385.

Klein,A.M. *et al.* (2015) Droplet barcoding for single-cell transcriptomics applied to

embryonic stem cells. *Cell*, **161**, 1187–1201.

Ko,D.K. and Brandizzi,F. (2020) Network-based approaches for understanding gene

regulation and function in plants. *Plant J.*, **104**, 302–317.

Krouk,G. *et al.* (2010) Predictive network modeling of the high-resolution dynamic

plant transcriptome in response to nitrate. *Genome Biol.*, **11**, R123.

Kumar,M.P. *et al.* (2018) Analysis of Single-Cell RNA-Seq Identifies Cell-Cell

Communication Associated with Tumor Characteristics. *Cell Rep.*, **25**, 1458-

1468.e4.

Li,H. and Durbin,R. (2009) Fast and accurate short read alignment with Burrows-
435   Wheeler transform. *Bioinformatics*, **25**, 1754–1760.

436   Li,Y. *et al.* (2020) Decode-seq: A practical approach to improve differential gene
437   expression analysis. *Genome Biol.*, **21**, 66.

438   Mathelier,A. *et al.* (2014) JASPAR 2014: An extensively expanded and updated open-
439   access database of transcription factor binding profiles. *Nucleic Acids Res.*, **42**,
440   D142–D147.

441   Matsumoto,H. *et al.* (2017) SCODE: an efficient regulatory network inference
442   algorithm from single-cell RNA-Seq during differentiation. *Bioinformatics*, **33**,
443   2314–2321.

444   Matys,V. *et al.* (2006) TRANSFAC and its module TRANSCompel: transcriptional
445   gene regulation in eukaryotes. *Nucleic Acids Res.*, **34**, D108.

446   Muir,P. *et al.* (2016) The real cost of sequencing: scaling computation to keep pace with
447   data generation. *Genome Biol.*, **17**, 53.

448   Ogami,K. *et al.* (2012) Computational gene network analysis reveals TNF-induced
449   angiogenesis. *BMC Syst. Biol.*, **6**, 3–8.

450   Patro,R. *et al.* (2017) Salmon provides fast and bias-aware quantification of transcript
451   expression. *Nat. Methods*, **14**, 417–419.

452   Pujato,M. *et al.* (2014) Prediction of DNA binding motifs from 3D models of
453   transcription factors; identifying TLX3 regulated genes. *Nucleic Acids Res.*, **42**,
454   13500–13512.

455   R Core Team (2017) R: A language and environment for statistical computing. R
456   Foundation for Statis-tical Computing, Vienna, Austria.

Sasagawa,Y. *et al.* (2013) Quartz-Seq: a highly reproducible and sensitive single-cell

RNA-Seq reveals non-genetic gene expression heterogeneity. *Genome Biol.*, **14**,

R31.

Sasagawa,Y. *et al.* (2018) Quartz-Seq2: A high-throughput single-cell RNA-sequencing

method that effectively uses limited sequence reads. *Genome Biol.*, **19**, 1–24.

Sebé-Pedrós,A. *et al.* (2013) Early eèolution of the T-box transcription factor family.

Proc. Natl. Acad. Sci. U. S. A., **110**, 16050–16055.

Sing,T. *et al.* (2005) ROCR: visualizing classifier performance in R. *Bioinformatics*, **21**,

3940–3941.

Skoda,A.M. *et al.* (2018) The role of the hedgehog signaling pathway in cancer: A

comprehensive review. *Bosn. J. Basic Med. Sci.*, **18**, 8–20.

Spies,D. and Ciaudo,C. (2015) Dynamics in Transcriptomics: Advancements in RNA-

seq Time Course and Downstream Analysis. *Comput. Struct. Biotechnol. J.*, **13**,

469–477.

Street,K. *et al.* (2018) Slingshot: Cell lineage and pseudotime inference for single-cell

transcriptomics. *BMC Genomics*, **19**, 477.

Stuart,T. *et al.* (2019) Comprehensive Integration of Single-Cell Data. *Cell*, **177**, 1888-

1902.e21.

Treutlein,B. *et al.* (2014) Reconstructing lineage hierarchies of the distal lung

epithelium using single-cell RNA-seq. *Nature*, **509**, 371–375.

Tritschler,S. *et al.* (2019) Concepts and limitations for learning developmental

trajectories from single cell genomics. *Dev.*, **146**.

Wang,S. *et al.* (2019) Cell lineage and communication network inference via

optimization for single-cell transcriptomics. *Nucleic Acids Res.*, **47**, 66.

Wei,G.H. *et al.* (2010) Genome-wide analysis of ETS-family DNA-binding in vitro and

in vivo. *EMBO J.*, **29**, 2147–2160.

Weirauch,M.T. *et al.* (2014) Determination and inference of eukaryotic transcription

factor sequence specificity. *Cell*, **158**, 1431–1443.

Weirauch,M.T. *et al.* (2013) Evaluation of methods for modeling transcription factor

sequence specificity. *Nat. Biotechnol.*, **31**, 126–134.

Zhang,J. *et al.* (2019) Differential regulatory network-based quantification and

prioritization of key genes underlying cancer drug resistance based on time-course

RNA-seq data. *PLOS Comput. Biol.*, **15**, e1007435.

Figures

Figure 1 Pseudo-time analysis for time-course individual bulk RNA-Seq of mouse

embryos in early development

(A) Transcriptomic trajectory of mouse embryos from E7.5 to E13.5. Each point on the

PC1-PC2 plane indicates each individual RNA-Seq result. The solid line indicates an

inferred trajectory by slingshot. (B) A scatterplot of pseudo-time for each sample. Each

point indicates each individual RNA-Seq result and is jittered along the Y-axis.

Figure 2. Inference of downstream genes based on the values in *A*

(A) Selection of a parameter *D* for SCODE. The residual sum of squares (RSS) for each

D (from 1 to 10 by 0.5) was calculated. In this study, we used *D* = 4. (B) Pearson's

correlation coefficients between the values of each *A* from 20 optimizations and the

mean*A,* which is the average of each value from all optimizations. (C) Calculation of

Pearson's correlations between the average expression level of each gene and the absolute

values in each row and column of *A*. (D) The violin plots of Pearson's correlations (E, F)

Scatter plot of the average expression level of each gene and the absolute values in the

column (E) and row (F) of *A* for *Sox8*. (G) An example of the definition of thresholds for

significant regulatory relationship between the regulator and downstream genes. A scatter

plot of the values of *A* for a regulator, *Sox8,* in the decreasing order. The solid line is a

513   regression line for the scatter plot. The dashed lines indicate the defined thresholds.

Figure 3. Validation of the inferred regulatory network of transcription factors with the TF2DNA database

(A) Scatter plot of area under the curve (AUC) target selection for each transcription factor (TF) based on the absolute values of A. (B) Scatter plots of the validated inferred target gene rate of *E2f3* and *Arid5b* in the TF2DNA database. Downstream genes of TFs were selected based on the absolute values of A in decreasing order. Solid lines indicate genes at the thresholds. (C) Bar graph of validated and non-validated downstream genes

of each TF in the TF2DNA database, in decreasing order of the total number of inferred

downstream genes. Only the top 30 TFs are shown. (D) Scatter plot of the validated target

gene rate of the inferred downstream genes of TFj and the background rate of target genes

in the TF2DNA database. Histograms show the distribution of validated target gene rates

of inferred downstream genes and the background rates of downstream genes in the

TF2DNA database.

Figure 4. Overlap of inferred genes downstream of ligand-related and receptor-related genes

(A) Bar plot of the number of inferred downstream genes that are common and unique for each ligand-receptor pair. (B-E) Upset plots of inferred downstream genes that are positively and negatively regulated by the representative ligand-related and receptor-related genes. (B) Retinoic acid signaling pathway. (C) Delta/Notch signaling pathway. (D) Hedgehog signaling pathway.

Supplementary Figure 1. Overlaps of inferred downstream genes for ligand-related

and receptor-related genes

Upset plots of inferred downstream genes that were positively and negatively regulated

by the representative ligand-related and receptor-related genes. (A) TNF signaling

pathway. (B) EGF signaling pathway. (C) TGFβ signaling pathway. (D) BMP signaling

pathway. (E) FGF signaling pathway.

Supplementary Figure 2. Schematic illustration of the retinoic acid and Hedgehog

signaling pathway

(A) In the absence of retinoic acid (RA), RA receptor (RARA) represses the target genes.

ALDH1A (ALDH1A3) synthesizes RA. In the presence of RA, RA receptor activates the

target genes. (B) In the absence of Hedgehog ligand (SSH2), GLI protein (GLI3) is

ubiquitinated to function as a repressor of the target genes. In the presence of Hedgehog

ligand, PATCHED (PTCHD4) inhibits GLI ubiquitination to activate the target genes.

Supplementary Figure 3. Overlaps of inferred downstream genes of *Wnt7a, Wnt8a,*

Fzd5,* and *Fzd6

Upset plots of inferred downstream genes that were positively and negatively regulated

by the representative *Wnt7a* and *Fzd5* (A), *Wnt8a* and *Fzd5* (B), and *Wnt7a* and *Fzd6* (C).

Supplementary Figure 4. Gene expression dynamics of *Fzd* genes

Violin plots for the relative gene expression of *Fzd* genes from E7.5 to E13.5.

565 **Supplementary Figure 5. Gene expression dynamics of *Wnt* genes**

566 Violin plots for the relative expression of *Wnt* genes from E7.5 to E13.5.