Discovery of an Antarctic ascidian-associated uncultivated *Verrucomicrobia* with antimelanoma

palmerolide biosynthetic potential.

Alison E. Murray[a*], Chien-Chi Lo[b], Hajnalka E. Daligault[b], Nicole E. Avalon[c], Robert W. Read[a],

Karen W. Davenport[b], Mary L. Higham[a], Yuliya Kunde[b], Armand E.K. Dichosa[b], Bill J. Baker[c*],

Patrick S.G. Chain[b*]

[a] Division of Earth and Ecosystem Science, Desert Research Institute, Reno, Nevada, USA.

[b] Bioscience Division, Los Alamos National Laboratory, Los Alamos, New Mexico, USA.

[c] Department of Chemistry, University of South Florida, Tampa, Florida, USA.

*Corresponding authors:

Alison E. Murray, Division of Earth and Ecosystem Sciences, Desert Research Institute, Reno,

NV 89512, USA. +1 775-673-7361. alison.murray@dri.edu.

Bill J. Baker, Department of Chemistry, University of South Florida, Tampa, FL 33620, USA.

+1-813-974-1967, bjbaker@usf.edu

Patrick S.G. Chain, Bioscience Division, Los Alamos National Laboratory, Los Alamos, NM

87545, USA. +1-505-665-4019. pchain@lanl.gov

Abstract

The Antarctic marine ecosystem harbors a wealth of biological and chemical innovation that has risen in concert over millennia since the isolation of the continent and formation of the Antarctic circumpolar current. Scientific inquiry into the novelty of marine natural products produced by Antarctic benthic invertebrates led to the discovery of a bioactive macrolide, palmerolide A, that has specific activity against melanoma and holds considerable promise as an anticancer therapeutic. While this compound was isolated from the Antarctic ascidian *Synoicum adareanum*, its biosynthesis has since been hypothesized to be microbially mediated, given structural similarities to microbially-produced hybrid non-ribosomal peptide-polyketide macrolides. Here, we describe a metagenome-enabled investigation aimed at identifying the biosynthetic gene cluster (BGC) and palmerolide A-producing organism. A 74 Kbp candidate BGC encoding the multi-modular enzymatic machinery (hybrid Type I-*trans*-AT polyketide synthase-non-ribosomal peptide synthetase and tailoring functional domains) was identified and found to harbor key features predicted as necessary for palmerolide A biosynthesis. Surveys of ascidian microbiome samples targeting the candidate BGC revealed a high correlation between palmerolide-gene targets and a single 16S rRNA gene variant (R=0.83 – 0.99). Through repeated rounds of metagenome sequencing followed by binning contigs into metagenome-assembled genomes, we were able to retrieve a near-complete genome (10 contigs) of the BGC-producing organism, a novel verrucomicrobium within the *Opitutaceae* family that we propose here as *Candidatus* Synoicihabitans palmerolidicus. The refined genome assembly harbors five highly similar BGC copies, along with structural and functional features that shed light on the host-associated nature of this unique bacterium.

Importance

Palmerolide A has potential as a chemotherapeutic agent to target melanoma. We interrogated

the microbiome of the Antarctic ascidian, *Synoicum adareanum,* using a cultivation-independent

high-throughput sequencing and bioinformatic strategy. The metagenome-encoded biosynthetic

machinery predicted to produce palmerolide A was found to be associated with the genome of a

member of the *S. adareanum* core microbiome. Phylogenomic analysis suggests the organism

represents a new deeply-branching genus, *Candidatus* Synoicihabitans palmerolidicus, in the

Opitutaceae family of the Verrucomicrobia phylum. The *Ca*. S. palmerolidicus 4.29 Mb genome

encodes a repertoire of carbohydrate-utilizing and transport pathways enabling its ascidian-

associated lifestyle. The palmerolide-producer's genome also contains five distinct copies of the

large palmerolide biosynthetic gene cluster that may provide structural complexity of

palmerolide variants.

Introduction

Across the world's oceans, marine benthic invertebrates harbor a rich source of natural products

that serve metabolic and ecological roles in situ. These compounds provide a multitude of

medicinal and biotechnological applications to science, health and industry. The organisms

responsible for their biosynthesis are often unknown (1, 2). Increasingly, these metabolites,

especially in the polyketide class (*trans*-AT in particular), are found to be produced by microbial

counterparts associated with the invertebrate host (3-5). Invertebrates including sponges, corals,

and ascidians for example, harbor a wealth of diverse microbes, few of which have been

cultivated (e.g., 6-8). Genomic tools, in particular, are revealing biochemical pathways

potentially critical in the host-microbe associations (9). Microbes that form persistent mutualistic

(symbiotic) associations provide key roles in host ecology, such as provision of metabolic

requirements, production of adaptive features such as photoprotective pigments,

bioluminescence, or antifoulants, and biosynthesis of chemical defense agents.

Antarctic marine ecosystems harbor species-rich macrobenthic communities (10-12), which

have been the subject of natural products investigations over the past 30 years resulting in the

identification of > 600 metabolites (13). Initially, it was not known whether the same selective

pressures (namely predation and competition, e.g., 14) that operate in mid and low latitudes

would drive benthic organisms at the poles to create novel chemistry (15). However, this does

appear to be the case, and novel natural products have been discovered across algae, sponges,

corals, nudibranchs, echinoderms, bryozoans, ascidians, and increasingly amongst

microorganisms (16) for which the ecological roles have been deduced in a number of cases (13,

17). Studies of Antarctic benthic invertebrate-microbe associations however, pale in comparison

to studies at lower latitudes, yet the few studies that have been reported suggest these

associations (i) harbor an untapped reservoir of biological diversity (18-21) including fungi (22),

(ii) are host species-specific (23, 24), (iii) provide the host with sources of nitrogen and fixed

carbon (25) and (iv) have biosynthetic functional potential (26, 27).

This study was specifically motivated by our desire to understand the biosynthetic origins of

a natural product, palmerolide A, given its potent anticancer activity (28), that is found to be

associated with the polyclinid Antarctic ascidian, *Synoicum adareanum* (Fig. 1A and B).

Ascidians are known to be rich sources of bioactive natural products (9). They have been found

to harbor polyketide, terpenoid, peptide, alkaloid and a few other classes of natural products, of

which the majority have cytotoxic and/or antimicrobial activities. In addition to palmerolide A, a

few other natural products derived from Antarctic ascidians have been reported (29-31).

Ascidian-associated microbes responsible for natural product biosynthesis have been shown to

be affiliated with bacterial phyla including Actinobacteria (which dominates the recognized

diversity), Cyanobacteria, Firmicutes, Proteobacteria (both Alphaproteobacteria and

Gammaproteobacteria) and Verrucomicrobia in addition to many fungi (32, 33). Metagenome-

enabled studies have been key in linking natural products to the organisms producing them in a

number of cases, e.g. patellamide A and C to Cyanobacteria-affiliated *Prochloron* spp. (4), the

tetrahydroisoquinoline alkaloid ET-743 to Gammaproteobacteria-affiliated *Candidatus*

Endoecteinascidia frumentensis (34), patellazoles to Alphaproteobacteria-affiliated *Candidatus*

Endolissoclinum faulkneri (35), and mandelalides to Verrucomicrobia-affiliated *Candidatus*

Didemnitutus mandela (36). However, this is most certainly an under-representation of the

diversity of ascidian-associated microorganisms with capabilities for synthesizing bioactive

compounds, given the breadth of ascidian biodiversity (37). These linkages have been yet to be

investigated for Antarctic ascidians.

Palmerolide A has anticancer properties with selective activity against melanoma when

tested in the National Cancer Institute 60 cell-line panel (28). This result is of particular interest,

as there are few natural product therapeutics for this devastating form of cancer. Palmerolide A

inhibits vacuolar ATPases, which are highly expressed in metastatic melanoma. Given the

current level of understanding that macrolides often have microbial biosynthetic origins, that the

holobiont metagenome has biosynthetic potential (26) and a diverse, yet persistent core

microbiome is found in palmerolide A-containing *S. adareanum* (27) we have hypothesized that

a microbe associated with *S. adareanum* is responsible for the biosynthesis of palmerolide A.

The core microbiome of the palmerolide A-producing ascidian *S. adareanum* in samples

collected across the Anvers Island archipelago (n=63 samples; 27) is comprised of five bacterial

phyla including Proteobacteria (dominating the microbiome), Bacteroidetes, Nitrospirae,

Actinobacteria and Verrucomicrobia. A few candidate taxa in particular, were suggested to be

likely palmerolide A producers based on relative abundance and biosynthetic potential

determined by analysis of lineage-targeted biosynthetic capability (genera: *Microbulbifer*,

Pseudovibrio, *Hoeflea*, and the family *Opitutaceae* (27). This motivated interrogation of the *S.*

adareanum microbiome metagenome, with the goals of determining the metagenome-encoded

biosynthetic potential, identifying candidate palmerolide A biosynthetic gene cluster (BGC)(s)

and establishing the identity of the palmerolide A producing organism.

Results and Discussion

Identification of a candidate palmerolide A biosynthetic gene cluster. Microbial-enriched

fractions of *S. adareanum* metagenomic DNA sequence from 454 and Ion Proton next generation

sequencing (NGS) libraries (almost 18 billion bases in all) were assembled independently, then

merged, resulting in ~ 145 Mbp of assembled bases distributed over 86,387 contigs (referred to

as CoAssembly 1; supplemental material, Table S1). As the metagenome sequencing effort was

focused on identifying potential BGCs encoding the machinery to synthesize palmerolide A, the

initial steps of analysis specifically targeted those contigs in the assembly that were > 40 Kbp

(102 contigs representing 0.12 % of all contigs, which represent 7.07 % of all reads mapped), as

the size of the macrolide ring with 24 carbons would require a large number of polyketide

modules to be encoded. This large fragment subset of CoAssembly 1 was submitted to

antiSMASH v.3 (38) and more recently to v.5 (39). The results indicated a heterogeneous suite

of BGCs, including a bacteriocin, two non-ribosomal peptide synthetases (NRPS), two hybrid

NRPS -Type I PKS, two terpenes, and three *trans*-AT-PKS hybrid NRPS clusters (supplemental

material, Table S2).

We predicted several functional characteristics of the BGC that would be required for

palmerolide A biosynthesis which aided our analysis (see (40) for detailed retrobiosynthetic

analysis). This included evidence of a hybrid nonribosomal peptide-polyketide pathway and

enzymatic domains leading to placement of two distinct structural features of the polyketide

backbone, a carbamoyl transferase that appends a carbamate group at C-11 of the macrolide ring,

and an HMGCoA synthetase that inserts a methyl group on an acetate C-1 position of the

macrolide structure (C-25). The antiSMASH results indicated that two of the three predicted

hybrid NRPS *trans*-AT-Type I PKS contained the predicted markers. Manual alignment of these

two contigs suggested near-identical overlapping sequence (36,638 bases) and, when joined, the

merged contig resulted in a 74,672 Kbp BGC (Fig. 1C). The cluster size was in the range of

other large *trans*-AT PKS encoding BGCs including pederin (54 Kbp; 41), leinamycin (135.6

Kbp; 42) as well as a *cis*-acting AT-PKS, jamaicamide (64.9 Kbp; 43). The combined contigs

encompassed what appeared to be a complete BGC that was flanked at the start with a

transposase and otherwise unlinked in the assembly to other contiguous DNA. The cluster lacked

phylogenetically informative marker genes from which putative taxonomic assignment could be

attributed.

The antiSMASH results suggested that the BGC appears to be novel with the highest degree

of relatedness to pyxipyrrolone A and B (encoded in the *Pyxidicoccus* sp. MCy9557 genome

(44), to which only 14% of the genes have a significant BLAST hit to genes in the metagenome-

encoded cluster. The ketosynthase (KS) sequences (13 in all) fell into three different sequence

groups (40). One was nearly identical (99% amino acid identity) to a previously reported

sequence from a targeted KS study of *S. adareanum* microbiome metagenomic DNA (26). The

other two were most homologous to KS sequences from *Allochromatium humboldtianum,* and

Dickeya dianthicola in addition to a number of hypothetical proteins from environmental

sequence data sets. The primary polyketide synthase (PKS) component of the BGC (labeled in

Fig. 4C, Contig 9) includes 11 of the 13 KS domains which is consistent with a polyketide

backbone which has 22 contiguous carbons since each elongation step adds a two-carbon

unit. This further supports the hypothesis that this candidate BGC is likely responsible for

palmerolide A biosynthesis. A detailed bioinformatic analysis into the step-wise biosynthetic

mechanism was conducted by Avalon et al. (40).

Taxonomic inference of palmerolide A BGC. Taxonomic attribution of the BGC was inferred

using a real time PCR strategy targeting three coding regions of the putative palmerolide A BGC

spanning the length of the cluster (acyltransferase, AT1; hydroxymethylglutaryl Co-A synthase,

HCS, and the condensation domain of the non-ribosomal peptide synthase NRPS, Fig. 1C) to

assay a *Synoicum* microbiome collection of 63 samples that have been taxonomically classified

using Illumina SSU rRNA gene tag sequencing (27). The three gene targets were present in all

samples ranging within and between sites at levels from $\sim 7 \times 10^1 - 8 \times 10^5$ copies per gram of

host tissue (Fig. 2A). The three BGC gene targets co-varied across all samples ($r^2 > 0.7$ for all

pairs), with the NRPS gene copy levels slightly lower overall (mean: 0.66 and 0.59 copies per ng

host tissue for NRPS:AT1 and NRPS:HCS respectively, n=63). We investigated the relationship

between BGC gene copies per ng host tissue for each sample and palmerolide A levels

determined for the same samples using mass spectrometry, however no correlation was found

(R<0.03, n=63; 27). We then assessed the semi-quantitative relationship between the occurrence

of SSU rRNA amplicon sequence variants (ASV, n=461; 27) and the abundance of the three

palmerolide A BGC gene targets. Here, we found a robust correlation (R=0.83 – 0.99) between

all 3 gene targets and a single amplicon sequence variant (ASV15) in the core microbiome (45).

This ASV is affiliated with the *Opitutaceae* family of the Verrucomicrobium phylum. The

Opitutaceae family ASV (SaM_ASV15) was a member of the core microbiome as it was

detected in 59 of the 63 samples surveyed at varying levels of relative abundance and displayed

strong correlations with the abundances of BGC gene targets (Fig. 2B, r^2 =0.68 with AT1, 0.97

with HCS, and 0.69 with NRPS, n=63 for all). The only other correlations R > 0.5 were ASVs

associated within the "variable" fraction of the microbiome, e.g., one low abundance ASV was

present in 24 of 63 samples (45).

This result supports the finding of Murray et al. (27) in which gene abundance and natural

product chemistry do not reflect a 1:1 ratio in this host-associated system. Neither the semi-

quantitative measure of ASV copies nor the real time PCR abundance estimates of the three

biosynthetic gene targets correlated with the mass-normalized levels of palmerolide A present in

the same samples. As discussed (27), this is likely a result of bioaccumulation in the ascidian

tissues. This result provided strong support that the genetic capacity for palmerolide A

production was associated with a novel member of the *Opitutaceae*, a taxonomic family with

representatives found across diverse host-associated and free-living ecosystems. Although the

biosynthetic capacity of this family is not well known (46), recent evidence (36) suggests this

family may be a fruitful target for cultivation efforts and natural product surveys.

Assembly of the palmerolide BGC-associated *Opitutaceae*-related metagenome assembled

genome (MAG). With metagenomes some genomes come together easily – while others present

compelling puzzles to solve. Assembly of the *pal* BGC-containing *Opitutaceae* genome was the

result of a dedicated effort of binning contigs, gene searches, additional sequencing of samples

with high BGC titer, and manual, targeted assembly. Binning efforts with CoAssembly 1 did not

result in association of the *pal* BGC with an associated metagenome assembled genome

(supplemental materials and methods, Table S3). Therefore, a further round of metagenome

sequencing using long read technology (Pacific Biosciences Sequel Systems technology; PacBio)

ensued.

The 16S rRNA gene ASV occurrence (27) and real time PCR data were used to guide *S.*

adareanum sample selection for sequencing. Two ascidian samples (Bon-1C-2011 and Del-2B-

2011) with high *Opitutaceae* ASV occurrences (ASV_015; > 1000 sequences each – relative

abundance of ~ 13.3-15.3 % compared to an overall average of $1.3 \pm 2.77\%$ across the 63

samples respectively, (27) and high BGC gene target levels (i.e., Bon-1C-2011 (copies per ng $\pm$

s.d., n=3): 6.93 ± 1.48 x 10^5 NRPS; 7.52 ± 1.29 x 10^5 AT1; 5.32 ± 1.22 x 10^5 HCS and Del 2B-

2011 (copies per ng $\pm$ s.d., n=3): 2.04 ± 0.10 x 10^5 NRPS; 2.57 ± 0.41 x 10^5 AT1; 5.84 ± 0.16 x

10^5 HCS) were selected for PacBio sequencing. This effort generated 28 GB of data that was

used to create a new hybrid CoAssembly 2 which combined all three sequencing technologies.

Similar to the assembly with the *Mycale hentscheli*-associated polyketide producers (47), the

long-read data set improved the assembly metrics, and subsequent binning resulted in a highly

resolved *Opitutaceae*-classified bin (45, supplemental material, Table S3). Interestingly

however, the palmerolide BGC contigs still did not cluster with this bin, which we later

attributed to binning reliance on sequence depth.

We used PacBio circular consensus sequence (CCS) reads to generate and manually edit the

assembly for our *Opitutaceae* genome of interest. The resulting 4.3 Mbp genome (Fig. 3A) had a

GC content of 58.7% and was resolved into a total of 10 contigs. Five of the contigs (Contigs 1-

5) were unique while contigs 6-10 represented highly similar repeated units of the *pal* BGC

(labeled *pal* BGC 1, 2, 3, 4, and 5) beginning and ending in linkage gaps. Support for the

assembly of a near-complete genome includes manual analysis of the ends of the five unique

contigs and underlying read data linking each end of all five contigs to parts of the repeated

contigs. The structure of the five repeated contigs was manually evaluated (Fig.1) and while each

copy had unique features, all were found to harbor portions of the palmerolide BGC. While the

exact placement of each of the five palmerolide contigs could not be positioned with respect to

the five unique contigs, the data support a single circular chromosome. Analyses further

supporting a complete genome includes the presence of a complete ribosomal operon and

estimated CheckM completeness of 96.04 % based on marker genes (presence of which were

identified using MetaERG annotation of the *Opitutaceae* genome, Table S3), and 45 tRNA

genes.

Alignment of all five repeated contigs to the longest palmerolide-containing BGC revealed a

long (36,198 base) repeated region that was shared between all 5 contigs with some substantial

differences at the beginning of the cluster and only minor differences at the end, indicating 3 full

length, and 2 shorter palmerolide BGC-containing contigs (Fig. 1 and 3). This was consistent

with coverage estimates based on read-mapping that suggested lower depth at the beginning of

the cluster (Fig. 3B). BGC 1 and 3 are nearly identical (over 86,135 bases) with only 2 single

nucleotide polymorphisms (SNPs) and an additional 1,468 bases in BGC1 (237 bases at the 5'

end and 1231 bases at the 3' end). BGC 4 is 13,470 bases shorter than BGC1 at the 5' end, and 5

bases longer than BGC 1 at the 3' end. Alignment of the real time PCR gene targets to the 5 *pal*

BGCs provided independent support for the different lengths of the 5 BGCs, as the region

targeted by the NRP primers was missing in two of the *pal* BGCs, thus explaining lower NRP:

AT or NRP:HCS gene dosages reported above.

Interestingly, precedent for naturally occurring multi-copy BGCs to our knowledge, has only

recently been found in one other bacterium, an ascidian (*Lissoclinum* sp.)-associated

Opitutaceae, *Candidatus* Didemnitutus mandela, linked to cytotoxic mandelalide polyketides

(36). Likewise, we can invoke a rationale similar to (36) that multiple gene clusters may be

linked to biosynthesis of different palmerolide derivatives; see Avalon et al. (40) for retro-

biosynthetic predictions of these clusters and annotation of putative enzymatic functions. Gene

duplication, loss, and rearrangement processes over evolutionary time, likely explain the source

of the multiple copies. At present we do not yet understand the regulatory controls, whether all

five are actively transcribed, in situ function and how this may vary amongst host microbiomes.

Phylogenomic characterization of the Opitutaceae-related MAG. The taxonomic relationship of

the *Opitutacae* MAG to other Verrucomicrobiota was assessed using distance-based analyses

with 16S rRNA and average amino acid identity (AAI). Then, it was classified using the GTDB-

Tk tool (48), and a phylogenomic analysis based on concatenated ribosomal protein markers.

Comparison of 16S rRNA gene sequences amongst other Verrucomicrobia with available

genome sequences (that also have 16S rRNA genes; supplemental material*,* Fig. S1) suggests

that the nearest relatives are *Cephaloticoccus primus* CAG34 (similarity of 0.9138), *Optitutus*

terrae PB90-1 (similarity of 0.9132) and *Geminisphaera coliterminitum* TAV2 (similarity of

0.9108). The *Opitutaceae*-affiliated MAG sequence is identical to a sequence (uncultured

bacterium clone Tun-3b A3) reported from the same host (*S. adareanum*) in a 2008 study (26);

bootstrapping supported a deep branching position in the *Opitutaceae* family. Thus, this

Opitutaceae-related MAG appears to be unique – its 16S rRNA gene sequence was not found in

bacterioplankton amplicon surveys from the Anvers Island archipelago (n=604 amplicon

sequences) and a culture collection reported previously (27), nor were any sequences with

identities higher than 95% found following against a large bacterioplankton amplicon data set

from further north in the Antarctic Peninsula (32,941 sequence clusters derived from 44.6

million sequences; NCBI Bioproject PRJNA316748). Likewise, when searched against the

global nr GenBank database representing environmental data sets the highest sequence identities

found so far are < 96% identity over 88% of the complete sequence.

When characterizing the MAG using AAI metrics (average nucleotide identity, ANI, found

no closely related genomes) the closest genomes were environmental metagenome assemblies

from the South Atlantic TOBG_SAT_155 (53.08 % AAI) and WB6_3A_236 (52.71 % AAI);

and the two closest isolate type genomes were *Nibricoccus aquaticus* str. NZ CP023344 (52.82

% AAI) and *Opitutus terrae* str. PB90 (52.75 % AAI). The Microbial Genome Atlas (MiGA)

support for the MAG belonging in the *Opitutaceae* family was weak (p-values of 0.5). Attempts

to classify this MAG using GTDB-Tk (47) were hampered by the fact we have no real

representative in the genome databases, resulting in low confidence predictions at the species or

genus levels (see the supplemental material for details).

Verrucomicrobia exhibit free-living and host-associated lifestyles in a multitude of terrestrial

and marine habitats on Earth. We performed a meta-analysis of Verrucomicrobia genomes, with

an emphasis on marine and host-associated *Opitutaceae*, to establish more confidence in the

phylogenetic position of the *Opitutaceae* MAG. The analysis was based on 24 conserved

proteins – 21 ribosomal proteins and three additional conserved proteins (InfB, lepA, pheS). The

diversity of the *Opitutaceae* family, and of Verrucomicrobia in general, is largely known from

uncultivated organisms in which there are 20 genera in GTDB (release 05-RS95), 2 additional

genera in the NCBI taxonomy database, and numerous unclassified single amplified genomes

(SAGs); in all, only eight genera have cultivated representatives. Given the uneven

representations of the 24 proteins across all (115) genomes assessed (MAGs and SAGs are often

incomplete), we selected a balance of 16 proteins across 48 genomes to assess phylogenomic

relatedness across the *Opitutaceae* (Fig. 4). Here too, as seen with the 16S rRNA gene

phylogenetic tree, the *S. adareanum-Opitutaceae* MAG held a basal position compared to the

other *Opitutaceae* genomes in the analysis.

***Ca.* S. palmerolidicus relative abundance estimates and ecological inference.** The relative

abundance of *Opitutaceae* bin 8 was estimated in the shotgun metagenomic samples by mapping

the NGS reads back to the assembled MAG across the four *S. adareanum* samples collected.

This indicated varying levels of genome coverage in the natural samples, with the two samples

selected based on real time PCR-quantified high BGC copy number being clearly enriched in this

strain (44.70 % of reads mapped to Bon-1C-2011 and 36.78 % to Del-2B-2011, Table 1). These

levels are higher than estimates of relative abundance derived from the 16S rRNA gene amplicon

surveys (estimated at 13.33 and 15.34 % respectively) for the same samples. This is likely a

result of the single-copy nature of the ribosomal operon in *Opitutaceae* bin 8 vs. other taxa with

multiple rRNA operon copies that could thus be over-represented in the core microbiome library

(e.g., *Pseudovibrio* sp. str. PSC04-5.I4 has 9 and *Microbulbifer* sp. is estimated at 4.1 ± 0.8

based on 9 finished *Microbulbifer* genomes available at the Integrated Microbial Genomes

Database). All host *S. adareanum* lobes surveyed (n=63) in the Anvers Island regional survey

contained high levels (0.49 – 4.06 mg palmerolide A x g^{-1} host dry weight) of palmerolide A

(27), and variable, yet highly concordant levels of the *pal* BGCs and 16S rRNA ASV levels (Fig.

2). Despite the natural population structure sampled here (four single host lobes), the bin-level

sequence variation was low (ranging from 72-243 SNPs) when the PacBio reads were mapped

back to the Opitutaceae bin 8 (Table 1). This suggests maintenance of a relatively invariant

population at the spatial and temporal scales of this coastal Antarctic region while highlighting

our limited understanding of the biogeographical extent of the *S. adareanum*-symbiont-

palmerolide relationship across a larger region of the Southern Ocean.

Several questions remain with regard to the in situ function of palmerolide A (a eukaryotic

V-ATPase inhibitor in human cell line assays (28) in this cryohabitat: how and why is it

bioaccumulated by the host? Overall, the study of natural products in high latitude marine

ecosystems is in its infancy. This palmerolide producing, ascidian-associated, *Opitutaceae*

provides the first Antarctic example in which a well-characterized natural product has been

linked to the genetic information responsible for its biosynthesis. Gaining an understanding of

environmental and biosynthetic regulatory controls, establishing integrated transcriptomic,

proteomic, and secondary metabolome expression in the environment will also reveal whether

the different clusters are expressed in situ. In addition to ecological pursuits, the path to clinical

studies of palmerolide will require genetic or cultivation efforts. At present, we hypothesize that

cultivation of *Opitutaceae* bin 8 may be possible, given the lack of genome reduction or of other

direct evidence for host-associated dependencies (e.g., a number of central carbohydrate and

energy metabolism pathways appear to be present).

***Candidatus* Synoicihabitans palmerolidicus genome attributes**. The Antarctic ascidian,

Synoicum adareanum, harbors a dense community of bacteria that has a conserved core set of

taxa (27). The near complete ~4.30 Mbp *Opitutaceae* bin 8 metagenome assembled genome (Fig.

3) represents one of the core members. This MAG is remarkable in that it encodes for five 36-74

Kbp copies of the candidate BGCs that are implicated in biosynthesis of palmerolide A and

possibly other palmerolide compounds. Intriguingly, this genome does not seem to show

evidence of genome reduction as found in *Candidatus* Didemnitutus mandela (36); the other

ascidian-associated *Opitutaceae* genome currently known to encode multiple BGC gene copies.

This is the first *Opitutaceae* genome characterized from a permanently cold, ~ -1.8 - 2 °C, often

ice-covered ocean ecosystem. This genome encodes one rRNA operon, 45 tRNA genes, and an

estimated 5058 coding sequences. Based on the low (< 92%) SSU rRNA gene identity and low

(< 54% AAI) values to other genera in the *Opitutaceae*, along with the phylogenomic position of

the *Opitutaceae* bin 8, the provisional name "*Candidatus* Synoicihabitans palmerolidicus" (*Ca*.

S. palmerolidicus) is proposed for this novel verrucomicrobium. The genus name

Synoicihabitans (Syn.o.i.ci.ha'bitans. N.L. neut. N. *Synoicum* a genus of ascidians; L. pres. part

habitans inhabiting; N.L. masc. n.) references this organism as an inhabitant of the ascidian

genus *Synoicum*. The species name *palmerolidicus* (pal.me.ro.li'di.cus. N.L. neut.

n. *palmerolidum* palmerolide; N.L. masc. adj.) designates the species as pertaining to

palmerolide.

The GC content of 58.7% is rather high compared to other marine *Opitutaceae* genomes

(ave. 51.49 s.d. 0.02, n=12), yet is ~ average for the family overall (61.58 s.d. 0.06, n=69;

supplemental material, Table S4). MetaERG includes metagenome assembled genomes available

in the GTDB as a resource for its custom GenomeDB that new genomes are annotated against.

This was a clear advantage in annotating the *Ca*. S. palmerolidicus genome as Verrucomicrobia

genomes are widely represented by uncultivated taxa. Likewise, antiSMASH was an invaluable

tool for *pal* BGC identification and domain structure annotation. This formed the basis to derive

a predicted step-wise mechanism of *pal* biosynthesis (40).

_Ca_. S. palmerolidicus genome structure, function and host-associated features. Beyond the

pal BGCs, the *Ca*. S. palmerolidicus genome encodes a variety of additional interesting

structural and functional features that provide insight its lifestyle. Here we provide but a brief

synopsis. In addition to the repeated BGCs, three additional repeated elements with two nearly

identical copies each (15.3 Kbp, 17.0 Kbp, 27.4 Kbp) were identified during the assembly

process (Fig. 3A). These coded for 20, 25 and 41 CDSs respectively, were in some cases flanked

by transposase/integrases (both internal and proximal) and had widespread homology with

Verrucomicrobia orthologs. The contents of the three repetitive elements were not shared

amongst each other.

Annotations were assigned to a little more than half of the CDSs in the 15.3 Mbp repeat

which is predicted to encode for xylose transport, two sulfatases, two endonucleases and a

MacB-like (potential macrolide export) periplasmic core protein. Xylan might be sourced from

seaweeds (49) or even the ascidian, as it is a minor component of the tunic cellulose (50).

Related to this, an endo-1,4-beta-xylanase which has exoenzyme activity in some

microorganisms (51) was identified elsewhere in the genome suggesting the potential for xylose

metabolism. Altogether, eight sulfatase copies were identified in this host-associated genome

(four in the 15.5 Mbp repeat elements). These may be involved in catabolic activities of

sulfonated polysaccharides, and possibly as *trans*-acting elements in palmerolide biosynthesis

(40). In addition to the MacB-like CDSs found in this repeat, 13 other MacB-homologs were

present in the genome – none of which were associated with the *pal* BGCs (supplemental

material, Fig. S2a). MacB is a primary component of the macrolide tripartite efflux pump that

operates as a mechanotransmission system involved both in antibiotic resistance and antibiotic

export depending on the size of the macrolide molecule (52). However, two additional elements

required for this pump to be functional, an intramembrane MacA and an outer membrane protein

TolC, were not co-located elsewhere in the genome. MacA may be missing, as searches against

the of *Ca*. S. palmerolidicus genome with to two other verrucomicrobia-associated MacA CDSs

were not identified using BLAST queries (Peat Soil MAG SbV1 SBV1_730043 and *Ca*.

Udaeobacter copiosis KAF5408997.1; (53). At least nine MacB CDSs were flanked by a FstX-

like permease family protein; the genomic structures of which were quite complex including

several with multiple repeated domains. Detailed transporter modeling is beyond the scope of

this work, but it is likely that these proteins are involved in signaling of cell division machinery

rather than macrolide transport (54).

Predicted CDSs in the 17.0 Mbp repeat included sugar binding and transport domains, as

well as domains encoding rhamnosidase, arabinofuranosidase, and other carbohydrate catabolism

functions. About half the proteins encoded in the 27.4 Mbp repeat were unknown in function,

and of the remaining characterized suggested diverse potential functional capacities. For

example, a zinc carboxypeptidase (1 of 3 in the genome), multidrug and toxic compound

transporter (MatE/NorM), and an exodeoxyribonuclease were identified.

The *Ca.* S. palmerolidicus genome has a number of features that suggest it is adapted to a

host-associated lifestyle, and several of these features were reported recently for two related

sponge-associated Opitutales metagenome bins (*Petrosia ficiformis*-associated bins 0 and 01,

Fig. 4; (55). These include identification of a bacterial microcompartment (BMC) 'super locus'.

Such loci were recently reported to be enriched in host-associated Opitutales genomes when

compared to free-living relatives. The structural proteins for the BMC were present as were other

conserved Planctomyces-Verrucomicrobia BMC genes (56). As in the sponge *Pectoria ficiformis*

metagenome bins, enzymes for carbohydrate (rhamnose) catabolism and modification were

found adjacent to the BMC locus (supplemental material*,* Fig. S3), in addition to the two that

were found in the 27.4 Mbp repeat. The genome did not appear to encode the full complement of

enzymes required for fucose metabolism, though a few alpha-L-fucosidases were identified.

Further evidence for carbohydrate metabolism was supported through genome similarity

searches with the CAZY database (57), including 7 identified carbohydrate binding modules, a

carbohydrate esterase, 14 glycoside hydrolases, 6 glycosyl transferases and a polysaccharide

lyase. In addition, three bacterial cellulases (PF00150, cellulase family A; glycosyl hydrolase

family 5) were identified, all with the canonical conserved glutamic acid residue. These appear to

have different evolutionary histories in which each variant has nearest neighbors in different

bacterial phyla (supplemental material, Fig. S2b) matching between 68% identity for Protein

J6386 03765 to *Lacunisphaera limnophila*, 57.5% identity for Protein J6386 22340 with a

cellulase from a shipworm symbiont *Alteromonadaceae* (*Terridinibacter* sp.), and 37.5%

sequence identity to a Bacteroidetes bacterium. This suggests the potential for cellulose

degradation – which is consistent with ascidians being the only animals known to produce

cellulose for its skeletal structure (50). In addition to the BGCs, the enzymatic resources in this

genome (e.g., xylan and cellulose hydrolysis) are a treasure trove rich with biotechnological

potential. Support for a Type II secretion system (i.e. GspD, GspE, GspG and GspO), common to

gram-negative bacteria which secrete folded proteins (e.g. hydrolytic enzymes required for

survival in host environments) from the periplasm into the extracellular environment (58) were

detected in the *Ca*. S. palmerolidicus genome.

Chemotaxis and flagellar biosynthesis are factors required for horizontally acquired

symbionts, as has been established with the *Vibrio fischeri-Euprymna scolopes* symbiosis (59).

The *Ca*. S. palmerolidicus genome encodes a chemotaxis system (e.g. CheA, CheW, CheR,

CheB, Che C, methyl-accepting chemotaxis domains) with flagellar motors in addition to a

number of other elements of flagellar biosynthesis, which is consistent with horizontal

acquisition by the host. The methyl-accepting chemotaxis system in *V. fischeri* responds to

chitobiose production by *E. scolopes* (59) – it will be interesting to further unravel the details of

Ca. S. palmerolidicus' cellular biology to understand the chemotaxis stimulants, preferential

cellular localization of palmerolide production and resistance mechanisms of the host to the

potent vacuolar ATPase, as well as products made by others in the *S. adareanum* microbiome.

Other indicators of host-association and palmerolide production include T-A domains,

multidrug exporters, and the potential for palmerolide transport and cofactor biosynthesis. T-A

domains were also prevalent in the *Petrosia ficiformis*-associated bins 0 and 01 (55). The *Ca*. S.

palmerolidicus genome encoded at least 22 TA-related genes including multiple MazG and

AbiEii toxin type IV TA systems, AbiEii-Phd_YefM type II toxin-antitoxin systems, along with

genes coding for PIN domains, Zeta toxin, RelB, HipA, MazE and MraZ. In addition to the MatE

(found in the 27.4 Mbp repeat) two other multidrug export systems with homology to MexB and

MdtB were identified. This analysis also resulted in identifying a putative AbiEii toxin

(PF13304) with homology to SyrD, a cyclic peptide ABC type transporter that was present in all

5 BGCs (Fig. 1C; with 52.7% BLAST percent identity to a *Desulfamplus* sp. homolog over the

full length of the protein, and a variety of other bacteria including an *Opitutaceae*-related strain

with similar levels of identity). This transporter is encoded downstream of the large polyketide

gene clusters following the acyl transferase domains and precedes the predicted *trans*-acting

domains at the 3' end of the BGC. Given its genomic position, this protein is a candidate for

palmerolide transport. The *Ca*. S. palmerolidicus genome also encodes the potential for

pantothenate biosynthesis via ilvD, ilvE, panB, ilvC, panD, and panC (60), which is consistent

with palmerolide biosynthesis in which the 4'-phosphopantetheinyl prosthetic group interacts

with acyl carrier proteins for multimodular assembly. Some symbionts, e.g. *Ca*. Entotheonella

sp., are auxotrophic for pantothenate and likely acquire it from other microbiome members (61).

Unlike in *Ca*. D. mandela (36), there does not appear to be ongoing genome reduction, which

may suggest that the *S. adareanum*-*Ca*. S palmerolidicus relationship is more recent, and/or that

the relationship is commensal rather than interdependent. Likewise, we suspect that the

pseudogene content may be high as several CDS appear to be truncated, in which redundant CDS

of varying lengths were found in several cases (including the MacB). There is evidence of lateral

gene transfer acquisitions of cellulase and numerous other enzymes that may confer ecological

advantages through the evolution of this genome. Similarly, the origin of the *pal* BGCs and how

recombination events play out in the success of this Antarctic host-associated system in terms of

adaptive evolution (62), not to mention the ecology of *S. adareanum* is a curiosity. This phylum

promises to be an interesting target for further culture-based and cultivation-free studies –

particularly in the marine environment, in which they have been less well studied compared to

terrestrial free-living and host associated systems.

Together, it appears that the genome of *Ca*. S. palmerolidicus is equipped for life in this host-

associated interactive ecosystem that stands to be one of the first high latitude marine

invertebrate-associated microbiomes with a genome-level understanding – and one that produces

a highly potent natural product, palmerolide A. This system holds promise for future research

now that we have identified the producing organism and *pal* BGC. We still have much to learn

about the ecological role of palmerolide A – if is it involved predation avoidance, antifouling,

antimicrobial defense or some other yet to be recognized aspect of life in the frigid, often ice-

covered and seasonally light-limited waters of the Southern Ocean.

Materials and Methods

Sample Collection. *S. adareanum* lobes were collected in the coastal waters off Anvers Island,

Antarctica and stored at -80 °C until processing (supplemental material, Table S1). See the

supplemental material and (27) for details of sample collection, microbial cell preparation and

DNA extraction.

Metagenome sequencing. Three rounds of metagenome sequencing were conducted, the details

of which are in the supplemental material. This included an initial 454 pyrosequencing effort

with a bacterial-enriched metagenomic DNA preparation from *S. adareanum* lobe (Nor2c-2007).

Next, an Ion Proton System was used to sequence a metagenomic DNA sample prepared from *S.*

adareanum lobe Nor2a-2007. Then two additional *S. adareanum* metagenome DNA samples

(Bon-1C-2011 and Del-2b-2011) selected based on high copy numbers of the palmerolide A

BGC (see real time PCR Methods, supplemental material) and sequenced using Pacific

Biosciences Sequel Systems technology.

Metagenome assembly, annotation and binning. Raw 454 metagenomic reads (1,570,137

single end reads, 904,455,285 bases) were assembled by Newbler (63) v2.9 (Life Technologies,

Carlsbad, CA, flags: -large -rip -mi 98 -ml 80), while Ion Proton metagenomic reads (89,330,870

reads, 17,053,251,055 bases) were assembled using SPAdes (64) v3.5 (flags: --iontorrent). Both

assembled datasets were merged with MeGAMerge (65) v1.2 and produced 86,387 contigs with

a maximum contig size 153,680 and total contig size 144,953,904 bases (CoAssembly 1). To

achieve more complete metagenome coverage and facilitate metagenome assembled genome

assembly, a Circular Consensus Sequence (CCS) protocol (PacBio) was used to obtain high

quality long reads on two samples Bon-1C-2011 and Del-2b-2011. The 5,514,426 PacBio reads

were assembled with aforementioned assembled contigs (CoAssembly 1) on EDGE

Bioinformatics using wtdbg2 (66), a fast and accurate long-read assembler. The contigs were

polished with three rounds of polishing by Racon (67) into a second Coassembly (CoAssembly

2) which has 4,215 contigs with a maximum contig size 2,235,039 and total size 97,970,181

bases. Lastly, a manual approach was implemented to arrive at assembly of the MAG of interest

the details of which are described in the supplemental material.

The contigs from both co-assemblies 1 and 2 were submitted initially to the EDGE

bioinformatics platform (68) for sequence annotation using Prokka (69) v1.13 and taxonomy

classification using BWA (70) mapping to NCBI RefSeq (Version: NCBI 2017 Oct 3).

Bioinformatic predictions of natural product potential was performed using the antibiotics and

secondary metabolite analysis shell (antiSMASH, bacterial versions 3.0, 4.0 and 5.0 (38, 39, 71).

This tool executed contig identification, annotation and analysis of secondary metabolite

biosynthesis gene clusters on both CoAssemblies 1 and 2 (> 1 Kbp and > 40 Kbp data sets). As

most of our attention was focused on analysis the *Ca*. S. palmerolidicus assembled metagenome,

we also used MetaERG (72) as the primary pipeline for metagenome annotation of the ten final

contigs in addition to NCBI's PGAP pipeline (see (45) for GFF dataset). There were 5186 coding

sequences predicted in the MetaERG annotation and 5186 in NCBI's PGAP annotation.

MaxBin (73) and MaxBin2 (74) were used to form metagenome bins for both CoAssembly 1

and 2. CheckM v1.1.11 (75) and v.1.1.12, and GTDB-Tk v.1.0.2 (48) were used to verify bin

quality and taxonomic classification. See supplemental material for details. In order to assess the

representation of assembled *Opitutaceae* genome across the 4 environmental samples used for

metagenome sequencing (resulting from MaxBin2 binning of CoAssembly 2), we used BWA to

map the CCS reads to each metagenome data set.

Real time PCR. Gene targets (non-ribosomal peptide synthase, acyltransferase, and 3-

hydroxymethylglutaryl coenzyme A synthase) were selected at different positions along the

length of the candidate BGC. Supplemental material, Table S5 lists the primer and the GBlocks

synthetic positive control sequence. Metagenomic DNA extracts from a large *S. adareanum*

sample set (n=63 *S. adareanum* lobes from 21 colonies), all containing high levels of

palmerolide A (27), were screened with the real time PCR assays on a Quant Studio 3 (Thermo

Fisher Scientific, Inc.; see supplemental material text for details of controls and analysis).

Phylogenomic analyses. A phylogenomic analysis of the assembled *Opitutaceae* MAG was

conducted based on shared ribosomal RNA and ribosomal proteins amongst 46 and 48 reference

genomes respectively, out of 115 genomes in total, mined from various databases (NCBI, GTDB

and IMG) for uncultivated and cultivated microorganisms identified in the Verrucomicrobia

phylum (supplemental material, Table S6). The details of these analyses are described in the

supplemental material. In addition, we used MiGA (NCBI Prokaryotic taxonomy and the

environmental TARA Oceans (Tully) databases; accessed August 2020) and GTDB-Tk (ver.

1.3.0) tools for MAG taxonomic classification.

Phylogenetic analysis of the MacB CDS sequences were retrieved from MetatERG annotated

Ca. S. palmerolidicus contigs and homologs were retrieved from the NCBI based on BLAST

results. Maximum likelihood analysis was conducted on 994 aligned (MUSCLE) positions using

RAxML v.8.2.12 using the PROTGAMMALG model and 550 bootstrap replicates. For the

phylogenetic analysis of the cellulase CDS, homologs were retrieved from the NCBI based on

BLAST results resulting in 19 sequences and 496 aligned positions (ClustalOmega) was also

conducted using RAxML v.8.2.12 under the PROTGAMMALG model of evolution with 1000

bootstraps.

Data Availability

The data (biosamples, SRA depositions) associated with this study is associated with NCBI

BioProject PRJNA662631. The *Ca.* S. palmerolidicus biosample ID is SAMN18473105 and

genome accession JAGGDC000000000. An annotation data set (.gff) for the Ca. S.

palmerolidicus MAG resulting from the metaERG pipeline is available (45).

Acknowledgments

The assistance of several collaborators and students are acknowledged including C. Amsler,

M. Amsler, L. Bishop, J. Cuce, B. Dent, N. Ernster, C. Gleasner, A. Maschek, A. Shilling, L.

Siao, S. Thomas, and the Palmer Station science support staff. We especially would like to

acknowledge the help of A. Oren in assisting with bacterial nomenclature proposed here.

Likewise, we thank J.T. Hollibaugh and A.L. Reysenbach for comments on previous drafts of the

manuscript.

References

1.   Simmons TL, Coates RC, Clark BR, Engene N, Gonzalez D, Esquenazi E, Dorrestein PC,
581   Gerwick WH. 2008. Biosynthetic origin of natural products isolated from marine
582   microorganism-invertebrate assemblages. Proc Natl Acad Sci USA 105:4587-4594.
583   2.   Morita M, Schmidt EW. 2018. Parallel lives of symbionts and hosts: chemical mutualism
584   in marine animals. Nat Prod Rep 35:357-378.
585   3.   Piel J, Hui DQ, Fusetani N, Matsunaga S. 2004. Targeting modular polyketide synthases
586   with iteratively acting acyltransferases from metagenomes of uncultured bacterial
587   consortia. Environ Microbiol 6:921-927.
588   4.   Schmidt EW, Nelson JT, Rasko DA, Sudek S, Eisen JA, Haygood MG, Ravel J. 2005.
589   Patellamide A and C biosynthesis by a microcin-like pathway in *Prochloron didemni*, the
590   cyanobacterial symbiont of *Lissoclinum patella*. Proc Natl Acad Sci USA 102:7315-
591   7320.
592   5.   Piel J. 2009. Metabolites from symbiotic bacteria. Natural Product Reports 26:338-362.
593   6.   Sunagawa S, Woodley CM, Medina M. 2010. Threatened corals provide underexplored
594   microbial habitats. Plos One 5:e9554.
595   7.   Behrendt L, Larkum AWD, Trampe E, Norman A, Sorensen SJ, Kuhl M. 2012. Microbial
596   diversity of biofilm communities in microniches associated with the didemnid ascidian
597   Lissoclinum patella. ISME J 6:1222-1237.
598   8.   Webster NS, Taylor MW. 2012. Marine sponges and their microbial symbionts: love and
599   other relationships. Environ Microbiol 14:335-346.
600   9.   McFall-Ngai M, Hadfield MG, Bosch TCG, Carey HV, Domazet-Loso T, Douglas AE,
601   Dubilier N, Eberl G, Fukami T, Gilbert SF, Hentschel U, King N, Kjelleberg S, Knoll
602   AH, Kremer N, Mazmanian SK, Metcalf JL, Nealson K, Pierce NE, Rawls JF, Reid A,
603   Ruby EG, Rumpho M, Sanders JG, Tautz D, Wernegreen JJ. 2013. Animals in a bacterial
604   world, a new imperative for the life sciences. Proc Natl Acad Sci USA 110:3229-3236.
605   10.   Gray J. 2001. Antarcic marine benthic biodiversity in a world-wide latitudinal context.
606   Polar Biol 24:633-641.
607   11.   Clarke A. 2008. Antarctic marine benthic diversity: patterns and processes. J Exp Mar
608   Biol Ecol 366:48-55.

609   12.   Piepenburg D, Archambault P, Ambrose WG, Jr., Blanchard AL, Bluhm BA, Carroll ML,
610   Conlan KE, Cusson M, Feder HM, Grebmeier JM, Jewett SC, Levesque M, Petryashev
611   VV, Sejr MK, Sirenko BI, Wlodarska-Kowalczuk M. 2011. Towards a pan-Arctic
612   inventory of the species diversity of the macro- and megabenthic fauna of the Arctic shelf
613   seas. Mar Biodivers 41:51-70.
614   13.   von Salm J, Schoenrock K, McClintock J, Amsler C, Baker B. 2018. The status of marine
615   chemical ecology in Antarctica: form and function of unique high-latitude chemistry, p
616   69, Chemical Ecology: The ecological impacts of marine natural products, 1st ed
617   doi:10.1201/9780429453465. CRC Press, Boca Raton.
618   14.   McClintock JB, Amsler CD, Baker BJ. 2001. Introduction to the symposium: Antarctic
619   marine biology. Am Zool 41:1-2.
620   15.   Bakus GJ, Green G. 1974. Toxicity in sponges and holothurians - geographic pattern.
621   Science 185:951-953.
622   16.   Soldatou S, Baker BJ. 2017. Cold-water marine natural products, 2006 to 2016. Nat Prod
623   Rep 34:585-626.
624   17.   Avila C, Taboada S, Nunez-Pons L. 2008. Antarctic marine chemical ecology: what is
625   next? Mar Ecol-Evol Persp 29:1-71.
626   18.   Webster NS, Negri AP, Munro M, Battershill CN. 2004. Diverse microbial communities
627   inhabit Antarctic sponges. Environ Microbiol 6:288-300.
628   19.   Webster NS, Bourne D. 2007. Bacterial community structure associated with the
629   Antarctic soft coral, *Alcyonium antarcticum*. FEMS Microbiol Ecol 59:81-94.
630   20.   Murray AE, Rack FR, Zook R, Williams MJM, Higham ML, Broe M, Kaufmann RS,
631   Daly M. 2016. Microbiome composition and diversity of the ice-dwelling sea anemone,
632   *Edwardsiella andrillae*. Integr Comp Biol 56:542-555.
633   21.   Rodriguez-Marconi S, De la Iglesia R, Diez B, Fonseca CA, Hajdu E, Trefault N. 2015.
634   Characterization of bacterial, archaeal and eukaryote symbionts from Antarctic sponges
635   reveals a high diversity at a three-domain level and a particular signature for this
636   ecosystem. Plos One 10:e0138837.
637   22.   Godinho VM, de Paula MTR, Silva DAS, Paresque K, Martins AP, Colepicolo P, Rosa
638   CA, Rosa LH. 2019. Diversity and distribution of hidden cultivable fungi associated with
639   marine animals of Antarctica. Fungal Biol 123:507-516.
640   23.   Steinert G, Wemheuer B, Janussen D, Erpenbeck D, Daniel R, Simon M, Brinkhoff T,
641   Schupp PJ. 2019. Prokaryotic diversity and community patterns in Antarctic continental
642   shelf sponges. Front Mar Sci 6:297.
643   24.   Sacristan-Soriano O, Criado NP, Avila C. 2020. Host species determines symbiotic
644   community composition in Antarctic sponges (Porifera: *Demospongiae*). Front Mar Sci
645   7:474.
646   25.   Moreno-Pino M, Cristi A, Gillooly JF, Trefault N. 2020. Characterizing the microbiomes
647   of Antarctic sponges: a functional metagenomic approach. Sci Rep UK 10:645.
648   26.   Riesenfeld CS, Murray AE, Baker BJ. 2008. Characterization of the microbial
649   community and polyketide biosynthetic potential in the palmerolide-producing tunicate,
650   *Synoicum adareanum*. J Nat Prod 71:1812-1818.
651   27.   Murray AE, Avalon NE, Bishop L, Davenport KW, Delage E, Dichosa AEK, Eveillard
652   D, Higham ML, Kokkaliari S, Lo C-C, Riesenfeld CS, Young RM, Chain PSG, Baker BJ.
653   2020. Uncovering the core microbiome and distributions of palmerolide in *Synoicum*
654   *adareanum* across the Anvers Island archipelago, Antarctica. Mar Drugs 18:298.

28. Diyabalanage T, Amsler CD, McClintock JB, Baker BJ. 2006. Palmerolide A, a cytotoxic
macrolide from the Antarctic tunicate *Synoicum adareanum*. J Am Chem Soc 128:5630-
5631.
29. Franco LH, Joffe EBD, Puricelli L, Tatian M, Seldes AM, Palermo JA. 1998. Indole
alkaloids from the tunicate *Aplidium meridianum*. J Nat Prod61:1130-1132.
30. Miyata Y, Diyabalanage T, Amsler CD, McClintock JB, Valeriote FA, Baker BJ. 2007.
Ecdysteroids from the antarctic tunicate *Synoicum adareanum*. J Nat Prod 70:1859-1864.
31. Seldes AM, Brasco MFR, Franco LH, Palermo JA. 2007. Identification of two
meridianins from the crude extract of the tunicate *Aplidium meridianum* by tandem mass
spectrometry. Nat Prod Rep21:555–563.
32. Chen L, Hu JS, Xu JL, Shao CL, Wang GY. 2018. Biological and chemical diversity of
ascidian-associated microorganisms. Mar Drugs 16:362.
33. Dou X, Dong B. 2019. Origins and Bioactivities of Natural Compounds Derived from
Marine Ascidians and Their Symbionts. Mar Drugs 17:670.
34. Rath CM, Janto B, Earl J, Ahmed A, Hu FZ, Hiller L, Dahlgren M, Kreft R, Yu FA,
Wolff JJ, Kweon HK, Christiansen MA, Hakansson K, Williams RM, Ehrlich GD,
Sherman DH. 2011. Meta-omic characterization of the marine invertebrate microbial
consortium that produces the chemotherapeutic natural product ET-743. ACS Chem Biol
6:1244-1256.
35. Kwan JC, Donia MS, Han AW, Hirose E, Haygood MG, Schmidt EW. 2012. Genome
streamlining and chemical defense in a coral reef symbiosis. Proc Natl Acad Sci USA
109:20655-20660.
36. Lopera J, Miller IJ, McPhail KL, Kwan JC. 2017. Increased biosynthetic gene dosage in a
genome-reduced defensive bacterial symbiont. mSystems 2:e00096-17.
37. Shenkar N, Swalla BJ. 2011. Global diversity of Ascidiacea. PLoS One 6:e20657.
38. Weber T, Blin K, Duddela S, Krug D, Kim HU, Bruccoleri R, Lee SY, Fischbach MA,
Muller R, Wohlleben W, Breitling R, Takano E, Medema MH. 2015. antiSMASH 3.0-a
comprehensive resource for the genome mining of biosynthetic gene clusters. Nuc Acids
Res 43:W237-W243.
39. Blin K, Shaw S, Steinke K, Villebro R, Ziemert N, Lee SY, Medema MH, Weber T.
2019. antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline. Nuc
Acids Res 47:W81-W87.
40. Avalon N, Murray A, Daligault H, Lo C, Davenport K, Dichosa A, Chain P, Baker B.
Bioinformatic and mechanistic analysis of the palmerolide PKS-NRPS biosynthetic
pathway from the microbiome of an Antarctic ascidian.
bioRxiv:doi.org/10.1101/2021.04.05.438531.
41. Piel J. 2002. A polyketide synthase-peptide synthetase gene cluster from an uncultured
bacterial symbiont of *Paederus* beetles. Proc Natl Acad Sci USA 99:14002-7.
42. Cheng YQ, Tang GL, Shen B. 2002. Identification and localization of the gene cluster
encoding biosynthesis of the antitumor macrolactam leinamycin in *Streptomyces*
atroolivaceus S-140. J Bacteriol 184:7013-24.
43. Edwards DJ, Marquez BL, Nogle LM, McPhail K, Goeger DE, Roberts MA, Gerwick
WH. 2004. Structure and biosynthesis of the jamaicamides, new mixed polyketide-
peptide neurotoxins from the marine cyanobacterium *Lyngbya majuscula*. Chem Biol
11:817-33.

44. Kjaerulff L, Raju R, Panter F, Scheid U, Garcia R, Herrmann J, Muller R. 2017.
Pyxipyrrolones: structure elucidation and biosynthesis of cytotoxic myxobacterial
metabolites. Angew Chem Int E. Engl 56:9614-9618.
45. Murray, AE, Lo, C-C, Daligault, HE, Avalon, NE, Read, RW, Davenport, KW, Higham,
ML, Kunde, Y, Dichosa, AEK, Baker, BJ, Chain, PSG 2021. Supplementary information
provided with Murray et al. "Discovery of an Antarctic ascidian-associated uncultivated
Verrucomicrobia with antimelanoma palmerolide biosynthetic potential.", Dryad,
Dataset, https://doi.org/10.5061/dryad.8sf7m0cpp.
46. Blin K, Andreu VP, de los Santos ELC, Del Carratore F, Lee SY, Medema MH, Weber
T. 2019. The antiSMASH database version 2: a comprehensive resource on secondary
metabolite biosynthetic gene clusters. Nuc Acids Res 47:D625-D630.
47. Storey MA, Andreassend SK, Bracegirdle J, Brown A, Keyzers RA, Ackerley DF,
Northcote PT, Owen JG. 2020. Metagenomic exploration of the marine sponge *Mycale*
hentscheli uncovers multiple polyketide-producing bacterial symbionts. Mbio 11:e02997-
19.
48. Chaumeil P-A, Mussig A, Hugenholtz P, Parks D. 2020. GTDB-Tk: a toolkit to classify
genomes with the Genome Taxonomy Database. Bioinformatics 36:1925–1927.
49. Qeshmi FI, Homaei A, Fernandes P, Hemmati R, Dijkstra BW, Khajeh K. 2020.
Xylanases from marine microorganisms: A brief overview on scope, sources, features
and potential applications. BBA-Proteins Proteom 1868:140312.
50. Zhao Y, Li J. 2014. Excellent chemical and material cellulose from tunicates: diversity in
cellulose production yield and chemical and morphological structures from differnet
tunicate species. Cellul Chem Technol 21:3427-3441.
51. Juturu V, Wu JC. 2014. Microbial exo-xylanases: a mini review. Appl Biochem
Biotechnol 174:81-92.
52. Greene NP, Kaplan E, Crow A, Koronakis V. 2018. Antibiotic resistance mediated by the
MacB ABC transporter family: a structural and functional perspective. Front Microbiol
9:950.
53. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment
search tool. J Mol Biol 215:403-410.
54. Crow A, Greene NP, Kaplan E, Koronakis V. 2017. Structure and mechanotransmission
mechanism of the MacB ABC transporter superfamily. Proc Natl Acad Sci USA
114:12572-12577.
55. Sizikov S, Burgsdorf I, Handley KM, Lahyani M, Haber M, Steindler L. 2020.
Characterization of sponge-associated Verrucomicrobia: microcompartment-based sugar
utilization and enhanced toxin-antitoxin modules as features of host-associated
Opitutales. Environ Microbiol 22:4669-4688.
56. Erbilgin O, McDonald KL, Kerfeld CA. 2014. Characterization of a planctomycetal
organelle: a novel bacterial microcompartment for the aerobic degradation of plant
saccharides. Appl Environ Microbiol 80:2193-2205.
57. Lombard V, Ramulu HG, Drula E, Coutinho P, Henrissat B. 2014. The carbohydrate-
active enzymes database (CAZy) in 2013. Nucleic Acids Res 42:D490–D495.
58. Costa TRD, Felisberto-Rodrigues C, Meir A, Prevost MS, Redzej A, Trokter M,
Waksman G. 2015. Secretion systems in Gram-negative bacteria: structural and
mechanistic insights. Nat Rev Microbiol 13:343-359.

59.   Visick KL, Stabb EV, Ruby EG. A lasting symbiosis: how Vibrio fischeri finds a squid
partner and persists within its natural host. Nat Rev Microbiol doi:10.1038/s41579-021-
00557-0.
60.   Leonardi R, Jackowski S. 2007. Biosynthesis of pantothenic acid and coenzyme A.
EcoSal Plus 2:10.1128/ecosalplus.3.6.3.4.
61.   Lackner G, Peters EE, Helfrich EJN, Piel J. 2017. Insights into the lifestyle of uncultured
bacterial natural product factories associated with marine sponges. Proc Natl Acad Sci
USA 114:E347-E356.
62.   Chevrette MG, Gutierrez-Garcia K, Selem-Mojica N, Aguilar-Martinez C, Yanez-Olvera
A, Ramos-Aboites HE, Hoskisson PA, Barona-Gomez F. 2020. Evolutionary dynamics
of natural product biosynthesis in bacteria. Nat Prod Rep 37:566-599.
63.   Chaisson MJ, Pevzner PA. 2008. Short read fragment assembly of bacterial genomes.
Genome Res 18:324-330.
64.   Nurk S, Bankevich A, Antipov D, Gurevich AA, Korobeynikov A, Lapidus A, Prjibelski
AD, Pyshkin A, Sirotkin A, Sirotkin Y, Stepanauskas R, Clingenpeel SR, Woyke T,
McLean JS, Lasken R, Tesler G, Alekseyev MA, Pevzner PA. 2013. Assembling single-
cell genomes and mini-metagenomes from chimeric MDA products. J Comput Biol
20:714-737.
65.   Scholz M, Lo CC, Chain PSG. 2014. Improved assemblies using a source-agnostic
pipeline for metagenomic assembly by merging (MeGAMerge) of contigs. Sci Rep
4:6480.
66.   Ruan J, Li L. 2019. Fast and accurate long-read assembly with wtdbg2. Nat Methods
17:155-158.
67.   Vaser R, Sovic I, Nagarajan N, Sikic M. 2017. Fast and accurate de novo genome
assembly from long uncorrected reads. Genome Res 27:737-746.
68.   Li P-E, Lo C-C, Anderson JJ, Davenport KW, Bishop-Lilly KA, Xu Y, Ahmed S, Feng
SH, Mokashi VP, Chain PSG. 2017. Enabling the democratization of the genomics
revolution with a fully integrated web-based bioinformatics platform. Nuc Acids Res
45:67-80.
69.   Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics
30:2068-2069.
70.   Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics 25:1754-1760.
71.   Blin K, Wolf T, Chevrette MG, Lu XW, Schwalen CJ, Kautsar SA, Duran HGS, Santos
E, Kim HU, Nave M, Dickschat JS, Mitchell DA, Shelest E, Breitling R, Takano E, Lee
SY, Weber T, Medema MH. 2017. antiSMASH 4.0-improvements in chemistry
prediction and gene cluster boundary identification. Nuc Acids Res 45:W36-W41.
72.   Dong XL, Strous M. 2019. An integrated pipeline for annotation and visualization of
metagenomic contigs. Front Genetics 10:999.
73.   Wu YW, Tang YH, Tringe SG, Simmons BA, Singer SW. 2014. MaxBin: an automated
binning method to recover individual genomes from metagenomes using an expectation-
maximization algorithm. Microbiome 2:26.
74.   Wu YW, Simmons BA, Singer SW. 2016. MaxBin 2.0: an automated binning algorithm
to recover genomes from multiple metagenomic datasets. Bioinformatics 32:605-607.

75.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the
790   quality of microbial genomes recovered from isolates, single cells, and metagenomes. Genome
791   Res 25:1043-1055.
792

793 **Table 1.** Metagenomic reads from 4 different samples were mapped back to the *Ca.* S.
794 palmerolidicus MAG.
795
796

| | *Synoicum adareanum* samples | | | |
	Bon-1c-2011	Del-2b-2011	Nor-2c-2007	Nor-2a-2007
Technology	PacBio CCS reads (1 cell)	PacBio CCS reads (3 cells)	454	Ion Torrent Proton
Number of reads	48,298	9,576	1,570,126	89,330,870
Mapped reads	21,591	3,522	23,993	15,979,084
Mapped reads (%)	44.70	36.78	1.53	17.89
Base coverage	99.98	99.89	90.15	99.98
Average fold	58.38	8.43	2.79	708.79
Gaps	2	3	1,734	8
Gap bases	644	4,618	422,870	774
SNPs	72	196	168	243
Indels	126	64	17	68

797
798

Figure 1. Palmerolide A, a cytotoxic, macrolide with anti-melanoma activity is found in the
tissues of *Synoicum adareanum* in which a candidate biosynthetic gene cluster has been
identified. (A) *S. adareanum* occurs on rocky coastal seafloor habitats in the Antarctic; this study
focused on the region off-shore of Anvers Island in the Antarctic Peninsula. (B) Palmerolide A,
is the product of a hybrid PKS-NRPS system in which biosynthesis begins with a PKS starter
unit followed by incorporation of a glycine subunit by an NRPS module. Subsequent elongation,
cyclization and termination steps follow. Two additional features of the molecule include a
methyl group on C-25 and a carbamate group on C-11. (C) Five repeats encompassing candidate
palmerolide biosynthetic gene clusters were identified. The BGC (in blue) is defined as starting
with the NRP unit and ending at the carbamoyltransferase (green). Candidate palmerolide A
biosynthetic gene cluster BGC4 was identified from initial metagenome library assemblies. The
other four clusters were identified following a third round of sequencing, assembly and manual
finishing. Primary BGC coding sequences (CDS) and a conserved tailoring cassette are in blue.
Light blue CDS are an ATP transporter with homology to an antibiotic transporter, SyrD. All
black CDS are repeated amongst the BGCs. Orange CDS are transposase/integrase domains.
Gray CDS are unique, non-repeated; and in BGC2 and 5, the unique CDS encode transposases,
distinct from the predicted amino acid sequences of those in orange. The red lines associated
with Contig 9 indicate targeted quantitative PCR regions.

Figure 2. Abundances of real time PCR-targeted coding regions in the candidate pal biosynthetic
gene cluster in Antarctic ascidian samples. (A) Gene copies estimated for three targeted coding
regions (Acyltransferase, AT1; 3-hydroxy-methyl-glutaryl coenzyme A synthase, HCS; and the
condensation domain of a non-ribosomal peptide synthase; NRPS) in the candidate *pal*
biosynthetic gene cluster surveyed over 63 DNA extracts derived from microbial cell
preparations enriched from the Antarctic ascidian *Synoicum adarenum.* Nine samples were
collected at each of seven sites: Bon, Bonaparte Point; Del, Delaca Island; Jan, Janus Island, Kil,
Killer Whale Rocks; Lag, Laggard Island, Lit, Litchfield Island; Nor, Norsel Point (27). (B)
Relationship between gene copy number for the three gene targets and the 16S rRNA gene ASV
occurrences of Opitutaceae-related ASV_15 across a 63 *S. adareanum* microbial DNA sample
set. * indicates samples Bon-1C-2011 and Del-2b-2011 that were selected for PacBio
sequencing.

Figure 3. Genome maps of assembled MAG, *Candidatus* Synoicohabitans palmerolidicus and
evidence of multi-copy biosynthetic gene clusters. (a) The 4,297,084 bp gene map is oriented to
dnaA at the origin. One possible assembly scenario of the *Ca.* S. palmerolidicus genome is
shown as the order of the contigs and palmerolide BGC's are not currently known. In addition to
the 5 BGCs, three other internally repetitive regions were identified (15.3, 17.0 and 27.4 Kbp).
The genes and orientation are shown with blue and tRNA indicated in red. (b) To demonstrate
depth of coverage outside and inside the BGC regions, CCS reads from sample Bon-1C-2011
and Del-2b-2011 were mapped to a 167.6 Kbp region. The profile extends 40 Kbp into the
genome on either side of the BGC where depth of coverage averages 60 fold, while in the BGC
depth of coverage varies across the BGC given differences in cover across the BGC, the highest
cover is 5X, or ~ 300 fold, supporting the finding of 5 repeats encoding the BGC.

Figure. 4. Maximum likelihood phylogenomic tree showing 48 verrucomicrobia genomes.
Phylogenomic relationship of *Candidatus* Synoicohabitans palmerolidicus (*Opitutaceae* bin 8)
with respect to other mostly marine, and host-associated verrrucomicrobia subdivision 4 and
other genomes. The tree is based on 16 concatenated ribosomal proteins (5325 amino acids)
common across 48 Verrucomicrobia genomes. Distance was estimated with RAxML with 300
bootstrap replicates. Symbols designate environmental origins of the organisms: free-living are
represented by circles: light blue - marine, green– freshwater, red – hydrothermal mud, brown –
soils. Host-associated taxa from marine systems with blue diamonds, and from terrestrial systems
with black diamonds.
854
855 **Supplemental Material**
856
Supplemental Materials and Methods
Details concerning ascidian sample collections, sample processing and high molecular weight
DNA extraction, metagenome sequencing, metagenome binning, bin taxonomic and functional
classification, Real Time PCR, manual assembly procedures, annotation and phylogenomic
analyses and associated references are provided in the supplemental materials and methods text
section.
863
864 **Supplemental Figure Legends**
865
Figure S1. Ribosomal RNA-based phylogenetic tree. Sequences were selected from
Verrucomicrobia (Phylum) and mostly Opitutaceae (family) genomes represented by isolates,
metagenome assembled genomes and single cell assembled genomes from host associated and
marine ecosystems. RAXML tree is based on 1,636 bases; 250 bootstraps were run, and values >
49 are shown at nodes. *Ca*. S. palmerolidicus (*Opitutaceae* bin 8) is indicated with an orange
diamond symbol. Symbols designate environmental origins of the organisms: free-living are
represented by circles: light blue - marine, green– freshwater, red – hydrothermal mud, brown –
soils. Host-associated taxa from marine systems with blue diamonds, and from terrestrial systems
with black diamonds.
875
Figure S2. Phylogenetic relationships of homologs of (a) MacB CDS and (b) celluase
glycosylhydrolase family 5 enzymes identified in the Ca. S. palmerolidicus genome and closest
neighboring sequences that were sourced from GenBank using BLAST. Maximum likelihood
analysis was conducted using RAxML v. 8.2.12 using the PROTGAMMALG model with 550
bootstrap replicate trees calculated for (a) and 1000 for (b). Bootstrap values > 49 are shown
which represent percent of times the topology was found.
882
Figure S3. Bacterial microcompartment (BMC) super loci identified in the *Ca*. S. palmerolidicus
MAG. Conserved structural elements identified with Pfam and Swiss-Prot annotations were
consistent with other PV-BMC loci which are associated carbohydrate utilization domains.
Transcriptional regulatory genes are indicated in green, conserved PV-BMC enzymes in blue, the
BMC proteins in red, enzymes associated with rhamnose metabolism in violet and those with
lactate utilization in purple.
889
890 **Supplemental Table Legends**

891
892 **Table S1.** Metagenome sequencing metadata for *S. adareanum* microbiome. Host lobe identities
893 indicate sample site (Nor, Norsel, Bon, Bonaparte Point, Del, DeLaca Island; followed by lobe
894 designation, and year sampled). Assembly 1 joined two data sets using mega-merged contigs
895 from 454 and Ion Proton sequence data sets. Assembly 2 included Assembly 1 and PacBio
896 assemblies in addition to the manually assembled *Ca.* S. palmerolidicus MAG.
897
898 **Table S2**. Assembly 1 antiSMASH results for the 102 contigs > 40 Kbp. The 72.1 and 78.1
899 regions encoded biosynthetic gene clusters presented two putative clusters that were further
900 investigated using QPCR to target samples with high BGC copy numbers for a subsequent round
901 of metagenome sequencing.
902
Table S3. Quality checks and taxonomic classification using CheckM and GTDB-Tk for
Opitutaceae-associated bins at different stages of analysis (most recent, highest quality on the
left-most column and initial Coassembly 1 and bin 4 in the right-most column). The *Ca.* S.
palmerolidicus-2 column reflects identifications (MetaERG annotation) of many of the missing
markers in *Ca.* S palmerolidicus-1. The Opitutaceae bin 8 – clean CoAssembly 2 data set reflects
an effort to remove non-Opitutaceae sequences from the bin (see supplemental material,
methods). CoAssembly 1 bins 1 and 2 included the palmerolide A biosynthetic gene cluster and
were otherwise populated by short contigs, and were taxonomically unidentified. CoAssembly 1
bin 4 included Opitutales-related contigs including a SSU rRNA gene sequence.
912
913 **Table S4**. GC percent for Opitutaceae-family representatives. Different genera are represented
914 with different colors. Marine Optitutaceae metagenome assembled genomes are indicated with *.
915 Source and species nomenclature: GTDB release 05-RS95. The GC content for *Ca.*
916 Synoicihabitans palmerolidicus is 58.7%.
917
918 **Table S5.** Quantitative PCR gene targets and primers. Primers targeting three gene targets were
919 used in this study Primer IDs are based on bases from the beginning of the CDS. A GBlocks
920 synthetic DNA positive control was used for estimating copy number[†].
921
922 **Table S6**. Conserved markers used in phylogenomic analysis. Integrated microbial genomes
923 (IMG) and/or GenBank accession numbers are listed. Shaded markers were those used for
924 phylogenomic analysis (Fig. 4).
925

A

B

C

Contig 6
BGC 1; 73,849 bases

Contig 7
BGC 2; 36,130 bases

Contig 8
BGC 3; 73,849 bases

Contig 9
BGC 4; 71,939 bases

CT   HCS   AT   PKS   NRP

Contig 10
BGC 5; 67,501 bases

10,000   20,000   30,000   40,000   50,000   60,000   70,000   80,000

A

B

A

B