Optimized Quantification of Intrahost Viral Diversity in SARS-CoV-2 and Influenza Virus Sequence Data

Running title: Optimized Quantification of Intrahost Viral Diversity

Roder AE*[1], Johnson KEE*[2], Knoll M*[2], Khalfan M[2], Wang B[2], Schultz-Cherry S[3], Banakis S[1], Kreitman A[1], Mederos C[1], Youn J-H[4], Mercado R[4], Wang W[1], Ruchnewitz D[5], Samanovic MI[6], Mulligan MJ[6], Lassig M[5], Łuksza M[7], Das S[4], Gresham D[2#], Ghedin E[1,2#].

Author Affiliations:

[1]Systems Genomics Section, Laboratory of Parasitic Diseases, DIR, NIAID, NIH, Bethesda, MD 20894, USA

[2]Center for Genomics and Systems Biology, Department of Biology, New York University, New York, NY, 10003, USA.

[3]Department of Infectious Diseases, St Jude Children Research Hospital, Memphis, TN

[4]Department of Laboratory Medicine, NIH, Bethesda, MD 20894, USA

[5]Institute for Biological Physics, University of Cologne, Cologne, Germany

[6]New York University Langone Vaccine Center, Department of Medicine, New York, NY 10016, USA

[7]Department of Oncological Sciences, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

*Contributed equally to this article, ordered by contributions to the manuscript

#Corresponding authors: elodie.ghedin@nih.gov; dgresham@nyu.edu

ABSTRACT

High error rates of viral RNA-dependent RNA polymerases lead to diverse intra-host viral

populations during infection. Errors made during replication that are not strongly deleterious to

the virus can lead to the generation of minority variants. However, accurate detection of minority

variants in viral sequence data is complicated by errors introduced during sample preparation and

data analysis. We used synthetic RNA controls and simulated data to test seven variant calling

tools across a range of allele frequencies and simulated coverages. We show that choice of

variant caller, and use of replicate sequencing have the most significant impact on single

nucleotide variant (SNV) discovery and demonstrate how both allele frequency and coverage

thresholds impact both false discovery and false negative rates. We use these parameters to find

minority variants in sequencing data from SARS-CoV-2 clinical specimens and provide guidance

for studies of intrahost viral diversity using either single replicate data or data from technical

replicates. Our study provides a framework for rigorous assessment of technical factors that

impact SNV identification in viral samples and establishes heuristics that will inform and improve

future studies of intrahost variation, viral diversity, and viral evolution.

IMPORTANCE

When viruses replicate inside a host, the virus replication machinery makes mistakes.

Over time, these mistakes create mutations that result in a diverse population of viruses inside

the host. Mutations that are neither lethal to the virus, nor strongly beneficial, can lead to minority

variants that are minor members of the virus population. However, preparing samples for

sequencing can also introduce errors that resemble minority variants, resulting in inclusion of false

positive data if not filtered correctly. In this study, we aimed to determine the best methods for

identification and quantification of these minority variants by testing the performance of seven

commonly used variant calling tools. We used simulated and synthetic data to test their

performance against a true set of variants, and then used these studies to inform variant

identification in data from clinical SARS-CoV-2 clinical specimens. Together, analyses of our data

provide extensive guidance for future studies of viral diversity and evolution.

INTRODUCTION

Large population sizes, high replication rates, and error prone polymerases all contribute to the generation of sequence diversity found in viral infections (1-5). Natural selection acts on this diversity, contributing to viral evolution. RNA viruses have some of the highest mutation rates among viruses (1, 6, 7). To replicate their genomes, RNA viruses must encode their own RNA-dependent RNA-polymerases (RdRp), which often lack proofreading capabilities. Coronaviruses are a notable exception, as they possess a distinct protein with 3'-5' exonuclease capability (1, 8, 9). Most errors made during replication—up to 40% in RNA viruses—are lethal (10, 11). Beneficial mutations make up a much smaller proportion, and these, along with neutral mutations, comprise the substitution rate. This substitution rate can be used to estimate the viral evolutionary rate, an important calculation in considering viral spread, pandemic potential, and vaccine design (4, 12).

Due to the large population sizes of RNA viruses, intrahost bottlenecks, and genetic drift, genetic diversity within a host is dynamic, with frequencies of mutations constantly rising and falling (13). Mutations can lead to changes in the consensus sequence, e.g., where the allele frequency (AF) is greater than 50%, and these specific sets of mutations separate globally circulating virus populations into clades. Mutations in the virus genomes that are not the majority within an infected host (i.e., present at lower than 50% frequency) represent minority variants. Deep sequencing enables the capture of intrahost variation, both at the majority and minority level, enabling the identification of variants and estimation of their frequency. Studying intrahost variation can help in tracking viral spread, estimating population bottleneck sizes, and identifying key amino acid changes that differentiate new viral strains (14-17). Additionally, minority variants can highlight regions of the genome under selection or regions with increased mutational tolerance, as well as allow for detection of subtle population shifts within the infected host and discovery of possible drug resistance mutations (18, 19). Thus, information gleaned from studying intrahost viral diversity has major implications for vaccine, monoclonal antibody, and drug development.

Given the many applications of studying intra-host viral diversity, accurately identifying

and quantifying viral variants is essential. Precise identification of viral variants, especially those

at low frequencies, is complicated by the fact that viral genome sequencing often requires reverse

transcription and amplification, which, along with library preparation and the sequencing process,

are error prone. Thus, distinguishing true sequence variation from technical and experimental

noise is challenging. Typically, several ad hoc metrics are used to filter variants, such as applying

frequency and coverage cutoffs to sequencing data, however, the frequency at which identified

variants are considered valid can vary widely (20-27). Most studies using large sample cohorts,

or performing analyses on publicly available data, generally use single replicate data, despite

evidence suggesting that replicate sequencing may be essential for filtering false positive minority

variants (27). Despite the large number of studies analyzing minority variants in virus data, there

is no consensus on what coverage cutoffs and allele frequency cutoffs to use, and no large-scale

studies have been performed to determine what thresholds lead to the highest confidence in

variant identification.

In addition to the diversity of cutoffs used for single nucleotide variant (SNV) identification,

there is also great diversity in the variant calling software available. Variant callers are often

designed with specific functions in mind, such as identifying germline or somatic mutations in

cancer genomes or single nucleotide variants in viral populations (28, 29). The function for which

a variant caller is designed can have a significant impact on the statistics used and assumptions

made by the software. Tools designed for detection of germline mutations, such as

HaplotypeCaller and freebayes, must consider the very large reference genome, higher frequency

variants, the diploid nature of the genome, the possibility of copy number variation, and long

repetitive regions or large insertions or deletions (30-35). In these instances, local realignment of

haplotypes may be most effective (28). By contrast, software used for somatic mutations in

tumors, such as Mutect2 and Varscan, or for viral diversity, such as iVar and timo (a variant caller

developed in our lab), use base by base comparisons, or a combination of this with haplotype-

based alignment, to find lower frequency variants (27, 33-36). These tools also may need

alternative methods to reduce false positive calls to account for PCR errors introduced during

amplification of the viral genome (29). Due to the differences in bioinformatic and statistical

approaches used by each variant calling tool, identifying the tool that is the best fit for the specific

research question being studied is essential. Some tools have been tested in pairwise

comparisons (27, 35), however little work has been done to extensively test the performance of

the many available tools on different viruses, across sequence coverages, and at various allele

frequencies in viral deep sequencing data.

Here, we tested seven variant callers on simulated, synthetic, and clinical deep

sequencing data. We tested each tool across a range of coverages, allele frequencies, and

experimental designs to determine the optimal parameters that should be used to decrease false

positive variant identification, without sacrificing true positive data. To compare performance

between a small RNA virus with a high mutation rate, and a large RNA virus with proofreading

capability, we tested the variant callers on two viruses of particular interest in the viral diversity

field, influenza and SARS-CoV-2. We find that choice of variant caller, and use of replicate

sequencing have the most significant impact on SNV discovery and demonstrate how both allele

frequency and coverage thresholds impact both false discovery and false negative rates. We also

provide guidance on best practices for leveraging deep sequencing data from public repositories

for intrahost studies. These analyses provide a resource for studies aiming to assess intrahost

viral diversity in SARS-CoV-2 or influenza, and they lay the groundwork for similar studies in other

viruses.

MATERIALS AND METHODS

Extended methods are available in the supplementary materials.

Generation of Simulated Data

Reads were simulated using NEAT (v2.0) by constructing a mutation, error, fragment

length, and GC model for each viral type (37). The models were provided to NEAT genReads.py

along with reference fasta files and a mutation rate of 0.009 (0.9%) for influenza and 0.0045

(0.45%) for SARS-CoV-2 to produce a "golden VCF" containing a defined number of SNVs in

each virus. Simulated random PCR errors were also added to each replicate using

genReads.py (NEAT). Several copies of the replicate golden VCFs were made, each with the

same variants but with differing allele frequencies (AF). These VCFs were used to simulate

paired end fastq libraries at 100,000X genome coverage and down-sampling was used to

simulate lower coverages.

Sequences were trimmed using trimmomatic v0.36 (38), aligned to the respective

reference genome with BWA mem v0.7.17 (39), and duplicate reads were marked using GATK

MarkDuplicatesSpark v4.1.7.0 (40). Variants were called in each replicate with seven different

tools, using multiple parameter configurations for each tool (**Table S1).** A VCF file containing the

intersection of the two replicates was generated using bcftools isec (v1.9) (39). The pipeline used

for data simulation, sequence processing, variant calling, and analysis is available at

https://github.com/gencorefacility/MAD2.

Synthetic RNA generation, library preparation, and data processing

Synthetic, 'wildtype' (WT), influenza genomic RNA and variant RNA (created by adding

18, 14 and 14 known nucleotide changes into the WT PB2, HA, and NA segments respectively)

were synthesized as double-stranded DNA (gBlocks). The DNA was *in vitro*-transcribed with the

HiScribeTM T7 High Yield RNA Synthesis Kit (Invitrogen). RNA samples were diluted to an equal

copy number concentration of 6×10^8 copies/μL. WT and variant RNA were mixed at equal

molarity. The pools were mixed at various frequencies (50%, 25%, 12.5%, 6.25%, 3.13%, 1.56%,

0.78%, 0.39%), and diluted to various copy number concentrations (6×10^6 - 6×10^3 copies/µL).

cDNA was generated and libraries were prepared using the Nextera XT library preparation

kit (Nextera), with all volumes scaled down to 0.25x of the manufacturer's instructions, cleaned

with AMPure beads, and pooled at equal molarity. Libraries were sequenced on the Miseq 300

Cycle v2 using 2 x 75 pair-end reads. Samples were amplified and sequenced in duplicate and

analyzed with the pipeline described above, with the addition of adapter trimming. Synthetic

SARS-CoV-2 data from a similarly designed study was downloaded from SRA (PRJNA682212)

and processed as above (26).

SARS-CoV-2 clinical sample preparation, processing, and variant calling

Total RNA was extracted from 300 µL of nasopharyngeal (NP) or mid-turbinate (MT)

swabs collected at the NIH Clinical Center as part of diagnostic testing between 07/24/2020 and

03/31/2021 (**Table S2**). All samples were de-identified and anonymized.

RNA from samples was extracted using the NUCLISENS easyMAG automated nucleic

acid extractor and the viral genome was amplified using a modified version of the ARTIC protocol

(https://artic.network/ncov-2019) and the methods described at

https://github.com/GhedinSGS/SARS-CoV-2_analysis. All libraries were prepared as above and

sequenced on either the Illumina MiSeq or the Illumnia NextSeq500 using either the 2x150 bp or

2x300 bp paired end protocol. All samples were processed in duplicate.

Samples were processed with the pipeline available and described above, with the

addition of merging the two SAM files (from A and B primer pools) for each biological sample into

one alignment file using Picard Tools MergeSamFiles v2.17.11. Variants were called as above

using the standard parameters for each tool (**Table S1**).

Data Availability

Synthetic influenza data (bioproject PRJNA865369) and SARS-CoV-2 data from clinical

samples are available in NCBI GenBank and SRA. Accession IDs can be found in **Table S2**. All

downstream analysis files are available at https://github.com/GhedinSGS/Optimized-

Quantification-of-Intrahost-Viral-Diversity.

Ethics Statement

All samples were anonymized and obtained with consent as part of SARS-CoV-2

diagnostic testing.

RESULTS

Simulated and synthetic data provide a 'true' set of minority variants to assess variant caller

performance.

To test the ability of each variant caller to accurately identify variants, it is essential to

know the 'true set' of variants within the data, which is not possible with real sequence data. With

this in mind, we tested the ability of six popular variant-calling software packages (Freebayes,

HaplotypeCaller (hc), iVar, Lofreq, Mutect2, and Varscan) and one in-house pipeline (timo) to

accurately identify minority variants in simulated and synthetic sequencing data (**Fig. S1**) (19-24).

Single nucleotide variants (SNVs) were simulated across three influenza virus genomes (A/H1N1,

A/H3N2, B/Victoria) and one coronavirus genome (SARS-CoV-2) at both defined and random

allele frequencies and across a range of downsampled coverages (**Fig. S1A-B**). Further, synthetic

RNA controls of three influenza virus segments (PB2, HA, and NA) containing known SNVs were

mixed in varying amounts at various dilutions to create a range of allele frequencies and genome

copy numbers (**Methods**) (**Fig. S1C-D**). Combined, we used these synthetic and simulated data

sets to test variant caller performance on a known set of SNVs.

We found that all callers performed poorly on our data using their default parameters (**Fig.**

S2A). Therefore, to compare all callers equally, we used a standard set of permissive input

parameters (min coverage = 1x, allele frequency cutoff = 0.01 (1%)) throughout our testing (**Table**

S1). When assessing the F1 statistic across a range of simulated frequencies, most variant callers

performed well at low frequencies (< 0.05 (5%)) when the coverage was high (downsampling

fraction >= 0.005 or ~500X coverage). Conversely, high frequencies were necessary for accurate

variant detection at small downsampling fractions where the average coverage was low (**Fig. 1A**).

We did not observe significant differences between the four viruses in terms of variant calling

accuracy (**Fig. 1A, Fig. S2A**). Most of the differences in performance between the variant callers

could be seen at downsampling fractions < 0.005 and allele frequencies below 0.05 (5%) (**Fig**

1A). A closer look at precision and recall for each tool at downsampling fractions of 0.002 and

0.003 indicated that many tools trade recall for precision at low frequencies (**Fig. 1B**). Some tools,

such as iVar, timo, and varscan tended to be extremely conservative, especially at low

frequencies. However, the stringency of these tools can be reduced by decreasing the input

frequency parameter from 0.01 (1%) to 0.001 (0.1%) (**Fig. S2** *custom input parameters*, **Table**

S1). The use of simulated data for initial testing showed that under ideal conditions, all variant

callers have the ability to perform well on viral sequence data. This is important for establishing

baseline performance against which to compare the variant caller performance on synthetic data

and data from clinical specimens.

While simulated data provides an ideal set of variants against which to compare the variant

calls for each tool, it lacks the reverse transcription, amplification, and sequence library

preparation steps involved in the generation of data from clinical specimens. To assess how these

sample preparation steps, along with duplicate sequencing, and SNV thresholds may impact

variant caller performance, we tested each tool on data from the synthetic RNA dataset (**Fig. S1C-**

D). The average read depth across gene segments was greater than 1000x and had similar

coverage distributions to our simulated datasets at downsampling fractions of 0.01 and 0.1 (**Fig.**

S3A). At this coverage in simulated data, we observed high F1, precision, and recall scores across

all variant callers for SNVs > 1% frequency (**Fig. 1A-B**). Observed frequencies differed from

expectation (**Fig. S3B**), likely due to mixing errors during sample preparation, but were consistent

between viral copy numbers. Comparing the observed allele frequency to the median observed

allele frequency revealed that most callers agreed on the frequency of identified variants;

however, there is considerable variance in the AF estimations despite the fact that all SNVs are

linked (**Fig. 1C, S3B**). We also tested the variant callers on a previously published synthetic

SARS-CoV-2 data set (26). Here, HaplotypeCaller and timo found more true variants than the

other callers, especially at higher viral load (**Fig. S3C**). Across all callers, the frequencies of

observed variants in the SARS-CoV-2 data were less accurate than in the synthetic flu data i.e.

the variance in allele frequencies is higher) (**Fig. S3B-C**). This is potentially due to the overall low

coverage of these samples and the amplicon-based sequencing method required for larger RNA

genomes, suggesting that allele frequency estimation may be less reliable in SARS-CoV-2 data.

Frequency thresholds and sequencing replicates reduce false positive SNVs

Previous studies have reported the necessity of establishing frequency and coverage

thresholds as well as having replicate sequencing to decrease false-positive SNVs in a data set

(27, 40). Given that most publicly available data consist of single replicate sequencing data, we

aimed to establish coverage and frequency thresholds that would minimize the false discovery

rate (FDR) and false negative rate (FNR) to levels comparable to those observed using two

replicates. To do this, we used both simulated and synthetic datasets with standard input

parameters (**Fig. S2A**) and ignored the 'binocheck' requirement from timo (requiring variants to

be found in near equal numbers of forward and reverse reads), allowing us to test the performance

of timo on low frequency SNVs.

In both the synthetic and simulated data, false positive SNVs were found across read

depths but were primarily at allele frequencies less than 0.03 (**Fig. 2A-2B**). Therefore, applying

frequency thresholds to single replicate data lowered the false discovery rate (FDR) for all callers

(**Fig. 3A, S4A**). While establishing coverage cutoffs did not drastically impact the number of false

positive calls in either the simulated or synthetic dataset, coverage and library size are important

factors when considering SNV recall (**Fig. 1A, 2A-B, S5**). However, using frequency thresholds

did come at the cost of significantly increasing the FNR, as true SNVs found at low frequencies

were filtered from the data. In contrast, keeping only SNVs shared between the two replicates

dramatically decreased the FDR, while maintaining relatively low FNRs (**Fig. 3, S4**).

HaplotypeCaller, LoFreq, and Mutect2 called notably higher numbers of false positive SNVs in

synthetic data, including many that were maintained even after merging replicates—indicating

that these callers are making consistently incorrect SNV calls. Further, these three callers had

multiple instances where true positive SNVs were identified at high frequencies (AF > 0.05) in one

replicate and were entirely absent in the other (**Fig. S6A**). Importantly, replicates also increased

the accuracy of allele frequency estimation of true positive variants in simulated data. The effect

of sequencing replicates on allele frequency is especially pronounced for low coverage data,

where the percent error of allele frequency estimation is pointedly lower for all tools when using

replicates (**Fig. S6B**). This effect was less pronounced in synthetic data, likely due to the high

coverage of these samples, again demonstrating the importance of read coverage for accurate

allele frequency estimation (**Fig. S6C**).

Together, these results indicate that using replicate sequencing with less stringent

frequency cutoffs (AF >= 0.01) is the best combination to reduce the FDR while maintaining a low

FNR. However, when replicate sequencing is unavailable, high coverage across the genome and

strict frequency cutoffs (AF >= 0.03) are necessary.

Choice of variant caller significantly impacts set and frequency of identified variants in real SARS-

CoV-2 data using single replicate data.

While simulated and synthetic data allow for testing minority variant callers and cutoffs in

a controlled setting, real data will always be more unpredictable. Thus, after using simulated and

synthetic data to assess variant caller performance across frequencies and coverages, we tested

how the callers performed on SARS-CoV-2 sequence data from diagnostic samples. Based on

the simulated and synthetic data testing, we determined that a coverage cutoff of 200X and an

allele frequency cutoff of 0.03 in single replicate data minimized false positive calls without

sacrificing large amounts of true positive data with most variant calling tools. To test the variant

calling tools on high-quality data, we used only samples where at least 80% of the genome had

a read depth over 200x coverage cutoff in both sequencing replicates (**Fig. S7A**). We used each

variant calling tool to identify minority variants in these samples and filtered them using a read

depth cutoff of 200 and an allele frequency cutoff of 0.03.

We were interested in how similar the set of variants was that was identified by each caller.

As a proof of principle, we filtered the set of variants for those present above an allele frequency

of 0.5 and at read depths greater than 5x to identify consensus changes (AF >=50%, or major

variants) within the data. As expected, the tools largely agreed on the consensus changes within

the data (**Fig. S7B**). There was a small set of major variants that the callers did disagree on,

however, most of which were a result of differences in the way some callers identify indels or

handle variant at consecutive nucleotide positions. For the purposes of this study, indels were

excluded from the analysis. These data indicate that even at high allele frequencies, the variant

callers disagree to some extent on the set of variants present in clinical data, an important

consideration when choosing how to define consensus sequences from SARS-CoV-2 data.

We then analyzed the intersection of the minority variants (allele frequencies between

0.03 and 0.5) identified by each tool. The total number of variants identified varied greatly between

the callers, with Varscan calling the fewest variants by far, followed by timo and Lofreq, in line

with the more conservative nature of these callers observed in the previous analyses (**Fig. 4A**).

Of note, we found that replicate 2 data had much higher numbers of minority variants, particularly

at very low frequencies, regardless of Ct value or date of sequencing. This suggests that freeze

thawing samples may impact minor variant numbers (**Fig. S7C**). When comparing the set of

minority variants identified by each of the seven tools, there was significant disagreement

between the variants. Mutect2 and HaplotypeCaller identified many variants that other callers did

not, particularly in replicate 1, and missed several variants identified by the other callers. This was

similar to the performance of these callers on the synthetic data sets (**Fig. S7D**). Given the high

number of false positives identified by HaplotypeCaller, Mutect2 and Lofreq in the simulated and

synthetic datasets, we focused on the intersection of minority SNVs found in just the other four

variant callers: Freebayes, iVar, Varscan, and timo. Of all the minority variants found in the data,

104 from replicate 1, and 142 from replicate 2 were identified by all four of the variant callers (**Fig.**

4B). Overall, choice of variant caller appears to have a significant impact on the set of minority

variants identified in SARS-CoV-2 data from clinical specimens.

Many studies of minority variants investigate frequency of minority variants to calculate

selection, bottleneck size, and potential for transmission (24, 41). We were interested in how well

the variant callers agreed on the frequency at which variants were identified. We plotted the

frequency of a variant in one caller against the frequency in each other caller and found that most

of the minority variant callers were strikingly similar in their frequency calls of shared variants.

Timo, Freebayes and iVar all showed almost complete agreement on frequency of the variants

(**Fig. 4C**). Varscan showed more variation in frequency, generally calling variants at a lower

frequency than the other three tools (**Fig. 4C**). Of interest, variants called by one caller but not

another spanned a frequency range of 0.03 all the way to 0.5, indicating that even high frequency

minority variants were often not agreed upon by variant callers. These data show that choice of

variant caller not only affects the set of the minority variants that are identified in a data set, but

also the frequency of those variants.

Most minority variants in data from SARS-CoV-2 clinical specimens are not reproducible across

sequencing replicates

In our sequencing data, the number of variants identified in each replicate by each tool

was markedly different, suggesting that many of the identified minor variants may not be true

variants introduced through viral replication, but instead technical artifacts (**Fig. 4A, S7C-D**). As

was shown with our simulated and synthetic data, errors introduced through PCR, library

preparation, and sequencing are mostly random, and therefore less likely to reappear and be

identified across multiple sequencing replicates, particularly when using Freebayes, iVar, timo, or

Varscan (**Fig. 3**). To find high confidence minority variants, we looked at the intersection of

variants between the two replicates using each caller, using a lower allele frequency threshold,

0.01, as established in synthetic data for merged replicates (**Fig. 3**). iVar and Freebayes called

the highest number of reproducible variants, while timo called the fewest number of reproducible

variants, however the percentage of reproducible variants compared to variants identified in single

replicates was the highest (**Fig. 5A**). Timo is one of the most conservative callers that was tested;

these data indicate that being conservative may lead to increased reproducibility and therefore

increased confidence when used on single replicate data. It is however important to note that the

relatively low percentages of reproducible variants are likely skewed by the high numbers of low

frequency variants found in replicate 2 (**Fig. 5A, S7C**). When we looked at the intersection of only

the variants found by each tool in both replicates, only 80 SNVs were found by all callers across

replicates, suggesting again that variant callers do not agree on the set of minority variants

present (**Fig. 5B**). Together, these data suggest that most minority variants are not reproducible

across replicates and support the idea that more than any other criteria, sequencing replicate has

the highest impact on the set of minority variants identified (**Fig. 5B, S7C-D**).

Variants identified by all variant callers show the most reproducible frequencies

Using synthetic data, we showed that in a controlled setting, SNVs that were found in both

sequencing replicates generally showed reproducible frequencies (**Fig. 2C**). Given that frequency

is an important metric in most analyses performed using minority variant data, we wanted to test

if this held true in clinical samples. Plotting the frequency in replicate 1 against the frequency in

replicate 2 revealed that while some variants showed consistent frequencies, some differed

drastically—identified at 5-10% in one replicate and as high as 45-50% in the other replicate (**Fig.**

5C). These data were striking as they reveal that averaging frequency across replicates, or

performing only one sequencing replicate, could drastically alter downstream analyses performed

using these numbers. Interestingly, when we looked at the variants that were reproducible across

replicates and found by most, or all the variant callers, frequency tended to be much more

consistent than those identified only in a single replicate, or by a single caller (**Fig. 5C,** dark red

points). Together, these data suggest that confidence in each variant and its frequency is
increased with replicate sequencing and identification by many variant callers.

Since replicate sequencing data is not always available, we investigated what frequency
cutoff could be applied such that single replicate data closely resembled the merged replicate
data. To do this, we looked at the intersection of SNVs called in both replicates by Freebayes,
iVar, timo, and Varscan (80 variants shown in **Fig. 5B**) and compared those to the intersection of
SNVs called by the same four callers in each individual replicate (**Fig. 4A**). We then applied allele
frequency cutoffs between 0.01 and 0.1 to determine the best cutoff for use on single replicate
data. Here, we identify a true positive as a variant present in the reproducible set, and a false
positive as any other variant found in a single replicate. As was noted previously, we find that
replicate 2 data shows an increased number of SNVs, perhaps due to freeze/thawing of samples
between preparations (**Fig. S7C, 5D**). As such, replicate 1 is likely more representative of what
single replicate data may typically look like. At an allele frequency cutoff of 0.01, all true positives
were found, but the number of false positives was very high, while a frequency cutoff of 0.05 or
0.1 removed an outsized number of true positives from the dataset (**Fig. 5D**). Based on these
data, we suggest an allele frequency cutoff of 0.03 when only single replicate data is available, a
cutoff that was also confirmed in the simulated and synthetic data sets (**Fig. 3**). We further suggest
using the intersection of multiple variant callers to increase confidence in the data, especially
when estimating SNV frequency (**Fig. 5B**). Using all variant callers for analysis would likely be
tedious and unrealistic, thus we looked at the intersection of just two callers, iVar and timo, and
we found a similar trade-off in true positive and false positive data when using a single replicate
and a cutoff of 0.03 (**Fig. 5E**). Based on these data, it is clear that there are many considerations
necessary when performing minority variant analyses, and parameters and cutoffs should thus
be chosen carefully and thoughtfully depending on the data available. In general, using replicate
data and multiple callers provides the highest confidence set of SNVs and the most accurate
frequencies.

DISCUSSION

It has long been understood that intrahost viral populations are heterogeneous in nature, however capturing and measuring this viral diversity is complicated due to errors introduced during preparation and sequencing. We set out to identify the optimal tools, parameters, and filtering methods necessary for accurate variant identification. To accomplish this goal, we used a combination of simulated and synthetic influenza and SARS-CoV-2 sequence data to test the technical and experimental challenges and limitations of minority variant analyses. We found that sequencing depth and choice of variant caller has a significant impact on sensitivity of minor variant calls, and most false positive SNVs were detected at either low allele frequency, or low read depth. Additionally, our results show that replicate sequencing allows for the use of lower frequency thresholds, and this combination provides the best results, keeping the false discovery rate low, without sacrificing true positive data. We tested our optimized frequency and coverage cutoffs using SARS-CoV-2 sequence data from clinical infections. Most variant callers did not agree on the set of minor variants in the virus sequence data from clinical samples, and most minority variants were not reproducible across replicates. Ultimately, we determined that using a combination of sequencing replicates and multiple variant callers, along with moderate allele frequency and coverage cutoffs, can increase confidence in SNV calls. Our results outline the main considerations for minority variant analyses and highlight the intricacies and difficulties of studies of this nature. Further, we provide a framework for designing minority variant analyses, which will ultimately lead to more accurate conclusions surrounding viral transmission and evolution.

Using a standardized set of parameters, most callers performed relatively similarly on high coverage simulated data, having both high precision and high recall. The main differences in caller performance were seen in lower coverage data or at low frequencies. As many minority variants are found at low frequencies, understanding how tools perform under these conditions is more relevant to analyses of real sequencing data. Timo had the lowest recall at lower coverages and

simulated frequencies due to its rigid requirements for SNVs to be above the 0.01 threshold

parameter, while many other callers found SNVs at or below this frequency, regardless of setting

a 0.01 AF cutoff. Timo, iVar, and Varscan all have the functionality to drop the input frequency

parameter down to 0.001. Decreasing this parameter did not change the accuracy of iVar and

Varscan but did increase the recall of timo. These data highlight the importance of optimizing

bioinformatic tools to one's own data.

As previously observed by our group and others, the best method for filtering out errors

generated during sample processing is to sequence each sample twice and only keep the SNVs

found in both replicates. Sequencing replicates removed nearly all false positive calls in simulated

data and significantly reduced the number of false positive SNVs in the synthetic datasets.

However, for the synthetic datasets, the number of false positive SNVs was highly dependent on

the variant caller used. HaplotypeCaller, Lofreq, and Mutect2 were all made and optimized for

identifying variants in cancer cell datasets and had significantly higher false discovery rates than

tools designed for viral use, particularly at low allele frequencies. Adjusting the filtering or input

parameters on these callers may better optimize them for their use on viral data. For example,

HaplotypeCaller suggests additional filtering of output data, however, when performed on this

dataset, SNV detection was significantly reduced. Without this additional filtering, most variants

are identified (high recall) but high numbers of false positives are included, suggesting additional

optimization could improve performance.

The use of simulated and synthetic data allows tools to be tested against a true set of

minority variants. This is invaluable given that it is not possible to know which SNVs are real, and

which are not, in clinical sequencing data. However, each of these controlled settings comes with

both benefits and limitations. Simulated data allows for testing of various coverages using random

downsampling, showing how low coverage data impacts caller performance and allele frequency

estimation. Since high coverage is often the goal in sequencing, and coverage is unpredictable

with real samples (42), this is a variable that can only be truly tested through simulation.

Conversely, while error models in simulated data can imitate PCR and sequencing errors, there

are limitations to how 'real' these errors look to variant callers. Synthetic data allows for real error

to be introduced during sample preparation and ultimately be easily separated from real variation

during analysis. This gives a more accurate depiction of false positives that would exist in clinical

data and variant caller performance on synthetic data may be more representative of true

performance. However, both methods are limited in their ability to mimic sequence data from

clinical samples where the RNA is not as homogeneous and intact as with synthetic RNA, and

likely contains human contamination from sample collection and potential degradation from

freezing, thawing, and handling (43). This is evident in the disagreement between variant callers

and replicates, both on the set of minority variants and on their frequencies, in real data.

Combined, the simulated, synthetic, and clinical data sets show that there will always be a trade-

off between inclusion of the maximum number of true variants, and inclusion of false positive data.

Our study provides an extensive framework for studying minority variants in sequence

data from clinical samples, outlining major considerations around choice of variant caller,

application of frequency and coverage thresholds, and use of replicate sequencing. Further, we

have established a pipeline that can be used for further testing and optimization of parameters,

or for other viruses. This work will inform and improve future studies of intrahost variation and

estimates surrounding viral diversity and viral evolution.

FIGURE LEGENDS

Figure 1. Variant caller performance on simulated and synthetic data with variants at set

allele frequencies. (A) F1 statistic for each variant caller across a range of downsampling

fractions and allele frequency values. Values shown are mean and standard deviation of the four

viruses (A/H1N1, A/H3N2, B/Victoria, SARS-CoV-2) using standard input parameters (**Table S1**).

Color represents the variant caller used. **(B)** Precision/Recall graphs of each variant caller across

allele frequencies (point shape) for downsampling fractions 0.003 (top left) and 0.002 (top right)

or across downsampling fractions (point shape) for allele frequencies (AF) 0.02 (bottom left) and

0.03 (bottom right). Color represents variant caller. Mean and standard deviation are shown

across the four viruses for precision and recall scores. **(C)** Scatter plot showing median observed

frequency (x-axis) versus individual observed frequency (y-axis) for each synthetic influenza gene

segment and SARS-CoV-2 genome. Black dots indicate the median value for each expected

frequency. Color represents the variant caller used.

Figure 2. Frequency and coverage of false positive variants in synthetic and simulated

data. (A-B) Scatter plots and associated histograms showing number of false positive SNVs

across allele frequencies and read depths in synthetic (**A**) or simulated (**B**) data. Dotted lines are

drawn at allele frequency = 0.03 and read depth = 200x. Color represents the variant caller used.

Figure 3. Effect of frequency cutoffs and sequencing replicates on false discovery rate and

false negative rate in synthetic and simulated data. (A) False discovery rate (FDR) of synthetic

data using either single replicate data (colored points and lines) with applied frequency cutoffs

(line type) or merged two replicate data without cutoffs (solid black points and lines). Lines

represent mean across viruses. **(B)** False negative rate (FNR) of synthetic data using either single

replicate data (colored points and lines) with applied frequency cutoffs (line type) or merged two

replicate data without cutoffs (solid black points and lines) as above.

Figure 4. Effect of variant caller on identification and allele frequency estimation of SNVs in SARS-CoV-2 data from clinical samples. (A) Bar plot showing raw number of variants identified by each variant caller in replicate 1 (left bar) or replicate 2 (right bar). **(B)** Upset plot showing agreement of minority variants between variant callers in each replicate using an allele frequency cutoff of 0.03 and coverage cutoff of 200X. Vertical bars indicate the size of the shared set of variants while dots and connecting lines show which callers share a given set of identified variants. **(C)** Scatter plot showing the output frequency of minority variants identified by two different variant callers. Color represents replicate. Variants with frequency of 0 were not identified by that variant caller.

Figure 5. Reproducibility of minority variants across sequencing replicates. (A) Bar plot showing number of reproducible variants across sequencing replicates by each variant caller. Percentages shown are the percentage of total individual variants that were reproducible. Background bars indicate the total number of variants found by each tool in each replicate (r1-left, r2-right). **(B)** UpsetR plot showing overlap of reproducible variants across Freebayes, iVar, timo, and Varscan, using a frequency cutoff of 0.01 and coverage cutoff of 200x. Vertical bars indicate the size of the shared set of variants while dots and connecting lines show which callers share a given set of reproducible variants. **(C)** Scatter plot showing frequency of variants across sequencing replicates with frequency in replicate 1 on the x-axis and frequency in replicate 2 on the y-axis. Color represents reproducibility of each variant across variant callers and replicates. **(D,E)** Line graph showing the number of 'true positive' (D) and 'false positive' (E) in single replicate data across allele frequency cutoffs compared to merged replicate data. A TP variant is defined as a SNV found by the selected callers in both replicates (80 variants shown in **B**) and a FP is defined as any other variant found in an individual replicate by the selected callers. Color represents sequencing replicate.

SUPPLEMENTAL FIGURES

Supplemental Figure 1. Experimental setup of simulated and synthetic data generation. (A) Schematic of the SNV simulation pipeline. **(B)** Nucleotide positions of SNVs in simulated data across the genomes of A/H1N1 (n=121), A/H3N2 (n=110), B/Victoria (n=118), and SARS-CoV-2 (n=144). Gene segments for influenza A and B strains are ordered largest (PB2) to smallest (NS), left to right. **(C)** Schematic of allele frequency and viral loads used for mixing WT and variant RNA for library preparation and sequencing. **(D)** Location of synthetic SNVs across the three influenza gene segments. Gray lines represent the designed SNVs (PB2 (n=18), HA (n=14), NA (n=14)) and red lines represent the pre-mRT-PCR errors, likely generated during template preparation or *in vitro*-transcription.

Supplemental Figure 2. Variant caller performance using default, standard, and custom parameters. (A) F1 statistic for each variant caller across a range of downsampling fractions and allele frequencies. Values shown are mean and standard deviation of four viruses (A/H1N1, A/H3N2, B/Victoria, SARS-CoV-2) using default (top), standard (middle) or custom (bottom) input parameters (**Table S1**). Color represents the variant caller used. Dark timo points and lines indicate timo output with the removal of the binomial check option.

Supplemental Figure 3. Coverage and Expected vs. Observed frequency of variants in synthetic data samples (A) Coverage plots showing log10 read depth for each synthetic

influenza control across the PB2, HA, and NA gene segments. Color represents individual

samples. **(B)** Expected (x-axis) versus observed frequency (y-axis) for synthetic influenza SNVs

(replicate one only). Color represents the copy number. **(C)** Expected (x-axis) versus observed

frequency (y-axis) for synthetic SARS-CoV-2 SNVs. Dashed black lines represent y=x.

Supplemental Figure 4. Effect of frequency cutoffs and sequencing replicates on false

discovery rate and false negative rate in simulated data. (A) False discovery rate (FDR) of

simulated data using either single replicate data (colored points and lines) with applied frequency

cutoffs (line type) or merged two replicate data without cutoffs (solid black points and lines).

Background points represent values for individual viruses. Lines represent mean across viruses.

(B) False negative rate (FNR) of simulated data using either single replicate data (colored points

and lines) with applied frequency cutoffs (line type) or merged two replicate data without cutoffs

(solid black points and lines) as above.

Supplemental Figure 5. Effect of coverage cutoffs and sequencing replicates on false

discovery rate and false negative rate in synthetic and simulated data. (A, B) False discovery

rate (FDR) (**A**) of false negative rate (FNR) (**B**) of synthetic data using either single replicate data

(colored points and lines) with applied coverage cutoffs (line type) or merged two replicate data

without cutoffs (solid black points and lines). Background points show individual values for each

of the four viruses. Lines represent mean across viruses. (**C, D**) False discovery rate (FDR) (**C**)

of false negative rate (FNR) (**D**) of synthetic data using either single replicate data (colored points

and lines) with applied coverage cutoffs (line type) or merged two replicate data without cutoffs

(solid black points and lines) as above.

557
Supplemental Figure 6. Effect of sequencing replicates on the accuracy of allele frequency

estimation. (A) Scatter plot showing the frequency of variants in synthetic influenza data across

sequencing replicates with frequency in replicate 1 on the x-axis and frequency in replicate 2 on

the y-axis. Color represents the SNV type. **(B, C)** Percent error ($\frac{(observed-expected)}{expected} * 100$) of

single replicate (greyed boxes) or merged replicate (open boxes) SNVs in simulated data across

downsampling fractions (**B**) or synthetic data across expected frequencies and viral loads (**C**).

Color represents the variant caller used.

Supplemental Figure 7. Quantification of majority and minority variants identified in data from

SARS-CoV-2 clinical specimens (A) Scatter plot showing Ct value against percentage of genome

with coverage over 200x after filtering for only samples with 80% of the genome over the 200x

cutoff. **(B)** Upset plot showing agreement of consensus changes between variant callers in each

replicate using an allele frequency cutoff of 0.5 and coverage cutoff of 5X. Vertical bars indicate

the size of the shared set of changes while dots and connecting lines show which callers share a

given set of identified changes. **(C)** Box and whisker plot showing number of minor variants with

indicated allele frequency cutoffs found in replicate 1 and replicate 2 sequencing data. Points

represent individual samples. Boxes and whiskers show min, first quartile, median, third quartile

and max for each replicate. **(D)** Upset plot showing agreement of minority variants between all

variant callers in each replicate using an allele frequency cutoff of 0.03 and coverage cutoff of

200x. Vertical bars indicate the size of the shared set of variants while dots and connecting lines

show which callers share a given set of identified variants.

REFERENCES

1.   Arnold JJ, Cameron CE. 2004. Poliovirus RNA-dependent RNA polymerase (3Dpol): pre-

steady-state kinetic analysis of ribonucleotide incorporation in the presence of Mg2+.

Biochemistry 43:5126-37.

2.   Sanjuan R. 2012. From molecular genetics to phylodynamics: evolutionary relevance of

mutation rates across viruses. PLoS Pathog 8:e1002685.

3.   Duffy S, Shackelton LA, Holmes EC. 2008. Rates of evolutionary change in viruses:

patterns and determinants. Nat Rev Genet 9:267-76.

4.   Peck KM, Lauring AS. 2018. Complexities of Viral Mutation Rates. J Virol 92.

5.   Domingo E. 2002. Quasispecies Theory in Virology. J Virol 76:463-465.

6.   Sanjuan R, Domingo-Calap P. 2016. Mechanisms of viral mutation. Cell Mol Life Sci

73:4433-4448.

7.   Lynch M, Ackerman MS, Gout JF, Long H, Sung W, Thomas WK, Foster PL. 2016. Genetic

drift, selection and the evolution of the mutation rate. Nat Rev Genet 17:704-714.

8.   Gorbalenya AE, Enjuanes L, Ziebuhr J, Snijder EJ. 2006. Nidovirales: evolving the largest

RNA virus genome. Virus Res 117:17-37.

9.   Smith EC, Sexton NR, Denison MR. 2014. Thinking Outside the Triangle: Replication

Fidelity of the Largest RNA Viruses. Annu Rev Virol 1:111-32.

10.   Sanjuan R, Moya A, Elena SF. 2004. The distribution of fitness effects caused by single-

nucleotide substitutions in an RNA virus. Proc Natl Acad Sci U S A 101:8396-401.

11.   Visher E, Whitefield SE, McCrone JT, Fitzsimmons W, Lauring AS. 2016. The Mutational

Robustness of Influenza A Virus. PLoS Pathog 12:e1005856.

12.   Peck KM, Chan CH, Tanaka MM. 2015. Connecting within-host dynamics to the rate of

viral molecular evolution. Virus Evol 1:vev013.

13.   Wang Y, Wang D, Zhang L, Sun W, Zhang Z, Chen W, Zhu A, Huang Y, Xiao F, Yao J, Gan

M, Li F, Luo L, Huang X, Zhang Y, Wong SS, Cheng X, Ji J, Ou Z, Xiao M, Li M, Li J, Ren P,

Deng Z, Zhong H, Xu X, Song T, Mok CKP, Peiris M, Zhong N, Zhao J, Li Y, Li J, Zhao J.

2021. Intra-host variation and evolutionary dynamics of SARS-CoV-2 populations in

COVID-19 patients. Genome Med 13:30.

14.   Martin MAK, Katia. 2021. Reanalysis of deep-sequencing data from Austria points

towards a small SARS-COV-2 transmission bottleneck on the order of one to three

virions. bioRxiv doi:https://doi.org/10.1101/2021.02.22.432096.

15.   McCrone JT, Lauring AS. 2018. Genetic bottlenecks in intraspecies virus transmission.

Curr Opin Virol 28:20-25.

16.   Wang D, Wang Y, Sun W, Zhang L, Ji J, Zhang Z, Cheng X, Li Y, Xiao F, Zhu A, Zhong B,

Ruan S, Li J, Ren P, Ou Z, Xiao M, Li M, Deng Z, Zhong H, Li F, Wang WJ, Zhang Y, Chen W,

Zhu S, Xu X, Jin X, Zhao J, Zhong N, Zhang W, Zhao J, Li J, Xu Y. 2021. Population

Bottlenecks and Intra-host Evolution During Human-to-Human Transmission of SARS-

CoV-2. Front Med (Lausanne) 8:585358.

17.   Lythgoe KA, Hall M, Ferretti L, de Cesare M, MacIntyre-Cockett G, Trebes A, Andersson

M, Otecko N, Wise EL, Moore N, Lynch J, Kidd S, Cortes N, Mori M, Williams R, Vernet G,

Justice A, Green A, Nicholls SM, Ansari MA, Abeler-Dorner L, Moore CE, Peto TEA, Eyre

DW, Shaw R, Simmonds P, Buck D, Todd JA, Oxford Virus Sequencing Analysis G, Connor

TR, Ashraf S, da Silva Filipe A, Shepherd J, Thomson EC, Consortium C-GU, Bonsall D,

Fraser C, Golubchik T. 2021. SARS-CoV-2 within-host diversity and transmission. Science

doi:10.1126/science.abg0821.

18.   Pybus OG, Rambaut A. 2009. Evolutionary analysis of the dynamics of viral infectious

disease. Nat Rev Genet 10:540-50.

19.   Rockett R, Basile K, Maddocks S, Fong W, Agius JE, Johnson-Mackinnon J, Arnott A,

Chandra S, Gall M, Draper J, Martinez E, Sim EM, Lee C, Ngo C, Ramsperger M, Ginn AN,

Wang Q, Fennell M, Ko D, Lim HL, Gilroy N, O'Sullivan MVN, Chen SC, Kok J, Dwyer DE,

Sintchenko V. 2022. Resistance Mutations in SARS-CoV-2 Delta Variant after Sotrovimab

Use. N Engl J Med 386:1477-1479.

20.   Martinez-Gonzalez B, Soria ME, Vazquez-Sirvent L, Ferrer-Orta C, Lobo-Vega R, Minguez

P, de la Fuente L, Llorens C, Soriano B, Ramos-Ruiz R, Corton M, Lopez-Rodriguez R,

Garcia-Crespo C, Somovilla P, Duran-Pastor A, Gallego I, de Avila AI, Delgado S, Moran F,

Lopez-Galindez C, Gomez J, Enjuanes L, Salar-Vidal L, Esteban-Munoz M, Esteban J,

Fernandez-Roblas R, Gadea I, Ayuso C, Ruiz-Hornillos J, Verdaguer N, Domingo E, Perales

C. 2022. SARS-CoV-2 Mutant Spectra at Different Depth Levels Reveal an Overwhelming

Abundance of Low Frequency Mutations. Pathogens 11.

21.   Dinis JM, Florek KR, Fatola OO, Moncla LH, Mutschler JP, Charlier OK, Meece JK,

Belongia EA, Friedrich TC. 2016. Deep Sequencing Reveals Potential Antigenic Variants

at Low Frequencies in Influenza A Virus-Infected Humans. J Virol 90:3355-65.

22.   Xue KS, Bloom JD. 2020. Linking influenza virus evolution within and between human

hosts. Virus Evol 6:veaa010.

23.   Valesano AL, Taniuchi M, Fitzsimmons WJ, Islam MO, Ahmed T, Zaman K, Haque R,

646   Wong W, Famulare M, Lauring AS. 2021. The Early Evolution of Oral Poliovirus Vaccine Is

647   Shaped by Strong Positive Selection and Tight Transmission Bottlenecks. Cell Host

648   Microbe 29:32-43 e4.

649   24.   Lauring AS. 2020. Within-Host Viral Diversity: A Window into Viral Evolution. Annu Rev

650   Virol 7:63-81.

651   25.   McCrone JT, Lauring AS. 2016. Measurements of Intrahost Viral Diversity Are Extremely

652   Sensitive to Systematic Errors in Variant Calling. J Virol 90:6884-95.

653   26.   Valesano ALR, Kalee E; Dimcheff, Derek E; Blair, Christopher N; Fitzsimmons, William J;

654   Petrie, Joshua G; Martin, Emily T; Lauring, Adam S. 2021. Temporal dynamics of SARS-

655   CoV-2 mutation accumulation within and across infected hosts. bioRxiv

656   doi:https://doi.org/10.1101/2021.01.19.427330.

657   27.   Grubaugh ND, Gangavarapu K, Quick J, Matteson NL, De Jesus JG, Main BJ, Tan AL, Paul

658   LM, Brackney DE, Grewal S, Gurfield N, Van Rompay KKA, Isern S, Michael SF, Coffey LL,

659   Loman NJ, Andersen KG. 2019. An amplicon-based sequencing framework for accurately

660   measuring intrahost virus diversity using PrimalSeq and iVar. Genome Biol 20:8.

661   28.   Koboldt DC. 2020. Best practices for variant calling in clinical sequencing. Genome Med

662   12:91.

663   29.   Olson ND, Lund SP, Colman RE, Foster JT, Sahl JW, Schupp JM, Keim P, Morrow JB, Salit

664   ML, Zook JM. 2015. Best practices for evaluating single nucleotide variant calling

665   methods for microbial genomics. Front Genet 6:235.

666   30.   Stead LF, Sutton KM, Taylor GR, Quirke P, Rabbitts P. 2013. Accurately identifying low-

667   allelic fraction variants in single samples with next-generation sequencing: applications

668   in tumor subclone resolution. Hum Mutat 34:1432-8.

669   31.   Auwera Gvd, O'Connor BD. 2020. Genomics in the cloud : using Docker, GATK, and WDL

670   in Terra, First edition. ed. O'Reilly Media, Sebastopol, CA.

671   32.   Cibulskis K, Lawrence MS, Carter SL, Sivachenko A, Jaffe D, Sougnez C, Gabriel S,

672   Meyerson M, Lander ES, Getz G. 2013. Sensitive detection of somatic point mutations in

673   impure and heterogeneous cancer samples. Nat Biotechnol 31:213-9.

674   33.   Garrison EM, Gabor. 2012. Haplotype-based variant detection from short-read

675   sequencing. ArXiv 1207.3907v2.

676   34.   Benjamin D, Sato T, Cibulskis K, Getz G, Stewart C, L. L. 2019. Calling Somatic SNVs and

677   Indels with Mutect2. bioRxiv doi:https://doi.org/10.1101/861054.

678   35.   Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, Miller CA, Mardis ER, Ding

679   L, Wilson RK. 2012. VarScan 2: somatic mutation and copy number alteration discovery

680   in cancer by exome sequencing. Genome Res 22:568-76.

681   36.   Wilm A, Aw PP, Bertrand D, Yeo GH, Ong SH, Wong CH, Khor CC, Petric R, Hibberd ML,

682   Nagarajan N. 2012. LoFreq: a sequence-quality aware, ultra-sensitive variant caller for

683   uncovering cell-population heterogeneity from high-throughput sequencing datasets.

684   Nucleic Acids Res 40:11189-201.

685   37.   Wybo WA, Jordan J, Ellenberger B, Marti Mengual U, Nevian T, Senn W. 2021. Data-

686   driven reduction of dendritic morphologies with preserved dendro-somatic responses.

687   Elife 10.

688   38.   Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina

689   sequence data. Bioinformatics 30:2114-20.

690   39.   Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler

691   transform. Bioinformatics 25:1754-60.

692   40.   Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A,

693   Jordan T, Shakir K, Roazen D, Thibault J, Banks E, Garimella KV, Altshuler D, Gabriel S,

694   DePristo MA. 2013. From FastQ data to high confidence variant calls: the Genome

695   Analysis Toolkit best practices pipeline. Curr Protoc Bioinformatics 43:11 10 1-11 10 33.

696   41.   Zhao L, Illingworth CJR. 2019. Measurements of intrahost viral diversity require an

697   unbiased diversity metric. Virus Evol 5:vey041.

698   42.   Lambisia AW, Mohammed KS, Makori TO, Ndwiga L, Mburu MW, Morobe JM, Moraa

699   EO, Musyoki J, Murunga N, Mwangi JN, Nokes DJ, Agoti CN, Ochola-Oyier LI, Githinji G.

700   2022. Optimization of the SARS-CoV-2 ARTIC Network V4 Primers and Whole Genome

701   Sequencing Protocol. Front Med (Lausanne) 9:836728.

702   43.   Li L, Li X, Guo Z, Wang Z, Zhang K, Li C, Wang C, Zhang S. 2020. Influence of Storage

703   Conditions on SARS-CoV-2 Nucleic Acid Detection in Throat Swabs. J Infect Dis 222:203-

704   205.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5