1 **Cecropins contribute to *Drosophila* host defence against fungal and**
2 **Gram-negative bacterial infection**
3

4 A. Carboni*, M.A. Hanson*, S.A. Lindsay†, S.A. Wasserman†, and B. Lemaitre*
5

6 * Global Health Institute, School of Life Science, École Polytechnique Fédérale de Lausanne
7 (EPFL), Lausanne, Switzerland. † Division of Biological Sciences, University of California
8 San Diego (UCSD), La Jolla, California, United States of America.

10 **Cecropins in *Drosophila* innate immunity**
11

Keywords:

Drosophila melanogaster, innate immunity, antimicrobial peptides, resistance,

CRISPR/Cas9, immune effectors, humoral immunity, Cecropin

Corresponding author(s):

Alexia Carboni (alexia.carboni@epfl.ch, +4121 693 18 33)

Bruno Lemaitre (bruno.lemaitre@epfl.ch, +4121 693 18 31)

Address

EPFL SV / GHI / UPLEM

SV 3838

Station nº19

CH-1015 Lausanne

ORCID IDs:

Carboni: https://orcid.org/0000-0002-5028-5724

Hanson: https://orcid.org/0000-0002-6125-3672

Wasserman: https://orcid.org/0000-0003-1680-3011

Lemaitre: https://orcid.org/0000-0001-7970-1667

ABSTRACT

Cecropins are small helical secreted peptides with antimicrobial activity that are widely distributed among insects. Genes encoding Cecropins are strongly induced upon infection, pointing to their role in host-defence. In *Drosophila*, four *Cecropin* genes clustered in the genome (*CecA1, CecA2, CecB* and *CecC*) are expressed upon infection downstream of the Toll and Imd pathways. In this study, we generated a short deletion *ΔCec^{A-C}* removing the whole *Cecropin* locus. Using the *ΔCec^{A-C}* deficiency alone or in combination with other antimicrobial peptide (AMP) mutations, we addressed the function of Cecropins in the systemic immune response. *ΔCec^{A-C}* flies were viable and resisted challenge with various microbes as wild-type. However, removing *ΔCec^{A-C}* in flies already lacking ten other AMP genes revealed a role for Cecropins in defence against Gram-negative bacteria and fungi. Measurements of pathogen loads confirm that Cecropins contribute to the control of certain Gram-negative bacteria, notably *Enterobacter cloacae* and *Providencia heimbachae*. Collectively, our work provides the first genetic demonstration of a role for Cecropins in insect host defence, and confirms their *in vivo* activity primarily against Gram-negative bacteria and fungi. Generation of a fly line (*ΔAMP14*) that lacks fourteen immune inducible AMPs provides a powerful tool to address the function of these immune effectors in host-pathogen interactions and beyond.

INTRODUCTION

In the late nineteen seventies, as immunologists were characterizing the antibody immune response of mammals, pioneering studies revealed that insects could resist infection by fearsome human pathogens despite lacking an adaptive immune system. Eventually a landmark discovery by Hans Boman and colleagues showed that insects

produced antimicrobial peptides (AMPs) following infection (Steiner *et al.* 1981),

invigorating interest in innate immunity (Ganz *et al.* 1985; Lemaitre 2004). These AMPs

differed from other previously identified immune effectors in their small size, cationic

charge, and amphipathic structure, which allowed them to directly disrupt the negatively

charged membrane of microbes. In contrast to another class of well-known immune

effectors, the lysozymes, AMPs lack enzymatic activity and require concentrations into

the micromolar range to achieve their microbicidal effects (Imler and Bulet 2005; Seo *et*

al. 2012; Hanson and Lemaitre 2020). Research has now shown that AMPs are common

across the tree of life, with similar molecules contributing to host defence in both plants

and animals (Broekaert *et al.* 1995). While they contribute to local defence in barrier

epithelia of vertebrates, insect AMPs are most famous for being secreted upon systemic

infection from the fat body into the hemolymph, where they reach potent concentrations

(Bulet *et al.* 1999). The characterization of a plethora of AMPs with diverse modes of

action has enriched our understanding of these immune effectors. However, the

functional study of AMPs was limited until recently due to technical challenges in

mutating the small AMP genes using traditional genetic approaches. This challenge has

now been overcome with the advent of CRISPR/Cas9 technology.

Cecropins were the first inducible AMPs to be isolated, found in the hemolymph of

infected pupae of the moth *Hyalophora cecropia* (Lepidoptera) (Hultmark *et al.* 1980;

Steiner *et al.* 1981). The helical conformation of Cecropins is thought to promote their

interaction with negatively charged bacterial membranes, contributing to pore formation

and membrane destabilization, and resulting in the bacterial lysis (Steiner *et al.* 1988). *In*

vitro studies have shown that Cecropins have high efficacy against a large panel of gram-

negative bacteria at concentrations below the levels induced in insects upon infection

(10µM), as well as against some filamentous fungi (DeLucca *et al.* 1997; Ekengren and

Hultmark 1999; Ouyang *et al.* 2015). Heterologous expression of *Cecropin* in transgenic

rice has also been shown to confer resistance to the rice blast fungus *Magnaporthe oryzae*

(Coca *et al.* 2004). Cecropins *in vitro* do not display antimicrobial activities against Gram-

positive bacteria (Smeianov *et al.* 2000). However some studies have reported an activity

of Cecropins against tumor cells, bacterial biofilms and viruses (Chiou *et al.* 2002;

Suttmann *et al.* 2008; Deslouches and Di 2017; Kalsy *et al.* 2020).

AMP regulation and function has been extensively studied in the model insect *Drosophila*

melanogaster. The *Drosophila* genome encodes four *Cecropin* genes (*CecA1* and *A2*, *CecB*

and *CecC*) and two pseudogenes (*Cec-Ψ1* and *Cec2*) that are clustered at position 99E2 at

the tip of the right arm of the third chromosome (Kylsten *et al.* 1990; Samakovlis *et al.*

1990; Sackton *et al.* 2007). The *Cecropin* locus is adjacent to another gene named

Andropin, which encodes a related antibacterial peptide expressed in the ejaculatory duct

(Samakovlis *et al.* 1991). CecA1 and CecA2 are identical at the protein level, differing only

by a few silent mutations at the nucleotide level, suggesting that they emerged from a

recent duplication. The four *Drosophila Cecropin* genes are strongly induced in the fat

body and hemocytes upon systemic infection. During systemic infection, *Cecropin* genes

are regulated mostly by the Imd pathway while also receiving minor input from the Toll

pathway (Lemaitre *et al.* 1996; Hedengren *et al.* 1999). Functional studies analyzing the

role of Cecropins *in vivo* are scarce. Overexpression of *CecA* in an otherwise *Imd, Toll*

immune-deficient background failed to detect a clear protective effect for *CecA* against a

battery of pathogens (Tzou *et al.* 2002). Other studies using overexpression approaches

have pointed to a role for *CecA* in the regulation of the gut microbiota (Ryu *et al.* 2008).

Transgenic mosquitoes overexpressing both Cecropin and Defensin under the control of

the vitellogenin promoter displayed an increased resistance to *Pseudomonas aeruginosa*

infection, indicating that these AMPs could act cooperatively against this pathogenic

bacterium (Kokoza *et al.* 2010).

We have previously generated fly strains with deletions in 10 *Drosophila* AMP genes

including: *Defensin*, two *Diptericins* (*DptA* and *B*), *Drosocin*, four *Attacins* (*AttA, B, C,* and

D), *Metchnikowin*, and *Drosomycin* (Hanson *et al.* 2019a). This study revealed that AMPs

play an important role in defence against Gram-negative bacteria and also somewhat in

defence against fungi. In contrast, another family of host defence peptides with no overt

antimicrobial activity *in vitro*, the Bomanins, plays a major role in the elimination of

Gram-positive bacteria and fungi (Clemmons *et al.* 2015; Lindsay *et al.* 2018).

Importantly, Hanson et al. (Hanson *et al.* 2019a) revealed evidence for synergy and

additivity, but also remarkable specificity in the action of AMPs against certain pathogens.

However, this study did not address the function of the four Cecropins due to a failure to

generate a proper *Cecropin* locus deletion. In the present study, we have generated fly

lines carrying a small deletion that removes the four immune *Cecropin* genes, and by

using flies carrying this deletion alone or in combination with other AMP mutations, we

address for the first time the role of Cecropins in the systemic immune response.

MATERIALS AND METHODS

Fly stocks and genetics

The w^{1118} DrosDel isogenic (iso w^{1118}) wild type was used as the genetic background for

mutant isogenization, as described by Ferreira and colleagues (Ferreira *et al.* 2014). The

ΔCec^{A-C} mutation was generated using CRISPR with two gRNAs and a homology directed

repair vector by cloning 5' and 3' region-homologous arms into the pHD-DsRed vector,

and consequently ΔCec^{A-C} flies express DsRed in their eyes, ocelli and abdomen. The

ΔCec^{A-C} mutation was generated by Cas9 mediated injection into the iso Mtk^{R1}; Drs^{R1}

background. Following this, two rounds of backcrossing were performed to replace the

1st and 2nd chromosome with the *iso DrosDel* 1st and 2nd chromosome, and to recombine

the ΔCec^{A-C} mutation away from other mutations. The resulting stock is here called iso

ΔCec^{A-C}. Afterwards, the ΔCec^{A-C} mutation was recombined independently with Drs^{R1} and

$AttD^{SK1}$ on chromosome 3, and introgressed alongside the other AMP mutations on

chromosome 2 (Hanson *et al.* 2019a) to generate *$\Delta AMP14$* flies lacking the 14 classical

AMP genes from the *Defensin, Drosocin, Attacin, Diptericin, Metchnikowin, Drosomycin*,

and *Cecropin* gene families. The iso $Bom^{\Delta 55C}$ and iso $Relish^{E20}$ flies are the same as used in

Hanson et al. (Hanson *et al.* 2019a). Finally, the *$\Delta AMP10$* genotype is described in Hanson

et al. ; however we removed the aberrant *Cecropin* locus (Cec^{SK6}) detected in this line to

avoid any potential effects this locus could have on Cecropin-mediated resistance to

infection (see Hanson et al. correction notice (Hanson *et al.* 2019b)).

Microbial culture conditions

Bacteria were grown overnight on a shaking plate at 200 RPM in their respective growth

media and at their optimal temperature conditions. They were then pelleted by

centrifugation (4000 RPM) at 4°C. The bacterial pellets were diluted to the desired optical

density at 600nm (OD_{600}).

Pectobacterium carotovorum carotovorum 15 (*Ecc15*) and *Micrococcus luteus* were grown

in LB media at 29°C. *Escherichia coli* strain 1106, *Providencia burhodogranariea*,

Providencia rettgeri and *Providencia heimbachae* were grown in LB media at 37°C.

Enterococcus faecalis, Listeria monocytogenes and *Enterobacter cloacae* were cultured in

BHI media at 37°C. *Streptococcus pneumoniae* was grown as described by Krejčová and

colleagues (Krejčová *et al.* 2019). *Candida albicans* strain ATCC 2001 was cultured in YPG

media at 37°C. *Aspergillus fumigatus* was grown at 37°C on Malt Agar; spores were

collected in sterile PBS, pelleted by centrifugation and resuspended at the desired OD.

Beauveria bassiana strain R444 and *Metharizium rileyi* strain PHP1705 commercial

spores were produced by Andermatt Biocontrol (products: BB-PROTEC and NOMU-

PROTEC).

Infection experiments and survival

Systemic infections with *P. carotovorum carotovorum 15* (*Ecc15*) (Basset *et al.* 2000), *M.*

luteus, *E. coli* strain 1106, *P. burhodogranariea*, *P. rettgeri, P. heimbachae* (Galac and

Lazzaro 2011)*, E. faecalis*, *L. monocytogenes*, *E. cloacae* and *C. albicans* were performed

as followed: 3-5 day old adult females were pricked in the thorax with a 100μm thick

needle dipped into a concentrated pellet of bacteria at the desired OD_{600}. Infected flies

were then maintained at 25 or 29°C and survivals were tracked. Systemic infection with

S. pneumoniae (Krejčová *et al.* 2019) or *M. rileyi* was performed by injecting 50nL of a

concentrated pellet of bacteria or suspension of fungal spores using a nanoinjector and

glass capillary needles.

Natural infections with *B. bassiana* were performed by shaking anesthetized flies in a vial

with 200mg of spores. Flies were flipped into fresh vials one day after fungal inoculation.

A minimum of two replicate survival experiments were performed for each infection with

20 flies per vial on standard fly medium without yeast. Survival was scored daily.

Bacterial load of flies

Flies were infected (systemic infection) with bacteria at the desired OD_{600}. At the

indicated time post-infection, flies were anaesthetized using CO_2, surface sterilized by

washing briefly in 70% EtOH, and blotted dry. Pools of 5 flies were transferred in 200μL

of sterile PBS and macerated using a pestle. The homogenates were centrifuged at 8,000

RPM for 3 minutes. The supernatants were serially diluted and 7μL droplets (6

replicates) were inoculated on LB agar overnight at 29°C. Colony-forming units (CFUs)

were manually counted the following day.

Gene expression levels by RT-qPCR

Flies that either were unchallenged or were infected systemically by pricking in the

thorax with a needle dipped in a pellet of *Ecc15* or *M. luteus* (OD_{600}=200) were frozen at

-20°C 6h or 12h post-infection, respectively. Gene expression measurements were then

performed by RT-qPCR as previously described (Hanson *et al.* 2019a). Briefly, 5 whole

flies were homogenized and their RNA was extracted using TRIzol reagent and

resuspended in RNase-free water. Reverse transcription was carried out using

PrimeScript RT (TAKARA) with random hexamers and oligo dTs. Quantitative PCRs were

performed on a LightCycler 480 (Roche) using PowerUp SYBR Green Master Mix.

Cecropin A injection

Commercially available *Hyalophora cecropia* Cecropin A (Sigma-Aldrich) was diluted in

PBS to a concentration of 50μM. Fifty nL of Cecropin was injected into the thorax using a

nanoinjector and glass capillary. Flies were left to recover for 2 hours and then pricked

with the desired pathogen.

MALDI-TOF

Hemolymph samples were collected from either unchallenged flies, or flies pricked with

a 1:1 cocktail of *E. coli* and *M. luteus* (OD=200) in 0.1% TFA, as described previously

(Hanson *et al.* 2019a). Samples were then added to an acetonitrile universal matrix and

subjected to desorption and ionization. Representative spectra are shown. Immune

induced peaks were identified based on previous studies (Uttenweiler-Joseph *et al.* 1998;

Levy *et al.* 2004) to confirm the absence of AMP-associated peaks, and presence of

immune-induced peptides not affected by the various AMP mutations. Spectra were

visualized using mMass and figures were additionally prepared using Inkscape v0.92.

Statistical analysis

Survival analyses were performed using a Cox proportional hazards (CoxPH) multiple

comparison model, with Benjamini-Hochberg corrections for p-values, in R 3.6.3. Survival

curves included at least one survival experiment with 2 cohorts of 20 flies per treatment.

Statistics were represented using a Compact Letter Display (CLD) graphical technique:

groups were assigned the same letter if they were not significantly different (p>.05).

Quantitative PCR data were compared by one-way ANOVA with Holm-

Šídák multiple test correction in Prism R7. Bacterial load values were transformed as

$\log_{10}(value+1)$ to allow graphical representation of the absence (0) of CFUs. Bacterial load

data were compared by one-way ANOVA with Holm- Šídák multiple test correction in

Prism 7. Statistics were represented using a CLD graphical technique.

 RESULTS

Generation and characterization of *Cecropin* mutants

We generated a fly line lacking the four *Cecropin* genes, *CecA1, CecA2, CecB*, and *CecC,* which are clustered at 99E and are inducible during the systemic response. For this, we used the CRISPR/Cas9 editing method to generate a 6kb deletion (referred as ΔCec^{A-C}) that removes the four inducible *Cecropins* but leaves the neighboring *Andropin* gene intact (Fig. 1A). The ΔCec^{A-C} mutation was generated by Cas9 mediated injection in the *w, Drosdel* (referred to as w^{1118}) background. The background of the ΔCec^{A-C} mutation was then cleaned by 2 successive crosses to the w^{1118} *iso* background to remove potential off-target alterations. To confirm the absence of *Cecropin* genes in ΔCec^{A-C} flies, we performed RT-qPCR of *CecA*1 and *CecC* genes, which are located at the two extremities of the *Cecropin* locus. Expression of *CecA1* or *CecC*, readily observed in the wild type, was not detected in ΔCec^{A-C} flies (Fig. 1B-C).

We previously generated a fly line in the w^{1118} *iso* background here referred to as "$\Delta AMP10$", harbouring six mutations that remove ten antimicrobial peptide genes: *Defensin*, *Metchnikowin*, the four Attacins (*A/B/C/D*), *Drosomycin*, two Diptericins (*A/B*), and *Drosocin* (Hanson et al., 2019). We recombined the iso ΔCec^{A-C} mutation with the iso $\Delta AMP10$ mutations to generate an iso fly line lacking all 14 'classical' antimicrobial peptides (referred to as "$\Delta AMP14$"). MALDI-TOF and RT-qPCR analysis confirms the absence of these 14 antimicrobial peptides in $\Delta AMP10$ and $\Delta AMP14$ flies (Fig. 1B-C and Fig. S1). The ΔCec^{A-C}, $\Delta AMP10$ and $\Delta AMP14$ flies were healthy and showed no morphological defects. We also confirmed that the two central NF-κB signaling pathways, Toll and Imd, were functional, as demonstrated by measuring expression of genes characteristic of each of these pathways (Fig. 1D-F). Furthermore, MALDI-TOF proteomic

analysis of hemolymph from infected flies 24 hours post infection (hpi) reveals a wild type-like induction of peaks associated with other NF-κB effectors (e.g., Bomanins, Daishos, and Baramicin A) (Fig. S1). Collectively, our study indicates that we have generated a fly line lacking all the Drosophila 'classical' AMPs, and that deleting these AMPs does not impact the production of other NF-κB effectors.

Cecropins contribute to survival against certain Gram-negative bacterial infections

We used wild-type, ΔCec^{A-C}, $\Delta AMP10$ and $\Delta AMP14$ flies to explore the role that Cecropins play in defence against pathogens during systemic infection. By performing survival analyses with wild-type and ΔCec^{A-C} flies, we assessed if the absence of the four Cecropins is sufficient to cause an immune deficiency. Likewise, any difference in survival rates between $\Delta AMP10$ and $\Delta AMP14$ flies would suggest a contribution of Cecropins that is only apparent in the absence of other AMPs. We first focused our attention on Gram-negative bacterial infections, as cecropins were initially identified for their activity against this class of bacteria, and the four *Cecropin* genes are mostly regulated by the Imd pathway that responds to Gram-negative bacterial infection (Lemaitre and Hoffmann 2007). We challenged wild-type, ΔCec^{A-C}, $\Delta AMP10$ and $\Delta AMP14$ flies with six different Gram-negative bacterial species, using inoculation doses (given as OD_{600}) selected such that Imd deficient, iso Rel^{E20} mutant control flies were killed. Our survival experiments did not reveal an overt contribution of Cecropins to resistance against the Gram-negative bacteria *Escherichia coli, Pectobacterium carotovorum carotovorum (Ecc15), Providencia rettgeri* or *Providencia burhodogranariea* (Fig 2A-D). In all cases, ΔCec^{A-C} flies survived as well as wild-type flies, while $\Delta AMP10$ flies were as susceptible as $\Delta AMP14$. One exception

was found for *P. burhodogranariea* infection: death of *ΔAMP10* flies was delayed by one

day compared to *ΔAMP14* flies, suggesting a contribution of Cecropins in combatting this

bacterium early in infection.

Interestingly, we did identify a prominent role for Cecropins against two Gram-negative

bacterial strains: *Enterobacter cloacae and Providencia heimbachae*. Although *ΔCec^{A-C}* flies

survived *E. cloacae* infection like wild-type flies and *ΔAMP10* flies had only a slight

reduction in survival, *ΔAMP14* flies infected with *E. cloacae* instead behaved like *Rel^{E20}*

mutants lacking Imd signaling entirely (Fig. 3A). This result suggests that the presence of

the four *Cecropin* genes confers a mild protective effect against this bacterium in flies that

lack ten other AMP genes. A significant difference in CFUs between *ΔAMP10* and *ΔAMP14*

flies at 8h post-infection (hpi) confirmed a role of Cecropins in limiting the growth of *E.*

cloacae (Fig. 3B). We also observed a consistently higher bacterial load in *ΔCec^{A-C}* flies

compared to wild-type controls, though this was not significant (p = .18). Moreover,

ΔAMP14 and *Rel^{E20}* fly CFUs were similar, consistent with survival data showing complete

mortality of these genotypes within 24 hours. These results indicate that the knock-out

of the 'classical' AMPs in the *ΔAMP14* line fully explains the susceptibility of Imd pathway

mutants to *E. cloacae* infection. Next, we attempted to rescue the susceptibility of *ΔAMP10*

and *ΔAMP14* flies using commercially available Cecropin A. We injected 50 nL of 50μM

Hyalophora cecropia Cecropin A (Sigma-Aldrich) or PBS (control) two hours before

challenging flies with *E. cloacae*. Interestingly, when we injected Cecropin A two hours

prior to *E. cloacae* infection, *ΔAMP10* flies survived as well as wild-type flies previously

injected with PBS (Fig. 3C). This result suggests that priming the fly defence by increasing

circulating levels of Cecropin is sufficient to combat *E. cloacae* infection, even in flies

lacking a broad range of other AMPs. However we did not succeed in rescuing the

susceptibility of *ΔAMP14* flies using the same approach. This suggests the rescue effect

we observed using *ΔAMP10* flies relies on the total Cecropin levels, which includes both

endogenously produced Cecropin and the supplemental Cecropin A we injected.

Collectively, our *in vivo* analysis is consistent with previous *in vitro* studies that showed

commercial Cecropin A from *Hyalophora cecropia* has activity against *E. cloacae* (Brey

and Ashida 1993).

Similarly, we observed a contribution of Cecropins against the Gram-negative bacterium

P. heimbachae in flies lacking other AMPs (fig. 3D-F). While *ΔAMP10* flies were able to

survive this infection as wild type at OD_{600}=50, *ΔAMP14* flies again behaved like *Rel^E20*

mutants and suffered complete mortality (Fig. 3D); the *ΔCec^{A-C}* mutation alone did not

increase susceptibility. Bacterial load measurement performed on flies collected 24 hpi

revealed a contribution of Cecropins both in the presence and absence of other AMPs (Fig.

3E). We again injected commercial *H. cecropia* Cecropin A in an attempt to rescue the

susceptibility of *ΔAMP10* and *ΔAMP14* flies to *P. heimbachae* (Fig. 3F). Using this bacterial

infection model, previous injection of PBS increased the susceptibility of wild-type flies

to *P. heimbachae*. Strikingly, however, injection of Cecropin A prior to infection rescued

survival of *ΔAMP10* flies to a level comparable to uninjured wild-type flies. Moreover,

injection of *H. cecropia* Cecropin A improved the survival of *ΔAMP14* flies against *P.*

heimbachae, extending *ΔAMP14* lifespan compared to PBS controls.

In summary, our results reveal that Cecropins contribute to *Drosophila* host defence

against a subset of Gram-negative bacteria, and that this contribution is more readily

apparent when other AMPs are also lacking.

Cecropins are not involved in the resistance to Gram-positive bacteria

Previous work with the *ΔAMP10* flies did not reveal a role of *Drosophila* AMPs against Gram-positive bacteria, indicating instead that other immune effectors – notably the bomanins – play a predominant role against this class of microbes (Hanson *et al.* 2019a; Lin *et al.* 2020). Therefore, we were curious if the added loss of Cecropins would reveal a cryptic contribution of *Drosophila* AMPs to defence against Gram-positive bacteria. For this, we challenged wild-type, *ΔCec^{A-C}*, *ΔAMP10* and *ΔAMP14* flies with three Gram-positive bacteria: *E. faecalis*, *S. pneumoniae* and *L. monocytogenes* (Fig. 4A-C). *E. faecalis* and *S. pneumoniae* contain Lysine-type peptidoglycan that is known to predominantly activate the Toll pathway while *L. monocytogenes* has DAP-type peptidoglycan, and is known to activate both the Toll and Imd pathways (Leulier *et al.* 2003). In these experiments, we included iso *Bom^{Δ55C}* control flies, which lack ten *Bomanin* genes and are known to be susceptible to Gram-positive bacterial and fungal infections (Clemmons *et al.* 2015). Our survival experiments did not reveal a major role of Cecropins individually or alongside other AMPs in combating these Gram-positive bacterial species, but confirmed the importance of bomanins (Fig. 4A-C).

Cecropins can contribute to antifungal defence

While Cecropins were initially identified as antibacterial peptides, further *in vitro* studies have suggested an antifungal activity as well (Ekengren and Hultmark 1999; Andrä *et al.* 2001). We therefore investigated the contribution of Cecropins to resistance upon septic injury with any of four fungal species: the entomopathogenic fungi *Metarhizium rileyi* and *Beauvaria bassiana* R444, the opportunistic mould *Aspergillus fumigatus,* and the yeast *Candida albicans*. Survival analysis did not reveal a prominent role for AMPs or Cecropins against *M. rileyi*, *A. fumigatus* or *C. albicans,* as *ΔCec^{A-C}*, *ΔAMP10* and *ΔAMP14* flies

generally survived as the wild-type (Fig. 5A-C). However, *ΔAMP14* flies suffered greater

mortality to *B. bassiana* using a natural infection mode, compared to wild type flies (Fig.

5D). We additionally introduced *B. bassiana* spores directly into the hemolymph by septic

injury for more controlled fungal infection kinetics, and measured fungal load at 48h hpi

by RT-qPCR. Monitoring pathogen load revealed that in *ΔAMP14* flies, *B. bassiana* loads

were higher than levels found in wild type (p=.07) and *ΔAMP10* flies (Fig. 5E). Taken

together, these results show a contribution of Cecropins to defence against the

entomopathogenic fungus *B. bassiana*.

352   **DISCUSSION**
353

In this study, we generated flies lacking the four-immune inducible *Cecropin* genes to

address their function alone or in combination with other *AMP* gene mutations. *ΔCec*[A-C]

and *ΔAMP14* flies were viable, fertile and did not show any morphological defect.

Moreover, they display normal activation of the Imd and Toll pathways, suggesting that

the classical *Drosophila* AMPs do not contribute to immune signaling, in contrast to

mammalian AMPs (Mookherjee *et al.* 2020).

Our survival analyses reveal a role of Cecropins in the defence against certain Gram-

negative bacterial species. However, we could not identify a bacterial species or context

for which flies mutant for *Cecropin* genes alone succumb faster than wild-type. Studies of

other AMPs have revealed that certain AMPs exhibit a high degree of specificity in

determining host-pathogen interactions, as illustrated by the requirement of *Diptericin*

in defence against *P. rettgeri*, *Drosocin* in defence against *E. cloacae*, and the recently-

described *Daisho* and *Baramicin A* genes in defence against *Fusarium oxysporum* and

Beauveria bassiana fungi, respectively (Unckless *et al.* 2016; Hanson *et al.* 2019a, 2021;

Cohen *et al.* 2020). Further studies may reveal bacteria for which the presence of

Cecropins is essential for survival.

The most striking phenotype in the present study was that loss of Cecropins has a

marked effect on *E. cloacae* and *P. heimbachae* infection in flies also lacking other AMP

genes. As such, we reveal an important but cryptic contribution of Cecropins in defence

against these bacteria. Generation of flies lacking refined subsets of AMPs might narrow

down the specific groups of peptides key to defence against *E. cloacae* and *P. heimbachae*.

The enhanced growth of *E. cloacae* in AMP mutants that also lack Cecropins is a

particularly striking demonstration of their importance. In this infection model, the

presence of Cecropins dictates whether AMP mutant flies initially suppress bacterial

growth, or phenocopy *Rel^{E20}* flies deficient for Imd signaling. Cecropins are induced with

faster kinetics than most other AMPs, with a peak expression as early as 3hpi (Lemaitre

et al. 1997; De Gregorio *et al.* 2002). As *Cecropins* encode simple helical peptides that do

not require extensive post-translational modification, it is tempting to speculate that they

become functional more rapidly, and play an important role in combatting bacteria

specifically at this early phase of infection.

Our study also reveals that endogenous Cecropins can play a role in defence against

fungi, but not against Gram-positive bacteria. Thus, our *in vivo* study corroborates the

antifungal and antibacterial activities of Cecropins previously observed with *in vitro*

studies (Samakovlis *et al.* 1990; DeLucca *et al.* 1997; Ekengren and Hultmark 1999).

While Imd is crucial for the expression of the four Cecropins, the Cecropin response to

infection relies in part on Toll signaling (De Gregorio *et al.* 2002). As such, the

contribution of Cecropins to defence against fungi could help explain the regulation of the

Cecropin locus by both Toll and Imd pathways.

The observations that AMP genes are induced to an incredible extent, reach high

peptide concentrations in the hemolymph, and display *in vitro* microbicidal activity are

all consistent with their role as immune effectors. Use of both *ΔAMP10* and *ΔAMP14* flies

has confirmed the important contribution of AMPs to host defence against Gram-negative

bacteria and fungi, but not against Gram-positive bacteria. *Drosophila* AMPs also regulate

the gut microbiota downstream of the Imd pathway, a function consistent with their

bactericidal activity (Marra *et al.* 2021). However, recent studies have suggested that

AMP-like genes may play more subtle roles in other processes like memory formation

(Barajas-Azpeleta *et al.* 2018), an erect wing response upon infection (Hanson *et al.*

2021), tumor control (Parvy *et al.* 2019; Araki *et al.* 2019), or regulation of JNK signaling

in the salivary gland (Krautz *et al.* 2020). While we confirm a primary importance for

Cecropins and other AMPs in the systemic immune response, exploring the functions of

AMPs including Cecropins in non-canonical roles is an exciting future direction of

research.

Our study and others contribute to the rapid progress made towards understanding

the roles of *Drosophila* immune effectors. Research on the effector response has

stagnated for over a decade, but recent functional characterizations by loss of function of

key effectors (*Cecropins, Defensin, Attacins, Diptericins, Drosocin, Drosomycin,*

Metchnikowin, Bomanins, Daishos, and *Baramicin*) has greatly advanced our

understanding of the roles of these effectors (Lindsay *et al.* 2018; Hanson *et al.* 2019a;

Cohen *et al.* 2020; Huang *et al.* 2020). Most importantly, these studies correct the

assumptions of the previous "cocktail" model for AMP-pathogen interactions (Yan and

Hancock 2001; Lazzaro 2008; Zdybicka-Barabas *et al.* 2012; Rahnamaeian *et al.* 2016),

revealing some AMPs to be general effectors against most pathogens, while others act as

"silver bullets" specifically required for defence against certain pathogens. The

susceptibility of Toll and Imd pathway mutants to specific pathogens can now be directly linked to the susceptibility of mutants for immune effectors regulated by these pathways (Hanson and Lemaitre 2020). As new genetic techniques allow greater characterization of the roles of known immune effectors, many of them yet remain to be characterized, notably a number of short peptide genes highlighted by old and recent transcriptomic studies (Gregorio *et al.* 2001; Troha *et al.* 2018; Tattikota *et al.* 2020; Schlamp *et al.* 2021). However, we are likely still exploring inside the box when assuming a unique immune role for these peptides.

Our study also highlights the power of multiple mutation analysis, as the role of C*ecropins* would not have been uncovered *in vivo* by mutating individual genes. While we have begun exploring the combinatorial potential of AMPs in defence against infection, future studies will benefit from probing the interaction of immune effectors like AMPs with other mechanisms of host defence such as phagocytosis or melanization. With the advent of CRISPR/Cas9 technology and many recently described mutants, the interactions of AMPs in defence are just the tip of the iceberg in developing a global framework to understand the *Drosophila* immune response.

ACKNOWLEDGEMENTS

We are thankful to EPFL Proteomics Core Facility for help on the MALDI-TOF analysis. Brian Lazzaro generously provided *Providencia* species (Galac and Lazzaro 2011) and Adam Bajgar (Krejčová *et al.* 2019) provided *Streptococcus pneumoniae* species used in this study. We thank Hannah Westlake for editing of the manuscript. We are grateful for the SNF Sinergia grant CRSII5_186397 for the funding of this study.

REFERENCES

Andrä J., O. Berninghausen, and M. Leippe, 2001 Cecropins, antibacterial peptides from

insects and mammals, are potently fungicidal against Candida albicans. Med

Microbiol Immunol 189: 169–173. https://doi.org/10.1007/s430-001-8025-x

Araki M., M. Kurihara, S. Kinoshita, R. Awane, T. Sato, *et al.*, 2019 Anti-tumour effects of

antimicrobial peptides, components of the innate immune system, against

haematopoietic tumours in Drosophila mxc mutants. Dis Model Mech 12.

https://doi.org/10.1242/dmm.037721

Barajas-Azpeleta R., J. Wu, J. Gill, R. Welte, C. Seidel, *et al.*, 2018 Antimicrobial peptides

modulate long-term memory. PLoS Genet 14.

https://doi.org/10.1371/journal.pgen.1007440

Basset A., R. S. Khush, A. Braun, L. Gardan, F. Boccard, *et al.*, 2000 The phytopathogenic

bacteria Erwinia carotovora infects Drosophila and activates an immune response.

PNAS 97: 3376–3381. https://doi.org/10.1073/pnas.97.7.3376

Brey P. T., and M. Ashida, 1993 Role of the integument in insect immunity: Epicuticular

abrasion and induction of cecropin synthesis in cuticular epithelial cells. Proc. Natl.

Acad. Sci. USA 5.

Broekaert W. F., G. Terras', A. Cammue, and R. W. Osborn, 1995 Plant Defensins: Nove1

Antimicrobial Peptides as Components of the Host Defense System'. 108: 6.

Bulet P., C. Hetru, J.-L. Dimarcq, and D. Hoffmann, 1999 Antimicrobial peptides in insects;

structure and function. Developmental & Comparative Immunology 23: 329–344.

https://doi.org/10.1016/S0145-305X(99)00015-4

Chiou P. P., C.-M. Lin, L. Perez, and T. T. Chen, 2002 Effect of Cecropin B and a Synthetic

Analogue on Propagation of Fish Viruses In Vitro. Mar. Biotechnol. 4: 294–302.

https://doi.org/10.1007/s10126-002-0021-1

Clemmons A. W., S. A. Lindsay, and S. A. Wasserman, 2015 An effector Peptide family

required for Drosophila toll-mediated immunity. PLoS Pathog 11: e1004876.

https://doi.org/10.1371/journal.ppat.1004876

Coca M., C. Bortolotti, M. Rufat, G. Peñas, R. Eritja, *et al.*, 2004 Transgenic Rice Plants

Expressing the Antifungal AFP Protein from Aspergillus Giganteus Show Enhanced

Resistance to the Rice Blast Fungus Magnaporthe Grisea. Plant Mol Biol 54: 245–259.

https://doi.org/10.1023/B:PLAN.0000028791.34706.80

Cohen L. B., S. A. Lindsay, Y. Xu, S. J. H. Lin, and S. A. Wasserman, 2020 The Daisho Peptides

Mediate Drosophila Defense Against a Subset of Filamentous Fungi. Front. Immunol.

11. https://doi.org/10.3389/fimmu.2020.00009

De Gregorio E., P. T. Spellman, P. Tzou, G. M. Rubin, and B. Lemaitre, 2002 The Toll and Imd

pathways are the major regulators of the immune response in Drosophila. The EMBO

Journal 21: 2568–2579. https://doi.org/10.1093/emboj/21.11.2568

DeLucca A. J., J. M. Bland, T. J. Jacks, C. Grimm, T. E. Cleveland, *et al.*, 1997 Fungicidal

activity of cecropin A. Antimicrobial Agents and Chemotherapy 41: 481–483.

https://doi.org/10.1128/AAC.41.2.481

Deslouches B., and Y. P. Di, 2017 Antimicrobial peptides with selective antitumor

mechanisms: prospect for anticancer applications. Oncotarget 8: 46635–46651.

https://doi.org/10.18632/oncotarget.16743

Ekengren S., and D. Hultmark, 1999 Drosophila cecropin as an antifungal agent. Insect

Biochemistry and Molecular Biology 29: 965–972. https://doi.org/10.1016/S0965-

1748(99)00071-5

Ferreira Á. G., H. Naylor, S. S. Esteves, I. S. Pais, N. E. Martins, *et al.*, 2014 The Toll-Dorsal

Pathway Is Required for Resistance to Viral Oral Infection in Drosophila. PLOS

Pathogens 10: e1004507. https://doi.org/10.1371/journal.ppat.1004507

Galac M. R., and B. P. Lazzaro, 2011 Comparative pathology of bacteria in the genus

Providencia to a natural host, Drosophila melanogaster. Microbes and Infection 13:

673–683. https://doi.org/10.1016/j.micinf.2011.02.005

Ganz T., M. E. Selsted, D. Szklarek, S. S. Harwig, K. Daher, *et al.*, 1985 Defensins. Natural

peptide antibiotics of human neutrophils. J Clin Invest 76: 1427–1435.

https://doi.org/10.1172/JCI112120

Gregorio E. D., P. T. Spellman, G. M. Rubin, and B. Lemaitre, 2001 Genome-wide analysis of

the Drosophila immune response by using oligonucleotide microarrays. PNAS 98:

12590–12595. https://doi.org/10.1073/pnas.221458698

Hanson M. A., A. Dostálová, C. Ceroni, M. Poidevin, S. Kondo, *et al.*, 2019a Synergy and

remarkable specificity of antimicrobial peptides in vivo using a systematic knockout

approach. eLife 8. https://doi.org/10.7554/eLife.44341

Hanson M. A., A. Dostálová, C. Ceroni, M. Poidevin, S. Kondo, *et al.*, 2019b Correction:

Synergy and remarkable specificity of antimicrobial peptides in vivo using a

systematic knockout approach. eLife 8: e48778. https://doi.org/10.7554/eLife.48778

Hanson M. A., and B. Lemaitre, 2020 New insights on Drosophila antimicrobial peptide

function in host defense and beyond. Current Opinion in Immunology 62: 22–30.

https://doi.org/10.1016/j.coi.2019.11.008

Hanson M. A., L. B. Cohen, A. Marra, I. Iatsenko, S. A. Wasserman, *et al.*, 2021 The

Drosophila Baramicin polypeptide gene protects against fungal infection. bioRxiv

2020.11.23.394148. https://doi.org/10.1101/2020.11.23.394148

Hedengren M., BengtÅsling, M. S. Dushay, I. Ando, S. Ekengren, *et al.*, 1999 Relish, a Central

Factor in the Control of Humoral but Not Cellular Immunity in Drosophila. Molecular

Cell 4: 827–837. https://doi.org/10.1016/S1097-2765(00)80392-5

Huang J., Y. Lou, J. Liu, P. Bulet, R. Jiao, *et al.*, 2020 The BaramicinA gene is required at

several steps of the host defense against Enterococcus faecalis and Metarhizium

robertsii in a septic wound infection model in Drosophila melanogaster. bioRxiv

2020.11.23.394809. https://doi.org/10.1101/2020.11.23.394809

Hultmark D., H. Steiner, T. Rasmuson, and H. G. Boman, 1980 Insect Immunity. Purification

and Properties of Three Inducible Bactericidal Proteins from Hemolymph of

Immunized Pupae of Hyalophora cecropia. European Journal of Biochemistry 106: 7–

16. https://doi.org/10.1111/j.1432-1033.1980.tb05991.x

Imler J.-L., and P. Bulet, 2005 Antimicrobial Peptides in Drosophila: Structures,Activities and

Gene Regulation. Mechanisms of Epithelial Defense 86: 1–21.

https://doi.org/10.1159/000086648

Kalsy M., M. Tonk, M. Hardt, U. Dobrindt, A. Zdybicka-Barabas, *et al.*, 2020 The insect

antimicrobial peptide cecropin A disrupts uropathogenic Escherichia coli biofilms. npj

Biofilms and Microbiomes 6: 1–8. https://doi.org/10.1038/s41522-020-0116-3

Kokoza V., A. Ahmed, S. Woon Shin, N. Okafor, Z. Zou, *et al.*, 2010 Blocking of Plasmodium

transmission by cooperative action of Cecropin A and Defensin A in transgenic Aedes

aegypti mosquitoes. Proc Natl Acad Sci U S A 107: 8111–8116.

https://doi.org/10.1073/pnas.1003056107

Krautz R., D. Khalili, and U. Theopold, 2020 Tissue-autonomous immune response regulates

stress signaling during hypertrophy. Elife 9. https://doi.org/10.7554/eLife.64919

Krejčová G., A. Danielová, P. Nedbalová, M. Kazek, L. Strych, *et al.*, 2019 Drosophila

macrophages switch to aerobic glycolysis to mount effective antibacterial defence.

eLife 8. https://doi.org/10.7554/eLife.50414

Kylsten P., C. Samakovlis, and D. Hultmark, 1990 The cecropin locus in Drosophila; a

compact gene cluster involved in the response to infection. The EMBO Journal 9:

217–224. https://doi.org/10.1002/j.1460-2075.1990.tb08098.x

Lazzaro B. P., 2008 Natural selection on the Drosophila antimicrobial immune system. Curr

Opin Microbiol 11: 284–289. https://doi.org/10.1016/j.mib.2008.05.001

Lemaitre B., E. Nicolas, L. Michaut, J.-M. Reichhart, and J. A. Hoffmann, 1996 The

Dorsoventral Regulatory Gene Cassette spätzle/Toll/cactus Controls the Potent

Antifungal Response in Drosophila Adults. Cell 86: 973–983.

https://doi.org/10.1016/S0092-8674(00)80172-5

Lemaitre B., J.-M. Reichhart, and J. A. Hoffmann, 1997 Drosophila host defense: Differential

induction of antimicrobial peptide genes after infection by various classes of

microorganisms. PNAS 94: 14614–14619. https://doi.org/10.1073/pnas.94.26.14614

Lemaitre B., 2004 The road to Toll. Nature Reviews Immunology 4: 521–527.

https://doi.org/10.1038/nri1390

Lemaitre B., and J. Hoffmann, 2007 The Host Defense of Drosophila melanogaster. Annual

Review of Immunology 25: 697–743.

https://doi.org/10.1146/annurev.immunol.25.022106.141615

Leulier F., C. Parquet, S. Pili-Floury, J.-H. Ryu, M. Caroff, *et al.*, 2003 The Drosophila immune

system detects bacteria through specific peptidoglycan recognition. Nature

Immunology 4: 478–484. https://doi.org/10.1038/ni922

Levy F., P. Bulet, and L. Ehret-Sabatier, 2004 Proteomic Analysis of the Systemic Immune

Response of Drosophila*. Molecular & Cellular Proteomics 3: 156–166.

https://doi.org/10.1074/mcp.M300114-MCP200

Lin S. J. H., L. B. Cohen, and S. A. Wasserman, 2020 Effector specificity and function in

Drosophila innate immunity: Getting AMPed and dropping Boms. PLOS Pathogens

16: e1008480. https://doi.org/10.1371/journal.ppat.1008480

Lindsay S. A., S. J. H. Lin, and S. A. Wasserman, 2018 Short-Form Bomanins Mediate Humoral

Immunity in Drosophila. J Innate Immun 10: 306–314.

https://doi.org/10.1159/000489831

Marra A., M. A. Hanson, S. Kondo, B. Erkosar, and B. Lemaitre, 2021 Drosophila

antimicrobial peptides and lysozymes regulate gut microbiota composition and

abundance. bioRxiv 2021.03.19.436153. https://doi.org/10.1101/2021.03.19.436153

Mookherjee N., M. A. Anderson, H. P. Haagsman, and D. J. Davidson, 2020 Antimicrobial

host defence peptides: functions and clinical potential. Nat Rev Drug Discov 19: 311–

332. https://doi.org/10.1038/s41573-019-0058-8

Ouyang L., X. Xu, S. Freed, Y. Gao, J. Yu, *et al.*, 2015 Cecropins from Plutella xylostella and

Their Interaction with Metarhizium anisopliae. PLOS ONE 10: e0142451.

https://doi.org/10.1371/journal.pone.0142451

Parvy J.-P., Y. Yu, A. Dostalova, S. Kondo, A. Kurjan, *et al.*, 2019 The antimicrobial peptide

defensin cooperates with tumour necrosis factor to drive tumour cell death in

Drosophila. eLife 8. https://doi.org/10.7554/eLife.45061

Rahnamaeian M., M. Cytryńska, A. Zdybicka-Barabas, and A. Vilcinskas, 2016 The functional

interaction between abaecin and pore-forming peptides indicates a general

mechanism of antibacterial potentiation. Peptides 78: 17–23.

https://doi.org/10.1016/j.peptides.2016.01.016

Ryu J.-H., S.-H. Kim, H.-Y. Lee, J. Y. Bai, Y.-D. Nam, *et al.*, 2008 Innate immune homeostasis

by the homeobox gene caudal and commensal-gut mutualism in Drosophila. Science

319: 777–782. https://doi.org/10.1126/science.1149357

Sackton T. B., B. P. Lazzaro, T. A. Schlenke, J. D. Evans, D. Hultmark, *et al.*, 2007 Dynamic

evolution of the innate immune system in Drosophila. Nat Genet 39: 1461–1468.

https://doi.org/10.1038/ng.2007.60

Samakovlis C., D. A. Kimbrell, P. Kylsten, A. Engström, and D. Hultmark, 1990 The immune

response in Drosophila: pattern of cecropin expression and biological activity. The

EMBO Journal 9: 2969–2976. https://doi.org/10.1002/j.1460-2075.1990.tb07489.x

Samakovlis C., P. Kylsten, D. A. Kimbrell, A. Engström, and D. Hultmark, 1991 The andropin

gene and its product, a male-specific antibacterial peptide in Drosophila

melanogaster. The EMBO Journal 10: 163–169. https://doi.org/10.1002/j.1460-

2075.1991.tb07932.x

Schlamp F., S. Y. N. Delbare, A. M. Early, M. T. Wells, S. Basu, *et al.*, 2021 Dense time-course

gene expression profiling of the Drosophila melanogaster innate immune response.

BMC Genomics 22: 304. https://doi.org/10.1186/s12864-021-07593-3

Seo M.-D., H.-S. Won, J.-H. Kim, T. Mishig-Ochir, and B.-J. Lee, 2012 Antimicrobial Peptides

for Therapeutic Applications: A Review. Molecules 17: 12276–12286.

https://doi.org/10.3390/molecules171012276

Smeianov V., K. Scott, and G. Reid, 2000 Activity of cecropin P1 and FA-LL-37 against

urogenital microflora. Microbes Infect 2: 773–777. https://doi.org/10.1016/s1286-

4579(00)90359-9

Steiner H., D. Hultmark, A. Engström, H. Bennich, and H. G. Boman, 1981 Sequence and

specificity of two antibacterial proteins involved in insect immunity. Nature 292:

246–248. https://doi.org/10.1038/292246a0

Steiner H., D. Andreu, and R. B. Merrifield, 1988 Binding and action of cecropin and cecropin

analogues: Antibacterial peptides from insects. Biochimica et Biophysica Acta (BBA) -

Biomembranes 939: 260–266. https://doi.org/10.1016/0005-2736(88)90069-7

Suttmann H., M. Retz, F. Paulsen, J. Harder, U. Zwergel, *et al.*, 2008 Antimicrobial peptides

of the Cecropin-family show potent antitumor activity against bladder cancer cells.

BMC Urology 8: 5. https://doi.org/10.1186/1471-2490-8-5

Tattikota S. G., B. Cho, Y. Liu, Y. Hu, V. Barrera, *et al.*, 2020 A single-cell survey of Drosophila

blood. Elife 9. https://doi.org/10.7554/eLife.54818

Troha K., J. H. Im, J. Revah, B. P. Lazzaro, and N. Buchon, 2018 Comparative transcriptomics

reveals CrebA as a novel regulator of infection tolerance in D. melanogaster. PLOS

Pathogens 14: e1006847. https://doi.org/10.1371/journal.ppat.1006847

Tzou P., J.-M. Reichhart, and B. Lemaitre, 2002 Constitutive expression of a single

antimicrobial peptide can restore wild-type resistance to infection in

immunodeficient Drosophila mutants. Proc Natl Acad Sci U S A 99: 2152–2157.

https://doi.org/10.1073/pnas.042411999

Unckless R. L., V. M. Howick, and B. P. Lazzaro, 2016 Convergent Balancing Selection on an

Antimicrobial Peptide in Drosophila. Current Biology 26: 257–262.

https://doi.org/10.1016/j.cub.2015.11.063

Uttenweiler-Joseph S., M. Moniatte, M. Lagueux, A. V. Dorsselaer, J. A. Hoffmann, *et al.*,

1998 Differential display of peptides induced during the immune response of

Drosophila: A matrix-assisted laser desorption ionization time-of-flight mass

spectrometry study. PNAS 95: 11342–11347.

https://doi.org/10.1073/pnas.95.19.11342

Yan H., and R. E. W. Hancock, 2001 Synergistic Interactions between Mammalian

Antimicrobial Defense Peptides. Antimicrobial Agents and Chemotherapy 45: 1558–

1560. https://doi.org/10.1128/AAC.45.5.1558-1560.2001

Zdybicka-Barabas A., P. Mak, A. Klys, K. Skrzypiec, E. Mendyk, *et al.*, 2012 Synergistic action

of Galleria mellonella anionic peptide 2 and lysozyme against Gram-negative

bacteria. Biochimica et Biophysica Acta (BBA) - Biomembranes 1818: 2623–2635.

https://doi.org/10.1016/j.bbamem.2012.06.008

Figure legends

642
Figure 1 : Description and validation of *ΔCecA-C*, *ΔAMP10* and *ΔAMP14* mutants.

(**A**) Schema of the *Cecropin* locus chromosomal deletion removing *CecA1* and *A2, CecB* and

CecC, plus 2 pseudo genes, *Cec2* and *Cec-Ψ1* clustered at position 99E2 (III) (**B,C**) RT-

qPCR of *CecA* (B) and *CecC* (C) expression in w^{1118}, *ΔCec^{A-C}*, *ΔAMP10* and *ΔAMP14* flies 6

hours post *Ecc15* infection. (**D-F**) The Imd (D-E) and Toll (F) pathways are functional in

ΔCec^{A-C}, ΔAMP10 and *ΔAMP14* flies after challenge as revealed by expression of target

genes upon septic injury with *Ecc15* or *M. luteus. PGRP-LB* and *Pirk* were used as readouts

for Imd pathway and *Bomanin (BomBc3)* for Toll pathway. Expression was normalized

with w^{1118} UC to a value of 1. Statistical significance levels were represented as: * = P≤0.05

; ** = P≤0.01 ; *** = P≤0.001 ; **** = P≤0.0001.

Figure 2 : Cecropins are not involved in resistance to a broad spectrum of Gram-

negative bacteria.

w^{1118} were used as wild-type flies and *Rel^{E20}* as susceptible flies lacking the Imd pathway

for all survival experiments to Gram-negative bacterial infection. Female w^{1118}, *ΔCec^{A-C}*,

ΔAMP10, ΔAMP14 and *Rel^{E20}* flies were pricked in the thorax with an inoculum of (**A**) *P.*

rettgeri, (**B**) *Ecc15,* (**C**) *E. coli* or (**D**) *P. burhodogranariea.* Cecropins were not critically

involved in combating infection with any bacteria presented here as *ΔCec^{A-C}* survived as

well as w^{1118} flies, while *ΔAMP10* and *ΔAMP14* mutant flies died as fast as *Rel^{E20}* mutants.

Bacterial concentrations are indicated in the figure.

Figure 3 : Cecropins are essential in the absence of other AMPs to resist *E. cloacae*

and *P. heimbachae* infection.

(**A**) Survival experiments upon infection with *E. cloacae* reveal that AMP deficient flies

having Cecropins (*ΔAMP10*) are significantly more resistant than those without

Cecropins (*ΔAMP14*). (**B**) Bacterial loads (CFU counts) of w^{1118}, *ΔCec^{A-C}, ΔAMP10, ΔAMP14*

and *Rel^{E20}* flies 8 hours post infection reveal a significant role for Cecropins in clearing

and controlling *E. cloacae.* (**C**) Commercial Cecropin A injection (50 nL at 50 μM) 2 hours

prior to *E. cloacae* infection increases the resistance of *ΔAMP10* mutant flies. However,

CecA injection did not rescue the susceptibility of *ΔAMP14* flies to *E. cloacae.* (**D-F**)

Survival analysis (D) bacterial load measurements 24 hours post infection (E) and

Cecropin supplementation experiments (F) in w^{1118}, *ΔCec^{A-C}, ΔAMP10, ΔAMP14* and *Rel^{E20}*

flies upon infection with *P. heimbachae* (as described for A-C).

Figure 4 : Cecropins are not involved in resistance to Gram-positive bacteria

w^{1118} were used as wild-type flies and $Bom^{\Delta55C}$ as susceptible flies lacking 10 *Bomanin*

genes for all survival experiments to Gram-positive bacterial infection. w^{1118}, *ΔCec^{A-C}*,

ΔAMP10, ΔAMP14 and $Bom^{\Delta55C}$ flies were pricked in the thorax with an inoculum of (**A**)

E. faecalis, (**B**) *S. pneumoniae* and (**C**) *L. monocytogenes*. Cecropins were not involved in

combating infection of 3 bacterial species: *ΔCec^{A-C}, ΔAMP10* and *ΔAMP14* flies survived as

well as w^{1118} flies. Bacterial concentrations are indicated in the figure.

Figure 5: Cecropins contribute to antifungal defence against *B. bassiana*

(**A-C**) w^{1118} were used as wild-type flies and Bom^{D55C} as susceptible flies lacking 10

Bomanin genes for all survival experiments to fungal infections. Cecropins were not

involved in combating infection of (**A**) *M. rileyi,* (**B**) *A. fumigatus* and (**C**) *C. albicans*, as

w^{1118}, *ΔCec^{A-C}, ΔAMP10* and *ΔAMP14* flies survived as well as w^{1118} flies. (**D**) Survival upon

(**D**) natural infection with *B. bassiana* reveals a significant increase in resistance of

ΔAMP10 flies compared to *ΔAMP14* flies, suggesting an important role for cecropins in

combating *B. bassiana*. (**E**) *B. bassiana* load (measured by *B. bassiana* 18S expression

related to *D. melanogaster RpL32*) is increased (p = .07) in *ΔAMP14* flies compared to

w^{1118}, *ΔCec^{A-C}* and *ΔAMP10* flies 48 hours post septic infection. Fungal concentrations are

indicated in the figure.

Figure S1: *ΔAMP14* flies retain the induction of non-AMP Immune Molecules (IMs)

upon systemic infection. Pools of ~40 mixed male and female flies were pricked with a

1:1 mix of OD_{600}=200 *E. coli* and *M. luteus* bacteria, and ~40nL of total hemolymph was

extracted by nanoject 24hpi, and directly placed in 5uL of 0.1% TFA. Peptide samples

were run on an acetonitrile matrix as described previously (Hanson *et al.* 2019a).

Hemolymph from infected w^{1118} wild-type in blue (top) and *ΔAMP14* flies in pink

(bottom) are shown. Peaks corresponding to AMP products in the wild-type are labelled

in pink, including IM7 whose sequence is unknown, but is absent in *ΔAMP14* flies. Of note,

a novel immune-induced peak was observed at ~3027 Da in *AMP* mutants. This peak

corresponds to an unknown peptide that is masked by Metchnikowin (~3026 Da) in the

wild-type. Reflectron analysis confirms a different spectrum profile for this peak.

Drosophila IMs are described in (Levy *et al.* 2004) and (Uttenweiler-Joseph *et al.* 1998)

Figure 1

Figure 2

Figure 3

Figure 4

A

B

C

Figure 5

A

M. rileyi OD$_{600}$= 4

Survival (%)

Time (days)

- — w^{1118} A
- — ΔCec^{A-C} A
- — $\Delta AMP10$ A
- — $\Delta AMP14$ A
- ⋯ $Bom\Delta^{55C}$ B

B

A. fumigatus OD$_{600}$= 200

Survival (%)

Time (days)

- — w^{1118} A
- — ΔCec^{A-C} A
- — $\Delta AMP10$ A
- — $\Delta AMP14$ A
- ⋯ $Bom\Delta^{55C}$ B

C

C. albicans OD$_{600}$= 200

Survival (%)

Time (days)

- — w^{1118} A
- — ΔCec^{A-C} A
- — $\Delta AMP10$ AB
- — $\Delta AMP14$ B
- ⋯ $Bom\Delta^{55C}$ C

D

B. bassiana 200mg

Survival (%)

Time (days)

- — w^{1118} A
- — ΔCec^{A-C} A
- — $\Delta AMP10$ A
- — $\Delta AMP14$ B
- ⋯ $Bom\Delta^{55C}$ C

E

B. bassiana OD$_{600}$=5 48 hpi

B. bassiana 18S related to D. melanogaster RpL32

w^{1118} ΔCec^{A-C} $\Delta AMP10$ $\Delta AMP14$ $Bom\Delta55C$