1 **Extracting socio-spatial networks from photo-ID data using multilevel**

2 **multinomial models**

4 Tyler R. Bonnell [a,b,c], Robert Michaud [d], Angélique Dupuch [a,b], Véronique Lesage[e], Clément

5 Chion [a,b]

[a] Department of Natural Sciences, Université du Québec en Outaouais, Gatineau, Québec,

Canada

[b] Institut des Sciences de la Forêt Tempérée, Université du Québec en Outaouais, Ripon,

Québec, Canada

[c] Department of Psychology, University of Lethbridge, 4401 University Drive, Alberta T1K

3M4, Canada

[d] Groupe de Recherche et d'Éducation sur les Mammifères Marins (GREMM), 870 avenue

Salaberry, Bureau R24, Québec, QC G1R 2T9, Canada

[e] Maurice Lamontagne Institute, Fisheries and Oceans Canada, P.O. Box 1000, 850 route de

la Mer, Mont-Joli, QC G5H 3Z4, Canada

Abstract:

Photo identification of individuals within a population is a common data source that is becoming more common given technological advances and the use of computer vision and machine learning to re-identify individuals. These data are collected through hand-held cameras, drones, and camera traps, and often come with biases in terms of sampling effort and distribution. In spite of these biases, a common goal of collecting these datasets is to better understand the habitat use pattern of individuals and populations. Here, we examine the potential for multilevel multinomial models to generate socio-spatial networks that capture the similarities in individual users across the spatial distribution of a species. We use this approach with 18 years of photo-ID data to better understand population structuring of beluga whales in the St. Lawrence River. We show using permuted and simulated data that this approach can identify community network structures within populations in a way that accounts for biases in collections methods. Applying this method to the entire 18 years dataset for SLE beluga, we found three spatially distinct communities. These results suggest that within the population's summer range individuals are moving within restricted areas (i.e., home ranges), and have implications for the estimated impacts of localized anthropogenic stressors, such as chemical pollution or acoustic disturbances on animal populations. We conclude that multilevel multinomial models can be effective at estimating socio-spatial networks that describe community structuring within wildlife populations.

Keywords: Multinomial Model, Beluga, Photo ID, Socio-Spatial Network, Bayesian Network, Community Detection

1. Introduction

An understanding of the spatial and temporal distribution of a species of concern is of central importance to conservation and management (Evans & Hammond, 2004). Increasingly, photo and video are being used to monitor individuals within populations (hereafter photo-ID data), providing a view of within population social mixing and habitat use (Koivuniemi et al., 2016). The increased use of machine learning to identify individuals from these data streams has greatly facilitated the use of these photo-ID data (Schneider et al., 2019). These individual identifications have facilitated the use of novel statistical and computational methods to quantify within population structures, such as social network analysis (Perryman et al., 2019; Schilds et al., 2019).

It is often the case, however, that efforts when collecting photo-ID are not evenly distributed. This differentiation in effort can heavily bias estimates of both habitat usage and population distribution estimates (Hupman et al., 2018). Here we propose the novel use of multilevel multinomial models to account for these biases and to estimate socio-spatial structures within populations.

The existence of social structuring within populations, such as communities, can have important ecological and management implications. If a population as a whole can be considered as highly mixed, i.e., with individuals showing no strong patterns of home range use or sub-structuring within the large population, then all individuals are equally likely to feel the impacts of local changes in the environment. In contrast, if the population cannot be considered to be highly mixed, and shows strong sub-structuring and site-fidelity patterns within the larger population, local stressors might have a disproportionate impact on subsections of the population. For example, if noise pollution increased in only one sector, in a highly mixed population all individuals would be lightly impacted, but in a structured population a subset of the population would be highly impacted. These differences in spatial

structuring of populations can lead to biased estimation of the likelihood and magnitude of

impacts from local stressors both at the individual and population levels (DeFur et al., 2007).

The multilevel multinomial modeling approach does not have a large body of

literature to draw on for use with photo-ID data. However, it does have some unique

advantages (Koster & McElreath, 2017). For instance, if sampling effort is biased in different

regions, the mean probability of being seen within highly sampled regions will be biased

upwards. By taking advantage, however, of the multilevel structure of the model it is possible

to extract individual deviations in the probability of being seen within a particular region.

Decisively, these individual level deviations from the mean probability are not biased by

changes in the sampling effort. That is the mean probability will increase with sampling

effort, but the relative difference between individuals within the sector will not. High users of

a particular region of a habitat will consistently be higher compared to low users of that

habitat, and this difference between high and low users will not be biased by sampling effort.

Furthermore, by comparing the individual differences between regions it is possible to see if

the high/low users of one region are similarly the high/low users of another region. The

similarity, or dissimilarity, between regions can then provide information about which

regions share similar user profiles. We suggest that by using the correlations between these

individual level deviations in high/low users between regions it is possible to generate socio-

spatial networks and identify social structuring within the population. In particular it can help

to identify spatial communities, i.e., a set of regions that share similar usage patterns and that

differ from other regions.

To evaluate the use of multilevel multinomial models to identify socio-spatial

structuring within a population, we make use of a long term photo-ID dataset of beluga

whales in the St. Lawrence Estuary, Canada. This population has undergone a drastic

decrease from around 10,000 in the late 1800s to less than 1,000 today, and is currently

considered as endangered in Canada according to the Species at Risk Act (COSEWIC 2014;

Fisheries and Oceans Canada, 2012; Mosnier et al., 2015). The population is part of a larger

study on the mitigation of noise pollution due to marine traffic (Chion et al., 2017; Lesage et

al., 2014; McQuinn et al., 2011; Parrott et al., 2011).

In this paper we first evaluate the performance of the multilevel multinomial models

using simulated data, testing if the method correctly estimates no community structuring

when none is present, and identifies the correct structure when it is present. In both cases we

use the 18 year beluga photo-ID dataset, randomly permuting uniquely identified individuals

to generate unstructured datasets, and randomly placing individuals within pre-specified

communities to generate structured datasets. We then apply the method to the observed data

and quantify community structures within the population's summer range in the St. Lawrence

Estuary. Finally, we discuss some potential extensions to the multilevel multinomial

modeling approach.

2.   Material and Methods

2.1 Data

Photo-ID data were collected using a handheld camera onboard a boat that was able to

navigate near to beluga, hereafter referred to as an encounter. Once near beluga a photo ID

protocol was then followed to generate photographs used to attempt to identify individuals.

Due to the logistical difficulty of covering a large body of water, the sampling effort was

unequally distributed across 14 sectors within the St. Lawrence Estuary (Fig. 1). The photo-

ID dataset used in this study was collected from 1989-2007 and are part of an ongoing

project. The data is stored in a database that facilitates the identification and association

between photos to help track the individual identification process. This resulted in a dataset

of 7,525 individual encounters where the individual was successfully identified by photo

sampling (hereafter referred to as a photo-ID), and where a GPS point was taken and the

sector recorded. This resulted in 821 unique individuals being successfully identified, with a

mean number of photo-IDs per individual of 9 (min = 1, max = 90) (Fig. 1).

Figure 1: Spatial distribution of photo-ID data (red points) within the St. Lawrence Estuary,

Quebec, Canada (red square in the inset map). The 14 sectors are outlined and labeled in

white, and covers the summer habitat of this beluga population.

2.2 Statistical Analysis

Our aim was to use photo-ID data to estimate the probability of seeing an individual

in each delineated sector of the St. Lawrence Estuary, and to use these individual

probabilities to estimate socio-spatial structures within the beluga population (Fig. 1). To

accomplish this aim we used a multilevel multinomial model, where the dependent variable

was the number of times an individual was captured photographically (i.e., photo-identified)

in each sector. The use of a multilevel model structure allows for the estimation of both the

mean probability of photo-identifying an individual in each sector, and the individual level

differences in this probability by using individual ID as a random intercept. If we take, as an

example, a case where there is only two sectors, then the log-odds of finding individual i in a

sector other than the reference sector can be modeled using a multilevel multinomial

following (Koster & McElreath, 2017) as:

$$log\left(\frac{p_{1,i}}{p_{r,i}}\right) = \mu_1 + v_{1i}$$
$$log\left(\frac{p_{2,i}}{p_{r,i}}\right) = \mu_2 + v_{2i}$$

Where $p_{1,i}$ is the probability of seeing beluga i in sector 1, whereas $p_{r,i}$ is the

probability of seeing beluga i in the reference sector. The μ_1 and μ_2 are the intercepts, i.e., the

mean probability of seeing a beluga in sectors 1 and 2. This mean probability represents

preference/avoidance of the specified sector, however, it is very likely to be biased due to

variation in sampling effort. Finally, the v_{1i} and v_{2i} are the estimated individual differences

(i.e., random intercepts) from the mean probability of capture in sectors 1 and 2, respectively.

These individual differences from the mean probability capture if an individual beluga is a

high/low user of that sector, and is not biased by variation in sampling. It is then possible to

model the covariance of the individual differences between two sectors using a multivariate

normal distribution:

$$\begin{bmatrix} v_{1i} \\ v_{2i} \end{bmatrix} \sim Multi\ Normal(0, \Omega_v)$$

$$\Omega_v = \begin{bmatrix} \sigma_{1,1} & \sigma_{1,2} \\ \sigma_{2,1} & \sigma_{2,2} \end{bmatrix}$$

This multivariate normal distribution has a mean of 0 and a covariance matrix Ω_v.

Here the diagonal entries in the covariance matrix (e.g., $\sigma_{1,1}$) represent the magnitude of

individual differences within a sector, i.e., are their high and low users in a sector or are all

individuals equally likely to be seen? The off-diagonal entries (e.g., $\sigma_{2,1}$) are the covariance

estimates between sectors, i.e., do sectors share similar high and low users? By estimating the

correlation of individual differences between sectors, this multilevel modeling approach

quantifies how much information individual differences in one sector can provide about

another sector. Positive correlations suggest that the high/low users in one sector are similarly

high/low users in another sector, while negative correlations suggest high/low users in one

sector are the low/high users in another sector.

This model can be fit using brms in the R environment using a multivariate syntax:

bf(y | trials(n) ~ 1 + (1|q|ID)) + multinomial(). Here, y is a set of column vectors where each

column is a sector and each row is an individual. The values in this column vector indicate

how many times each individual was seen in each sector. The n is the total number of times

an individual was captured, and q is an arbitrary character choice that allows correlations

between the estimates of random intercepts for each sector (Bürkner, 2017).

2.2.1 Addition of a common reference sector

Tests using simulated data suggested that adding a preset reference category (i.e., a

fifteenth sector used as a reference sector) to the data was required to estimate the correlation

between sectors (Fig. S1). To create this reference category, we tally up the observations for

each individual and add that number of observations to the new reference sector. This

essentially sets the probability of a capture in the reference sector to 0.5 for all individuals,

i.e., equal to the probability of being captured outside of this reference sector. This ensures

that all individuals have the same baseline probability in the reference category, and as the

parameters in the multinomial model measure deviations away from the reference sector, we

gain better estimates of the relative deviations between individuals (Fig. S1).

2.2.2 Dealing with biases in photo-ID datasets

This multilevel multinomial approach accounts for repeated sampling of individuals,

and provides an estimate of whether some individuals are found more or less often than the

mean probability of captures in each sector. We are particularly interested in the estimates of

individual differences from the mean probability of capture (i.e., the random intercepts) as

these estimates are not impacted by bias in sampling effort among sectors. This is not the

case for estimates of the mean probability of capture for each sector, which are expected to

increase in highly sampled sectors. For example, the Saguenay River is over-sampled

compared to the other sectors (SAG in Fig. 1), increasing the mean probability of capturing

individuals in that sector. However, over-sampling should not affect the individual

differences in the probability of being captured, i.e., all individuals' chances of being captured

go up or down equally.

Similarly, potential biases due to ease of recognition, e.g., some beluga or age classes

might have more distinctive markings, are minimized using a multilevel multinomial

approach where the differences in the probability of being seen between sectors is the main

focus. For example, if juveniles are less likely to be successfully identified by photo, then

they might have a reduced number of photo-IDs compared to other age classes, but the

difference in distribution of these fewer photo-IDs across sectors will not be impacted. For

example, if both an adult and juvenile spend twice as much time in the SAG sector compared

to all other sectors, you might expect a photo-ID distribution (Sag:not-sag) of 10:5 and 2:1,

respectively. In both cases the probability of being captured in the SAG sector is twice that of

the remaining sector. Due to the adaptive partial pooling properties of multilevel models,

individuals with few photo-IDs will, however, be less likely to show differences to the mean

probability, i.e., they contain less information. This means that if an age class has very little

chance of being identified by photo-ID, they will likely contribute less to the estimated socio-

spatial structures estimated by the multilevel multinomial approach.

By using a multilevel modeling approach we also reduce the chance of false positives

when making comparisons between many different individuals in many different sectors (i.e.,

problem of multiple comparisons). For example, if we were to estimate the differences in the

probability of each sector separately for each individual, the risk of false positives would be

increased. By using a multilevel approach to estimating the differences we can make effective

use of partial pooling of information to reduce extreme values, especially where the number

of photo IDs is not equal between individuals. Furthermore, by running this in a Bayesian

framework we are able to place priors on the individual differences within sectors that start

the model assuming that there are no differences between individuals in their use of each

sector, e.g., student_t(3,0,1).

2.3 Social Network Analysis

Social networks are often used when visualizing and quantifying social structures

within populations, with individuals often represented as nodes and their interactions as edges

between these nodes. In our case, we use sectors as nodes, and the similarities in user profiles

between sectors as edges. The correlations between sectors estimated from the multilevel

multinomial model can be used to create a network where the posterior predictions of each

correlation parameter corresponds to an edge weight in the network. In this way, each edge

has a posterior distribution and can be used to create many networks from which a

distribution of network metrics can be generated, e.g., the distribution of node strength values

can be calculated for each sector. The advantage of having distributions of network measures

is that the measures can be readily compared, e.g., does one sector have a higher node

strength than another? It is also possible to use the distribution of edge weights, and a chosen

threshold (e.g., 95% credible interval), to highlight only the edges where the sign of the

correlation is known with a particular range of certainty. In this paper, we used this latter

approach to generate a signed network (i.e., a network with positive and negative edges) and

use a simple signed-edge rule to define communities: where a distinct community is a set of

nodes that share positive edges but no negative edges. We also made use of signed

blockmodeling, an algorithm that can also be used to identify blocks of nodes that maximize

within block positive edges and minimize within block negative edges (Doreian & Mrvar,

2015). While the signed-edge rule generally provides relatively intuitive results with simple

networks, using the signed blockmodeling is likely to be particularly advantageous when

dealing with large networks.

2.4 Testing data

To assess the accuracy of the multilevel multinomial modelling approach, we generated

test datasets from the observed photo-ID data. We ensured that the test datasets contained the

same number of unique individuals, distribution of sightings (i.e., some individuals are seen

more than others), and overall number of photo-IDs as the observed dataset. We, however,

varied the spatial mixing of the test datasets. To test if the proposed method correctly

detected no pattern when none existed, we created a completely random test dataset by

permuting the sector associated with each photo-ID in the observed dataset. The expected

result was to find no correlations between sectors given that the sectors for each photo-ID had

been randomly permuted. To then test whether the proposed method could also correctly

identify patterns when a known pattern existed, we generated a structured test dataset by

randomly assigning each individual to four equally populated communities with the

following and hypothetical home range of adjacent sectors: community 1-BSM, SAG, CTN,

community 2- CTN, CTO, AMN, community 3- AVO, AVS, AVN, and community 4-AME,

CTS, CTE. Following this, we altered the sector of where the individual photo-IDs were

taken so as to fall within sectors associated with an individual's community, i.e., one of their

home range sectors. We did this by choosing a sector for each photo-ID based on the

individual's assigned community 80% of the time; a random sector was chosen for the other

20% of the time, introducing noise in the assignment of sectors. We then tested whether the

model correctly identified the correlations between sectors that defined the home range of

each of the communities.

3. Results

3.1 Testing data

When the multinomial multilevel model was fit to the data with sectors randomly

permuted between all photo-IDs, the model found no evidence for positive/negative

correlations between sectors (Fig. 2a). Similarly, when we simulated data with some known

structure, i.e., when we artificially created spatially distinct communities, we found that the

model accurately estimated the correlations between sectors that defined these artificial

communities (Fig. 2b). The simple signed-edge rule and blockmodelling algorithm applied to

the simulated datasets both revealed the four artificially generated communities, though the

blockmodeling algorithm had difficulty with the multi-membership node as it could not

assign a node to two blocks (i.e., the CTN node that was shared between communities 1 and

2).

Figure 2: Similarity and dissimilarity between sectors in the simulated datasets: a) randomly

permuted data, where there are no spatial communities, and b) structured data, where there

are four distinct communities. In b) the simulated communities are represented by color codes

for each of their sectors (Note: CTN is part of the orange and yellow communities). The

green edges between two sectors signify that the sectors share high/low users, while red

edges signify that they have dissimilar high/low users. The lack of an edge signifies that the

high/low users of one sector does not provide information about the high/low users of other

sectors. Nodes represent sectors, and are coloured based on the communities imposed when

simulating the data.

 3.2 Observed data

The results from our multilevel multinomial model found that within sectors there

were consistent individual differences in how likely it was to see individual beluga (Table 1).

That is, within sectors, there were some beluga that used the sector heavily, while others did

not. The model also found that between sectors these individual differences were correlated

(Table S1). These correlations quantify the magnitude of similarity/dissimilarity between

sectors in terms of which beluga are using those sectors heavily or rarely. If we take two

sectors as examples, e.g., the SAG and CTE sectors, representing, respectively, a tributary to

the St. Lawrence Estuary and a sector on the opposite side close to the South shore of the

Estuary, and we look at the top 10 estimated high users (i.e., relatively high probability of

being found there) within the SAG, we find that they are found to be low users in the CTE

sector (see blue dots in Fig 3 a) and b)).

296
Figure 3: Estimate of the relative use (i.e., deviation from mean use) for each individual

within the SAG (a) and CTE sectors (b) of the St. Lawrence beluga summer habitat. The

values are deviations from the mean probability of observing individuals within a sector and

are on the logit scale. The red dashed line represents the mean use, black points represent the

estimated deviation from the mean, while the horizontal grey lines represent the 95% credible

interval. To highlight how correlations are estimated between sectors, the estimated top 10

users of the SAG sector are represented by blue dots (panel a), and those same individuals are

also highlighted in blue in the CTE sector (panel b).

The use of a multilevel model also allowed us to estimate the magnitude of individual

differences in each sector, i.e., the extent to which there are high/low beluga users in a sector.

Our model found that the CTN sector showed very little individual differences in use (Table

1, i.e., low "sd" value) compared to other sectors, suggesting very little differences in high

and low users of that sector. While the BSM sector showed large individual differences, with

some very high/low users of that sector (Table 1).

Table 1: Parameter estimates from the multilevel multinomial model predicting the

probability of capturing a photo-ID by sector. Estimated magnitudes of individual differences

(sd) are presented for each sector. Higher 'sd' estimates indicate more individual differences

in individual use of that sector, whereas lower estimates indicate individuals are using the

sector at very similar levels. To facilitate interpretation we have ordered the table by lowest

to highest estimates of individual differences. As the number of parameters in the model is

large, the overall mean by sector, and estimated correlations between individual differences,

are presented in the supplementary section (Table S1).

Parameter	Estimate	SD	l-95% CI	u-95% CI
sd(mu_CTN)	0.33	0.03	0.27	0.39
sd(mu_AMO)	0.65	0.38	0.05	1.41
sd(mu_AMN)	0.76	0.29	0.14	1.31

sd(mu_AVO)	0.78	0.05	0.68	0.88
sd(mu_CTE)	0.78	0.05	0.68	0.89
sd(mu_AVS)	0.83	0.07	0.70	0.97
sd(mu_AVE)	1.16	0.20	0.78	1.57
sd(muy_SAG)	1.16	0.08	1.01	1.31
sd(mu_CTS)	1.18	0.11	0.97	1.40
sd(mu_CTO)	1.27	0.14	1.00	1.55
sd(mu_AVN)	1.30	0.17	0.98	1.65
sd(mu_BSM)	1.30	0.11	1.09	1.52
sd(mu_AMS)	1.53	0.22	1.10	1.98
sd(mu_AME)	2.00	0.19	1.64	2.38

Using the between sector correlations to generate a signed network overlaid on top of

the sectors in the St. Lawrence, suggests, for example, that individuals that are seen in the

SAG sector a lot, are also seen in the BSM and CTN sectors a lot, but are seen very little in

the CTE and CTS sectors (Fig. 4). Applying the simple-signed rule and the blockmodeling

algorithm to delineate communities, both find that there are three distinct communities (Fig.

4). Though, in the case of AVS the simple sign-rule suggested multi-membership, while the

blockmodeling algorithm found AVS to be part of the cluster containing (AVO, AVN, AVE)

or that the two clusters (orange and purple in fig. 4) merged into one depending on the choice

of weighting parameter (i.e., emphasizing positive or negative edges).

Figure 4: Similarity and dissimilarity between sectors in the beluga whale population of the

St. Lawrence. The green edges between two sectors signify that the sectors share high/low

users, while red edges signify that they have dissimilar high/low users. The lack of an edge

signifies that the high/low users of one sector does not provide information about the

high/low users of other sectors. Nodes represent sectors, and are coloured based on shared

communities: i.e., shared green edges, and no red edges. Node sizes represent the magnitudes

of individual differences in use within the sector, i.e., larger nodes suggest larger differences

between high and low users.

4. Discussion

Here we've shown that using photo-ID data with a multilevel multinomial model it is possible to estimate socio-spatial networks, identifying spatial communities while controlling for sampling biases. Our results suggest that the beluga population shows a non-random social-spatial structuring within the summer range.

Within sectors of the St. Lawrence, our model suggests that subsets of belugas are heavily using some sectors, while other sectors show little evidence of differences in use. The magnitude of individual differences in each sector, i.e., how much individuals differ in their probability of being observed in a particular sector, shows that the CTN sector, in particular, has very little in the way of individual differences in the probability of being seen in that sector (Table 1). This result suggests that the CTN sector is used similarly by most individuals, and represents a potential high mixing zone for the population. In contrast, the AME sector shows the highest level of individual differences, suggesting that there are large differences in how beluga are using this sector. These results suggest that the population is not randomly mixing with the St. Lawrence, and that there are belugas that make use of some sectors more than other belugas.

Between sectors of the St. Lawrence, our results add to the evidence that the beluga population cannot be assumed to be randomly mixing within its summer habitat. Rather, comparing the individual differences in beluga usage patterns within sectors suggests similar/dissimilar user populations across sectors (Fig. 4). By using correlations between sector usage patterns to create a socio-spatial network, and running community detection algorithms, our results found that there are spatially distinct communities that make use of particular regions of the St. Lawrence and the Saguenay River (Fig. 4). We found that the beluga population in the St. Lawrence could be separated into three distinct communities: 1)

The lower St. Lawrence (AVO, AVS, AVE, AVN), the Saguenay River and mouth (BSM,

SAG, CTN), and the upper and eastern portion of the St. Lawrence (CTE, CTS, CTO, AME,

AMS) (Fig. 4).

Our findings have direct implications for estimating the impacts of anthropogenic

disturbances on this population. As the population shows evidence of spatially restricted

habitat use, disturbances to particular regions can have a disproportionate impact on

particular segments of the larger population. In particular, the cumulative impacts over time

are likely to be greatly increased in some segments while reduced in other segments of the

population, altering estimations of the distribution of impacts. If cumulative impacts, such as

noise, or environmental contaminants, have a threshold beyond which individual survival is

greatly reduced, properly estimating the distribution of cumulative impacts can have large

implications for conservation management. Our results add to the current understanding of

socio-spatial structuring within this population (Michaud, 1993, 2005), and suggest that more

empirical data, e.g., photo-ID data, movement data, aerial surveillance, should be collected to

better refine socio-spatial mixing in this population.

The modeling approach presented in this paper relies on defined sectors within a

particular spatial range, e.g., SAG sector, CTN sector… etc (Fig. 1). In some cases, these

delineations can be justified as they identify management zones, but in other cases, the

delineation and scale of these sectors can be delineated somewhat arbitrarily. Future work

could assess the use of continuous random effects (as opposed to categorical) where

individual differences in the probability of being seen could be on a continuous surface. Point

estimates of individual differences could then be estimated at any location, and correlations

between individual differences obtained between any two points in continuous space. This

approach could avoid reliance on user-defined sectors and facilitate a means of looking at the

results at different scales (e.g., grids of points at various scales could be used when estimating

correlations). Similarly, given that the method requires repeated sampling of individuals to

obtain probability of being seen in any one sector, there is a reliance on longitudinal data. In

well sampled populations, it could be feasible to estimate the change in time of individuals

being seen in particular sectors. Here differences in being seen in any particular area could

be explicitly modeled to capture any temporal changes in community substructures, or

developmental trajectories related to habitat use.

In terms of implementing the multilevel multinomial model on other photo-ID

datasets, the use of test datasets should hold a prominent role in the analysis. The use of

permutation/randomization methods to both generate structured and unstructured datasets,

while maintaining the sample size distribution of the original datasets, can be very valuable in

helping to set model priors and to interpret the final model results. The use of permutation

approaches is common in social network analysis (Croft et al., 2011; Farine, 2017), and is

becoming more common in statistical workflows more generally (Gelman et al., 2013;

McElreath, 2020).

5. Conclusions

We have introduced the use of multilevel multinomial modeling to estimate socio-

spatial networks from photo-ID data. We've shown, using testing datasets, that the proposed

method is effective at detecting socio-spatial structures. When applied to 18 years of photo-

ID data from an endangered population of beluga whales in the St. Lawrence, our results

suggest strong evidence that the population has three distinct spatial communities. We

suggest that multilevel multinomial models can be effective in extracting socio-spatial

structuring within animal populations monitored by photo-ID, and can have direct

implications for conservation management.

6. Acknowledgment

We would like to thank all those who worked with and supported the GREMM in collecting these long term data.

7. Funding

Funding was provided by Ministère des Forêts, de la Faune et des Parcs du Québec and Secrétariat à la stratégie maritime du Québec.

8. Data accessibility

The permuted and simulated photo ID datasets are available on github (github.com/tbonne/photoID_multinomial), along with code used in the analysis.

9. Author Contributions

RM collected the data; TRB conceived the analytical methodology and performed the analysis; all authors contributed critically to the drafts and gave final approval for publication.

References:

Chion, C., Lagrois, D., Dupras, J., Turgeon, S., McQuinn, I. H., Michaud, R., Ménard, N., &

Parrott, L. (2017). Underwater acoustic impacts of shipping management measures:

Results from a social-ecological model of boat and whale movements in the St.

Lawrence River Estuary (Canada). *Ecological Modelling*, *354*, 72–87.

https://doi.org/10.1016/j.ecolmodel.2017.03.014

Croft, D. P., Madden, J. R., Franks, D. W., & James, R. (2011). Hypothesis testing in animal

social networks. *Trends in Ecology & Evolution*, *26*(10), 502–507.

DeFur, P. L., Evans, G. W., Hubal, E. A. C., Kyle, A. D., Morello-Frosch, R. A., & Williams,

D. R. (2007). Vulnerability as a function of individual and group resources in

cumulative risk assessment. *Environmental Health Perspectives*, *115*(5), 817–824.

Doreian, P., & Mrvar, A. (2015). Structural balance and signed international relations.

Journal of Social Structure, *16*, 1.

Evans, P. G., & Hammond, P. S. (2004). Monitoring cetaceans in European waters. *Mammal

Review*, *34*(1–2), 131–156.

Farine, D. R. (2017). A guide to null models for animal social network analysis. *Methods in

Ecology and Evolution*, *8*(10), 1309–1320.

Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., & Rubin, D. B. (2013).

Bayesian data analysis. CRC press.

Hupman, K., Stockin, K. A., Pollock, K., Pawley, M. D. M., Dwyer, S. L., Lea, C., &

Tezanos-Pinto, G. (2018). Challenges of implementing Mark-recapture studies on

poorly marked gregarious delphinids. *PLOS ONE*, *13*(7), e0198167.

https://doi.org/10.1371/journal.pone.0198167

Koivuniemi, M., Auttila, M., Niemi, M., Levänen, R., & Kunnasranta, M. (2016). Photo-ID

as a tool for studying and monitoring the endangered Saimaa ringed seal. *Endangered*

Species Research, *30*, 29–36.

Koster, J., & McElreath, R. (2017). Multinomial analysis of behavior: Statistical methods.

Behavioral Ecology and Sociobiology, *71*(9), 138.

Lesage, V., McQuinn, I. H., Carrier, D., Gosselin, J.-F., & Mosnier, A. (2014). Exposure of

the beluga (Delphinapterus leucas) to marine traffic under various scenarios of transit

route diversion in the St. Lawrence Estuary. *DFO Can. Sci. Advis. Sec., Res. Doc.*

2013/125. Iv, 28.

McElreath, R. (2020). *Statistical rethinking: A Bayesian course with examples in R and Stan*.

CRC press.

McQuinn, I. H., Lesage, V., Carrier, D., Larrivée, G., Samson, Y., Chartrand, S., Michaud,

R., & Theriault, J. (2011). A threatened beluga (Delphinapterus leucas) population in

the traffic lane: Vessel-generated noise characteristics of the Saguenay-St. Lawrence

Marine Park, Canada. *The Journal of the Acoustical Society of America*, *130*(6),

3661–3673. https://doi.org/10.1121/1.3658449

Michaud, R. (1993). *Distribution estivale du béluga du Saint-Laurent: Synthèse 1986 à 1992*.

Ministère des pêches et des océans, Direction de la gestion des pêches et de ….

Michaud, R. (2005). Sociality and ecology of the odontocetes. *Sexual Segregation in*

Vertebrates, 303–326.

Parrott, L., Chion, C., Martins, C. C. A., Lamontagne, P., Turgeon, S., Landry, J. A., Zhens,

B., Marceau, D. J., Michaud, R., Cantin, G., Ménard, N., & Dionne, S. (2011). A

decision support system to assist the sustainable management of navigation activities

in the St. Lawrence River Estuary, Canada. *Environmental Modelling & Software*,

26(12), 1403–1418. https://doi.org/10.1016/j.envsoft.2011.08.009

Perryman, R. J. Y., Venables, S. K., Tapilatu, R. F., Marshall, A. D., Brown, C., & Franks, D.

W. (2019). Social preferences and network structure in a population of reef manta

rays. *Behavioral Ecology and Sociobiology*, *73*(8), 114.

https://doi.org/10.1007/s00265-019-2720-x

Schilds, A., Mourier, J., Huveneers, C., Nazimi, L., Fox, A., & Leu, S. T. (2019). Evidence

for non-random co-occurrences in a white shark aggregation. *Behavioral Ecology and*

Sociobiology, *73*(10), 138. https://doi.org/10.1007/s00265-019-2745-1

Schneider, S., Taylor, G. W., Linquist, S., & Kremer, S. C. (2019). Past, present and future

approaches using computer vision for animal re-identification from camera trap data.

Methods in Ecology and Evolution, *10*(4), 461–470. https://doi.org/10.1111/2041-

210X.13133

495
496

Supplementary material:

Use of a preset reference sector:

We use the simulated dataset with structure, presented in the main text, to fit a

multilevel multinomial model with and without a preset reference sector. That is, in the

model with the preset reference sector we duplicated each individual's photo-IDs and placed

them in the reference sector. This results in a probability of 0.5 for being seen in the reference

sector for all individuals. The results suggest that using a preset reference sector where all

individuals have the same probability of being seen is required to estimate correlations

between sectors and produces appropriate socio-spatial networks (Fig S1).

Fig S1: The estimate socio-spatial networks for a multilevel multinormial model a) without a

preset reference sector, i.e., AME is used as the reference, and b) with a preset reference

sector. The green edges between two sectors signify that the sectors share high/low users,

while red edges signify that they have dissimilar high/low users. The lack of an edge signifies

that the high/low users of one sector does not provide information about the high/low users of

other sectors. Nodes represent sectors, and are coloured based on the communities imposed

when simulating the data.

Full model table:

Table S1: Parameter estimates from the multilevel multinomial model predicting probability

of capturing a photo-ID by sector. Estimated mean probability (logit scale), individual

differences in mean probability ('sd'), and correlation of individual differences between the

different sectors are presented along with an estimate of their 95% credible intervals. Positive

correlations suggest that the high/low users in one sector are similarly high/low users in

another sector, while negative correlations suggest high/low users in one sector are the

low/high users in another sector.

Type	Parameter	Estimate	SD	l-95% CI	u-95% CI
Mean probability (logit scale)					
	mu_AME	-5.85	0.28	-6.43	-5.33
	mu_AMN	-5.60	0.24	-6.15	-5.17
	mu_AMO	-6.68	0.34	-7.41	-6.08
	mu_AMS	-6.33	0.33	-7.00	-5.73
	mu_AVE	-5.47	0.24	-5.99	-5.02
	mu_AVN	-5.06	0.22	-5.52	-4.65
	mu_AVO	-1.82	0.05	-1.93	-1.72
	mu_AVS	-2.90	0.07	-3.04	-2.76
	mu_BSM	-3.78	0.13	-4.04	-3.52
	mu_CTE	-2.20	0.05	-2.30	-2.10
	mu_CTN	-1.24	0.03	-1.30	-1.18
	mu_CTO	-4.58	0.16	-4.92	-4.28

mu_CTS	-4.09	0.13	-4.35	-3.85
mu_SAG	-2.78	0.09	-2.96	-2.61

Individual Differences

sd(mu_AME)	2.00	0.19	1.64	2.38
sd(mu_AMN)	0.76	0.29	0.14	1.31
sd(mu_AMO)	0.65	0.38	0.05	1.41
sd(mu_AMS)	1.53	0.22	1.10	1.98
sd(mu_AVE)	1.16	0.20	0.78	1.57
sd(mu_AVN)	1.30	0.17	0.98	1.65
sd(mu_AVO)	0.78	0.05	0.68	0.88
sd(mu_AVS)	0.83	0.07	0.70	0.97
sd(mu_BSM)	1.30	0.11	1.09	1.52
sd(mu_CTE)	0.78	0.05	0.68	0.89
sd(mu_CTN)	0.33	0.03	0.27	0.39
sd(mu_CTO)	1.27	0.14	1.00	1.55
sd(mu_CTS)	1.18	0.11	0.97	1.40
sd(mu_SAG)	1.16	0.08	1.01	1.31

Correlations between individual differences

cor(mu_AME,mu_AMN)	0.39	0.19	-0.03	0.71
cor(mu_AME,mu_AMO)	0.27	0.24	-0.26	0.67
cor(mu_AMN,mu_AMO)	0.14	0.24	-0.36	0.57
cor(mu_AME,mu_AMS)	0.63	0.11	0.39	0.82
cor(mu_AMN,mu_AMS)	0.25	0.21	-0.19	0.62
cor(mu_AMO,mu_AMS)	0.27	0.24	-0.26	0.67
cor(mu_AME,mu_AVE)	0.18	0.16	-0.14	0.46
cor(mu_AMN,mu_AVE)	0.15	0.21	-0.28	0.56

cor(mu_AMO,mu_AVE)	0.10	0.23	-0.36	0.52
cor(mu_AMS,mu_AVE)	0.05	0.19	-0.32	0.41
cor(mu_AME,mu_AVN)	-0.04	0.15	-0.33	0.25
cor(mu_AMN,mu_AVN)	-0.02	0.21	-0.42	0.39
cor(mu_AMO,mu_AVN)	0.00	0.22	-0.42	0.44
cor(mu_AMS,mu_AVN)	-0.07	0.18	-0.41	0.28
cor(mu_AVE,mu_AVN)	0.50	0.14	0.19	0.75
cor(mu_AME,mu_AVO)	-0.21	0.10	-0.40	0.00
cor(mu_AMN,mu_AVO)	0.03	0.19	-0.35	0.40
cor(mu_AMO,mu_AVO)	-0.06	0.21	-0.45	0.36
cor(mu_AMS,mu_AVO)	-0.28	0.14	-0.54	0.01
cor(mu_AVE,mu_AVO)	0.48	0.13	0.22	0.71
cor(mu_AVN,mu_AVO)	0.50	0.10	0.28	0.69
cor(mu_AME,mu_AVS)	0.13	0.12	-0.10	0.36
cor(mu_AMN,mu_AVS)	0.21	0.18	-0.16	0.53
cor(mu_AMO,mu_AVS)	0.13	0.21	-0.31	0.50
cor(mu_AMS,mu_AVS)	0.03	0.15	-0.26	0.32
cor(mu_AVE,mu_AVS)	0.40	0.14	0.12	0.66
cor(mu_AVN,mu_AVS)	0.38	0.12	0.14	0.60
cor(mu_AVO,mu_AVS)	0.54	0.08	0.38	0.69
cor(mu_AME,mu_BSM)	-0.57	0.09	-0.75	-0.38
cor(mu_AMN,mu_BSM)	-0.37	0.17	-0.66	0.00
cor(mu_AMO,mu_BSM)	-0.22	0.23	-0.62	0.26
cor(mu_AMS,mu_BSM)	-0.37	0.13	-0.62	-0.10
cor(mu_AVE,mu_BSM)	-0.34	0.13	-0.59	-0.06
cor(mu_AVN,mu_BSM)	-0.22	0.12	-0.44	0.02

cor(mu_AVO,mu_BSM)	-0.16	0.08	-0.32	0.01
cor(mu_AVS,mu_BSM)	-0.66	0.07	-0.80	-0.51
cor(mu_AME,mu_CTE)	0.54	0.10	0.34	0.72
cor(mu_AMN,mu_CTE)	0.34	0.18	-0.04	0.65
cor(mu_AMO,mu_CTE)	0.21	0.23	-0.29	0.60
cor(mu_AMS,mu_CTE)	0.38	0.14	0.10	0.62
cor(mu_AVE,mu_CTE)	0.07	0.14	-0.21	0.36
cor(mu_AVN,mu_CTE)	-0.11	0.12	-0.35	0.14
cor(mu_AVO,mu_CTE)	-0.10	0.08	-0.26	0.07
cor(mu_AVS,mu_CTE)	0.52	0.08	0.34	0.67
cor(mu_BSM,mu_CTE)	-0.80	0.06	-0.89	-0.68
cor(mu_AME,mu_CTN)	-0.26	0.12	-0.49	-0.03
cor(mu_AMN,mu_CTN)	-0.24	0.19	-0.59	0.16
cor(mu_AMO,mu_CTN)	-0.12	0.21	-0.50	0.32
cor(mu_AMS,mu_CTN)	-0.09	0.15	-0.39	0.21
cor(mu_AVE,mu_CTN)	-0.53	0.13	-0.77	-0.26
cor(mu_AVN,mu_CTN)	-0.44	0.12	-0.67	-0.18
cor(mu_AVO,mu_CTN)	-0.75	0.07	-0.86	-0.60
cor(mu_AVS,mu_CTN)	-0.63	0.09	-0.79	-0.44
cor(mu_BSM,mu_CTN)	0.47	0.09	0.28	0.64
cor(mu_CTE,mu_CTN)	-0.25	0.10	-0.46	-0.05
cor(mu_AME,mu_CTO)	0.63	0.09	0.43	0.79
cor(mu_AMN,mu_CTO)	0.28	0.20	-0.15	0.63
cor(mu_AMO,mu_CTO)	0.16	0.22	-0.30	0.56
cor(mu_AMS,mu_CTO)	0.55	0.13	0.27	0.78
cor(mu_AVE,mu_CTO)	-0.12	0.17	-0.45	0.22

cor(mu_AVN,mu_CTO)	-0.24	0.15	-0.54	0.07
cor(mu_AVO,mu_CTO)	-0.55	0.09	-0.72	-0.36
cor(mu_AVS,mu_CTO)	-0.16	0.12	-0.39	0.07
cor(mu_BSM,mu_CTO)	-0.33	0.11	-0.53	-0.12
cor(mu_CTE,mu_CTO)	0.39	0.11	0.17	0.60
cor(mu_CTN,mu_CTO)	0.19	0.12	-0.06	0.42
cor(mu_AME,mu_CTS)	0.70	0.08	0.53	0.85
cor(mu_AMN,mu_CTS)	0.42	0.19	0.00	0.73
cor(mu_AMO,mu_CTS)	0.21	0.23	-0.27	0.61
cor(mu_AMS,mu_CTS)	0.42	0.14	0.14	0.67
cor(mu_AVE,mu_CTS)	0.16	0.16	-0.15	0.45
cor(mu_AVN,mu_CTS)	0.06	0.14	-0.23	0.33
cor(mu_AVO,mu_CTS)	-0.04	0.10	-0.23	0.15
cor(mu_AVS,mu_CTS)	0.42	0.11	0.20	0.61
cor(mu_BSM,mu_CTS)	-0.72	0.08	-0.85	-0.55
cor(mu_CTE,mu_CTS)	0.64	0.09	0.46	0.80
cor(mu_CTN,mu_CTS)	-0.37	0.11	-0.57	-0.14
cor(mu_CTO,mu_CTS)	0.48	0.11	0.25	0.68
cor(mu_AME,mu_SAG)	-0.43	0.09	-0.61	-0.24
cor(mu_AMN,mu_SAG)	-0.36	0.17	-0.66	0.00
cor(mu_AMO,mu_SAG)	-0.17	0.22	-0.56	0.29
cor(mu_AMS,mu_SAG)	-0.22	0.13	-0.47	0.03
cor(mu_AVE,mu_SAG)	-0.44	0.12	-0.68	-0.19
cor(mu_AVN,mu_SAG)	-0.37	0.10	-0.57	-0.16
cor(mu_AVO,mu_SAG)	-0.48	0.06	-0.60	-0.35
cor(mu_AVS,mu_SAG)	-0.80	0.06	-0.90	-0.68

cor(mu_BSM,mu_SAG)	0.83	0.04	0.74	0.91
cor(mu_CTE,mu_SAG)	-0.68	0.06	-0.79	-0.55
cor(mu_CTN,mu_SAG)	0.67	0.07	0.52	0.80
cor(mu_CTO,mu_SAG)	-0.12	0.11	-0.32	0.09
cor(mu_CTS,mu_SAG)	-0.62	0.08	-0.77	-0.45

Note on interpreting low sd within sectors:

Similar to the CTN sector, the sector AMO is also estimated to have a low magnitude of

individual differences. However, it has a large uncertainty in this estimate. This highlights

that it is possible to have little individual differences in a sector due to either: 1) limited data,

resulting in all individuals being pooled to the mean value, and 2) limited data is not a factor,

but individuals are using this sector relatively equally. Care should therefore be taken when

interpreting the magnitude of individual differences, nevertheless, the estimated uncertainty

around magnitude estimates is one way to help identify sectors with limited data.